Integrative analysis of hexaploid wheat roots identifies signature components during iron starvation

Gazaldeep Kaur[1,2*], Vishnu Shukla[1,3*], Anil Kumar[1,2], Mandeep Kaur[1,2], Parul Goel[1],

Palvinder Singh[1], Anuj Shukla[1], Varsha Meena[1], Jaspreet Kaur[3], Jagtar Singh[2], Shrikant

Mantri[1], Hatem Rouached[4] and Ajay Kumar Pandey[1#]

Author affiliations

[1]National Agri-Food Biotechnology Institute (Department of Biotechnology), Sector 81,

Knowledge City, Mohali-140306, Punjab, India.

[2] Department of Biotechnology, Panjab University, Chandigarh, India.

[3]University Institute of Engineering and Technology, Panjab University, Chandigarh, India

[4]BPMP, Université de Montpellier, INRA, CNRS, Montpellier SupAgro, 34060

Montpellier, France.

[*]*Authors contributed equally*

[#]*Corresponding author*

Dr. Ajay K Pandey, Scientist-E.

National Agri-Food Biotechnology Institute (Department of Biotechnology), Sector 81,

Knowledge City, Mohali-140306, Punjab, India.

Telephone: +91-1724990124

Email: pandeyak@nabi.res.in; pandeyak1974@gmail.com

ORCID iD: https://orcid.org/0000-0003-1064-139X

Running Title: Fe starvation response in hexaploid wheat roots

Highlight: During Fe starvation, wheat roots show prolific expression of core components

involved in Strategy-II mode of Fe uptake along with significant changes in the metabolome

including enhanced GST activity.

Abstract

Iron is an essential micronutrient for all organisms. In crop plants, iron deficiency can decrease crop yield significantly, however our current understanding of how major crops respond to iron deficiency remains limited. Herein, the effect of Fe deprivation at both the transcriptomic and metabolic levels in hexaploid wheat was investigated. Genome-wide gene expression reprogramming was observed in wheat roots subjected to Fe starvation, with a total of 5854 genes differential expressed. Homoeolog and subgenome specific analysis unveiled induction bias contribution from the A and B genomes. In general, the predominance of genes encoding for nicotianamine synthase, yellow stripe like transporters, metal transporters, ABC transporters and zinc-induced facilitator-like protein was noticed. Expression of genes related to the strategy-II mode of Fe uptake was predominant as well. Our transcriptomic data were in agreement with the GC-MS analysis that showed the enhanced accumulation of various metabolites such as fumarate, malonate, succinate and xylofuranose, which could be contributing to Fe-mobilization. Interestingly, Fe starvation leads to significant temporal increase of glutathione-S-transferase both at transcriptional and in enzymatic activity levels, which indicates the involvement of glutathione in response to Fe stress in wheat roots. Taken together, our result provides new insight into the wheat response to Fe starvation at molecular level and lays foundation to design new strategies for the improvement of Fe nutrition in crops.

Key words: iron starvation, *Triticum aestivum*, glutathione metabolism, transcriptome, gene expression, genome bias.

Introduction

Iron (Fe) is among the essential micronutrients in plants that participate as catalytic cofactors in several key processes including photosynthesis, respiration, chlorophyll biosynthesis, and nitrogen fixation (Kim and Rees, 1992; Morrissey and Guerinot, 2009; Li et al., 2017). The bioavailability of Fe in the soil is strongly dependant on its solubility, aerobic and calcareous soil condition, pH, and the presence of natural ligands secreted by plant roots (Marschner H., 1995; Morrissey and Guerinot, 2009; Thomine et al., 2013). To circumvent the above challenges for Fe uptake by the roots, plants have adapted two modes of uptake strategies. Strategy-I, mostly predominant in Eudicot species, primarily relies on the enrichment of rhizospheric regions with protons (H^+) and other reducing agents (Hell and Stephan, 2003; Santi et al., 2005; Santi and Schmidt 2009; Kobayashi and Nishizawa 2012). In contrast, graminaceous species follow the Strategy-II mode of uptake, which involves the transport of Fe in the complex-chelated form (Mori et al., 1999; Kobayashi and Nishizawa, 2012; Connorton et al., 2017). The primary components involved in chelation are the phytosiderophores (PS) secreted by plant cells in the rhizospheric region mainly by efflux transporters (Morrissey and Gueirnot, 2009; Kobayashi and Nishizawa, 2012). One of the main components of these secreted siderophores involved in Fe chelation is referred to as mugineic acids (MAs). The transporters involved in the secretion of the MAs are identified as transporter of mugineic (TOM) acid proteins (Nozoye et al., 2011). The complex chelated form of Fe^{3+}-PS is taken up by the specific root transporter referred to as yellow stripe-like transporter proteins (YSL) (Curie et al., 2001; Gross et al., 2003, Yordem et al., 2011). Subsequently, Fe is transported in the plant organelles by multiple partners including specialized long distance, tissue-specific transporters as reported in monocots (e.g rice, maize) and eudicots (e.g. Arabidopsis) (Kim et al., 2006; Waters et al., 2011). In other monocots such as maize and rice, the presence of genetic components for both Strategy-I and II were reported (Ishimaru et al., 2006; Inoue et al., 2009; Lee et al., 2009; Zanin et al., 2017). At the metabolome level, analytical approaches utilizing GC-MC and LC-MS were utilized to study the components of Fe starvation in plants (Palmer et al., 2014; Kabir et al., 2013). However, the molecular and metabolic activity predominant for Fe uptake by the roots of hexaploid wheat still remains to be elucidated.

Microarrays have been successfully used to investigate the global transcriptional changes in Arabidopsis plants grown in Fe starved conditions (Thimm et al., 2001; Buckhout et al., 2009; Yang et al., 2010). Significant changes in the expression pattern of the genes primarily involved in Strategy-II mode of uptake were observed in roots of maize and rice

subjected to Fe-deficiency stress (Li et al., 2014; Quinet et al., 2012; Kobayashi et al., 2014;

Bashir et al., 2014). The transcriptome analysis of Fe-starved plants is characterised by an

important change in gene expression of several transcriptional regulators such as transcription

factors (TFs) (Colangelo et al., 2004; Kobayashi et al., 2009; Long et al., 2010; Li et al.,

2016; Connorton et al., 2017) or key genes related to phytohormone homeostasis (Schmidt et

al., 2000; Hindt and Guerinot, 2012). Fe starvation also leads to changes in the abundance of

transcripts related to plant metabolism and genes involved in signalling pathways that

modulate nutrient uptake. Additionally, genes involved in ethylene/auxin signalling or linked

with certain other macronutrients like nitrogen, sulphur and phosphorus are also significantly

expressed (Romera et al., 1994; Zheng et al., 2009; Romera et al., 2011; Borlotti et al., 2012;

Zuchi et al., 2015; Lin et al., 2016; Zanin et al., 2017; Garnica et al., 2018). Therefore,

transcriptome analysis is a powerful approach to help understand the network involved in

plants response to Fe.

115   Hexaploid wheat is an important crop that is also a good, affordable source of

nutrition. Fe deficiency strongly affects the growth of cereal crops and productivity (Yousfi

et al., 2009; Bocchini et al., 2015). The development of transcriptome technology such as

RNA sequencing-RNAseq and the availability of genome sequence for hexaploid wheat

(genome A, B, D) (International Wheat Genome Consortium, 2014) combined with

metabolomic approaches will help improve our knowledge of wheat response to nutritional

stress (e.g. –Fe) (Borrill et al., 2018). Our current knowledge on how the wheat genome

responds to Fe limiting condition remains obscure and unanswered. In addition, how the

different genomes could impact the homoeolog based expression of the transcripts,

categorically under different stress conditions, remains to be explored (Ramirez-Gonzalez et

al., 2018). In the current study, RNA-seq based approach was utilized to address the

molecular components involved in wheat roots during Fe starved condition. Further, genome-

based expression study was performed that provided preliminary clue on coordinated

expression of the homoeologs. Metabolic profiling result support the data from the

transcriptome study and pinpoint the role of glutathione mediated response. Collectively our

results provide the first insight on molecular and biochemical responses of hexaploid wheat

under Fe starvation.

Materials and methods

Plant materials, starvation conditions and plant sampling

Hexaploid bread wheat variety 'C-306' was used for the experimental purpose. Wheat grains

were subjected to stratification in dark at 4 ^{0}C for overnight and were allowed to germinate

for 5 days on the Petri plates containing 2-3 layers of wet Whatman filter paper. The

endosperms were removed from the developing seedlings once they started browning.

Subsequently, the seedlings were transferred in PhytaBox^{TM} and grown for 7 days in

Hoagland's nutrient solution. A total of forty-eight seedlings were kept equally in four

PhytaBoxes, used for each treatment. After 7 days, nutrient solutions were replaced on the

basis of different treatments. For Fe starvation (–Fe) 2 µM Fe (III) EDTA was used as the Fe

source. For control plants (+Fe) concentrations of nutrients were unchanged in above-

mentioned Hoagland's solution containing (20 µM Fe (III) EDTA). Treated seedlings were

grown in the described medium for 20 days in a growth chamber set at 20±1 °C, 50-70%

relative humidity and photon rate of 300 µmol quanta m^{-2} s^{-1} with 16 h day/8 h night

cycle. For sampling, roots and shoots were collected at different time points after starvation

(5, 10, 15 and 20 days). On the basis of distinct phenotype, samples collected at 20 days after

starvation (DAS) were used for transcriptome analysis. A total of four biological replicates

(each containing 10-12 seedlings) were used. Subsequently, RNA extractions were performed

independently for each of the pools. Prior to RNA sequencing, quality of RNA was checked

and extractions derived from two replicates were pooled together thereby generating two

experimental samples for each of the respective conditions. Remaining samples were snap

frozen in liquid nitrogen and stored at -80°C. To distinctively observe primary root and 1st

order lateral root, individual plants were moved onto a 150 mm wide petri plate filled with

distilled water and characteristics were manually examined. To ascertain root characteristics,

6-8 wheat seedlings were used for each of the treatments with two experimental replicates.

RNA extraction and Illumina sequencing

Total RNA was extracted from the treated root samples along with the control by Trizol

(Invitrogen, ThermoFisher, USA) as per the manufacturer's instruction. The quality and

quantity were checked on the 1% denaturing RNA agarose gel and Nanodrop respectively.

Subsequently, all the RNA used for library preparations were checked for their RNA integrity

number ≥ 8.5 using Bioanalyzer (Agilent, USA). The quality control (QC) passed RNA

samples were then processed for library preparation. The Paired-ended (PE) libraries were

prepared from the total RNA using Illumina TruSeq stranded mRNA library prep kit as per

the instructions (Illumina Inc*.,* USA). The generated mean of the libraries fragment size

distribution was 559 bp, 584 bp, 546 bp and 604 bp for the samples. The generated libraries

were sequenced on the NextSeq 500 using 2 X 150 bp chemistry. The raw reads were

processed further before the sorted high-quality reads were mapped to the reference genome.

RNA-Seq analysis

Adapter clipping and quality trimming of the raw reads were performed using Trimmomatic-

0.35. The sequenced raw reads were processed to obtain high quality clean reads. Ambiguous

reads (reads with unknown nucleotides "N" larger than 5%) and low-quality sequences (more

than 10% quality threshold QV<20 phred score) were removed. A minimum length of 100

nts after trimming was applied. Finally, high-quality (QV>20), paired-end reads were used

for reference based read mapping. The genome of *Triticum aestivum* L. was taken as

reference for analysis. The Gene Feature Format files were downloaded from Ensembl Plants

(TGACv1.37). The reads were mapped to the reference genome using TopHat v2.1.1 with

default parameters. Cufflinks v2.2.1 program assembled the transcriptome data from RNA-

seq data and quantified their expression. Mapped reads were subjected to Cufflinks, followed

by Cuffmerge and Cuffdiff. Log2 Fold Change (FC) values greater than one was considered

up-regulated whereas less than 1 were considered as down-regulated. These genes were

further categorized on the basis of statistical significance (p<0.05) and the False Discovery

Rate (FDR 0.05) after Benjamin-Hochberg corrections for multiple testing for their

significant expression.

Comprehensive gene annotation of wheat sequences was done using KOBAS 3.0 (Xie

et al., 2011) annotate module by alignment with Rice sequences (BLASTP, cutoff 1e-5).

MapMan was used to visualize the pathways involving wheat differentially expressed genes

(DEGs). Pathway enrichment analysis was performed using KOBAS standalone tool. MeV

was used to construct heatmaps for selected DEGs using the normalized expression values of

genes. The data generated from this study has been deposited in the NCBI Sequence Read

Archive (SRA) database and is accessible with the submission ID- SUB5206887 and

BioProjectID-PRJNA529036.

Gene annotation filtering and functional enrichment analysis

Significant sets of DEGs under iron starvation were further mapped using GO and MapMan.

The GO annotation was downloaded from ensembl plants

(https://plants.ensembl.org/biomart/martview). Mercator (Lohse et al., 2014) was used to

build MapMan mapping file for TGACv1 sequences and DEGs were visualized in MapMan

v3.1.1 tool (Thimm et al., 2004). For functional categorization of DEGs that were positively

and negatively correlated with iron starvation, BINGO version 3.0.3 (Maere et al., 2005) was

used to perform GO enrichment analysis with hypergeometric test and significant terms with

an FDR value below 0.05 were considered. For gene ontology mapping, GO_full.obo

ontology file was downloaded from GO consortium. MapMan classification was used to

categorize DEGs into transcriptional factors. The results were visualized as network using

EnrichmentMap version 2.2.1 (Merico et al., 2010) and gene expression overview in various

pathways were visualized in MapMan tool.

Homoeolog specific expression analysis for genome bias

To identify wheat homoeologous triplets; ensembl biomart TGACv1 was used to extract all

possible homoeologous relations. This led to 86830 pairwise homoeologous relations. An in-

house script was used to select only the triplets where contribution from each genome was

1:1:1 (A:B:D), thus generating 16850 triplets. Further, triplets resulting from potential

translocation events were not considered, i.e., only homoeolog triplets from same

chromosome (eg., 1A, 1B, 1D triplet is accepted, whereas, 2A, 3B, 3D is rejected) were taken

for analysis. Finally, 15604 triplets (15604*3 = 46812 genes) were used for studying genome

induction biasness in response to iron stress. Paired end reads were aligned to the reference

(selected scaffolds from genome that harbour 15604 triplets) using TopHat v2.1.1 with a

specific argument (--b2-very-sensitive) (Powell et al., 2017), which leads to more stringent

alignments as required for homoeologs. The Cufflinks pipeline was used to obtain FPKM

values and differentially expressed genes.

Further, relative abundance levels and expression bias for homoeologs was studied by

considering the homoeologous gene triplets with expression of FPKM ≥ 1 each in both the

control as well as Fe starved conditions. For this, the normalised relative expression for each

homoeolog within a triad was calculated. For example, the relative expression from A will be

represented as:

$$Relative\ expression_A = \frac{\text{FPKM}_A}{FPKM_A + FPKM_B + FPKM_D}$$

Thus, relative expression levels of A, B and D homoeologs within each triad were calculated

similarly. Seven homoeolog expression bias categories were defined as described earlier by

Ramirez-Gonzalez et al., (2018). In total, seven categories defined as one balanced category

and six unbalanced homoeolog-suppressed/homoeolog-dominant (from either of the

genomes) were listed for the ideal relative expression of A, B and D. Eucledian distance

(using cdist function from rdist package, R3.3.2) was calculated between normalised relative

expression for each triad and the seven ideal categories. The shortest distance was used as a

deciding factor to group the triads into the seven respective categories.

Quantitative real time-PCR (qRT-PCR)

For validation of the gene expression, qRT-PCR analysis was performed. Total RNA (2 µg)

isolated from the above experiments were used to validate the expression by using qRT-PCR

method. Genomic DNA present in trace amount was removed by DNaseI treatment using a

Turbo DNA-free kit (Invitrogen, ThermoFisher, USA). Further, cDNA was synthesized from

two micrograms of DNA-free RNA using Superscript III first strand (Invitrogen,

ThermoFisher, USA) with random hexamer primers following the manufacturer's guidelines.

For qPCR reaction, gene-specific primers of each gene were designed from conserved

regions of all three homeolog sequences (Supplementary Table S1). qRT-PCR was

performed using QuantiTect SYBR Green RT-PCR mastermix (Qiagen, USA) with programs

recommended by the manufacturer in the ABI 7700 sequence detector (Applied Biosystems,

USA). ADP-ribosylation factor (ARF) and Actin were used as internal controls. Two

independent experimental replicates with four technical replicates were performed for each

sample. The relative amount of gene expression was calculated using $2^{-\Delta\Delta CT}$ method.

Metabolite extraction and GC-MS profiling

Extraction of total metabolites was performed similarly as previously described (Wang et al.,

2018). Roots of plants grown under –Fe and +Fe were sampled in triplicate manner and dried

for 1 week. Each of 50 mg crushed samples was extracted in a mixture of solvents

(precooled; 300 µl methanol, 30 µl 2 mg ml^{-1} nonadecanoic acid methylester and 30 µl 0.2

mg ml^{-1} sorbitol) for 15 minutes (70°C, 1000rpm). Further, at room temperature 200 µl

chloroform was added and shaken for 5 min (37°C, 1000rpm). To obtain phase separation,

400 µl H$_2$O was added to each sample, vortexed and centrifuged (10 min, 13,000 rpm). From

the upper polar phase approximately 200 µl was finally aliquoted for complete drying.

For GC-MS analysis, metabolites were subjected to methoxyamination and

trimethylsilylation. Dried polar phase were shaken for 1.5 hr at 30°C in 40 µl of MeOX (40

mg ml^{-1} methoxyaminhydrochloride in pyridine) followed by 30 min shaking at 37°C in 80 µl

BSTFA mixture (70 µl N,O-bis(trimethylsilyl)trifluoroacetamide + 10 µl alkane mix). The

derivatized metabolites were subjected to GC-MS analysis (Agilent technologies 7890, USA)

coupled with mass spectrometry. Measurement from an injection volume of 1 µl was taken in

split-less mode in DB-5 column (30 m × 0.25 mm, 0.25 µm film thickness, Agilent) using

helium as carrier gas. Metabolites were separated as described by Wagner et al. (2013).
Qualitative analysis of chromatograms was performed in MassHunter Qualitative analysis
Sp1 workstation (Agilent, USA). Identification and annotation of each compound was
supervised manually using AMDIS software and NIST08 database
(http://www.nist.gov/srd/mslist.html). Data were normalized to sample weight and internal
control (sorbitol). Statistical analysis was performed as described earlier (Quanbeck et al.,
2012). Log2 ratio of metabolite abundances in –Fe was plotted against +Fe. Delta method
approximation was used to calculate standard errors (se) of log-ratio, se log-ratio = 1/ln
$2\sqrt{[(SE_T/T)^2 + (SE_C/C)^2]}$, where SE_T and SE_C are standard errors of average –Fe and +Fe
metabolite abundances.

Measurement of glutathione-S-transferase (GST) activity and iron mobilization assay for PS
estimation

Activity measurement of GST was performed in the wheat roots subjected to –Fe for 10, 15
and 20 days of treatment along with the control plants (no stress) as mentioned in plant
materials. The estimation was done using Glutathione-S-transferases assay kit (Sigma, USA).
Briefly, 100 mg of tissue was used for total protein extraction. Equal amount of total protein
(25 μg) was used as a source of enzyme and 1-Chloro-2,4-dinitrobenzene (DNB) was used as
a substrate. The resulting GS-DNB conjugate was measured at 340 nm wavelength during the
time course of the reaction. The direct increase in absorption was measured and GST activity
was calculated as described in the manufacturer instruction kit.

For PS release, iron mobilization assay was performed as described earlier (Takagi et
al., 1976). Briefly, 10 seedlings (each for –Fe and +Fe for 10 days treatment) were used for
the experiment. After treatment the seedlings were subjected to PS release after 2 hours of
onset of light period in aerobic condition for 3 hrs in 20 ml deionised water with 10 mgl^{-1}
Micropur (Katadyn, Switzerland). Next, 2 ml of freshly precipitated solution of Fe(OH)$_3$ and
0.5 ml of 0.5 M Na acetate buffer (pH 5.6) was added to 8 ml of the collection solution. The
solution was shaken for 2 hrs and subsequently was filtered (Whatman #1) into 0.2 ml of
6NHCl. Ferric iron was reduced by addition of 0.5 ml of 8% hydroxylamine-hydrochloride
and heating to 60 °C for 20 min. The total concentration of ferrous iron was calculated by
measuring absorbance at 562 nm after adding 0.2 ml of 0.25%ferrozine and 1 ml of 2 M Na-
acetate buffer (pH 4.7).

Elemental analysis using inductive coupled plasma-mass spectroscopy and nitrate estimation
Elemental analysis in roots and shoots was performed using Inductive Coupled Plasma-MS
(ICP-MS). Metal analysis was performed as described earlier (Bhati et al., 2016; Aggarwal,
2018). Briefly, the samples were ground to fine powder and subsequently subjected to the
microwave-digested with HNO_3 as described earlier (SuraPure[TM], Merck). Respective metal
standards were also prepared for analysis. Three independent replicates from the experiments
were used for metal analysis.

Nitrate content in wheat roots was measured according to method described
previously (Cataldo et al., 1975). Briefly, 1 g of fresh tissue was homogenized in 6 ml of
deionised water and centrifuged at 30,000g for 15 min. The 100 µl of supernatant was added
to 400 µl of salicylic acid (w/v dissolved in conc. H_2SO_4). After mixing the reaction was kept
at room temperature for 20 min. 2N NaOH (9.5ml) was then added slowly to raise the pH
above 12. The samples were allowed to cool and readings were taken at 410nm in
spectrophotometer.

Results

Fe starvation affects wheat growth capacity and nutrient uptake
Fe starvation is known to affect plant growth capacity. In order to determine the effect of Fe
starvation, one-week old wheat seedlings, grown on complete medium (presence of Fe) were
transferred to Fe starvation media for additional days. After 20 days of starvation (DAS) the
wheat seedlings started showing strong phenotypic symptoms including visible chlorosis and
therefore detailed study was performed for this specific time point. In response to Fe
starvation, plants showed decrease in the shoot biomass with an enhanced chlorosis
phenotype and shortening of the root system compared to control wheat seedlings (Fig. 1A
and B). Roots of Fe-starved wheat seedlings showed decrease in number of lateral roots and
significant reduction in the primary root length in comparison to control plants (Fig. 1C and
D). Earlier studies have suggested that changes in the root system and Fe supply not only
affect the Fe accumulation capacity but also impacts the uptake of other nutrients such as Zn,
Cd etc. (Sperotto et al., 2012; Shukla et al., 2017). Therefore, effect of Fe starvation stress on
the uptake of Zn, Mn, Cu, and Mg in wheat undergoing Fe stress was studied (Table 1). Out
data indicated increased uptake of nutrient elements such as Zn, Mn, Cu, and Mg in roots but
accumulation in shoots was either unaltered or decreased. This further support the importance

to study root response to Fe starvation to gain insight on the molecular mechanism evolved

by wheat to cope with nutritional stress.

Differential expression analysis, homoeolog (A, B and D) induction and expression bias

during –Fe response

The effect of Fe starvation on the wheat root transcriptome has not been investigated till date.

To perform this study, RNAseq technology was used to identify the changes in the transcripts

of wheat roots, where plants were grown in presence or absence of optimal Fe (20 DAS). Our

analysis resulted in 87 million quality filtered reads, with an average of nearly 22 million

reads from each sample (more than 87% reads had a quality score greater than Q30). Filtered

reads from the four libraries had a mapping rate ranging from 81.7% to 85.4% when mapped

against release-37 of the wheat genome using TopHat (Supplementary Table S2). As quality

check, a strong correlation within the two biological replicates from each condition was

observed, while a clear variation was seen between the two conditions (Fig. 2A). We

thereafter analysed the expression values as FPKMs (Fragment Per Kilobase of transcript per

Million mapped reads), calculated by using Cufflinks software. Differentially expressed

genes (DEGs) were then identified by calculating logFC (log2 fold change) and performing

statistical tests between FPKM values from control and stressed samples using CuffDiff.

50610 genes with an FPKM of greater than or equal to 1 in at least one of the two conditions

were considered to be "expressed transcripts". In all, 7221 genes had logFC > 0 and 8010 had

logFC< 0 (Fig. 2B). On setting up a criterion of logFC of more than 1 for up-regulated genes,

and that of less than -1 for down-regulated genes and an FDR < 0.05, a total of 3478 genes

were highly expressed, whereas, 2376 were down- regulated under –Fe condition in wheat

roots (Fig. 2C). Interestingly, 45 genes were also induced exclusively during starvation

condition when compared to the control samples (Fig. 2C, Supplementary Table S3).

Our data allowed us to analyse the chromosomal distribution of the DEGs under the –

Fe condition. While all chromosomes contributed the DEGs, the highest number of genes was

mapped on chromosome 2 of the wheat genome (Fig. 3A). Equal representation of transcripts

was observed for the chromosome 7 and 5. The remaining chromosomes, 1, 3, 4 and 6

showed 15, 13, 12 and 10% distribution of DEGs, respectively. In polyploidy crops like

wheat, homoeolog induction bias could impact plant response to various stresses (Liu et al.,

2015; Powell et al., 2017). To determine the extent of induction bias (from A, B and D sub-

genomes) in wheat during Fe starvation, homoeolog specific expression analysis was

performed. Starting from a list of 8473 gene triplets present/expressed selected as 'accepted

triplets' (Supplementary Table S4), most of homoeolog triplets (8349) showed no significant

biasness in expression (A = B = D). Out of these, 8321 triplets appeared to be unaffected by –

Fe stress, while 22 and 6 homoeologous triplets were up- and down-regulated respectively.

Homoeolog expression bias was observed in 124 homoeolog triplets. Eighty-seven triplets

had only one of the homoeologs differentially expressed (up- or down-regulated). Sub-

genome specific contribution towards this biasness within triplets is depicted in left panel of

Fig. 3B. These include forty-seven in the category '1UP' with A > B = D, B > A = D and D >

A = B, 40 in category named '1DOWN' having A < B = D, B < A = D or D < A = B. Table 2

provides the list of genes with significant induction predominance occurring from the A and

B genomes in response to Fe starvation . Few of the prominent transcripts exclusively

induced by these two genomes include transcripts related to MYB TFs, metal transporters,

zinc transporters, RINGH-H2 type proteins, genes belonging to major facilitator superfamily

proteins etc. Additionally, 37 triplets had two of the homoeologs differentially expressed

while the third showed normal expression even under Fe stress (Fig. 3B; right panel). '2UP'

category includes AB > D, AD > B and BD > A while '2DOWN' includes AB < D, AD < B

and BD < A (Fig. 3B). This suggests that during Fe starvation, the additive homoeolog

contribution from either A or B sub-genome was the highest.

For polyploid genomes such as wheat, the interaction of its sub-genomes is known to

affect the final phenotype or contribute towards trait development (Borrill et al., 2015). To

check the homoeolog/sub-genome expression biasness under –Fe condition, effect on the

expression of transcripts from the genomes was performed by comparing the relative

normalised expression for each homoeolog within a triad. This resulted seven combinations

including one balanced category and six homoeolog specific dominance or suppression

categories. Our analysis reflected that most of the triads were falling under the balanced

category (Table 3 and Fig.3C). This category was represented by 77% and 77.89% of the

total triads for control and Fe starvation condition. Triads with unbalanced expression varied

in the range of 0.94 to 7.09 for all the 6 sub-categories across the two conditions (Table 3 and

Fig.3C). Our analysis revealed that maximum genome specific expression biasness was

observed for A and B on both the conditions. Interestingly, D genome was least suppressed

with a representation of 5.15% of the total triads taken in consideration in control when

compared to 4.93% in Fe starvation. 89% triads showed conserved balanced/unbalanced

contribution across both conditions. Overall, a higher relative abundance of D genome

(Control: 33.94%; –Fe: 33.95%) was noted as compared to the A (Control 33.06%; –Fe:

33.00%) and B (Control: 33.04%; –Fe: 33.95%) genomes.

Identification of DEGs pinpoints the prolific expression of genes involved in Strategy-II mode

of Fe uptake

To identify the transcripts differentially expressed in roots in response to Fe starvation, top 50

genes either up-regulated or down-regulated were shortlisted (Fig. 4). Expression of most of

the highly up-regulated transcripts ranged from 12 to 4.8 log2 fold change (Supplementary

Table S3) indicating their higher fold accumulation under Fe starvation compared to the

control conditions. As observed in the top 50 upregulated genes, the predominance of

pathway genes encoding the components for Strategy-II mode Fe uptake was observed across

all the 3478 upregulated genes. Categorically, these highly induced genes are known to be

involved during Strategy-II mode of iron uptake, including the sub-family of nicotinamine

synthase (NAS) and deoxymugineic acids (DMA) biosynthesis genes (Fig. 4, left panel).

Other important genes include genes belonging to the Major Facilitator Superfamily

including an ABC transporter, zinc-induced facilitator like transporters (ZIFL), sulphate

transporters. Therefore, under Fe starvation, the DMAS biosynthesis genes were highly

induced. Similarly, YSL genes were significantly induced under Fe starvation condition. All

the homoeologs of *TaYSL9* and *TaYSL1A* showed especially high expression under starvation

condition (Supplementary Table S5). On the similar lines, genes encoding for NRAMP also

showed high transcript abundance in roots subjected to Fe starvation (Supplementary Table

S5). The validation of the expression of few Strategy-II uptake genes was also done by

quantitative real time-PCR (qRT-PCR). Our results for qRT-PCR validate our inference from

the RNAseq analysis. Almost all of the Strategy-II genes tested for their expression showed a

very high fold expression in Fe starved roots as compared to the control (Supplementary Fig.

S1). Highest expression was obtained for ZIFL4, DMAS1, NAAT1, NAS1 those are the

prime components for Strategy-II mediated uptake. Other transcripts encoding for thaumatin

like proteins etc. were also induced under Fe limiting condition.

Transcripts of genes involved in the Strategy-I mode of Fe uptake were also analysed

(Supplementary Table S6). An important component of Strategy-I pathway that includes H^+-

ATPase (AHA) subfamily genes was not differentially expressed under Fe limiting condition

(Supplementary Table S6). This was in agreement with the qRT-PCR analysis where, either

no change or down-regulation of AHA genes was observed (Supplementary Fig. S1) in wheat

roots. Metallo-reductases are important components of Strategy-I represented by ferric-

chelate reductase (FRO). Most of the transcripts encoding for wheat FROs do not show

significant changes in starved roots as compared to control except for one transcript.

Interestingly, iron regulated transporters (IRT;

TRIAE_CS42_7DS_TGACv1_622068_AA2032200;

TRIAE_CS42_4AL_TGACv1_289466_AA0971640) were significantly expressed in Fe

starved wheat roots, suggesting their involvement in wheat under metal stress. The high

expression of *TaIRT1* was also confirmed by qRT-PCR (Supplementary Fig. S1). This suggest

that IRT function might be conserved among the plant species.

In addition to this, forty-five genes showed exclusive transcript abundance under Fe

starvation (Supplementary Table S3). No transcript for any of these genes was detected in the

control root samples, suggesting their high specificity for Fe response. Some of the

transcripts responding to Fe starvation encode for metallothionine, metal transporters,

vacuolar iron transporters (VIT1) and nicotianamine synthase 2 (NAS2). This suggests that in

wheat few components involved in Fe uptake could actually respond exclusive to Fe

starvation. The fold expression levels of genes those were highly down-regulated ranged

from -7.1 to -2.85-fold at the level of log2 scale (Supplementary Table S3). Interestingly,

multiple genes encoding for nitrate transporters were highly down-regulated under Fe

limiting condition (Fig. 4, right panel). Two genes encoding for cytochrome-P450 also

showed down-regulation. Among the others, genes encoding for Histone deacetylase,

peptidases A1 containing domain protein, cinnamyl alcohol dehydrogenase, dirigent protein

show significant down-regulation.

Functional enrichment network of Fe starvation related genes

Gene ontology annotations and classification of DEGs was performed to get the overview of

processes those are representation of cellular, molecular and biological functions. Analysis

was further extended to cluster analysis using Cytoscape plugins, BINGO and Enrichment

Map (Maere et al*.,* 2005; Merico et al*.,* 2010). A total of 5854 DEGs were found to be

differentially-expressed in response to Fe starvation; significant GO categories were assigned

to all DEGs. The DEGs annotated for GO terms were visualized using WEGO tool (Fig. 5A,

Supplementary Table S7). Most enriched GO terms in the "cellular component" category

were membrane and intracellular organelle. Other overrepresented terms included catalytic

activity and ion as well as organic compound binding for the "molecular function" category,

while metabolic processes related to nitrogen as well as other cellular processes were

observed for "biological process" category (Fig. 5A). Overall, catalytic activity and binding

activities were most significantly enriched GO terms in –Fe condition (Fig. 5B). Further

mapping of upregulated genes to databases such as, Kyoto encyclopedia of genes and
genomes (KEGG) pathway (Xie et al., 2011) and MapMan (Thimm et al., 2004) revealed
enrichment related to phenyl-propanoid biosynthesis, amino acid biosynthesis and carbon
metabolism, and glutathione metabolism (Fig. 5C and Supplementary Fig. S2). The role of
glutathione in response to Fe starvation is intriguing and deserves further investigations.

GO category "integral to membrane" was found to be associated with pathways
clustered into "metal ion and trans-membrane transport" and "photosynthesis" related GO
categories (Fig. 6A). Clustered pathways involved in "response to nutrient levels" and
"Sulphur amino acid metabolism" was also enriched (Fig. 6A). A few other up-regulated
clustered pathway genes involved fall into the category "response to phosphate starvation". In
the enriched network of down-regulated DEGs, genes involved in iron and other metal
homeostasis were associated with two different clusters, mainly included genes related to
lipid, ketone and carboxylic metabolism, in addition to those involved in nitric acid and
salicylic acid response (Fig. 6B). A second cluster contains transcripts related to purine and
adenine nucleotide binding and pyrophosphatase activity. Thus, whole set of significant up-
regulated and down-regulated DEGs were clustered in distinct cellular processes suggesting
differential transcriptional response under Fe starvation.

Identification of transcriptional regulatory genes during Fe starvation
To address an important question on how the –Fe signal is connected to the transcriptional
machinery in wheat, genes encoding TFs were identified. Earlier the TFs involved in the
response to Fe starvation were identified in the model plant Arabidopsis and rice including
POPEYE (PYE), basic helix-loop helix (bHLH), ethylene insensitive-3 (EIN3), EIN3-like1
(EIL1) (Long et al., 2010, Mai et al., 2016, Bauer et al., 2011, Ivanov et al., 2012; Ogo et al.,
2011; Bashir et al., 2014). Consistently, our transcriptome analysis showed that all
homoeologs for PYE encoding a bHLH transcription factor and BRUTUS-like (BTS) known
to encode a putative hemerythrin E3 ligase protein in Arabidopsis and rice were highly
expressed in wheat roots under Fe starvation (Supplementary Table S8). Analysis of these
category of genes led to identification of 41 significantly up-regulated TF family members
(>log2FC) (Supplementary Table S8). The TF family members belong to categories like
APETALA2 ethylene-responsive element binding proteins (AP2/EREBP), WRKY, C2H2,
Zinc finger proteins (including C3HC4), NAM, bHLH hemerythrin and U-box (Fig. 7A). Out
of these, transcripts encoding for AP2/EREBP followed by WRKY and bHLH were the most
abundant in wheat. Genes encoding for AP2-EREBP, WRKY and C2H2 type of TFs also

showed high predominance (Supplementary Table S8 and Supplementary Fig. S3). However,
genes encoding for TFs such as MADS, PHD and homeobox (HB) domain containing
proteins were highly represented in the list of down-regulated transcripts. Finally, to ascertain
regulatory functions under iron starvation, enrichment mapping was performed for the
significantly expressed DEGs those were classified into transcriptional factors. Interestingly,
genes involved in regulation of gene expression were clustered with primary and nucleotide
metabolism related genes in both positive and negative correlation. This functional cluster
was in positive correlation with TFs involved in gene ontology (GO) categories such as
"vacuolar transport" and "DNA topoisomerase III activity". However, two separate clusters
"cellular response to auxin stimulus and signalling" and "cell-fate specification" were in
negative correlation with "regulation of gene expression" and "primary and nucleotide
metabolism" (Fig. 7B). This suggested an important regulatory role of TFs involved in auxin
signalling and cell-fate specification in Fe starvation response control, thereby modulating
the network of Fe homeostasis.
519
Glutathione-S-transferases are involved in the response to Fe starvation in wheat
Glutathione mediated conjugation of multiple metabolites plays important role during the
metal stress (Zhang et al., 2013). Our transcriptome data revealed the enhanced expression of
multiple glutathione-S-transferases (GST) in Fe-starved wheat root as compared to control
(Supplementary Table S9). To correlate the expression response with its enzymatic activity,
temporal response of GST was measured in wheat roots under –Fe conditions. Using the GST
functional assay, the activity was determined in wheat roots of plants grown for 10, 15 and 20
days under Fe starvation. Our activity assays showed significant increase in the GST activity
under –Fe condition, which peaked at 10 and 20 days after the beginning of Fe starvation
compared to control plants (Fig. 8A). Therefore, our experiments validate the increased GST
transcript abundance with enhanced glutathione activity. These results indicate an important
role of glutathione in response to Fe starvation in wheat roots.
532
Fe starvation causes an accumulation of organic acids and polyhydroxy acids in wheat
To obtain a comparative insight of the metabolite profile of wheat roots of plants grown in
absence or presence or Fe; GC-MS profiling analysis was performed. Metabolites were
extracted from the roots of three replicate pools of plants each containing eight seedlings.
Qualitative processing of each chromatogram for peak area and identification was performed
in MassHunter version B.05.00 software coupled with NIST11 compound library. The

compound annotation was determined by comparing individual resolved peaks to library
searches based on mass spectra and compounds chromatographic retention indices.
Interestingly, analysis resulted in the identification of 54 metabolites and further 39 annotated
metabolites were analysed for their response ratio (Supplementary Table S10). To compare
the change under Fe-starvation, metabolite abundances (–Fe roots/control roots) were
calculated and expressed in Log2 fold change values. Fe-starvation significantly affected
accumulation of 22 metabolites that includes organic acids, polyhydroxy acids, amino acids
and some of the sugars, fatty acids and phosphates (Fig. 8B). Amongst treatment-specific
changes, few organic acids such as fumaric acid, acetic acid and malonic acid showed higher
level of accumulation in Fe starved roots as compared to the control. In contrast, citric acid,
malic acid, valeric acid and aconitic acid were significantly lowered in Fe starved roots when
compared to control samples. In Fe-starved roots, the accumulation of amino acids mainly, L-
Valine showed significant increase while hydroxyl butyric acid and pyroglutamic acid was
lowered in comparison with control roots. Polyhydroxy acids like gluconic acid and glyceric
acid were also significantly high in Fe starved roots, whereas level of hexonic acid and
arabinoic acid was found to be low. Taken together, our results showed that during Fe
starvation wheat roots undergo reprograming for metabolic changes to maintain the Fe-flux.

Discussion

How plants maintain nutrient homeostasis is a fascinating question in plant biology. In this
direction, *A. thaliana*, with its fully sequenced small genome has provided some basic
information. Developing our knowledge on nutrient homeostasis in crops, especially those
having complex genomes, including hexaploid wheat is more challenging. Recently, the
release of genome sequence, generation of transcriptome data and the advancement of
different functional tools have provided much needed impetus in this direction (Borrill et al.,
2015). The current study was undertaken to gain insight into the response of hexaploid wheat
exposed to Fe starvation that is known to severely affect crop production. In this study,
transcriptome of wheat (cultivar C306) in response to Fe starvation was generated. Our
analysis revealed that; (a) wheat utilizes primarily Strategy-II mode of Fe uptake, (b)
accumulates transcripts encoding for methionine-salvage pathway coupled with enhanced
GST activity and (c) accumulates specific metabolites including fumarate, acetate, malonate
and xylofuranose, to efficiently mobilize soil Fe. Interestingly, our systematic analysis of
expression data revealed that transcripts show an induction biasness for A and B genomes of

wheat in response to –Fe condition. Overall, this work provides the first comprehensive
insight at the molecular level for wheat roots during Fe-deficiency.

Wheat plants subjected to Fe starvation show physiological defects such as a decrease
in the root growth. This phenotype was consistent with the previous report showing negative
impact on the root growth of wheat seedlings under Fe stress (Garnica et al., 2018). Our
analysis also suggested down-regulation of few nitrate transporters for the time point studied.
Homology searches indicated the identified nitrate transporters in wheat as
TaNRT2.2 and *TaNRT2.3* (Buchner and Hawkesford, 2014). These observations suggest that
Fe starvation represses nitrate transporters, which in turn leads to lower accumulation of
nitrate levels in the roots (Supplementary Fig. S4). Although impact of Fe starvation on the
nitrate metabolism has not been studied in detail, yet similar decrease in nitrate levels were
also observed in cucumber shoots subjected to –Fe condition (Borlotti et al., 2012). This
observation further reinforces the existence of an interaction between macro and
micronutrients in plants, which has recently gained attention (Rouached and Rhee, 2017;
Bouain et al., 2019).

Due to the low availability of micronutrients in soil, plants are equipped in recruiting
components that could participate in the two major strategies for Fe uptake (Kobayashi and
Nishizawa, 2012). Cereals such as maize and rice predominantly utilize Strategy-II mode of
uptake, unlike Arabidopsis, that employs Strategy-I mode of Fe uptake (Li et al., 2014; Zanin
et al., 2017; Hell and Stephan, 2003). Nevertheless, a comprehensive study to identify the
molecular players involved during Fe limiting conditions in wheat is lacking. Our RNAseq
based analysis, strongly suggested an increase in the transcript abundance of genes for
Strategy-II mode of Fe uptake. With the exception of conserved genes like *IRT1* (iron
regulated transporter), prime genes for Strategy-I uptake mechanism like *FRO* (Ferric chelate
reductase) and proton *H-ATPase* (AHA-like) were also present during starved condition but
at very low abundance and no FRO activity has been reported for cereals. During this study,
wheat root subjected to Fe starvation also does not show any FRO activity (data not shown).
Therefore, our results reveal that wheat utilize Strategy-II mediated uptake of Fe and induces
IRT genes that might be conserved for their function across the plant kingdom.

The series of events during Strategy-II uptake mechanism used by Gramineae
involves secretion of PS that facilitate conjugation of Fe to form Fe-complexes. These Fe-PS
complexes are imported in roots for remobilization to the different developing organs. As
represented in Figure 9, the transport of these Fe-complexes occurs via membrane
transporters via ZIFL genes encoding for TOM proteins (Nozoye et al., 2011; Nozoye et al.,

2015). Multiple wheat ZIFL show high transcript accumulation and are closest homolog of

OsTOM1 from rice, thereby speculating its role in Fe acquisition. Utilizing gain- and loss of

function approaches OsTOM1 has been demonstrated to be a DMA effluxer for its role in

enhancing mobilization and thereby improving Fe uptake (Nozoye et al., 2011). Altogether,

based upon the high similarity and response under Fe starvation, it is tempting to propose

wheat ZIFL (ZIFL4) as a functional transporter of DMA. Although, other *ZIFLs* were also

significantly expressed but functional wheat transporter for DMA needs to be deciphered. PS

secretion was also observed during our analysis for the roots of C306 exposed to Fe

starvation. In this study a high PS release of ~40-45 $nmolg^{-1}$ root biomass $(3\ hrs)^{-1}$ was

observed, reinforcing our RNAseq data that DMA biosynthesis genes and its transporters

were highly upregulated. Control plants show accumulation of ~1-2 $nmolg^{-1}$ suggesting that

wheat releases a basal level of PS to maintain the constant flux of Fe in roots. Upon careful

analysis of the RNAseq data, it was observed that multiple such efflux transporters are

represented in the list that are highly abundant under Fe starvation. During our study, we

identified *TaDMAS1* encoding for deoxymugineic acid synthase that was highly responsive

for Fe starvation (Supplementary TableS5). Earlier, *TaDMAS1* was reported as a gene that

was broadly expressed across the tissue and was regulated during Fe-starved condition. These

observations support the notion that TaDMAS1 has potential to enhance the seed iron storage

capacity (Beasley et al., 2017). During our study, GO term enrichment was also observed for

DMA biosynthesis pathway, transmembrane transporters and cellular response that reinforce

importance of these functional categories. Surprisingly, genes pertaining to photosynthesis

were also observed. This anomaly could be explained by the presence of active photosystem

in the basal roots of our collected wheat tissue, similar to that observed in Arabidopsis due to

the changes in the auxin/cytokinin ratio (Kobayashi et al., 2012). Secondly, during this study

the wheat roots under Fe starvation show significant accumulation of two Golden-2 like

(GLK) TF's (~logFC 1.54 and 1.55; Supplementary Table S3). GLK's are known to improve

phototrophic performance of roots and thereby enhancing root photosynthesis (Kobayashi et

al., 2013). These reasons may account for the enhanced accumulation of transcripts related to

photosynthesis along with other metabolic related genes.

Wheat is a hexaploid crop with three genomes and therefore studying the homoeolog

induction and expression bias could provide an insight on the regulation of the transcript

under a given biotic and abiotic stress. Therefore, we subjected our RNAseq data to these

analyses. In depth analysis confirms the minimal suppression for the expression bias for the

D-genome derived transcripts as compared to A or B. These observations support the

previous analysis wherein, suppression of D genome was significantly less frequent in

multiple tissue (Ramirez-Gonzalez et al., 2018). Earlier, homoeolog expression was also

studied for wheat under infection with *Fusarium pseudograminearum*, wherein the

expression bias was observed for B and D subgenomes (Powell, et al., 2017). Utilizing

RNAseq data, changes in the pattern of the homoeologous gene expression were also

reported in bread wheat (Leach et al., 2014). Therefore, such studies are important to pin-

point actively expressed/induced homoeologs that could be targets for genetic improvement.

However, the genome biasness and its association with the secretion of siderophores,

genotypic variability in a complex wheat genome has yet to be investigated. Previous studies

have shown that the phytosiderophores release mechanism in wheat is effective when the

three genomes A, B and D contribute synergistically to induce its biosynthesis and release. In

general, genome induction biasness for A and B genome was observed for the hexaploid-

wheat under Fe stress (Fig. 3B). Interestingly, these genomes were also suggested to be

important during starvation of other micronutrients like Zn (Tolay et. al., 2001). These

observations provide clue for the importance of A and B genomes during micronutrient

starvation in hexaploid wheat when compared to their progenitor. The PS release intensity is

also dependent on complementary action of the genotypic ploidy with the following the

pattern, *T. aestivum* (BBAADD) > *T. dicoccum* (BBAA) > *T. monococcum* (AA) > *Ae.*

tauschii (DD) (Ma et al., 1999; Tolay et al., 2001).

Multiple genes for the salvage pathway were highly up-regulated including *NAS1*,

NAS2 and *NAAT1* (Supplementary Table S3 and Supplementary Table S5). The contribution

of these transcripts arises from different genomes suggesting that all the genomes of wheat

are capable of responding to the starvation condition. Based on our RNAseq analysis it is

likely that the gene encoding for *TaYS1A* and *TaYSL9* could be the putative transporter for

Fe-siderophore complex. Previous, studies have indicated the high response of *TaYS1A* under

Fe starvation in roots and shoots (Kumar et al., 2019). Out of the closest orthologs of wheat-

YS1A, HvYS1A and *ZmYS1A*, *HvYS1A* has been shown to be involved in specific transport of

Fe whereas, ZmYS1 was reported to have broad substrate specificity (Schaaf et al., 2004).

Our RNAseq data also revealed a conserved function of genes from monocots and dicots

during the Fe regulation and known important Fe regulators such as PYE and BRUTUS also

showed differential expression in wheat roots (Ivanov et al., 2012). Same for TFs spanning

multiple gene families such as MYB (Rubio et al., 2001), *bHLH* (Colangelo et al., 2004;

Jakoby et al., 2004; Ogo et al., 2007), *C2H2, NAC, AP2-ERB* (Kim et al., 2012), and *WRKY*

(Devaiah et al., 2007) were characterized in Arabidopsis and rice for their involvement in

nutrient uptake which were found to be highly represented in root of Fe-starved wheat. Similarly, hemerythrin motif-containing protein were also overexpressed in our study. As in case of other graminaceous plants like rice, hemerythrin domain containing proteins including OsHRZ homolog for Arabidopsis BRUTUS was also differentially perturbed in – Fe (Kobayashi et al., 2014). Overall, based on the information obtained from our studies future research focus should be able to gain better insights of the molecular functions of these genes using tilling population in wheat and/or a heterologous system.

Changes in metabolite content in response to Fe-deficiency can lead the plant to undergo adaptive processes to maintain iron homeostasis (Schmidt et al., 2014). A schematic representation of data provides a comprehensive summary of the metabolic and transcriptional changes that occur during Fe starvation in wheat roots (Fig. 9). The mobilization and transport of Fe inside the plant tissue depends on its speciation and complexation form. Depending on the tissue, Fe could form a complex with multiple metabolites such as nicotianamine, malate, citrate that facilitates the transport through the phloem and xylem (Grillet et al., 2014; Palmer et al., 2013). Our analysis for the metabolome of roots suggested predominance of additional organic compounds such as fumarate, acetate, glycerate etc. thereby, suggesting that they could be also involved in Fe-complexation to assist in long distance transport and remobilization to different tissues (Fig. 9). These metabolites are reported for their ability to complex with Fe(III) state during the transport in the xylem vessels as detected during the X-ray absorption near-edge structure spectroscopy based studies (Terzano, et al., 2013). Similarly, the accumulation of gylcerate could be responsible for added thrust towards glycerate-serine pathway during Fe starvation (Figure 9). Glycerate-serine pathway are important to generate methionine and precursor (S-adenosyl methionine) of DMA biosynthesis. Consistent with the metabolome data, the transcripts encoding these proteins were highly upregulated in our study as presented by the bins (Figure 9). Members of ABC-B transporters were also reported to participate in oxidative stress that was linked with metal stress including Fe starvation (Kispal et al., 1997; Schaedler et al., 2014). During our analysis we have identified a TAP type, ABCB subfamily transporter referred to as *TaABCB25* that was highly upregulated during Fe starvation (Fig. 4). The closest ortholog for wheat ABCB25 transporters were identified in rice (OsABCB25) and Arabidopsis (AtABCB27) (Supplementary Fig. S5A). The temporal-induction of *TaABCC25* was also re-confirmed in roots subjected to Fe starvation (Supplementary Fig. S5B). *Arabidopsis thaliana* ABCB27 was shown to have the ability to transport glutathione conjugate metabolite precursors for the Fe-S cluster assembly (Schaedler et al., 2014). Based

on the presence of domain structure these ABCB transporters from rice, Arabidopsis and
wheat qualify as half-size transporter (Supplementary Fig. S5C). In graminaceous species,
glutathione related activity was observed to be highly upregulated under Fe deficient
condition supporting our observation in the current study (Bashir et al., 2007). Enhanced
GST activity in wheat under Fe starvation and the abundance of *TaABCB25* in roots, warrants
further study of this gene for its role in mobilizing micronutrient uptake. Glutathione and
GST plays an essential role during Fe starvation related responses in *Arabidopsis* and
also quench reactive molecules to protect the cell from oxidative damage and prevent
chlorophyll loss (Ramirez et al., 2013; Shanmugam et al., 2015; Kumar and Trivedi, 2018).
Our analysis also suggests that high transcript and enzyme activity of GST could be linked
with the primary metabolism since it accumulates fumarate (Fig. 9). In *Arabidopsis*, it has
been shown that GST catalyses Glutathione-dependent isomerization of maleyl-acetoacetate
into fumaryl-acetoacetate eventually leading to accumulation of fumarate and acetoacetate
(Dixon et al., 2002). Additionally, GST is also linked with the conjugation of the reduced
glutathione (GSH) forms to various substrate to make it water soluble. The sulphur derived
metabolite and glutathione are well known to be involved in biosynthesis methionine, a well-
known precursor for DMA biosynthesis (Forieri et al., 2013). Altogether, this supports the
notion that in wheat role of glutathione is important during Fe starvation and ABC
transporters might play a significant role in mobilization of metabolite/organic acid or
specific siderophores.

In conclusion, in this study, the core components for Fe starvation response in
hexaploid wheat have been identified so as to provide a better understanding of the molecular
events that participate during Fe starvation in wheat roots. In particular, we provide line of
evidence for the role of GST in the response to –Fe in roots. Complementary approaches,
analytical and transcriptome analysis reinforce the importance of primary metabolism for
reprogramming organic acids and amino acids thereby leading to the Fe homeostasis in
wheat. The information here not only help to design strategies to improve plant response to
Fe starvation in wheat, but will also foster Fe uptake and accumulation studies, which are
required to boost productivity and grain nutritional quality through Fe biofortification
programs.

Acknowledgement
Authors thank Executive Director, NABI for facilities and support. This research was funded
by the NABI-CORE grant to AKP and partial support from DST-SERB grant

(PDF/2016/001355) to PG. GK and AK acknowledge NABI-SRF Fellowships. Technical

help from Jagdeep Singh for the GC-MS analysis is highly appreciated. DBT-eLibrary

Consortium (DeLCON) is acknowledged for providing timely support and access to e-

resources for this work. The wheat genome resources developed by International Wheat

Genome Sequencing Consortium are highly appreciated. We also acknowledge the critical

comments given by Dr. Kerry Pedley, USDA, USA. We would like to thank Abhijeet Panwar

(CDAC-Pune) for his help in the preparation of the Sankey plot.

Reference

Aggarwal S, Kumar A, Bhati KK, Kaur G, Shukla V, Tiwari S, Pandey AK. 2018. RNAi-mediated downregulation of inositol pentakisphosphate kinase (IPK1) in wheat grains decreases phytic acid levels and increases Fe and Zn accumulation. Frontiers in Plant Science **9,** 259.

Bashir K, Hanada K, Shimizu M, Seki M, Nakanishi H and Nishizawa NK. 2014. Transcriptomic analysis of rice in response to iron deficiency and excess. Rice **7**, 18.

Bashir K, Nagasaka S, Itai RN, Kobayashi T, Takahashi M, Nakanishi H, Mori S, Nishizawa NK. 2007. Expression and enzyme activity of glutathione reductase is upregulated by Fe-deficiency in graminaceous plants. Plant Molecular Biology **65**, 277–84.

Bauer P, Blondet E. 2011. Transcriptome analysis of *ein3 eil1* mutants in response to iron deficiency. Plant Signaling & Behavior **6**, 1669–1671.

Beasley JT, Bonneau JP, Johnson AAT. 2017. Characterisation of the nicotianamine aminotransferase and deoxymugineic acid synthase genes essential to Strategy II iron uptake in bread wheat (*Triticum aestivum* L.) PLoS ONE **12**, e0177061.

Bhati KK, Alok A, Kumar A, Kaur J, Tiwari S, Pandey AK. 2016. Silencing of *ABCC13* transporter in wheat reveals its involvement in grain development, phytic acid accumulation and lateral root formation. Journal of Experimental Botany **67**, 4379–4389.

Bocchini M, Bartucca ML, Ciancaleoni S, Mimmo T, Cesco S, Pii Y, Albertini E, Del Buono D. 2015. Iron deficiency in barley plants: phytosiderophore release, iron translocation, and DNA methylation. Frontiers in Plant Science **6**, 514.

Borlotti A, Vigani G, Zocchi G. 2012. Iron deficiency affects nitrogen metabolism in cucumber (*Cucumis sativus* L.) plants. BMC Plant Biology **12**, 189.

Borrill P, Adamski N and Uauy C. 2015. Genomics as the key to unlocking the polyploid potential of wheat. New Phytologist **208**, 1008-1022.

Borrill P, Harrington SA, Uauy C. 2018. Applying the latest advances in genomics and phenomics for trait discovery in polyploid wheat. The Plant Journal **97**, 56-72.

Bouain N, Krouk G, Lacome B, Rouached H. 2019. Getting to the root of plant mineral nutrition: combinatorial nutrient stresses reveal emergent properties. Trends in Plant Science **24**, 542-552

Buchner P, Hawkesford MJ. 2014. Complex phylogeny and gene expression patterns of members of the NITRATE TRANSPORTER 1/PEPTIDE TRANSPORTER family (NPF) in wheat. Journal of Experimental Botany **65**, 5697–710.

Buckhout TJ, Yang TJ, Schmidt W. 2009. Early iron-deficiency-induced transcriptional changes in Arabidopsis roots as revealed by microarray analyses. BMC Genomics **10**, 147.

Cataldo DA, Maroon M, Schrader LE, Youngs VL. 1975. Rapid colorimetric determination of nitrate in plant tissue by nitration of salicylic acid. Communications in Soil Science and Plant Analysis **6**, 71–80.

Colangelo EP, Guerinot ML. 2004. The essential Basic Helix-Loop-Helix protein FIT1 is required for the iron deficiency response. The Plant Cell **16**, 3400–3412.

Connorton JM, Balk J, Rodríguez-Celma J. 2017. Iron homeostasis in plants – a brief overview. Metallomics **9**, 813–823.

Connorton JM, Jones ER, Rodríguez-Ramiro I, Fairweather-Tait S, Uauy C, Balk J. 2017. Wheat vacuolar iron transporter TaVIT2 transports Fe and Mn and is effective for biofortification. Plant Physiology **174**, 2434–2444.

Curie C, Panaviene Z, Loulergue C, Dellaporta SL, Briat JF, Walker EL. 2001. Maize yellow stripe1 encodes a membrane protein directly involved in Fe(III) uptake. Nature **409**, 346–349.

Devaiah BN, Karthikeyan AS, Raghothama KG. 2007. WRKY75 transcription factor is a modulator of phosphate acquisition and root development in Arabidopsis. Plant Physiology **143**, 1789–1801.

Dixon DP, Davis BG, Edwards R. 2002. Functional divergence in the glutathione transferase superfamily in plants. Journal of Biological Chemistry **277**, 30859–30869.

Forieri I, Wirtz M and Hell R. 2013. Toward new perspectives on the interaction of iron and sulfur metabolism in plants. Frontiers in Plant Science **4**, 357.

Garnica M, Bacaicoa E, Mora V, San Francisco S, Baigorri R, Zamarreño AM, Garcia-Mina JM. 2018. Shoot iron status and auxin are involved in iron deficiency-induced phytosiderophores release in wheat. BMC Plant Biology **18**, 105.

Grillet L, Ouerdane L, Flis P, Hoang MTT, Isaure MP, Lobinski R, Curie C and Mari S. 2014. Ascorbate efflux as a new strategy for iron reduction and transport in plants. Journal of Biological Chemistry **289**, 2515-2525.

Gross J, Stein RJ, Fett-Neto AG, Fett JP. 2003. Iron homeostasis related genes in rice. Genetics and Molecular Biology **26**, 477–497.

Hell R, Stephan UW. 2003. Iron uptake, trafficking and homeostasis in plants. Planta **216**, 541–51.

Hindt MN, Guerinot ML. 2012. Getting a sense for signals: regulation of the plant iron deficiency response. Biochimica et Biophysica Acta **1823**, 1521–30.

Inoue H, Kobayashi T, Nozoye T, Takahashi M, Kakei Y, Suzuki K, Nakazono M, Nakanishi H, Mori S, Nishizawa NK. 2009. Rice OsYSL15 is an iron-regulated iron (III)-deoxymugineic acid transporter expressed in the roots and is essential for iron uptake in early growth of the seedlings. The Journal of Biological Chemistry **284**, 3470–9.

Ishimaru Y, Suzuki M, Tsukamoto T, *et al.* 2006. Rice plants take up iron as an Fe^{3+}-phytosiderophore and as Fe^{2+}. The Plant Journal **45**, 335–346.

Ivanov R, Brumbarova T, Bauer P. 2012. Fitting into the harsh reality: regulation of iron-deficiency responses in dicotyledonous plants. Molecular Plant **5**, 27–42.

Jakoby M, Wang HY, Reidt W, Weisshaar B, Bauer P. 2004. *FRU* (*BHLH029*) is required for induction of iron mobilization genes in *Arabidopsis thaliana*. FEBS Letters **577**, 528–534.

Kabir AH, Paltridge NG, Roessner U, Stangoulis JCR. 2013. Mechanisms associated with Fe-deficiency tolerance and signaling in shoots of *Pisum sativum*. Physiologia Plantarum **147**, 381–395.

Kim J, Rees DC. 1992. Structural models for the metal centers in the nitrogenase molybdenum-iron protein. Science **257**, 1677–82.

Kim MJ, Ruzicka D, Shin R, Schachtman DP. 2012. The Arabidopsis AP2/ERF transcription factor RAP2.11 modulates plant response to low-potassium conditions. Molecular Plant **5**, 1042–1057.

Kim SA, Punshon T, Lanzirotti A, Li L, Alonso JM, Ecker JR, Kaplan J, Guerinot ML. 2006. Localization of iron in Arabidopsis seed requires the vacuolar membrane transporter VIT1. Science **314**, 1295–1298.

Kispal G, Csere P, Guiard B, Lill R. 1997. The ABC transporter Atm1p is required for mitochondrial iron homeostasis. FEBS Letters **418**, 346–50.

Kobayashi K, Baba S, Obayashi T, *et al.* 2012. Regulation of root greening by light and auxin/cytokinin signaling in Arabidopsis. The Plant Cell **24, 1081–1095.

Kobayashi K, Sasaki D, Noguchi K, *et al.* 2013. Photosynthesis of root chloroplasts developed in Arabidopsis lines overexpressing GOLDEN2-LIKE transcription factors. Plant and Cell Physiology **54, 1365–1377.

Kobayashi T, Itai RN, Ogo Y, Kakei Y, Nakanishi H, Takahashi M, Nishizawa NK. 2009. The rice transcription factor IDEF1 is essential for the early response to iron deficiency, and induces vegetative expression of late embryogenesis abundant genes. The Plant Journal **60**, 948–961.

Kobayashi T, Nakanishi Itai R, Nishizawa NK. 2014. Iron deficiency responses in rice roots. Rice **7**, 27.

Kobayashi T, Nishizawa NK. 2012. Iron uptake, translocation, and regulation in higher plants. Annual Review of Plant Biology **63**, 131–152.

Kumar A, Kaur G, Goel P, Bhati KK, Kaur M, Shukla V, Pandey AK. 2019. Genome-wide analysis of oligopeptide transporters and detailed characterization of yellow stripe transporter genes in hexaploid wheat. Functional & Integrative Genomics **19**, 75-90.

Kumar S, Trivedi PK. 2018. Glutathione S-Transferases: role in combating abiotic stresses including arsenic detoxification in plants. Frontiers in Plant Science **9**, 751.

Leach LJ, Belfield EJ, Jiang C, Brown C, Mithani A and Harberd NP. 2014. Patterns of homoeologous gene expression shown by RNA sequencing in hexaploid bread wheat. BMC Genomics **15**, 276.

Lee S, Chiecko JC, Kim SA, Walker EL, Lee Y, Guerinot ML, An G. 2009. Disruption of OsYSL15 leads to iron inefficiency in rice plants. Plant Physiology **150**, 786–800.

Li W, Lan P. 2017. The understanding of the plant iron deficiency responses in strategy I plants and the role of ethylene in this process by omic approaches. Frontiers in Plant Science **8**, 40.

Li X, Zhang H, Ai Q, Liang G, Yu D. 2016. Two bHLH transcription Factors, bHLH34 and bHLH104, regulate iron homeostasis in *Arabidopsis thaliana*. Plant Physiology **170**, 2478–2493.

Li Y, Wang N, Zhao F, Song X, Yin Z, Huang R, Zhang C. 2014. Changes in the transcriptomic profiles of maize roots in response to iron-deficiency stress. Plant Molecular Biology **85**, 349–363.

Lin XY, Ye YQ, Fan SK, Jin CW, Zheng SJ. 2016. Increased sucrose accumulation regulates iron-deficiency responses by promoting auxin signaling in arabidopsis plants. Plant Physiology **170**, 907–920.

Liu Z, Xin M, Qin J, Peng H, Ni Z, Yao Y, Sun Q. 2015. Temporal transcriptome profiling reveals expression partitioning of homeologous genes contributing to heat and drought acclimation in wheat (*Triticum aestivum* L.). BMC Plant Biology **15**, 152.

Lohse M, Nagel A, Herter T, May P, Schroda M, Zrenner R, Tohge T, Fernie AR, Stitt M, Usadel B. 2014. Mercator: a fast and simple web server for genome scale functional annotation of plant sequence data. Plant, Cell & Environment **37**, 1250–1258.

Long TA, Tsukagoshi H, Busch W, Lahner B, Salt DE, Benfey PN. 2010. The bHLH transcription factor POPEYE regulates response to iron deficiency in Arabidopsis roots. The Plant Cell **22**, 2219–2236.

Ma JF, Taketa S, Chang YC, Takeda K and Matsumoto H. 1999. Biosynthesis of phytosiderophores in several *Triticeae* species with different genomes. Journal of Experimental Botany **50**, 723-726.

Maere S, Heymans K, Kuiper M. 2005. BiNGO: a Cytoscape plugin to assess overrepresentation of gene ontology categories in biological networks. Bioinformatics **21**, 3448–3449.

Mai HJ, Pateyron S, Bauer P. 2016. Iron homeostasis in Arabidopsis thaliana: transcriptomic analyses reveal novel FIT-regulated genes, iron deficiency marker genes and functional gene networks. BMC Plant Biology **16**, 211.

Marschner H. 1995. Mineral nutrition of higher plants. Second edition. Academic Press Edition London.

Mayer KFX, Rogers J, Dole el J, *et al.* 2014. A chromosome-based draft sequence of the hexaploid bread wheat (*Triticum aestivum*) genome. Science **345**, 1251788–1251788.

Merico D, Isserlin R, Stueker O, Emili A, Bader GD. 2010. Enrichment Map: A network-based method for gene-set enrichment visualization and interpretation. PLoS ONE **5**, e13984.

Mori S. 1999. Iron acquisition by plants. Current Opinion in Plant Biology **2**, 250–253.

Morrissey J, Guerinot ML. 2009. Iron uptake and transport in plants: the good, the bad, and the ionome. Chemical Reviews **109**, 4553–4567.

Nishizawa NK. 2007. Expression and enzyme activity of glutathione reductase is upregulated by Fe-deficiency in graminaceous plants. Plant Molecular Biology **65**, 277–84.

Nozoye T, Nagasaka S, Kobayashi T, Sato Y, Uozumi N, Nakanishi H, Nishizawa NK. 2015. The Phytosiderophore efflux transporter TOM2 is involved in metal transport in rice. Journal of Biological Chemistry **290**, 27688–27699.

Nozoye T, Nagasaka S, Kobayashi T, Takahashi M, Sato Y, Sato Y, Uozumi N, Nakanishi H, Nishizawa NK. 2011. Phytosiderophore efflux transporters are crucial for iron acquisition in graminaceous plants. Journal of Biological Chemistry **286**, 5446–5454.

Ogo Y, Itai RN, Kobayashi T, Aung MS, Nakanishi H, Nishizawa NK. 2011. OsIRO2 is responsible for iron utilization in rice and improves growth and yield in calcareous soil. Plant Molecular Biology **75**, 593–605.

Ogo Y, Nakanishi Itai R, Nakanishi H, Kobayashi T, Takahashi M, Mori S, Nishizawa NK. 2007. The rice bHLH protein OsIRO2 is an essential regulator of the genes involved in Fe uptake under Fe-deficient conditions. The Plant Journal **51**, 366–377.

Palmer CM, Hindt MN, Schmidt H, Clemens S and Guerinot ML. 2013. MYB10 and MYB72 are required for growth under iron-limiting conditions. PLoS Genetics **9**, 1003953.

Palmer L, Dias D, Boughton B, Roessner U, Graham R, Stangoulis JC. 2014. Metabolite profiling of wheat (*Triticum aestivum* L.) phloem exudate. Plant Methods **10**, 27.

Powell JJ, Fitzgerald TL, Stiller J, Berkman PJ, Gardiner DM, Manners JM, Henry RJ, Kazan K. 2017. The defence-associated transcriptome of hexaploid wheat displays homoeolog expression and induction bias. Plant Biotechnology Journal **15**, 533–543.

Quanbeck SM, Brachova L, Campbell AA, *et al*. 2012. Metabolomics as a hypothesis-generating functional genomics tool for the annotation of *Arabidopsis thaliana* genes of "Unknown Function". Frontiers in Plant Science **3**, 15.

Quinet M, Vromman D, Clippe A, Bertin P, Lequeux H, Dufey I, Lutts S and Lefevre I. 2012. Combined transcriptomic and physiological approaches reveal strong differences between short☐and long☐term response of rice (*Oryza sativa*) to iron toxicity. Plant, Cell & Environment **35**, 1837-1859.

Ramírez L, Bartoli CG, Lamattina L. 2013. Glutathione and ascorbic acid protect Arabidopsis plants against detrimental effects of iron deficiency. Journal of Experimental Botany **64**, 3169–78.

Ramírez-González RH, Borrill P, Lang D, et al. 2018. The transcriptional landscape of polyploid wheat. Science **361**, 6089.

Romera FJ, Alcantara E. 1994. Iron-deficiency stress responses in cucumber (*Cucumis sativus* L.) roots (a possible role for ethylene?). Plant Physiology **105**, 1133–1138.

Romera FJ, García MJ, Alcántara E, Pérez-Vicente R. 2011. Latest findings about the interplay of auxin, ethylene and nitric oxide in the regulation of Fe deficiency responses by Strategy I plants. Plant Signaling & Behavior **6**, 167–70.

Rouached H, Rhee SY. 2017. System-level understanding of plant mineral nutrition in the big data era. Current Opinion in Systems Biology **4**, 71–77.

Rubio V. 2001. A conserved MYB transcription factor involved in phosphate starvation signaling both in vascular plants and in unicellular algae. Genes & Development **15**, 2122–2133.

Santi S, Cesco S, Varanini Z, Pinton R. 2005. Two plasma membrane H+-ATPase genes are differentially expressed in iron-deficient cucumber plants. Plant Physiology and Biochemistry **43**, 287–292.

Santi S, Schmidt W. 2009. Dissecting iron deficiency-induced proton extrusion in Arabidopsis roots. New Phytologist **183**, 1072–1084.

Schaaf G, Ludewig U, Erenoglu BE, Mori S, Kitahara T, von Wirén N. 2004. ZmYS1 functions as a proton-coupled symporter for phytosiderophore- and nicotianamine-chelated metals. Journal of Biological Chemistry **279**, 9091–9096.

Schaedler TA, Thornton JD, Kruse I, Schwarzländer M, Meyer AJ, van Veen HW, Balk J. 2014. A conserved mitochondrial ATP-binding cassette transporter exports glutathione polysulfide for cytosolic metal cofactor assembly. Journal of Biological Chemistry **289**, 23264–23274.

Schmidt H, Günther C, Weber M, Spörlein C, Loscher S, Böttcher C, Schobert R and Clemens S. 2014. Metabolome analysis of Arabidopsis thaliana roots identifies a key metabolic pathway for iron acquisition. PloS ONE **9**, 102444.

Schmidt W, Tittel J, Schikora A. 2000. Role of hormones in the induction of iron deficiency responses in Arabidopsis roots. Plant Physiology **122**, 1109–18.

Shanmugam V, Wang Y-W, Tsednee M, Karunakaran K, Yeh K-C. 2015. Glutathione plays an essential role in nitric oxide-mediated iron-deficiency signaling and iron-deficiency tolerance in Arabidopsis. The Plant Journal **84**, 464–77.

Shukla D, Rinehart CA, Sahi SV. 2017. Comprehensive study of excess phosphate response reveals ethylene mediated signaling that negatively regulates plant growth and development. Scientific Reports **7**, 3074.

Sipos K, Lange H, Fekete Z, Ullmann P, Lill R, Kispal G. 2002. Maturation of cytosolic iron-sulfur proteins requires glutathione. The Journal of Biological Chemistry **277**, 26944–9.

Sperotto RA, Vasconcelos MW, Grusak MA, Fett JP. 2012. Effects of different Fe supplies on mineral partitioning and remobilization during the reproductive development of rice (*Oryza sativa* L.). Rice **5**, 27.

Takagi SI. 1976. Naturally occurring iron-chelating compounds in oat-and rice-root washings: I. Activity measurement and preliminary characterization. Soil Science and Plant Nutrition **22**, 423-433.

Terzano R, Mimmo T, Vekemans B, Vincze L, Falkenberg G, Tomasi N, Ramos MS, Pinton R and Cesco S. 2013. Iron (Fe) speciation in xylem sap by XANES at a high brilliant synchrotron X-ray source: opportunities and limitations. Analytical and Bioanalytical Chemistry **405**, 5411-5419.

Thimm O, Bläsing O, Gibon Y, Nagel A, Meyer S, Krüger P, Selbig J, Müller LA, Rhee SY, Stitt M. 2004. MAPMAN: a user-driven tool to display genomics data sets onto diagrams of metabolic pathways and other biological processes. The Plant Journal **37**, 914–39.

Thimm O, Essigmann B, Kloska S, Altmann T, Buckhout TJ. 2001. Response of Arabidopsis to iron deficiency stress as revealed by microarray analysis. Plant Physiology **127**, 1030–43.

Thomine S, Vert G. 2013. Iron transport in plants: better be safe than sorry. Current Opinion in Plant Biology **16**, 322–7.

Tolay I, Erenoglu B, Römheld V, Braun HJ, Cakmak I. 2001. Phytosiderophore release in *Aegilops tauschii* and *Triticum* species under zinc and iron deficiencies. Journal of Experimental Botany **52**, 1093–9.

Wagner DS, Pirhalla JL, Bowers GD. 2013. Metabolite structure analysis by high-resolution MS: supporting drug-development studies. Bioanalysis **5**, 463–479.

Wang Y, Lysøe E, Armarego-Marriott T, Erban A, Paruch L, van Eerde A, Bock R, Liu-Clarke J. 2018. Transcriptome and metabolome analyses provide insights into root and root-released organic anion responses to phosphorus deficiency in oat. Journal of Experimental Botany **69**, 3759–3771.

Waters BM, Sankaran RP. 2011. Moving micronutrients from the soil to the seeds: Genes and physiological processes from a biofortification perspective. Plant Science **180**, 562–574.

Xie C, Mao X, Huang J, Ding Y, Wu J, Dong S, Kong L, Gao G, Li C-Y, Wei L. 2011. KOBAS 2.0: a web server for annotation and identification of enriched pathways and diseases. Nucleic Acids Research **39**, W316-22.

Yang TJW, Lin WD, Schmidt W. 2010. Transcriptional profiling of the Arabidopsis iron deficiency response reveals conserved transition metal homeostasis networks. Plant Physiology **152**, 2130–2141.

Yordem BK, Conte SS, Ma JF, Yokosho K, Vasques KA, Gopalsamy SN, Walker EL. 2011. *Brachypodium distachyon* as a new model system for understanding iron homeostasis in grasses: phylogenetic and expression analysis of Yellow Stripe-Like (YSL) transporters. Annals of Botany **108**, 821–833.

Yousfi S, Rabhi M, Abdelly C, Gharsalli M. 2009. Iron deficiency tolerance traits in wild (*Hordeum maritimum*) and cultivated barley (*Hordeum vulgare*). Comptes Rendus Biologies **332**, 523–533.

Zanin L, Venuti S, Zamboni A, Varanini Z, Tomasi N, Pinton R. 2017. Transcriptional and physiological analyses of Fe deficiency response in maize reveal the presence of Strategy I components and Fe/P interactions. BMC Genomics **18**, 154.

Zhang Y, Liu J, Zhou Y, Gong T, Wang J, Ge Y. 2013. Enhanced phytoremediation of mixed heavy metal (mercury)–organic pollutants (trichloroethylene) with transgenic alfalfa co-expressing glutathione S-transferase and human P450 2E1. Journal of Hazardous Materials **260**, 1100–1107.

Zheng L, Huang F, Narsai R, *et al.* 2009. Physiological and transcriptome analysis of iron and phosphorus interaction in rice seedlings. Plant Physiology **151**, 262–274.

Zuchi S, Watanabe M, Hubberten H-M, *et al.* 2015. The interplay between sulfur and iron nutrition in tomato. Plant Physiology **169**, 2626-2639.

Table 1: Metal concentration (mg/g of tissue dry weight) in roots and shoots of wheat seedlings subjected to –Fe stress.

Treatments	ROOTS					SHOOTS				
	Fe	Zn	Mn	Mg	Cu	Fe	Zn	Mn	Mg	Cu
+Fe+P	106.73 ± 14.2	14.6 ± 3.2	15.8 ± 2.0	542 ± 70	7.4 ± 1.1	72.9 ± 1.7	26.8 ± .09	17.4 ± .05	1031 ± 2.3	10.4 ± 0.6
–Fe +P	66.8 ± 17	**40.4** ± 12	**38.2** ± 8.9	**912** ± 213	**17.9** ± 3.7	28.6 ± 1.6	22.3 ± 1.05	22.1 ± 1.9	1013 ± 48	5.1 ± 0.3

Table 2: List of genes showing genome induction bias contribution from either A or B sub-genome. Expression changes within triads w.r.t. iron stress were observed to identify the presence of differential regulation among homoeologs from each triad, thus contributing towards genome induction bias. Table gives descriptive list of homoeologs that were up/down-regulated under iron starvation only in the A sub-genome, followed by those only in the B sub-genome.

Gene	logFold change	RAP-DB description
A genome		
TRIAE_CS42_3AL_TGACv1_197291_AA0665720	3.01	Myb transcription factor domain containing protein.
TRIAE_CS42_2AL_TGACv1_094922_AA0304900	2.8	Similar to Prolyl endopeptidase (EC 3.4.21.26) (Post-proline cleaving enzyme) (PE).
TRIAE_CS42_5AL_TGACv1_379416_AA1256390	2.44	Similar to Solute carrier family 35, member F1.
TRIAE_CS42_2AL_TGACv1_094606_AA0300340	2.18	RmlC-like jelly roll fold domain containing protein.
TRIAE_CS42_3AS_TGACv1_211026_AA0683370	3.36	Protein of unknown function DUF1399 family protein.
TRIAE_CS42_7AL_TGACv1_557470_AA1781720	3.47	Heavy metal-transporting P1B-ATPase, Root-to-shoot cadmium (Cd) translocation
TRIAE_CS42_6AL_TGACv1_471682_AA1512850	2.26	Similar to CRT/DRE binding factor 1.
TRIAE_CS42_6AS_TGACv1_485332_AA1543320	1.85	Zinc/iron permease family protein.
TRIAE_CS42_2AL_TGACv1_092944_AA0267670	2.43	Heavy metal transport/detoxification protein domain containing protein.
TRIAE_CS42_5AL_TGACv1_375378_AA1220660	2.21	Similar to SUSIBA2 (WRKY protein).
TRIAE_CS42_1AL_TGACv1_000708_AA0017440	2.57	Chlorophyll a-b binding protein 2, chloroplast precursor (LHCII type I CAB-2) (LHCP).
TRIAE_CS42_2AL_TGACv1_093154_AA0273510	1.86	C2 domain containing protein.
TRIAE_CS42_5AL_TGACv1_374025_AA1188170	2.66	Nodulin-like domain containing protein.
TRIAE_CS42_3AL_TGACv1_197522_AA0666570	2.22	Lipase, GDSL domain containing protein.
TRIAE_CS42_7AL_TGACv1_556100_AA1755640	2.34	Bifunctional inhibitor/plant lipid transfer protein/seed storage domain containing protein.
TRIAE_CS42_4AS_TGACv1_306527_AA1009640	2.2	Multi antimicrobial extrusion protein MatE family protein.
TRIAE_CS42_7AS_TGACv1_570336_AA1834060	1.98	Major facilitator superfamily protein.
TRIAE_CS42_2AL_TGACv1_093456_AA0280470	2.15	Glycolipid transfer protein domain domain containing protein.
TRIAE_CS42_1AL_TGACv1_002205_AA0039730	-2.09	Similar to IAA8 (Fragment).

TRIAE_CS42_4AL_TGACv1_290815_AA0989640	-2	Lipase, class 3 family protein.
TRIAE_CS42_5AL_TGACv1_374561_AA1203290	-2.24	Cinnamyl alcohol dehydrogenase (EC 1.1.1.195).
TRIAE_CS42_5AL_TGACv1_374359_AA1197930	-2.36	Similar to Lipoxygenase L-2 (EC 1.13.11.12).
TRIAE_CS42_6AL_TGACv1_472625_AA1524260	-1.89	RAG1-activating protein 1 homologue domain containing protein.
TRIAE_CS42_7AL_TGACv1_558250_AA1791610	-2.32	Similar to Pleiotropic drug resistance protein 3.
TRIAE_CS42_2AL_TGACv1_097246_AA0323500	-1.72	Similar to Peroxidase (EC 1.11.1.7).
TRIAE_CS42_7AL_TGACv1_556210_AA1758330	-2.21	Similar to Kaurene synthase A (Fragment).
TRIAE_CS42_1AL_TGACv1_000555_AA0014640	-1.84	No apical meristem (NAM) protein domain containing protein.
TRIAE_CS42_6AL_TGACv1_471077_AA1502240	-1.79	Similar to OSIGBa0145M07.8 protein.
TRIAE_CS42_7AL_TGACv1_559906_AA1801190	-1.68	Similar to H0801D08.12 protein.
TRIAE_CS42_7AL_TGACv1_558101_AA1790150	-1.76	Similar to Acyl-ACP thioesterase (Fragment).
TRIAE_CS42_6AL_TGACv1_472321_AA1520860	-2.06	Similar to Subtilisin-like protease (Fragment).
B Genome		
TRIAE_CS42_5BS_TGACv1_423346_AA1374840	2.19	Similar to Calmodulin NtCaM13.
TRIAE_CS42_3B_TGACv1_224030_AA0791180	2.2	Similar to IN2-2 protein.
TRIAE_CS42_2BL_TGACv1_129296_AA0377070	2.46	Similar to OSIGBa0127A14.7 protein.
TRIAE_CS42_6BL_TGACv1_499646_AA1588130	2.32	TGF-beta receptor, type I/II extracellular region family protein.
TRIAE_CS42_7BS_TGACv1_592587_AA1940840	2.25	Similar to RING-H2 finger protein ATL1R (RING-H2 finger protein ATL8).
TRIAE_CS42_2BS_TGACv1_146290_AA0461590	2.64	NA
TRIAE_CS42_5BL_TGACv1_404610_AA1306090	1.84	Similar to Senescence-associated protein SAG102.
TRIAE_CS42_5BL_TGACv1_405319_AA1324840	2.77	Similar to Transporter associated with antigen processing-like protein.
TRIAE_CS42_7BL_TGACv1_577614_AA1879540	2.69	Peptidase A1 domain containing protein.
TRIAE_CS42_4BS_TGACv1_329166_AA1098520	2.32	Similar to Alcohol dehydrogenase.
TRIAE_CS42_4BL_TGACv1_321683_AA1064050	2.56	Protein of unknown function DUF1262 family protein.
TRIAE_CS42_7BL_TGACv1_591489_AA1920550	NA	Similar to zinc transporter 4.
TRIAE_CS42_7BL_TGACv1_577301_AA1871590	2.41	Delta-tonoplast intrinsic protein.
TRIAE_CS42_4BS_TGACv1_329309_AA1100040	NA	Similar to Major facilitator superfamily antiporter.
TRIAE_CS42_2BL_TGACv1_129348_AA0379680	2	Hypothetical conserved gene.
TRIAE_CS42_2BS_TGACv1_148847_AA0495340	4.34	Helix-loop-helix DNA-binding domain containing protein.
TRIAE_CS42_2BL_TGACv1_130820_AA0418390	1.88	NA
TRIAE_CS42_1BL_TGACv1_031794_AA0120680	-2.79	Divalent ion symporter domain containing protein.
TRIAE_CS42_5BL_TGACv1_406838_AA1350360	-2.29	Similar to anther-specific proline-rich protein APG.
TRIAE_CS42_2BL_TGACv1_130182_AA0405730	-2.09	Protein of unknown function DUF3741 domain containing protein.
TRIAE_CS42_4BS_TGACv1_330685_AA1108490	-2.83	Serine/threonine protein kinase-related domain containing protein.
TRIAE_CS42_2BS_TGACv1_147909_AA0488910	-2.28	Delayed-early response protein/equilibrative nucleoside transporter family protein.
TRIAE_CS42_3B_TGACv1_224656_AA0799480	-1.83	Glyoxalase/bleomycin resistance protein/dioxygenase domain containing protein.
TRIAE_CS42_7BL_TGACv1_577476_AA1876230	-2.93	Protein kinase, catalytic domain domain containing protein.
TRIAE_CS42_6BL_TGACv1_499809_AA1592210	-1.88	Cupredoxin domain containing protein.
TRIAE_CS42_7BL_TGACv1_576755_AA1853830	-2.04	Cellulase (EC 3.2.1.4).

TRIAE_CS42_2BL_TGACv1_130686_AA0416110	-2.3	Similar to OSIGBa0096P03.3 protein.
TRIAE_CS42_2BS_TGACv1_146035_AA0453750	NA	Conserved hypothetical protein.
TRIAE_CS42_1BL_TGACv1_030699_AA0098220	-1.91	Lipase, class 3 family protein.
TRIAE_CS42_2BL_TGACv1_129634_AA0391150	-1.78	2OG-Fe(II) oxygenase domain containing protein.

Table 3: Percentage of homoeolog triads categorised into ideal genome expression bias categories in control and Fe starved conditions.

	Control	–Fe
Balanced	77.07%	77.89%
A-suppressed	6.90%	6.90%
B-suppressed	7.09%	6.82%
D-suppressed	5.15%	4.93%
A-dominant	1.12%	0.94%
B-dominant	1.24%	1.18%
D-dominant	1.43%	1.32%

Legends for Figures:

Fig. 1. Effect of Fe-starvation (–Fe) on the growth parameters of wheat seedlings post 20 days after starvation. (A) Phenotype of wheat seedlings exposed to Fe starvation. (B) Total biomass of roots and shoots of wheat seedlings after 20 DAS. 12-15 seedlings were collected for calculating the fresh tissue weight (in grams). (C) Number of first order lateral roots in roots subjected to –Fe condition and control plants (+Fe). (D) Primary root length of wheat roots. 10-12 seedlings were used for measuring the total primary root length of wheat seedlings under –Fe and +Fe condition. # indicate significant difference at $p<0.05$.

Fig. 2. Analysis of RNAseq data from the wheat roots during Fe starvation. (A) Principal component analysis of samples from control (Control_0, Control_1) and –Fe (Single1_0, Single1_1) conditions. (B) Volcano plot of DEGs, the x-axis shows the log2 fold change difference in the expression of genes in iron starved condition w.r.t. control, and the y-axis indicates the negative log of p-value (pval) for the differences in expression. Genes without significant differences are indicated by grey dots. Significant genes with logFC > 0 are represented by red dots, and those with logFC < 0 are represented by green dots in the scatter plot. (C) Number of DEGs in –Fe. Up: Upregulated under –Fe (logFC > 1), Down: Downregulated under –Fe (logFC < -1), Exc–Fe: exclusively expressed in –Fe, ExcCtrl: Exclusively expressed in control condition.

Fig. 3. Genomic distribution and homoeolog bias studies during Fe starvation. (A) Chromosomal inclination of DEGs. (B) Pie charts showing (left panel) sub-genomic distribution of genome induction bias in triads where one of the homoeologs was up/down-regulated. A, B and D depict the sub-genome to which the DE homoeolog belongs; (right Panel) distribution of triads for which two of the homoeologs were differentially expressed and the third one had normal expression. AB refers to the triads for which up/down regulation was observed in the homoeologs belonging to A and B sub-genomes, while the D sub-genome homoeolog behaved normal w.r.t control. (C) Sankey diagram depicting the homoeolog expression bias in Control condition and Fe starvation. Homoeolog triads were classified into the defined seven categories based on relative normalised expression within each triad. Nodes flowing from Control to –Fe (Fe starvation) represent the triads with same (flow to same category) as well as changed (flow to a different category) expression patterns across both conditions. Distinct colors represent the flow of triads belonging to the seven categories form control condition into same category under –Fe or transition into a different category under –Fe.

Fig. 4. Top up-regulated and down-regulated genes in –Fe condition, annotated via KOBAS using rice RAP-DB/RefSeq annotations as reference. The heat map shows top 50 genes those are highly up-regulated (red-left panel) and down-regulated (green-right panel) identified in wheat roots under –Fe condition with respect to control. For expression analysis, FPKM values were obtained using Cufflinks, and CuffDiff was used to identify DEGs by calculating significant changes in transcript expression between the stressed and normal samples (FDR≤0.05).

Fig. 5. Gene ontology (GO) categorization of the differentially expressed genes and its analysis. (A) WEGO plot describing GO annotation and classification of DEGs, with y-axis showing the percentage of genes belonging to respective GO terms (grey bars for down-regulated genes and red bars for up-regulated genes) as well as the right plane ticks for y-axis depicting the number of both up- and down-regulated genes represented by the respective percentages on the left plane. Percentage and number of genes were calculated for the three broad main categories listed on the x-axis; (B) Enriched GO terms in DEGs under –Fe condition, y-axis depicting the significance of GO term enrichment. (C) Figure showing top 20 enriched KEGG pathways represented by the up-regulated genes under iron starvation condition. x-axis showing names of the pathways, y-axis representing the number of genes enriched in respective pathways.

Fig. 6. Co-expression/hub genes and function enrichment network for identified DEGs. Function enrichment network for DEGs associated under Iron starvation in wheat roots with high significance (FDR ≤ 0.05) for (A) Up-regulated and (B) Down-regulated genes. Enriched GO functional categories are clustered with correlated DEGs and represented by node circles.

Fig. 7. Transcriptional factors (TFs) significantly associated with Fe starvation (FDR ≤ 0.05) in wheat roots. (A) List of TFs differentially expressed in response to Fe starvation stress. Blue bars represent up-regulated and red bars represent down-regulated TFs. (B) Co-expression/hub genes and network analysis using Fe responsive TFs (FDR ≤ 0.05) in wheat roots. Circles indicate the processes associated with it (green for down-regulated genes; Blue for up-regulated genes). Enriched GO functional categories are clustered with correlated TFs and represented by node circles.

Fig. 8. Measurement of GST activity and metabolite profiling of wheat roots subjected to Fe starvation. (A) Glutathione-S-transferase activity of wheat roots under Fe starvation (–Fe) and control (+Fe) condition. (B) Metabolite profiling of amino acids, sugars, polyols, organic acids and related compounds. Change in abundance of significant (P < 0.05) metabolites identified by GC-MS in Fe-starved roots. Abundance variation of each metabolite is represented in Log2 fold values of response ratio (–Fe /+Fe) of metabolite concentrations. Values are means of three biological replicates with bar representing Log ratio of standard error. * indicate significant difference at p<0.01; # indicate significant difference at p<0.05.

Fig. 9. Schematic representation describing the core components involved in Fe starvation. The red font indicates the genes/metabolite those were highly up-regulated/showed high-accumulation during our study; whereas the green font indicates down-regulated/low accumulation. Methionine salvage pathway bins indicate the level of the gene expression levels for the transcript indicated next to it. Red and blue bins near pathways and steps represent the up- and down-regulation of related genes, respectively. (Abbreviation used: FBP: fructose-1,6-bisphosphatase I, PFP: diphosphate-dependent phosphofructokinase, ALDO: fructose-bisphosphate aldolase, GAPDH: glyceraldehyde 3-phosphate dehydrogenase, PGAM: 2,3-bisphosphoglycerate-dependent phosphoglycerate mutase, PDC: pyruvate decarboxylase, GLYK: D-glycerate 3-kinase, GlyA: glycine hydroxymethyltransferase, ltaE: threonine aldolase, BMT2: homocysteine S-methyltransferase, metE: homocysteine methyltransferase, metK: S-adenosylmethionine synthetase, SamDC: S-adenosylmethionine decarboxylase, SRM: spermidine synthase, MTN: 5'-methylthioadenosine nucleosidase, mtnK: 5-methylthioribose kinase, mtnA: methylthioribose-1-phosphate isomerase, DEP1: enolase-phosphatase E1, mtnC: enolase-phosphatase E1, mtnD: 1,2-dihydroxy-3-keto-5-methylthiopentene dioxygenase, TyrAT: tyrosine aminotransferase, NAS: nicotianamine synthase, NAAT: nicotianamine aminotransferase, DMAS: 3"-

deamino-3"-oxonicotianamine reductase, YSL: Yellow Stripe Like, maiA: maleylacetoacetate isomerase, AAT: aspartate aminotransferase, TAT: tyrosine aminotransferase, HPD: 4-hydroxyphenylpyruvate dioxygenase, GGT: gamma-glutamyltranspeptidase, AOS3: hydroperoxide dehydratase, OPR: 12-oxophytodienoic acid reductase, ACX: acyl-CoA oxidase, MFP2: enoyl-CoA hydratase, ACAT: acetyl-CoA acyltransferase 1)

Supporting information

Supplementary Fig. S1: qRT-PCR validation of selected genes from the DEGs during Fe-deficient roots after 20 days of starvation. A total of 2 µg of RNA (DNA free) was used for cDNA synthesis and qRT-PCR was performed using gene specific primers (Supplementary Table S1). C_t values were normalized against wheat *ARF1* as an internal control.

Supplementary Fig. S2: Overview of genes modulated by the Fe starvation in wheat roots. MapMan overview demonstrating differentially expressed transcripts under iron starvation in general metabolic pathways. Log2 fold change values of DEGs were imported into MapMan. Read and blue bins represent up-regulation and down-regulation, respectively, in terms of log2 fold change, as shown by the scale.

Supplementary Fig. S3: MapMan visualization depicting the differentially expressed transcription factors families, with red and blue colored bins for up and down-regulated transcripts, respectively. Numbers in the scale represent fold changes in expression levels expressed as Log2.

Supplementary Fig. S4: Estimation of Nitrate levels under Fe-starvation using salicylic acid method. Three Biological replicate root samples of 10, 15 and 20 days after starvation (DAS) were completely dried for the extraction. Yellow coloration (in test tubes: lower panel) represents level of nitrate in the sample. Potassium Nitrate was used as standard in 0-70 µg concentrations.

Supplementary Fig. S5: Characterization of wheat ABCB25 transporter. (A) Phylogeny analysis of TaABCB25 along with its closest orthologs from rice and arabidopsis. (B) Expression analysis of *TaABCC25* in roots of wheat seedlings subjected to Fe starvation. Wheat seedlings (5-7 days old) were subjected to Fe stress and samples were harvested after 12 hours, 3, 6, 9 and 15 days(d) post starvation. The relative qRT-PCR was performed using wheat ADP-ribosylation factor (ARF) as an internal control gene. Fold accumulation was calculated with respect to the control roots. (C) Schematic comparision of different domains of AtABCB27, OsABCB25 and TaABC25. The number indicates the predicted amino acid position of the TM domains.

Supplementary Table S1: List of primers used in the current study. *wheat genes named according to rice RAP-DB/RefSeq based on KOBAS annotation.

Supplementary Table S2: Summary of filtered and mapped reads for each sample. Obtained RNA-seq reads were quality filtered using Trimmomatic v0.35. TopHat was used to map the obtained reads to the wheat genome (TGACv1).

Supplementary Table S3: DEGs in response to Fe starvation in wheat roots List of up-regulated genes, downregulated genes (sheet 2), genes exclusively expressed in response to Fe starvation (sheet 3). Table enlists Control and Fe starvation expression values, logFC for Fe starvation wrt Control samples. Each DEG is annotated with information like rice ortholog, gene definition, KEGG Orthology, Pathways and Pfam domains, which were obtained through KOBAS 3.0 stand-alone tool, using *Oryza sativa* RAP-DB and RefSeq as reference.

Supplementary Table S4: List of 8473 homoeolog triads that were used for homoeolog induction and expression biasness analysis. 8473 homoeolog triplets that were expressed in A, B as well as, D genome were selected for homoeolog specific analysis after filtering all the homoeolog triads obtained from from ensembl.

Supplementary Table S5: Expression profiles of genes/gene families involved in Strategy-II mode of Fe uptake. Strategy-II components of Fe uptake were identified and classified based on screening wheat genes annotated by KOBAS for respective KO IDs for NAS, NAAT, YSL and DMAS genes.

Supplementary Table S6: Expression profiles of genes/gene families involved in Strategy-I mode of Fe uptake. Strategy-I components of Fe uptake were identified and classified based on screening wheat genes annotated by KOBAS for respective KO IDs for AHA, IRT, FRO, PEZ genes.

Supplementary Table S7: Gene Ontology analysis of up- and down-regulated genes in response to Fe starvation. WEGO tool was used to categorize DEGs into GO categories and identify significant GO terms. Table lists the number and percentage of up- and down-regulated genes as well as p-values for each GO term.

Supplementary Table S8: Expression profiles of genes/gene of different transcription factors those are differentially up- and down regulated. MapMan was used to identify TFs and categorize them into TF families. Table gives logFC value for starvation *vs* control for

each TF showing significantly altered expression. A gradient of red and green is used for up-regulated and down-regulated TFs respectively.

Supplementary Table S9: Expression profiles of genes involved during the process of glutathione mediated detoxification process. DEGs mapped to glutathione metabolism were identified using KOBAS. Table lists expression values and annotation for genes. Red color denotes up-regulated genes and green color represent down-regulation.

Supplementary Table S10: GC-MS analysis of wheat roots subjected to 20 days of Fe starvation. Each metabolite is represented with concentrations in three independent replicate manner. For concentration calculation, Individual metabolite area was normalized to sample weight and area of internal control (sorbitol). Metabolites with no detectable area in any of the conditions were considered to be the metabolite with minimum area. Delta method approximation was used to calculate standard errors (se) of log-ratio, se log-ratio = 1/ln 2√[(SET /T)2 + (SEC /C)2], where SET and SEC are standard errors of average –Fe and +Fe metabolite abundances. Metabolites with significant (p-value <0.05) differential abundance were plotted.

Figure 1

(A)

(B)

(C)

Figure 2

(A)

-Fe DEGs distribution

(B)

(C)

Figure 3

Figure 4

Figure 5

Figure 6

(A)

(B)

Figure 7

(A)

(B)

Sugar conjugates	α-D-Glucopyranoside
	α-D-Mannopyranoside
Phosphates	Phosphoric acid
Amino Acids	β-Amino isobutyric acid
	Hydroxybutyric acid
	pyroglutamic acid
	L- l-Valine
Sugars	D-Turanose
	Maltose
	D-Erythro-Pentopyranose
	D-Xylofuranose
	2-Desoxy-pentos-3-ulose
	4-Ketoglucose
	Hexopyranose
	Galactose
	Glucose
	Fructose
	β-DL-Arabinopyranose
	β-D-Xylopyranose
Polyols	Glycerol
	Ribitol
	Myo-Inositol
Fatty Acids	Linoleic acid
	Linolenic acid
Polyhydroxy Acids	Gluconic acid
	Glyceric acid
	Arabino-Hexonic acid
	Arabinonic acid
Acids	Citric acid
	Erythro-Pentonic acid
	Pentanoic acid
	Aconitic acid
	Malic acid
	Lactic acid
	Succinic acid
	Malonic acid
	Acetic acid
	Fumaric acid

Log2 fold (-Fe roots/control roots)

Figure 8

Figure 9