# Memory-guided microsaccades

Konstantin-Friedrich Willeke[1,2,3]†, Xiaoguang Tian[1,2,3]†, Antimo Buonocore[1,2]†,
Joachim Bellet[1,2,3], Araceli Ramirez-Cardenas[1,3], Ziad M. Hafed[1,2]*

[1]Werner Reichardt Centre for Integrative Neuroscience, Tuebingen University, Tuebingen, Germany 72076
[2]Hertie Institute for Clinical Brain Research, Tuebingen University, Tuebingen, Germany 72076
[3]Graduate School of Neural and Behavioural Sciences, International Max Planck Research School, Tuebingen University, Tuebingen, Germany 72074

*Correspondence to: ziad.m.hafed@cin.uni-tuebingen.de

†Equal contributions

**Corresponding author address:**

Ziad M. Hafed
Werner Reichardt Centre for Integrative Neuroscience
and
Hertie Institute for Clinical Brain Research
Otfried-Mueller Str. 25
Tuebingen, 72076
Germany
Phone: +49 7071 29 88819

**Abstract**

Microsaccades are overwhelmingly described as involuntary eye movements. Here we show in both human subjects and monkeys that individual microsaccades of any direction can easily be triggered: (1) "on demand", based on an arbitrary instruction, (2) without any special training, (3) without visual guidance by a stimulus, and (4) in a spatially and temporally accurate manner. Subjects voluntarily generated instructed "memory-guided" microsaccades readily, and similarly to how they made normal visually-guided ones. In two monkeys, we also observed midbrain superior colliculus neurons that exhibited movement-related activity bursts exclusively for memory-guided microsaccades, but not for similarly-sized visually-guided movements. Our results demonstrate behavioral and neural evidence for voluntary control over individual microsaccades, supporting recently discovered functional contributions of individual microsaccade generation to visual performance alterations and covert visual selection.

**Introduction**

A vast majority of currently published research on microsaccades, as well as almost all public discourse on them, refers to these tiny fixational eye movements, which occur when we attempt to fix our gaze, as "involuntary" or "spontaneous" (Supplementary Fig. 1). However, likelihoods of microsaccade direction, amplitude, and frequency are systematically modulated under a variety of conditions [1-18], and microsaccade generation is causally affected by activity in the midbrain superior colliculus (SC) [19-22] and the cortical frontal eye fields (FEF) [23], both involved in voluntary eye movement control [24]. Microsaccades can also be suppressed either voluntarily [25] or when perceptually challenging discrimination stimuli are expected [7,8]. Finally, two highly experienced human subjects were described as being able to generate voluntary horizontal or vertical saccades as small as microsaccades [26].

The persistence of descriptions of microsaccades as being involuntary and spontaneous (Supplementary Fig. 1), despite the above evidence, stems from a severe lack of explicit demonstration that these eye movements, like larger saccades, can be generated by naïve subjects "at will", based on abstract task-defined instructions, and without visual guidance. However, such explicit demonstration is critically needed given that the occurrence of any individual microsaccade strongly alters visual performance [3,4,27-32], providing an almost-deterministic link between covert visual attentional effects and the execution of such tiny eye movements [3-5]. Here, we provide such demonstration; we show in both naïve human subjects and monkeys that tiny eye movements thought to be involuntary can be generated at will, based on arbitrary instruction, and without any visual guidance. We also show that these eye movements are governed by highly specific neural circuitry in the SC.

**Results**

*Memory-guided microsaccades are generated "at will"*

We first asked three male rhesus macaque monkeys to perform a memory-guided

microsaccade task [33,34] (Methods). The monkeys were initially trained to maintain

fixation while a brief (58 ms) flash was presented at an eccentricity $\geq 5^o$, greater than

an order of magnitude larger than microsaccade amplitudes (most frequently defined

as $<1^o$, ref. [35], but often significantly smaller, ref. [19]). The monkeys had to remember

flash location for 300-1100 ms, after which the fixation spot was removed. The

monkeys used fixation spot disappearance as the abstract instruction to voluntarily

execute a timely eye movement towards the remembered flash location, and the

target only reappeared after the eye movement had ended (Methods); there was no

specific instruction to re-fixate the reappearing target, but the monkeys often did so

nonetheless. Our task therefore required voluntary eye movement generation based

on an arbitrary instruction that we enforced by task design (fixation spot

disappearance) and without any visual guidance [33,34], and corrective re-fixation

saccades were visually-guided.

After initial training with large saccades, we tested the monkeys on pseudorandomly

chosen stimulus locations, including at small eccentricities typically associated with

microsaccades. We were struck by the fact that the monkeys very easily generated

memory-guided movements even for target eccentricities as low as 6 min arc ($0.1^o$),

and without any need for special training. They simply generalized the rule that they

had learned for target eccentricities $\geq 5^o$. Figure 1a demonstrates this by way of an

example trial from monkey N. The monkey fixated steadily between flash onset (at a

location 6 min arc left of and 12 min arc below fixation) and the instruction to trigger a

movement (the "go signal"). The monkey then successfully generated an oblique

downward-leftward microsaccade ~206 ms after the "go" signal (fixation spot

disappearance). After target reappearance, the monkey corrected the remaining eye

position error (due to overshoot in the memory-guided microsaccade) with a second

visually-guided microsaccade. Note how the reaction time of the corrective, visually-

guided microsaccade (from target reappearance) was comparable (albeit slightly

shorter) to that of the voluntarily generated memory-guided microsaccade (from

fixation spot disappearance), and that the latter movement was directionally accurate

(directed towards the remembered foveal flash location). Thus, spatially and

temporally accurate microsaccades can be triggered "at will" based on arbitrary

instruction and without visual guidance (also see Supplementary Movies 1, 2).

**Figure 1 Memory-guided microsaccades in monkeys and humans. (a)** Eye position measurements from an example trial in monkey N. Upward deflections in each trace mean rightward or upward eye position displacements, respectively, and the position scale bar denotes 15 min arc. A brief target flash appeared 6 min arc to the left of and 12 min arc below fixation. After a "memory interval", the fixation spot disappeared, instructing a memory-guided eye movement towards the remembered flash location. The monkey generated a spatially-accurate memory-guided microsaccade after a reaction time of ~206 ms from fixation spot disappearance. When the target reappeared, a smaller, corrective movement (which was now visually-guided) was triggered after a shorter reaction time. **(b)** Similar observations from an example trial in a human subject. In this case, the target flash occurred at 18 min arc to the right of and 6 min arc above fixation. Note that the human data were recorded with a video-based eye tracker; monkey N data were obtained from a scleral search coil (Methods). Figures 2-4 and Supplementary Figs. 1-4 demonstrate the robustness of the observations in this figure across the population, and Figs. 6-7 and Supplementary Fig. 5 highlight underlying neural mechanisms.

Timely voluntary memory-guided microsaccade generation was consistent in all three

monkeys, and also in 7 human subjects (Methods; Fig. 1b shows an example trial

from one human subject similar in format to that in Fig. 1a). For all flash eccentricities

<$1^o$ (the most typically cited amplitude threshold for microsaccades [19,35]; Methods),

we plotted histograms of all microsaccade latencies after the go signal (fixation spot

disappearance; Fig. 2a, d, g, j). This allowed us to evaluate the reaction times of

instructed, memory-guided microsaccades. We also plotted the histograms of

microsaccade latencies after target reappearance (Fig. 2m, p, s, v), for evaluating the

reaction times of corrective, visually-guided microsaccades. In all cases, memory-

guided microsaccades had reaction times consistent with these movements being

successfully triggered after the instruction to do so [36,37]. Moreover, the latency

distributions that we observed were similar to those of larger memory-guided

saccades with flash locations >1$^{o}$ (Supplementary Fig. 2). Corrective, visually-guided

microsaccades expectedly had shorter reaction times than memory-guided

microsaccades (monkey N: 146 ms vs. 203 ms; monkey M: 160 ms vs. 212 ms;

monkey P: 189 ms vs. 245 ms; humans: 189 ms vs. 252 ms; all $p<10^{-50}$, t-test).

Therefore, timely, voluntary microsaccades were generated "at will" after the abstract

instruction to do so (fixation spot disappearance).

**Figure 2 Memory-guided microsaccades had reaction times and success rates consistent with being genuine responses to task instruction. (a)** Reaction time distribution for memory-guided microsaccades in monkey N. The data shows all trials with target eccentricities <1° (Methods). The monkey had a reaction time distribution typical of saccadic responses. Error bars denote 95% confidence intervals, and the histogram was normalized by the total number of trials. **(b)** Cumulative probability of the same data, demonstrating a plateau of >80% success rate; the monkey successfully generated a memory-guided microsaccade on the great majority of trials. Error bars denote 95% confidence intervals. **(c)** Same data as in **b** but separated by target eccentricity (inset legend). Even the smallest target eccentricities were associated with a majority of successful reactions. **(d-f)** Similar observations from monkey M. **(g-i)** Similar observations from monkey P. **(j-l)** Similar observations from our human subjects (Methods). **(m-x)** We repeated the same analyses above, but now for corrective, visually-guided microsaccades occurring after target reappearance. Reaction times (**m**, **p**, **s**, **v**) were faster than for memory-guided microsaccades, and often exhibited express reactions (e.g. arrow in **m**). However, success rates were significantly lower than in the instructed memory-guided microsaccades

It may be argued that while memory-guided microsaccades did occur in a timely

fashion after the go signal, their likelihood of occurrence across many trials was

simply low and random as opposed to reflecting a deterministic, voluntary reaction to

an explicit "go" instruction. After all, irrelevant abrupt stimulus onsets can reflexively

reset microsaccade generation rhythms [38-40], suggesting that some aspects of

microsaccade behavior may appear to be reflexive in nature. However, when we

plotted cumulative probabilities of memory-guided microsaccades after the go signal

(i.e. all trials with target eccentricity <1$^{o}$), we found that almost every single instructed

trial was associated with a microsaccadic response (Fig. 2b, e, h, k; erroneous trials

were primarily ones with reaction times larger than our upper limit for acceptance,

Methods); this was not the case for, say, microsaccades after flash onset at the

beginnings of trials when monkeys were instructed to maintain fixation

(Supplementary Fig. 3b, e, h, k). These early flash-induced microsaccades were not

instructed, and therefore did not happen on every single trial (e.g. they were missing

in the example trials of Fig. 1); their occurrence depended on a variety of other

factors related to the quality of fixation control [3,29]. More importantly, the "success"

rate for the instructed memory-guided microsaccades (i.e. the fraction of trials in

which a movement was successfully generated within a reasonable reaction time;

Methods; Fig. 2b, e, h, k) was equal to or significantly higher than that for the

corrective, visually-guided microsaccades, which were not instructed (Fig. 2n, q, t, w)

(monkey N: p=0.028; monkey M: p=0.5; monkey P: p<10$^{-8}$; humans: p<10$^{-50}$, $\chi^2$-test).

This indicates that task instruction was a strong impetus to generate the voluntary

microsaccadic eye movements in our task. Expectedly, the lowest success rate for

both memory- and visually-guided (corrective) microsaccades was for the smallest

target eccentricities (Fig. 2c, f, i, l, o, r, u, x).

Overall, these results indicate that memory-guided microsaccades can indeed be

generated explicitly following abstract instruction (i.e. one that is learned from task

design), and even more successfully than uninstructed corrective, visually-guided

movements. These are hallmarks of voluntary behavior. Interestingly, when visually-

guided microsaccades were explicitly instructed in a separate condition (the delayed,

visually-guided saccade task; Methods), their success rates were also less than

those of the memory-guided microsaccades (Supplementary Fig. 4a, d). This is

largely due to the fact that the monkeys reflexively fixated the visually-persistent

target in this task paradigm even before the "go" signal.

*Memory-guided microsaccades are spatially accurate*

High success was also evident in the directional accuracy of memory-guided

microsaccades. For the same data in Fig. 2, we plotted the angular directional

difference between memory-guided microsaccades and target locations (Fig. 3b, f, j,

n). We also plotted the same measure for the uninstructed corrective, visually-guided

microsaccades (Fig. 3c, g, k, o); in this case, we compared corrective microsaccade

direction to the direction of the reappearing target relative to gaze position after the

memory-guided microsaccade had ended (Methods). Once again, in both monkeys

and humans, memory-guided microsaccades were significantly more directionally

accurate than the uninstructed corrective, visually-guided movements. We confirmed

this statistically by comparing the distribution of absolute directional errors for

memory-guided microsaccades to that of absolute directional errors for corrective,

visually-guided microsaccades. In monkey N, the former was $20.4^o$ +/- $5.9^o$ 95% c. i.

and the latter was $27.1^o$ +/- $1.3^o$ 95% c. i. ($p < 10^{-4}$, Watson-Williams Test, circular

analog of the two-sampled t-test); for monkey M, the results were $23.4^o$ +/- $2.5^o$ 95%

c. i. and $31.5^o$ +/- $1.1^o$ 95% c. i. ($p < 10^{-10}$); and for monkey P, they were $33.8^o$ +/- $2.4^o$

95% c. i. and $51.2^o$ +/- $1.4^o$ 95% c. i. ($p < 10^{-50}$). Similarly, for humans, the directional

errors were $23.8^o$ +/- $0.8^o$ 95% c. i. versus $43.1^o$ +/- $0.9^o$ 95% c. i. ($p < 10^{-50}$). When

instructed visually-guided microsaccades were tested separately (during the delayed,

visually-guided saccade task; Methods), directional accuracy was restored to the

levels of memory-guided microsaccades (Supplementary Fig. 4b, e), consistent with

prior results [41].

**Figure 3 Memory-guided microsaccades had better directional accuracy than either microsaccades during fixation or corrective, visually-guided microsaccades. (a)** We plotted the angular distribution of microsaccade direction error (difference between target direction and microsaccade direction; Methods). This is shown for all microsaccades from monkey N occurring in the final 250 ms of fixation before fixation spot disappearance. Microsaccades could occur either towards or away from the remembered target location. The histogram was normalized by the total number of observations. **(b)** The same monkey exhibited highly accurate microsaccade directions towards the remembered target location when explicitly instructed to do so. **(c)** After target reappearance, corrective, visually-guided microsaccades were also target-directed, but their direction errors were significantly more variable than the instructed movements in **b** (see statistics in text). Thus, instructed memory-guided microsaccades were highly directionally accurate. **(d)** Each group of colored arrows shows the directions of 40 randomly selected memory-guided microsaccades when remembered target directions were drawn from the underlying corresponding-colored faint cone. Directionally accurate microsaccades could be generated for all oblique target directions. **(e-h)** Similar observations for monkey M. **(i-l)** Similar observations for monkey P. **(m-p)** Similar observations for our human subjects. In all cases, directional accuracy was the highest for the instructed, memory-guided movements and not for fixational microsaccades (**a**, **e**, **i**, **m**) or corrective, visually-guided ones (**c**, **g**, **k**, **o**); also see Supplementary Fig. 3 for flash-induced microsaccades. For the leftmost column, n=1432, 2875, 4870, 2831 microsaccades for monkey N, monkey M, monkey P, and the human subjects, respectively. The numbers of trials for memory-guided or corrective, visually-guided microsaccades are the same as in Fig. 2.

It may also be argued that the directional accuracy described above was not

necessarily a result of a voluntary "effort of will" to follow task instruction, but instead

an outcome of covert attentional shifts. For example, monkey [7] and human [1,2]

microsaccades exhibit direction biases in covert attentional tasks, and it could be the

case that the requirement to remember target location was enough to bias

microsaccade directions near the ends of trials (perhaps due to sustained covert

attention). However, if so, such a bias should have also appeared during the memory

interval itself. We therefore analyzed microsaccade directions in the final 250 ms of

the memory interval, during instructed fixation (Fig. 3a, e, i, m). Directional accuracy,

relative to target location, was much worse than in the instructed memory-guided

microsaccades (Fig. 3b, f, j, n) ($p<10^{-50}$ for comparing absolute directional error in

memory-guided microsaccades and microsaccades at the end of the memory interval

in each of the monkeys and also in the human data). In fact, the fixational

microsaccades (at the end of the memory interval) were biased either away

(monkeys) or towards (humans) the target location. This difference in directional bias

between species (for the same trial-duration distributions) likely reflects expected

differences in the temporal dynamics of microsaccadic oscillations between humans

and monkeys [3,30,39].

We also analyzed microsaccades immediately after target flash onset, to compare

potential reflexive stimulus-induced effects (perhaps due to exogenous covert

attentional effects after target flash onset) to instructed movements after fixation spot

disappearance (Supplementary Fig. 3c, f, i, l): other than an expected minority of

express visually-driven reactions immediately after flash onset [29], the directions of

flash-induced microsaccades were much more variable than the directions of

instructed memory-guided microsaccades. Therefore, even transient, exogenous

covert attention shifts [1,2] are not sufficient to explain the directional accuracy of the

instructed memory-guided microsaccades (Fig. 3b, f, j, n).

It is also important to note that voluntary, memory-guided microsaccades could be

triggered in all directions. Thus, directional accuracy was not restricted to, say, only

horizontal or only vertical directions. To illustrate this, Fig. 3d shows raw plots of

example memory-guided microsaccade direction vectors, color-coded by target

locations in each of the four quadrants from monkey N. The other two monkeys and

the human subjects also generated directionally accurate memory-guided

microsaccades in all directions (Fig. 3h, l, p). Thus, not only did all subjects and

monkeys successfully generate untrained voluntary eye movements for small-

eccentricity target locations, but these eye movements were also highly directionally

accurate. The directional accuracy was, once again, better than in the uninstructed

corrective, visually-guided movements (Fig. 3c, g, k, o).

We next investigated memory-guided microsaccade amplitudes. We found that such

amplitudes also scaled with remembered target eccentricity, consistent with the

notion that there was voluntary generation of spatially-accurate (in both direction and

amplitude) microsaccadic eye movements in the task. For example, in Fig. 4a, c, e,

g, we plotted the amplitude of memory-guided microsaccades as a function of

remembered target eccentricity (this figure also included larger saccades and

eccentricities for comparison). We used logarithmic scaling to magnify the small

eccentricities associated with microsaccades [20]. There was a small overshoot in

movement amplitude for the smallest target eccentricities, but the amplitude of eye

movements systematically increased with increasing eccentricity of the remembered

target location. That is, despite the initial overshoot, there was scaling of

microsaccade amplitudes with remembered target eccentricity. This scaling of

memory-guided microsaccade amplitude was true in all 3 monkeys (Fig. 4a, c, e), as

well as in the human subjects (Fig. 4g). Interestingly, when the monkeys or subjects

made uninstructed corrective, visually-guided microsaccades in the same task, the

amplitudes of these visually-guided movements (relative to target eccentricity) were

worse than for the instructed memory-guided microsaccades (i.e. did not scale

appropriately; Fig. 4b, d, f, h). Larger corrective movements were accurate in

amplitude. Similarly, when visually-guided microsaccades were instructed (in the

delayed, visually-guided saccade task; Methods), the amplitudes were more accurate

(as expected [41]), and overshoot did not occur as strongly for the tiniest of

eccentricities as in the case of the memory-guided microsaccades (Supplementary

Fig. 4c, f). These results are consistent with all other results above, demonstrating

willful generation of individual eye movements of microsaccadic sizes, based on

abstract task-defined instruction to do so (fixation spot disappearance).

**Figure 4 Memory-guided microsaccade amplitudes scaled better with target eccentricity than corrective, visually-guided microsaccades. (a)** For all target eccentricities in monkey N, we plotted memory-guided movement amplitude as a function of target eccentricity. The data below 1° target eccentricity (vertical dashed line) are from the same trials described in Figs. 1-3 for reaction times, success rates, and directional accuracy. Here, we show their amplitude scaling. Memory-guided microsaccades increased in amplitude with increasing target eccentricity, consistent with them being genuine responses to task instruction, but they showed systematic overshoot (e.g. Fig. 1). The overshoot disappeared for larger eye movements (also see Fig. 5 for a potential explanation). Target eccentricities were binned at the following non-overlapping bin-center values: 0.05, 0.15, 0.25, 0.35, 0.55, 0.75, 1.05, 1.45, 2, 3, 5, 7.5, 10, 15; error bars denote 95% confidence intervals. **(b)** For similar small target eccentricities, corrective, visually-guided microsaccades overshot the target more than the instructed memory-guided microsaccades, and their amplitudes did not scale with target eccentricity as well as the memory-guided movements (despite the presence of a visual target). Larger corrective movements were more accurate. **(c, d)** Similar observations in monkey M.  **(e, f)** Similar observations in monkey P. **(g, h)** Similar observations in the human subjects. In all cases, despite the amplitude

overshoot for the smallest memory-guided microsaccades (see Fig. 5), instructed memory-guided
microsaccades exhibited better amplitude scaling with increasing target eccentricity than corrective,
visually-guided microsaccades (also see Fig. 5). n=2947, 4931, 3972, 10890 for memory-guided
movements of all amplitudes in monkey N, monkey M, monkey P, and the human subjects; n=2528,
4086, 2674, 7002 for corrective, visually-guided movements of all amplitudes in monkey N, monkey M,
monkey P, and the human subjects.

It is also interesting that the latency distributions for the corrective, visually-guided

microsaccades in the same task (Fig. 2m, p, s, v) showed suggestive evidence of the

oculomotor system being implicitly aware of residual errors after the previous

memory-guided saccades or microsaccades (e.g. potential overshoot). Specifically,

there was a distribution of "express" microsaccades with very short reaction times of

<120 ms after target reappearance. Since express microsaccades are most likely

when a movement plan is already well on its way at the time of stimulus onset [29], the

occurrence of such express microsaccades in the corrective, visually-guided

movements after target reappearance indicates that the previously-occurring

memory-guided movements were indeed internally accounted for. This provides

further evidence that memory-guided microsaccades are voluntarily triggered.

*"Expansion" of foveal space in spatial working memory even with manual responses*

Despite the timeliness (Fig. 2a, d, g, j), success rate (Fig. 2b, e, h, k), and directional

accuracy (Fig. 3b, f, j, n) of memory-guided microsaccades, their amplitude

overshoots for some eccentricities (Fig. 4a, c, e, g) may be interpreted as a failure to

make genuine target-directed voluntary eye movements of small amplitude.

However, an alternative possibility is that the overshoots were instead a property of

the encoding of foveal locations in spatial working memory, and not an execution

limitation by the oculomotor system. To investigate this alternative, we ran our human

subjects on two control tasks not explicitly requiring an eye movement to report the

perceived location of the remembered target flash (Methods). Rather, we asked the

subjects to perceptually report the remembered target location, and they did so via

button or computer-mouse presses (Methods). In one variant, the button-press task,

subjects were free to move their eyes while responding, and they had up to 20 s to

finalize their response; in another variant, the mouse pointer task, subjects pointed to

the remembered location (with mouse cursor) while still maintaining fixation. In both

cases, amplitude overshoot for the smallest target eccentricities was still clearly

evident (Fig. 5), much like when eye movements were used to perform the task (the

memory-guided microsaccade task). Therefore, memory-guided microsaccade

amplitude-overshoot (Fig. 4a, c, e, g) was not a failure of the oculomotor system to

voluntarily generate small eye movements. Instead, it reflected an intriguing "foveal

expansion" in the representation of space during working memory (Fig. 5). Consistent

with this, and as stated above, microsaccade amplitude overshoot was less evident

with instructed delayed, visually-guided microsaccades (Supplementary Fig. 4c, f).

We should also note here that directional accuracy was also very high in the

memory-guided manual tasks, similar to Fig. 3.

391
**Figure 5 Overshoot in memory-guided microsaccades was not restricted to eye movement**
**responses; perceptual localization of foveal remembered target locations was also expanded in**
**an illusory manner. (a)** In humans, we tested perceptual localization of remembered target locations
without explicitly requiring an eye movement response. In the button-press task, subjects used a button
box to move a displayed cursor to the remembered target location (Methods). We analyzed their
perceptual reports similarly to eye movement reports. The same overshoot for small target eccentricities
was observed. Error bars denote 95% confidence intervals. **(b)** We also ran a variant of the task (the
mouse pointer task; Methods), in which we also maintained a visual reference at fixation (the fixation
spot). The same overshoot for foveal target eccentricities was still observed. Therefore, the overshoot
of memory-guided microsaccades in Fig. 4 was likely a general property of representing foveal space in
spatial working memory, as opposed to an intrinsic inability to generate small memory-guided
microsaccades. n=6405 and 9062 trials for the button-press and mouse pointer tasks, respectively.
404

*Memory-guided microsaccades recruit neural resources from superior colliculus*

*neurons*

The strongest evidence that we obtained for voluntary control over microsaccades

emerged when we recorded SC neural activity. In monkeys N and M, we recorded

visual, visual-motor, and motor (saccade-related) SC neurons (Methods) during the

memory-guided saccade task. We specifically aimed to sample sites in the rostral

SC, in which microsaccade-related activity is expected [19,20,22], but we also sampled

slightly more caudally (i.e. more eccentrically) for completeness (Methods). We

inserted multi-electrode linear arrays into the SC, and we used offline sorting to

isolate individual putative single units (Methods). Our first goal was to ask whether

SC neurons responded at all during memory-guided microsaccades. For example,

some SC neurons do not emit saccade-related responses in the absence of a visual

saccade target [42-44], and it is not clear whether this can happen for microsaccades or

not.

Individual SC neurons did indeed emit saccade-related motor bursts for memory-

guided microsaccades. For example, in Fig. 6a, we plotted neural activity from a

sample neuron from monkey M. We identified a set of memory-guided

microsaccades of amplitude $<1^{o}$ and direction towards the neuron's movement

response field (RF), shown in Fig. 6b (neural RF's were assessed as described in

Methods, and the movements included in Fig. 6a were directed towards the RF

hotspot location and with angular direction within +/- 90 deg from the vector

connecting gaze center to this location). We plotted radial eye position aligned on

microsaccade onset and also firing rate (individual rows of tick marks indicate action

potential times, with each row representing an individual movement; this formatting is

similar to that in Fig. 1C of ref. [19]). Memory-guided microsaccade-related SC

discharge was clearly present even for memory-guided movements as small as 6 min

arc, and similar to expected saccade-related discharge in general. Therefore,

individual voluntary, memory-guided microsaccades are associated with SC neural

discharge.

**Memory-guided microsaccades**

**Figure 6 Superior colliculus (SC) activity during memory-guided microsaccades. (a)** The top panel
shows radial eye position aligned on microsaccade onset (for memory-guided microsaccades <1°). All
traces were aligned to start (on the vertical axis) from the same eye position. Most movements had a
radial amplitude <30 min arc (0.5°). These movements were selected relative to the response field (RF)
as assessed from the analysis in **b**. The bottom panel shows microsaccade-aligned firing rate. Classic
SC microsaccade-related activity [19] was observed for memory-guided microsaccades. Each row of tick
marks shows the neuron's discharge for an individual movement from the top panel, and the rows are
sorted by memory-guided microsaccade amplitude (black for <30 min arc; gray for 30-60 min arc). Even
the smallest memory-guided microsaccades were associated with SC microsaccade-related discharge
[19]. The gray bar on the x-axis indicates the analysis window used in **b**. Error bars denote 95% confidence
intervals. **(b)** Movement RF of the same neuron. For each movement displacement, plotted in log-polar
coordinates [20], we plotted mean firing rate from the measurement interval in **a**. The neuron, from the
right SC, exhibited a movement RF on the contralateral side along an oblique upward direction. The
neuron clearly responded for memory-guided microsaccades as small as 0.1° (6 min arc). The origin of
the log-polar plot denotes 0.03° radial amplitude. The inset shows spike waveforms from every 100th
spike recorded in the task (gray; 682 waveforms); red shows the average spike waveform (and 95%
confidence intervals) across all recorded spikes.

Not only did we find microsaccade-related responses for memory-guided

microsaccades (Fig. 6), but there were also neurons that discharged *exclusively* for

memory-guided microsaccades and not for similarly sized corrective, visually-guided

microsaccades. Figure 7a shows an additional example neuron from the same

monkey as in Fig. 6, now demonstrating this unexpected phenomenon. We plotted

an estimate of the neuron's movement field on log-polar coordinates up to 1°

eccentricity. In the left panel, we plotted the movement field of the neuron when

memory-guided microsaccades were generated. Clear discharge for specific

movement vectors was evident. In the right panel, we plotted the movement field of

the same neuron when corrective, visually-guided microsaccades of similar

amplitudes and directions were generated. The neuron now did not emit any

movement-related discharge at all (compare to the left panel), even though

movements of similar sizes were executed. These results were also obvious when

we plotted firing rates as a function of time from microsaccade onset in the two

conditions (Fig. 7b). Activity was present when memory-guided microsaccades were

executed, but not when visually-guided movements of the same amplitude and

direction range were generated. Activity for these latter movements looked

indistinguishable from activity during baseline fixation before trial onset (Fig. 7b,

black curve; Methods define baseline intervals). Therefore, not only was there strong

behavioral evidence that naïve monkeys and humans can generate voluntary,

memory-guided microsaccades (Figs. 1-4), but there were also individual SC

neurons showing exclusive responses only for memory-guided microsaccades but

not for similarly sized visually-guided ones.

**Figure 7 Double dissociation of SC microsaccade-related discharge for visually- and memory-guided microsaccades. (a)** Movement RF's of an example SC neuron. Left shows peri-movement discharge (like in Fig. 6b) for memory-guided microsaccades; right shows the same for corrective, visually-guided microsaccades. The neuron was silent for the visually-guided movements, but not for similar-sized memory-guided ones. **(b)** Peri-movement discharge of the same neuron under the two conditions. The black curve (when visible) shows the neuron's activity during fixation before trial onset; the neuron only discharged for memory-guided, but not visually-guided, microsaccades. Error bars denote 95% confidence intervals. **(c, d)** Same as **a**, **b** but for an example neuron only showing microsaccade-related discharge for visually-guided movements (similar to visually-dependent saccade-related, VDSR, neurons [42]). The inset shows (as an example) the analyzed eye movements (like Fig. 6a) in the two conditions, demonstrating clear overlap (similar overlap was present in **a**, **b**). **(e)** Distribution of the different neuron types (V=Visual; VM=Visual-motor; M=Motor; VDSR=Visually-dependent saccade-related; ER=Exclusively responding for memory-guided microsaccades). **(f)** Movement RF hotspots for the neurons. ER-Neurons tended to be more eccentric than other neurons exhibiting microsaccade-related discharge, consistent with a foveal expansion for memory-guided microsaccades (see text and Supplementary Fig. 5). **(g, h)** Peri-stimulus (**g**) or peri-microsaccadic (**h**) discharge for all of our encountered neuron types during microsaccade generation. Each neuron's response was first normalized before averaging responses across neurons. V- and VM-Neurons exhibited strong visual responses; VM- and M-Neurons showed movement-related activity regardless of stimulus; VDSR-Neurons only showed movement-related activity in the presence of a visual target; ER-Neurons only showed movement-related activity in the absence of a visual target. In all cases, we analyzed neural activity for microsaccades (<1°) towards the RF. Error bars denote 95% confidence intervals.

We found 41 SC neurons exhibiting exclusive responses for memory-guided

microsaccades; these constituted 22% of SC neurons with saccade-related

discharge in either memory or visual condition (Fig. 7e, f). Our criteria for including

neurons in this accounting was the observation of visual and/or movement discharge

for eccentricities <1$^\circ$ (Methods). We referred to these neurons as ER-Neurons, for

"**E**xclusively **R**esponding" (Methods). Their average firing rate modulations for foveal

visual target onsets and for microsaccades are shown in Fig. 7g, h (rightmost

column), and these modulations are consistent with those in the example neuron of

Fig. 7a, b. Interestingly, for larger saccades, ER-Neurons did exhibit saccade-related

activity for visually-guided eye movements (Supplementary Fig. 5), suggesting that

memory-guided microsaccades may have recruited additional neural resources from

SC saccade-related neurons that would normally be recruited for slightly larger eye

movements when visual targets are present (Supplementary Fig. 5). Consistent with

this, RF hotspots for ER-Neurons tended to be more eccentric than other neuron

types in our database (Fig. 7f).

Interestingly, we also found all other classic saccade-related SC responses to be

present as well for memory-guided microsaccades. For example, we found visual-

only neurons responding to foveal flash onset in the beginning of a trial but not during

the actual memory-guided (or visually-guided) eye movements (Fig. 7g, h, leftmost

column). We also found visual-motor neurons, exhibiting both a visual response and

either a memory- or visually-guided microsaccade response (Fig. 7g, h, second

column). Motor neurons that responded for both memory- and visually-guided

microsaccades, but not to flash or target onset, were also present (Fig. 7g, h, third

column). Finally, in addition to the ER-Neurons (e.g. Fig. 7a, b), we discovered

neurons exhibiting the exact opposite profile: these were visually-dependent

microsaccade-related neurons, extending earlier findings of SC visually-dependent

saccade-related neurons (VDSR-Neurons [42-44]) to the realm of microsaccades. One

example such neuron is shown in Fig. 7c, d. The neuron exhibited microsaccade-

related activity exclusively during visually-guided microsaccades, but not during

similarly sized memory-guided microsaccades. This neuron type (Fig. 7g, h, fourth

column) was not described previously in refs. [19,20]. Overall, in the memory-guided

microsaccade task, the different classes of neurons that we encountered (Fig. 7g, h)

were distributed in our database (Methods) as shown in Fig. 7e.

Our results demonstrate that memory-guided microsaccades are generated "at will":

based on abstract task-defined instruction, without visual guidance, with high fidelity

in spatial accuracy, and with underlying SC circuitry that supports all different aspects

of memory- or visually-guided behavior. These results demonstrate that

overwhelming descriptions of microsaccadic eye movements as being involuntary

and spontaneous (Supplementary Fig. 1) are too simplistic and can mask the

significance of these tiny eye movements in modulating perceptual and attentional

performance.

## Discussion

We demonstrated clear voluntary control over individual microsaccades even in naïve human subjects and monkeys. We were struck by how easily memory-guided microsaccades could be generated, with no need for special training. Monkeys simply generalized the task rule that they had learned for larger saccades, and humans were only verbally instructed to "look at the remembered target location". We believe that our results overturn an overwhelming discourse on microsaccades, even in public understanding of these eye movements, as being involuntary and spontaneous (Supplementary Fig. 1). Indeed, despite evidence like that cited in **Introduction**, a surprisingly large fraction of references to microsaccades in the literature (especially in the last two decades) asserts that these eye movements are involuntary eye movements (Supplementary Fig. 1).

We believe that explicit demonstration of behavioral and neural evidence for voluntary control over individual microsaccades, as we have done in this study, is absolutely necessary to clarify continued debates about the functional roles of these eye movements in influencing brain activity and behavior. Consider, for example, the field of covert visual attention. A long history of research has assumed for several decades that microsaccades either do not happen in these tasks at the time of task-relevant events, or that they happen in a random and spontaneous manner that should not influence the interpretation of results. However, it is now known that this view is too simplistic [1-4,27-31]. For this important revelation to become appreciated more widely, and in other fields related to visual and cognitive neuroscience, explicit demonstration of voluntary control over individual microsaccades, like we have shown in this study, is firmly needed.

The strongest evidence that we found for voluntary control over individual

microsaccades is related to how the SC is involved in generating memory-guided

versions of these eye movements. We were particularly intrigued by the ER-Neurons

that we found (Fig. 7). These neurons are SC neurons that discharge exclusively for

memory-guided microsaccades but not for similarly sized visually-guided

microsaccades. It would be highly interesting to investigate the mechanisms

associated with these ER-Neurons in more detail in future studies. For example, the

fact that these neurons discharge (i.e. exhibit saccade-related activity) for larger

visually-guided saccades might suggest an expansion of RF's towards the fovea

when spatial working memory is engaged (Supplementary Fig. 5). Prior work is

inconclusive about how SC RF's change during spatial working memory tasks [43,45],

and there is no systematic investigation, to our knowledge, of foveal spatial

representation in working memory anyway. Therefore, this issue needs to be

approached and investigated in future follow-up studies. This can allow

understanding the entire transformation from visual input (flash onset) to memory

representation in the memory interval, and finally to movement generation. Relevant

questions could include whether RF's change during the memory interval as well, or

only at the time of memory-guided microsaccade generation (like in Fig. 7a,b,

Supplementary Fig. 5). Importantly, studying this in the SC would clarify a sub-

cortical involvement in working memory, which is a topic that is generally investigated

more strongly from a cortical perspective [33,46]. Studying this in the SC would also link

SC activity to mechanisms of spatial updating of memory representations across eye

movements [47,48].

We were equally intrigued by the VDSR-Neurons [42-44] that we found for

microsaccades (Fig. 7c, d, g, h). Very few studies have mentioned these neurons,

which only exhibit saccade-related discharge if saccades are directed towards a

visual target. Moreover, these few mentions were only made from the perspective of

larger saccades. Here, we found that VDSR-Neurons also exist in the rostral SC

needed [19] to generate microsaccades. That is, we found microsaccade-related

neurons that only showed movement-related discharge if there was a visual target

guiding microsaccade generation; the same neurons did not discharge when the

microsaccades were generated towards a blank display area. We find these neurons

interesting because they suggest that the oculomotor system, even for

microsaccades, incorporates information about visual features at the time of saccade

execution. It would be interesting to relate such incorporation of visual information in

saccade commands to cortical interactions between saccades and visual

representations [49] and to shape guidance of saccades [49] and microsaccades [39].

Because SC neurons exhibit visual-feature tuning in visual responses (e.g. [50-52]), it

would also be interesting to explore how these VDSR-Neurons behave during

saccades to different visual objects or features.

In all, our neural results with visual, visual-motor, and motor neurons all being

involved in microsaccade generation confirm views that microsaccades are

essentially indistinguishable from larger saccades in terms of neural mechanisms [21].

This is consistent with a variety of neurophysiological evidence in multiple areas

[20,22,23,53,54], and it also gives further credence to the notion that these eye movements

are indeed voluntary movements, despite the opposite description in a large portion

of the literature (Supplementary Fig. 1). The visual responses, in particular, are worth

further investigation, both for understanding foveal vision in the SC in general, and

also for exploring the speeded reactions to target reappearance in the corrective,

visually-guided microsaccades (Fig. 2m, p, s, v).

Behaviorally, there was a remarkable difference between instructed and uninstructed

movements in general. Most strikingly, the corrective, visually-guided microsaccades

in the memory-guided saccade task were both less successful (i.e. less often

generated) and also sloppier in direction and amplitude (e.g. Figs. 2-4) than the

instructed memory-guided movements, even though these latter movements did not

have explicit visual guidance. Only when we instructed visually-guided saccades (in

the delayed, visually-guided saccade task) did visually-guided microsaccades appear

more spatially precise (Supplementary Fig. 4b, c, e, f). This observation is perhaps

expected, but it also supports our interpretation that our memory-guided

microsaccades were generated voluntarily. Further support for this conclusion is the

presence of some express microsaccades after target reappearance (Fig. 2). As

stated earlier, because express microsaccades are more likely when a movement

plan is already under way [29], the presence of these express microsaccades after

target reappearance indicates that the oculomotor system was already planning a

corrective movement for any residual errors from previous memory-guided

microsaccades.

Finally, our results pave the way for interesting investigations of spatial working

memory at the microscopic scale of the fovea. A wealth of research has been

performed on the mechanisms of working memory [33,46] in general. However, just like

subcortical mechanisms are less investigated, how foveal space is represented in

spatial working memory remains unknown. For example, our perceptual control

experiments in humans (Fig. 5) suggest that the overshoot in microsaccade

amplitudes that we observed (Fig. 4) was not a deficit in oculomotor control per se.

Instead, it likely reflected an expansion of foveal space when working memory was

engaged (Fig. 5). This might be a complement of foveal biases in peripheral visual

perception [55-57]. Specifically, because of foveal magnification of neural tissue in the

visual system [58], it could be that engaging spatial working memory in the fovea

causes distortions in the representation of space, such that foveal locations get

mislocalized "outward" (i.e. overshoot) whereas peripheral locations get mislocalized

"inward" (i.e. undershoot). Interestingly, perceptual mislocalizations around the time

of microsaccades [27] are also consistent with "outward" distortions in the fovea and

"inward" distortions in peripheral vision. This might suggest that there may be very

generalized underlying mechanisms of perceptual distortions of different kinds

(whether due to eye movements, working memory, attention, or otherwise), likely

being caused by how individual brain circuits represent visual space in anatomical

tissue, and how multiple such circuits (with potentially different levels and kinds of

representational magnification) may interact with each other [59].

## Methods

*Ethics approvals*

All monkey experiments were approved by ethics committees at the

Regierungspräsidium Tübingen. The experiments were in line with European Union

directives, and German laws, governing animal research.

Human experiments were approved by ethics committees at the Medical Faculty of

Tuebingen University. Subjects provided written, informed consent, and they were

financially compensated for their participation. Our experiments were in accordance

with the Declaration of Helsinki.

*Laboratory setups*

Monkey experiments were performed in the same laboratory as that described

recently [50,51,60]. Human experiments were done in the laboratory described in [27,59].

Eye movements were recorded at 1kHz using magnetic induction [61,62] (for monkeys

N and M) or video-based eye tracking (for monkey P and the human subjects). Head

position in the human subjects was stabilized using a custom head-holder [27].

*Animal preparation*

We collected behavioral data from 3 adult, male rhesus macaques (macaca mulatta).

Monkeys N and M (aged 10 and 7 years, respectively, and weighing 11.5 and 8 kg,

respectively) were implanted with scleral search coils to allow measuring eye

movements using the magnetic induction technique [61,62]. The eye movements of

Monkey P (aged 10 years and weighing 8.6 kg) were recorded using a video-based

eye tracker (EyeLink 1000, SR Research). All 3 monkeys were implanted with a head

holder to stabilize head position during experiments, with details on all implant

surgeries provided earlier [60,63].

We also recorded superior colliculus (SC) neural activity from monkeys N and M. The

SC was approached through recording chambers cranially-implanted on the midline

and aimed at a point 1 mm posterior of and 15 mm above the inter-aural line. The

chambers were tilted backwards from vertical by an angle of $35^o$ in monkey N and

$38^o$ in monkey M. Other details for chamber implant surgeries were provided

elsewhere [28,31,51].

*Monkey behavioral tasks*

Monkeys performed a memory-guided saccade task [33,34]. In each trial, a central white

fixation spot (72 cd/m$^2$ for monkey N; 86 cd/m$^2$ for monkey M; 48.1 cd/m$^2$ for monkey

P) was presented over a uniform gray background (21 cd/m$^2$ for monkey N; 29.7

cd/m$^2$ for monkey M; 4.4 cd/m$^2$ for monkey P). The fixation spot was a square of 5.3

x 5.3 min arc dimensions. After 300-1200 ms, a brief target flash (identical to the

fixation spot) occurred at some location on the display (duration: ~58 ms), while the

fixation spot remained on; this flash represented the location to be memorized by the

monkeys for an upcoming saccadic eye movement. After the flash, the fixation spot

remained visible for 300-1100 ms; this period constituted the memory interval of the

task, after which the fixation spot disappeared. Fixation spot disappearance was the

abstract instruction for the monkeys to generate a spatially-accurate saccadic eye

movement to the remembered flash location. Because of the disappearance of the

fixation spot and the flash before it, the saccade that was triggered was generated

without any visual guidance. Monkeys had to trigger this saccade within <700 ms

from fixation spot disappearance (reaction times were actually much shorter as

shown by the distributions in Fig. 2, Supplementary Fig. 2). After gaze entered within

a virtual window around the true target location by 300 ms, the target reappeared for

400-500 ms. During this time, a visually-guided saccade, which we also referred to

as a corrective saccade, was sometimes triggered to correct for any remaining error

between the memory-guided saccade and the actual target position. This corrective

movement happened readily, and we did not specifically instruct the monkeys to

generate it. It was, however, useful for analysis because individual trials often had

both a memory movement as well as a "control" visually-guided one. The radius of

the virtual window around the invisible target position was <3$^o$ and was varied

depending on target eccentricity. The monkeys were rewarded with water or apple

juice at the end of every trial. If they broke fixation during the fixation interval or failed

to remain within the virtual window around target location (after the instruction to

make the memory-guided saccade), the trial was aborted and repeated.

Across trials, we varied target location (i.e. flash location) between 6 min arc and 16$^o$

throughout the entire display. During SC recordings, we made sure to include within

our sampling of target locations positions that were in and around the response fields

(RF's) of neurons around our recording electrodes, as assessed online using our

neurophysiological data acquisition system (see below). We collected a total of 6428

trials in this task from monkey N, 9631 from monkey M, and 14666 from monkey P.

In monkeys N and M, we also compared memory-guided saccades to those obtained

in a task with explicit visual guidance. In this delayed, visually-guided saccade task,

stimulus events were identical to those described above, except that the target flash

was now replaced by target persistence until trial end. In other words, the target

remained on during the memory interval (now called the delay interval) and also after

fixation spot disappearance. Thus, the instructed saccade (i.e. after the offset of the

fixation spot) was visually-guided. We collected 6147 trials in this task from monkey

N and 5871 trials from monkey M.

*Human behavioral tasks*

Human subjects performed the same memory-guided saccade task described above.

Target locations were chosen, across trials, from a total of 480 possible pre-defined

display positions ranging in eccentricity from 6 min arc to 12$^o$. 288 of these 480

possible target positions were within a rectangle of +/- 0.8$^o$ x +/- 0.8$^o$ size in order to

allow us to explore memory-guided microsaccades in particular detail; the remaining

locations were equally spaced in the rest of the display. Seven subjects (4 naïve and

3 untrained authors) performed this experiment, with each subject completing 480-

600 trials per session and 4-5 sessions (total of 16844 trials overall). Ten practice

trials (with target eccentricities >1$^o$) were performed at the beginning of every

session. This task allowed us to relate the monkey behavioral and neurophysiological

results to human performance. The subjects were aged 23-39, and 3 were female.

We also ran two additional control experiments in the humans, in order to understand

whether intricacies of memory-guided microsaccade behavior (e.g. Fig. 4) were

specific to a particular motor modality (eye movements) or more related to the

engagement of spatial working memory per se. We therefore required a perceptual

assessment of remembered flash location without the use of an instructed eye

movement. In the first control task, which we refer to as the button-press task, trials

were identical to the memory-guided saccade task described above up until the end

of the memory interval. At this point, instead of fixation spot disappearance, the

fixation spot was replaced with a cross-shaped cursor of 59 min arc dimensions.

Subjects used a response box with four buttons allowing them to move the cursor at

individual pixel resolution (1.46 min arc per click) rightward, leftward, upward, or

downward. Individual button presses moved the cursor by individual pixels;

prolonged button presses moved the cursor at one of two faster speeds depending

on the duration of the press. Once subjects were satisfied with the cursor position as

reflecting, as accurately as possible, the remembered flash location, they pressed a

fifth button confirming their choice. The target then reappeared at its true position

with the cursor disappearing simultaneously. Subjects were free to move their eyes

at will when the cursor was visible, and they were given up to 20 s to move the cursor

and decide on their perceptual localization of the remembered target location.

Therefore, this task allowed assessing the quality of target recall independent of

making an instructed memory-guided saccade. Seven subjects (ages: 25-39 years; 1

female) participated in this experiment, four of which had performed the original

saccade version of the task. Each subject participated in 5-6 sessions (250-350 trials

per session), for an overall total of 11249 trials across subjects.

The second control experiment, called the mouse pointer task, was similar to the

button-press task except that the cursor was replaced with a computer mouse

pointer, which appeared at a random position $1^o$-$5^o$ away from the true flashed target

location in every trial. Critically, in this version of the control task, we maintained the

fixation spot on during the manual response interval (i.e. until subjects pressed a

mouse button), so that an explicit foveal reference frame was always present when

subjects were pointing to the remembered target location (similar to how eye

movements presumably relied on the initial reference frame of fixation for

localization). Subjects were instructed to fixate the fixation spot throughout the

response interval until they pressed a mouse button, and they were given up to 15 s

to make their response. This task tested whether an explicit foveal reference frame

altered the overall pattern of behavior from the button-press task or from the original

memory-guided saccade task (Fig. 5). Seven subjects (ages: 25-39 years; 1 female)

performed this task, 4 of which had also participated in the button-press and

memory-guided saccade versions of it. We collected 380-500 trials per session for

each subject. We ran 5 sessions per subject, with the exception of one participant

who completed only 4 sessions (for a total of 16213 trials overall).

*Monkey neurophysiological recordings*

We recorded SC activity from monkeys N and M using multi-electrode linear arrays

(16-channel V-Probes with 150 $\mu$m inter-electrode spacing, Plexon). We performed

25 experiments in monkey N and 16 experiments in monkey M. In each experiment,

we advanced either one or two V-probes towards SC surface. We monitored the

deepest electrode contact in a probe and compared physiological landmarks picked

up on this contact to anatomical landmarks in MRI's obtained from the same animals

before implant surgeries. Once the deepest electrode contact was in the SC, we

advanced the probes further, such that as many electrode contacts as possible were

recording SC activity. We used online spike sorting to guide our approach and

maximize the number of channels that were recording SC activity. When we inserted

two simultaneous V-Probes (23/25 sessions in monkey N), one probe was in the right

rostral SC (rostral being the region where microsaccade-related activity is expected

[19,20]) and the other was at a slightly more caudal position in the left SC (to allow

sampling multiple neuronal preferred eccentricities from the SC's topographic map).

When we inserted a single probe, it could be in the right or left SC and at a variety of

rostral and slightly more caudal sites.

Once the probes were in the SC, we ran a series of experimental tasks, including the

memory-guided and delayed, visually-guided saccade tasks described above. We

also ran a variety of other tasks that were not directly relevant to the present study.

Our neurophysiology system sampled electrical activity in each contact at 40 kHz.

Spike times during online sorting were sampled at 1 kHz.

*Monkey and human behavioral analyses*

We detected saccades and microsaccades using established methods in our

laboratory [63,64]. We manually inspected all trials to correct for false alarms or misses

by the automatic algorithms. We also marked blinks or noise artifacts (which were

more likely in video-based eye tracking data) for later removal. We analyzed all eye

movements occurring in the interval between fixation onset and trial end in the

memory-guided and delayed, visually-guided saccade tasks. In the button-press

control task, subjects were free to move their eyes after cross-shaped cursor

appearance, so we only analyzed eye movements up to such appearance. Individual

subjects had idiosyncratic patterns of cursor trajectories and accompanying eye

movements that did not appear related in systematic ways to task performance; we

thus elected not to analyze eye movements during the response interval in this task.

In the mouse pointer task, we monitored eye movements after cursor appearance to

ensure that subjects maintained fixation while responding.

In this paper, we report each individual monkey's behavioral performance separately.

For the human data, we combined results from all subjects into aggregate

measurements. We felt justified in doing so because the human effects exhibited

very strong similarities to the monkey behavior, which was a primary reason for

performing the human experiments in the first place. Moreover, trial numbers were

uniformly distributed across the individual human subjects as much as possible;

therefore, no single subject biased the pooled results.

In all tasks, we first identified valid trials that we accepted for subsequent analysis.

Valid trials did not have any saccades larger than $1^o$ during the memory or delay

interval (to ensure that fixation was maintained properly), and also no blinks or noise

artifacts within +/- 100 ms from target flash time (or target onset time in the delayed,

visually-guided version of the task). This latter criterion ensured that subjects could

see the target flash well in the memory version of the tasks. We also removed trials if

there were blinks or noise artifacts from -100 ms to +400 ms from the instruction to

generate a memory- or visually-guided saccade. Because microsaccades alter

visibility and modulate visual sensitivity in general [3,30,32,65], we wanted to ensure that

subjects could detect target flash (or onset) and fixation spot disappearance (or other

instruction to respond) with high sensitivity. We therefore required that valid trials

also did not have any microsaccades (or large saccades) within +/- 50 ms from flash

(target) onset or fixation spot disappearance (or any other instruction to respond).

Finally, memory-guided saccades had to land within $2^o$ of the target; this constraint

was irrelevant for memory-guided microsaccades (which were much smaller), but it

helped ensure that the eye movements were accurate when large target

eccentricities were used.

Based on the detected saccades and microsaccades, we computed measures of

saccade amplitude, latency (reaction time), direction, starting point, and endpoint. We

calculated the starting point as the average eye position in the interval 30 ms before

the start of the saccade. The end point was simply the very last sample of the

detected saccade. To obtain saccade direction error (i.e. saccade direction relative to

target direction from initial fixation spot position), we computed the counter-clockwise

angular difference between saccade and target direction. For corrective saccades

after target reappearance in the memory-guided saccade task, the target reappeared

at a position that was variable relative to gaze position (depending on the landing

error of the previous memory-guided movement). We therefore redefined the position

of the reappearing target relative to corrective saccade starting eye position. We then

calculated saccade direction error as above. We used a similar procedure for target

eccentricity. Note that after large memory-guided saccades, target position error for

the corrective movements was often small (i.e. large memory-guided saccades were

relatively accurate as expected). We therefore had significantly more data for

corrective, visually-guided microsaccades than for memory-guided microsaccades

(e.g. see the numbers of trials in the legend of Fig. 2).

In the memory-guided saccade task, we defined as the memory-guided saccade (or

memory-guided microsaccade) of interest as the first saccade occurring with a

latency of at least 50 ms after fixation spot disappearance and less than 50 ms after

target reappearance. Similarly, the corrective, visually-guided saccade was defined

as the first saccade that was executed with latency >50 ms after target

reappearance. Because most studies define microsaccades as smaller than $1^{o}$ in

amplitude and we were interested in comparing these to voluntarily generated

movements, we treated trials with target eccentricities less than $1^{o}$ as memory-

guided microsaccade trials.

The above analyses were also applied in the delayed, visually-guided saccade task.

For the control tasks, we treated button and mouse press locations similarly to how

we analyzed saccade endpoints.

In all analyses, we defined trials with target eccentricities $<1^{o}$ as the trials involving

microsaccades, as stated above. Corrective, visually-guided saccades were also

classified as microsaccades when target eccentricity (relative to eye position at target

reappearance) was $<1^{o}$. As stated above, we used this threshold because it was a

typical threshold used in the literature [19,35], but our results (also from other studies)

confirm that there is indeed a continuum of saccade generation across different

amplitudes.

*Spike sorting for neurophysiological data analyses*

For all neurophysiological data analyses, we performed offline sorting to isolate

individual single units. We isolated single units from wide-band (40 kHz) electrode

data by adapting the Klusta toolbox [66]. As an additional input to the toolbox, we

recreated the electrode geometry (16 channels with linear spacing) such that

neighboring channels could share spike timing and waveform information (as in,

"virtual" tetrodes). From filtered neural data (750-5000 Hz), the toolbox performed

automated spike detection and classification. We inspected the resulting cluster

assignments (putative units) using the Kwik-gui interface included in the Klusta

toolbox. Based on auto- and cross-correlograms, as well as waveform shapes, we

manually curated each cluster in order to obtain well-isolated units. Clusters that

showed no characteristic auto-correlogram, or that produced atypical spike

waveforms, were clustered together, labeled as noise, and subsequently excluded

from further analysis. We estimated single unit isolation quality of all resulting units

using established metrics [67,68]. Specifically, we computed an estimate of the false

positive rate based on refractory period violations (inter-spike interval violations).

Furthermore, we obtained an isolation distance measure for each cluster, by

calculating the Mahalanobis distance between any two units. All isolated units that

exceeded an estimated false positive rate of 10% or had an isolation distance below

30 were excluded from further analysis; all other units are hereafter termed putative

neurons (297 neurons in total; we selected for analysis all 188 neurons that showed

clear SC visual and/or saccade-related RF's encompassing microsaccadic amplitude

ranges, as assessed by the procedures described below). Spike trains of all putative

neurons were then convolved with an appropriately-scaled Gaussian ($\sigma$ = 10 ms) to

obtain an estimation of the neuron's firing rate in spikes/s.

*Neurophysiological data analyses*

We classified SC neurons as visual, visual-motor, or motor depending on visual and

saccade-related responses. We defined several 100-ms intervals based on key trial

events. The visual response interval comprised the time from 50 to 150 ms after

target onset. The motor interval was set to be 100 ms centered around saccade

onset. We also measured this saccade-related interval for corrective saccades in the

memory-guided saccade task. Since we were especially interested in characterizing

microsaccade-related discharge, we classified movement-related aspects of neurons

based on responses associated with saccades <1$^o$ in amplitude. To estimate

baseline activity during fixation without any target or saccade onset, we used the100-

ms steady fixation interval before target onset. We avoided contamination of baseline

and visual neural activity by saccades or microsaccades; we excluded all trials in

which such movements occurred during the measurement intervals. Similarly, we

excluded trials in which there was more than one movement within the motor interval.

In all cases, we measured average firing rate within a given measurement interval, which resulted in a population of measurements across trials; we only included neurons with >10 repetitions per measurement interval. We compared each of visual, memory-guided microsaccade, corrective microsaccade, or visually-guided microsaccade (in the delayed version of the task) activity to baseline activity using one-tailed Wilcoxon-ranksum tests ($\alpha = 0.05$). We then grouped neurons into five categories: (1) Visual (V-Neuron), possessing only a visual increase relative to baseline; (2) Visual-motor (VM-Neuron), possessing a visual increase and either memory- or corrective-(visual) saccade response; (3) Motor (M-Neuron), exhibiting both memory- and corrective-(visual) saccade response but no visual response; (4) Visually-dependent (VDSR-Neuron), possessing a saccade-related response only when a visual target was present for the eye movement [42-44]; (5) Exclusively-responding (ER-Neuron), exhibiting a microsaccade-related response only in the memory-guided microsaccade task but not for similarly sized visually-guided microsaccades.

RF hotspot locations (e.g. Fig. 7f) were estimated based on all of the sampled saccade vector amplitudes (and corresponding flash locations for visual responses) in the memory-guided saccade task (even >1°); our estimates may therefore include neurons that may potentially have responded even more strongly had we obtained even larger eccentricities (at the very least, all neurons in Fig. 7f were related in one form or another to microsaccades). For every isolated neuron that had a motor response, we also estimated the saccadic amplitudes that caused the neuron to discharge, also known as the neuron's movement field. For every neuron, we compared the population of baseline values as described above to the trial-by-trial responses for saccades. We then calculated the movement field by a linear-

interpolation of all z-transformed saccade-related responses. Locations with z-scores

>1.96 were considered to be part of the movement field. We plotted movement fields

using log-polar coordinates, to magnify small eccentricities associated with

microsaccades, as described previously [20]. Movement-related hotspots were

identified based on the stronger of either the memory-guided or the visually-guided

movement-related responses.

*Statistical analyses*

We performed statistical tests (behavioral and neural) with an $\alpha$ level of 0.05 unless

otherwise noted. When the assumptions of normality were not met for a two-sampled

t-test, we used a Wilcoxon ranksum test instead. For tests related to saccade

directions, we used the CircStat toolbox [69]. When comparing proportions (for

example, for success rate), we used a $\chi^2$-test.

*Literature meta-analysis*

We searched for and read articles published between 1965 and November 2017. We

used the Web of Science search database, and we employed the keyword

"microsaccade*". We then selected, out of 467 research articles in the global results,

a total of 342 articles that were genuinely related to microsaccades and that were

also written in English. Any article that referred to microsaccades as being "voluntary"

(20/31 articles), "controlled" (6/31 articles), or "suppressed voluntarily" (5/31 articles)

was classified in the "voluntary" category (31 articles). Even though "suppressed

voluntary" is conceptually different from the voluntary control that we were interested

in for the current study, we nonetheless included this description in the voluntary

category to give this category the benefit of the doubt in our meta-analysis when

comparing to "involuntary" descriptions. In contrast to 31 articles, there were 148

articles in the "involuntary" category (almost 5 times as many). For the "involuntary"

category (148 articles), we included any article using the following descriptors for

microsaccades: "involuntary" (134/148 articles), "spontaneous" (7/148 articles),

"unaware" (3/148 articles), "automatic" (2/148 articles), "unconscious" (1/148

articles), "stochastic" (1/148 articles).

1013

1014    *Data availability*

1015    All data presented in this paper are stored in institute computers and are available

1016    upon reasonable request.

1017

1018

**Acknowledgements**

1019

1020 We were funded by the Werner Reichardt Centre for Integrative Neuroscience (CIN).

1021 The CIN is an Excellence Cluster (EXC307) funded by the Deutsche

1022 Forschungsgemeinschaft (DFG). We were also supported by the Hertie Institute for

1023 Clinical Brain Research at Tuebingen University, and DFG-funded Research Unit

1024 (FOR1847; project: HA6749/2-1).

1025

1026 **Author contributions**

1027 XT, AB, ZMH collected neurophysiological data. XT, JB, ZMH collected monkey

1028 behavioral data. KW, AR collected human behavioral data. KW, AB, ZMH analyzed

1029 data. ZMH wrote manuscript. KW, XT, AB, AR, ZMH edited manuscript.

1030

1031 **Declaration of interests**

1032 The authors declare no competing interests.

1033

1034

**References**

1035

1036

1037     1      Hafed, Z. M. & Clark, J. J. Microsaccades as an overt measure of covert
1038             attention shifts. *Vision Res* **42**, 2533-2545, doi:S0042698902002638 [pii]
1039             (2002).

1040     2      Engbert, R. & Kliegl, R. Microsaccades uncover the orientation of covert
1041             attention. *Vision Res* **43**, 1035-1045, doi:S0042698903000841 [pii] (2003).

1042     3      Tian, X., Yoshida, M. & Hafed, Z. M. A Microsaccadic Account of Attentional
1043             Capture and Inhibition of Return in Posner Cueing. *Frontiers in systems*
1044             *neuroscience* **10**, 23, doi:10.3389/fnsys.2016.00023 (2016).

1045     4      Hafed, Z. M., Chen, C.-Y. & Tian, X. Vision, perception, and attention through
1046             the lens of microsaccades: mechanisms and implications. *Frontiers in systems*
1047             *neuroscience* **9**, 167, doi:10.3389/fnsys.2015.00167 (2015).

1048     5      Tian, X., Yoshida, M. & Hafed, Z. M. Dynamics of fixational eye position and
1049             microsaccades during spatial cueing: the case of express microsaccades. *J*
1050             *Neurophysiol*, doi:10.1152/jn.00752.2017 (2018).

1051     6      Rolfs, M. Microsaccades: small steps on a long way. *Vision Res* **49**, 2415-
1052             2441, doi:S0042-6989(09)00369-1 [pii]
1053    10.1016/j.visres.2009.08.010 (2009).

1054     7      Hafed, Z. M., Lovejoy, L. P. & Krauzlis, R. J. Modulation of microsaccades in
1055             monkey during a covert visual attention task. *Journal of Neuroscience* **31**,
1056             15219-15230, doi:10.1523/JNEUROSCI.3106-11.2011 (2011).

1057     8      Pastukhov, A. & Braun, J. Rare but precious: microsaccades are highly
1058             informative about attentional allocation. *Vision Res* **50**, 1173-1184, doi:S0042-
1059             6989(10)00163-X [pii]
1060    10.1016/j.visres.2010.04.007 (2010).

1061     9      Engbert, R. Microsaccades: A microcosm for research on oculomotor control,
1062             attention, and visual perception. *Prog Brain Res* **154**, 177-192, doi:S0079-
1063             6123(06)54009-9 [pii]
1064    10.1016/S0079-6123(06)54009-9 (2006).

1065    10      Laubrock, J., Engbert, R. & Kliegl, R. Microsaccade dynamics during covert
1066             attention. *Vision Res* **45**, 721-730, doi:S0042-6989(04)00495-X [pii]
10.1016/j.visres.2004.09.029 (2005).

11    Laubrock, J., Engbert, R. & Kliegl, R. Fixational eye movements predict the
perceived direction of ambiguous apparent motion. *J Vis* **8**, 13 11-17,
doi:10.1167/8.14.13
/8/14/13/ [pii] (2008).

12    Kliegl, R., Rolfs, M., Laubrock, J. & Engbert, R. Microsaccadic modulation of
response times in spatial attention tasks. *Psychol Res* **73**, 136-146,
doi:10.1007/s00426-008-0202-2 (2009).

13    White, A. L. & Rolfs, M. Oculomotor inhibition covaries with conscious
detection. *J Neurophysiol* **116**, 1507-1521, doi:10.1152/jn.00268.2016 (2016).

14    Hicheur, H., Zozor, S., Campagne, A. & Chauvin, A. Microsaccades are
modulated by both attentional demands of a visual discrimination task and
background noise. *J Vis* **13**, 18, doi:10.1167/13.13.18 (2013).

15    Watanabe, M., Matsuo, Y., Zha, L., MacAskill, M. R. & Kobayashi, Y.
Fixational saccades alter the gap effect. *Eur J Neurosci* **39**, 2098-2106,
doi:10.1111/ejn.12566 (2014).

16    Wang, C. A., Blohm, G., Huang, J., Boehnke, S. E. & Munoz, D. P.
Multisensory integration in orienting behavior: Pupil size, microsaccades, and
saccades. *Biological psychology* **129**, 36-44,
doi:10.1016/j.biopsycho.2017.07.024 (2017).

17    Watanabe, M., Matsuo, Y., Zha, L., Munoz, D. P. & Kobayashi, Y. Fixational
saccades reflect volitional action preparation. *J Neurophysiol* **110**, 522-535,
doi:10.1152/jn.01096.2012 (2013).

18    Yuval-Greenberg, S., Tomer, O., Keren, A. S., Nelken, I. & Deouell, L. Y.
Transient induced gamma-band response in EEG as a manifestation of
miniature saccades. *Neuron* **58**, 429-441, doi:10.1016/j.neuron.2008.03.027
(2008).

19    Hafed, Z. M., Goffart, L. & Krauzlis, R. J. A neural mechanism for
microsaccade generation in the primate superior colliculus. *Science* **323**, 940-
943, doi:323/5916/940 [pii]
10.1126/science.1166112 (2009).

20    Hafed, Z. M. & Krauzlis, R. J. Similarity of superior colliculus involvement in
microsaccade and saccade generation. *J Neurophysiol* **107**, 1904-1916,
doi:10.1152/jn.01125.2011 (2012).

21      Hafed, Z. M. Mechanisms for generating and compensating for the smallest possible saccades. *Eur J Neurosci* **33**, 2101-2113, doi:10.1111/j.1460-9568.2011.07694.x (2011).

22      Hafed, Z. M., Lovejoy, L. P. & Krauzlis, R. J. Superior colliculus inactivation alters the relationship between covert visual attention and microsaccades. *Eur J Neurosci* **37**, 1169-1181, doi:10.1111/ejn.12127 (2013).

23      Peel, T. R., Hafed, Z. M., Dash, S., Lomber, S. G. & Corneil, B. D. A Causal Role for the Cortical Frontal Eye Fields in Microsaccade Deployment. *PLoS Biol* **14**, e1002531, doi:10.1371/journal.pbio.1002531 (2016).

24      Krauzlis, R. J. The control of voluntary eye movements: new perspectives. *Neuroscientist* **11**, 124-137, doi:11/2/124 [pii]
10.1177/1073858404271196 (2005).

Steinman, R. M., Cunitz, R. J., Timberlake, G. T. & Herman, M. Voluntary control of microsaccades during maintained monocular fixation. *Science* **155**, 1577-1579 (1967).

Haddad, G. M. & Steinman, R. M. The smallest voluntary saccade: implications for fixation. *Vision Res* **13**, 1075-1086 (1973).

Hafed, Z. M. Alteration of visual perception prior to microsaccades. *Neuron* **77**, 775-786, doi:10.1016/j.neuron.2012.12.014 (2013).

Chen, C. Y., Ignashchenkova, A., Thier, P. & Hafed, Z. M. Neuronal Response Gain Enhancement prior to Microsaccades. *Curr Biol* **25**, 2065-2074, doi:10.1016/j.cub.2015.06.022 (2015).

Tian, X., Yoshida, M. & Hafed, Z. M. Dynamics of fixational eye position and microsaccades during spatial cueing: the case of express microsaccades. *J Neurophysiol* **119**, 1962-1980, doi:10.1152/jn.00752.2017 (2018).

Bellet, J., Chen, C. Y. & Hafed, Z. M. Sequential hemifield gating of alpha and beta behavioral performance oscillations after microsaccades. *J Neurophysiol*, jn 00253 02017, doi:10.1152/jn.00253.2017 (2017).

Chen, C. Y. & Hafed, Z. M. A Neural Locus for Spatial-Frequency Specific Saccadic Suppression in Visual-Motor Neurons of the Primate Superior Colliculus. *J Neurophysiol*, jn 00911 02016, doi:10.1152/jn.00911.2016 (2017).

32   Hafed, Z. M. & Krauzlis, R. J. Microsaccadic suppression of visual bursts in
the primate superior colliculus. *J Neurosci* **30**, 9542-9547, doi:30/28/9542 [pii]
10.1523/JNEUROSCI.1137-10.2010 (2010).

33   Funahashi, S., Bruce, C. J. & Goldman-Rakic, P. S. Mnemonic coding of
visual space in the monkey's dorsolateral prefrontal cortex. *J Neurophysiol* **61**,
331-349, doi:10.1152/jn.1989.61.2.331 (1989).

34   Hikosaka, O. & Wurtz, R. H. Visual and oculomotor functions of monkey
substantia nigra pars reticulata. III. Memory-contingent visual and saccade
responses. *J Neurophysiol* **49**, 1268-1284 (1983).

35   Martinez-Conde, S., Otero-Millan, J. & Macknik, S. L. The impact of
microsaccades on vision: towards a unified theory of saccadic function. *Nat*
*Rev Neurosci* **14**, 83-96, doi:10.1038/nrn3405 (2013).

36   Wyman, D. & Steinman, R. M. Letter: Latency characteristics of small
saccades. *Vision Res* **13**, 2173-2175 (1973).

37   Kalesnykas, R. P. & Hallett, P. E. Retinal eccentricity and the latency of eye
saccades. *Vision Res* **34**, 517-531 (1994).

38   Rolfs, M., Kliegl, R. & Engbert, R. Toward a model of microsaccade
generation: the case of microsaccadic inhibition. *J Vis* **8**, 5 1-23,
doi:10.1167/8.11.5
/8/11/5/ [pii] (2008).

39   Hafed, Z. M. & Ignashchenkova, A. On the dissociation between
microsaccade rate and direction after peripheral cues: microsaccadic inhibition
revisited. *J Neurosci* **33**, 16220-16235, doi:10.1523/JNEUROSCI.2240-
13.2013 (2013).

40   Buonocore, A. *et al.* Alteration of the microsaccadic velocity-amplitude main
sequence relationship after visual transients: implications for models of
saccade control. *J Neurophysiol*, jn 00811 02016, doi:10.1152/jn.00811.2016
(2017).

41   Ko, H. K., Poletti, M. & Rucci, M. Microsaccades precisely relocate gaze in a
high visual acuity task. *Nat Neurosci* **13**, 1549-1553, doi:nn.2663 [pii]
10.1038/nn.2663 (2010).

42   Mohler, C. W. & Wurtz, R. H. Organization of monkey superior colliculus:
intermediate layer cells discharging before eye movements. *J Neurophysiol*
**39**, 722-744 (1976).

43    Edelman, J. A. & Goldberg, M. E. Dependence of saccade-related activity in
the primate superior colliculus on visual target presence. *J Neurophysiol* **86**,
676-691, doi:10.1152/jn.2001.86.2.676 (2001).

44    Mays, L. E. & Sparks, D. L. Dissociation of visual and saccade-related
responses in superior colliculus neurons. *J Neurophysiol* **43**, 207-232 (1980).

45    Stanford, T. R. & Sparks, D. L. Systematic errors for saccades to remembered
targets: evidence for a dissociation between saccade metrics and activity in
the superior colliculus. *Vision Res* **34**, 93-106 (1994).

46    Wimmer, K., Nykamp, D. Q., Constantinidis, C. & Compte, A. Bump attractor
dynamics in prefrontal cortex explains behavioral precision in spatial working
memory. *Nat Neurosci* **17**, 431-439, doi:10.1038/nn.3645 (2014).

47    Inaba, N. & Kawano, K. Neurons in cortical area MST remap the memory
trace of visual motion across saccadic eye movements. *Proc Natl Acad Sci U
S A* **111**, 7825-7830, doi:10.1073/pnas.1401370111 (2014).

48    Semework, M., Steenrod, S. C. & Goldberg, M. E. A spatial memory signal
shows that the parietal cortex has access to a craniotopic representation of
space. *eLife* **7**, doi:10.7554/eLife.30762 (2018).

49    Moore, T. Shape representations and visual guidance of saccadic eye
movements. *Science* **285**, 1914-1917 (1999).

50    Chen, C. Y. & Hafed, Z. M. Orientation and Contrast Tuning Properties and
Temporal Flicker Fusion Characteristics of Primate Superior Colliculus
Neurons. *Front Neural Circuits* **12**, 58, doi:10.3389/fncir.2018.00058 (2018).

51    Chen, C. Y., Sonnenberg, L., Weller, S., Witschel, T. & Hafed, Z. M. Spatial
frequency sensitivity in macaque midbrain. *Nat Commun* **9**, 2852,
doi:10.1038/s41467-018-05302-5 (2018).

52    Hafed, Z. M. & Chen, C.-Y. Sharper, stronger, faster upper visual field
representation in primate superior colliculus. *Current Biology*
http://dx.doi.org/10.1016/j.cub.2016.04.059,
doi:http://dx.doi.org/10.1016/j.cub.2016.04.059 (2016).

53    Van Horn, M. R. & Cullen, K. E. Coding of microsaccades in three-dimensional
space by premotor saccadic neurons. *J Neurosci* **32**, 1974-1980,
doi:10.1523/JNEUROSCI.5054-11.2012 (2012).

54 Arnstein, D., Junker, M., Smilgin, A., Dicke, P. W. & Thier, P. Microsaccade
control signals in the cerebellum. *J Neurosci* **35**, 3403-3411,
doi:10.1523/JNEUROSCI.2458-14.2015 (2015).

55 Sheth, B. R. & Shimojo, S. Compression of space in visual memory. *Vision*
*Res* **41**, 329-341 (2001).

56 Kerzel, D. Memory for the position of stationary objects: disentangling foveal
bias and memory averaging. *Vision Res* **42**, 159-167 (2002).

57 van der Heijden, A. H., van der Geest, J. N., de Leeuw, F., Krikke, K. &
Musseler, J. Sources of position-perception error for small isolated targets.
*Psychol Res* **62**, 20-35 (1999).

58 Dow, B. M., Snyder, A. Z., Vautin, R. G. & Bauer, R. Magnification factor and
receptive field size in foveal striate cortex of the monkey. *Exp Brain Res* **44**,
213-228 (1981).

59 Grujic, N., Brehm, N., Gloge, C., Zhuo, W. & Hafed, Z. M. Peri-saccadic
perceptual mislocalization is different for upward saccades. *J Neurophysiol*,
doi:10.1152/jn.00350.2018 (2018).

60 Skinner, J., Buonocore, A. & Hafed, Z. M. The transfer function of the rhesus
macaque oculomotor system for small-amplitude slow motion trajectories. *J*
*Neurophysiol*, doi:10.1152/jn.00437.2018 (2018).

61 Judge, S. J., Richmond, B. J. & Chu, F. C. Implantation of magnetic search
coils for measurement of eye position: an improved method. *Vision Res* **20**,
535-538 (1980).

62 Fuchs, A. F. & Robinson, D. A. A method for measuring horizontal and vertical
eye movement chronically in the monkey. *J Appl Physiol* **21**, 1068-1070
(1966).

63 Chen, C. Y. & Hafed, Z. M. Postmicrosaccadic enhancement of slow eye
movements. *The Journal of neuroscience : the official journal of the Society for*
*Neuroscience* **33**, 5375-5386, doi:10.1523/JNEUROSCI.3703-12.2013 (2013).

64 Bellet, M. E., Bellet, J., Nienborg, H., Hafed, Z. M. & Berens, P. Human-level
saccade detection performance using deep neural networks. *J Neurophysiol*,
doi:10.1152/jn.00601.2018 (2018).

65 Herrington, T. M. *et al.* The effect of microsaccades on the correlation
between neural activity and behavior in middle temporal, ventral intraparietal,
and lateral intraparietal areas. *J Neurosci* **29**, 5793-5805, doi:29/18/5793 [pii]
10.1523/JNEUROSCI.4412-08.2009 (2009).

66 Rossant, C. *et al.* Spike sorting for large, dense electrode arrays. *Nat Neurosci*
**19**, 634-641, doi:10.1038/nn.4268 (2016).

67 Hill, D. N., Mehta, S. B. & Kleinfeld, D. Quality metrics to accompany spike
sorting of extracellular signals. *J Neurosci* **31**, 8699-8705,
doi:10.1523/JNEUROSCI.0971-11.2011 (2011).

68 Schmitzer-Torbert, N., Jackson, J., Henze, D., Harris, K. & Redish, A. D.
Quantitative measures of cluster quality for use in extracellular recordings.
*Neuroscience* **131**, 1-11, doi:10.1016/j.neuroscience.2004.09.066 (2005).

69 Berens, P. CircStat: a MATLAB toolbox for circular statistics. *Journal of*
*Statistical Software* **31**, 1-21 (2009).
1243

1244

1245

 **Supplementary information**

 **Supplementary Movie 1 Example trial showing the instantaneous eye position of**
 **a human subject generating a voluntary, memory-guided microsaccade towards**
 **an invisible foveal location.** The cyan cross depicts instantaneous eye position, and
 the green dashed circle around the central fixation point (white dot) is a circle of 1°
 radius. The scale bar on the bottom left describes visual dimensions of the displayed
 area, and the bottom right text indicates elapsed time from trial onset. The movie is
 running in slow motion to clarify the specific task events, which are highlighted by the
 text on the top left corner of each frame. The movie pauses briefly at target flash onset
 (normally a very brief event) in order to demonstrate the small eccentricity associated
 with the memory-guided microsaccade that is successfully generated later. The subject
 generated a spatially accurate microsaccade to the invisible target location after the
 fixation spot disappearance ("Go" command). Supplementary Movie 2 shows the same
 movie in real-time.

 **Supplementary Movie 2 Same as Supplementary Movie 1, but now running in**
 **real-time.** Despite the rapid sequence of events in the trial, the subject was
 successfully able to voluntarily generate a spatially accurate memory-guided
 microsaccade.

1267

1268

**Supplementary Figure 1 Overwhelming description of microsaccades as being**
**"involuntary" eye movements in the literature.** In the top panel, we performed a
meta-analysis of all published articles on microsaccades between 1965 and 2017. We
classified articles based on how they referred to microsaccades with respect to volition
(see Methods). We grouped the articles into 3 categories (Methods): explicitly referring
to microsaccades as being either "voluntary" (blue) or "involuntary" (red) eye
movements, or as making no explicit mention regarding volition (dark gray). Each time
slice (x-axis) shows the distribution of these 3 categories in published research (y-
axis). The bottom histogram shows the number of published articles in 2-year bins
within the same time period as in the top panel; it shows a pattern consistent with Rolfs'
meta-analysis [6], in which there was almost a complete gap of research on
microsaccades in the final two decades of the previous century. In practically all time
slices of the top panel, there was a majority of references to microsaccades as being
"involuntary" compared to "voluntary". In the current century (dashed vertical line and
rightward block arrows), there was a significant increase in microsaccade research
(bottom histogram) with a concomitant introduction of a large bias to refer to
microsaccades as being involuntary. The large fraction of "gray" papers (with no
explicit mention regarding volition) in the current century is attributed to the emergence
of new questions regarding microsaccades [21], such as their links with cognition.

1288

1289

**Reaction times**

1290

1291

**Supplementary Figure 2 Similarity of memory-guided microsaccade reaction
times to the reaction times of larger memory-guided saccades.** In each panel, we
show data similar in format to Fig. 2a, d, g, j. Here, we included large saccades, and
we also binned data according to memory-guided saccade/microsaccade size (see
color legend in the top panel). Memory-guided microsaccadic reaction times (the first
two colors in the color legend; movements with <60 min arc amplitudes) had similar
distributions to those of larger memory-guided saccades. Note that there were
quantitative differences in reaction times (particularly evident in the human subjects in
which we had larger numbers of trials overall) that reflect factors other than the intrinsic
ability to react to the instruction to generate an eye movement (for example, increases
in reaction times for smaller target eccentricities [37]).

1303

1304

**Supplementary Figure 3 Memory-guided microsaccades were more successful and directionally accurate than flash-induced microsaccades at the beginnings of trials.** For the same data as in Figs. 2, 3, we plotted reaction time (**a**, **d**, **g**, **j**), success rate (**b**, **e**, **h**, **k**), and direction error (**c**, **f**, **i**, **l**) for the first microsaccade to occur after target flash onset (flash-induced microsaccades). The monkeys exhibited express reactions on a minority of trials (**a**, **d**, **g**, **j**), similar to the corrective, visually-guided movements in Fig. 2m, p, s, but then barely any eye movement responses at all at the times in which the instructed memory-guided microsaccades would have happened (the light blue histograms show the memory-guided microsaccade reaction time distributions for easy comparison). Humans had no express reactions, but they again showed barely any responses to target flash onset in the later interval in which memory-guided microsaccades were triggered (also see Fig. 1). This means that the overall success rates after target flash onset were significantly lower than for the instructed memory-guided microsaccades (**b**, **e**, **h**, **k**). Therefore, the latter movements did not reflect transient reflexes, perhaps due to covert attentional shifts [1,2] associated with the task. In terms of direction errors, we plotted angular distributions (as in Fig. 3) for flash-induced microsaccades (**c**, **f**, **i**, **l**). In the left column of each panel, we plotted direction errors for express reactions (occurring 50-120 ms after target flash onset), and in the right column of each panel, we plotted the direction errors for reaction times 120-300 ms after target flash onset (similar to the range of reaction times for instructed memory-guided microsaccades). Express flash-induced microsaccades were directionally accurate, as expected [29]; the later responses were significantly less directionally accurate than instructed memory-guided microsaccades (shown in transparent light blue for easier comparison).

1333

1334

**Supplementary Figure 4 Delayed, visually-guided microsaccades had lower**
**success rates than memory-guided microsaccades, but they were spatially**
**accurate in both direction and amplitude, as expected. (a)** Success rate as in Fig.
2b for monkey N in the delayed, visually-guided saccade task. The shown data is for
all trials with target eccentricities <1°. The monkey often failed to generate the
instructed movement within the specified time, resulting in a plateau success rate of
only ~50% (significantly less than for instructed memory-guided microsaccades). This
could reflect an overall difficulty in withholding eye movement responses for prolonged
periods with the continuous presence of a visual target; the monkey often just fixated
the persistently present visual target before the go instruction. **(b)** However, the
successful microsaccades that were generated in the task were directionally accurate.
We plotted the distribution of angular direction errors between target locations and
evoked eye movements (as in Fig. 3). A distribution centered near 0 direction error
indicates directionally accurate eye movements. **(c)** The successful microsaccades
were also accurate in terms of their amplitude relative to target eccentricity. No
overshoot was observed as in the case of memory-guided microsaccades and
uninstructed corrective, visually-guided microsaccades (Fig. 4) (and also memory-
guided perceptual judgements; Fig. 5). **(d, e, f)** Similar observations for monkey M.
n=581 trials for **a**, **b**; n=2843 trials for **c**; n=314 trials for **d**, **e**; n=2476 trials for **f**.

1354

1355

**Memory-guided saccades**

**Delayed, visually-guided saccades**

Monkey M

**Supplementary Figure 5 The activity of the neuron of Fig. 7a, b in different tasks.** The left panel shows the neuron's activity from the memory-guided saccade task, including for large saccades, and the right panel shows the neuron's activity from the delayed, visually-guided saccade task (and with a similar range of tested saccade directions and amplitudes). All formatting conventions are similar to Figs. 6, 7. During the memory-guided saccade task (left), the neuron's movement RF expanded foveally, allowing the neuron to respond exclusively for memory-guided microsaccades but not for similarly sized visually-guided microsaccades (Fig. 7a, b). The insets show individual spike waveforms from the neuron for every tenth spike in each task (light gray), with the red waveforms being the averages (of ~700 spikes each). We ensured that the neuron met criteria for being sorted as the same neuron in the two tasks (e.g. inter-spike intervals and waveform shape criteria; Methods). Each inset shows ~70 waveforms (gray curves) per task.