Androgens enhance adult hippocampal neurogenesis in males but not females in an age-

dependent manner

Paula Duarte-Guterman[1], Dwayne K. Hamson[1], Steven R. Wainwright[1], Carmen Chow[1], Jessica

Chaiton[1], Stephanie Lieblich[1], Neil V. Watson[2], Liisa A.M. Galea[1]

1. Djavad Mowafaghian Centre for Brain Health and Department of Psychology, University of

British Columbia, Vancouver, BC, Canada

2. Department of Psychology, Simon Fraser University, Burnaby, BC, Canada

Address all correspondence and requests for reprints to:

L. A. M. Galea, PhD,

Djavad Mowafaghian Centre for Brain Health

2215 Wesbrook Mall

Vancouver, British Columbia

V6T 1Z3, Canada

2136 West Mall, Vancouver,

British Columbia, V6T 1Z4, Canada.

E-mail: lgalea@psych.ubc.ca.

Abstract

Androgens (testosterone and dihydrotestosterone) increase adult hippocampal

neurogenesis by increasing new neuron survival in male rats and mice via an androgen receptor

pathway, but it is not known whether androgens regulate neurogenesis in females and whether

the effect is age-dependent. We investigated the effects of dihydrotestosterone, a potent

androgen, on neurogenesis in adult and middle-aged males and females. Rats were

gonadectomized and injected with the DNA synthesis marker, bromodeoxyuridine (BrdU). The

following day rats began receiving daily injections of oil or DHT for 30 days. We evaluated cell

proliferation (Ki67) and new neuron survival (BrdU and BrdU/NeuN) in the hippocampus of

male and female rats using immunohistochemistry. As expected, DHT increased new neuron

survival in young males but surprisingly not in middle-aged male rats. In females, DHT did not

significantly affect adult neurogenesis in young or middle age. Our results indicate that DHT

regulates adult hippocampal neurogenesis in a sex- and age-dependent manner.

Main text

Neurogenesis, the production of new neurons, in the hippocampus continues through the

life span of most mammals studied to date (Kempermann et al., 2018). Sex hormones (estrogens

and androgens) regulate different aspects of hippocampal neurogenesis: e.g. proliferation and/or

survival of these new neurons in rodents (reviewed in Duarte-Guterman et al., 2015b; Mahmoud

et al., 2016). There is also evidence of sex differences in how hormones regulate neurogenesis.

For example, estradiol regulates new neuron survival and cell proliferation in female but not

male rodents (Barker and Galea, 2008). We have previously shown that androgens (testosterone

and dihydrotestosterone) increase the survival of new neurons but not cell proliferation in the

hippocampus of male rats via an androgen receptor (AR) pathway (Spritzer and Galea, 2007;

Hamson et al., 2013) and mice (Swift-Gallant et al., 2018). However it is not known whether

androgens regulate any aspects of adult neurogenesis in females. In addition to sex, age can also

modulate the effects of hormones on hippocampal neurogenesis. In middle age, the hippocampus

loses its ability to respond to estrogens in female rats. For instance, estradiol increases cell

proliferation in the hippocampus in young but not middle-aged nulliparous female rats (Chiba et

al., 2007; Barha and Galea, 2011). The objective of this study was to investigate the effects of

dihydrotestosterone (DHT) on hippocampal neurogenesis (proliferation and new neuron

survival) in young and middle-aged male and female rats.

At 2 months (~70 days old, young) and 11-12 months of age (middle-aged),

gonadectomized male and female Sprague–Dawley rats received a single intraperitoneal

injection of bromodeoxyuridine (BrdU; 200mg/kg) to label dividing cells and their progeny

(Hamson et al., 2013). The following day, males and females were injected subcutaneously with

either 0.25mg dihydrotestosterone (DHT in 0.1ml of sesame oil) or an equivalent volume of

sesame oil for 30 days. Twenty-four hours after the final injection, animals were overdosed with

sodium pentobartitol and brains perfused and collected for BrdU (survival of 30 day old cells),

Ki67 (cell proliferation marker), androgen receptor (AR), and BrdU/NeuN (new neurons using

NeuN, marker for mature neurons) immunohistochemistry (Hamson et al., 2013; Swift-Gallant et

al., 2018). Thus, in this experiment, BrdU+ cells were 30 day-old daughter cells from progenitor

cells that had been synthesizing DNA for a 2-hour period 31 days before euthanasia. All

protocols were approved by the Animal Care Committee at the University of British Columbia

and conformed to the guidelines set out by the Canadian Council for Animal Care.

A researcher blind to experimental conditions counted BrdU+ and Ki67-immunoreactive

(ir) cells in the entire rostrocaudal extent of the granule cell layer (GCL) including the sub

granular zone, defined as the 50 µm band between the GCL and the hilus. An estimate of the

total number of cells in the GCL and hilus was calculated by multiplying the number of cells

counted by 10 to account for the fact that we used 1/10 series of sections (Tanapat et al., 2005;

Spritzer and Galea, 2007; Swift-Gallant et al., 2018). GCL and hilus volumes were quantified

from digitized images using Cavelier's principle, multiplying the sum of the area of each section

by the section thickness (40 µm Gundersen et al., 1988). Androgen receptor immunoreactivity

was characterized using a relative scale: robust (+++), intermediate (++), light (+), or absent (0)

in the GCL, CA1, and CA3 regions in DHT and oil treated groups as we have done before

(Hamson et al., 2013). To determine whether BrdU+ cells were of a neuronal phenotype, in a

subset of brains, all BrdU+ cells within a section were examined for co-labeling with NeuN

(neuronal marker) for 50 cells (young adult group) or all cells (middle-aged group) were

phenotyped. All analyses were performed using Statistica v.8.0 (StatSoft Inc, Tulsa, OK).

Volume of the dentate gyrus (GCL and hilus) and density of BrdU+ cells (total cells per unit

volume) were analyzed using repeated measures analysis of variance (ANOVA) with age

(young, middle-aged), sex (male, female), and treatment (DHT, oil) as between subjects factors,

and region (GCL, hilus) as a within subjects factor. Density of Ki67-ir cells and proportion of

BrdU/NeuN co-labeled cells were analyzed using ANOVA with age (young, middle-aged), sex

(male, female), and treatment (DHT, oil) as between subjects factors. When appropriate, post-

hoc analysis used the Neuman-Keul's procedure. Test statistics were considered significant if $p \leq$

0.05.

Regardless of age and treatment, the GCL and hilus were larger in males compared to

females (main effect of sex; $F(1,42)=4.42$; $P<0.05$; Table 1). The volume of the GCL and hilus

increased with age irrespective of sex and treatment (main effect of age; $F(1,42)=47.37$;

$P<0.0001$). In the case of the hilus, the volume increased with aging more so in males than in

females (interaction between region, age, and sex; $F(1,42)=20.20$; $P<0.0001$). As expected, the

volume of the hilus was larger than the volume of the GCL (main effect of region;

$F(1,44)=1450.14$; $P<0.0001$). Treatment with DHT did not significantly affect GCL or hilus

volumes (all P's>0.08). To account for the sex differences in GCL and hilus volumes, we present

BrdU and Ki67 counts as densities (total cells per unit volume).

DHT treatment increased the density of BrdU+ cells in the GCL in young males

($P<0.001$) but not in young females or middle-aged animals of both sexes (all P's>0.9;

interaction between age, sex, treatment and region; $F(1,42)=4.03$; $P=0.05$; Figure 1). The density

of BrdU+ cells in the GCL was significantly higher in the young compared to the middle-aged

rats irrespective of sex and treatment, as expected (main effect of age; $F(1,42)=647.85$;

$P<0.0001$). In the hilus, the density of BrdU+ cells was not significantly affected by age, sex or

treatment (all P's>0.9). To determine how many BrdU+ cells were neurons, we examined the

colabelling of BrdU and NeuN (a neuronal marker; BrdU/NeuN) in the GCL. Aging decreased

the proportion of BrdU/NeuN colabeled cells in the dentate gyrus (main effects of age;

$F_{(1,39)}=27.98$; $P<0.0001$). In the young group, sex and treatment did not affect the percent of

BrdU/NeuN colabeled cells (all P's>0.18; Table 2). In middle age, DHT treatment increases the

proportion of BrdU/NeuN cells irrespective of sex (interaction between age and treatment;

$F_{(1,39)}=4.77$; $P=0.03$).

DHT treatment did not affect the density of Ki67-ir cells in young or middle age males

and females (all P's>0.25; Figure 1). However, as expected, there were more Ki67-ir cells in the

GCL of the young compared to middle-aged rats (main effect of age; $F_{(1,43)}=97.18$; $P<0.0001$)

irrespective of sex and treatment.

Finally, we performed a qualitative analysis on AR expression in the hippocampus. In

gonadectomized oil treated animals, AR-ir cells were absent throughout the GCL in young and

middle-aged male and female rats. In the CA1, we found low to intermediate levels of AR-ir and

in the CA3 low to absent levels of AR-ir in young and middle-aged rats of both sexes. DHT

treatment increased the expression of AR in the CA1 to intermediate or robust levels in both

sexes (Table 3; Figure 1).

In the present study, we found that chronic (30 days) DHT increased new neuron survival

but not cell proliferation in the hippocampus of gonadectomized young adult male rats,

consistent with our previous research in male rats and mice (Spritzer and Galea, 2007; Hamson

et al., 2013; Swift-Gallant et al., 2018). This is also in line with previous work showing that

castration decreases new neuron survival 24-30 days after BrdU injection but has no effect on

cell proliferation (Spritzer and Galea, 2007; Wainwright et al., 2011). Shorter testosterone

treatment (3, 15 or 21 days) has no effect on hippocampal neurogenesis (Zhang et al., 2010;

Spritzer et al., 2011; Carrier and Kabbaj, 2012; Wainwright et al., 2016) indicating that a longer

exposure (30 days) to androgens is required to increase neurogenesis in the hippocampus at least

in physiological doses as higher doses can decrease neurogenesis (Brännvall et al., 2005; Zhang

et al., 2014). Thus, collectively while longer term exposure to androgens increases new neuron

survival in the dentate gyrus, androgens do not appear to influence cell proliferation in male rats

(this study; Spritzer and Galea, 2007; Hamson et al., 2013), mice (Swift-Gallant et al., 2018), or

voles (Fowler et al., 2003).

In contrast to young adult males, DHT did not affect new neuron survival or cell

proliferation in gonadectomized young adult female rats. We previously found that estradiol

modulates cell proliferation and new neuron survival in young adult female rats (Ormerod et al.,

2003; Mazzucco et al., 2006; Barker and Galea, 2008; Barha et al., 2009; Duarte-Guterman et al.,

2015a) but has no effect on neurogenesis in adult male rats (Spritzer and Galea, 2007; Barker

and Galea, 2008). Together this suggests that sex steroids have sex-specific effects on

hippocampal neurogenesis with androgens modulating neurogenesis in young adult males and

estrogens modulating neurogenesis in young adult females. While these findings may not seem

surprising, it is important to understand that both sexes have ARs and estrogen receptors (ERs)

but these receptors are responding to respective hormones in a sex-specific way to modulate

neurogenesis in the hippocampus.

Perhaps surprisingly, the effect of DHT to modulate survival of new neurons was absent

in middle-aged males and females. This effect is consistent with the work of Moser et al. (2019)

who found that testosterone did not affect the number of immature neurons (using doublecortin),

in middle-aged (13 months) or aged (23 months) male rats. This would suggest that with aging,

the dentate gyrus loses its ability to respond to sex steroids. Indeed, in females, estradiol

increases cell proliferation in young but not middle-aged rats (Barha et al., 2009; Barha and

Galea, 2011). Intriguingly in females, previous reproductive experience (pregnancy and

motherhood) can rescue the hippocampus' response to estrogens later in middle-age, as acute

estrogens increased cell proliferation in multiparous rats (Barha and Galea, 2011). Thus, it is

possible that experience, reproductive or otherwise, may restore the ability of androgens to

upregulate hippocampal neurogenesis in males in middle age. However, Moser et al. (2019) did

not see an influence of high fat diet on the androgen modulation of neurogenesis. Intriguingly in

our study, the proportion of BrdU/NeuN colabelled cells was affected by age and treatment.

DHT increased the proportion of BrdU/NeuN colabelled cells, an effect mostly driven by the

middle-aged females, and overall as expected, the proportion of new cells that express mature

neuronal protein decreased with age. In middle-aged female mice letrozole, an aromatase

inhibitor blocking the conversion of androgens to estrogens, increases hippocampal neurogenesis

(using the immature marker doublecortin; Chaiton et al., 2018). Together with our findings, this

suggests that the hippocampus may still be able to respond to sex steroid hormones in middle

age, an effect that varies by sex as in both studies middle-aged females showed increased

neurogenesis levels with androgens.

We found that ARs were expressed in the CA1 and CA3 regions of the hippocampus. We

did not find expression of ARs in the granule cell layer in male rats and this is consistent with

previous research (Kerr et al., 1995; Xiao and Jordan, 2002; Hamson et al., 2013); although there

is conflicting evidence which may be due to strain and species differences. In Wistar male rats,

Moghadami et al. (2016) and Brännvall et al. (2005) found AR expression in the granule cell

layer in young male rats. In gonadectomized male mice, ARs are not expressed in the granule

cell layer but in males receiving DHT treatment, granule cells express AR (Swift-Gallant et al.,

218). In other rat strains (Bruce-Spruce Long-Evans, Fischer 344 and Sprague Dawley), ARs are

not expressed in the granule cell layer (Xiao and Jordan 2002; Kerr et al., 1995; Hamson et al.,

2013). We also found that DHT increases the expression of ARs in the CA1 in males in line with

previous research in young male rats (Hamson et al., 2013) and male mice (Swift-Gallant et al.,

2018). In females and males, 2 days of testosterone treatment also showed increased AR

expression in the CA1 region (Xiao and Jordan, 2002) but to our knowledge no other studies

have investigated AR expression after chronic DHT treatment in females. Previous work has

found that androgens increase neurogenesis via the androgen receptor (AR). Blocking the AR

with flutamide abolishes the increase in new neuron survival in male rats (Hamson et al., 2013).

However, in male rats and mice, ARs are not found in immature neurons (doublecortin

expressing cells) in the granule cell layer of the hippocampus (Hamson et al., 2013; Swift-

Gallant et al., 2018). We have previously proposed that one mechanism of action is that

androgens bind to ARs in the CA3 region and this initiates a retrograde response of a survival

factor that targets newborn neurons in the granule cell layer (Galea et al., 2013). In the current

study, we found ARs were expressed in the CA1 and CA3 regions in young and middle-aged

male and female rats and their expression increases after DHT treatment in both sexes at both

ages. However, we only observed an effect of DHT on new neuron survival in young males. In

young females and middle-aged males and females, ARs are expressed in the CA1 and CA3 but

binding of DHT to these ARs does not result in the modulation of neurogenesis, possibly

indicating different downstream mechanisms of bound AR to modulate neurogenesis in young

adult females, and middle-aged males. Interestingly, we found higher expression of AR in the

CA1 of middle-aged compared to young rats of both sexes. AR mRNA expression increases in

the hippocampus with aging in males (Kerr et al., 1995). DHT did not increase new neuron

survival in mice overexpressing AR in the CA1, indicating that high doses of AR result in an

abolished response (Swift-Gallant et al., 2018). It is possible that the increase in AR expression

with aging is responsible for the lack of DHT mediated increase in new neuron survival.

To summarize, DHT increased survival of new neurons in young adult male rats but not

in female or middle-aged male and female rats. However, DHT treatment increased the

proportion of surviving cells expressing the mature neuronal protein in middle-aged females and

AR expression in both sexes, regardless of age, suggesting that ARs have the potential to

respond to DHT. Our current work together with our previous research (reviewed in Mahmoud et

al., 2016) indicate that androgens and estrogens have sex-specific and age-specific effects on

hippocampal neurogenesis.

Acknowledgments

This work was funded by the Canadian Institutes of Health Research (MOP102568) to

LAMG. We thank Kim Go for assistance with research and Dr. Mark Martindale (Whitney

Laboratory, University of Florida) for access to an epifluorescent microscope.

References

Barha CK, Galea LAM. 2011. Motherhood alters the cellular response to estrogens in the
hippocampus later in life. Neurobiology of Aging 32:2091–2095.

Barha CK, Lieblich SE, Galea LAM. 2009. Different Forms of Oestrogen Rapidly Upregulate
Cell Proliferation in the Dentate Gyrus of Adult Female Rats. Journal of
Neuroendocrinology 21:155–166.

Barker JM, Galea LAM. 2008. Repeated estradiol administration alters different aspects of
neurogenesis and cell death in the hippocampus of female, but not male, rats.
Neuroscience 152:888–902.

Brännvall K, Bogdanovic N, Korhonen L, Lindholm D. 2005. 19-Nortestosterone influences
neural stem cell proliferation and neurogenesis in the rat brain. European Journal of
Neuroscience 21:871–878.

Carrier N, Kabbaj M. 2012. Testosterone and imipramine have antidepressant effects in socially
isolated male but not female rats. Hormones and Behavior 61:678–685.

Chaiton J, Wong S, Galea L. 2018. Letrozole increases hippocampal neurogenesis in middle-
aged female mice. bioRxiv:434662.

Chiba S, Suzuki M, Yamanouchi K, Nishihara M. 2007. Involvement of Granulin in Estrogen-
Induced Neurogenesis in the Adult Rat Hippocampus. Journal of Reproduction and
Development 53:297–307.

Duarte-Guterman P, Lieblich SE, Chow C, Galea LAM. 2015a. Estradiol and GPER Activation
Differentially Affect Cell Proliferation but Not GPER Expression in the Hippocampus of
Adult Female Rats. PLoS ONE 10:e0129880.

Duarte-Guterman P, Yagi S, Chow C, Galea LAM. 2015b. Hippocampal learning, memory, and
neurogenesis: Effects of sex and estrogens across the lifespan in adults. Horm Behav
74:37–52.

Fowler CD, Freeman ME, Wang Z. 2003. Newly proliferated cells in the adult male amygdala
are affected by gonadal steroid hormones. J Neurobiol 57:257–269.

Galea LAM, Wainwright SR, Roes MM, Duarte-Guterman P, Chow C, Hamson DK. 2013. Sex,
Hormones and Neurogenesis in the Hippocampus: Hormonal Modulation of
Neurogenesis and Potential Functional Implications. Journal of Neuroendocrinology
25:1039–1061.

Gundersen HJ, Bendtsen TF, Korbo L, Marcussen N, Møller A, Nielsen K, Nyengaard JR,
Pakkenberg B, Sørensen FB, Vesterby A. 1988. Some new, simple and efficient
stereological methods and their use in pathological research and diagnosis. APMIS
96:379–394.

Hamson DK, Wainwright SR, Taylor JR, Jones BA, Watson NV, Galea LAM. 2013. Androgens
Increase Survival of Adult-Born Neurons in the Dentate Gyrus by an Androgen Receptor-
Dependent Mechanism in Male Rats. Endocrinology 154:3294–3304.

Kempermann G, Gage FH, Aigner L, Song H, Curtis MA, Thuret S, Kuhn HG, Jessberger S,
Frankland PW, Cameron HA, Gould E, Hen R, Abrous DN, Toni N, Schinder AF, Zhao
X, Lucassen PJ, Frisén J. 2018. Human Adult Neurogenesis: Evidence and Remaining
Questions. Cell Stem Cell 23:25–30.

Kerr JE, Allore RJ, Beck SG, Handa RJ. 1995. Distribution and hormonal regulation of androgen
receptor (AR) and AR messenger ribonucleic acid in the rat hippocampus. Endocrinology
136:3213–3221.

Mahmoud R, Wainwright SR, Galea LAM. 2016. Sex hormones and adult hippocampal
neurogenesis: Regulation, implications, and potential mechanisms. Frontiers in
Neuroendocrinology 41:129–152.

Mazzucco CA, Lieblich SE, Bingham BI, Williamson MA, Viau V, Galea LAM. 2006. Both
estrogen receptor α and estrogen receptor β agonists enhance cell proliferation in the
dentate gyrus of adult female rats. Neuroscience 141:1793–1800.

Moghadami S, Jahanshahi M, Sepehri H, Amini H. 2016. Gonadectomy reduces the density of
androgen receptor-immunoreactive neurons in male rat's hippocampus: testosterone
replacement compensates it. Behavioral and Brain Functions [Internet] 12. Available
from: http://behavioralandbrainfunctions.biomedcentral.com/articles/10.1186/s12993-
016-0089-9

Moser VA, Christensen A, Liu J, Zhou A, Yagi S, Beam CR, Galea L, Pike CJ. 2019. Effects of
aging, high-fat diet, and testosterone treatment on neural and metabolic outcomes in male
brown Norway rats. Neurobiol Aging 73:145–160.

Ormerod BK, Lee TT-Y, Galea LAM. 2003. Estradiol initially enhances but subsequently
suppresses (via adrenal steroids) granule cell proliferation in the dentate gyrus of adult
female rats. Journal of Neurobiology 55:247–260.

Spritzer MD, Galea LAM. 2007. Testosterone and dihydrotestosterone, but not estradiol,
enhance survival of new hippocampal neurons in adult male rats. Developmental
Neurobiology 67:1321–1333.

Spritzer MD, Ibler E, Inglis W, Curtis MG. 2011. Testosterone and social isolation influence
adult neurogenesis in the dentate gyrus of male rats. Neuroscience 195:180–190.

Swift-Gallant A, Duarte-Guterman P, Hamson DK, Ibrahim M, Monks DA, Galea LAM. 2018.
Neural androgen receptors affect the number of surviving new neurones in the adult
dentate gyrus of male mice. Journal of Neuroendocrinology 30:e12578.

Tanapat P, Hastings NB, Gould E. 2005. Ovarian steroids influence cell proliferation in the
dentate gyrus of the adult female rat in a dose- and time-dependent manner. J Comp
Neurol 481:252–265.

Wainwright SR, Lieblich SE, Galea LAM. 2011. Hypogonadism predisposes males to the
development of behavioural and neuroplastic depressive phenotypes.
Psychoneuroendocrinology 36:1327–1341.

Wainwright SR, Workman JL, Tehrani A, Hamson DK, Chow C, Lieblich SE, Galea LAM.
2016. Testosterone has antidepressant-like efficacy and facilitates imipramine-induced
neuroplasticity in male rats exposed to chronic unpredictable stress. Hormones and
Behavior 79:58–69.

Xiao L, Jordan CL. 2002. Sex Differences, Laterality, and Hormonal Regulation of Androgen
Receptor Immunoreactivity in Rat Hippocampus. Hormones and Behavior 42:327–336.

Zhang W, Cheng J, Vagnerova K, Ivashkova Y, Young J, Cornea A, Grafe MR, Murphy SJ,
Hurn PD, Brambrink AM. 2014. Effects of androgens on early post-ischemic
neurogenesis in mice. Transl Stroke Res 5:301–311.

Zhang Z, Yang R, Zhou R, Li L, Sokabe M, Chen L. 2010. Progesterone promotes the survival
of newborn neurons in the dentate gyrus of adult male mice. Hippocampus 20:402–412.

 FIGURES

Figure 1. Chronic dihydrotestosterone (DHT) treatment increases the density of BrdU+ cells in

the dentate gyrus of young male adult rats but has no effects in middle-aged males and young

and middle-aged female rats. (**A**) Photomicrographs of a representative section of dentate gyrus

with the granule cell layer (GCL), hilus, CA1 and CA3 regions and representative BrdU+ and

Ki67-ir cells in the subgranular zone. (**B**) Representative images of androgen receptor (AR)

expression in the CA1 region of the hippocampus in male and female rats treated with oil or

DHT. AR is lowly expressed in the CA1 of gonadectomized oil treated animals. DHT increases

the expression of AR in both male and female rats. (**C**) Mean ± SEM total number of BrdU+

cells in the granule cell layer of the dentate gyrus in young and middle-aged male and female

rats. In young males, DHT increased the density of BrdU+ cells relative to the oil treatment

group ($P < 0.001$). (**D**) Mean ± SEM total number of Ki67-ir cells in the granule cell layer of the

dentate gyrus in young and middle-aged male and female rats. Chronic DHT treatment did not

affect cell proliferation in the hippocampus of young or middle-aged male and female rats.

Circles represent individual data points.

TABLES

Table 1. Mean (SEM) volume of the granule cell layer (GCL) and the hilus (mm^3) and BrdU

density in the hilus in gonadectomized young and middle-aged male and female rats treated with

oil or dihydrotestosterone (DHT). Males had a larger volume of GCL and hilus and the volume

of the hilus increased with aging more so in males than females. DHT treatment did not have an

effect on the volume of the GCL and hilus.

Age	Sex and Treatment	Sample size	GCL volume (mm^3)	Hilus volume (mm^3)	Hilus BrdU density (per mm^3)
Young	*Females*				
	Oil	7	2.51 (0.12)	6.09 (0.19)	85.14 (15.34)
	DHT	8	2.46 (0.08)	6.00 (0.28)	76.06 (7.06)
	Males				
	Oil	5	2.31 (0.19)	5.32 (0.49)	96.18 (23.29)
	DHT	6	2.62 (0.21)	5.37 (0.51)	134.70 (17.38)
Middle Age	*Females*				
	Oil	6	2.33 (0.09)	7.67 (0.42)	20.69 (7.89)
	DHT	6	2.45 (0.15)	7.33 (0.28)	6.40 (2.49)
	Males				
	Oil	6	2.94 (0.25)	9.05 (0.79)	6.06 (1.41)
	DHT	6	2.87 (0.22)	10.05 (0.60)	4.65 (1.63)

322 **Table 2.** Mean (SEM) percentage of cells co-expressing BrdU and NeuN in the GCL in

323 gonadectomized young and middle-aged male and female rats treated with oil or

324 dihydrotestosterone (DHT). The proportion of BrdU+ cells colabelled with NeuN was

325 significantly higher in young compared to middle-aged animals. In middle age, DHT increases

326 the proportion of BrdU/NeuN colabelled cells relative to the oil treatment group.

Age	Sex and Treatment	Sample size	BrdU/NeuN %
Young	*Females*		
	Oil	5	87.89 (2.27)
	DHT	8	87.16 (1.10)
	Males		
	Oil	4	88.68 (1.94)
	DHT	5	84.24 (3.12)
Middle Age	*Females*		
	Oil	6	63.80 (3.43)
	DHT	6	81.32 (5.82)
	Males		
	Oil	6	67.03 (5.39)
	DHT	6	70.88 (7.00)

328   **Table 3.** Androgen receptor (AR) expression in the hippocampus using a relative rating scaling:

329   absent (0), light (+), intermediate (++), and robust (+++) in the granule cell layer (GCL), CA1,

330   and CA3 regions in young and middle-aged male and female rats treated with oil or

331   dihydrotestosterone (DHT).

Age	Sex and Treatment	AR expression		
		GCL	CA1	CA3
Young	*Females*			
	Oil	0	0/+	0
	DHT	0	+/++	0/+
	Males			
	Oil	0	0	0
	DHT	0	+/++	0/+
Middle Age	*Females*			
	Oil	0	+/++	+
	DHT	0	+++	+
	Males			
	Oil	0	+/++	0/+
	DHT	0	+++	0/+