Androgens enhance adult hippocampal neurogenesis in males but not females in an age-

dependent manner

Paula Duarte-Guterman[1], Dwayne K. Hamson, Stephanie Lieblich[1], Steven R. Wainwright[1],

Carmen Chow[1], Jessica Chaiton[1], Neil V. Watson[2], Liisa A.M. Galea[1]

1. Djavad Mowafaghian Centre for Brain Health and Department of Psychology, University of

British Columbia, Vancouver, BC, Canada

2. Department of Psychology, Simon Fraser University, Burnaby, BC, Canada

Address all correspondence and requests for reprints to:

L. A. M. Galea, PhD

Djavad Mowafaghian Centre for Brain Health

2215 Wesbrook Mall

Vancouver, BC

V6T 1Z3, Canada

E-mail: lgalea@psych.ubc.ca.

Short title: Androgens enhance neurogenesis in young males

Keywords: dihydrotestosterone; adult neurogenesis; androgen receptor; aging; sex differences

Financial support: Canadian Institutes of Health Research (MOP102568) to LAMG.

Disclosure summary: The authors have nothing to disclose.

Abstract

Androgens (testosterone and dihydrotestosterone) increase adult hippocampal

neurogenesis by increasing survival of new neurons in male rats and mice via an androgen

receptor pathway, but it is not known whether androgens regulate neurogenesis in females and

whether the effect is age-dependent. We investigated the effects of dihydrotestosterone (DHT), a

potent androgen, on neurogenesis in young adult and middle-aged males and females. Rats were

gonadectomized and injected with the DNA synthesis marker, bromodeoxyuridine (BrdU). The

following day rats began receiving daily injections of oil or DHT for 30 days. We evaluated cell

proliferation (Ki67) and survival of new neurons (BrdU and BrdU/NeuN) in the hippocampus of

male and female rats using immunohistochemistry. As expected, DHT increased the number of

BrdU+ cells in young males but surprisingly not in middle-aged male rats or in young and

middle-aged females. In middle age, DHT increased the proportion of BrdU/NeuN cells, an

effect driven by females. AR expression also increased with aging in both female and male rats,

which may contribute to a lack of DHT neurogenic effect in middle age. Our results indicate that

DHT regulates adult hippocampal neurogenesis in a sex- and age-dependent manner.

Main text

Neurogenesis, the production of new neurons, in the hippocampus continues through the life span of most mammals studied to date (1). Sex hormones (estrogens and androgens) regulate different aspects of hippocampal neurogenesis: e.g. proliferation and/or survival of these new neurons in rodents (reviewed in (2,3)). There is also evidence of sex differences in how hormones regulate neurogenesis. For example, estradiol regulates cell proliferation and survival of new neurons in female but not male rats (4). We have previously shown that androgens (testosterone and dihydrotestosterone) increase the survival of new neurons but not cell proliferation in the hippocampus of male rats and mice via an androgen receptor (AR) pathway (5,6,7). However it is not known whether androgens regulate any aspects of adult neurogenesis in females, even though ARs are expressed in the female hippocampus (8). In addition to sex, age can also modulate the effects of hormones on hippocampal neurogenesis. In middle age, neurogenesis decreases (9) while reduction of corticosteroid levels (by adrenalectomy) (10) and exercise (11) restore neurogenesis levels in aged rodents. With aging, the hippocampus also loses its ability to respond to estrogens in female rats (12,13). For instance, estradiol increases cell proliferation in the hippocampus in young but not middle-aged nulliparous female rats (12,13). The objective of this study was to investigate the effects of dihydrotestosterone (DHT) on hippocampal neurogenesis (proliferation and survival of new neurons) in young and middle-aged male and female rats.

At 2 months (~70 days old, young) and 11-12 months of age (middle-aged), male and female Sprague–Dawley rats were gonadectomized and allowed to recover for one week (n=5-8/group) (4–6,14). One week allows for circulating gonadal hormone levels to decrease to very low or undetectable levels (4,5,15,16). One day after ovariectomy, estradiol levels are

undetectable in female rats (15) and approximately 5 days after gonadectomy circulating

estradiol and testosterone decrease to approximately 10% of their original levels or to

undetectable levels in males (17,18). We chose 11-12 months as middle-age as rats can live up to

~24 months, at 12 months the levels of neurogenesis are substantially decreased compared to

young adults (9,19–21), and at 12 months sexual motivation and fecundity is significantly

reduced in both sexes (22–24). After the one week recovery period, all animals received a single

intraperitoneal injection of bromodeoxyuridine (BrdU; 200 mg/kg) to label dividing cells and

their progeny (6). The following day, chronic hormone or vehicle treatment began. Males and

females were injected subcutaneously with either 0.25 mg dihydrotestosterone (DHT in 0.1 ml of

sesame oil) or an equivalent volume of sesame oil for 30 days. The dose of DHT chosen in this

study was the lowest dose examined that increased neurogenesis in castrated young adult male

rats (5,6). Twenty-four hours after the final injection, animals were overdosed with sodium

pentobarbitol, and perfused with 4% paraformaldehyde, then brains were collected, sectioned

using a freezing microtome and processed for BrdU (survival of 30 day old cells), Ki67 (cell

proliferation marker), androgen receptor (AR), and colabelled for BrdU/NeuN (new neurons

using NeuN, marker for mature neurons) immunohistochemistry. The following primary

antibodies were used with diaminobenzidine (DAB) chromogen: mouse anti-BrdU monoclonal

(1:200; Roche Cat#11170376001 (25)), rabbit anti-Ki67 polyclonal (1:3000; Vector Laboratories

Cat# VP-K451 (26)), and rabbit anti-androgen receptor monoclonal (1:100; Abcam Cat#

ab133273 (27)). For fluorescence double labelling the following primary antibodies were used:

rat anti-BrdU (1:500; Bio-Rad/ABD Serotec Cat#OBT0030S (28)) and mouse anti-NeuN (1:250;

Millipore Cat# MAB377 (29)). Detailed protocols are found in (6,7). Thus, in this experiment,

BrdU+ cells were 30 day-old daughter cells from progenitor cells that had been synthesizing

DNA for a 2-hour period 31 days before euthanasia. A subset of samples were used to measure

DHT levels in serum collected on the day of perfusion (stored at -20°C) using a commercial

ELISA kit (IBL-America, Cat#IB59116 (30)). All samples were run in duplicate following the

manufacturer's protocol. The DHT antibody is highly specific with 8.7% cross-reactivity with

testosterone and 0.2% cross-reactivity with androstenedione; the sensitivity is 6.0 pg/ml.

Average intra-assay coefficients of variation were <15%. All protocols were approved by the

Animal Care Committee at the University of British Columbia and conformed to the guidelines

set out by the Canadian Council on Animal Care.

93   A researcher blind to experimental conditions counted all BrdU+ and Ki67+ cells in both

hemispheres for each section for the entire rostrocaudal extent of the granule cell layer (GCL)

including the sub granular zone, defined as the 50 μm band between the GCL and the hilus (13-

15 sections in total per animal). We used a modified optical fractionator method (31,32) to

estimate the total number of BrdU+ and Ki67+ cells in the GCL and hilus, as has been used

before (5,7,33–37). Total cells were calculated by multiplying the total number of cells counted

by 10 to account for the fact that we used 1/10 series of sections for each immunohistochemistry

procedure. GCL and hilus volumes were quantified from digitized images using Cavelier's

principle, multiplying the sum of the area of each section by the section thickness (40 μm) (38).

Densities of BrdU+ and Ki67+ were calculated by dividing the total number of cells by the GCL

volume. Densities (total cells per unit volume) were used as there are sex differences in the

volume of the dentate gyrus in rats (39,40). We assessed expression of AR by quantitative

densiometric analysis using ImageJ (U. S. National Institutes of Health, Bethesda, Maryland,

USA, http://imagej.nih.gov/ij/). Photomicrographs of four hippocampal sections per animal were

taken at 40X magnification using the same exposure and gain settings. Optical density (OD) was

assessed in the CA1, CA3, and GCL regions by placing six circles (40 µm diameter) along these

regions. OD levels were corrected for background levels using areas with no immunoreactivity

(i.e. molecular layer of the dentate gyrus and radiatum layer of the hippocampus). To determine

whether BrdU+ cells were of a neuronal phenotype, in a subset of brains, BrdU+ cells were

examined for co-labeling with NeuN (neuronal marker) for 50 cells (young adult group) or all

cells (middle-aged group). All analyses were performed using Statistica v.8.0 (StatSoft Inc,

Tulsa, OK). Volume of the dentate gyrus (GCL and hilus) and density of BrdU+ cells were

analyzed using repeated measures analysis of variance (ANOVA) with age (young, middle-

aged), sex (male, female), and treatment (DHT, oil) as between subjects factors, and region

(GCL, hilus) as a within subjects factor. AR OD was analyzed using repeated measures ANOVA

with age (young, middle-aged), sex (male, female), and treatment (DHT, oil) as between subjects

factors, and region (CA1, CA3, GCL) as a within subjects. Density of Ki67+ cells, proportion of

BrdU/NeuN co-labeled cells, and serum DHT concentrations were analyzed using ANOVA with

age (young, middle-aged), sex (male, female), and treatment (DHT, oil) as between subjects

factors. When appropriate, post-hoc analysis used the Neuman-Keul's procedure. Test statistics

were considered significant if $p \leq 0.05$.

Regardless of age and treatment, the volume of the GCL and hilus were larger in males

compared to females (main effect of sex; $F(1,42)=4.42$; $P<0.05$; Table 1). The volume of the

GCL and hilus increased with age irrespective of sex and treatment (main effect of age;

$F(1,42)=47.37$; $P<0.0001$). In the case of the hilus, the volume increased with aging more so in

males than in females (interaction between region, age, and sex; $F(1,42)=20.20$; $P<0.0001$). As

expected, the volume of the hilus was larger than the volume of the GCL (main effect of region;

$F(1,44)=1450.14$; $P<0.0001$). Treatment with DHT did not significantly affect GCL or hilus

volumes (all P's>0.08). To account for the sex differences in GCL and hilus volumes, we present

BrdU+ and Ki67+ cell counts as densities (total cells per unit volume).

As expected, DHT treatment increased serum levels of DHT in males and females

regardless of age (main effect of treatment; $F_{(1,17)}=36.67$; $P<0.0001$). The mean (±SEM) DHT

level in oil treated rats was 38.55 ± 2.99 pg/ml and in DHT-treated rats was 73.64 ± 8.70 pg/ml.

DHT levels were significantly higher in females than males regardless of treatment (main effect

of sex; $F_{(1,17)}=26.6$; $P<0.0001$). The mean (±SEM) DHT level in DHT-treated females was

69.58 ± 8.66 pg/ml and in DHT-treated males was 40.02 ± 3.79 pg/ml. No significant differences

were detected in DHT levels with age (P's>0.3).

DHT treatment increased the density of BrdU+ cells in the GCL in young males

($P<0.001$) but not in young females or middle-aged rats of both sexes (all P's>0.9; interaction

between age, sex, treatment and region; $F_{(1,42)}=4.03$; $P=0.05$; Figure 1). The density of BrdU+

cells in the GCL was significantly higher in the young compared to the middle-aged rats

irrespective of sex and treatment, as expected (main effect of age; $F_{(1,42)}=647.85$; $P<0.0001$). In

the hilus, the density of BrdU+ cells was not significantly affected by age, sex or treatment (all

P's>0.9; Table 1). To determine how many BrdU+ cells were neurons, we examined the

colabelling of BrdU and NeuN (a mature neuronal marker; BrdU/NeuN) in the GCL. Aging

decreased the proportion of BrdU/NeuN colabeled cells in the dentate gyrus (main effects of age;

$F_{(1,39)}=27.98$; $P<0.0001$). In the young group, sex and treatment did not affect the percent of

BrdU/NeuN colabeled cells (all P's>0.18; Table 2). In middle age, DHT treatment increased the

proportion of BrdU/NeuN cells (interaction between age and treatment; $F_{(1,39)}=4.77$; $P=0.03$),

and this effect was driven by the middle-aged females ($P=0.005$) compared to the middle-aged

males ($P=0.48$).

DHT treatment did not affect the density of Ki67+ cells in young or middle-aged males

and females (all P's>0.25; Figure 1). However, as expected, there were more Ki67+ cells in the

GCL of the young compared to middle-aged rats (main effect of age; $F(1,43)=97.18$; $P<0.0001$)

irrespective of sex and treatment.

Finally, we performed a qualitative and quantitative (OD) analysis of AR expression in

the hippocampus and both methods obtained similar results (Table 3; Figure 2, respectively). AR

OD increased significantly with aging in the CA1, CA3, and GCL, regardless of sex, and DHT

treatment increased the expression of AR in the CA1 in middle-aged rats of both sexes

(interaction between region, age and treatment; $F(2,48)=8.1$; $P<0.001$; Figure 2). However, DHT

treatment increased AR OD in the CA1, regardless of age, more so in females ($P=0.0001$) than in

males ($P=0.056$; interaction between region, sex and treatment; $F(2,48)=3.3$; $P<0.05$; Figure 2).

Qualitatively, in oil treated animals, AR-ir cells were absent throughout the GCL in young males

and female rats but were expressed at low levels in middle-aged male and female rats. In the

CA3, we found low to absent levels of AR-ir cells in young rats of both sexes and expression

increased in middle-aged rats. In the CA1, we found low to intermediate levels of AR-ir in oil

treated young rats and levels increased with aging and DHT treatment in both sexes (Figure 2).

In the present study, we found that chronic (30 days) DHT increased survival of new

neurons but not cell proliferation in the hippocampus of gonadectomized young adult male rats,

consistent with our previous research in male rats and mice (5–7). This is also in line with

previous work showing that castration decreases survival of new neurons 24-30 days after BrdU

injection but has no effect on cell proliferation (5,41). Shorter testosterone treatment (3, 15 or 21

days) has no effect on hippocampal neurogenesis (42–45) indicating that a longer exposure (30

days) to androgens is required to increase neurogenesis in the hippocampus at least in

physiological doses as higher doses can decrease neurogenesis (46,47). Thus, collectively while

longer term exposure to androgens increases survival of new neurons in the dentate gyrus,

androgens do not appear to influence cell proliferation in male rats (this study and (5,6)), mice

(7), or voles (48).

In contrast to young adult males, DHT did not affect survival of new neurons or cell

proliferation in gonadectomized young adult female rats. We previously found that estradiol

modulates cell proliferation and survival of new neurons in young adult female rats (4,49–52)

but has no significant effect on neurogenesis in adult male rats (4,5). We found that females had

higher serum DHT levels than males regardless of treatment or age. The higher DHT levels in

females were likely due to the dose being somewhat higher in females compared to males. So it

is possible that these higher circulating DHT levels resulted in an eliminated response to

neurogenesis, although we observed an increase in cell fate (BrdU/NeuN) with DHT in middle-

aged females (discussed below). Together our results suggest that sex steroids have sex-specific

effects on hippocampal neurogenesis with androgens modulating neurogenesis in young adult

males and estrogens modulating neurogenesis in young adult females. While these findings may

not seem surprising, it is important to understand that both sexes have ARs and estrogen

receptors (ERs) but these receptors are responding to respective hormones in a sex-specific way

to modulate neurogenesis in the hippocampus. Both ERα and ERβ are expressed in the

hippocampus (CA1, CA3, GCL) in males and females and no sex differences exist in their

expression (53,54). Intriguingly, there are more ERs in the GCL than there are ARs in both

sexes, and ERs have been detected on proliferating cells (Ki67+ or BrdU+ cells) or immature

neurons (doublecortin expressing cells) in adult male and female rats (50,55,56) but to our

knowledge ARs have not been detected on proliferating cells or immature neurons in the dentate

gyrus in male rats and mice (6,7). Our findings are unlikely to involve ERs (α or β) due to the

use of DHT in this study. DHT is a non-aromatizable androgen that binds with high affinity to

AR. However, 5α-androstane-3β, 17β-diol (3β-Adiol), a DHT metabolite, has been shown to

function as an ERβ ligand (57). In order to avoid any possible ERβ activation, we used the

lowest possible dose of DHT that increases neurogenesis in young males (5). From previous

work, estradiol does not increase neurogenesis in males and decreases neurogenesis (survival of

new neurons) in females (4). In addition, in young male rats, the same dose of DHT regulates

neurogenesis via ARs as blocking AR with flutamide eliminates the DHT-induced increase in

new neuron survival (6). Together this suggests that the DHT regulation of hippocampal

neurogenesis is mediated by the AR.

Perhaps surprisingly, the effect of DHT to modulate survival of new neurons was absent

in middle-aged males and females. This effect is consistent with the recent work of Moser et al.

(58) who found that testosterone did not affect the number of immature neurons (using

doublecortin), in middle-aged (13 months) or aged (23 months) male rats. This would suggest

that with aging, the dentate gyrus loses its ability to respond to sex steroids. Indeed, in females,

acute estradiol increases cell proliferation in young but not middle-aged nulliparous rats (12,13).

Intriguingly in females, previous reproductive experience (pregnancy and motherhood) can

rescue the hippocampus' response to estrogens later in middle-age, as acute estrogens increased

cell proliferation in multiparous rats (13). Thus it is possible that experience, reproductive or

otherwise, may restore the ability of androgens to upregulate hippocampal neurogenesis in males

in middle age. However, Moser et al. (58) did not see an influence of high fat diet on the

androgen modulation of neurogenesis. Intriguingly in our study, the proportion of BrdU/NeuN

colabelled cells was affected by age and treatment. DHT increased the proportion of BrdU/NeuN

colabelled cells only in middle-aged females, and to our knowledge the first description of such a

change in cell fate with aging and DHT. Overall as expected, the proportion of new cells that

express mature neuronal protein decreased with age. In middle-aged female mice, letrozole, an

aromatase inhibitor blocking the conversion of androgens to estrogens, increases hippocampal

neurogenesis (using the immature marker doublecortin; (59)). Together with our findings, this

suggests that the hippocampus may still be able to respond to sex steroid hormones in middle

age, an effect that varies by sex as in both studies middle-aged females showed increased

neurogenesis levels with androgens.

Surprisingly, we found that AR expression increased with aging in both sexes in all

hippocampal regions, which to our knowledge has not been reported before. We found ARs were

expressed at high and moderate levels in the CA1 and CA3 regions of the hippocampus,

respectively, in both young and middle-aged rats. But, consistent with previous research, we did

not find expression of ARs in the GCL in young gonadectomized male rats (6,8,60); although

there is conflicting evidence which may be due to strain, species, and age differences, as we

report AR expression in the GCL at middle age but not in young adults (see below). In Wistar

rats, Moghadami et al. (61) and Brännvall et al. (46) found AR expression in the GCL in young

gonadectomized and intact male rats. In gonadectomized male mice, ARs are not expressed in

the GCL but in males receiving DHT treatment, granule cells did express AR (7). In other rat

strains (Bruce-Spruce Long-Evans, Fischer 344 and Sprague Dawley), most studies have found

that ARs are not expressed in the GCL (6,8,60) but one study found low to medium AR

expression using qualitative analysis in the GCL of intact young male Sprague Dawley rats (62).

We also found that DHT increases the expression of ARs in the CA1 in males in line with

previous research in young male rats (6) and male mice (7) and in other brain regions (preoptic–

hypothalamic regions) (24). In females and males, 2 days of testosterone treatment also showed

increased AR expression in the CA1 region (8) but to our knowledge no other studies have

investigated AR expression in the female hippocampus after chronic DHT treatment. In female

rats, AR expression varies with the estrous cycle (62) as AR expression was highest when

estradiol levels are low in the CA1, CA3, and dentate gyrus (62). Furthermore, DHT treatment

for 7 days decreased AR expression in most brain regions in intact female rats (62), suggesting

that DHT in the presence of gonadal hormones downregulates AR expression in the female

hippocampus. In the current study, DHT increased AR expression in gonadectomized females (as

well as males), suggesting that gonadal hormones interact to regulate brain levels of AR. As

discussed above, previous work has found that androgens increase neurogenesis via the AR.

However, in young male rats and mice, ARs are not found in immature neurons (doublecortin

expressing cells) in the GCL of the hippocampus (6,7) but instead ARs are expressed in mature

neurons (NeuN expressing cells) in the hippocampus (62). We have previously proposed that one

mechanism of action is that androgens bind to ARs in the CA3 region and this initiates a

retrograde response of a survival factor that targets newborn neurons in the GCL (63). In the

current study, we found ARs were expressed in the CA1 and CA3 regions in young and middle-

aged male and female rats and their expression increases after DHT treatment in both sexes at

both ages. However, we only observed an effect of DHT on new neuron survival in young males.

In young females and middle-aged males and females, ARs are expressed in the CA1 and CA3

but binding of DHT to these ARs does not result in the modulation of neurogenesis, possibly

indicating different downstream mechanisms of bound AR to modulate neurogenesis in young

adult females, and middle-aged males. Interestingly, we found higher expression of AR in the

hippocampus of middle-aged compared to young rats of both sexes. In addition, in contrast to

young rats, we detected AR-ir cells in the GCL in middle-aged rats. AR mRNA expression also

increases in the hippocampus with aging in males (60) but to our knowledge this is the first study

showing that hippocampal AR expression increases with aging in females. It may seem

paradoxical that although DHT increases neurogenesis via the AR in males (6), despite the

increase in AR expression with middle age, DHT no longer increases neurogenesis in middle-

aged rats. However, we have previously found that overexpression of the AR in the brain

(Nestin-AR) resulted in a failure of DHT to increase neurogenesis in young adult male mice (7).

This suggests a 'dose-response' of AR expression, with optimal levels needed for DHT to

increase neurogenesis. In the preoptic area, the number of AR-ir cells increase with aging in

intact male rats (64), but not in castrated males (24), and these are negatively correlated with

circulating testosterone levels (64). In young rats, AR expression is dependent on androgen

levels and gonadectomy decreases AR expression in the hippocampus (8,60,61). Circulating

androgens decrease with age in male rats (65) and therefore we would expect AR expression to

decrease with age. Our findings and the ones by Wu et al. (64) are surprising and it is possible

that with aging the relationship between androgens and DHT changes. Indeed, we observed that

DHT treatment increased AR expression in the CA1 in middle-aged males and females

indicating that exogenous regulation of AR expression is similar in both ages. As outlined above,

high doses of AR in the brain (in transgenic male mice) result in an abolished neurogenic

response to DHT (7). It is possible that the increase in AR expression with aging is responsible

for the lack of DHT mediated increase in new neuron survival.

To summarize, our study demonstrates that DHT increases neurogenesis in young adult

males but not in young adult females and that aging eliminates the ability of DHT to enhance

neurogenesis in males showing sex and age differences in the neurogenic response to DHT.

However, in middle-aged rats, DHT treatment increased the proportion of surviving cells

expressing the mature neuronal protein (an effect seen in the females only). AR expression in the

hippocampus increases with aging and DHT treatment in both sexes, suggesting that ARs do

respond to DHT but this does not result in a neurogenic response in females and middle-aged

males. Altogether our current and previous research (reviewed in (3)) indicates that androgens

and estrogens have sex-specific and age-specific effects on hippocampal neurogenesis.

Acknowledgments

This work was funded by the Canadian Institutes of Health Research (MOP102568) to

LAMG. We thank Dr. Mark Martindale (Whitney Laboratory for Marine Bioscience, University

of Florida) for access to an epifluorescent microscope.

References

1. Kempermann G, Gage FH, Aigner L, Song H, Curtis MA, Thuret S, Kuhn HG, Jessberger
S, Frankland PW, Cameron HA, Gould E, Hen R, Abrous DN, Toni N, Schinder AF, Zhao
X, Lucassen PJ, Frisén J. Human Adult Neurogenesis: Evidence and Remaining Questions.
Cell Stem Cell 2018;23(1):25–30.

2. Duarte-Guterman P, Yagi S, Chow C, Galea LAM. Hippocampal learning, memory, and
neurogenesis: Effects of sex and estrogens across the lifespan in adults. *Horm Behav*
2015;74:37–52.

3. Mahmoud R, Wainwright SR, Galea LAM. Sex hormones and adult hippocampal
neurogenesis: Regulation, implications, and potential mechanisms. *Frontiers in*
Neuroendocrinology 2016;41:129–152.

4. Barker JM, Galea LAM. Repeated estradiol administration alters different aspects of
neurogenesis and cell death in the hippocampus of female, but not male, rats. *Neuroscience*
2008;152(4):888–902.

5. Spritzer MD, Galea LAM. Testosterone and dihydrotestosterone, but not estradiol, enhance
survival of new hippocampal neurons in adult male rats. *Developmental Neurobiology*
2007;67(10):1321–1333.

6.   Hamson DK, Wainwright SR, Taylor JR, Jones BA, Watson NV, Galea LAM. Androgens
Increase Survival of Adult-Born Neurons in the Dentate Gyrus by an Androgen Receptor-
Dependent Mechanism in Male Rats. *Endocrinology* 2013;154(9):3294–3304.

7.   Swift-Gallant A, Duarte-Guterman P, Hamson DK, Ibrahim M, Monks DA, Galea LAM.
Neural androgen receptors affect the number of surviving new neurones in the adult dentate
gyrus of male mice. *Journal of Neuroendocrinology* 2018;30(4):e12578.

8.   Xiao L, Jordan CL. Sex Differences, Laterality, and Hormonal Regulation of Androgen
Receptor Immunoreactivity in Rat Hippocampus. *Hormones and Behavior* 2002;42(3):327–
336.

9.   Kuhn H, Dickinson-Anson H, Gage F. Neurogenesis in the dentate gyrus of the adult rat:
age-related decrease of neuronal progenitor proliferation. *The Journal of Neuroscience*
1996;16(6):2027–2033.

10. Cameron HA, McKay RD. Restoring production of hippocampal neurons in old age. *Nat.*
Neurosci. 1999;2(10):894–897.

11. Praag H van, Shubert T, Zhao C, Gage FH. Exercise Enhances Learning and Hippocampal
Neurogenesis in Aged Mice. *J Neurosci* 2005;25(38):8680–8685.

12. Chiba S, Suzuki M, Yamanouchi K, Nishihara M. Involvement of Granulin in Estrogen-
Induced Neurogenesis in the Adult Rat Hippocampus. *Journal of Reproduction and*
Development 2007;53(2):297–307.

13. Barha CK, Galea LAM. Motherhood alters the cellular response to estrogens in the
hippocampus later in life. *Neurobiology of Aging* 2011;32(11):2091–2095.

14. Barha CK, Lieblich SE, Chow C, Galea LAM. Multiparity-induced enhancement of
hippocampal neurogenesis and spatial memory depends on ovarian hormone status in
middle age. *Neurobiology of Aging* 2015;36(8):2391–2405.

15. Woolley CS, McEwen BS. Roles of estradiol and progesterone in regulation of
hippocampal dendritic spine density during the estrous cycle in the rat. *J. Comp. Neurol.*
1993;336(2):293–306.

16. Barker JM, Galea LAM. Sex and regional differences in estradiol content in the prefrontal
cortex, amygdala and hippocampus of adult male and female rats. *General and*
Comparative Endocrinology 2009;164(1):77–84.

17. Andò S, Giacchetto C, Canonaco M, Aquila S, Valenti A, Beraldi E, Piro A, Dessì-Fulgheri
F. Effects of castration on androstenedione, testosterone and dihydrotestosterone plasma
levels in adult male rats. *Horm. Res.* 1986;23(2):122–127.

18. Ma L, Li W, Zhu H-P, Li Z, Sun Z-J, Liu X-P, Zhao J, Zhang J-S, Zhang Y-Q. Localization
and Androgen Regulation of Metastasis-Associated Protein 1 in Mouse Epididymis. *PLoS*
One 2010;5(11). doi:10.1371/journal.pone.0015439.

19. Rao MS, Hattiangady B, Abdel-Rahman A, Stanley DP, Shetty AK. Newly born cells in the
ageing dentate gyrus display normal migration, survival and neuronal fate choice but
endure retarded early maturation. *European Journal of Neuroscience* 2005;21(2):464–476.

20. Rao MS, Hattiangady B, Shetty AK. The window and mechanisms of major age-related
decline in the production of new neurons within the dentate gyrus of the hippocampus.
Aging Cell 2006;5(6):545–558.

21. Eid RS, Chaiton JA, Lieblich SE, Bodnar TS, Weinberg J, Galea LAM. Early and late
effects of maternal experience on hippocampal neurogenesis, microglia, and the circulating
cytokine milieu. *Neurobiology of Aging* 2019;78:1–17.

22. Chambers KC, Phoenix CH. Sexual behaviors of aging female rats: influence of age and
hormonal state of male partners. *Neurobiology of Aging* 1986;7(3):165–171.

23. Mattheij JA, Swarts JJ. Quantification and classification of pregnancy wastage in 5-day
cyclic young through middle-aged rats. *Lab. Anim.* 1991;25(1):30–34.

24. Wu D, Gore AC. Changes in androgen receptor, estrogen receptor alpha, and sexual
behavior with aging and testosterone in male rats. *Hormones and Behavior*
2010;58(2):306–316.

25. RRID:AB_514483. https://scicrunch.org/resolver/RRID:AB_514483.

26. RRID:AB_2314701. https://scicrunch.org/resolver/RRID:AB_2314701.

27. RRID:AB_11156085. https://scicrunch.org/resolver/RRID:AB_11156085.

28. RRID:AB_609570. https://scicrunch.org/resolver/RRID:AB_609570.

29. RRID:AB_2298772. https://scicrunch.org/resolver/RRID:AB_2298772.

30. RRID:AB_2801465, https://scicrunch.org/resolver/RRID:AB_2801465.

31. West MJ, Slomianka L, Gundersen HJ. Unbiased stereological estimation of the total
number of neurons in thesubdivisions of the rat hippocampus using the optical fractionator.
Anat. Rec. 1991;231(4):482–497.

32. West MJ. New stereological methods for counting neurons. *Neurobiology of Aging*
1993;14(4):275–285.

33. Kempermann G, Kuhn HG, Gage FH. More hippocampal neurons in adult mice living in an
enriched environment. *Nature* 1997;386(6624):493–495.

34. Eadie BD, Redila VA, Christie BR. Voluntary exercise alters the cytoarchitecture of the
adult dentate gyrus by increasing cellular proliferation, dendritic complexity, and spine
density. *Journal of Comparative Neurology* 2005;486(1):39–47.

35.   Tanapat P, Hastings NB, Gould E. Ovarian steroids influence cell proliferation in the
390   dentate gyrus of the adult female rat in a dose- and time-dependent manner. *J. Comp.*
391   *Neurol.* 2005;481(3):252–265.

392   36.   Snyder JS, Choe JS, Clifford MA, Jeurling SI, Hurley P, Brown A, Kamhi JF, Cameron
393   HA. Adult-born hippocampal neurons are more numerous, faster maturing, and more
394   involved in behavior in rats than in mice. *J. Neurosci.* 2009;29(46):14484–14495.

395   37.   Leuner B, Caponiti JM, Gould E. Oxytocin stimulates adult neurogenesis even under
396   conditions of stress and elevated glucocorticoids. *Hippocampus* 2012;22(4):861–868.

397   38.   Gundersen HJ, Bendtsen TF, Korbo L, Marcussen N, Møller A, Nielsen K, Nyengaard JR,
398   Pakkenberg B, Sørensen FB, Vesterby A. Some new, simple and efficient stereological
399   methods and their use in pathological research and diagnosis. *APMIS* 1988;96(5):379–394.

400   39.   Chow C, Epp JR, Lieblich SE, Barha CK, Galea LAM. Sex differences in neurogenesis and
401   activation of new neurons in response to spatial learning and memory.
402   *Psychoneuroendocrinology* 2013;38(8):1236–1250.

403   40.   Yagi S, Chow C, Lieblich SE, Galea LAM. Sex and strategy use matters for pattern
404   separation, adult neurogenesis, and immediate early gene expression in the hippocampus.
405   *Hippocampus* 2016;26(1):87–101.

406   41.   Wainwright SR, Lieblich SE, Galea LAM. Hypogonadism predisposes males to the
407   development of behavioural and neuroplastic depressive phenotypes.
408   *Psychoneuroendocrinology* 2011;36(9):1327–1341.

409   42.   Zhang Z, Yang R, Zhou R, Li L, Sokabe M, Chen L. Progesterone promotes the survival of
410   newborn neurons in the dentate gyrus of adult male mice. *Hippocampus* 2010;20(3):402–
411   412.

412   43.   Spritzer MD, Ibler E, Inglis W, Curtis MG. Testosterone and social isolation influence adult
413   neurogenesis in the dentate gyrus of male rats. *Neuroscience* 2011;195:180–190.

414   44.   Carrier N, Kabbaj M. Testosterone and imipramine have antidepressant effects in socially
415   isolated male but not female rats. *Hormones and Behavior* 2012;61(5):678–685.

416   45.   Wainwright SR, Workman JL, Tehrani A, Hamson DK, Chow C, Lieblich SE, Galea LAM.
417   Testosterone has antidepressant-like efficacy and facilitates imipramine-induced
418   neuroplasticity in male rats exposed to chronic unpredictable stress. *Hormones and*
419   *Behavior* 2016;79:58–69.

420   46.   Brännvall K, Bogdanovic N, Korhonen L, Lindholm D. 19-Nortestosterone influences
421   neural stem cell proliferation and neurogenesis in the rat brain. *European Journal of*
422   *Neuroscience* 2005;21(4):871–878.

47. Zhang W, Cheng J, Vagnerova K, Ivashkova Y, Young J, Cornea A, Grafe MR, Murphy
SJ, Hurn PD, Brambrink AM. Effects of androgens on early post-ischemic neurogenesis in
mice. *Transl Stroke Res* 2014;5(2):301–311.

48. Fowler CD, Freeman ME, Wang Z. Newly proliferated cells in the adult male amygdala are
affected by gonadal steroid hormones. *J. Neurobiol.* 2003;57(3):257–269.

49. Ormerod BK, Lee TT-Y, Galea LAM. Estradiol initially enhances but subsequently
suppresses (via adrenal steroids) granule cell proliferation in the dentate gyrus of adult
female rats. *Journal of Neurobiology* 2003;55(2):247–260.

50. Mazzucco CA, Lieblich SE, Bingham BI, Williamson MA, Viau V, Galea LAM. Both
estrogen receptor α and estrogen receptor β agonists enhance cell proliferation in the
dentate gyrus of adult female rats. *Neuroscience* 2006;141(4):1793–1800.

51. Barha CK, Lieblich SE, Galea LAM. Different Forms of Oestrogen Rapidly Upregulate
Cell Proliferation in the Dentate Gyrus of Adult Female Rats. *Journal of
Neuroendocrinology* 2009;21(3):155–166.

52. Duarte-Guterman P, Lieblich SE, Chow C, Galea LAM. Estradiol and GPER Activation
Differentially Affect Cell Proliferation but Not GPER Expression in the Hippocampus of
Adult Female Rats. *PLoS ONE* 2015;10(6):e0129880.

53. Weiland NG, Orikasa C, Hayashi S, McEwen BS. Distribution and hormone regulation of
estrogen receptor immunoreactive cells in the hippocampus of male and female rats.
Journal of Comparative Neurology 1997;388(4):603–612.

54. Kalita K, Szymczak S, Kaczmarek L. Non-nuclear estrogen receptor β and α in the
hippocampus of male and female rats. *Hippocampus* 2005;15(3):404–412.

55. Isgor C, Watson SJ. Estrogen receptor α and β mRNA expressions by proliferating and
differentiating cells in the adult rat dentate gyrus and subventricular zone. *Neuroscience*
2005;134(3):847–856.

56. Pérez SE, Chen E-Y, Mufson EJ. Distribution of estrogen receptor alpha and beta
immunoreactive profiles in the postnatal rat brain. *Developmental Brain Research*
2003;145(1):117–139.

57. Lund TD, Hinds LR, Handa RJ. The Androgen 5α-Dihydrotestosterone and Its Metabolite
5α-Androstan-3β, 17β-Diol Inhibit the Hypothalamo–Pituitary–Adrenal Response to Stress
by Acting through Estrogen Receptor β-Expressing Neurons in the Hypothalamus. *J.
Neurosci.* 2006;26(5):1448–1456.

58. Moser VA, Christensen A, Liu J, Zhou A, Yagi S, Beam CR, Galea L, Pike CJ. Effects of
aging, high-fat diet, and testosterone treatment on neural and metabolic outcomes in male
brown Norway rats. *Neurobiol. Aging* 2019;73:145–160.

59. Chaiton JA, Wong SJ, Galea LA. Chronic aromatase inhibition increases ventral
hippocampal neurogenesis in middle-aged female mice. *Psychoneuroendocrinology*
2019;106:111–116.

60. Kerr JE, Allore RJ, Beck SG, Handa RJ. Distribution and hormonal regulation of androgen
receptor (AR) and AR messenger ribonucleic acid in the rat hippocampus. *Endocrinology*
1995;136(8):3213–3221.

61. Moghadami S, Jahanshahi M, Sepehri H, Amini H. Gonadectomy reduces the density of
androgen receptor-immunoreactive neurons in male rat's hippocampus: testosterone
replacement compensates it. *Behavioral and Brain Functions* 2016;12(1).
doi:10.1186/s12993-016-0089-9.

62. Feng Y, Weijdegård B, Wang T, Egecioglu E, Fernandez-Rodriguez J, Huhtaniemi I,
Stener-Victorin E, Billig H, Shao R. Spatiotemporal expression of androgen receptors in the
female rat brain during the oestrous cycle and the impact of exogenous androgen
administration: A comparison with gonadally intact males. *Molecular and Cellular*
Endocrinology 2010;321(2):161–174.

63. Galea LAM, Wainwright SR, Roes MM, Duarte-Guterman P, Chow C, Hamson DK. Sex,
Hormones and Neurogenesis in the Hippocampus: Hormonal Modulation of Neurogenesis
and Potential Functional Implications. *Journal of Neuroendocrinology* 2013;25(11):1039–
1061.

64. Wu D, Lin G, Gore AC. Age-related changes in hypothalamic androgen receptor and
estrogen receptor α in male rats. *Journal of Comparative Neurology* 2009;512(5):688–701.

65. Rosario ER, Chang L, Beckett TL, Carroll JC, Murphy MP, Stanczyk FZ, Pike CJ. Age-
related changes in serum and brain levels of androgens in male Brown Norway rats.
Neuroreport 2009;20(17):1534–1537.

 FIGURES

 Figure 1. Chronic dihydrotestosterone (DHT) treatment increases the density of BrdU+ cells in

 the dentate gyrus of young male adult rats but has no effects in middle-aged males and young

 and middle-aged female rats. (**A**) Photomicrographs of a representative section of dentate gyrus

 with the granule cell layer (GCL), hilus, CA1 and CA3 regions and representative BrdU+ and

 Ki67+ cells in the subgranular zone. (**B**) Mean ± SEM total number of BrdU+ cells in the

 granule cell layer of the dentate gyrus in young and middle-aged male and female rats. In young

 males, DHT increased the density of BrdU+ cells relative to the oil treatment group (*** $P <$

 0.001; interaction between age, sex, treatment, and region; $P=0.05$). (**C**) Mean ± SEM total

 number of Ki67+ cells in the granule cell layer of the dentate gyrus in young and middle-aged

 male and female rats. Chronic DHT treatment did not affect cell proliferation in the hippocampus

 of young or middle-aged male and female rats. Circles represent individual data points (number

 of individuals).

 Figure 2. Chronic dihydrotestosterone (DHT) treatment increases AR optical density in the CA1

 region of the hippocampus in young and middle-aged male and female rats. (**A**) Representative

 images of androgen receptor (AR) expression in the hippocampus of young and middle-aged

 gonadectomized oil treated female and male rats. AR expression increased with aging in the

 CA1, CA3 and GCL in both sexes. (**B**) Representative images of androgen receptor (AR)

 expression in the CA1 region of the hippocampus in young male and female rats treated with oil

 or DHT. AR is lowly expressed in the CA1 of gonadectomized oil treated animals. DHT

 increases the expression of AR in both female and male rats. Optical density (OD) for AR (mean

 ±SEM) was measured in the CA1 (**C**), CA3 (**D**), and GCL (**E**) regions. AR OD increased

significantly with aging in the CA1, CA3, and GCL, regardless of sex, and DHT treatment

increased the expression of AR in the CA1 in young and middle-aged rats of both sexes

(interaction between region, age and treatment; $P<0.001$). DHT treatment increased AR OD in

the CA1, regardless of age, more so in females than in males (interaction between region, sex

and treatment; $P<0.05$). Asterisks denote significant differences between oil and DHT treatment

(*$P\leq0.05$; ***$P<0.001$). Circles represent individual data points (number of individuals).

TABLES

Table 1. Mean (SEM) volume of the granule cell layer (GCL) and the hilus (mm^3) and BrdU

density in the hilus in gonadectomized young and middle-aged male and female rats treated with

oil or dihydrotestosterone (DHT). The volume of the hilus was larger than the volume of the

GCL (main effect of region; P<0.0001). Males had a larger volume of GCL and hilus than

females (main effect of sex; P<0.05). The volume of the GCL and hilus increased with age

(main effect of age; P<0.0001) and in the hilus, the volume increased with aging more so in

males than females (interaction between region, age, and sex; P<0.0001). DHT treatment did not

affect the volume of the GCL and hilus (all P's>0.08). The density of BrdU+ cells was not

significantly affected by age, sex or treatment (all P's>0.9). Sample size is number of

individuals.

Age	Sex and Treatment	Sample size	GCL volume (mm^3)	Hilus volume (mm^3)	Hilus BrdU density (per mm^3)
Young	*Females*				
	Oil	7	2.51 (0.12)	6.09 (0.19)	85.14 (15.34)
	DHT	8	2.46 (0.08)	6.00 (0.28)	76.06 (7.06)
	Males				
	Oil	5	2.31 (0.19)	5.32 (0.49)	96.18 (23.29)
	DHT	6	2.62 (0.21)	5.37 (0.51)	134.70 (17.38)
Middle Age	*Females*				
	Oil	6	2.33 (0.09)	7.67 (0.42)	20.69 (7.89)
	DHT	6	2.45 (0.15)	7.33 (0.28)	6.40 (2.49)
	Males				
	Oil	6	2.94 (0.25)	9.05 (0.79)	6.06 (1.41)
	DHT	6	2.87 (0.22)	10.05 (0.60)	4.65 (1.63)

Table 2. Mean (SEM) percentage of cells co-expressing BrdU and NeuN in the GCL in

gonadectomized young and middle-aged male and female rats treated with oil or

dihydrotestosterone (DHT). The proportion of BrdU+ cells colabelled with NeuN was

significantly higher in young compared to middle-aged animals (main effects of age; P<0.0001.

In middle age, but not in young adults, DHT increased the proportion of BrdU/NeuN colabelled

cells relative to the oil treatment group (interaction between age and treatment) but this effect

was driven by the middle-aged females (P=0.005) compared to the middle-aged males (P=0.48).

** P<0.01 relative to oil treatment group. Sample size is number of individuals.

Age	Sex and Treatment	Sample size	BrdU/NeuN %
Young	*Females*		
	Oil	5	87.89 (2.27)
	DHT	8	87.16 (1.10)
	Males		
	Oil	4	88.68 (1.94)
	DHT	5	84.24 (3.12)
Middle Age	*Females*		
	Oil	6	63.80 (3.43)
	DHT	6	81.32 (5.82)**
	Males		
	Oil	6	67.03 (5.39)
	DHT	6	70.88 (7.00)

533   **Table 3.** Androgen receptor (AR) expression in the hippocampus using a relative rating scaling:

534   absent (0), light (+), intermediate (++), and robust (+++) in the granule cell layer (GCL), CA1,

535   and CA3 regions in young and middle-aged male and female rats treated with oil or

536   dihydrotestosterone (DHT).

Age	Sex and Treatment	AR expression		
		GCL	CA1	CA3
Young	*Females*			
	Oil	0	+	0/+
	DHT	0	++/+++	0
	Males			
	Oil	0	0/+	0
	DHT	0	++	0
Middle Age	*Females*			
	Oil	0/+	++	+
	DHT	+	+++	+/++
	Males			
	Oil	0/+	++	+
	DHT	+	++/+++	+

Figure 1

Figure 2