Energy balance implications of floral traits involved in pollinator attraction and water balance

Adam B. Roddy*

School of Forestry & Environmental Studies, Yale University, New Haven, CT, USA
*Corresponding author: adam.roddy@yale.edu

Abstract

For most angiosperms, flowers are critical to reproduction because they increase rates of outcrossing. Flowers are highly variable in numerous traits, including size, shape, and color. Most of this variation is thought to have arisen due to selection by pollinators. Yet, non-pollinator selection is increasingly being recognized as contributing to floral trait evolution. One such non-pollinator agent of selection that often opposes pollinator selection includes the physiological and resource costs of producing and maintaining flowers. Here I (1) summarize recent studies on the macroevolution of floral physiological traits and (2) apply an energy balance model to examine how two pollination traits (flower color and flower size) can interact with hydraulic traits to influence flower physiology. These modeling results show that under certain conditions flower color variation can overwhelm the effects of floral transpiration and flower size variation on flower temperature. Additionally, the range of flower size most common in the California flora is the range in which complex, non-linear dynamics in flower energy balance occur. These results suggest that floral traits under selection by biotic agents can have large implications for flower physiology, and only some of these potentially deleterious effects can be offset by flower hydraulic traits. These complex interactions between pollination traits (flower size and color) and physiological traits (surface conductance to water vapor) suggests that a more unified framework for understanding the evolution of floral form and function would simultaneously consider the interaction between physiological traits and traits under biotic selection.

Introduction

It is commonly assumed that an organism's phenotype is the result of adaptation. Selection, it is thought, hones the various traits of an organism to best maximize fitness in the ecological context in which the organism thrives. The fundamental challenge facing an organism is, therefore, one of allocation: how to allocate its limited resources in order to maximize survival and reproduction. But simply observing the diversity of plant life makes obvious the fact that there is no single solution to this problem. Allocation even to reproductive structures–whose function is most immediately tied to fitness–can vary, in terms of biomass, from 1-60% among plant species (Bazzaz et al. 1987). One way to reconcile the presumption of maximizing fitness with the obvious diversity of phenotypes has been with optimality models (e.g. Parkhurst and Loucks 1972; Givnish and Vermeij 1976; Ashman and Schoen 1994). These models define cost and benefit functions, which, when solved under varying conditions, yield a variety of optimal solutions. In evolutionary models, the number of equally fit solutions increases and the maximum fitness of any one of them decreases as the number of constraints or selective agents increases (Niklas 1994). Put simply, phenotypic diversity increases as organisms must perform more tasks.

Defining the costs of plant structures for optimality models is inherently a question of physiology–what are the costs of producing and maintaining the structure? The most common currency of cost is carbon, which is a basic component of plant structures and is necessary for respiration. For photosynthetic structures, such as leaves, water is also considered a cost because the diffusion of CO_2 from the atmosphere into the leaf requires water to evaporate. As a result, optimal leaf size and shape can be predicted for different environmental conditions based on the carbon and water costs of construction and maintenance, the carbon benefits of photosynthesis, and the various environmental factors (light, temperature, humidity) that influence the photosynthetic efficiency of leaves (Parkhurst and Loucks 1972; Givnish and Vermeij 1976). Thus, the factors that generate phenotypic diversity are those that influence the cost and benefit functions, which are typically in terms of carbon and water.

For flowers, however, benefit functions are defined by ovule fertilization and pollen

dissemination (e.g. Ashman and Schoen 1994). Since the recognition and articulation of the

role of flowers in plant reproduction over two centuries ago (Sprengel 1793; Vogel 1996),

there has, rightfully, been a strong emphasis on characterizing the diversity of floral form

and the adaptive value of various floral traits. These studies have elaborated the numerous

floral traits that can be involved in pollinator interactions (Kevan and Baker 1983; Fenster

et al. 2004), have detailed how different floral morphological traits are associated with

different pollinators (e.g. Whittall and Hodges 2007), and have shown how certain

combinations of floral characters are likely more advantageous than others (Stebbins

1951a; O'Meara et al. 2016). Yet, the costs of flowers–the resources associated with their

production and maintenace (Reekie and Bazzaz 1987a,b)–have received notably less

attention (Gleason 2018), despite their key role in determining fundamental traits, such as

longevity (Ashman and Schoen 1994) and despite the tradeoff between resource

investments in reproduction and leaf function (i.e. gas exchange and growth; Bazzaz et al.

1987; Reekie and Bazzaz 1987c; Galen et al. 1999). Furthermore, including the

maintenance costs of flowers–e.g. rates of respiration–along with their static production

costs can improve predictions of future reproductive effort (Ashman 1994). Thus,

characterizing the adaptive accuracy of flowers for their pollination function may ignore

important physiological factors that may also exert substantial influence over floral form

and evolution.

Although pollinators are still considered the primary drivers of floral traits (Fenster et al.

2004), over the last few decades the role of resource availability has been increasingly

acknowledged as an important factor shaping floral traits (Sapir 2017; Sapir and Ghara

2017; Caruso et al. 2018). Because non-pollinator agents of selection often oppose

pollinator selection, there is often a tradeoff between these competing agents (Galen 1999;

Strauss and Whittall 2006). Despite the importance of these non-pollinator agents of

selection, little effort has been made to fully characterize the physiological costs of flowers

and the evolution of traits underlying these costs. While studies focused on one or a few

species help to elucidate physiological mechanisms (e.g. Galen et al. 1999; Chapotin et al.

2003; Roddy and Dawson 2012; Teixido and Valladares 2014a, 2015; Roddy et al. 2018), a

comparative, macroevolutionary framework is critical to understanding the long-term, net

impacts of variation in floral physiological traits on evolutionary dynamics and to

distinguishing the developmentally possible from the selectively advantageous.

Furthermore, the climate is changing faster than even the most aggressive models have

predicted (Raupach et al. 2007), driving major shifts in phenology (Fitter and Fitter 2002;

Parmesan and Yohe 2003) and plant investment in reproduction (Wright and Calderon

2006). Because these phenological shifts have physiological underpinnings and

implications, understanding the physiology of flowers is, therefore, critical to

understanding the causes and consequences of shifting flowering time on plant function

and evolution.

Here I have two aims. First, I summarize recent research elucidating the physiological traits

of flowers, focusing primarily on hydraulic architecture, and layout a series of hypotheses

about the evolution of hydraulic structure-function relationships in flowers. That abiotic

agents of selection on flowers can overwhelm the effects of biotic agents (Caruso et al.

2018) motivates a more mechanistic and more comparative understanding of the

physiology of flowers. Second, I apply an energy balance modeling approach to show how

traits important to pollinator interactions can have physiological implications. I focus on

two traits, flower color and flower size, both of which experience selection by pollinators.

My approach here is, by no means, comprehensive; rather, I aim to showcase how a

physiological perspective can elucidate new insight into floral form, function, and

evolution.

Centering metabolism in macroevolutionary studies of flowers

The origin and early diversification of the flowering plants may not have presaged the

grandeur of their future. The first angiosperms were most likely ruderal and adapted to

living under the dominant ferns and gymnosperms (Stebbins 1951b; Feild et al. 2004). Yet

the angiosperms rapidly diversified, accumulating novel traits–new reproductive

structures and new anatomical features that facilitated higher rates of metabolism–that

enabled them to dominate ecosystems globally and further promoted their diversity and

dominance (Bond 1989; Crepet and Niklas 2009; Brodribb and Feild 2010; de Boer et al.

2012; Simonin and Roddy 2018). Traditionally, innovations in floral reproductive traits

were thought to have been the primary drivers of angiosperm diversity (Sanderson and

Donoghue 1994; Crepet and Niklas 2009), but more recent work has suggested that their

ecological dominance–and also possibly their diversity–may be due to factors linked to

metabolism (Brodribb and Feild 2010; de Boer et al. 2012; Simonin and Roddy 2018).

While the earliest angiosperms were likely drought-intolerant with low physiological rates

(Feild and Arens 2007; Feild et al. 2009a), subsequent lineages rapidly developed smaller,

more densely packed cells that allowed unprecedented rates of primary (Simonin and

Roddy 2018).

Central to this emerging perspective about early angiosperm evolution and their

subsequent dominance is that the need to supply resources for primary metabolism

(photosynthesis) is a main selective agent driving the structure and organization of cells

and tissues. Because the diffusion of CO_2 into the leaf mesophyll requires that stomata on

the leaf surface be open, the wet internal surfaces of the leaf are exposed to a desiccating

atmosphere. In this way CO_2 uptake is mechanistically coupled to the transport of water

throughout the leaf such that traits associated with water supply and loss are coordinated

(Sack et al. 2003; Brodribb et al. 2005, 2007; Scoffoni et al. 2016). However, given that

most flowers are heterotrophic and assimilate little carbon, selection for high rates of

resource transport has likely been relaxed, potentially leading to novel combinations of

anatomical and physiological traits (Olson and Pittermann 2019).

Because flowers are heterotrophic and short-lived, the higher rates of metabolism that

smaller, more densely packed cells enable (Simonin and Roddy 2018) are unneccessary in

flowers. Flowers have lower vein and stomatal densities and larger cells than their

conspecific leaves (Lipayeva 1989; Feild et al. 2009b; Roddy et al. 2013, 2016; Zhang et al.

2018). The interspecific correlations between traits are similar within leaves and within

flowers, for both anatomical and physiological traits (Roddy et al. 2016, 2019; Zhang et al.

2018), suggesting that similar scaling relationships define flower and leaf trait covariation

despite divergent selection regimes acting on flowers and leaves. This divergence in flower

and leaf hydraulic traits points to the importance of developmental modularity, which has

allowed flowers and leaves to occupy difference regions of trait space due to their different

selective constraints (Berg 1960; Roddy et al. 2013, 2019; Olson and Pittermann 2019).
Furthermore, greater modularity of foliar and floral traits may have occurred among more
recently derived groups. While ANA grade and magnoliid leaves have lower vein and
stomatal densities than other angiosperms, their flowers have relatively high vein and
stomatal densities (Figure 1; Roddy et al. 2016). The hydraulic traits of ANA grade and
magnoliid flowers, therefore, seem to be more similar to those of leaves, while leaf and
flower hydraulic traits of the monocots and eudicots are more different from their
conspecific leaves. Though these patterns are based on relatively few species, they do
suggest that metabolism has been a central agent of selection shaping flower physiology,
just as it has shaped leaf physiology.

Reducing the abundance of stomata may have been a key physiological innovation with
cascading consequences on other floral traits. In leaves, stomata function to regulate water
loss while CO_2 diffuses into the leaf mesophyll where it is assimilated into sugars. At the
same time, transpired water carries with it latent heat which helps to cool the leaf and
prevent it from overheating. Volatile organic compounds (VOCs), which can function in
communication, defense, and pollinator attraction, can also be released through leaf
stomata (Baldwin 2010). Flower corollas, in constrast, are generally heterotrophic, and,
therefore, may be relaxed from the need to transpire high rates of water to support
photosynthesis. Relaxing this constraint would have allowed flowers to reduce water loss
by eliminating stomata. While stomata may be important in VOC release, other structures,
such as osmophores, crenulated epidermal cells, and trichomes, can also be sources of VOC
release in flowers (Baldwin 2010). Eliminating stomata would have, in turn, allowed the
densities of veins to be reduced (Roddy et al. 2013). This transition would have reduced
both water and carbon costs of flowers and possibly would have allowed alternative modes
of hydration. While *Illicium* (Schisandraceae) and magnoliid flowers are hydrated by the
xylem (Feild et al. 2009a,b; Roddy et al. 2018), some have argued based on water potentials
that eudicot flowers are hydrated by the phloem (Trolinder et al. 1993; Chapotin et al.
2003). Such a transition in the basic plumbing of a flower would have been a critical event
in floral evolution, but unequivocal evidence for this pattern remains lacking. Further,

other physiological properties, such as high hydraulic capacitance could generate similar

patterns as those considered indicative of phloem-hydration (Roddy et al. 2018).

On an area basis, these transitions in both water and carbon costs seem to have occurred in

the common ancestor of monocots and eudicots, although better sampling of these

divergences is strongly needed (Figure 1). While reducing stomatal densities would have

reduced maximum transpiration rate, stomata play a critical role in dynamically regulating

water loss in leaves and flowers (Roddy et al. 2018). Without stomata, water loss from

flowers may be controlled solely by epidermal and cuticular traits. The non-stomatal

epidermal conductance of leaves (commonly termed the minimum epidermal conductance,

g_{min}) is increasingly being considered an important trait that may influence drought

tolerance (Duursma et al. 2019), and in flowers it correlates with vein density and whole-

flower hydraulic conductance (K_{flower}), suggesting that it is key to flower water balance as

well (Roddy et al. 2016). If selection has driven reduced water loss from flowers, then the

loss of stomata would have drastically reduced maximum transpiration rates and any

further selection on reduced water loss may have revealed epidermal and cuticular traits to

selection on water balance.

The macroevolutionary patterns in floral hydraulic traits suggest that innovations in floral

water balance may have been a key dimension of floral evolution among early

angiosperms, but is flower water balance under contemporary selection as well?

Correlations between traits and climate are often taken as strong evidence of natural

selection; within species, larger flowers and floral displays occur in wetter habitats

(Lambrecht 2013). Because larger flowers can more readily heat up, the disproportionate

maintenance costs of large flowers can lead to shorter longevities (Teixido and Valladares

2014a, 2015), suggesting that physiological traits related to water balance interact with

other traits linked to pollination success (Galen 1999, 2000; Teixido and Valladares 2013,

2014b). Insect pollinators frequently prefer warmer flowers particularly in cool climates

(Kevan 1975; Dyer et al. 2006), yet flowers can overheat particularly in warm climates

(Patiño and Grace 2002; Patiño et al. 2002). With morphological and physiological traits

flowers must balance the need to prevent overheating and the benefit of being warmer to

attract pollinators. These early results suggest that the need to maintain water balance

may influence key pollination traits and affect floral form and function over both
contemporary and historical timescales.

Physiological implications of pollination traits

In cold habitats, flower shape and orientation are under selection to increase the
temperature of the flower microenvironment in order to attract pollinators (Kevan 1975;
Stanton and Galen 1989; Atamian et al. 2016). In many cases, then, elevated flower
temperatures are beneficial, as they also can be involved in the release and dissemination
of floral volatiles (Jakobsen and Olsen 1994; Sagae et al. 2008; Schiestl 2015), a component
of pollination interactions that likely predates the angiosperms (Terry et al. 2007; Peñalver
et al. 2012). But flower temperatures cannot exceed critical temperatures that might
hinder pollination and seed development, and so there is a tradeoff between elevating
temperatures to attract and reward pollinators and maintaining temperatures cool enough
to prevent overheating.

Whether tall, canopy trees or small, herbaceous annuals, plants commonly place their
flowers above their leaves, where air temperature is warmest and humidity is lowest. This
placement can have important implications on the energy and water balance of flowers
because of high solar radiation and exposure to high windspeeds. For example, in
seasonally dry tropical forests, most species flower in the dry season, when rainfall and
atmospheric humidity are low and air temperature and solar radiation high, creating
conditions of high evaporative demand (Wright and Van Schaik 1994; Wright and Calderon
1995; Wright and Calderon 2006). In the understory, conditions are generally thought to be
more stable, but even understory flowers can still overheat to temperatures that induce
mortality in carpels and pollen (Patiño and Grace 2002).

One way to regulate temperature is through latent heat loss due to the evaporation of
water. Leaves transpire substantial amounts of water through their stomata, but the
relatively low abundance of stomata on flower petals (Lipayeva 1989; Roddy et al. 2016;
Zhang et al. 2018) suggests their ability to regulate water loss and temperature is limited.
Yet, stomatal densities and, thus, maximum transpiration rates, have been measured on

relatively few species. Furthermore, many traits that influence energy balance (e.g. color,

shape) are more variable in flowers than in leaves, suggesting that the dynamics of floral

temperature may be more complicated, particularly because of the multiple selective

pressures acting on flowers. One first step in understanding these dynamics would be to

determine the sensitivity of flower energy balance to variation in traits linked to pollinator

preference. The lower vein and stomatal densities of flowers suggest that they have lower

rates of transpiration, which would hinder their ability to dissipate energy through latent

heat loss. Using energy balance as a framework, I focus here on two flower traits important

to pollinator interactions: flower color and flower size. In order to be seen and visited by a

pollinator, a flower must stand out against the green backdrop of leaves and be distinct

from other flowers in the community or from closely related species (Spaethe et al. 2001;

Peter and Johnson 2008; Hopkins and Rausher 2012; Muchhala et al. 2014). Differences

between pollinators in their ability to see colors can help to drive diversity in flower color,

often in association with other morphological traits (Whittall and Hodges 2007), but

pollinators are not the only selective agent on flower color. Similarly, flower size is

commonly under positive selection from visually-oriented pollinators because larger

flowers are more easily visible to pollinators, yet the physiological costs of producing large

flowers can oppose pollinator selection (Galen 1999; Strauss and Whittall 2006)

Energy balance model

To test the sensitivity of flower energy balance to variation in flower color and flower size, I

modeled energy balance using the R package 'tealeaves' (Muir 2019). Because these models

are generally applied only to leaves, there is an implicit assumption that the structure being

modeled is planar, and so the temperature modeled is most applicable to a flat petal. In

reality, more complex, three-dimensional shapes would cause deviation from the modeled

temperatures, and these deviations could either increase or decrease temperature. For

example, tubular flowers with fused petals would probably thicken the boundary layer,

reduce transpiration, and elevate flower temperature (T_{flower}) above that modeled

assuming a flat flower. Other morphological traits, such as petal curvature, the presence of

trichomes, and flower orientation would also influence energy balance. As a first

approximation, however, I have assumed flowers to be flat and oriented perpendicular to

the sky in order to estimate how variation in flower color and flower size may impact

energy balance.

In all model runs, I kept constant solar radiation (660 W m^{-2}, which is approximately 1500

µmol quanata m^{-2} s^{-1}), air temperature (T_{air}, 25^{o}C), relative humidity (50%), windspeed (2

m s^{-1}), and atmospheric pressure (100 kPa). Variation in each of these parameters would

influence transpiration, latent heat loss, and thus, the calculated flower temperature

(T_{flower}). The modeled T_{flower} results, therefore, reflect the relative effects of the varied

parameters (color, size, surface conductance to water vapor) and not the absolute

temperatures the flowers may experience.

Because such modeling is common in studies of leaves, I focus not on the effects of these

environmental parameters but instead on the effects of traits important to flower function:

color affects absorptivity of shortwave radiation, the total surface conductance to water

vapor (g_t) affects transpiration rate and latent heat loss, and flower size has complex effects

on multipe parameters that affect energy balance. Two of these traits, color and size, are

commonly under pollinator selection, and I examined how variation in these traits affect

energy balance in combination with variation in g_t. I parameterized the model with

published values of g_t (5, 25, 50, 100 mmol m^{-2} s^{-1}, which are equivalent to 0.05, 0.25, 0.5,

and 1 µmol m^{-2} s^{-1} Pa^{-1} at 100 kPa atmospheric pressure) that span the range of stomatal

and minimum epidermal conductances (g_{min}) measured from a phylogenetically diverse set

of species (Roddy et al. 2016, 2018). For flowers lacking stomata, the epidermal

conductance is the primary contributor to g_t. Further details about the energy balance

calculations (e.g. characteristic dimension, absorptance) are discussed in the relevant

sections below.

Flower color

The optical properties of flowers are influenced by a variety of factors, including surface

texture (e.g. cell shape and cuticle structure; Whitney et al. 2009), internal structure of the

mesophyll (van der Kooi et al. 2016), which can scatter absorbed light, and the type and

distribution of pigments throughout the petal (Kay et al. 1981). These factors that influence

floral color profiles can also influence flower energy balance (van der Kooi et al. 2017) and

water relations. The surface texture of petals can have a significant impact on the amount

of light–and, by extension, energy–that is reflected or absorbed (Gorton and Vogelmann

1996; van der Kooi et al. 2015, 2017; Vignolini et al. 2015). In general, darker pigmentation

leads to higher absorptance, elevating flower temperature, but latent heat lost to

transpiration can prevent overheating of the floral microenvironment (Patiño and Grace

2002).

To test for the effects of flower color in combination with variation in g_t on energy balance,

I modeled energy balance using the fixed parameters stated above and varied the

absorptance of shortwave radiation from 1% to 90% at a fixed flower size of approximately

28 cm^2 (equivalent to a characteristic dimension of 3 cm, assuming circular flowers). This

variation in absorptance spanned almost the entire range of possible absorptances, and so

to contextualize it, I measured reflectance and transmittance profiles of three *Dodecatheon*

sp. (Primulaceae) flower morphs that differ in flower color (white, pink, red) in order to

compare how intraspecific variation in flower color can affect absorptance while

controlling for variation in other anatomical and physiological traits. I used an Ocean

Optics S2000 spectrometer (Ocean Optics, Largo, Florida, USA) with sensitivity between

400 and 900 nm interfaced with an integrating sphere to measure white, pink, and red

flowers. Reflectance and transmittance spectra were calculated from measurements on

individual petals (approx. 1-cm^2), having subtracted out the background spectrum with no

illumination and normalized by a 99.9% reflectance standard. Absorptance profiles were

calculated as the difference between unity and the sum of the reflectance and

transmittance profiles. While the white flower had near uniform absorptance,

transmittance, and reflectance across the visible waveband, the red and pink flowers had

higher absorptance below 650 nm (Figure 2a-c). Red flowers had near complete

absorptance in the 400-600 nm band. These differences in absorptance were expected to

drive differences in temperature. Using a representative spectrum for incoming solar

radiation measured on a sunny day, I calculated a weighted average absorptance as:

$$\overline{A} = \frac{\sum(A \cdot \%E_e)}{\sum(\%E_e)}$$

where A is the absorptance at each wavelength and %E_e is the intensity of that wavelength

scaled to the maximum intensity of the incoming solar radiation. The weighted average

absorptances for the white, pink, and red flowers were 0.29%, 26.8%, and 64.2%,

respectively. The spectral profiles for these flowers are simply examples to show how

color variation can affect absorptance.

While darker flowers were more absorptive, which elevated T_{flower}, higher g_t somewhat

reduced T_{flower} (Figure 2d). Darkening flowers from white to red caused T_{flower} to increase

by 6-7°C. However, varying g_t twenty-fold caused T_{flower} to change less than 1°C for white

and pink flowers and approximately 1.3°C for red flowers. Therefore, the effects of flower

color on energy balance can overwhelm the effects of latent heat loss due to transpiration,

under the conditions modeled here. The relatively small effects of g_t on T_{flower}, relative to

the effects of flower color, may explain why flowers of the monocots and eudicots have few

stomata. Increasing stomatal abundance may help to prevent their flowers from

overheating, but the effects would be relatively small and the benefits offset by the cost of

losing water and by the potential benefit of increasing temperature to volatilizeorganic

compounds. In certain habitats, however, high g_t may be beneficial, such as in humid

tropical environments where high humidity and low windspeeds would otherwise reduce

transpiration rates and lead to elevated T_{flower} (Patiño and Grace 2002).

Flower size

Despite pollinator selection for larger flowers, flowers are not universally large, and the

complexity of their shape has hindered progress toward understanding the evolution of

flower size. Methods for estimating flower size vary depending on the taxonomic group

and the perceived importance of the sizes of different floral organs to animal pollinators. In

regional floras, flower size is commonly reported as either petal length or flower diameter,

yet these ways of estimating size are not necessarily physiologically meaningful. First, I

tested the utility of flower size estimates available from floras by comparing measurements

of the surface area of flowers to reports of flower size from the regional flora of California

(Jepson eFlora, 2015). Second, I used this variation in flower size to parameterize the

characteristic dimension used in the energy balalnce model. I combined variation in flower

size with variation in flower color and g_t to highlight the range of thermal outcomes when
all three traits are allowed to covary.

First, I measured the two-dimensional, projected surface area of floral perianths on fully
expanded flowers of 43 species from 29 families growing at the University of California
Botanical Garden. Flowering shoots were excised, the cut shoot surfaces immediately recut
underwater, and the shoots transported back to the laboratory where flowers were
dissected. The calyx and corolla structures were individually placed on a flatbed scanner,
and their areas measured using ImageJ (Rueden et al. 2017), totaled per flower and
averaged per species. This provided two potential estimates of flower size: one in which
the corolla and calyx structures were summed and one in which only the corolla structures
were included. For flowers lacking a differentiated perianth, such as monocots and
magnoliids, I included all tepals. For these 43 species, I extracted estimates of flower size
from the Jepson eFlora (2015), which were reported as either petal length or flower
diameter and often included a minimum and maximum. Because measurements of flower
size were in units of area (i.e. square centimeters), reports from the flora needed to be
converted into similar units for comparison. I averaged the minimum and maximum
measurements reported in the flora and made two assumptions to convert these estimates
into units of area: (1) that flowers are circular, and (2) that the length of a petal was
equivalent to the radius of this circle. These assumptions were probably safer for
actinomorphic flowers than zygomorphic flowers. I used standard major axis regression (R
package "smatr"; Warton et al. 2012) to compare the slopes and intercepts in the
relationships between predicted and measured flower size. Flower size estimated from the
flora was a strong predictor of actual flower size, for whole flowers (slope = 0.94 [0.79,
1.11]; R^2 = 0.63, P < 0.0001; Figure 3a) and for corollas (slope = 1.03 [0.88, 1.22]; R^2 = 0.65,
P < 0.0001; Figure 3b). For neither relationship was the slope significantly different from
unity (P = 0.45 and P = 0.69, respectively), suggesting that sampling a sufficient number of
species allows estimates of flower size from floras to reliably reflect physiologically
meaningful estimates of flower size. This result enables mining published floras for data on
flower size, a trait that is known to be important to pollination but for which there has been
no consistent method for measurement.

Second, I used these measurements of flower size to to parameterize the energy balance

model. Most species in the dataset had relatively small flowers, below 20 cm² (Figure 4a). I

varied flower size in the energy balance model by assuming that the characteristic

dimension that represents size was equivalent to flower diameter. I varied this

characteristic dimension from 0.5 cm to 12 cm, which represents variation in corolla

surface area from approximately 0.2 cm² to 113 cm², the range observed among the

California flora (Figure 4a). I did this for three values of absorptance, equivalent to the

white, pink, and red flowers used above, and in combination with variation in g_t between

0.05 and 1 μmol m^{-2} s^{-1} Pa^{-1}.

All three vairables had strong impacts on T_{flower} (Figure 4b). Consistent with the previous

results (Figure 3d), flower color had the greatest overall effect on T_{flower}. Increasing g_t

universally decreased T_{flower}, and this effect was greater for darker flowers. Above

approximately 15 cm² in surface area, increasing flower size had relatively little effect on

T_{flower}. Among small flowers, however, flower size variation had large and inconsistent

effects on T_{flower}. While reducing flower size among small, white flowers elevated

temperatures, reducing flower size among pink and red flowers decreased T_{flower}.

Coincidentally, these large effects of small changes in flower size on T_{flower} occurred in the

range of flower sizes that are most common among the species sampled from the California

flora (Figure 4a).

The morphological complexity of flowers complicates more accurate modeling of flower

energy balance. For these simulations, I assumed flowers were planar, but many flowers

are curved or tubular, both of which would likely reduce latent heat loss by increasing

boundary layer resistance, causing an increase in T_{flower}. In contrast, dissection (e.g. by

having unfused, dissected petals) would likely reduce the boundary layer thickness and

increase latent heat loss. Despite the morphological complexity among flowers, as a first

approximation, this sensitivity analysis shows that nonlinear dynamics in the energy

balance model may be particularly important for small-flowered species, which are

predominant in certain habitats like the Mediterranean-type climate of California.

Conclusions

Despite their ephemerality, flowers are a critical component of the plant life cycle, and their physiology is intimately tied to their performance. The physiological costs of flower production and maintenance often oppose pollinator selection and, therefore, can directly impact their form and function. Here I have used a modeling approach to examine how two traits under pollinator selection, flower color and flower size, can interact with physiological traits to impact flower energy balance. Flower color can have a large impact on flower temperature, overwhelming flower size and flower hydraulics. Variation in these traits have interactive effects, showcasing how multiple optimal phenotypes are possible. In doing so, I have shown that physiologically relevant estimates of flower size can be reliably determined from regional floras, enabling new approaches to studying flower size evolution. Overall, these results suggest that centering metabolism in our study of flowers can yield new insights into the evolution of floral form and function. For example, early evidence from macroevolutionary patterns of floral physiological traits suggests that reducing the costs of flowers would have relaxed the strength of non-pollinator selection and allowed floral traits to more rapidly and more closely track pollinator preference. These results highlight the need to more fully characterize the diverse tradeoffs between pollination traits and physiological traits. Only by more fully understanding the physiological traits of flowers will we be able to more completely understand the many dimensions of their past and future evolution in response to a changing climate.

Acknowledgments

T.A. Kursar gave me an incredible opportunity and taught me most of the ecophysiological methods I still use; his impact is long-lasting. I thank N.M. Holbrook for the initial suggestion to work on flowers. T.E. Dawson, D.D. Ackerly, D.D. Baldocchi, P.V.A. Fine, K.A. Simonin, C.R. Brodersen, and S. Mambelli have provided valuable feedback and insight into thinking about plant form, function, and evolution. C.M. Guilliams, T. Lilittham, J. Farmer, V. Wormser, and T. Pham helped with data collection of flower size. J.H. Matthes helped with measurements of floral spectral. K.P. Tu has guided me in thinking about energy balance

435   considerations, among other topics. I thank the organizers of the symposium from which

436   this special issue developed, particularly Y. Sapir who gave useful suggestions on a

437   previous draft. C.J. van der Kooi, C. D. Muir, and K. Prats also provided valuable feedback on

438   earlier drafts.

References

Ashman T-L 1994 A dynamic perspective on the physiological cost of reproduction in plants. The American Naturalist 144:300–316.

Ashman T-L, DJ Schoen 1994 How long should flowers live? Nature 371:788–791.

Atamian HS, NM Creux, EA Brown, AG Garner, BK Blackman, SL Harmer 2016 Circadian regulation of sunflower heliotropism, floral orientation, and pollinator visits. Science 353:587–590.

Baldwin IT 2010 Plant volatiles. Current Biology 20:R392-R397.

Bazzaz FA, NR Chiariello, P Coley, LF Pitelka 1987 Allocating resources to reproduction and defense. BioScience 37:58–67.

Berg RL 1960 The ecological significance of correlation pleiades. Evolution 14:171–180.

Boer HJ de, MB Eppinga, MJ Wassen, SC Dekker 2012 A critical transition in leaf evolution facilitated the cretaceous angiosperm revolution. Nature Communications 3:1221.

Bond WJ 1989 The tortoise and the hare: Ecology of angiosperm dominance and gymnosperm persistence. Biological Journal of the Linnean Society 36:227–249.

Brodribb TJ, TS Feild 2010 Leaf hydraulic evolution led a surge in leaf photosynthetic capacity during early angiosperm diversification. Ecology Letters 13:175–183.

Brodribb TJ, TS Feild, GJ Jordan 2007 Leaf maximum photosynthetic rate and venation are linked by hydraulics. Plant Physiology 144:1890–1898.

Brodribb TJ, NM Holbrook, MA Zwieniecki, B Palma 2005 Leaf hydraulic capacity in ferns, conifers and angiosperms: Impacts on photosynthetic maxima. New Phytologist 165:839–846.

Caruso CM, KE Eisen, RA Martin, N Sletvold 2018 A meta-analysis of the agents of selection on floral traits. Evolution 73:4-14.

Chapotin SM, NM Holbrook, S Morse, MV Gutiérrez 2003 Water relations of tropical dry forest flowers: Pathways for water entry and the role of extracellular polysaccharides. Plant, Cell & Environment 26:623–630.

Crepet WL, KJ Niklas 2009 Darwin's second "abominable mystery": Why are there so many angiosperm species? American Journal of Botany 96:366–381.

Duursma RA, CJ Blackman, R Lopéz, NK Martin-StPaul, H Cochard, BE Medlyn 2019 On the minimum leaf conductance: Its role in models of plant water use, and ecological and environmental controls. New Phytologist 221:693–705.

Dyer AG, HM Whitney, SE Arnold, BJ Glover, L Chittka 2006 Behavioural ecology: Bees associate warmth with floral colour. Nature 442:525.

Feild TS, NC Arens 2007 The ecophysiology of early angiosperms. Plant, Cell &
Environment 30:291–309.

Feild TS, NC Arens, JA Doyle, TE Dawson, MJ Donoghue 2004 Dark and disturbed: A new
image of early angiosperm ecology. Paleobiology 30:82–107.

Feild TS, DS Chatelet, TJ Brodribb 2009a Ancestral xerophobia: A hypothesis on the whole
plant ecophysiology of early angiosperms. Geobiology 7:237–264.

Feild TS, DS Chatelet, TJ Brodribb 2009b Giant flowers of southern magnolia are hydrated
by the xylem. Plant Physiology 150:1587–1597.

Fenster CB, WS Armbruster, P Wilson, MR Dudash, JD Thomson 2004 Pollination
syndromes and floral specialization. Annual Review of Ecology, Evolution, and
Systematics 35:375–403.

Fitter A, R Fitter 2002 Rapid changes in flowering time in British plants. Science 296:1689–
1691.

Galen C 1999 Why do flowers vary? BioScience 49:631–640.

Galen C 2000 High and dry: drought stress, sex allocation trade-offs, and selection on
flower size in the alpine wildflower *Polemonium viscosum* (Polemoniaceae). The
American Naturalist 156:72-83.

Galen C, RA Sherry, AB Carroll 1999 Are flowers physiological sinks or faucets? Costs and
correlates of water use by flowers of *Polemonium viscosum*. Oecologia 118:461–470.

Givnish TJ, GJ Vermeij 1976 Sizes and shapes of liane leaves. The American Naturalist
110:743–778.

Gleason SM 2018 A blooming interest in the hydraulic traits of flowers. Plant, Cell &
Environment 41:2247–2249.

Gorton HL, TC Vogelmann 1996 Effects of epidermal cell shape and pigmentation on optical
properties of antirrhinum petals at visible and ultraviolet wavelengths. Plant
Physiology 112:879–888.

Hopkins R, MD Rausher 2012 Pollinator-mediated selection on flower color allele drives
reinforcement. Science 335:1090–1092.

Jakobsen HB, CE Olsen 1994 Influence of climatic factors on emission of flower volatiles *in*
situ. Planta 192:365–371.

Jepson Flora Project (eds.) 2015 *Jepson eFlora*. http://ucjeps.berkeley.edu/IJM.html
[accessed 2015].

Kay Q, H Daoud, C Stirton 1981 Pigment distribution, light reflection and cell structure in
petals. Botanical Journal of the Linnean Society 83:57–83.

Kevan P, H Baker 1983 Insects as flower visitors and pollinators. Annual Review of
Entomology 28:407–453.

Kevan PG 1975 Sun-tracking solar furnaces in high arctic flowers: Significance for
pollination and insects. Science 189:723–726.

Lambrecht SC 2013 Floral water costs and size variation in the highly selfing *Leptosiphon*
bicolor (Polemoniaceae). International Journal of Plant Sciences 174:74–84.

Lipayeva LI 1989 On the anatomy of petals in angiosperms. Botanicheskii Zhurnal 74:9–18.

Muchhala N, S Johnsen, SD Smith 2014 Competition for hummingbird pollination shapes
flower color variation in Andean Solanaceae. Evolution 68:2275–2286.

Muir CD 2019 tealeaves: An R package for modelling leaf temperature using energy
budgets. bioRxiv.

Niklas KJ 1994 Morphological evolution through complex domains of fitness. Proceedings
of the National Academy of Sciences 91:6772–6779.

Olson ME, Pittermann J 2019 Cheap and attractive: water relations and floral adaptation.
New Phytologist. doi: 10.1111/nph.15749

O'Meara B, S Smith, WS Armbruster, L Harder, CR Hardy, L Hileman, L Hufford, A Litt, S
Magallón, S Smith, et al. 2016 Non-equilibrium dynamics and floral trait interactions
shape extant angiosperm diversity. Proceedings of the Royal Society B: Biological
Sciences 283:20152304.

Parkhurst DF, O Loucks 1972 Optimal leaf size in relation to environment. Journal of
Ecology 60:505–537.

Parmesan C, C Yohe 2003 A globally coherent fingerprint of climate change impacts across
natural systems. Nature 421:37–42.

Patiño S, J Grace 2002 The cooling of convolvulaceous flowers in a tropical environment.
Plant, Cell & Environment 25:41–51.

Patiño S, C Jeffree, J Grace 2002 The ecological role of orientation in tropical
convolvulaceous flowers. Oecologia 130:373–379.

Peñalver E, CC Labandeira, E Barron, X Delclos, P Nel, A Nel, P Tafforeau, C Soriano 2012
Thrips pollination of mesozoic gymnosperms. Proceedings of the National Academy
of Sciences 109:8623–8628.

Peter CI, SD Johnson 2008 Mimics and magnets: The importance of color and ecological
facilitation in floral deception. Ecology 89:1583–1595.

Raupach MR, G Marland, P Ciais, C Le Quere, JG Canadell, G Klepper, CB Field 2007 Global
and regional drivers of accelerating CO_2 emissions. Proceedings of the National
Academy of Sciences 104:10288–10293.

Reekie EG, FA Bazzaz 1987a Reproductive effort in plants. 1. Carbon allocation to
reproduction. The American Naturalist 129:876–896.

——— 1987b Reproductive effort in plants. 2. Does carbon reflect the allocation of other
resources? The American Naturalist 129:897–906.

——— 1987c Reproductive effort in plants. 3. Effect of reproduction on vegetative activity.
The American Naturalist 129:907–919.

Roddy AB, CR Brodersen, TE Dawson 2016 Hydraulic conductance and the maintenance of
water balance in flowers. Plant, Cell & Environment 39:2123–2132.

Roddy AB, TE Dawson 2012 Determining the water dynamics of flowering using miniature
sap flow sensors. Acta Horticulturae 951:47–54.

Roddy AB, CM Guilliams, T Lilittham, J Farmer, V Wormser, T Pham, PVA Fine, TS Feild, TE
Dawson 2013 Uncorrelated evolution of leaf and petal venation patterns across the
angiosperm phylogeny. Journal of Experimental Botany 64:4081–4088.

Roddy AB, Jiang G.-F., K-F Cao, KA Simonin, CR Brodersen 2019 Hydraulic traits are more
diverse in flowers than in leaves. New Phytologist. doi: 10.1111/nph.15749

Roddy AB, KA Simonin, KA McCulloh, CR Brodersen, TE Dawson 2018 Water relations of
Calycanthus flowers: Hydraulic conductance, capacitance, and embolism resistance.
Plant, Cell & Environment 41:2250–2262.

Rueden C, J Schindelin, M Hiner, B DeZonia, A Walter, E Arena, K Eliceri 2017 ImageJ2:
ImageJ for the next generation of scientific image data. BMC Bioinformatics 18:529.

Sack L, P Cowan, N Jaikumar, N Holbrook 2003 The 'hydrology'of leaves: Co-ordination of
structure and function in temperate woody species. Plant, Cell & Environment
26:1343–1356.

Sagae M, N Oyama-Okubo, T Ando, E Marchesi, M Nakayama 2008 Effect of temperature on
the floral scent emission and endogenous volatile profile of *Petunia axillaris*.
Bioscience, Biotechnology, and Biochemistry 72:110–115.

Sanderson M, MJ Donoghue 1994 Shifts in diversification rate with the origin of
angiosperms. Science 264:1590–1593.

Sapir Y 2017 Pollinator-mediated selection is better detected when controlling for resource
limitation. Functional Ecology 31:7–8.

Sapir Y, M Ghara 2017 The (relative) importance of pollinator-mediated selection for
evolution of flowers. American Journal of Botany 104:1787–1789.

Schiestl FP 2015 Ecology and evolution of floral volatile-mediated information transfer in
plants. New Phytologist 206:571–577.

Scoffoni C, DS Chatelet, J Pasquet-kok, M Rawls, MJ Donoghue, EJ Edwards, L Sack 2016
Hydraulic basis for the evolution of photosynthetic productivity. Nature Plants
2:16072.

Simonin KA, AB Roddy 2018 Genome downsizing, physiological novelty, and the global
dominance of flowering plants. PLoS Biology 16:e2003706.

Spaethe J, J Tautz, L Chittka 2001 Visual constraints in foraging bumblebees: Flower size
and color affect search time and flight behavior. Proceedings of the National
Academy of Sciences 98:3898–3903.

Sprengel C 1793 Das entdeckte Geheimniss der Natur im Bau und in der Befruchtung der
Blumen. Berlin: Friedrich Vieweg dem aeltern.

Stanton ML, C Galen 1989 Consequences of flower heliotropism for reproduction in an
alpine buttercup (*Ranunculus adoneus*). Oecologia 78:477–485.

Stebbins GL 1951a Natural selection and the differentiation of angiosperm families.
Evolution 5:299–324.

——— 1951b Aridity as a stimulus to plant evolution. The American Naturalist 86:33–44.

Strauss S, J Whittall 2006 Non-pollinator agents of selection on floral traits. In: *Ecology and*
Evolution of Flowers eds. LD Harder and SCH Barrett. New York: Oxford University
Press. pp. 120–138.

Teixido AL, F Valladares 2013 Large and abundant flowers increase indirect costs of
corollas: A study of coflowering sympatric mediterranean species of contrasting
flower size. Oecologia 173:73–81.

Teixido AL, F Valladares 2014a Disproportionate carbon and water maintenance costs of
large corollas in hot mediterranean ecosystems. Perspectives in Plant Ecology,
Evolution and Systematics 16:83–92.

——— 2014b Pollinator-mediated selection does not always modulate flower size and
number in the large-flowered mediterranean shrub *Cistus ladanifer* (Cistaceae).
Botanical Journal of the Linnean Society 176:540–555.

——— 2015 Temperature-limited floral longevity in the large-flowered mediterranean
shrub *Cistus ladanifer* (Cistaceae). International Journal of Plant Sciences 176:131–
140.

Terry I, GH Walter, C Moore, R Roemer, C Hull 2007 Odor-mediated push-pull pollination in
cycads. Science 318:70–70.

Trolinder N, B McMichael, D Upchurch 1993 Water relations of cotton flower petals and
fruit. Plant, Cell & Environment 16:755–760.

van der Kooi CJ, AG Dyer, DG Stavenga 2015 Is floral iridescence a biologically relevant cue
in plant–pollinator signaling? New Phytologist 205:18–20.

van der Kooi CJ, JTM Elzenga, J Dijksterhuis, DG Stavenga 2017 Functional optics of glossy
614   buttercup flowers. Journal of the Royal Society Interface 14: 20160933.

615   van der Kooi CJ, JTM Elzenga, M Staal, DG Stavenga 2016 How to colour a flower: On the
616   optical principles of flower coloration. Proceedings of the Royal Society B
617   283:20160429.

618   Vignolini S, E Moyroud, T Hingant, H Banks, PJ Rudall, U Steiner, BJ Glover 2015 The flower
619   of *Hibiscus trionumis* both visibly and measurably iridescent. New Phytologist
620   205:97–101.

621   Vogel S 1996 Christian Konrad Sprengel's theory of the flower: The cradle of floral ecology.
622   In: Lloyd D; Barrett S, editors. Springer. pp. 44–62.

623   Warton DI, RA Duursma, DS Falster, S Taskinen 2012 smatr 3–an R package for estimation
624   and inference about allometric lines. Methods in Ecology and Evolution 3:257–259.

625   Whitney HM, M Kolle, P Andrew, L Chittka, U Steiner, BJ Glover 2009 Floral iridescence,
626   produced by diffractive optics, acts as a cue for animal pollinators. Science 323:130–
627   133.

628   Whittall J, SA Hodges 2007 Pollinator shifts drive increasingly long nectar spurs in
629   columbine flowers. Nature 447:706–709.

630   Wright IJ, PB Reich, M Westoby, DD Ackerly, Z Baruch, F Bongers, J Cavender-Bares, T
631   Chapin, JH Cornelissen, M Diemer, et al. 2004 The worldwide leaf economics
632   spectrum. Nature 428:821.

633   Wright SJ, O Calderon 1995 Phylogenetic patterns among tropical flowering phenologies.
634   Journal of Ecology 83:937–948.

635   Wright SJ, O Calderon 2006 Seasonal, El Niño and longer term changes in flower and seed
636   production in a moist tropical forest. Ecology Letters 9:35–44.

637   Wright SJ, CP Van Schaik 1994 Light and the phenology of tropical trees. The American
638   Naturalist 143:192–199.

639   Zhang F-P, MR Carins Murphy, AA Cardoso, GJ Jordan, TJ Brodribb 2018 Similar geometric
640   rules govern the distribution of veins and stomata in petals, sepals and leaves. New
641   Phytologist 219:1224–1234.

Figure legends

Figure 1. Variation in floral physiological and structural traits among extant angiosperm
clades (modified from Roddy et al. (2016) with permission from the publisher). Traits are
standardized and include whole-flower hydraulic conductance (K_{flower}), vein density,
minimum epidermal conductance to water vapor (g_{min}), stomatal pore area index (a
function of stomatal density and stomatal size), and corolla dry mass per projected surface
area. Images show stomata (left) and veins (right) for leaves and flowers of two species,
(top) *Calycanthus occidentalis* (Calycanthaceae) and (bottom) *Rhododendron*
davidsonianum (Ericaceae). Outlines and colors indicate clade affinity (grey = ANA grade,
blue = magnoliids, orange = monocots, red = eudicots). Scale bars for images of stomata are
100 μm and for images of veins are 1 mm.

Figure 2. (a) Absorptance, (b) transmittance, and (c) reflectance spectra for three
Dodecatheon sp. flowers that differ in flower color (grey lines = white flower, pink lines =
pink flower, red lines = red flower). (d) The modeled differences between between flower
temperature (T_{flower}) and air temperature (T_{air}) as a function of absorptance and the total
surface conductance to water vapor (g_t). Numbers indicate the value of the g_t isoclines
(0.05, 0.25, 0.50, 1.00 μmol m^{-2} s^{-1} Pa^{-1}), and colored points represent the values for the
three flowers whose spectra are shown in (a-c).

Figure 3. The scaling of measured flower surface area and the surface area predicted from
descriptions for 43 species in the California flora. (a) The measured surface area is the
total surface area of the calyx and corolla structures. (b) The measured surface area of only
the corolla structures. In both figures, point colors correspond to clade membership (blue
= magnoliids, orange = monocots, red = eudicots), the solid line is the 1:1 relationship, the
dashed line and the grey shading represent the estimated standard major axis regression
and its 95% confidence intervals.

Figure 4. (a) Histogram of the two-dimensional projected surface area of flowers from the
California flora (n = 43 species). (b) The modeled differences between flower temperature
(T_{flower}) and air temperature (T_{air}) as a function of flower size and the total surface
conductance to water vapor (g_t). Numbers indicate the value of the g_t isoclines (0.05, 0.50,
1.00 μmol m^{-2} s^{-1} Pa^{-1}).