Evolutionary dynamics of *de novo* mutations and mutant lineages arising in a simple, constant

environment

Margie Kinnersley[1,¶], Katja Schwartz[3,¶], Jacob Boswell[2], Dong-Dong Yang[2], Gavin Sherlock[3,*]

Frank Rosenzweig[1,2,*]

[1] Division of Biological Sciences, The University of Montana, Missoula, MT 59812

[2] School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA 30332

[3] Department of Genetics, Stanford University School of Medicine, Stanford, CA 94305-5120

[¶] These authors contributed equally to this work.

[*] Co-corresponding authors: frank.rosenzweig@biology.gatech.edu and gsherloc@stanford.edu

Keywords: *E. coli,* adaptive evolution, chemostat, metagenomics, population sequencing, clone

sequencing, clonal interference, parallelism

Running Title: Parallelism and clonal interference in evolving bacterial populations

Abstract

A large, asexual population founded by a single clone evolves into a population teeming with

many, whether or not its environment is structured, and whether or not resource levels are constant

or fluctuating. The maintenance of genetic complexity in such populations has been attributed to

balancing selection, or to either clonal interference or clonal reinforcement, arising from

antagonistic or synergistic interactions, respectively. To distinguish among these possibilities, to

identify targets of selection and establish when and how often they are hit, as well as to gain insight

into how *de novo* mutations interact, we carried out 300-500 generation glucose-limited chemostat

experiments founded by an *E. coli* mutator. To discover all *de novo* mutations reaching ≥1%

frequency, we performed whole-genome, whole-population sequencing at ~1000X-coverage every

50 generations. To establish linkage relationships among these mutations and depict the dynamics

of evolving lineages we sequenced the genomes of 96 clones from each population when allelic

diversity was greatest. Operon-specific mutations that enhance glucose uptake arose to high

frequency first, followed by global regulatory mutations. Late-arising mutations were related to

energy conservation as well as to mitigating pleiotropic effects wrought by earlier regulatory

changes. We discovered extensive polymorphism at relatively few loci, with identical mutations

arising independently in different lineages, both between and within replicate populations. Out of

more than 3,000 SNPs detected in nearly 1,800 genes or intergenic regions, only 17 reached a

frequency ≥ 98%, indicating that the evolutionary dynamics of adaptive lineages was dominated

by clonal interference. Finally, our data show that even when mutational input is increased by an

ancestral defect in DNA repair, the spectrum of beneficial mutations that reach high frequency in

a simple, constant resource-limited environment is narrow, resulting in extreme parallelism where

many adaptive mutations arise but few ever go to fixation.

Author Summary

Microbial evolution experiments open a window on the tempo and dynamics of evolutionary change in asexual populations. High-throughput sequencing can be used to catalog *de novo* mutations, determine in which lineages they arise, and assess allelic interactions by tracking the fate of those lineages. This *adaptive genetics* approach makes it possible to discover whether clonal interactions are antagonistic or synergistic, and complements genetic screens of induced deleterious/loss-of-function mutants. We carried out glucose-limited chemostat experiments founded by an *E. coli* mutator and performed whole-genome, whole-population sequencing on 300-500 generation evolutions, cataloging 3,346 *de novo* mutations that reached ≥1% frequency. Mutations enhancing glucose uptake rose to high frequency first, followed by global regulatory changes that modulate growth rate and limiting resource assimilation, then by mutations that favor energy conservation or mitigate pleiotropic effects of earlier regulatory changes. We discovered that a few loci were highly polymorphic, with identical mutations arising independently in different lineages, both between and within replicate populations. Thus, when mutational input is increased by an ancestral defect in DNA repair, the spectrum of beneficial mutations that arises under constant resource-limitation is narrow, resulting in extreme parallelism where many adaptive mutations arise but few ever become fixed.

Introduction

Evolution experiments using microbes have enlarged our understanding of the tempo and dynamics of evolutionary change, as well as how selection, drift and historical contingency influence evolutionary trajectories. Combined with high throughput sequencing, experimental microbial evolution (EME) can now be used to identify substantial numbers of *de novo* beneficial mutations in laboratory populations, to determine in which lineages they arise and the fate of those lineages, and to evaluate the sign and strength of possible epistatic interactions [1-3]. This approach, adaptive genetics, based on analyzing cohorts of spontaneous beneficial mutations to determine how their frequencies fluctuate over time, constitutes a mode of inquiry that complements traditional genetic screening of induced deleterious/loss-of-function mutants (e.g., [4] and [5] among others). Adaptive genetics also expands the possibilities for discovering constraints on protein structure and function and for discerning the architecture and malleability of networks that regulate nutrient-sensing and cell division.

Microbial populations were once thought to evolve by periodic selection as a succession of adaptive clones, each fitter than its antecedent, replacing one another over time [6-9]. This model was consistent with Muller and Haldane's view of how beneficial mutations spread in large asexual populations [10-12] under conditions governed by competitive exclusion [13]. Today we know that large, initially clonal populations rapidly accumulate and retain genetic variation, much of which is beneficial [14-18]. In fact, the amount of adaptive genetic variation observed in EME populations can be enormous, owing large population sizes with a continuous input of beneficial mutations and the subsequent competition among new adaptive lineages, which gives rise to clonal interference [14,16,19,20].

Clonal interference can occur within a larger framework of stable subpopulation structure

[21] when microbial lineages come under balancing selection [22-25] or specialize to exploit

niches created either by the culture conditions [23,26,27], or by the organisms themselves [28-30].

In a simple constant environment like a chemostat the persistence of subpopulations likely depends

on founder genotype, the emergence of specific key mutations, and availability of the limiting

nutrient [31]. Ferenci and colleagues never observed stable subpopulation structure in glucose-

limited evolutions originating from *E. coli* K12 strain BW2952 [32], whereas Adams and

colleagues, using a different strain, often did [30,33]. Unlike BW2952, the K12-derived ancestor

used by Adams, JA122 [30] harbors a supE44 *glnX* tRNA nonsense suppressor as well as nonsense

mutations in housekeeping and stationary-phase transcription factors, RpoD and RpoS

respectively, and mismatch repair enzyme MutY. The JA122 ancestor's defect in DNA repair

increases mutational load on its descendants [28,34], while the nonsense suppressor mitigates the

effect of mutations that create premature stop codons. Such a suppressor would likely make the

blunt instrument of *de novo* nonsense mutations a less effective agent of adaptive change, possibly

resulting in a more nuanced spectrum of beneficial mutations than would otherwise occur among

mutators.

To understand the impact that a mutator/suppressor founder has on the spectrum and fate

of new beneficial mutations, and on the dynamics of population structure, we repeated Adams *et*

al. classic evolution experiments using the same ancestral strain and culture conditions [30]. We

monitored, at 50-generation intervals, the incidence of mutations that reached at least 1%

frequency over the course of 300-500 generations, identifying mutations that were either

transiently beneficial or hitch-hiking with mutations that were. To determine which mutations co-

occurred within a given lineage we sequenced 96 clones from each population at the time-point

where we observed greatest allelic diversity. We uncovered no evidence for stable sub-population

structure, but instead saw pervasive clonal interference, with only 17 out of 3,346 mutations going

to near fixation across replicate experiments. The temporal order in which certain mutations rose

to high frequency was predictable, reflecting a high degree of parallelism both within and between

replicates. In general, mutations that enhanced glucose assimilation arose early, followed by

mutations in global regulators and mutations that either increase efficiency of limiting resource

utilization or mitigate the deleterious effects of certain earlier mutations. Altogether, our results

show that even in bacterial populations founded by an ancestor having a high mutation rate and

the capacity to tolerate many *de novo* mutations, the spectrum of genomic changes that rise to

appreciable frequency and the adaptive outcome of replicate evolutions are limited when those

populations evolve in a simple constant environment.

Results

Experimental design. Evolution experiments were carried out in triplicate under continuous

nutrient limitation using Davis Minimal Medium [30], with glucose (0.0125% w/v) as the sole

source of carbon for energy and growth. Chemostats (300 mL working volume) were run under

aerobic conditions for 300-500 generations at constant temperature (30°C) and at constant dilution

rate (D=0.2 hr^{-1}). Under these conditions, population density reaches ~10^8 cells mL^{-1} at steady

state. The *E. coli* strain used to initiate these experiments, JA122, is distinguished from *E. coli*

K12 by alleles likely to influence the spectrum of mutations arising during adaptive evolution

(**Table S1**; [28]). Among these is a nonsense mutation in MutY (Leu299*) that results in a 10-fold

greater mutation rate and GC→TA transversion bias [28], nonsense mutations in the genes that

encode stationary phase sigma factor RpoS (Gln33*) [35] and 'housekeeping' sigma factor RpoD

(Glu26*), as well as a suppressor mutation in the *glnX* tRNA known to suppress amber, ochre and

opal mutations (**Table S1)** [36].

To identify the mutations that arose during the evolutions, we performed whole genome, whole

population sequencing every 50 generations on each of the three chemostat populations. We

generated approximately 50 million 2x100bp paired end reads per sample, yielding coverage of up

to ~1000x for each time point (inserts were selected to be short enough such that forward and

reverse reads overlapped, which while reducing coverage, increases quality; see Methods). Based

on this level of coverage, we were able to identify mutations that rose to an allele frequency of

~1% of greater. Given an effective population size of $>10^{10}$ and 300-500 generations of selection

it is highly improbable that any allele could reach such a frequency by drift alone [23]. We can

therefore assume that every mutation recovered was either under positive selection or hitch-hiking

along with one that was.

Population sequencing shows general patterns of mutation that are consistent across

independent evolutions. Across all samples, 3,326 SNPs were detected in 2,083 unique genes or

intergenic regions (**File S1**). The overwhelming majority (97.5%) of these SNPs were GC→TA

transversions, as expected given the ancestral strain's defect in the mismatch repair protein MutY,

which encodes adenine glycosylase [37]. Consistent with the protein coding density of *E. coli*

(87.8%) [38], 85% (2,854) of SNPs occurred in coding regions. On average, 69.2% of these created

a missense mutation, 23.4% resulted in a synonymous mutation and 7.4% caused a nonsense

mutation (**Fig. 1**). Relative to proportions observed in mutation accumulation experiments carried

out using wild-type *E. coli* [39], we observed more nonsynonymous and nonsense mutations.

Small deletions were rarely detected (one single-nucleotide deletion in each of vessel 1 and vessel

2, and none detected in vessel 3), but we observed a single large ~150kb duplication in vessel 2.

The overall number of mutations in each population increased linearly over time and at

approximately the same rate across replicates (**Fig. 1**), as would be expected with a mutator

phenotype.

Comparison of population level mutations reveals clonal interference and widespread

parallelism. Despite the large number of SNPs detected, only 17 alleles arose above a frequency

of 98% across replicate evolutions ranging from 300-500 generations. Moreover, the maximum

frequency of most alleles never exceeded 10% (**Fig. 2A**), and the vast majority of alleles were

present at lower frequency in the final time-point than they were at their maximum (**Fig. 2B**).

Together, the foregoing observations suggest that in each evolution experiment population

dynamics was largely driven by clonal interference [40]. A small number of loci were recurrently

mutated above what would be expected by chance, indicating that variants at these loci were likely

beneficial (**Table 1**, **Table S2**). For example, a total of 212 mutations arose in the 10 most

significantly mutated genes in the population sequencing data, with each gene receiving at least

five mutations (**Table 1**). Moreover, 30 and 14 distinct allelic variants were discovered in just two:

the genes encoding the DNA binding repressor GalS and the RNA-binding protein Hfq,

respectively (**Table S3**). High-resolution population sequencing also revealed that 13 SNPs not

present at the start of the experiment reached at least 1% frequency in all three vessels at various

time-points, while 52 SNPs recurred in two out of three chemostats (**Table S4**). Thus, our data

also provide compelling evidence for substantial parallel evolution at the genic level.

Clonal sequencing further clarifies lineage relationships and parallelism To determine

linkage relationships between the novel alleles, we sequenced 96 individual clones from each

vessel. In each case, the 96 clones were isolated at random from the time-point at which we

detected the greatest number of mutant alleles at $\geq$ 5% frequency. To assess whether the frequency

estimates from population sequencing were reasonable, and whether the isolated clones constituted

a reasonable subsample, we compared frequencies of mutations identified in both datasets at the

corresponding time-point and found that they correlate well (**Fig. 3**).

For each set of 96 clones, we constructed a phylogeny to represent their putative

evolutionary relationships (**Fig. 4**). Inspection of the mutations and trees from each vessel (i.e.

each independent evolution) revealed several instances in which exactly the same mutation arose

not only in different vessels, but often more than once in the same vessel on distinct branches of a

given tree. In the most extreme case, 6 of the 11 *hfq* alleles detected via clone sequencing were

identified in clones from different vessels, indicating independent parallel origins (**Fig. 4, File S2**).

Furthermore, 7 of the 11 appear to have arisen more than once within the same vessel.

Clonal dynamics are shaped by relationships among de novo alleles, hard and soft selective

sweeps, and absence of periodic selection Combining population allele frequency data with

linkage information derived from clonal sequencing makes it possible to depict lineage dynamics

using Muller diagrams (**Fig. 5, Files S3-S5**). In general, we observe early, hard sweeps of highly

beneficial mutations related to limiting nutrient influx, followed by soft sweeps [41-43] and

multiple-origin soft sweeps that may fine-tune glucose uptake or utilization later in the experiment

when diversity was higher [44-46]. Hard sweeps consistently involved mutations in regulators

(*galS* in chemostat 1, transcriptional terminator *rho* in chemostats 1 and 3) or regulatory regions

(upstream of *dnaG* in chemostat 1, upstream of *mglB* in chemostats 1, 2 and 3 – See supplementary

Files 2, 3 and 4 for detailed dynamics), while soft sweeps were comprised of both regulatory and

operon-specific mutations (e.g. *hfq* and *opgH* in chemostats 1, 2 and 3, upstream of *adhE* in

chemostat 1, *pgi* in chemostat 3) (**Fig. 5, S1, Files S3-S5)** [42,47]. Here, we note that multiple-

origin soft sweeps may be especially prevalent in our experiments due to the ancestral mutator

allele at *mutY*, as the likelihood of concurrent identical mutations in the same gene should increase

with mutation rate.

With regard to periodic selection, rather than favorable alleles arising within a set of lineages

that successively replace one another over time, we observe groups or cohorts of mutations co-

evolving, with widespread clonal interference among lineages that carry different beneficial

mutations [48]. For example, in chemostat 1, a spreading lineage with a cohort of mutations

upstream of *mglB*/*lptA*/*opgH* (pink) is checked by the emergence of lineages carrying mutations

in *hfq* (green) (**File S3**). All of these phenomena – hard and soft sweeps, cohorts of mutations that

increase or decrease in frequency together, and clonal interference – have been observed in yeast

[14,19,49] and *E. coli* [23] populations evolving in the laboratory, as well as in *Pseudomonas*

aeruginosa evolving in the cystic fibrosis lung [50].

Early sweeps are related to influx of the limiting nutrient glucose. For specific growth rates

between $\sim \mu =0.1$ hr^{-1} and $\mu =0.9$ hr^{-1}, glucose is most efficiently transported using a combination

of the maltoporin LamB and the galactose transporter MglBAC, and glucose limitation frequently

selects for mutations that increase expression of these proteins [45,51-59]. As expected, 7 of the

top 10 frequently mutated genes/gene regions we observed (*galS*, upstream *mglB*, *malT*, *malK*,

hfq, *rho* and upstream *dnaG*) play a role in transcriptional regulation of *lamB* or *mglBAC*, either

directly or through their interactions with global regulators (**Table 1**, **Fig. 6**).

MUTATIONS IN *GALS* AND UPSTREAM OF *MGLB*. Thirty different alleles of *galS* (encoding GalS,

a negative regulator of *mglBAC* transcription) were detected over the course of our experiments.

These spanned the length of the gene, and the majority caused missense amino acid changes likely

to disrupt *mglBAC* transcriptional repression and augment glucose flux across the inner membrane

(**Fig. 6**) [60]. Despite the large number of alleles we observed, few persisted beyond generation 50

or attained a final frequency greater than 5%, demonstrating high clonal diversity early in the

experiment. In chemostats 2 and 3, no clear "winner" *galS* genotype emerged, though in chemostat

1, a GalS allele (Arg146Leu) swept to near fixation (89.6% of the population at generation 50).

Early-arising *GalS* mutant genotypes were rapidly displaced by clones carrying highly-

beneficial mutations in the *mgl* operator sequence upstream of *mglB* (**Fig. 5**). This sequence of

events, like the mutations themselves, has been observed elsewhere [45,61]. The most successful

mutation upstream of *mglB* (bp 2,238,647 C→A) occurred early in every vessel, and in every case

increased in frequency to over 90% of the population (**Table S5**, **File S3-S5**). Notably, this same

mutation was observed in chemostat-grown *E. coli* by Notley-McRobb et al. (1999) as well as in

the experimental population described by Helling et al. (1987), where it was found to be the only

SNP shared by all members of a cross-feeding consortium [34,54].

The dynamics of *galS* replacement illustrates the effect that clonal interference can have

on the fate of different alleles. In chemostat 1, clones carrying GalS Arg146Leu rapidly dropped

in frequency when lineages emerged with a mutation upstream of *mglB* (position 2,238,647), but

were not completely displaced until generation 400 and even enjoyed brief periods of expansion.

In chemostat 2, clones with the same mutation upstream of *mglB* were present by generation 50,

but did not surpass a 90% threshold for another 250 generations due to competition from 22

different *galS* lineages and a lineage carrying a different upstream of *mglB* allele (2,238,648 G→T)

(**Fig. 5, S1B, File S4**). By contrast, in chemostat 3, a lineage with the upstream *mglB* mutation

(2,238,647) experienced little competition and was almost fixed by generation 150 **(Table S5).**

Over the remainder of the experiment only three other mutations upstream of *mglB*

mutations reached the threshold for detection: two were within 2 base-pairs of the first mutation

and did not rise to high frequency, while the third (chemostat 1, 2,238,630 C→A) located in the

CRP activator binding site, co-occurred with 2,238,647 C→A and increased to ca. 80% frequency

by generation 500 (**Figs. 5, S1A, File S3**). This dynamic suggests additional mutations that affect

GalS repressor binding are not of great benefit after the preferred allele has swept the population,

whereas mutations that modulate the activity of other regulators (i.e. CRP) can act synergistically.

THE DYNAMICS OF LAMB REGULATION. LamB glycoporin overexpression is a hallmark

feature of *E. coli* populations adapted to glucose-limited chemostat growth [34,45,53,55,56,62].

Previous experiments have shown that under glucose limitation, overexpression of LamB can be

the result of any one of the following: constitutive activation of transcriptional regulator MalT,

disruption of the MalT inhibitor MalK, mutation of the RNA chaperone Hfq, alteration of sigma

factor dynamics (σ^S/σ^D ratio), or mutation of the *malT* repressor Mlc, (**Fig. 6**) [34,45,53-56,62,63].

Across the three replicate evolutions, we observed 19 unique *malT* alleles and 14 unique

malK alleles (**Figs. 6, S2, Table S3**). Over half of the mutations in *malT* (10 out of 19) are known

either to cause MalT to become constitutively active, or to occur in amino acids involved in

MalT/MalK interaction [45,64,65]. A single MalK mutation (Ala296Asp) rose to high frequency

early (94% by generation 100) in chemostat 1 (**Fig. S2**, **Table S5**). This SNP is in a regulatory

domain likely to be the site of MalK/MalT interaction [66]. Alteration of a neighboring residue

(Asp297) has been previously shown to allow unregulated transcription of the *mal* operon [66].

malT/*malK* allele dynamics differ among experimental populations. As mentioned above

and shown in **Fig. S2**, MalK Ala296Asp sweeps early in chemostat 1, whereas in chemostat 3,

early MalK mutations (blue) are displaced by later mutations in MalT (green). In chemostat 2, the

picture is quite different: clones with either *malK* or *malT* mutations co-exist through all 500

generations. The reason for this contrast in allele frequency dynamics cannot be attributed to

emergence of a single "most fit" allele, as the majority types from chemostats 1 and 3 arose

independently in chemostat 2, but did not sweep. Despite the importance of MalT and MalK as

high-value targets for selection during adaptation to glucose limitation, other advantageous

mutations (upstream *mglB*, *rho* and *hfq*, discussed below) may have ultimately carried "winning"

mal alleles in chemostats 1 and 3 to higher frequency, purging allelic diversity at this locus.

Interestingly, although we observed 30 *malT* and 22 *malK* mutations in the population sequencing

data (**Table 1**), in only 5 out of the 288 sequenced clones do mutant alleles of these two genes co-

occur, suggesting that there may be no additional advantage or even some disadvantage to having

both. In the Helling et al. evolution experiments [30], which were founded by the same ancestor

used here, secondary resource specialists share a mutation in MalT, whereas the primary resource

(glucose) specialist that feeds those clones carries a mutation in MalK [34].

SELECTION OF MUTATIONS IN RNA CHAPERONE HFQ THAT AFFECT TRANSLATION OF LAMB

AND STATIONARY PHASE TRANSCRIPTION FACTOR RPOS. Hfq is a global regulatory protein that

facilitates translation and/or RNA degradation by mediating ncRNA-mRNA interactions. It

participates in a diverse range of cellular processes including nutrient uptake, motility and

metabolism [67]. *hfq* mutations identified in other glucose-limited evolution experiments exhibit

pleiotropic physiological effects: they appear to increase translation of LamB glycoporin, reduce

levels of stationary phase transcription factor RpoS, inhibit cellular aggregation, and enhance

glucose transport via PtsG [63,68].

Hfq is one of the most frequently mutated genes observed in our experiments: 24 *hfq*

mutations, resulting in 14 distinct *hfq* alleles, were detected via population sequencing; by the end

of our evolutions >50% of each population carried a mutation in *hfq* (**Table S3**). (**Table 1**). Two

of these alleles arose independently in all three vessels (same nucleotide position, same SNP), and

six additional alleles were observed in two of three vessels (**Table S4**). The frequency of and

parallelism exhibited in *hfq* mutations is particularly curious in the context of experiments by

Maharjan et al. in which *hfq* mutations arise, but are at low frequency and subject to negative

frequency-dependent selection and epistatic interaction with mutations in *rpoS* [32,52,62,63].

The dynamics of hfq mutations are variable across evolutions and may depend on which

other beneficial alleles are present in the same lineage or in the same population. In chemostats 1

and 2, a large number of *hfq* alleles (10 in chemostat 1 and 11 in chemostat 2) appear after

generation 250 and are preceded by mutations in *malK* or *malE* and the *opg* operon. The most

successful Hfq allele in chemostat 1, Val62Phe, occurred in a sweeping lineage with a secondary

mutation upstream of *mglB* (discussed above) and may have been carried along by association. In

chemostat 3, a single *hfq* mutation arises early (Ser60Tyr, present by generation 100), sweeps to

near fixation alongside MalT Met311Ile and is closely followed by mutations in *opgH* (**Fig. S1,**

File S5).

RECURRENT MUTATIONS OCCUR IN RHO. Early-arising mutations in the *rho* termination factor

are a conspicuous feature of chemostats 1 and 3 (**Figs. 5, S1**). Rho is required for transcriptional

termination of up to 50% of cellular mRNAs [69,70] and can participate in gene regulation via

intragenic terminators [71]. Mutagenesis and ChIP-chip analyses have identified Rho-dependent

terminators within multiple genes relevant to glucose limitation, specifically *lamB*, *mglA*, and

mglC and downstream of *malT* and *mglC* [71,72]. In fact, it has long been known that defective

LamB expression in MalT activator mutants can be restored via compensatory mutations in *rho*

[73]. In chemostats 1 and 3, Rho mutations fix or nearly fix early and do so in concert with

mutations in MalK (chemostat 1 Ala296Asp) and mutations upstream of *mglB* (chemostats 1 and

3, bp 2,238,647) (**Figs. 5, S1, Table S3, Table S5, Supplementary Files 2 and 4**). Conversely, in

chemostat 2 only three *rho* alleles were detected, none of which rose in frequency to >6% of the

population (**Fig. S1, Table S3**).

Mutations that impact energy conservation, membrane biogenesis and cell adhesion are

late arising targets of selection PHOSPHOGLUCOSEISOMERASE (PGI) is an abundantly expressed

central metabolic enzyme responsible for converting glucose-6-phosphate into fructose-6-

phosphate. Knockdown of *pgi* mRNA alleviates catabolite repression [74], favoring increased

expression of CRP-regulated genes such as *lamB* and *mglBAC*. Twenty-four unique *pgi* alleles

were detected over the course of our three replicate evolutions. However, few rose to appreciable

frequency before generation 200, suggesting their benefit may be contingent on the presence of

other mutations or some aspect of the chemostat environment that consistently changed after this

time point. *Pgi* alleles were least successful in chemostat 1, which was also the only replicate in

which a large fraction of clones (79% by generation 500) acquired a second mutation upstream of

mglB. This observation suggests that *pgi* mutations and mutations in the CRP-binding site of the

mglBAC promoter may be functionally redundant.

MEMBRANE GLYCOSYLTRANSFERASE OPGH is involved in the synthesis of periplasmic glucans,

highly branched oligosaccharides made from β-linked glucose monomers. While we do not

observe *opgH* mutations earlier than generation 100, they rapidly increase in frequency once they

appear, usually either just before or just after *hfq* mutations (**Fig. S1, S4, File S3-S5**). Novel *opgH*

alleles, especially the nonsense mutations that we frequently observe, may constrain glucan

production and serve as a glucose conservation measure. A "moonlighting" function has also

recently been reported for OpgH: the glucosyltransferase interacts with the tubulin-like cell

division protein FtsZ to delay cell division when levels of UDP-glucose are low [75]. Thus,

mutations in OpgH may augment the rate of cell division, and thereby provide a fitness advantage

under slow-growth chemostat conditions. The only *opg* operon mutation identified among strains

in previous Adams et al. experiments occurred in *opgG* of the glucose scavenger, CV103 (E487*)

[28].

MUTATIONS IN RHO-INDEPENDENT TERMINATOR T_1 THAT ALLOW RUN-THROUGH

TRANSCRIPTION MAY TIP THE BALANCE BETWEEN COMPETING SIGMA FACTORS. Sigma factor RpoD

(σ^{70}) is the predominant sigma factor associated with RNA polymerase during exponential growth.

As cells enter stationary phase, transcription of the gene for alternate sigma factor RpoS (σ^S)

increases [76]. *rpoS* mutations are often selected for under continuous glucose limitation as they

allow continued transcription from promoters negatively regulated by σ^S but required for glucose

uptake and metabolism (e.g. [77,78]).

In chemostat 1, a mutation in the *rpsU-dnaG-rpoD* macromolecular synthesis operon

upstream of *dnaG* (bp 3,209,081 G→T) was present in over 90% of the population by generation

50 (**Table S5**). This SNP decreases the stability of the rho-independent terminator T_1 situated

between *rpsU* and *dnaG*, and thus may be expected to increase expression of RpoD [79] and as a

result operons positively controlled by σ^{70} (e.g. *mglBAC* and *malK-lamB-malM*). A T1 mutation

(bp 3,209,075 C→A) was also shared among Helling et al. strains, defining the lineage that gave

rise to three of four consortium members [30,34]. In addition, in Chemostat 2, we observed an

~150kb duplication that included *rpoD* and in chemostat 3, eight clones out of 96 carried intragenic

suppressor mutations of the ancestral nonsense allele (*26Asp and *26Gln) in RpoD.

FIMBRIAL PROTEIN GENES (*FIM*) Genes associated with production/function of type 1 fimbriae,

particularly *fimH* (fimbrial adhesion), were an unexpected and frequent target of mutation in all

three chemostats (**Table 1**, **Figs. 4**, **S1, Table S3, Files S3-S5**). Though novel *fim* alleles were

transient in vessels 2 and 3, in chemostat 1 a FimH Asn54Lys variant rose to a frequency of 70%

by generation 150, temporarily displacing high-fitness alleles in *rho*, *malK* and upstream *mglB*

(**File S3**). Because *fimH* mutants demonstrated an increased capacity for biofilm formation (data

not presented), a recurrent issue in chemostat experiments, but did not acquire any of the mutations

expected to enhance glucose metabolism, *fimH* mutations were likely related to chemostat

persistence rather than to competition for limiting substrate.

Discussion and Conclusions

History matters: ancestry influences evolutionary trajectory The tempo and trajectory of a clonal

population depend on its genetic point of departure. Our departure point was a founder that

harbored nonsense mutations in mismatch repair (MutY, Leu299*), and in housekeeping and

stationary phase sigma factors (RpoD, Glu26* and RpoS, Gln33*), but also carried an

amber/ochre/opal nonsense tRNA suppressor. Populations originating from such a founder would

not only have an increased mutational load but also the capacity to tolerate those mutations, in

particular nonsense mutations that would otherwise result in complete loss-of-function.

Laboratory evolution studies have borne out the idea that loss-of-function mutations can

be significant drivers of adaptation [20,80-82]. Metabolic network re-programming via modulation

of existing function can occur much faster than the evolution of new pathways via mutation [81],

and in many cases nonsense mutations or deletions confer greater fitness benefit than missense

mutations affecting the same gene [82]. However, loss of function often comes at the expense of

metabolic flexibility, limiting the ability of evolved clones to compete in alternative environments

[20]. RpoS has been shown to be a high-value target of selection under nutrient limitation: under

low-nutrient conditions RpoS normally outcompetes RpoD for binding to RNA polymerase,

repressing genes required for growth and cell division and activating those required to enter

stationary phase [78,83]. *rpoS* mutants thus continue to divide under conditions where wild-type

cells arrest. In this respect, our genetic 'point of departure' could be viewed as being pre-adapted

to life under glucose limitation. However, the combined phenotypic effect of ancestral *rpoS* and

rpoD nonsense mutations in a suppressor background is murky and raises the question of whether

this combination of mutations is favorable under glucose limitation, merely tolerated or

detrimental. Despite the fact that many changes we observed (*galS*, upstream *mglB*, *hfq*) enhance

glucose assimilation, are predictable, occur repeatedly and rise to high frequency, we also saw the

persistence of clones with none of these mutations that instead carry intragenic suppressors of the

nonsense mutation in *rpoD* (*26→Asp and *26→Gln, chemostat 3 **Fig. 4**) or a duplication that

includes *rpoD*. If rapid adaptation can be driven by loss-of-function but occurs at the expense of

metabolic flexibility, nonsense mutations have a distinct advantage over deletions in that reversion

or suppression is possible should environmental conditions change [20].

395   Another ancestral allele that we expected to influence evolutionary trajectories was an

A→T CRP binding site mutation 224 bp upstream of the acetate scavenging enzyme, *acs* (acetyl-

CoA synthetase). This mutation alters regulation of the *acs-pta* operon such that the ancestor

poorly assimilates acetate excreted during growth under continuous glucose-limitation, opening

up a secondary resource for novel mutants that can [34]. Here, we uncovered no evidence for the

type of cross-feeding described in prior reports [29,30,33]. This result was not unanticipated, as

evidence for cross-feeding polymorphisms was observed in only half the evolution experiments

founded by this ancestor or its close relatives [33]. Moreover, a recent model [31] defining the

boundary conditions for cross-feeding to evolve in a chemostat showed that such an outcome is

sensitive to variation in dilution rate as well as to the relative fitness of *de novo* mutants that gain

access to secondary metabolites. Subtle differences in either of these parameters may account for

why we saw no evidence for acetate/glycerol/formate cross-feeding in our experiments. The

absence of such interactions may also be due to the fact that no variants arose at loci where

mutations have been implicated in cross-feeding evolution: *acs* (acetyl CoA synthetase)*, lpd*

(lipoamide dehydrogenase) and *ptsI* (phosphoenolpyruvate phosphotransferase).

As expected, the ancestral *acs-pta* defect resulted in appreciable levels of residual acetate

(~45-90 μM) at the onset of our experiments (**Fig. S3**). While we uncovered no evidence for the

evolution of secondary resource specialists [30,31,33] and refs therein), residual acetate levels

consistently fell below detection limit by generation 200. Thus, adaptive mutants arising here

found other ways than cross-feeding to metabolize all available carbon. One possible work-around

may involve the *pgi* locus, which was second only to *galS* in the total number of mutations

recovered (**Table S2**). Generation 200 coincides with the emergence of mutant *pgi* alleles in all

three populations. In chemostat experiments with *pgi* deletion mutants, Yao et al. found that in the

absence of Pgi, glucose uptake rate drops slightly compared to wild-type, but no overflow acetate

is produced and biomass yield is unchanged [84].

Population and clone sequencing open up a detailed view of the full spectrum of

beneficial mutations and how that spectrum changes over time High-coverage, whole-genome,

whole-population sequencing makes it possible to discover every new allele reaching ≥1%

frequency in a population of $>10^{10}$ cells. Because alleles are highly unlikely to reach such

frequencies by draft, all were either transiently beneficial or hitchhiking with alleles that were.

This depth of analysis opens up a richly-detailed view of the spectrum of beneficial mutations

arising in *E. coli* under constant resource limitation. Periodic whole-population sequencing allows

patterns to be discerned as to how these spectra change over time, while clone sequencing makes

it possible to establish linkage relations among novel alleles and represent their collective fate as

evolving lineages. Multiple patterns emerge from these analyses. *First,* new alleles accumulate in

replicate populations at similar rates, and the proportion of alleles that are missense, nonsense,

synonymous, or noncoding remains fairly constant; the great majority is either missense (60-70%)

or nonsense (5-10%). *Second,* the distribution of new mutations across the genome is skewed, with

only a few dozen of the more than 1,000 mutated genes having a significant number of mutations;

yet even among these most frequently mutated genes, few *de novo* mutations fix. *Third,* by clonal

sequencing we are able to determine that many, independent linages co-exist and compete within

the culture. Thus, evolutionary dynamics in these populations is governed by clonal interference

rather than by clonal replacement or reinforcement.

A *fourth* pattern to emerge is widespread parallelism in regulatory evolution. Both across

and within populations, the same genes are mutated again and again, often at exactly the same

nucleotide position in independent replicates, and sometimes in independent lineages co-evolving

in the same vessel. Many of these genes (*galS*, *malT*, *malK*, upstream *mglB*, *hfq*, *rho*) act in

processes related to the transport and assimilation of the limiting nutrient, glucose. However, in

most cases the mutations recovered alter regulation of these processes, and not the structural

proteins that carry them out.

A *fifth* pattern relates to the order of beneficial mutations and the influence that order has

on dynamics. Consistent with previous reports, mutations that increase glucose flux across the

inner membrane (*galS*, upstream *mglB*) occur early and precede those that increase flux across the

outer membrane (*malK*/*malT*, *hfq*, *rho*). In both cases, mutations in binding partners

(GalS/upstream *mglB* and MalT/MalK) rarely occur in the same clone, and the order in which they

occur can lead to either a sweep (upstream *mglB* clones quickly displace *galS* clones) or clonal

interference (*malT* and *malK* clones can co-exist). Other alleles appear to emerge later in the

experiment and nearly always together: clones with existing mutations in the *mal* operon acquire

subsequent mutations in *hfq* and *opgH*, regardless of which gene is altered first and which alleles

are already present in the population **(Fig. S1)**. These patterns are echoed by the genotypes reported

by Kinnersley et al. [22] in which glucose scavenger CV103 has mutations in *malK*, *opgG* and *hfq*

while acetate specialist CV101 only carries a mutation in *malT*.

457   Similar experiments carried out by Maharjan et al. [32] using *E. coli* BW2952

demonstrated that under continuous glucose limitation population-level phenotypic changes are

often the result of multiple soft sweeps by combinations of beneficial mutations. While we did not

assay clone phenotypes, multiple alleles of *galS*, *hfq* and *opgH* appear to sweep our populations in

concert suggesting a similar pattern in which a phenotypic effect (reduced expression of a

particular gene) is favored, but has different genetic bases in co-existing lineages. At the clone

level, BW2952 also exhibits sign epistasis between mutations in *rpoS*/*hfq* and *galS*/*malT* [32,52].

In our experiments, we did not uncover evidence of sign epistasis between the ancestral *rpoS* allele

and *hfq*: by generation 250, over 50% of clones in populations 1 and 3 carry mutations in both

genes. Maharjan et al (2013) proposed that fitness deficits exhibited by *rpoS*/*hfq* double mutants

may be the result of altered cell division [62,85] and that *hfq* mutations enhance glucose uptake

during slow growth, but diminish viability when cells are dividing rapidly. Hfq deletion mutants

exhibit cell division anomalies due to elevated expression of cell division proteins, including FtsZ

[86,87]. Recent work by Hill et al. (2013) has shown that during fast growth OpgH (which in our

experiments is nearly always mutated alongside *hfq*) binds FtsZ to postpone cell division. Thus, it

may be that negative fitness effects experienced by *hfq-rpoS* double mutants are the result of cell

division anomalies mitigated by mutations in *opgH*. It is noteworthy in this regard that whereas

cells in the Maharjan et al. experiments experienced a dilution rate of D=0.1 hr^{-1}, those in

evolutions performed by Adams, Helling and colleagues were doubling twice as fast (D=0.2 h^{-1}).

Thus, this discrepancy may be a manifestation of trade-offs between glucose uptake and cell

viability.

Finally, some mutations occur repeatedly and are likely beneficial, but their dynamics are

unpredictable (e.g. beneficial mutations in transcriptional terminator *rho* sweep when they co-

occur with beneficial mutations upstream of *mglB,* but otherwise remain at low frequency (**Table

S3, Fig. S1**). This dependence on genetic context, or "quasi-hitchhiking", of beneficial mutations

was previously observed by Lang et al. (2013) in evolving yeast populations and may be consistent

feature of microbial evolution experiments that becomes observable when populations are

sequenced at high depth of coverage and sufficient temporal resolution [14].

The evolution of population genetic complexity. Szostak, Hazen and others [88,89] argue that

a biological system's complexity should be evaluated in terms of its functional information

content. Although the total number of alleles in an evolving population at any given time-point is

information content, it is functional only in how it is integrated among the lineages co-existing at

that time-point. Our approach of integrating population sequencing with clone sequencing makes

it possible to estimate the pace and extent with which complexity, measured as lineage-specific

functional information [88,89], emerges in replicate evolving populations originating from a

common ancestor. Implicit in our perspective is the assumption that the sequence differences by

which lineages can be distinguished have physiological and fitness consequences. For each

population, we calculated at 50-generation intervals three measures of complexity: Shannon's

Entropy (H), equitability (H/H$_{max}$) and normalized population richness (Lineage counts) (**Fig. S5**).

All three measures increased during the course of evolution, but with a different tempo in each

population. Complexity increased in population 3 with no indication of reaching an asymptote by

generation 300 when the experiment was terminated. Populations 1 and 2 reached asymptotes after

~400 generations, following a steady increase in population 1, but a nearly-300 generation period

of stasis in Population 2. While longer experiments are clearly called for, our finding that longer-

term experiments appear to reach an asymptote in complexity is consistent with theoretical [90]

and empirical [91]observations that fitness plateaus when microbes are cultured by serial transfer

or in chemostats, even starting with mutator strains [92], and that complexity itself may plateau

when its evolution is simulated using RNA-like replicators [93]. It is intriguing to contemplate the

possibility that there may be a limit to the level of clonal interference that can be sustained in

asexual populations once all avenues for large fitness gains have been exhausted. Indeed, it was

recently shown using lineage tracking, that while fixation of an adaptive mutant causes a stochastic

crash in diversity, the generation of new adaptive mutants within such a fixing lineage is expected

to generate new diversity, such that a longer term steady state level of diversity will be achieved

[94].

Previous evolution experiments founded with the ancestor used here often resulted in stable

sub-populations supported by cross-feeding. In the present experiments neither the spectrum of

observed mutations nor the structure of clone phylogenies provides evidence for the evolution of

this type of trophic interaction. Instead, the possibility of a plateau in complexity, coupled with the

finding that every population has driven residual metabolites close to their detection limit, suggest

that these populations converge on an adaptive peak by diverse mechanisms but that clonal

interference keeps adaptive lineages off the summit, confined to exploring the many roads by

which the summit can be approached.

Materials and Methods

Strains, media and culture conditions. *Escherichia coli* JA122, population samples and clones were maintained as permanent frozen stocks and stored at -80°C in 20% glycerol. Davis minimal medium was used for all liquid cultures with 0.025% glucose added for batch cultures and 0.0125% for chemostats, as previously described [34]. Chemostat cultures were initiated using colonies picked from Tryptone Agar (TA) plate inoculated with JA122, and outgrown in Davis minimal batch medium overnight. Chemostats were maintained at 30°C with a dilution rate of ≈ 0.2 hr^{-1} for 300-500 generations. Every other day culture density was assessed by measuring absorbance spectrophotometrically at A_{550}. Every other day, population samples were archived at -80°C, and assayed for purity by plating serial dilutions on TA and examining Colony Forming Units (CFU) that arose following 24-hr incubation at 30°C. When necessary, chemostats were re-started from frozen stocks (chemostat 1: generation 217; chemostat 2: generation 410; chemostat 3 generation 251). At each sequencing time-point, 50 mL of culture was pelleted then stored at -80°C for DNA extraction. For clone sequencing, entire colonies were picked from TA plates inoculated from glycerol stocks, and re-archived in 96-well plate format.

Metabolite assays. 10 mL of sterile, cell-free chemostat filtrate was concentrated 20-fold by lyophilization (Labconco 4.5 Liter Freeze Dry System), then re-suspended in 0.5 mL sterile Millipore water. Residual glucose and residual acetate concentrations were determined on concentrated filtrate. Glucose was assayed enzymatically using the High Sensitivity Glucose Assay Kit (Sigma-Aldrich, Cat# MAK181), while acetate concentration was determined using the Acetate Colorimetric Assay Kit (Sigma-Aldrich, Cat# MAK086). Results presented in **Fig. S3** represent means ± SEM of duplicate assays.

Population sequencing. Bacterial DNA was prepared using the DNeasy Blood and Tissue Kit (Qiagen, cat. 69504) following the manufacturer's guidelines. For population sequencing, 5×10^{10}

cells, collected from every 50 generations in three chemostat vessels (up to 500 generations in

vessels 1 and 2, and up to 300 generations in vessel 3, 29 samples total) and frozen as pellets,

yielded 10-20μg of DNA. Following Proteinase K treatment, RNaseA treatment was used (20μL

10mg/mL RNAse A, 2 min at room temperature) to avoid degraded RNA from visually obscuring

size selection during library preparation. Samples were split into two columns to avoid

overloading. Bacterial DNA was sheared to a 150-200bp fragment size using a Covaris S2 series

sonicator (6min, Duty=5%, Intensity=3, Cycles/Burst=200), and was then ligated to barcoded

adapters as described [95], except that 200bp fragments were size selected after adapter ligation

(to maximize the fidelity of sequencing, by reading each fragment in both directions). Six barcoded

libraries were combined and sequenced on each lane of HiSeq 2000 Illumina Sequencer.

Variant calling from population sequencing with CLC Genomics Workbench 7.5 Illumina reads

were trimmed (removing adapters on both ends) and stringently mapped (Mismatch cost 2,

Insertion cost 3, Deletion cost 3, Length fraction 1.0, Similarity fraction 0.97) to the reference

sequence (WIS_MG1655_m56). Variants were called with the following parameters: minimum

frequency 1%, minimal coverage 100, minimum count 2, and base quality filtering (neighborhood

radius 5, minimum central quality 15 and minimum neighborhood quality 20). Sequencing data

uncovered low-level contamination of whole population samples with *Serratia liquifaciensis*. We

therefore first determined proportion of contaminating reads by mapping population sequencing

to *S. liquifaciensis* genome and then removed SNPs with frequency closely tracking percentage of

contamination (between 1 and 5%) that matched *S. liquifaciensis* sequence.

Selection of clones for sequencing. Allele frequencies for each chemostat were examined at each

time point, and the time-point at which there was the largest number of alleles present at 5% or

greater frequency was chosen for the isolation of clones for whole genome sequencing. The

rationale for this was that it would afford us the greatest opportunity to phase as many high

frequency alleles as possible.

Clonal DNA preparation. A colony was re-suspended in 300µL of sterile ddH$_2$0 with 17%

glycerol and stored in three aliquots at -80°C. 100 µL of glycerol stock were used for DNA

preparation. After removing glycerol (using MultiScreen High Volume Filter Plates with 0.45 µm

Durapore membrane, Millipore MVHVN4525), cell were resuspended in 500µL LB and grown

overnight at 30°C without shaking in deep well plates. Cells were collected again using filter plates

and subjected to DNeasy 96 Blood and Tissue Kit (Qiagen 69581) (yielding 4-15µg per strain).

Clonal libraries preparation and sequencing. Multiplexed sequencing libraries from clones were

prepared using the Nextera kit (Illumina catalog # FC-121-1031 and # FC-121-1012) as described

in [96], starting with 1-4ng of genomic DNA. Resulting libraries from each 96-well plate were

pooled at an equal volume. Resulting pooled libraries were analyzed on the Qubit and Bioanalyzer

platforms and sequenced on HiSeq 2000 (one lane per 96 clone pool). All raw sequencing data are

available from the SRA under BioProject ID PRJNA517527.

Variant calling from clonal sequencing with CLC Genomics Workbench 7.5 Short reads with

adapters removed were mapped to the reference with the same parameters as above, except the

length fraction was set to 0.5, and the similarity fraction to 0.8. Variants were called with a

minimum frequency 80%, minimum count 2 and the same base quality filtering as above.

Generation of phylogenies. For each chemostat, SNP and indel events for all 96 clones and the

ancestor JA122 were concatenated and re-coded as binary characters (i.e. presence/absence with

the ancestral state composed of all zeroes) and assembled into character matrices. PAUP ver.

4.0a149 was used to generate Camin-Sokal parsimony trees using the ancestor as the outgroup

under the assumption that reversions were extremely unlikely due to the extreme transversion bias

[97,98]. Tree files (.tre) were loaded into the Interactive Tree of Life (iTOL) web service for

character mapping and figure generation [99].

Determining genes with an excess of mutations. To identify genes with an excess of mutations,

we first determined the overall density of mutations as:

$\rho = M/L$, where M is the total number of mutations, and L is the length of the genome.

The probability of a given mutation landing in a segment of length l, is:

$\lambda = \rho \times l$

To calculate the p-value of n mutations landing in a segment of length l, we assume a Poisson

sampling process of a mutation landing in a given segment, and thus use:

599
$$p = \sum_{i=n}^{\infty} \frac{\lambda^i \times e^{-\lambda}}{i!}$$

though in practice, we capped i arbitrarily at 50, as continually summing at $i > 50$ does not

appreciably affect the calculated p-value. For a given segment, we calculated the number of

segments that would be expected to have p-value as good or better, as the number of tested

segments multiplied by the p-value. From this, we also determined a false positive rate.

Generation of Muller diagrams. Based on both the clonal sequencing we were able to determine

which mutations were in which lineages together, and from both the clonal and population

sequencing an approximate order of those mutations (though this was not exhaustive for all

mutations). Using these data, we developed a lineage file format that described which mutations

occurred in which lineages, and which lineages descended from one another, and used a custom

Perl script that combined this information with the allele frequencies over time from the population

sequencing to generate a graphical representation of the evolutionary dynamics, often referred to

as a Muller diagram.

Acknowledgements

The authors thank Matthew Herron and Eugene Kroll for their careful reading of the manuscript and their thoughtful suggestions for its improvement.

Author Contributions

Conceived and designed the experiments: GS MK FR. Performed the experiments: MK and KS. Analyzed the data: GS KS MK JB DY FR. Contributed reagents/materials/analysis tools: GS. Wrote the paper: MK GS FR

Main Figure legends

Fig. 1. Input of de novo mutations. The rate at which novel alleles appear, and the proportion of synonymous, missense and nonsense mutations, and non-coding mutations are consistent across all three replicate evolutions.

Fig. 2. Most *de novo* mutations only reach low allele frequencies, and experience pervasive clonal interference. (A) Histogram of maximum allele frequencies from three replicate evolutions, (B) Final versus maximum allele frequency for each *de novo* mutation, shows most mutations are at a lower frequency at the end of the experiment than they were at their maximum.

Fig. 3. Isolated clones are representative of the populations from which they are drawn. Mutation frequencies for population and clonal sequencing for mutations identified in both datasets at the same time-point shows similar frequencies.

Fig. 4. Clone phylogenies. Phylogenies depicting relationships among sequenced clones isolated from chemostats when allelic diversity attained its maximum. Distributions of different *malK*, *malT*, *fimH*, *hfq* and *opgH* alleles are indicated by colored bars. For each gene, all alleles observed in the dataset are numbered (see File S2 for details of which number corresponds to which allele for each gene). Underlined numbers denote alleles independently observed in more than one chemostat, while numbers marked with an asterisk appear to have arisen more than once within the same vessel. Grey shading delineates clades comprised of clones that have not acquired the standard mutations related to enhanced glucose uptake and instead carry variant *fimH* alleles that contribute to biofilm formation. Bracketed clones in chemostat 3 exhibited mutations expected to revert the ancestral nonsense mutations in the housekeeping gene encoding sigma factor RpoD.

Fig. 5. **Muller diagrams.** Evolutionary dynamics of adaptive lineages, deduced from combining

whole-population whole-genome sequence data and whole-genome sequence data of individual

clones isolated from each chemostat at the time-point where allelic diversity reached its maximum

value. Select genes are indicated in the plots – see Figure S1 and Supplementary Files S3-S5 for

additional details. Also note, most mutations that went extinct by the sampling timepoint are not

shown. See Figure S1 for their relative frequencies.

Fig. 6. **Overview of pathways relating some of the most frequently mutated genes to glucose**

transport and metabolism. Numbers in parentheses next to protein/gene names denote the

number of mutant alleles found in each chemostat population over the course of 300-500

generations (also see Table S3).

Supplementary Figure Legends

Fig. S1. Population-level dynamics of mutations in 10 frequently hit genes show consistent patterns. For each panel, elapsed generations are depicted on the x-axis and the height of each grey box represents a frequency of 100%. Cumulative frequencies for all alleles of a given gene present in the population at each time point were calculated and are represented as colored plots. (A) chemostat 1 (B) chemostat 2 (C) chemostat 3.

Fig. S2. MalK/MalT population dynamics. Mutant alleles of both LamB regulators, *malT* and *malK*, seldom co-occur in the same lineage, and when they do, those lineages fail to go to high frequency.

Fig. S3. **Cell density and residual metabolite concentrations**. Chemostat populations at steady state exhibit balanced growth, where population size remains constant and the limiting substrate, glucose is near its detection limit. As expected, populations initially produce the overflow metabolite acetate, as the founder carries a mutation that dysregulates acetyl CoA synthetase, the chief route by which *E. coli* assimilates low levels of this metabolite.

Fig. S4. Mutations in glucosyltransferase *opgH* occur repeatedly and go to high frequency.

Fig. S5. Patterns of change in population genetic complexity. Shannon's Entropy [H], Equitability [H/Hmax] and normalized population Richness [Lineage counts] were calculated at 50 generation intervals for each of three replicate evolution experiments. Shannon's entropy is an effective metric of population diversity as it accounts for both lineage richness (the number of

lineages observed) and the relative abundance of different lineages (evenness). Lineage richness

was normalized between zero and one by dividing the number of observed lineages, S, by the

maximum S observed over the course of each experiment.

Main Tables

Table 1. Characteristics of frequently mutated genes. Each asterisk indicates an allele that

arose more than once independently, either within or between vessels.

Supplementary Tables

Table S1. Key mutations that distinguish ancestral strain JA122 from K12 (MG1655)

Table S2. Beneficial alleles

Table S3. Population allele frequencies for frequently mutated genes

Table S4. Identical mutations arise within and among replicate evolution experiments.

Table S5. Fixed alleles among replicate populations ("fixed" defined as >98% at any time point

between generation 50 and 500).

Supplementary Data Files

File S1. Identity and frequencies of mutations detected via population sequencing.

File S2. Alleles mapped onto clone phylogenies represented in main Fig. 4.

File S3. Muller diagrams for novel alleles arising in chemostat 1, showing details for each lineage

File S4. Muller diagrams for novel alleles arising in chemostat 2, showing details for each lineage

File S5. Muller diagrams for novel alleles arising in chemostat 3, showing details for each lineage.

Table 1 Frequently mutated genes. Each asterisk indicates an allele that
arose more than once independently, either within or between vessels.

Rank	Gene	Observed Mutations	Expected Mutations	P-Value	FDR
Population sequencing					
1	galS***	38	0.78	6.55E-50	4.42E-45
2	hfq********	24	0.23	6.91E-40	2.33E-35
3	pgi********	35	1.23	4.54E-38	1.02E-33
4	opgH**	31	1.90	8.74E-27	1.48E-22
5	malT*********	30	2.02	8.10E-25	1.09E-20
6	malK*******	22	0.83	7.47E-24	8.40E-20
7	upstream mglB**	7	0.21	2.91E-09	2.81E-05
8	rho**	11	0.94	5.49E-09	4.64E-05
9	upstream dnaG	5	0.08	3.06E-08	2.30E-04
10	fimH***	9	0.68	4.38E-08	2.96E-04
Clonal sequencing					
1	hfq*******	26	0.1020	3.79E-53	3.68E-48
2	pgi****	17	0.5448	5.52E-20	2.68E-15
3	opgH***	17	0.8400	6.57E-17	2.13E-12
4	upstream mglB**	8	0.0925	1.22E-13	2.96E-09
5	fimH****	10	0.2982	1.17E-12	2.26E-08
6	ompR***	8	0.2377	2.05E-10	3.31E-06
7	upstream adhE*	6	0.1575	1.85E-08	0.000257
8	malT***	10	0.8935	3.98E-08	0.000482
9	proQ*	6	0.2308	1.72E-07	0.001858
10	pfkA**	6	0.3180	1.09E-06	0.010612

References

1. Lang GI, Desai MM (2014) The spectrum of adaptive mutations in experimental evolution. Genomics 104: 412-416.
2. Cvijovic I, Nguyen Ba AN, Desai MM (2018) Experimental studies of evolutionary dynamics in Microbes. Trends Genet 34: 693-703.
3. Cooper VS (2018) Experimental Evolution as a high-throughput screen for genetic adaptations. mSphere 3.
4. Garay E, Campos SE, Gonzalez de la Cruz J, Gaspar AP, Jinich A, et al. (2014) High-resolution profiling of stationary-phase survival reveals yeast longevity factors and their genetic interactions. PLoS Genet 10: e1004168.
5. Paradis-Bleau C, Kritikos G, Orlova K, Typas A, Bernhardt TG (2014) A genome-wide screen for bacterial envelope biogenesis mutants identifies a novel factor involved in cell wall precursor metabolism. PLoS Genet 10: e1004056.
6. Novick A, Szilard L (1950) Experiments with the chemostat on spontaneous mutations of bacteria. Proc Natl Acad Sci U S A 36: 708-719.
7. Novick A, Szilard L (1951) Experiments on spontaneous and chemically induced mutations of bacteria growing in the Chemostat. Cold Spring Harb Symp Quant Biol 16: 337-343.
8. Atwood KC, Schneider LK, Ryan FJ (1951) Periodic selection in Escherichia coli. Proc Natl Acad Sci U S A 37: 146-155.
9. Atwood KC, Schneider LK, Ryan FJ (1951) Selective mechanisms in bacteria. Cold Spring Harb Symp Quant Biol 16: 345-355.
10. Haldane JBS (1927) A Mathematical Theory of Natural and Artificial Selection, Part V: Selection and Mutation. Mathematical Proceedings of the Cambridge Philosophical Society 23: 838-844.
11. Fisher RAS (1930) The genetical theory of natural selection. Oxford: Clarendon Press.
12. Muller HJ (1932) Some genetic aspects of sex. The American Naturalist 66: 118-138.
13. Gause GF (1934) Experimental analysis of Vito Volterra's mathematical theory of the struggle for existence. Science 79: 16-17.
14. Lang GI, Rice DP, Hickman MJ, Sodergren E, Weinstock GM, et al. (2013) Pervasive genetic hitchhiking and clonal interference in forty evolving yeast populations. Nature 500: 571-574.
15. Marad DA, Buskirk SW, Lang GI (2018) Altered access to beneficial mutations slows adaptation and biases fixed mutations in diploids. Nat Ecol Evol 2: 882-889.
16. Levy SF, Blundell JR, Venkataram S, Petrov DA, Fisher DS, et al. (2015) Quantitative evolutionary dynamics using high-resolution lineage tracking. Nature 519: 181-186.
17. Good BH, McDonald MJ, Barrick JE, Lenski RE, Desai MM (2017) The dynamics of molecular evolution over 60,000 generations. Nature 551: 45-50.
18. Lauer S, Avecilla G, Spealman P, Sethia G, Brandt N, et al. (2018) Single-cell copy number variant detection reveals the dynamics and diversity of adaptation. PLoS Biol 16: e3000069.
19. Kao KC, Sherlock G (2008) Molecular characterization of clonal interference during adaptive evolution in asexual populations of Saccharomyces cerevisiae. Nat Genet 40: 1499-1504.
20. Kvitek DJ, Sherlock G (2013) Whole genome, whole population sequencing reveals that loss of signaling networks is the major adaptive strategy in a constant environment. PLoS Genet 9: e1003972.

21. Behringer MG, Choi BI, Miller SF, Doak TG, Karty JA, et al. (2018) Escherichia coli
cultures maintain stable subpopulation structure during long-term evolution. Proc Natl
Acad Sci U S A 115: E4642-E4650.
22. Rozen DE, Lenski RE (2000) Long-term experimental evolution in Escherichia coli. VIII.
Dynamics of a balanced polymorphism. Am Nat 155: 24-35.
23. Herron MD, Doebeli M (2013) Parallel evolutionary dynamics of adaptive diversification in
Escherichia coli. PLoS Biol 11: e1001490.
24. Maddamsetti R, Lenski RE, Barrick JE (2015) Adaptation, clonal interference, and
frequency-dependent jnteractions in a long-term evolution experiment with Escherichia
coli. Genetics 200: 619-631.
25. Charlesworth D (2006) Balancing selection and its effects on sequences in nearby genome
regions. PLoS Genet 2: e64.
26. Rainey PB, Travisano M (1998) Adaptive radiation in a heterogeneous environment. Nature
394: 69-72.
27. Rozen DE, Schneider D, Lenski RE (2005) Long-term experimental evolution in Escherichia
coli. XIII. Phylogenetic history of a balanced polymorphism. J Mol Evol 61: 171-180.
28. Kinnersley M, Wenger J, Kroll E, Adams J, Sherlock G, et al. (2014) Ex uno plures: clonal
reinforcement drives evolution of a simple microbial community. PLoS Genet 10:
e1004430.
29. Rosenzweig RF, Sharp RR, Treves DS, Adams J (1994) Microbial evolution in a simple
unstructured environment: genetic differentiation in Escherichia coli. Genetics 137: 903-
917.
30. Helling RB, Vargas CN, Adams J (1987) Evolution of Escherichia coli during growth in a
constant environment. Genetics 116: 349-358.
31. Gudelj I, Kinnersley M, Rashkov P, Schmidt K, Rosenzweig F (2016) Stability of cross-
feeding polymorphisms in microbial communities. PLoS Comput Biol 12: e1005269.
32. Maharjan RP, Liu B, Feng L, Ferenci T, Wang L (2015) Simple phenotypic sweeps hide
complex genetic changes in populations. Genome Biol Evol 7: 531-544.
33. Treves DS, Manning S, Adams J (1998) Repeated evolution of an acetate-crossfeeding
polymorphism in long-term populations of Escherichia coli. Mol Biol Evol 15: 789-797.
34. Kinnersley MA, Holben WE, Rosenzweig F (2009) E Unibus Plurum: genomic analysis of
an experimentally evolved polymorphism in Escherichia coli. PLoS Genet 5: e1000713.
35. Atlung T, Nielsen HV, Hansen FG (2002) Characterisation of the allelic variation in the rpoS
gene in thirteen K12 and six other non-pathogenic Escherichia coli strains. Mol Genet
Genomics 266: 873-881.
36. Singaravelan B, Roshini BR, Munavar MH (2010) Evidence that the supE44 mutation of
Escherichia coli is an amber suppressor allele of glnX and that it also suppresses ochre
and opal nonsense mutations. J Bacteriol 192: 6039-6044.
37. Au KG, Clark S, Miller JH, Modrich P (1989) Escherichia coli mutY gene encodes an
adenine glycosylase active on G-A mispairs. Proc Natl Acad Sci U S A 86: 8877-8881.
38. Blattner FR, Plunkett G, 3rd, Bloch CA, Perna NT, Burland V, et al. (1997) The complete
genome sequence of Escherichia coli K-12. Science 277: 1453-1462.
39. Lee H, Popodi E, Tang H, Foster PL (2012) Rate and molecular spectrum of spontaneous
mutations in the bacterium Escherichia coli as determined by whole-genome sequencing.
Proc Natl Acad Sci U S A 109: E2774-2783.

40. de Visser JA, Rozen DE (2006) Clonal interference and the periodic selection of new
beneficial mutations in Escherichia coli. Genetics 172: 2093-2100.
41. Hermisson J, Pennings PS (2005) Soft sweeps: molecular population genetics of adaptation
from standing genetic variation. Genetics 169: 2335-2352.
42. Pennings PS, Hermisson J (2006) Soft sweeps II--molecular population genetics of
adaptation from recurrent mutation or migration. Mol Biol Evol 23: 1076-1084.
43. Pennings PS, Hermisson J (2006) Soft sweeps III: the signature of positive selection from
recurrent mutation. PLoS Genet 2: e186.
44. Desai MM, Walczak AM, Fisher DS (2013) Genetic diversity and the structure of
genealogies in rapidly adapting populations. Genetics 193: 565-585.
45. Notley-McRobb L, Ferenci T (1999) The generation of multiple co-existing mal-regulatory
mutations through polygenic evolution in glucose-limited populations of Escherichia coli.
Environ Microbiol 1: 45-52.
46. Hermisson J, Pennings PS (2017) Soft sweeps and beyond: understanding the patterns and
probabilities of selection footprints under rapid adaptation. Methods in Ecology and
Evolution 8: 700-716.
47. Jensen JD (2014) On the unfounded enthusiasm for soft selective sweeps. Nat Commun 5:
5281.
48. Fogle CA, Nagle JL, Desai MM (2008) Clonal interference, multiple mutations and
adaptation in large asexual populations. Genetics 180: 2163-2173.
49. Buskirk SW, Peace RE, Lang GI (2017) Hitchhiking and epistasis give rise to cohort
dynamics in adapting populations. Proc Natl Acad Sci U S A 10.1073/pnas.1702314114.
50. Diaz Caballero J, Clark ST, Coburn B, Zhang Y, Wang PW, et al. (2015) Selective Sweeps
and Parallel Pathoadaptation Drive Pseudomonas aeruginosa Evolution in the Cystic
Fibrosis Lung. MBio 6: e00981-00915.
51. Ferenci T (2001) Hungry bacteria--definition and properties of a nutritional state. Environ
Microbiol 3: 605-611.
52. Maharjan RP, Ferenci T (2013) Epistatic interactions determine the mutational pathways and
coexistence of lineages in clonal Escherichia coli populations. Evolution 67: 2762-2768.
53. Manch K, Notley-McRobb L, Ferenci T (1999) Mutational adaptation of Escherichia coli to
glucose limitation involves distinct evolutionary pathways in aerobic and oxygen-limited
environments. Genetics 153: 5-12.
54. Notley-McRobb L, Ferenci T (1999) Adaptive mgl-regulatory mutations and genetic
diversity evolving in glucose-limited Escherichia coli populations. Environ Microbiol 1:
33-43.
55. Notley-McRobb L, Ferenci T (2000) Experimental analysis of molecular events during
mutational periodic selections in bacterial evolution. Genetics 156: 1493-1501.
56. Notley-McRobb L, Seeto S, Ferenci T (2003) The influence of cellular physiology on the
initiation of mutational pathways in Escherichia coli populations. Proc Biol Sci 270: 843-
848.
57. Ferenci T (1996) Adaptation to life at micromolar nutrient levels: the regulation of
Escherichia coli glucose transport by endoinduction and cAMP. FEMS Microbiol Rev
18: 301-317.
58. Death A, Ferenci T (1993) The importance of the binding-protein-dependent Mgl system to
the transport of glucose in Escherichia coli growing on low sugar concentrations. Res
Microbiol 144: 529-537.

845 59. Death A, Notley L, Ferenci T (1993) Derepression of LamB protein facilitates outer
846   membrane permeation of carbohydrates into Escherichia coli under conditions of nutrient
847   stress. J Bacteriol 175: 1475-1483.
848 60. Tweeddale H, Notley-McRobb L, Ferenci T (1999) Assessing the effect of reactive oxygen
849   species on Escherichia coli using a metabolome approach. Redox Rep 4: 237-241.
850 61. Geanacopoulos M, Adhya S (1997) Functional characterization of roles of GalR and GalS as
851   regulators of the gal regulon. J Bacteriol 179: 228-234.
852 62. Maharjan R, McKenzie C, Yeung A, Ferenci T (2013) The basis of antagonistic pleiotropy in
853   hfq mutations that have opposite effects on fitness at slow and fast growth rates. Heredity
854   (Edinb) 110: 10-18.
855 63. Maharjan R, Zhou Z, Ren Y, Li Y, Gaffe J, et al. (2010) Genomic identification of a novel
856   mutation in hfq that provides multiple benefits in evolving glucose-limited populations of
857   Escherichia coli. J Bacteriol 192: 4517-4521.
858 64. Richet E, Joly N, Danot O (2005) Two domains of MalT, the activator of the Escherichia coli
859   maltose regulon, bear determinants essential for anti-activation by MalK. J Mol Biol 347:
860   1-10.
861 65. Dardonville B, Raibaud O (1990) Characterization of malT mutants that constitutively
862   activate the maltose regulon of Escherichia coli. J Bacteriol 172: 1846-1852.
863 66. Bohm A, Diez J, Diederichs K, Welte W, Boos W (2002) Structural model of MalK, the
864   ABC subunit of the maltose transporter of Escherichia coli: implications for mal gene
865   regulation, inducer exclusion, and subunit assembly. J Biol Chem 277: 3708-3717.
866 67. Moller P, Overloper A, Forstner KU, Wen TN, Sharma CM, et al. (2014) Profound impact of
867   Hfq on nutrient acquisition, metabolism and motility in the plant pathogen
868   Agrobacterium tumefaciens. PLoS One 9: e110427.
869 68. Maharjan RP, Ferenci T, Reeves PR, Li Y, Liu B, et al. (2012) The multiplicity of divergence
870   mechanisms in a single evolving population. Genome Biol 13: R41.
871 69. Cardinale CJ, Washburn RS, Tadigotla VR, Brown LM, Gottesman ME, et al. (2008)
872   Termination factor Rho and its cofactors NusA and NusG silence foreign DNA in E. coli.
873   Science 320: 935-938.
874 70. Banerjee S, Chalissery J, Bandey I, Sen R (2006) Rho-dependent transcription termination:
875   more questions than answers. J Microbiol 44: 11-22.
876 71. Peters JM, Mooney RA, Kuan PF, Rowland JL, Keles S, et al. (2009) Rho directs widespread
877   termination of intragenic and stable RNA transcription. Proc Natl Acad Sci U S A 106:
878   15406-15411.
879 72. Ciampi MS (2006) Rho-dependent terminators and transcription termination. Microbiology
880   152: 2515-2528.
881 73. Colonna B, Hofnung M (1981) rho Mutations restore lamB expression in E. coli K12 strains
882   with an inactive malB region. Mol Gen Genet 184: 479-483.
883 74. Nakashima N, Ohno S, Yoshikawa K, Shimizu H, Tamura T (2014) A vector library for
884   silencing central carbon metabolism genes with antisense RNAs in Escherichia coli. Appl
885   Environ Microbiol 80: 564-573.
886 75. Hill NS, Buske PJ, Shi Y, Levin PA (2013) A moonlighting enzyme links Escherichia coli
887   cell size with central metabolism. PLoS Genet 9: e1003663.
888 76. Lange R, Hengge-Aronis R (1994) The cellular concentration of the sigma S subunit of RNA
889   polymerase in Escherichia coli is controlled at the levels of transcription, translation, and
890   protein stability. Genes Dev 8: 1600-1612.

77. King T, Ishihama A, Kori A, Ferenci T (2004) A regulatory trade-off as a source of strain
variation in the species Escherichia coli. J Bacteriol 186: 5614-5620.
78. Notley-McRobb L, King T, Ferenci T (2002) rpoS mutations and loss of general stress
resistance in Escherichia coli populations as a consequence of conflict between
competing stress responses. J Bacteriol 184: 806-811.
79. Versalovic J, Koeuth T, Britton R, Geszvain K, Lupski JR (1993) Conservation and
evolution of the rpsU-dnaG-rpoD macromolecular synthesis operon in bacteria. Mol
Microbiol 8: 343-355.
80. Hutchins PR, Miller SR (2017) Genomics of variation in nitrogen fixation activity in a
population of the thermophilic cyanobacterium Mastigocladus laminosus. ISME J 11: 78-
86.
81. Hottes AK, Freddolino PL, Khare A, Donnell ZN, Liu JC, et al. (2013) Bacterial adaptation
through loss of function. PLoS Genet 9: e1003617.
82. Venkataram S, Dunn B, Li Y, Agarwala A, Chang J, et al. (2016) Development of a
Comprehensive Genotype-to-Fitness Map of Adaptation-Driving Mutations in Yeast.
Cell 166: 1585-1596 e1522.
83. Phan K, Ferenci T (2013) A design-constraint trade-off underpins the diversity in
ecologically important traits in species Escherichia coli. ISME J 7: 2034-2043.
84. Yao R, Hirose Y, Sarkar D, Nakahigashi K, Ye Q, et al. (2011) Catabolic regulation analysis
of Escherichia coli and its crp, mlc, mgsA, pgi and ptsG mutants. Microb Cell Fact 10:
67.
85. Vecerek B, Rajkowitsch L, Sonnleitner E, Schroeder R, Blasi U (2008) The C-terminal
domain of Escherichia coli Hfq is required for regulation. Nucleic Acids Res 36: 133-
143.
86. Takada A, Wachi M, Nagai K (1999) Negative regulatory role of the Escherichia coli hfq
gene in cell division. Biochem Biophys Res Commun 266: 579-583.
87. Zambrano N, Guichard PP, Bi Y, Cayrol B, Marco S, et al. (2009) Involvement of HFq
protein in the post-transcriptional regulation of E. coli bacterial cytoskeleton and cell
division proteins. Cell Cycle 8: 2470-2472.
88. Hazen RM, Griffin PL, Carothers JM, Szostak JW (2007) Functional information and the
emergence of biocomplexity. Proc Natl Acad Sci U S A 104 Suppl 1: 8574-8581.
89. Szostak JW (2003) Functional information: Molecular messages. Nature 423: 689.
90. Gordo I, Campos PR (2013) Evolution of clonal populations approaching a fitness peak. Biol
Lett 9: 20120239.
91. de Visser JA, Lenski RE (2002) Long-term experimental evolution in Escherichia coli. XI.
Rejection of non-transitive interactions as cause of declining rate of adaptation. BMC
Evol Biol 2: 19.
92. Maharjan RP, Liu B, Li Y, Reeves PR, Wang L, et al. (2013) Mutation accumulation and
fitness in mutator subpopulations of Escherichia coli. Biol Lett 9: 20120961.
93. Takeuchi N, Hogeweg P (2008) Evolution of complexity in RNA-like replicator systems.
Biol Direct 3: 11.
94. Blundell JR, Schwartz K, Francois D, Fisher DS, Sherlock G, et al. (2018) The dynamics of
adaptive genetic diversity during the early stages of clonal evolution. Nat Ecol Evol
10.1038/s41559-018-0758-1.

95. Schwartz K, Wenger JW, Dunn B, Sherlock G (2012) APJ1 and GRE3 homologs work in
concert to allow growth in xylose in a natural Saccharomyces sensu stricto hybrid yeast.
Genetics 191: 621-632.
96. Kryazhimskiy S, Rice DP, Jerison ER, Desai MM (2014) Microbial evolution. Global
epistasis makes adaptation predictable despite sequence-level stochasticity. Science 344:
1519-1522.
97. Blount ZD, Barrick JE, Davidson CJ, Lenski RE (2012) Genomic analysis of a key
innovation in an experimental Escherichia coli population. Nature 489: 513-518.
98. Swofford DL (2002) PAUP*. Phylogenetic Analysis Using Parsimony (*and Other
Methods). 4 ed. Sunderland, Massachusetts: Sinauer Associates.
99. Letunic I, Bork P (2007) Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree
display and annotation. Bioinformatics 23: 127-128.
947

Fig. 1 Input of de novo mutations

Fig. 2 Most de novo mutations only reach low allele frequencies

Fig. 3 Isolated clones are representative of the populations from

Fig. 4 Clone phylogenies

Fig. 5 Muller diagrams

Fig. 6 Pathways containing frequently mutated genes