1   **Title:** Fingerprinting Adiposity and Metabolic Function in the Brains of Overweight and Obese
2   Humans
3

4   **Authors:** Michael C. Farruggia[a,b], Maria J. Van Kooten[b,c], Mary V. Burke[a,b], Dustin
5   Scheinost[d,e,f], R. Todd Constable[a,d,g], Dana M. Small[a,b,h,i]
6
Affiliations:
[a]Interdepartmental Neuroscience Program, Yale University, 333 Cedar Street, New Haven, CT,
U.S.
[b]Department of Psychiatry, Yale University School of Medicine, 300 George Street, New Haven,
CT 06511, USA.
[c]University of Groningen, Faculty of Medical Sciences, Antonius Deusinglaan 1, 9713 AV
Groningen, The Netherlands.
[d]Department Radiology and Biomedical Imaging, Yale University School of Medicine, New
Haven, CT, United States.
[e]Department Statistics and Data Science, Yale University, New Haven, CT, United States.
[f]Child Study Center, Yale School of Medicine, New Haven, CT, United States.
[g]Department of Neurosurgery, Yale University School of Medicine, New Haven, CT, United
States.
[h]Department of Psychology, Yale University, New Haven, CT, United States.
[i]fMEG Center, University of Tubingen, Tubingen, Germany.
22
23
24
Correspondence should be addressed to:
*Dana M Small, PhD
dana.small@yale.edu
(D.M.S.)
*Michael C Farruggia, MS
michael.farruggia@yale.edu
(M.C.F.)
1 Church Street, Suite 6a
New Haven, CT 06510
Phone: (954) 325-2660

One Sentence Summary: We find that brain connectivity in humans can best predict waist
circumference over other adiposity measures, and that this predictive network is largely distinct
from one that can predict fasting insulin, a marker of metabolic dysfunction.
38

Abstract: Evidence points to the brain as central in the pathophysiology of obesity. However,
none of the methods used to diagnose and quantify obesity such as body mass index (BMI), body
weight, percent body fat, waist circumference, and waist-hip ratio are brain measures. Here we
used Connectome-based Predictive Modeling (CPM), a data-driven approach, to determine
whether brain CPMs could be isolated for these measures of adiposity. CPM exploits individual
variation in whole-brain functional connectivity to predict a behavior or trait that varies across
individuals in a leave-one-out cross-validated framework. We used CPM to predict BMI, weight,
percent body fat, waist circumference, and waist/hip ratio in 67 overweight and obese individuals
(ages 18-45) using individual blood oxygen level dependent (BOLD) response to milkshake. A
CPM could only be identified for waist circumference, consisting of high-degree nodes in
prefrontal, parietal, limbic, cerebellar, subcortical, and temporal networks (r =0.354, p<0.01,
FDR corrected). In an exploratory analysis, we also found that CPM could predict fasting blood
insulin (r = 0.271, p = 0.047), a measure of insulin resistance. This CPM consisted of high-
degree nodes in cerebellar, temporal, and limbic networks and was largely separable from the
waist circumference CPM. Notably, CPM could not isolate a fingerprint for BMI, which is the
most frequently used measure to assess brain correlates of obesity. These findings suggest that
brain function is more tightly coupled to waist circumference than BMI and that adiposity and
glucose tolerance have distinct associations with brain function.

Introduction

Obesity is a neurobehavioral disorder resulting from a biologically vulnerable brain that is highly sensitive to its environment (*1*). Evidence for this derives from genome-wide association studies (GWAS), demonstrating that genetic variants associated with obesity tend to affect brain function. For example, high-risk variants in the FTO (fat mass and obesity associated) gene predispose individuals to weighing on average 3kg more and increasing their obesity risk 1.67x over those with low-risk variants (*2*, *3*). The FTO gene encodes the alpha-ketoglutarate-dependent dioxygenase enzyme, a protein that is highly expressed in the human brain, particularly in the hypothalamus and cerebellum as well as on dopamine receptors where it influences reward learning (*4 - 6*). Hypotheses of FTO's action in the human brain suggest that it may act to increase consumption of food and alter sensing of dietary macronutrient composition (*7*). Since the discovery of FTO, additional GWAS have found another 97 BMI-associated SNPs which encode for proteins that are significantly enriched in the human brain (*8*).

While the vast majority of evidence continues to point to the brain as integral in the pathophysiology of obesity, none of the adiposity measures we have to quantify it or characterize its extent are brain measures. Body mass index, or BMI, is the anthropometric measurement commonly used to classify obesity, with individuals having a BMI ≥ 30 characterized as obese (*9*). However, BMI as a measure of adiposity is often criticized for being an inaccurate reflection of fat mass, in particular in Asians and in athletes (*10*, *11*). Body fat percentage alone has also been criticized as an inaccurate predictor of negative health outcomes, as regional distribution of fat in the body influences risk for negative health consequences. For example, women with predominantly upper body adipose tissue are more likely to have diabetic glucose tolerance results and higher fasting plasma triglycerides than those with predominantly lower body adipose tissue (*12*). More recent studies have found that distribution of adipose tissue, particularly in the thighs, is inversely correlated with metabolic syndrome in obesity (*13*).

Ultimately, if obesity is a neurobehavioral disorder resulting from a vulnerable brain interacting with an obesogenic environment, the most informative adiposity measures should be predicted by brain function to some extent. Further, if an adiposity measure can be reliably predicted using features of brain function in a cross-validated framework, it follows that these features may represent a generalizable biomarker for adiposity risk. Thus, not only is it important that we are able to determine which adiposity measure is best predicted by whole-brain functional connectivity, but it also imperative to determine which anatomical regions (i.e. nodes) and which connections among them (i.e. edges) are influencing our predictive power.

With this in mind, we set out to determine which commonly-used adiposity measures, including BMI, body fat %, waist size, body weight, and waist/hip ratio, could be predicted by whole-brain function connectivity. To accomplish this, we use connectome-based predictive modeling (CPM), a data-driven approach that builds predictive models of brain-behavior relationships using connectivity data in a cross-validated framework (*14*, *15*). More specifically, CPM exploits individual variation in whole-brain functional connectivity to predict a behavior or trait that varies across individuals in a leave-one-out procedure. CPM has been used to identify brain "fingerprints" (*14*) as well as to predict personality traits (*16*), creative ability (*17*), and sustained attention (*18*). Here we used this technique to ask whether a predictive CPM could be identified for our different adiposity measures. As an exploratory aim, we also tested whether a CPM could be found for measures of glucose tolerance.

Results
CPM predicts individual differences in waist circumference
Of all measures, including BMI, body fat %, waist size, body weight, and waist/hip ratio, we
found that CPM significantly predicts waist circumference (combined network, r = 0.354, p =
0.005) after applying Benjamini-Hochberg FDR correction for multiple tests (Fig. 1). Overall,
the combined model accounts for approximately 12.5% of the variance in the waist
circumference of a novel subject.
Waist circumference CPM contains DMN, motor, visual, cerebellar, and subcortical
networks
The combined model consists of edges whose strengths are positively (positive network) or
negatively (negative network) correlated with our variable of interest, waist circumference.
These edges can be visualized as connections among nodes assigned to one of ten bilateral brain
regions (Fig. 2A), or as the density of edges among ten canonical networks normalized by
network size (Fig. 2B). Brain regions include Prefrontal, Motor, Insula, Parietal, Temporal,
Occipital, Limbic, Cerebellum, Subcortical, and Brainstem. We operationally define our high
degree nodes as the top N nodes that occur before multiple degree repetitions. High degree nodes
(degree ≥ 9), or nodes with the highest number of connections for the waist circumference CPM
were found in prefrontal, parietal, limbic, cerebellar, subcortical, and temporal regions (Fig. 2A,
see also 5A). AAL labels for these regions include left inferior frontal gyrus, left postcentral
gyrus, left amygdala, left culmen of cerebellum, right putamen and right fusiform gyrus,
respectively (Table 1).
Canonical networks emphasize different functional properties of the human brain and consist of
medial frontal (MF), frontoparietal (FP), default mode (DM), motor (MO), visual 1 (V1), visual
2 (V2), visual association (VA), cingulo-opercular (CO), subcortical (SC), and cerebellum (CE)
networks (*19*, *20*). The highest proportion of edges, that is, the edges used most in prediction for
the positive network exist between DM-V2, MO-CE, V2-VA, and CO-SC. In the negative
network, these are between DM-SC, MO-V2, MO-V1, V1-SC, VA-VA, and CE-CE (Fig. 2B).
Exploratory analysis: CPM predicts individual differences in fasting insulin
Due to the large scope of the dataset, we were able to conduct a post-hoc exploratory analysis of
metabolic markers to examine convergence or divergence of predictive networks as they relate to
the waist circumference CPM. For this analysis, we analyzed additional bloods data until we had
a complete metabolic panel from all 59 of 67 subjects. Fasting glucose, fasting insulin, and
HOMA-IR were input into CPM as our measures of interest. CPM was able to predict fasting
insulin (p < 0.05, uncorrected; Fig. 3A) but not fasting glucose or HOMA-IR, accounting for
approximately 7% of the variance in a novel individual's fasting insulin levels. Positive and
negative predictive networks for fasting insulin are notably sparser than those for waist
circumference (Fig. 3B). High degree nodes (degree ≥ 6) for these networks were found in
cerebellar (left crus and right lobule VIIB), temporal (left superior temporal pole and right
fusiform), and limbic (right posterior cingulate gyrus) regions (Fig. 3B, *see also 5B, Table 2*). In
the positive network, a high density of edges exists between MF-CE, DM-V2, MO-CE, V2-CE,
and V2-SC (Fig. 3C). In the negative network, these exist between MF-MF, FP-VA, FP-V1,
DM-CE, DM-SC, V1-SC, V1-V2, VA-CO, CO-CE, and SC-SC (Fig. 3C).
Waist circumference and fasting insulin CPMs are separable
To examine convergence and divergence of predictive networks, we subtracted the number of
negative edges from the number of positive edges between each canonical network and plotted
the resultant degree of polarity (Fig. 4A-B). For example, networks can have mostly positive or

negative edges connecting them (i.e. polarity), or an even distribution of positive and negative
edges (i.e. apolarity). Indeed, this technique demonstrates the degree to which positive or
negative edges are used in prediction and can help ascertain common or uncommon
neurobiological correlates between fasting insulin and waist circumference. The predictive
network for waist size shows a large degree of polarity (Fig. 4A), while the relative sparsity of
the predictive network for insulin is visible and there is notable apolarity particularly among
visual networks (Fig. 4B). Predictive networks for insulin and waist size are on average
dissimilar. Of the 55 network pairs, 26 network pairs (47%) exhibited similar degrees of polarity
between the waist circumference and insulin predictive networks. Most notably, patterns of
connectivity within cerebellar and subcortical networks showed the most similar patterns of
polarity between waist circumference and fasting insulin predictive networks (Fig. 4). In contrast
their polarity differs most in medial-frontal, frontoparietal, default mode, motor, visual, and
cingulo-opercular networks. To aid in comparison, we plotted the spatial extent of the top 10%
highest-degree predictive nodes for each network (Fig. 5A-B, Tables 1 & 2). Only nine of the top
10% of nodes (27 total) in each CPM are shared among each other, those being centered in left
posterior cingulate gyrus (node 227), right fusiform (59), right and left superior temporal pole
(92 and 188), left cerebellum (255), left and right lobule III of cerebellum (251 and 103), left
crus II of cerebellum (242), and midbrain (132). The 18 nodes (66%) that differ are depicted in
Fig. 6A-B.
The role of subject motion
We used frame-to-frame displacement (FTF) as a gross measure of head motion (*21*). FTF was
calculated for each run in each subject and averaged across runs. FTF significantly correlated
with waist size ($R^2 = 0.199$, $p < 0.001$), BMI ($R^2 = 0.115$, $p < 0.01$), body weight ($R^2 = 0.132$, p
< 0.01), waist/hip ratio ($R^2 = 0.103$, $p < 0.01$), and BF% ($R^2 = 0.107$, $p < 0.01$). FTF was notably
not correlated with fasting glucose ($r = 0.05$, $p = 0.68$), fasting insulin ($r = 0.15$, $p = 0.25$), or
HOMA-IR ($r = 0.15$, $p = 0.27$). While a correlation between any of our adiposity measures and
FTF suggests that motion should be accounted for using partial Pearson correlation, recent
evidence suggests that in-scanner head motion is heritable and phenotypically correlated with
BMI (*22*). Indeed, if head motion and adiposity contain shared genetic influences, then
controlling for motion in CPM would unnecessarily reduce its predictive power in this instance.
Instead we corrected for motion during preprocessing (see Methods) and report results following
partial Pearson correlation for mean FTF subject (fig. S2). Of note, CPMs for both waist
circumference and insulin were both significant, but did not survive FDR correction when
motion is included as a covariate.
183
Discussion
Our experiment produced several novel findings. First, we determined, using a data-driven
approach, that waist circumference could be predicted by whole-brain functional connectivity. In
particular, with CPM we can account for approximately 12.5% of the variance in this measure in
a novel individual. Notably, we could not predict BMI, the most commonly used measure to
define obesity, or other adiposity measures, such as body fat %, body weight, and waist/hip ratio.
Second, we found that a metabolic marker, fasting insulin, could be predicted from whole-brain
functional connectivity and that this predictive network was distinct from that of waist
circumference.
There is long-standing controversy regarding the optimal method to quantify adiposity. We
assert that, not only is it important to use a measure that captures an individual's risk for

morbidity and mortality, but it is equally important to capture obesity's underpinnings as a
neurobehavioral disorder. Waist circumference does both.
This proposal is consistent with a number of recent findings. Waist circumference has better
discriminatory capability for type 2 diabetes than either BMI or body adiposity index [BAI,
calculated as (hip circumference)/((height)$^{1.5}$)–18] in a northern Iranian population (*23*, *24*). A
systematic review found that waist circumference performed 3% better than BMI in
discriminating adults with hypertension, T2D, metabolic syndrome, dyslipidaemia, and general
cardiovascular outcomes (*25*). A meta-analysis of some 259,200 individuals found that high
waist circumference is a better predictor than high BMI for the development of diabetes across
all ethnicities tested and for the development of hypertension among Hispanics/Latinos in
particular (*26*). Additional evidence suggests that individuals who are classified as obese by
waist circumference have a greater risk for all-cause and cardiovascular disease-related death
than individuals who are classified as obese by BMI (*27*). In a separate cohort of 2266
hypertensive patients, waist circumference was able to predict the development of coronary
artery disease, while other measures of adiposity such as BMI and waist/hip ratio had no
independent prognostic value (*28*).
Although most neuroimaging studies use BMI to assess neural correlates of obesity, when waist
circumference is pitted against BMI, stronger and more extensive correlations are often found.
For example, waist circumference, is a better predictor of local gray matter volume
(*29*, *30*) and brain response to food cues (*31* – *33*). The current study confirms and extends these
findings by using functional connectivity and a whole-brain data-driven approach to show that
brain measures not only correlate most strongly with waist circumference, but also are able to
predict waist circumference. This indicates that this neural measure may act as a biomarker for
the measure of adiposity that best predicts obesity-related health outcomes (*34*).
More specifically, high-degree nodes for the waist circumference CPM were located in left
inferior frontal gyrus, left postcentral gyrus, left amygdala, the left culmen of the cerebellum,
right putamen, and right fusiform gyrus (Fig. 5A). Enhanced responses to food-related stimuli
can be found in many of these regions and responses in some regions have been associated with
weight gain (*35*, *36*). However, in these instances the relationship is correlative. For example,
amygdala response to food related stimuli when sated, but not hungry, correlates with 12-month
weight gain (*37*). This suggests that amygdala responsivity to food cues in the absence of hunger
is predictive of overeating and weight gain, but it is not a direct test of the hypothesis because the
amygdala response could be epiphenomenal. Likewise, response in the right putamen, during
milkshake consumption is predictive of weight change with direction depending upon genotype
(*38*). The current results go beyond these findings. CPM presents as a strength its cross-validated
framework based on data-driven model optimization. Here we were able to identify brain
features that predict and therefore may constitute a brain biomarker of waist circumference.
This result raises the question as to how the CPM we have identified contributes to the
development of increased waist circumference and conversely, how and why waist
circumference specifically influences the identified brain "fingerprint". Notably, although waist
circumference is the best adiposity predictor of glucose control, inspection of the anatomy of the
CPMs identified for waist circumference and fasting insulin, a marker of insulin resistance,
reveals that they are largely distinct (Fig. 5, 6) (*39* – *41*). In particular, the CPMs for waist
circumference and fasting insulin showed minimal overlap in the highest degree nodes. This
suggests there are distinct neuropathophysiological features. The distinction between the CPMs
is best captured by their relative polarity, or the number of positive versus negative edges

between networks. The cerebellar and subcortical networks show similar patterns of connectivity
while patterns in medial-frontal, frontoparietal, default mode, motor, visual, and cingulo-
opercular networks largely differ. Overlap between cerebellar and subcortical networks for
adiposity and metabolic dysfunction suggests that these regions may play an integral role in the
pathophysiology of obesity. These observations support future efforts to uncover the link
between the distinct networks and their contributions to the causes and consequences of obesity.
Our findings also have clinical implications. One important application of neuroimaging in
clinical practice, is to enable the prediction of clinical outcomes using brain structure, activity, or
connectivity (42). Previous methods in predicting brain-behavior relationships tended to be
correlational, and thus would overfit to data and fail to generalize to novel individuals (15). As
indicated above, CPM presents as a strength its cross-validated framework, which suggests that
the predictive features for waist circumference and fasting insulin are generalizable within novel
subjects in this dataset. In other words, it is a window into the brain features that contribute to
predicting adiposity and metabolic outcomes and as such may constitute biomarkers to index risk
or the efficacy of therapeutic interventions.
Further, to apply such a finding to clinical practice, network-based prediction models of obesity
must be shown to generalize across samples, populations, and experimental settings (42). A
promising avenue to then implement CPM as a tool for clinical use is to use real-time fMRI
biofeedback to change connectivity patterns or activation profiles within predictive networks
across subjects. Similar approaches have been successfully used in previous studies, namely
those attempting to alleviate contamination anxiety, enhance emotion regulation capabilities, and
mitigate auditory verbal hallucinations in schizophrenia, among others (43 – 45). Another
approach, similar to that of Rosenberg et al., 2016 (46), is to treat overweight and obesity using a
particular intervention and observe how the predictive networks change. We predict that more
efficacious treatments will reduce network strength within networks that predict high waist
circumference and fasting insulin, and increase the strength of antithetical networks. Recent
work suggests that networks obtained using CPM can predict abstinence during treatment for
cocaine dependence (47). If the obtained CPMs are relevant to the pathophysiology of obesity, it
is thus possible that they may predict weight loss during an intervention, which would be useful
for obese individuals seeking treatment. Finally, not only is waist circumference better than BMI
and other adiposity measures at predicting a variety of adverse health-related outcomes, but it is
also better predicted from whole-brain functional connectivity using CPM. This suggests that
including waist circumference as a measure of interest in a clinical setting is imperative to
accurately quantify the extent and progression of obesity.
Limitations and future directions
This study has a number of limitations to address. First, the sample size was relatively limited for
a study implementing CPM. It is quite possible that with more data we would be able to build a
more informative predictive model. Second, we only include overweight and obese individuals in
our dataset due to the nature and design of the longitudinal study for which this data was
originally collected. It is thus possible that these CPMs for waist circumference and fasting
insulin do not generalize to lean individuals. However, since biomarkers are needed to evaluate
therapeutic interventions it can be argued that an overweight/obese sample is most appropriate.
Third, because the study design was not specifically optimized for use in this analysis, it remains
to be seen whether inclusion of tasteless blocks reduced our predictive power. Tasteless may be
more similar to a classic rest condition than milkshake. Across all studies implementing CPM,
resting state is least able to maximize detection of individual differences (19). Fourth, we did not

use DEXA or whole-body MRI, which are considered the optimal measurement devices to
determine adiposity (*48*). Although, waist circumference and fasting insulin are more practical
measures, future work should validate our observations using the gold standard measures.
Finally, longitudinal studies are necessary to determine whether the obtained predictive networks
arise as a cause or consequence of increased fasting insulin and waist circumference.

Conclusions
Our analyses reveal that by characterizing whole brain connectivity during consumption of a
palatable energy-dense milkshake it is possible to predict an individual's waist circumference but
not BMI or percent body fat. This finding suggests that waist circumference is the phenotypic
marker of obesity that is most closely related to brain function during food consumption.
However, whether this brain biomarker is a cause or a consequence of waist size is unknown. We
were also able to predict fasting insulin but not glucose or HOMA-IR. Notably, although fasting
insulin reflects insulin resistance (*49*) and waist circumference has been shown to correlate with
insulin resistance, the CPMs predictive of waist circumference and fasting insulin were largely
distinct.
302
Materials and Methods
Participants
We recruited 100 overweight and obese individuals using media, flyer advertisements, and the
Yale Center for Clinical Investigation Help Us Discover database. Subjects were recruited to
participate in an ongoing clinical trial (NCT 01976156), but all related fMRI scans occurred at
baseline prior to any experimental manipulations. We screened participants via phone or online
to ensure they met the following criteria: right-handed, English-speaking, 18-45 years of age,
body mass index (BMI) of 25 or higher, nonsmokers, no psychiatric or neurological impairments
(including drug addiction, eating disorders), no obesity-related diseases such as type-2 diabetes,
no food allergies, no chemosensory impairments, and no MRI contraindications (pregnancy,
claustrophobia, metal implants, or prior head injury with loss of consciousness). In total, 67
participants (mean age = 30.7yrs, mean BMI = 33.3 ± 6.5) met these criteria and underwent
fMRI procedures. All participants provided written informed consent according to our protocol
that was approved by the Yale University School of Medicine Human Investigation Committee.
Measures
This study was part of a larger clinical trial (NCT 01976156). Below we outline the procedures
related to the baseline fMRI assessment, on which the analyses in this study are based.
Physical Characteristics and Diet
Participants arrived at the lab on the week of fMRI scanning for anthropometric measurements,
which included body weight, height, waist size, hip size, and adiposity. Adiposity, as described
by body fat percentage (BF%), was measured using air displacement plethysmography
(BodPod). Waist/Hip ratio and BMI (kg/cm^2) were subsequently calculated. Additional
participant characteristics are shown in fig. S1.
Bloods
Following an overnight fast of 10-12 hours, blood samples were collected from 59 out of 67
subjects. We were unable to collect bloods on eight subjects due to venipuncture difficulty.
Samples were collected, centrifuged immediately, and then frozen at either 80 °C (insulin) or 20
°C (glucose). Insulin protein concentrations were subsequently measured using a
radioimmunoassay kit (APLCO insulin ELISA Catalog #80-INSHU-E01.1, Salem, NH) and
plasma glucose concentrations were determined using a glucose oxidase analyzer (YSI

Incorporated, Yellow Springs, OH). Insulin sensitivity was calculated using the homeostasis
model of insulin resistance, or HOMA-IR, which is based on the following formula: [insulin
(mU/L) x glucose (mmol/L)]/22.5 (*50*).
Image protocols
All fMRI scans were scheduled during the week of blood draws, sometime between 8:00 AM
and 12:00 PM. Participants were instructed to arrive neither full nor hungry, and adherence to
these instructions were confirmed using internal state ratings. Structural and functional imaging
data were then collected on a Siemens 3.0T TIM Trio scanner at the Yale Magnetic Resonance
Research Center (MRRC), using the following scan parameters: high resolution T1-weighted
structural scan repetition time (TR) = 2530ms, echo time (TE) = 3.31ms, slice thickness = 1mm,
flip angle = 7°, field of view (FOV) = 256 x 256, 256 x 256 matrix, 176 slices; functional
milkshake run TR = 2000ms, TE = 20 ms, slice thickness = 3mm, flip angle = 80°, FOV = 220 x
220, matrix = 64 x 64, 40 contiguous slices.
fMRI scans consisted of two 9-minute runs that were divided into blocks of 4-8 cued
presentations of milkshake or tasteless solutions. These solutions were delivered in quantities of
0.5 mL over a 2s period followed by 7s to swallow. Different milkshake flavors were used to
reduce the possibility of stimulus habituation. After each block, participants rinsed their mouths
with 0.5mL of deionized water over a period of 2s and were given 3s to swallow. Blocks varied
in length from 36s to 72s and order of presentation was counterbalanced.
Stimuli
Milkshake stimuli were presented as two flavors, chocolate and strawberry. Chocolate
milkshakes were made by combining 12 fl oz of whole milk with equal parts Garelick Farms
Chug chocolate milkshake and Garelick Farms Chug cookies and cream milkshake. The
strawberry milkshake was made by combining 32 fl oz of whole milk with 6 fl oz of Hershey's
strawberry syrup. One subject did not like the taste of strawberry and instead was given a vanilla
milkshake created by mixing 32 fl oz whole milk with 6 fl oz of Eclipse vanilla syrup.
Milkshakes were presented in an interleaved manner to reduce the possibility of sensory
adaptation. Tasteless solutions were chosen individually by each participant, based on what they
perceived "tastes most like nothing." Tasteless solutions were one of either 25mM KCl /2.5mM
NaH CO3, 18.75mM KCl/1.875mM NaHCO3, 12.5mM KCl/1.25mM NaHCO3, 6.25mM
KCl/0.625mM NaHCO3, or 3.13mM KCl/0.313mM NaHCO3, dissolved in distilled water.
Stimulus Delivery
0.5 mL of milkshake was delivered into the mouth over 2s using an MRI-compatible gustometer
system. The gustometer system (*51*) is a portable device controlling 11 programmable BS-8000
syringe pumps (Braintree scientific, Braintree, Ma). Pumps are loaded with 60 mL syringes filled
with milkshake, which are infused into 25ft of Tygon beverage tubing (Saint-Gobain
Performance Plastics, Akron, OH) connected a Teflon gustatory manifold on the MRI headcoil.
The manifold consists of multiple channels that converge at a central point to allow delivery of a
liquid stimulus to the tongue tip.
Preprocessing
Neuroimaging data were preprocessed using SPM12 (Statistical Parametric Mapping, Wellcome
Department of Imaging Neuroscience, London, UK) and BioImage Suite (*52*). Participants'
functional images were corrected for geometric distortions resulting from susceptibility-induced
field inhomogeneities. The FieldMap toolbox in SPM12 was used to calculate static distortions
(*53*). Corrections were then made for both static distortions and changes in these distortions
resulting from head motion (*54, 55*). Images were realigned, unwarped, and coregistered to each

participant's anatomical T1 image and then processed using a method combining segmentation,
bias correction, and spatial normalization (*56*). Functional images were slice-time corrected,
normalized to a standard MNI template, and smoothed using a 6 mm FWHM isotropic Gaussian
kernel. Normalized, motion-corrected images were further preprocessed using BioImage Suite.
These additional steps included regression of the mean time courses of global signal, CSF, and
white matter, in addition to removal of the linear trend and low-pass filtering.

Construction of predictive networks
We performed Connectome-based Predictive Modeling (CPM; *14*, *15*, *18*) to determine whether
task-based functional connectivity during milkshake receipt can predict individual differences in
anthropometric measurements. To accomplish this, we first used a whole-brain functional atlas
(*57*) to parcellate the brains of each subject into 268 nodes (i.e spatial ROIs) that maximize the
similarity of voxel-wise time courses within each node. 268x268 connectivity matrices for each
run were subsequently created in BioImage Suite by averaging the BOLD signal of each voxel
within nodes (*52*). Connectivity matrices were Fisher Transformed to convert the skewed
distribution of Pearson r (or Spearman rho) values to an approximately normal distribution.
Matrices were then averaged across runs to create a mean matrix for each subject, which were
then collapsed across subjects to create 268x268xN-subject matrices.
In the N-1 training set, CPM uses a linear regression (i.e. Pearson's r or Spearman's Rho) to
correlate each edge (i.e. connection) in each connectivity matrix to a behavioral measure of
interest. Each edge, as a result, is then associated with a particular p-value and the most relevant
edges are selected for further use via thresholding (P = 0.01). Edges that are positively correlated
with behavior constitute a "positive prediction network," while edges whose strengths are
negatively correlated with behavior constitute a "negative prediction network." Single-subject
summary values are then calculated by separately summing the edge-strengths of positive and
negative prediction networks (*15*). These positive and negative "network strengths" are used in a
linear model with behavior and applied prospectively to the left-out subject's network strengths
to generate predicted behavioral scores.

Permutation Testing
To determine the significance of the obtained correlations between observed and predicted
behavioral scores, we conducted 1000 repetitions of CPM using randomly shuffled observed
scores to generate predicted scores and networks. The 1000 correlation coefficients obtained
from permutation testing were used to comprise a null distribution against which the correlation
coefficients in the Results section were tested for significance. Final p-values were calculated by
determining the number of permutations, out of 1000, that generated correlation coefficients
larger than those reported in Results.

Visualization
Figures were constructed using BioImage Suite (*52*) and custom scripts in MATLAB® 2016b.
Anatomical locations were isolated using the Wake Forest PickAtlas Toolbox v. 3.0.5 (Wake
Forest University, Winston-Salem, NC, USA) in SPM12 and the Automated Anatomical
Labeling (AAL) atlas (*58*).
419
420　**Supplementary Materials**
421　Table S1. Participant Characteristics of Full Sample.
422　Table S2. FTF-Corrected Correlations and *P*-Values.
423
424

References and Notes:

1. S. O'Rahilly, I. S. Farooqi, Human obesity: A heritable neurobehavioral disorder that is highly sensitive to environmental conditions. *Diabetes* **57**, 2905-2910 (2008).

2. T. M. Frayling, N. J. Timpson, M. N. Weedon, E. Zeggini, R. M. Freathy, C. M. Lindgren, J. R. B. Perry, K. S. Elliott, H. Lango, N. W. Rayner, B. Shields, L. W. Harries, J. C. Barrett, S. Ellard, C. J. Groves, B. Knight, A. Patch, A. R. Ness, S. Ebrahim, D. A. Lawlor, S. M. Ring, Y. Ben-Shlomo, M. Jarvelin, U. Sovio, A. J. Bennett, D. Melzer, L. Ferrucci, R. J. F. Loos, I. Barroso, N. J. Wareham, F. Karpe, K. R. Owen, L. R. Cardon, M. Walker, G. A. Hitman, C. N. A. Palmer, A. S. F. Doney, A. D. Morris, G. Davey Smith, The Wellcome Trust Case Control Consortium, A. T. Hattersley, M. I. McCarthy, A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. *Science* **316**, 889-894 (2007).

3. A. Scuteri, S. Sanna, W. M. Chen, M. Uda, G. Albai, J. Strait, S. Najjar, R. Nagaraja, M. Orrú, G. Usala, M. Dei, S. Lai, A. Maschio, F. Busonero, A. Mulas, G. B. Ehret, A. A. Fink, A. B. Weder, R. S. Cooper, P. Galan, A. Chakravarti, D. Schlessinger, A. Cao, E. Lakatta, G. R. Abecasis, Genome-wide association scan shows genetic variants in the FTO gene are associated with obesity-related traits. *PLoS Genetics* **3**, 1200-1210 (2007).

4. M. E. Hess, S. Hess, K. D. Meyer, L. A. Verhagen, L. Koch, H. S. Brönneke, M. O. Dietrich, S. D. Jordan, Y. Saletore, O. Elemento, B. F. Belgardt, T. Franz, T. L. Horvath, U. Rüther, S. R. Jaffrey, P. Kloppenburg, & J. C. Brüning, The fat mass and obesity associated gene (FTO) regulates activity of the dopaminergic midbrain circuitry. *Nature Neuroscience* **16**, 1042-1048 (2013).

5. S. Smemo, J. J. Tena, K. H. Kim, E. R. Gamazon, N. J. Sakabe, C. Gómez-Marín, I. Aneas, F. L. Credido, D. R. Sobreira, N. F. Wasserman, J. Lee, V. Puviindran, D. Tam, M. Shen, J. Son, N. Alizadeh Vakili, H. Sung, S. Naranjo, R. D. Acemel, M. Manzanares, A. Nagy, N. J. Cox, C. Hui, J. Gomez-Skarmeta, M. A. Nóbrega, Obesity-associated variants within FTO form long-range functional connections with IRX3. *Nature* **507**, 371-375 (2014).

6. M. Sevgi, L. Rigoux, A. B. Kühn, J. Mauer, L. Schilbach, M. E. Hess, T. O. J. Gruendler, M. Ullsperger, K. E. Stephan, J. C. Brüning, M. Tittgemeyer, An obesity-predisposing variant of the FTO gene regulates D2R-dependent reward learning. *Journal of Neuroscience* **35**, 12584-12592 (2015).

7. R. J. F. Loos, G. S. H. Yeo, The bigger picture of FTO—the first GWAS-identified obesity gene. *Nature Reviews Endocrinology* **10**, 51-61 (2014).

8. A. E. Locke, B. Kahali, S. I. Berndt, A. E. Justice, T. H. Pers, F. R. Day, C. Powell, S. Vedantam, M. L. Buchkovich, J. Yang, D. C. Croteau-Chonka, T. Esko, T. Fall, T. Ferreira, S. Gustafsson, Z. Kutalik, J. Luan, R. Mägi, J. C. Randall, T. W. Winkler, A. R. Wood, T. Workalemahu, J. D. Faul, J. A. Smith, J. Zhao, W. Zhao, J. Chen, R. Fehrmann, Å. K. Hedman, J. Karjalainen, E. M. Schmidt, D. Absher, N. Amin, D. Anderson, M. Beekman, J. L. Bolton, J. L. Bragg-Gresham, S. Buyske, A. Demirkan, G. Deng, G. B. Ehret, B. Feenstra, M. F. Feitosa,

K. Fischer, A. Goel, J. Gong, A. U. Jackson, S. Kanoni, M. E. Kleber, K. Kristiansson, U. Lim,
V. Lotay, M. Mangino, I. Mateo Leach, C. Medina-Gomez, S. E. Medland, M. A. Nalls, C. D.
Palmer, D. Pasko, S. Pechlivanis, M. J. Peters, I. Prokopenko, D. Shungin, A. Stančáková, R. J.
Strawbridge, Y. Sung, T. Tanaka, A. Teumer, S. Trompet, S. W. van der Laan, J. van Setten, J.
V. Van Vliet-Ostaptchouk, Z. Wang, L. Yengo, W. Zhang, A. Isaacs, E. Albrecht, J. Ärnlöv, G.
M. Arscott, A. P. Attwood, S. Bandinelli, A. Barrett, I. N. Bas, C. Bellis, A. J. Bennett, C. Berne,
R. Blagieva, M. Blüher, S. Böhringer, L. L. Bonnycastle, Y. Böttcher, H. A. Boyd, M.
Bruinenberg, I. H. Caspersen, Y. Ida Chen, R. Clarke, E. Warwick Daw, A. J. M. de Craen, G.
Delgado, M. Dimitriou, A. S. F. Doney, N. Eklund, K. Estrada, E. Eury, L. Folkersen, R. M.
Fraser, M. E. Garcia, F. Geller, V. Giedraitis, B. Gigante, A. S. Go, A. Golay, A. H. Goodall, S.
D. Gordon, M. Gorski, H. Grabe, H. Grallert, T. B. Grammer, J. Gräßler, H. Grönberg, C. J.
Groves, G. Gusto, J. Haessler, P. Hall, T. Haller, G. Hallmans, C. A. Hartman, M. Hassinen, C.
Hayward, N. L. Heard-Costa, Q. Helmer, C. Hengstenberg, O. Holmen, J. Hottenga, A. L.
James, J. M. Jeff, Å. Johansson, J. Jolley, T. Juliusdottir, L. Kinnunen, W. Koenig, M.
Koskenvuo, W. Kratzer, J. Laitinen, C. Lamina, K. Leander, N. R. Lee, P. Lichtner, L. Lind, J.
Lindström, K. Sin Lo, S. Lobbens, R. Lorbeer, Y. Lu, F. Mach, P. K. E. Magnusson, A.
Mahajan, W. L. McArdle, S. McLachlan, C. Menni, S. Merger, E. Mihailov, L. Milani, A.
Moayyeri, K. L. Monda, M. A. Morken, A. Mulas, G. Müller, M. Müller-Nurasyid, A. W. Musk,
R. Nagaraja, M. M. Nöthen, I. M. Nolte, S. Pilz, N. W. Rayner, F. Renstrom, R. Rettig, J. S.
Ried, S. Ripke, N. R. Robertson, L. M. Rose, S. Sanna, H. Scharnagl, S. Scholtens, F. R.
Schumacher, W. R. Scott, T. Seufferlein, J. Shi, A. Vernon Smith, J. Smolonska, A. V. Stanton,
V. Steinthorsdottir, K. Stirrups, H. M. Stringham, J. Sundström, M. A. Swertz, A. J. Swift, A.
Syvänen, S. Tan, B. O. Tayo, B. Thorand, G. Thorleifsson, J. P. Tyrer, H. Uh, L. Vandenput, F.
C. Verhulst, S. H. Vermeulen, N. Verweij, J. M. Vonk, L. L. Waite, H. R. Warren, D.
Waterworth, M. N. Weedon, L. R. Wilkens, C. Willenborg, T. Wilsgaard, M. K. Wojczynski, A.
Wong, A. F. Wright, Q. Zhang, The LifeLines Cohort Study, E. P. Brennan, M. Choi, Z. Dastani,
A. W. Drong, P. Eriksson, A. Franco-Cereceda, J. R. Gådin, A. G. Gharavi, M. E. Goddard, R.
E. Handsaker, J. Huang, F. Karpe, S. Kathiresan, S. Keildson, K. Kiryluk, M. Kubo, J. Lee, L.
Liang, R. P. Lifton, B. Ma, S. A. McCarroll, A. J. McKnight, J. L. Min, M. F. Moffatt, G. W.
Montgomery, J. M. Murabito, G. Nicholson, D. R. Nyholt, Y. Okada, J. R. B. Perry, R. Dorajoo,
E. Reinmaa, R. M. Salem, N. Sandholm, R. A. Scott, L. Stolk, A. Takahashi, T. Tanaka, F. M.
van 't Hooft, A. A. E. Vinkhuyzen, H. Westra, W. Zheng, K. T. Zondervan, The ADIPOGen
Consortium, The AGEN-BMI Working Group, The CARDIOGRAMplusC4D Consortium, The
CKDGen Consortium, The GLGC, The ICBP, The MAGIC Investigators, The MuTHER
Consortium, The MIGen Consortium, The PAGE Consortium, The ReproGen Consortium, The
GENIE Consortium, The International Endogene Consortium, A. C. Heath, D. Arveiler, S. J. L.
Bakker, J. Beilby, R. N. Bergman, J. Blangero, P. Bovet, H. Campbell, M. J. Caulfield, G.
Cesana, A. Chakravarti, D. I. Chasman, P. S. Chines, F. S. Collins, D. C. Crawford, L. A.
Cupples, D. Cusi, . Danesh, U. de Faire, H. M. den Ruijter, A. F. Dominiczak, R. Erbel, J.
Erdmann, J. G. Eriksson, M. Farrall, S. B. Felix, E. Ferrannini, J. Ferrières, I. Ford, N. G.
Forouhi, T. Forrester, O. H. Franco, R. T. Gansevoort, P. V. Gejman, C. Gieger, O. Gottesman,
V. Gudnason, U. Gyllensten, A. S. Hall, T. B. Harris, A. T. Hattersley, A. A. Hicks, L. A.
Hindorff, A. D. Hingorani, A. Hofman, G. Homuth, G. K. Hovingh, S. E. Humphries, S. C.
Hunt, E. Hyppönen, T. Illig, K. B. Jacobs, M. Jarvelin, K. Jöckel, B. Johansen, P. Jousilahti, J.
W. Jukema, A. M. Jula, J. Kaprio, J. J. P. Kastelein, S. M. Keinanen-Kiukaanniemi, L. A.
Kiemeney, P. Knekt, J. S. Kooner, C. Kooperberg, P. Kovacs, A. T. Kraja, M. Kumari, J.

Kuusisto, T. A. Lakka, C. Langenberg, L. Le Marchand, T. Lehtimäki, V. Lyssenko, S. Männistö, A. Marette, T. C. Matise, C. A. McKenzie, B. McKnight, F. L. Moll, A. D. Morris, A. P. Morris, J. C. Murray, M. Nelis, C. Ohlsson, A. J. Oldehinkel, K. K. Ong, P. A. F. Madden, G. Pasterkamp, J. F. Peden, A. Peters, D. S. Postma, P. P. Pramstaller, J. F. Price, L. Qi, O. T. Raitakari, T. Rankinen, D. C. Rao, T. K. Rice, P. M. Ridker, J. D. Rioux, M. D. Ritchie, I. Rudan, V. Salomaa, N. J. Samani, J. Saramies, M. A. Sarzynski, H. Schunkert, P. E. H. Schwarz, P. Sever, A. R. Shuldiner, J. Sinisalo, R. P. Stolk, K. Strauch, A. Tönjes, D. Trégouët, A. Tremblay, E. Tremoli, J. Virtamo, M. Vohl, U. Völker, G. Waeber, G. Willemsen, J. C. Witteman, M. Carola Zillikens, L. S. Adair, P. Amouyel, F. W. Asselbergs, Themistocles L. Assimes, M. Bochud, B. O. Boehm, E. Boerwinkle, S. R. Bornstein, E. P. Bottinger, C. Bouchard, S. Cauchi, J. C. Chambers, S. J. Chanock, R. S. Cooper, P. I. W. de Bakker, G. Dedoussis, L. Ferrucci, P. W. Franks, P. Froguel, L. C. Groop, C. A. Haiman, A. Hamsten, J. Hui, D. J. Hunter, K. Hveem, R. C. Kaplan, M. Kivimaki, D. Kuh, M. Laakso, Y. Liu, N. G. Martin, W. März, M. Melbye, A. Metspalu, S. Moebus, P. B. Munroe, I. Njølstad, B. A. Oostra, C. N. A. Palmer, N. L. Pedersen, M. Perola, L. Pérusse, U. Peters, C. Power, T. Quertermous, R. Rauramaa, F. Rivadeneira, T. E. Saaristo, D. Saleheen, N. Sattar, E. E. Schadt, D. Schlessinger, P. Eline Slagboom, H. Snieder, T. D. Spector, U. Thorsteinsdottir, M. Stumvoll, J. Tuomilehto, A. G. Uitterlinden, M. Uusitupa, P. van der Harst, M. Walker, H. Wallaschofski, N. J. Wareham, H. Watkins, D. R. Weir, H. Wichmann, J. F. Wilson, P. Zanen, I. B. Borecki, P. Deloukas, C. S. Fox, I. M. Heid, J. R. O'Connell, D. P. Strachan, K. Stefansson, C. M. van Duijn, G. R. Abecasis, L. Franke, T. M. Frayling, M. I. McCarthy, P. M. Visscher, A. Scherag, C. J. Willer, M. Boehnke, K. L. Mohlke, C. M. Lindgren, J. S. Beckmann, I. Barroso, K. E. North, E. Ingelsson, J. N. Hirschhorn, R. J. F. Loos, E. K. Speliotes, Genetic studies of body mass index yield new insights for obesity biology. *Nature* **518**, 197-206 (2015).

9. WHO, Physical Status: The Use and Interpretation of Anthropometry: Report of a World Health Organization (WHO) Expert Committee. Geneva, Switzerland: World Health Organization (1995).

10. V. Dudeja, A. Misra, R. M. Pandey, G. Devina, G. Kumar, N. K. Vikram, BMI does not accurately predict overweight in Asian Indians in northern India. *British Journal of Nutrition* **86,** 105-112 (2001).

11. J. J. Ode, J. M. Pivarnik, M. J. Reeves, J. L. Knous, Body mass index as a predictor of percent fat in college athletes and nonathletes. *Medicine and Science in Sports and Exercise* **39,** 403-409 (2007).

12. A. H. Kissebah, N. Vydelingum, R. Murray, D. J. Evans, R. K. Kalkhoff, P. W. Adams, Relation of body fat distribution to metabolic complications of obesity. *The Journal of Clinical Endocrinology & Metabolism* **54**, 254-260 (1982).

13. B. H. Goodpaster, S. Krishnaswami, T. B. Harris, A. Katsiaras, S. B. Kritchevsky, E. M. Simonsick, M. Nevitt, P. Holvoet, A. B. Newman, Obesity, regional body fat distribution, and the metabolic syndrome in older men and women. *Archives of Internal Medicine* **165**, 777-783 (2005).

563 14. E. S. Finn, X. Shen, D. Scheinost, M. D. Rosenberg, J. Huang, M. M. Chun, X.
564 Papademetris, R. T. Constable, Functional connectome fingerprinting: identifying individuals
565 using patterns of brain connectivity. *Nature Neuroscience* **18**, 1664-1671 (2015).
566
567 15. X. Shen, E. S. Finn, D. Scheinost, M. D. Rosenberg, M. M. Chun, X. Papademetris, R. T.
568 Constable, Using connectome-based predictive modeling to predict individual behavior from
569 brain connectivity. *Nature Protocols* **12,** 506-518 (2017).
570
571 16. W. T. Hsu, M. D. Rosenberg, D. Scheinost, R. T. Constable, M. M. Chun, Resting-state
572 functional connectivity predicts neuroticism and extraversion in novel individuals. *Social
573 Cognitive and Affective Neuroscience* **13,** 224-232 (2018).
574
575 17. R. E. Beaty, Y. N. Kenett, A. P. Christensen, M. D. Rosenberg, M. Benedek, Q. Chen, A.
576 Fink, J. Qiu, T. R. Kwapil, M. J. Kane, P. J. Silvia, Robust prediction of individual creative
577 ability from brain functional connectivity. *Proceedings of the National Academy of Sciences* **115**,
578 1087-1092 (2018).
579
580 18. M. D. Rosenberg, E. S. Finn, D. Scheinost, X. Papademetris, X. Shen, R. T. Constable, M.
581 M. Chun, A neuromarker of sustained attention from whole-brain functional connectivity. *Nature
582 Neuroscience* **19***,* 165-171 (2016).
583
584 19. A. S. Greene, S. Gao, D. Scheinost, R. T. Constable, Task-induced brain state manipulation
585 improves prediction of individual traits. *Nature Communications* **9**, 2807 (2018).
586
587 20. S. Noble, M. N. Spann, F. Tokoglu, X. Shen, R. T. Constable, D. Scheinost, Influences on
588 the test–retest reliability of functional connectivity MRI and its relationship with behavioral
589 utility. *Cerebral Cortex* **27**, 5415-5429 (2017).
590
591 21. T. D. Satterthwaite, M. A. Elliott, R. T. Gerraty, K. Ruparel, J. Loughead, M. E. Calkins, S.
592 B. Eickhoff, H. Hakonarson, R. C. Gur, R. E. Gur, D. H. Wolf, An improved framework for
593 confound regression and filtering for control of motion artifact in the preprocessing of resting-
594 state functional connectivity data. *Neuroimage* **64**, 240-256 (2013).
595
596 22. K. Hodgson, R. A. Poldrack, J. E. Curran, E. E. Knowles, S. Mathias, H. H. Göring, N. Yao,
597 R. L. Olvera, P. T. Fox, L. Almasy, R. Duggirala, D. M. Barch, J. Blangero, D.C. Glahn, Shared
598 genetic factors influence head motion during MRI and body mass index. *Cerebral Cortex* **27**,
599 5539-5546 (2016).
600
601 23. N. Motamed, B. Rabiee, H. Keyvani, G. R. Hemasi, M. Khonsari, F. S. Saeedian, M. Maadi,
602 F. Zamani, The best obesity indices to discriminate type 2 diabetes mellitus. *Metabolic Syndrome
603 and Related Disorders* **14**, 249-253 (2016).
604
605 24. R. N. Bergman, D. Stefanovski, T. A. Buchanan, A. E. Sumner, J. C. Reynolds, N. G.
606 Sebring, A. H. Xiang, R. M. Watanabe, A better index of body adiposity. *Obesity* **19**, 1083-1089
607 (2012).
608

25. M. Ashwell, P. Gunn, S. Gibson, Waist-to-height ratio is a better screening tool than waist
circumference and BMI for adult cardiometabolic risk factors: systematic review and
meta-analysis. *Obesity Reviews* **13**, 275-286 (2012).

26. D. C. Seo, S. Choe, M. R. Torabi, Is waist circumference ≥ 102/88 cm better than body mass
index ≥ 30 to predict hypertension and diabetes development regardless of gender, age group,
and race/ethnicity? Meta-analysis. *Preventive Medicine* **97**, 100-108 (2017).

27. S. K. Tanamas, W. L. Ng, K. Backholer, A. Hodge, P. Z. Zimmet, A. Peeters, Quantifying
the proportion of deaths due to body mass index-and waist circumference-defined obesity.
Obesity **24**, 735-742 (2016).

28. K. Dimitriadis, C. Tsioufis, A. Mazaraki, I. Liatakis, E. Koutra, A. Kordalis, A. Kasiakogias,
D. Flessas, N. Tentolouris, D. Tousoulis, Waist circumference compared with other obesity
parameters as determinants of coronary artery disease in essential hypertension: A 6-year follow-
up study. *Hypertension Research* **39**, 475-479 (2016).

29. F. Kurth, J. G. Levitt, O. R. Phillips, E. Luders, R. P. Woods, J. C. Mazziotta, A. W. Toga,
K. L. Narr, Relationships between gray matter, body mass index, and waist circumference in
healthy adults. *Human Brain Mapping* **34**, 1737-1746 (2013).

30. D. Janowitz, K. Wittfeld, J. Terock, H. J. Freyberger, K. Hegenscheid, H. Völzke, M. Habes,
N. Hosten, N. Friedrich, M. Nauck, G. Domanska, H. J. Grabe, Association between waist
circumference and gray matter volume in 2344 individuals from two adult community-based
samples. *NeuroImage* **122,** 149-157 (2015).

31. S. Luo, A. Romero, T. C. Adam, H. H. Hu, J. Monterosso, K. A. Page, Abdominal fat is
associated with a greater brain reward response to high-calorie food cues in hispanic women.
Obesity **21***,* 2029-2036 (2013).

32. E. Green, A. Jacobson, L. Haase, C. Murphy, Reduced nucleus accumbens and caudate
nucleus activation to a pleasant taste is associated with obesity in older adults. *Brain Research*
1386, 109-117 (2011).

33. S. Wallner-Liebmann, K. Koschutnig, G. Reishofer, E. Sorantin, B. Blaschitz, R. Kruschitz,
H. F. Unterrainer, R. Gasser, F. Freytag, C. Bauer-Denk, H. Mangge, Insulin and hippocampus
activation in response to images of high-calorie food in normal weight and obese adolescents.
Obesity **18**, 1552-1557 (2010).

34. I. Janssen, P. T. Katzmarzyk, R. Ross, Waist circumference and not body mass index
explains obesity-related health risk. *The American Journal of Clinical Nutrition* **79**, 379-384
(2004).

35. A. M. Jastreboff, R. Sinha, C. Lacadie, D. M. Small, R. S. Sherwin, M. N. Potenza, Neural
correlates of stress-and food cue–induced food craving in obesity: Association with insulin
levels. *Diabetes Care* **36**, 394-402 (2013).

36. H. R. Noori, A. C. Linan, R. Spanagel, Largely overlapping neuronal substrates of reactivity
to drug, gambling, food and sexual cues: A comprehensive meta-analysis. *European*
Neuropsychopharmacology **26**, 1419-1430 (2016).

37. X. Sun, N. B. Kroemer, M. G. Veldhuizen, A. E. Babbs, I. E. de Araujo, D. R. Gitelman, R.
S. Sherwin, R. Sinha, D. M. Small, Basolateral amygdala response to food cues in the absence of
hunger is associated with weight gain susceptibility. *Journal of Neuroscience* **35**, 7964-7976
(2015).

38. E. Stice, S. Spoor, C. Bohon, D. M. Small, Relation between obesity and blunted striatal
response to food is moderated by TaqIA A1 allele. *Science* **322**, 449-452 (2008).

39. Q. M. Nguyen, S. R. Srinivasan, J. H. Xu, W. Chen, G. S. Berenson, Distribution and
cardiovascular risk correlates of hemoglobin A1c in nondiabetic younger adults: the Bogalusa
Heart Study. *Metabolism* **57**, 1487-1492 (2008).

40. G. Vazquez, S. Duval, D. R. Jacobs Jr, K. Silventoinen, Comparison of body mass index,
waist circumference, and waist/hip ratio in predicting incident diabetes: A meta-analysis.
Epidemiologic Reviews **29**, 115-128 (2007).

41. Y. Wang, E. B. Rimm, M. J. Stampfer, W. C. Willett, F. B. Hu, Comparison of abdominal
adiposity and overall obesity in predicting risk of type 2 diabetes among men. *The American*
Journal of Clinical Nutrition **81**, 555-563 (2005).

42. C. W. Woo, L. J. Chang, M. A. Lindquist, T. D. Wager, Building better biomarkers: brain
models in translational neuroimaging. *Nature Neuroscience* **20**, 365-377 (2017).

43. D. Scheinost, T. Stoica, J. Saksa, X. Papademetris, R. T. Constable, C. Pittenger, M.
Hampson, Orbitofrontal cortex neurofeedback produces lasting changes in contamination anxiety
and resting-state connectivity. *Translational Psychiatry* **3**, e250 (2013).

44. Y. Koush, D. E. Meskaldji, S. Pichon, G. Rey, S. W. Rieger, D. E. J. Linden, D. Van De
Ville, P. Vuilleumier, F. Scharnowski, Learning control over emotion networks through
connectivity-based neurofeedback. *Cerebral Cortex* **27**, 1193-1202 (2015).

45. N. D. Orlov, V. Giampietro, O. O'Daly, S. L. Lam, G. J. Barker, K. Rubia, P. McGuire, S. S.
Shergill, P. Allen, Real-time fMRI neurofeedback to down-regulate superior temporal gyrus
activity in patients with schizophrenia and auditory hallucinations: a proof-of-concept study.
Translational Psychiatry **8**, 46 (2018).

46. M. D. Rosenberg, S. Zhang, W. T. Hsu, D. Scheinost, E. S. Finn, X. Shen, R. T. Constable,
C. R. Li, M. M. Chun, Methylphenidate modulates functional network connectivity to enhance
attention. *Journal of Neuroscience* **36**, 9547-9557 (2016).

47. S. W. Yip, D. Scheinost, M. N. Potenza, K. M. Carroll, Connectome-based prediction of
cocaine abstinence. *American Journal of Psychiatry*, appi-ajp (2019).

48. A. Shuster, M. Patlas, J. H. Pinthus, M. Mourtzakis, The clinical importance of visceral
adiposity: a critical review of methods for visceral adipose tissue analysis. *The British Journal of*
Radiology **85**, 1-10 (2012).

49. B. Singh, A. Saxena, Surrogate markers of insulin resistance: A review. *World Journal of*
Diabetes **1**, 36-47 (2010).

50. D. R. Matthews, J. P. Hosker, A. S. Rudenski, B. A. Naylor, D. F. Treacher, R. C. Turner,
Homeostasis model assessment: insulin resistance and β-cell function from fasting plasma
glucose and insulin concentrations in man. *Diabetologia* **28**, 412-419 (1985).

51. M. G. Veldhuizen, G. Bender, R. T. Constable, D. M. Small, Trying to detect taste in a
tasteless solution: modulation of early gustatory cortex by attention to taste. *Chemical Senses* **32**,
569-581 (2007).

52. A. Joshi, D. Scheinost, H. Okuda, D. Belhachemi, I. Murphy, L. H. Staib, X. Papademetris,
Unified framework for development, deployment and robust testing of neuroimaging algorithms.
Neuroinformatics **9**, 69-84 (2011).

53. C. Hutton, R. Deichmann, R.Turner, J. Andersson, Combined correction for geometric
distortion and its interaction with head movement in fMRI. In *Proc. of the 12th Scientific*
Meeting and Exhibition of the ISMRM **12**, 1084 (2004).

54. J. L. Andersson, C. Hutton, J. Ashburner, R. Turner, K. Friston, Modeling geometric
deformations in EPI time series. *NeuroImage* **13**, 903-919 (2001).

55. C. Hutton, A. Bork, O. Josephs, R. Deichmann, J. Ashburner, R. Turner, Image distortion
correction in fMRI: A quantitative evaluation. *NeuroImage* **16**, 217-240 (2002).

56. J. Ashburner, K. J. Friston, Unified segmentation. *NeuroImage* **26**, 839-851 (2005).

57. X. Shen, F. Tokoglu, X. Papademetris, R. T. Constable, Groupwise whole-brain parcellation
from resting-state fMRI data for network node identification. *NeuroImage* **82**, 403-415 (2013).

58. J. L. Lancaster, M. G. Woldorff, L. M. Parsons, M. Liotti, C. S. Freitas, L. Rainey, P.V.
Kochunov, D. Nickerson, S.A. Mikiten, P. T. Fox, Automated Talairach atlas labels for
functional brain mapping. *Human Brain Mapping* **10**, 120-131 (2000).

Acknowledgements: We would like to thank Corey L Horien and Abigail Greene for their help
with data visualization. **Funding:** This work was supported by the National Institutes of Health
training grant T32NS041228 (M.C.F.); the Yale Medical School Fellowship (M.C.F.); the
National Institutes of Health R01 CA180030 (D.M.S.). **Author Contributions:**
Conceptualization, D.M.S. and M.C.F.; Investigation D.M.S and M.C.F.; Software, R.T.C. and

D.S.; Formal Analysis, M.C.F. and M.J.V.; Visualization, M.C.F. and M.J.V.; Writing – Original
Draft, M.C.F. and M.J.V.; Writing – Review and Editing, M.C.F. and D.M.S.; Supervision,
R.T.C., D.S., and D.M.S., Funding Acquisition, D.M.S.
Competing Interests: The authors declare no competing interests.

Fig. 1. Relationship Between Observed and Predicted Waist Circumference.
Scatterplots relating observed versus predicted waist circumference in positive, negative, and combined networks. Positive networks consist of edges that are positively correlated with behavior, while negative networks are constructed using edges that are negatively correlated with behavior. Combined networks are constructed using a general linear model (GLM). Overall, approximately 13% of the variance in a novel individual's waist circumference can be accounted for using CPM.

Fig. 2. Visualization of Positive and Negative Predictive Networks.
Representations of positive (red) and negative (blue) predictive networks in (**A**). Each semicircle is a different hemisphere of the brain. Connections represent edges between nodes, with nodes organized among ten macroscale regions. Matrices (**B**) show the proportion of positive versus negative edges shared between network pairs among ten canonical networks normalized for network size (MF: medial frontal; FP: frontoparietal; DM: default mode; MO: Motor; V1: visual 1; V2: visual 2; VA: visual association; CO: cingulo-opercular; SC: subcortical; CE: cerebellar).

Fig. 3. Exploratory Analysis of Fasting Insulin
CPM predicts fasting insulin levels (**A**). Positive and negative predictive networks are shown as hemisphere plots in (**B**). Notably, the number of positive and negative edges are highest among cerebellar and frontoparietal regions and the rest of the brain (**C**).

Fig. 4. Difference Matrices of Waist Circumference and Insulin

Matrices express the polarity in the number of positive versus negative edges between canonical networks adjusted for network size. For example, (**A**) shows more edges in the positive network exist between cerebellar and medial frontal nodes, while more edges in the negative network exist for cerebellar-cerebellar connections. Taken together, (**A**) and (**B**) demonstrate the greatest degree of similarity in connectivity patterns of cerebellar and subcortical networks in waist circumference and fasting insulin CPMs.

Fig. 5. Top 10% High-Degree Nodes for Waist Circumference and Fasting Insulin CPMs
Top 10% high-degree nodes (27 nodes total) for waist circumference (**A**) encompass networks spanning the whole brain and include nodes varying in degree from 21 to 7. (**B**) The CPM for fasting insulin is similarly encompassing, with nodes varying in degree from 11 to 3.

Fig. 6. Difference of Top 10% Nodes between Waist Circumference and Fasting Insulin CPMs
(**A**) High-degree nodes (top 10%) participating in the waist circumference CPM but not the fasting insulin CPM (**B**) Nodes in the top 10% participating in the fasting insulin CPM but not the waist circumference CPM.

Table 1. Top 10% High-Degree Nodes in the Waist Circumference Network.

Degree	Node	Region	MNI Coordinates	AAI Label
21	172	L-Parietal	23.5, -31.6, 63.6	Postcentral gyrus
20	228	L-Limbic	-26.8, 2.4, -18.7	Left Amygdala
19	244	L-Cerebellum	-6.5, -50.2, -11.4	Left lobule IV,V Cerebellum
15	124	R-Subcortical	26.6, 6.3, 0.1	Right Putamen
15	59	R-Temporal	43.4, -26.5, -24.6	Right Fusiform
13	226	L-Limbic	-8.8, -42.6, 50.1	Left Midcingulate Area
12	169	L-Insula	-38.7, 8.1, -4.8	Left Insula
12	132	R-Brainstem	6.3, -24.9, -17.5	N/A (Midbrain)
10	255	L-Cerebellum	-6.6, -54.7, -26.3	N/A (Cerebellum)
9	153	L-Prefrontal	-32, 20.5, -16	Left Inferior Frontal Gyrus, Pars Orbitalis
8	251	L-Cerebellum	-10.3, -37.7, -25.1	Left Lobule III of Cerebellum
8	227	L-Limbic	-7.5, -42.1, 13.3	Left Posterior Cingulate Gyrus
8	151	L-Prefrontal	-46.2, 28.2, -7.1	Left Inferior Frontal Gyrus, Pars Orbitalis
8	111	R-Cerebellum	11.7, -84.1, -34.6	Right crus II of Cerebellum
8	101	R-Cerebellum	6.1, -50.7, -12.3	Lobule IV, V of Vermis
8	34	R-Insula	41.8, 5, -7.6	Right Insula
7	242	L-Cerebellum	-30.2, -80.2, -40.3	Left Crus II of Cerebellum
7	216	L-Occipital	-22.1, -66.7, 7.4	Left Calcarine Sulcus
7	215	L-Occipital	-6, -81.2, 12.2	Left Calcarine Sulcus
7	188	L-Temporal	-49.7, 6.4, -15.2	Left Superior Temporal Pole
7	128	R-Subcortical	5.5, -9.7, 5.2	Right Thalamus
7	103	R-Cerebellum	13.7, -40.1, -25.4	Right Lobule III of Cerebellum
7	95	R-Limbic	28, -28.4, -13.7	Right Parahippocampal Gyrus
7	92	R-Limbic	31.2, 3.7, -21.6	Right Superior

				Temporal Pole
7	91	R-Limbic	8.3, -39.9, 48.1	Right Midcingulate Area
7	81	R-Occipital	31.2, -91.8, -10.8	Right Inferior Occipital Lobe
7	58	R-Temporal	40.3, -11.3, -35.8	Right Fusiform Gyrus

Table 2. Top 10% High-Degree Nodes in the Fasting Insulin Network.

Degree	Node	Region	MNI Coordinates	AAL Label
11	238	L-Cerebellum	-37, -52.9, -31.1	Left Crus I of Cerebellum
8	186	L-Temporal	-34.6, 18.6, -32.2	Left Superior Temporal Pole
7	227	L-Limbic	-7.5, -42.1, 13.3	Left Posterior Cingulate Gyrus
7	59	R-Temporal	43.4, -26.5, -24.6	Right Fusiform
6	116	R-Cerebellum	41.9, -64, -49.2	Right Lobule VIIB of Cerebellum
5	103	R=Cerebellum	13.7, -40.1, -25.4	Right Lobule III of Cerebellum
5	102	R-Cerebellum	39.1, -74.9, -29.7	Right Crus I of Cerebellum
5	92	R-Limbic	31.2, 3.7, -21.6	Right Superior Temporal Pole
4	265	L-Brainstem	-5, -21.5, -15.8	N/A (Midbrain)
4	263	L-Subcortical	-4.9, -10.3, 5.8	Left Thalamus
4	208	L-Occipital	-16.8, -84.9, 33	Left Cuneus
4	188	L-Temporal	-49.7, 6.4, -15.2	Left Superior Temporal Pole
4	154	L-Prefrontal	-43, 42, 11	Left Inferior Frontal Gyrus Pars Triangularis
4	139	L-Prefrontal	-18.2, 57, -14.3	Left Superior Frontal Gyrus, Orbital part
4	99	R-Limbic	19.3, -7.7, -14.8	Right Hippocampus
4	36	R-Insula	37.5, 21.1, -10.1	Right Inferior Frontal Gyrus, Orbital part
3	255	L-Cerebellum	-6.6, -54.7, -26.3	N/A (Cerebellum)
3	254	L-Cerebellum	-21.2, -53.4, -23.6	Left Lobule VI of Cerebellum
3	253	L-Cerebellum	-26.3, -69.5, -30.6	Left Crus I of Cerebellum
3	252	L-Cerebellum	-46.4, -46.8, -42.9	Left Crus II of Cerebellum
3	251	L-Cerebellum	-10.3, -37.7, -25.1	Left lobule III of Cerebellum
3	246	L-Cerebellum	-42.6, -63.7, -46.3	Left Crus II of

				Cerebellum
3	242	L-Cerebellum	-30.2, -80.2, -40.3	Left Crus II of Cerebellum
3	202	L-Temporal	-30, -5.8, -40.9	Left Fusiform Gyrus
3	163	L-MotorStrip	-57, -3.4, 6.8	Left Superior Temporal Gyrus
3	135	L-Prefrontal	-18.2, 19, -21	Left Inferior Frontal Gyrus, Pars Orbitalis
3	132	R-Brainstem	6.3, -24.9, -17.5	N/A (Midbrain)