Luliconazole, a New Antifungal, vs Amphotericin B, Voriconazole, Posaconazole and Caspofungin against Clinical and Environmental *Aspergillus* Nigri Complex

Sahar Hivary[1], Mahnaz Fatahinia[1,2], Marzieh Halvaeezadeh[1], Ali Zarei Mahmoudabadi[1,2]

[1]Department of Medical Mycology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

[2]Infectious and Tropical Diseases Research Center, Health Research Institute, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Correspondence: Prof. Ali Zarei Mahmoudabadi, Infectious and Tropical Diseases Research Center, Health Research Institute, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran, Email: zarei40@hotmail.com; Tel: +986133330074; Fax: +986133332036

Running title: Luliconazole against black Aspergilli

ABSTRACT

Black Aspergilli are, the most causes of aspergillosis and *Aspergillus niger and A. tubingensis are two more frequently isolates. Although, amphotericin B was a gold standard for the treatment of invasive fungal infection for several decades, it replaced by several new antifungals. Furthermore, a novel antifungal, luliconazole, appears to offer the potential for improved therapy for aspergillosis. The aim of the present study was to compare the effect of a novel antifungal agent, luliconazole, with classical antifungal* against clinical and environmental strains of black Aspergilli. Sixty seven strains of black Aspergilli were identified using morphological and molecular tests (β-Tubulin gene). Antifungal susceptibility test was applied according to CLSI M38 A2. The results were reported as MIC range, MIC_{50}, MIC_{90} and MIC_{GM}. In the present study, *A. niger* was the common isolate followed by, *A. tubingensis* and 54.1% (clinical) and 30% (environmental) of isolates were resistant to caspofungin. The highest resistant rate was found in amphotericin B for both clinical (86.5%) and environmental (96.7%) strains. Clinical strains of *Aspergillus* were more sensitive to voriconazole (86.7%) than environmental strains (70.3%). On the other hand, 83.8% of clinical and 70% of environmental isolates were resistant to posaconazole, respectively. It is found that the lowest MIC range, MIC_{50}, MIC_{90}, and MIC_{GM} was attributed to luliconazole in clinical strains. In conclusion, luliconazole vs. routine antifungal is a potent antifungal for *A. niger* complex *in vitro*. The MIC range, MIC_{50}, MIC_{90} and MIC_{GM} of luliconazole against black Aspergilli were the lowest among the representative tested antifungals.

Keywords: Black Aspergilli, Luliconazole, Clinical and environmental isolates, Antifungal profile

Introduction 42

Aspergillus species are the commonest environmental saprophytic fungi that have worldwide 43

distribution. The Genus *Aspergillus* containing several sections, including, *Aspergillus* 44

section Nigri, section Flavi, section Terrei, section Fumigati, and section Clavati (1, 2). 45

Although, all sections are contributing in the involving aspergillosis, the most cases of disease 46

are usually caused by different species of *Aspergillus* in the sections of Nigri and Flavi. Black 47

Aspergilli (section Nigri) are often grown in different environmental conditions, whereas, they 48

are more prevalent at tropical and subtropical areas. 49

The section Nigri (*Aspergillus niger*, sensu lato) contains 19 species including, 50

Aspergillus niger, *A. tubingensis*, *A. awamory*, *A. welwitschiae*, *A. acidus*, *A. brasiliensis* etc. 51

(3-5). The Aspergilli in this section are comprised of several closely related species, and 52

accurate identification based on sequence analyses of β-tubulin gene (6). *Aspergillus niger* and 53

A. tubingensis are two more frequently isolates from clinical materials (4, 7-9). Black 54

Aspergilli cause several types of aspergillosis among predisposed patients. Out of them, 55

otomycosis is the most common cutaneous infection that are usually caused by black 56

Aspergilli. 57

The increasing of fungal opportunistic infections among receiving intensive therapy 58

immunocompromised, hematological malignancies and transplant patients during last decades 59

is remarkable (10-14). Invasive *Aspergillus* infections are one of the life threatening human 60

disease. On the other hand, some species of *Aspergillus* have inherent resistance to some 61

antifungal agents. Moreover, some species have raised minimum inhibitory concentration 62

(MIC) against specific antifungals. As a results, infection prevention consultant and the best 63

choice antifungal are common clinical challenges. 64

While, amphotericin B was a Gold standard in the first-line treatment of invasive fungal 65

infections for several decades, it has been replaced by several new antifungals including, 66

voriconazole, posaconazole and caspofungin (13). The randomized clinical trials were 67

considered voriconazole as the primary therapy for invasive pulmonary aspergillosis (15). 68

Further studies have shown that posaconazole is a useful antifungal for invasive fungal 69

infection including aspergillosis (13). On the other hand, during 2-3 last decades, caspofungin 70

was developed to improve the prognosis of invasive aspergillosis (16). 71

The availability of a novel antifungal, luliconazole, appears to offer the potential for 72

improved therapy for a wide range of invasive fungal infections, including aspergillosis. 73

Luliconazole was basically introduced as anti-dermatophytic infections in Japan and India (17, 74

18). However it is newly under evaluation by several researchers for *A. fumigatus* (6, 19), *A.* 75

terreus (6, 20), *A. flavus* (6, 21), *A. niger* (6) and *A. tubingensis* (6). 76

Objectives 77

The aim of the present study was to compare the effect of a novel antifungal agent, luliconazole, 78

with amphotericin B, voriconazole, posaconazole and caspofungin against clinical and 79

environmental strains of black Aspergilli. 80

Materials and Methods 81

Ethics statement 82

This project was approved by the ethical committee of Ahvaz Jundishapur University of 83

Medical Sciences (IR.AJUMS.REC.1396.1066). 84

Fungal isolates 85

Thirty seven clinical strains of black Aspergilli were previously isolated from clinical 86

materials, identified based on morphology characteristics and preserved at Medical Mycology 87

laboratory affiliated to Ahvaz Jundishapur University of Medical Sciences. Environmental 88

strains (30 strains) of black Aspergilli were trapped from airborne spores using 89

Sabouraud dextrose agar, SDA (BioLife, Italia) plates. Primary screening of black Aspergilli 90

strains was applied based on colony morphology (Black colony) and micromorphology. In the 91

present study all strains (clinical and environmental) were subcultured on SDA and re-identified using molecular tests.

DNA extraction

All strains (clinical and environment isolates) were cultured on SDA plates and incubated at 29ºC for 24-48 hours. Mycelia were collected in cryo-tubes containing 300 µL lysis buffer and 0.46 g glass beads and kept at 4ºC for 72 hours. The tube contents were homogenized using a SpeedMill PLUS Homogenizer (Analytikjena, Germany) for 6 minutes (3 cycles) and boiled at 100ºC for 20 minutes. 300 µL of sodium acetate (3 molar) was added to each tube and stored at -20ºC for 10 minutes. Supernatants were removed after a centrifugation at 12000 rpm for 10 min. DNA was purified using phenol-chloroform-isoamyl alcohol according to Makimura et al., protocol (22). Finally purified DNA was preserved at -20 ºC for further tests.

Molecular identification

In the present study, β-Tubulin gene was used for the molecular detection of strains using primers pair, βt2a (forward), 5' GGTAACCAAATCGGTGCTGCTTTC 3' and βt2b (reverse) 5' ACCCTCAGTGTAGTGACCCTTGGC 3' (23). PCR products subjected for sequence analysis and then sequences were manually verified by MEGA6 software package and aligned using the CLUSTALW algorithm. All sequences were compared to reference sequences in the GenBank (NCBI) and CBS database via the nucleotide BLAST ™ algorithm to obtain a definitive identification (similarity values ≥ 99%). Finally, all nucleotide sequences representative were deposited in the GenBank database.

Antifungal susceptibility assay

Twofold serial dilutions of antifungals including, luliconazole (APIChem Technology, China) (from 0.25 to 0.00012 µg/mL), amphotericin B (Sigma - Aldrich, Germany) (from 16 to 0.125 µg/mL), voriconazole (Sigma - Aldrich, Germany) (from 4 to 0.0078 µg/mL), posaconazole (Sigma - Aldrich, Germany) (from 4 to 0.0312 µg/mL), and caspofungin (Sigma - Aldrich,

Germany) (from 1 to 0.0078 µg/mL) were prepared in RPMI 1640 (Bio Idea, Iran). Antifungal 117

susceptibility test was applied according to CLSI M38 A2 (24). A standard suspension (0.5 118

McFarland) of 48 - 72 hours cultures on SDA was prepared in sterile saline (0.85%) with 0.2% 119

Tween 20 (Merck, Germany). Then, 100 µL of diluted suspension (1:50) and 100 µL of serial 120

dilutions of each antifungal purred in 96 well microplates. Microplates incubated at 35ºC for 121

24 - 72 h and results were recorded as MIC. Finally, MIC range, MIC_{50}, MIC_{90} and MIC_{GM} 122

were calculated. CLSI or EUCAST have not been defined any clinical or epidemiologic 123

breakpoints/cutoffs for amphotericin B, voriconazole, posaconazole, caspofungin and 124

Aspergillus species. In the present study, strains susceptibility / resistance to each antifungals 125

was evaluated according to proposed breakpoints by several researchers (Table 1) (25-30). 126

Table 1: Defined breakpoints of amphotericin B, voriconazole, posaconazole and caspofungin 127

for *Aspergillus niger* sensu lato 128

Antifungals	MIC (µg/mL)	
	Sensitive	**Resistance**
Amphotericin B	≤2	>2
Posaconazole	≤0.5	>0.5
Voriconazole	≤1	>1
Caspofungin	≤0.06	>0.06
Luliconazole	Undefined	Undefined

MIC, Minimum inhibitory concentration 129

Statistical analysis 131

The Chi-squared test using the Social Science Statistics software (Online) was applied to 132

determine the significant between variables and P value < 0.05 is considered as significance 133

level. 134

Results 135

Molecular detection of isolates 136

In the present study, 37 clinical strains of black Aspergilli were detected using molecular and 137

sequencing method. *Aspergillus niger* (21, 56.8%) was the common strain followed by, *A.* 138

tubingensis (11, 29.8%), *A. luchuensis* (1, 2.7%), and black *Aspergilli* (4, 10.8%) (Table 2). 139

Furthermore, out of 30 environmental black Aspergilli isolates, 15 (50%) was identified as *A.* 140

niger followed by, *A. tubingensis* (13, 43.3%), *A. piperis* (1, 3.3%) and black Aspergilli (1, 141

3.3%). However, we could not identified four clinical and one environmental black Aspergilli, 142

using molecular technique due to lack enough DNA. 143

Table 2: Clinical and environmental black Aspergilli with accession numbers 144

Sources	Morphological identification	Molecular identification	Accessions numbers
Clinical isolates (37 isolates)	*Aspergillus niger* sensu lato	*A. niger*, sensu stricto (21)	LC441155, LC441156, LC441157, LC441158, LC441159, LC441160, LC441161, LC441162, LC441163, LC441164, LC441165, LC441166, LC441167, LC456320, LC456323, LC456326, LC456335, LC456336, LC456337, LC456339, LC456341
		A. tubingensis (11)	LC441168 , LC441169, LC441170, LC441171, LC456297, LC456298, LC456301, LC456302, LC456303, LC456338, LC456340
		A. luchuensis (1)	LC456304
	Unidentified black *Aspergilli* (4)	***	***
Environmental isolates (30 isolates)	*Aspergillus niger* sensu lato	*A. niger*, sensu stricto (15)	LC456317, LC456318, LC456319, LC456321, LC456322, LC456324, LC456325, LC456327, LC456328, LC456329, LC456330, LC456331, LC456332, LC456333, LC456334
		A. tubingensis (13)	LC456299, LC456300, LC456306, LC456307, LC456308, LC456309, LC456310, LC456311, LC456312, LC456313, LC456314, LC456315, LC456316
		A. piperis (1)	LC456305
	Unidentified black *Aspergilli* (1)	***	***

Clinical isolates 146

The minimum effective concentration (MEC) range for all clinical *Aspergillus* species was 1 - 147

0.0078 μg/ml for caspofungin (Table 3). In addition, the 50% and 90% MEC (MEC_{50}, MEC_{90}) 148

values were 0.125 and 0.5 µg/ml for caspofungin. In the present study, 54.1% of isolates were 149

resistant to caspofungin. The results have shown that the MIC range of amphotericin B for 150

tested isolates was 16 - 0.25 µg/mL. However, MIC_{50}, MIC_{90} was similar, 8 µg/mL. The highest 151

resistant rate was found in amphotericin B (86.5%). All clinical isolates were inhibited at the 152

ranges 4 - 0.0078 and 4 - 0.0625 µg/mL of voriconazole and posaconazole, respectively. 153

However, the MIC_{GM} for voriconazole (0.77 µg/mL) was lower than posaconazole (1.45 154

µg/mL). In our study, 29.7% and 83.8% of isolates were resistant to voriconazole and 155

posaconazole, respectively. It is found that the lowest MIC range (0.125 - 0.00024 µg/mL), 156

MIC_{50} (0.00195 µg/mL), MIC_{90}, (0.125 µg/mL) and MIC_{GM} (0.00295 µg/mL) was attributed 157

to luliconazole. 158

Table 3: The antifungal susceptibility pattern of 37 clinical strains of black Aspergilli 159

Caspofungin	No	(μg/mL)				
		MEC range	MEC_{50}	MEC_{90}	MEC_{GM}	R (%)
A. niger	21	1-0.0078	0.125	0.5	0.099	11(52.4%)
A. tubingensis	11	0.5-0.032	0.125	0.5	0.133	7(63.6%)
A. luchuensis	1	0.032	-	-	-	-
Black *Aspergilli*	4	0.25-0.0625	-	-	-	2(50%)
Total	37	1-0.0078	0.125	0.5	0.107	20(54.1%)
Amphotericin B	No	MIC range	MIC_{50}	MIC_{90}	MIC_{GM}	R (%)
A. niger	21	8-0.25	8	8	4.56	17(81%)
A. tubingensis	11	16-4	8	8	8	11(100%)
A. luchuensis	1	1	-	-	-	-
Black *Aspergilli*	4	8-4	-	-	-	4(100%)
Total	37	16-0.25	8	8	5	32(86.5%)
Voriconazole	No	MIC range	MIC_{50}	MIC_{90}	MIC_{GM}	R (%)
A. niger	21	2-0.0625	1	2	0.99	5(23.8%)
A. tubingensis	11	4-0.5	1	2	1.20	4(36.4%)
A. luchuensis	1	0.0078	-	-	-	-
Black *Aspergilli*	4	2-0.5	-	-	-	2(50%)
Total	37	4-0.0078	1	2	0.77	11(29.7%)
Posaconazole	No	MIC range	MIC_{50}	MIC_{90}	MIC_{GM}	R (%)
A. niger	21	4-0.0625	2	2	1.26	17(81%)
A. tubingensis	11	4-0.125	2	4	2.13	10(90.9%)
A. luchuensis	1	0.5	-	-	-	-
Black *Aspergilli*	4	4-0.25	-	-	-	4(100%)
Total	37	4-0.0625	2	4	1.45	31(83.8%)
Luliconazole	No	MIC range	MIC_{50}	MIC_{90}	MIC_{GM}	R (%)

A. niger	21	0.125-0.00024	0.00195	0.125	0.00378	-
A. tubingensis	11	0.125-0.00024	0.00195	0.00391	0.00251	-
A. luchuensis	1	0.00098	-	-	-	-
Black *Aspergilli*	4	0.00391-0.00049	-	-	-	-
Total	37	0.125-0.00024	0.00195	0.125	0.00295	-

MEC, Minimum effective concentration; MIC, Minimum inhibitory concentration; R, Resistant

Environmental isolates

Table 4 summarizes the *in vitro* susceptibilities of 30 environmental *Aspergillus* Nigri against caspofungin. The MEC range, MEC_{50}, MEC_{90} and MEC_{GM} were 0.5 - 0.0078, 0.0625, 0.25, and 0.0507 µg/ml, respectively. Furthermore, 30% of environmental strains were resistant to caspofungin. As shown, the MIC range for amphotericin B was 16 - 2 µg/ml followed by, MIC_{50}, MIC_{90} and MIC_{GM} were 8, 8 and 6.063 µg/ml, respectively. Moreover, 96.7% of strains were resistant to antifungal. Totally, the MIC range voriconazole for environmental isolates of *Aspergillus* was 2 - 0.0625 µg/ml, whereas MIC_{90} 2 µg/ml, MIC_{50} 0.5 and MIC_{GM} 0.4665 µg/ml). Our results indicated that only 4(13.3%) of strains were resistant to voriconazole. In the present study, tested isolates were inhibited at MIC range 4 - 0.0625 µg/ml by posaconazole. Furthermore, the MIC_{50}, MIC_{90} and MIC_{GM} were 2, 4 and 1.2599 µg/ml, respectively. In addition, 70% of strains were resistant to posaconazole. The same as clinical isolates, the lowest MIC range was 0.00781 - 0.00049 µg/ml for luliconazole. Moreover, the MIC_{50}, MIC_{90} and MIC_{GM} were 0.00195, 0.00391 and 0.00195 µg/ml, respectively.

Table 4: The antifungal susceptibility pattern of 30 environmental strains of black Aspergilli

Caspofungin	No	(µg/mL)				
		MEC range	MEC_{50}	MEC_{90}	MEC_{GM}	R (%)
A. niger	15	0.25-0.0078	0.032	0.25	0.0412	3(20%)
A. tubingensis	13	0.5-0.0078	0.0625	0.5	0.0733	6(46.2%)
A. piperis	1	0.0625	-	-	-	-
Black *Aspergilli*	1	0.0078	-	-	-	-
Total	30	0.5-0.0078	0.0625	0.25	0.0507	9(30%)
Amphotericin B	**No**	**MIC range**	MIC_{50}	MIC_{90}	MIC_{GM}	R (%)
A. niger	15	16-2	8	16	6.964	14(93%)
A. tubingensis	13	8-4	4	8	5.508	13(100%)
A. piperis	1	4	-	-	-	-

	No	MIC range	MIC$_{50}$	MIC$_{90}$	MIC$_{GM}$	R (%)
Black *Aspergilli*	1	4	-	-	-	-
Total	30	16-2	8	8	6.063	29(96.7%)
Voriconazole	**No**	**MIC range**	**MIC$_{50}$**	**MIC$_{90}$**	**MIC$_{GM}$**	**R (%)**
A. niger	15	2-0.125	1	2	0.6300	2(13.3%)
A. tubingensis	13	2-0.0625	0.5	2	0.4261	2(15.4%)
A. piperis	1	0.125	-	-	-	-
Black *Aspergilli*	1	0.0625	-	-	-	-
Total	30	2-0.0625	0.5	2	0.4665	4(13.3%)
Posaconazole	**No**	**MIC range**	**MIC$_{50}$**	**MIC$_{90}$**	**MIC$_{GM}$**	**R (%)**
A. niger	15	4-0.5	2	4	1.8234	14(93%)
A. tubingensis	13	4-0.125	2	4	1.1125	7(53.8%)
A. piperis	1	0.5	-	-	-	-
Black *Aspergilli*	1	0.0625	-	-	-	-
Total	30	4-0.0625	2	4	1.2599	21(70%)
Luliconazole	**No**	**MIC range**	**MIC$_{50}$**	**MIC$_{90}$**	**MIC$_{GM}$**	**R (%)**
A. niger	15	0.00781-0.00098	0.00195	0.00391	0.00214	-
A. tubingensis	13	0.00781-0.00049	0.00195	0.00391	0.00195	-
A. piperis	1	0.00195	-	-	-	-
Black *Aspergilli*	1	0.00049	-	-	-	-
Total	30	0.00781-0.00049	0.00195	0.00391	0.00195	-

MEC, Minimum effective concentration; MIC, Minimum inhibitory concentration; R, Resistant

In the present study, caspofungin was significantly more effective against environmental than clinical strains (P = 0.048). However, the inhibitory effect of other antifungals (amphotericin B, posaconazole and voriconazole) against both strains (clinical and environmental) was similar (amphotericin B, P=0.147; voriconazole, P=0.109; posaconazole, P=0.178). When we compared the effect antifungals against *A. niger* and *A. tubingensis* among clinical and environmental strains, it is found that caspofungin was more effective on *A. niger* with environmental sources than clinical strains (P=0.0482). Whereas, the effect of other antifungals against both species was not significant.

Our results showed that 32 (86.5%) of clinical strains have multi drug resistance, 2(5.4%) resistance to one antifungal and 3(8.1%) were fully sensitive to antifungals (Table 5). Two strains of *A. tubingensis*, one *A. niger* and one black Aspergilli were resistant to all antifungals (except luliconazole). On the other hand, in environmental strains, 70% and 30% of strains were multi drug resistance (2-4 antifungals) and drug resistance (one antifungals),

respectively (Table 6). Two strains of *A. niger* and one *A. tubingensis* were resistant to all antifungals (except luliconazole).

Table 5: Drug resistance against tested antifungals among 37 clinical strains

Species (Clinical)	Antifungal drugs				
	LUL	POS	VOR	AMP	CAS
Aspergillus niger	0.125	R	S	R	R
A. niger	0.125	S	S	R	R
A. niger	0.125	S	S	R	R
A. niger	0.125	R	S	R	R
A. tubingensis	0.125	R	S	R	R
A. niger	0.01561	R	S	R	R
A. niger	0.00781	R	S	R	R
A. niger	0.00781	R	S	S	S
A. tubingensis	0.00391	R	R	R	R
Black Aspergilli	0.00391	R	S	R	S
A. tubingensis	0.00391	R	R	R	R
A. niger	0.00391	R	S	R	R
A. niger	0.00195	R	R	R	S
A. tubingensis	0.00195	R	R	R	S
A. tubingensis	0.00195	R	R	R	S
Black Aspergilli	0.00195	R	R	R	S
A. tubingensis	0.00195	R	S	R	R
A. niger	0.00195	R	S	R	R
A. tubingensis	0.00195	R	S	R	R
A. niger	0.00195	R	R	R	S
A. tubingensis	0.00195	R	S	R	S
A. niger	0.00195	R	R	R	S
Black Aspergilli	0.00195	R	R	R	R
A. niger	0.00195	R	S	R	S
A. tubingensis	0.00195	S	S	R	R
A. niger	0.00098	R	S	R	S
A. niger	0.00098	R	S	R	R
A. niger	0.00098	R	S	S	S
A. niger	0.00098	R	R	R	R
A. tubingensis	0.00098	R	S	R	S
A. niger	0.00098	R	R	R	S
A. niger	0.00098	R	S	R	R
A. luchuensis	0.00098	S	S	S	S
Black Aspergilli	0.00049	S	S	R	R

A. tubingensis	0.00024	R	S	R	R
A. niger	0.00024	S	S	S	S
A. niger	0.00024	S	S	S	S

LUL, Luliconazole; POS, Posaconazole; VOR, Voriconazole; AMP, Amphotericin B; CAS, 195
Caspofungin; R, Resistance: S, Sensitive 196
197

Table 6: Drug resistance against tested antifungals among 30 environmental strains 198

Species (Environmental)	Antifungal drugs				
	LUL	POS	VOR	AMP	CAS
Aspergillus niger	0.00781	R	S	R	S
A. tubingensis	0.00781	R	R	R	S
A. niger	0.00391	R	S	R	S
A. niger	0.00391	R	S	R	S
A. niger	0.00391	R	R	R	R
A. tubingensis	0.00391	R	R	R	R
A. niger	0.00391	R	S	R	S
A. niger	0.00195	R	S	R	S
A. tubingensis	0.00195	R	S	R	R
A. niger	0.00195	R	S	R	S
A. tubingensis	0.00195	S	S	R	S
A. niger	0.00195	R	S	R	S
A. niger	0.00195	R	S	S	S
A. tubingensis	0.00195	S	S	R	S
A. niger	0.00195	R	S	R	R
A. tubingensis	0.00195	S	S	R	S
A. tubingensis	0.00195	R	S	R	R
A. tubingensis	0.00195	R	S	R	R
A. tubingensis	0.00195	R	S	R	S
A. tubingensis	0.00195	R	S	R	R
A. niger	0.00195	R	S	R	S
A. tubingensis	0.00195	S	S	R	S
A. piperis	0.00195	S	S	R	S
A. niger	0.00098	S	S	R	S
A. niger	0.00098	R	S	R	S
A. tubingensis	0.00098	S	S	R	S
A. niger	0.00098	R	R	R	R
A. niger	0.00098	R	S	R	S
Black Aspergilli	0.00049	S	S	R	S
A. tubingensis	0.00049	S	S	R	R

199

LUL, Luliconazole; POS, Posaconazole; VOR, Voriconazole; AMP, Amphotericin B; CAS, Caspofungin; R, Resistance: S, Sensitive

Discussion

In the present study, the *Aspergillus* strains isolated from clinical and air borne samples were species typed using classical morphological features and molecular methods. Moreover, their susceptibilities to amphotericin B, caspofungin and three azole antifungals including luliconazole, voriconazole, and posaconazole were assayed. *Aspergillus tubingensis*, *A. luchuensis* and *A. piperis* were identified as the cryptic species of *A. niger* sensu lato by the sequence analysis of β-tubulin gene. Several reports have shown that *A. niger* is generally as common causative agent of otomycosis and one of the most important agent for invasive aspergillosis (8, 10, 31-33). However, new molecular techniques are indicating that this species comprises 19 cryptic species (4, 6, 9).

The most common cryptic species is *A. niger* (30.2 - 57.0%) sensu stricto, followed by *A. tubingensis* (31.6 - 53.5%) and other cryptic species of *A. niger* sensu lato (4.0 - 8.8%) (4, 6, 16). It is believed that the partial β-tubulin sequencing was superior to ITS sequencing for the species assignment of black Aspergilli (34, 35). Overall, our results were comparable to those of previous studies about the frequency of cryptic species. The most common clinical (86.5%) and environmental (93.3%) species in our data set are *A. niger* sensu stricto and *A. tubingensis*, respectively. Furthermore, with using β-tubulin sequencing, two rare species, *A. luchuensis* and *A. piperis* were respectively identified from clinical and environmental samples.

There are the limited data in *in vitro* efficacy of antifungals against the black Aspergilli both from clinical and environmental sources. While, the clinical and environmental strains had the same MIC ranges for caspofungin, the resistant to antifungal showed the clear differences between clinical and environmental strains (P = 0.048), where the clinical isolates

showed higher resistant rate than the environmental strains. In a report by Badali *et al*., only

6.1% of environmental strains of *A. niger* were resistant to caspofungin and all clinical isolates

ranged at 0.008–0.063 µg/ml (9).

The *in vitro* activities of posaconazole, voriconazole, and amphotericin B against

clinical *Aspergillus* strains have been reported by Arikan *et al*. (13). They reported that

voriconazole was the most active antifungal against *A. niger*. Comparable to our results,

voriconazole was more potent than the other tested antifungal (with exception luliconazole)

against both clinical and environmental strains. *Aspergillus tubingensis* resistant strains to

amphotericin B was very common both in environment and clinical settings, followed by

posaconazole, caspofungin, and voriconazole. However, the resistant rate was lower among

environmental than clinical strains. Hashimoto et al., finding suggests that *A. tubingensis* is

intrinsically resistant to azole antifungals (3). Antifungal susceptibility testing of our *A.*

tubingensis strains revealed 90.9% and 53.8% of clinical and environmental isolates were

resistant to posaconazole.

During the last decades, some studies have shown a high efficacy of luliconazole

against dermatophytes and onychomycosis agents both *in vivo* and *in vitro* (17, 18, 36-38).

Furthermore, recently a few studies examined the potency of luliconazole against different

species of *Candida*, *A. fumigatus*, *A. terreus* and *Fusarium* species (20, 21, 39). However, the

efficacy profile of luliconazole against *A. niger*, complex is unknown. Our results showed that,

although the MIC range for strains was externally low, this range for environmental strains

(0.00781-0.00049 µg/ml) was lower than clinical strains (0.125 - 0.00024 µg/ml). As shown in

table 5, only five clinical strains (*A. niger* sensu stricto, 4 isolates and *A. tubingensis*, 1 isolate)

have a MIC = 0.125 µg/ml. In the present study, 100% of environmental and 83.8% of clinical

strains had the lowest MICs (MICs < 0.00781 µg/ml) against luliconazole. Moreover, the

MIC_{GM} for environmental and clinical strains were 0.00195 and 0.00295 µg/ml, respectively.

Abastabar et al. (19) and Omran et al. (21) were respectively tested luliconazole against *A.* *fumigatus* and *A. flavus*, and found that the antifungal has the lowest MICs against *A. fumigatus* (0.016 µg/ml) and *A. flavus* (0.062 µg/ml).

In conclusion, luliconazole vs. amphotericin B, voriconazole, posaconazole and caspofungin is a potent antifungal for *A. niger* sensu lato *in vitro*. The MIC range, MIC_{50}, MIC_{90} and MIC_{GM} of luliconazole against black Aspergilli were the lowest among the representative tested antifungals. However, these results must be confirmed by the clinical trials for the efficacy of luliconazole as anti-aspergillosis infections.

Acknowledgments

We would like to thank the Infectious and Tropical Diseases Research Center, Health Research Institute, Ahvaz Jundishapur University of Medical Sciences for their support.

Authors' Contribution

Study concept and design, Ali Zarei Mahmoudabadi; isolation and preparing clinical and environmental isolates, Marzieh Halvaeezadeh and Sahar Hivary; conducting the experiments, Sahar Hivary; data analysis and interpretation of the results, Ali Zarei Mahmoudabadi, Mahnaz Fatahinia, and Sahar Hivary; drafting of the manuscript, Ali Zarei Mahmoudabadi; Critical editing Mahnaz Fatahinia.

Funding support

This study was a part of MSc thesis (Sahar Hivary) supported by a grant (No: OG-96148) from the Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

Conflict of interest

No conflict of interest declared.

References

1.   **Alastruey-Izquierdo A, Mellado E, Pelaez T, Peman J, Zapico S, Alvarez M, Rodriguez-Tudela JL, Cuenca-Estrella M, Group FS.** 2013. Population-based

survey of filamentous fungi and antifungal resistance in Spain (FILPOP Study). Antimicrob Agents Chemother **57:**3380-3387.

2. **Varga J, Due M, Frisvad JC, Samson RA.** 2007. Taxonomic revision of Aspergillus section Clavati based on molecular, morphological and physiological data. Stud Mycol **59:**89-106.

3. **Hashimoto A, Hagiwara D, Watanabe A, Yahiro M, Yikelamu A, Yaguchi T, Kamei K.** 2017. Drug Sensitivity and Resistance Mechanism in Aspergillus Section Nigri Strains from Japan. Antimicrob Agents Chemother **61**.

4. **Gautier M, Normand AC, L'Ollivier C, Cassagne C, Reynaud-Gaubert M, Dubus JC, Bregeon F, Hendrickx M, Gomez C, Ranque S, Piarroux R.** 2016. Aspergillus tubingensis: a major filamentous fungus found in the airways of patients with lung disease. Med Mycol **54:**459-470.

5. **D'Hooge E, Becker P, Stubbe D, Normand AC, Piarroux R, Hendrickx M.** 2018. Black aspergilli: A remaining challenge in fungal taxonomy? Med Mycol doi:10.1093/mmy/myy124.

6. **Hagiwara S, Tamura T, Satoh K, Kamewada H, Nakano M, Shinden S, Yamaguchi H, Makimura K.** 2018. The Molecular Identification and Antifungal Susceptibilities of Aspergillus Species Causing Otomycosis in Tochigi, Japan. Mycopathologia doi:10.1007/s11046-018-0299-9.

7. **Castro C, Galan-Sanchez F, Linares MJ, Tejero R, Ruiz M, Serrano ML, Rodriguez-Iglesias M, Martin-Mazuelos E, Group F-IS.** 2018. A prospective survey of Aspergillus spp. in respiratory tract samples: Species identification and susceptibility patterns. Med Mycol doi:10.1093/mmy/myy080.

8. **Ali K, Hamed MA, Hassan H, Esmail A, Sheneef A.** 2018. Identification of Fungal Pathogens in Otomycosis and Their Drug Sensitivity: Our Experience. Int Arch Otorhinolaryngol **22:**400-403.

9. **Badali H, Fakhim H, Zarei F, Nabili M, Vaezi A, Poorzad N, Dolatabadi S, Mirhendi H.** 2016. In Vitro Activities of Five Antifungal Drugs Against Opportunistic Agents of Aspergillus Nigri Complex. Mycopathologia **181:**235-240.

10. **Atchade E, Jean-Baptiste S, Houze S, Chabut C, Massias L, Castier Y, Brugiere O, Mal H, Montravers P.** 2017. Fatal invasive aspergillosis caused by Aspergillus niger after bilateral lung transplantation. Med Mycol Case Rep **17:**4-7.

11. **Peghin M, Monforte V, Martin-Gomez MT, Ruiz-Camps I, Berastegui C, Saez B, Riera J, Ussetti P, Sole J, Gavalda J, Roman A.** 2016. 10 years of prophylaxis with

nebulized liposomal amphotericin B and the changing epidemiology of Aspergillus spp. 310
infection in lung transplantation. Transpl Int **29:**51-62. 311

12. **Koehler P, Tacke D, Cornely OA.** 2014. Aspergillosis of bones and joints - a review 312
from 2002 until today. Mycoses **57:**323-335. 313

13. **Arikan S, Sancak B, Alp S, Hascelik G, McNicholas P.** 2008. Comparative in vitro 314
activities of posaconazole, voriconazole, itraconazole, and amphotericin B against 315
Aspergillus and Rhizopus, and synergy testing for Rhizopus. Med Mycol **46:**567-573. 316

14. **Martin-Mazuelos E, Peman J, Valverde A, Chaves M, Serrano MC, Canton E.** 317
2003. Comparison of the Sensititre YeastOne colorimetric antifungal panel and Etest 318
with the NCCLS M38-A method to determine the activity of amphotericin B and 319
itraconazole against clinical isolates of Aspergillus spp. J Antimicrob Chemother 320
52:365-370. 321

15. **Zhang M, Su X, Sun WK, Chen F, Xu XY, Shi Y.** 2014. Efficacy of the combination 322
of voriconazole and caspofungin in experimental pulmonary aspergillosis by different 323
Aspergillus species. Mycopathologia **177:**11-18. 324

16. **Gheith S, Saghrouni F, Bannour W, Ben Youssef Y, Khelif A, Normand AC,** 325
Piarroux R, Ben Said M, Njah M, Ranque S. 2014. In vitro susceptibility to 326
amphotericin B, itraconazole, voriconazole, posaconazole and caspofungin of 327
Aspergillus spp. isolated from patients with haematological malignancies in Tunisia. 328
Springerplus **3:**19. 329

17. **Koga H, Nanjoh Y, Makimura K, Tsuboi R.** 2009. In vitro antifungal activities of 330
luliconazole, a new topical imidazole. Med Mycol **47:**640-647. 331

18. **Koga H, Nanjoh Y, Kaneda H, Yamaguchi H, Tsuboi R.** 2012. Short-term therapy 332
with luliconazole, a novel topical antifungal imidazole, in guinea pig models of tinea 333
corporis and tinea pedis. Antimicrob Agents Chemother **56:**3138-3143. 334

19. **Abastabar M, Rahimi N, Meis JF, Aslani N, Khodavaisy S, Nabili M, Rezaei-** 335
Matehkolaei A, Makimura K, Badali H. 2016. Potent Activities of Novel Imidazoles 336
Lanoconazole and Luliconazole against a Collection of Azole-Resistant and - 337
Susceptible Aspergillus fumigatus Strains. Antimicrob Agents Chemother **60:**6916- 338
6919. 339

20. **Zargaran M, Taghipour S, Kiasat N, Aboualigalehdari E, Rezaei-Matehkolaei A,** 340
Zarei Mahmoudabadi A, Shamsizadeh F. 2017. Luliconazole, an alternative 341
antifungal agent against Aspergillus terreus. J Mycol Med **27:**351-356. 342

21. **Omran SM, Taghizadeh-Armaki M, Zarrinfar H, Hedayati MT, Abastabar M, Moqarabzadeh V, Ansari S, Saber S, Hoseinnejad A, Miri A, Verweij PE, Seyedmousavi S.** 2019. In-vitro antifungal susceptibility testing of lanoconazole and luliconazole against Aspergillus flavus as an important agent of invasive aspergillosis. J Infect Chemother **25:**157-160.

22. **Makimura K, Tamura Y, Mochizuki T, Hasegawa A, Tajiri Y, Hanazawa R, Uchida K, Saito H, Yamaguchi H.** 1999. Phylogenetic classification and species identification of dermatophyte strains based on DNA sequences of nuclear ribosomal internal transcribed spacer 1 regions. J Clin Microbiol **37:**920-924.

23. **Glass NL, Donaldson GC.** 1995. Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. Appl Environ Microbiol **61:**1323-1330.

24. **Anonymous.** 2008. Reference method for broth dilution antifungal susceptibility testing of filamentous fungi, approved standard. Second edition. M38-A2. . **28**.

25. **Lass-Florl C.** 2014. Susceptibility testing in Aspergillus species complex. Clin Microbiol Infect **20 Suppl 6:**49-53.

26. **Espinel-Ingroff A, Cuenca-Estrella M, Fothergill A, Fuller J, Ghannoum M, Johnson E, Pelaez T, Pfaller MA, Turnidge J.** 2011. Wild-type MIC distributions and epidemiological cutoff values for amphotericin B and Aspergillus spp. for the CLSI broth microdilution method (M38-A2 document). Antimicrob Agents Chemother **55:**5150-5154.

27. **Espinel-Ingroff A, Diekema DJ, Fothergill A, Johnson E, Pelaez T, Pfaller MA, Rinaldi MG, Canton E, Turnidge J.** 2010. Wild-type MIC distributions and epidemiological cutoff values for the triazoles and six Aspergillus spp. for the CLSI broth microdilution method (M38-A2 document). J Clin Microbiol **48:**3251-3257.

28. **Espinel-Ingroff A, Fothergill A, Fuller J, Johnson E, Pelaez T, Turnidge J.** 2011. Wild-type MIC distributions and epidemiological cutoff values for caspofungin and Aspergillus spp. for the CLSI broth microdilution method (M38-A2 document). Antimicrob Agents Chemother **55:**2855-2859.

29. **Pfaller MA, Boyken L, Hollis RJ, Kroeger J, Messer SA, Tendolkar S, Diekema DJ.** 2009. In vitro susceptibility of clinical isolates of Aspergillus spp. to anidulafungin, caspofungin, and micafungin: a head-to-head comparison using the CLSI M38-A2 broth microdilution method. J Clin Microbiol **47:**3323-3325.

30. **Blum G, Perkhofer S, Haas H, Schrettl M, Wurzner R, Dierich MP, Lass-Florl C.** 2008. Potential basis for amphotericin B resistance in Aspergillus terreus. Antimicrob Agents Chemother **52:**1553-1555.

31. **Kaya AD, Kiraz N.** 2007. In vitro susceptibilities of Aspergillus spp. causing otomycosis to amphotericin B, voriconazole and itraconazole. Mycoses **50:**447-450.

32. **Baddley JW, Marr KA, Andes DR, Walsh TJ, Kauffman CA, Kontoyiannis DP, Ito JI, Balajee SA, Pappas PG, Moser SA.** 2009. Patterns of susceptibility of Aspergillus isolates recovered from patients enrolled in the Transplant-Associated Infection Surveillance Network. J Clin Microbiol **47:**3271-3275.

33. **Vermeulen E, Maertens J, Meersseman P, Saegeman V, Dupont L, Lagrou K.** 2014. Invasive Aspergillus niger complex infections in a Belgian tertiary care hospital. Clin Microbiol Infect **20:**O333-335.

34. **Hendrickx M, Beguin H, Detandt M.** 2012. Genetic re-identification and antifungal susceptibility testing of Aspergillus section Nigri strains of the BCCM/IHEM collection. Mycoses **55:**148-155.

35. **Heo MS, Shin JH, Choi MJ, Park YJ, Lee HS, Koo SH, Lee WG, Kim SH, Shin MG, Suh SP, Ryang DW.** 2015. Molecular identification and amphotericin B susceptibility testing of clinical isolates of Aspergillus from 11 hospitals in Korea. Ann Lab Med **35:**602-610.

36. **Khanna D, Bharti S.** 2014. Luliconazole for the treatment of fungal infections: an evidence-based review. Core Evid **9:**113-124.

37. **Scher RK, Nakamura N, Tavakkol A.** 2014. Luliconazole: a review of a new antifungal agent for the topical treatment of onychomycosis. Mycoses **57:**389-393.

38. **Baghi N, Shokohi T, Badali H, Makimura K, Rezaei-Matehkolaei A, Abdollahi M, Didehdar M, Haghani I, Abastabar M.** 2016. In vitro activity of new azoles luliconazole and lanoconazole compared with ten other antifungal drugs against clinical dermatophyte isolates. Med Mycol **54:**757-763.

39. **Taghipour S, Kiasat N, Shafiei S, Halvaeezadeh M, Rezaei-Matehkolaei A, Zarei Mahmoudabadi A.** 2018. Luliconazole, a new antifungal against Candida species isolated from different sources. J Mycol Med **28:**374-378.