Optical measurement of voltage sensing by endogenous ion channels

Authors

Parashar Thapa[1‡], Robert Stewart[1‡], Rebecka J. Sepela[1], Oscar Vivas[2], Laxmi K. Parajuli[2], Mark Lillya[1], Sebastian Fletcher-Taylor[1,3], Bruce E. Cohen[3], Karen Zito[2], Jon T. Sack[1*]

[1]Department of Physiology and Membrane Biology, University of California, Davis, CA 95616

[2]Center for Neuroscience, University of California, Davis, CA 95616

[3]Molecular Foundry and Division of Molecular Biophysics & Integrated Bioimaging, Lawrence Berkeley National Laboratory, Berkeley, CA 94720

[‡]Authors contributed equally to this work

*Correspondence to jsack@ucdavis.edu

ABSTRACT

A primary goal of molecular physiology is to understand how conformational changes of

proteins affect the function of cells, tissues, and organisms. Here, we describe an imaging

method for measuring the conformational changes of the voltage sensors of endogenous ion

channel proteins within live tissue, without genetic modification. We synthesized GxTX-594, a

variant of the peptidyl tarantula toxin guangxitoxin-1E, conjugated to a fluorophore optimal

for two-photon excitation imaging through light-scattering tissue. GxTX-594 targets the voltage

sensors of Kv2 proteins, which form potassium channels and plasma membrane–endoplasmic

reticulum junctions. GxTX-594 dynamically labels Kv2 proteins on cell surfaces in response to

voltage stimulation. To interpret dynamic changes in labeling intensity, we developed a

statistical thermodynamic model that relates the conformational changes of Kv2 voltage sensors

to labeling intensity. We used two-photon excitation imaging of rat brain slices to image Kv2

proteins in neurons. This imaging method enabled identification of conformational changes of

endogenous Kv2 voltage sensors in tissue.

INTRODUCTION

To move the field of voltage-sensitive physiology forward, we need new tools that indicate when and where voltage-sensitive conformational changes in endogenous proteins occur. Many classes of transmembrane proteins have been found to be voltage sensitive (Bezanilla 2008). One important class of voltage-sensitive proteins is the voltage gated ion channels. Electrophysiological techniques have enabled remarkably precise studies of the voltage-sensitivity of ionic conductances, primarily under reductionist experimental conditions where the channels have been removed from their native tissue. In addition to their canonical function as ion-conducting channels, voltage gated ion channel proteins have nonconducting functions which are independent of their ion conducting functions (Tanabe, Beam et al. 1988, Kaczmarek 2006). These nonconducting protein functions are largely inaccessible to study by electrophysiology and are more poorly understood. Novel approaches are needed to learn more about voltage-sensing in intact tissues and to unlock the mysterious realm of nonconducting voltage-sensitive physiology. Here, we present a new approach to image the conformational changes of endogenous voltage-sensitive proteins.

We have developed a protein imaging tool, an <u>E</u>ndogenous <u>V</u>oltage-sensor <u>A</u>ctivity <u>P</u>robe (EVAP), which dynamically labels living tissue to reveal the conformational status of endogenous Kv2 proteins with subcellular resolution. Kv2 proteins form voltage-gated K^+ channels (Frech, VanDongen et al. 1989), bind endoplasmic reticulum proteins to form plasma membrane–endoplasmic reticulum junctions (Johnson, Leek et al. 2018, Kirmiz, Palacio et al. 2018), regulate a wide variety of physiological responses in tissues throughout the body (Bocksteins 2016), and integrate their response to voltage with many other cellular processes including phosphorylation (Murakoshi, Shi et al. 1997), sumoylation (Plant, Dowdell et al. 2011), oxidation (MacDonald, Salapatek et al. 2003), membrane lipid composition (Ramu, Xu et al. 2006), and auxiliary subunits (Gordon, Roepke et al. 2006, Peltola, Kuja-Panula et al. 2011).

Kv2 proteins are members of the voltage-gated cation channel superfamily. The voltage sensors of proteins in this superfamily are comprised of a bundle of four transmembrane helices termed S1-S4 (Long, Campbell et al. 2005, Long, Tao et al. 2007). The S4 helix contains positively charged arginine and lysine residues, gating charges, which respond to voltage changes by

moving through the transmembrane electric field (Aggarwal and MacKinnon 1996, Seoh, Sigg et al. 1996, Islas and Sigworth 1999). When voltage sensor domains encounter a transmembrane voltage that is more negative on the inside of the cell membrane, voltage sensors are biased towards resting conformations, or down states, in which gating charges are localized intracellularly. When voltage becomes more positive, gating charges translate towards the extracellular side of the membrane, and voltage sensors are progressively biased towards up states, in a process of voltage activation (Armstrong and Bezanilla 1973, Zagotta, Hoshi et al. 1994, Tao, Lee et al. 2010, Xu, Li et al. 2019). Channel pore opening is distinct from, but coupled to, voltage sensor movement. In some voltage-gated ion channel proteins, voltage sensor movement is coupled to nonconducting protein functions (Tanabe, Beam et al. 1988, Kaczmarek 2006). To study the functional outputs of voltage sensors it is essential to measure voltage sensor activation itself. Conformational changes of voltage sensors have been detected with electrophysiological measurements of gating currents (Armstrong and Bezanilla 1973, Schneider and Chandler 1973, Bezanilla 2018) or by optical measurements from fluorophores covalently attached to voltage sensors by genetic encoding (Lin and Schnitzer 2016) or by chemical engineering (Zhang, Zheng et al. 2015). However, experimental limitations prevent these existing techniques from measuring conformational changes of voltage sensors of most endogenous proteins: gating currents can only be measured when the proteins are expressed at high density in a voltage-clamped membrane; engineered proteins differ from endogenous channels; and introducing point mutations and conjugated fluorophores into voltage sensors alters structure and function. Here, we develop a different strategy, EVAPs that dynamically label endogenous voltage sensors of Kv2 proteins to reveal conformational states.

To image where the voltage sensors of Kv2 proteins adopt a specific resting conformation in tissue, we exploit the conformation-selective binding of the tarantula peptide guangxitoxin-1E (GxTX), which can be conjugated to fluorophores to report Kv2 conformational changes (Tilley, Eum et al. 2014). Here, we synthesize GxTX-594, a Ser13Cys GxTX variant conjugated to Alexa Fluor 594, a fluorophore compatible with two-photon excitation imaging through light-scattering tissue. This EVAP equilibrates with Kv2 proteins on the time scale of seconds, revealing the probability (averaged over time) that unbound voltage sensors are resting or active. We develop a method to calculate the average conformational status of unlabeled Kv2 proteins

from images of GxTX-594 fluorescence. We deploy the GxTX-594 probe in brain slices and image voltage-sensitive fluorescence changes that reveal conformational changes of endogenous neuronal Kv2 proteins. This EVAP approach provides an imaging technique to study conformational changes of voltage-sensitive proteins in samples that have not (or cannot) be genetically modified.

MATERIALS AND METHODS
GxTX-594 synthesis

The Ser13Cys GxTX peptide was synthesized as described (Tilley, Eum et al. 2014). Methionine 35 of GxTX was replaced by the oxidation-resistant noncanonical amino acid norleucine and serine 13 was replaced with cysteine to create a spinster thiol. Ser13Cys GxTX was labeled with a Texas red derivative (Alexa Fluor 594 C_5 maleimide, Thermo Fisher Scientific, Catalog # 10256) to form GxTX-594. Ser13Cys GxTX lyophilisate was brought to 560 μM in 50% ACN + 1 mM Na_2EDTA. 2.4 μL of 1M Tris (pH 6.8 with HCl), 4 μL of 10 mM of Alexa Fluor 594 C_5 maleimide in DMSO and 17.9 μL of 560 μM Ser13Cys GxTX were added for a final solution of 100 mM Tris, 1.6 mM Alexa Fluor 594 C_5 maleimide and 0.4 mM GxTX in 24 μL of reaction solution. Reactants were combined in a 1.5 mL low protein binding polypropylene tube (LoBind, Eppendorf, Catalog # 022431081) and mixed at 1000 RPM, 20°C for 4 hours (Thermomixer 5355 R, Eppendorf). After incubation, the tube was centrifuged at 845 RCF for 10 minutes at room temperature. A purple pellet was observed post-centrifugation. The supernatant was transferred to a fresh tube and centrifuged at 845 RCF for 10 minutes. After this 2nd centrifugation, no visible pellet was seen. The supernatant was injected onto a reverse-phase HPLC C_{18} column (Biobasic- 4.6mm RP-C_{18} 5 μm, Thermo Fisher Scientific, Catalog # 2105-154630) equilibrated in 20% ACN, 0.1% TFA at 1 mL/min and eluted with a protocol holding in 20% ACN for 2 minutes, increasing to 30% ACN over 1 minute, then increasing ACN at 0.31% per minute. HPLC effluent was monitored by fluorescence and an absorbance array detector. 1 mL fractions were pooled based on fluorescence (280 nm excitation, 350 nm emission) and absorbance (214 nm, 280 nm, and 594 nm). GxTX-594 peptide-fluorophore conjugate eluted at approximately 35% ACN, and mass was confirmed by mass spectrometry using a Bruker Ultra Flextreme MALDI-TOF/TOF (Fig 1). Samples for identification from HPLC eluant were mixed 1:1 in an aqueous solution of 25% MeOH and 0.05% TFA saturated with alpha-cyano-4-

hydrocinnamic acid, pipetted onto a ground-steel plate, dried under vacuum, and ionized with

60-80% laser power. Molecular species were detected using a reflector mode protocol and

quantitated using Bruker Daltonics flexAnalysis 3.4. Lyophilizate containing GxTX-594

conjugation product was dissolved in CE buffer (defined below) and stored at -80 °C. GxTX-594

concentration was determined by 280 nm absorbance using a calculated extinction coefficient of

18,900 A.U. M^{-1} cm^{-1}.

CHO cell methods

CHO cell culture and transfection

The CHO-K1 cell line (ATCC) and a subclone transfected with a tetracycline-inducible rat

Kv2.1 construct (Kv2.1-CHO) (Trapani and Korn 2003), were cultured as described previously

(Tilley, Eum et al. 2014). To induce Kv2.1 expression in Kv2.1-TREx-CHO cells, 1 μg/mL

minocycline (Enzo Life Sciences, Catalog # ALX-380-109-M050), prepared in 70% ethanol at 2

mg/mL, was added to the maintenance media to induce K^+ expression. Minocycline was added

40-48 hours before imaging and voltage clamp fluorometry experiments. Minocycline was added

1-2 hours before whole cell ionic current recordings, to limit K^+ conductance such that voltage

clamp could be maintained. Transfections were achieved with Lipofectamine 2000 (Life

Technologies, Catalog # 1668027). 1.1 μL of Lipofectamine was diluted, mixed, and incubated

in 110 μL of Opti-MEM (Gibco, Product Number 31985062, Lot Number 1917064) in a 1:100

ratio for 5 minutes at room temperature. Concurrently, 1 μg of plasmid DNA and 110 μL of

Opti-MEM were mixed in the same fashion. DNA and Lipofectamine 2000 mixtures were

combined, titurated, and let sit for 20 minutes. Then, the transfection cocktail mixture was added

to a 35 mm cell culture dish of CHO cells at approximately 40% confluency and allowed to settle

at 37°C 5% CO_2 for 4-6 hours before the media was replaced. Rat Kv2.1-GFP (Antonucci, Lim

et al. 2001), rat Kv2.2-GFP (Kirmiz, Vierra *et al.* 2018), rat Kv1.5-GFP (Li, Guo et al. 2001), rat

Kv4.2-GFP (Shibata, Misonou et al. 2003), mouse BK-GFP (Trimmer Lab, UC Davis), rat Kv2.1

S586A-GFP (Kirmiz, Palacio et al. 2018), rat AMIGO-YFP (Bishop, Guan et al. 2015), rat

KvBeta2 (Shibata, Misonou et al. 2003) and rat KCHiP2 (An, Bowlby et al. 2000) plasmids were

all kind gifts from James Trimmer, University of California, Davis. Identities of constructs were

confirmed by sequencing from their CMV promoter.

Cell transfection for electrophysiology

Prior to transfection, cells were plated at 40% confluency in culture media free of selection

agents. Transfections were achieved with Lipofectamine 2000 (Life Technologies, Catalog #

1668027) following the manufacturer's protocol. Transfection mixes were prepared in Opti-

MEM (Life Technologies, Product Number 31985062) and included 1 µg of peGFP. Cells were

incubated in the transfection cocktail for 6-8 hours before being returned to regular growth

media. Cells were given 40-48 hours recovery following transfection before being used for

experiments.

Confocal and Airy disk imaging

Confocal images were obtained with an inverted confocal system (Zeiss LSM 880 410900-247-

075) run by ZEN black v2.1. A 63x/1.40 NA Oil DIC objective (Zeiss 420782-9900-799) was

used for most imaging experiments; a 63x/1.2 NA Water DIC objective (Zeiss 441777-9970-

000) was used for voltage clamp fluorometry experiments. GFP and YFP were excited with the

488 nm line from an argon laser (3.2 mW at installation) powered at 0.5% unless otherwise

noted. GxTX-594 was excited with 594 nm helium-neon laser (0.6 mW at installation) powered

at 10% unless otherwise noted. WGA-405 was excited with a 405 nm diode laser, (3.5 mW at

installation) powered at 1% unless otherwise noted. Temperature inside the microscope housing

was 27-30 °C.

In confocal imaging mode, fluorescence was collected with the microscope's 32-detector gallium

arsenide phosphide detector array arranged with a diffraction grating to measure 400-700 nm

emissions in 9.6 nm bins. Emission bands were 495-550 nm for GFP and YFP, 605-700 nm for

GxTX-594, and 420-480 nm for WGA-405. Point spread functions were calculated using ZEN

black software using emissions from 0.1 µm fluorescent microspheres, prepared on a slide

according to manufacturer's instructions (Thermo Fisher Scientific, Catalog # T7279). The point

spread functions for confocal images with the 63x/1.40 Oil DIC objective in the X-Y direction

were 228 nm (488 nm excitation) and 316 nm (594 nm excitation).

In Airy disk imaging mode, fluorescence was collected with the microscope's 32-detector

gallium arsenide phosphide detector array arranged in a concentric hexagonal pattern (Zeiss

Airyscan 410900-2058-580). After deconvolution, the point spread functions for the 63x/1.40 NA objective with 488 nm excitation was 124 nm in X-Y and 216 nm in Z; with 594 nm excitation 168 nm in X-Y and 212 nm in Z. For the 63x/1.2 NA objective, the point spread function with 488 nm excitation was 187 nm in X-Y and 214nm in Z; the point spread function with 594 nm excitation was 210 nm in X-Y and 213 nm in Z.

Unless stated otherwise, cells were plated in uncoated 35 mm dishes with a 7 mm inset No. 1.5 coverslip (MatTek, Catalog # P35G-1.5-20-C). The CHO cell external (CE) solution used for imaging and electrophysiology contained (in mM): 3.5 KCl, 155 NaCl, 10 HEPES, 1.5 CaCl$_2$, 1 MgCl$_2$, adjusted to pH 7.4 with NaOH. Measured osmolarity was 315 mOsm. When noted, solution was supplemented with either 10 mM glucose (CEG) or 10 mM glucose and 1% BSA (CEGB).

For time-lapse, GxTX-594 dose-response experiments, Kv2.1-CHO cells were plated onto 22 x 22 mm No. 1.5H coverglass (Deckglaser). Prior to imaging, cell maintenance media was removed and replaced with CEGB, then coverslip was mounted on an imaging chamber (Warner Instrument, Catalog # RC-24E) with vacuum grease. During time-lapse imaging, indicated concentrations of GxTX-594 were applied for 10 minutes followed by manual wash-out with CEGB solution. 15 minutes after wash-out, the next GxTX-594 concentration was added. Solutions were added to the imaging chamber perfusion via a syringe at a flow rate of ~1 mL / 10 sec. Images were taken every 5 sec. Laser power was set to 0.5% for the 488 nm laser and 1.5% for the 594 nm laser. For colocalization experiments with GFP-tagged proteins, cells were incubated in 100 nM GxTX-594 for 5 minutes and then washed with 1 mL of CEGB three times before imaging.

Whole cell voltage clamp for CHO cell imaging
Kv2.1-TREx-CHO cells plated in glass-bottom 35 mm dishes were washed with CEGB, placed in the microscope, and then incubated in 100 μL of 100 nM GxTX594 for 5 minutes to label cells. Before patch clamp, the solution was diluted with 1 mL of CEG for a working concentration of 9 nM GxTX594 during experiments. Cells with obvious GxTX-594 surface staining were voltage-clamped in whole-cell mode with an EPC-10 patch clamp amplifier

(HEKA) run by Patchmaster software, v2x90.2 (HEKA). The patch pipette contained a

potassium-deficient Cs^+ internal pipette solution to limit outward current and reduce voltage

error: 70 mM CsCl, 50 mM CsF, 35mM NaCl, 1 mM EGTA, 10 mM HEPES, brought to pH 7.4

with CsOH. Osmolarity was 310 mOsm. The liquid junction potential was calculated to be 3.5

mV and was not corrected. Borosilicate glass pipettes (Sutter Instruments, Catalog # BF 150-

110-10HP) were pulled with blunt tips to resistances less than 3.0 MΩ in these solutions. Cells

were held at -80 mV (unless noted otherwise) and stepped to indicated voltages. The voltage step

stimulus was maintained until any observed change in fluorescence was complete. For stimulus-

frequency dependence experiments, cells were given 2 ms steps to 40 mV at stated frequencies

(0.02, 5, 10, 20, 50, 100, 150, or 200 Hz). Images for voltage clamp fluorometry were taken in

Airy disk imaging mode with the settings described above.

K^+ channel-GFP ionic current recordings

Whole-cell voltage clamp was used to measure currents from CHO cells expressing Kv2.1-GFP,

Kv2.2-GFP, Kv1.5-GFP, Kv4.2-GFP, BK-GFP or GFP that was transfected as described above.

Cells were plated on glass bottom dishes. Cells were held at -80 mV, then 100 ms voltage steps

were delivered ranging from -80 mV to +80 mV in +5 mV increments. Pulses were repeated

every 2 seconds. The external (bath) solution contained CE solution. The internal (pipette)

solution contained (in mM): 35 KOH, 70 KCl, 50 KF, 50 HEPES, 5 EGTA adjusted to pH 7.2

with KOH. Liquid junction potential was calculated to be 7.8 mV and was not corrected.

Borosilicate glass pipettes (Sutter Instruments, Cat # BF150-110-10HP) were pulled into pipettes

with resistance less than 3 MΩ for patch clamp recording. Recordings were at room temperature

(22–24 °C). Voltage clamp was achieved with an Axon Axopatch 200B amplifier (MDS

Analytical Technologies) run by Patchmaster software, v2x90.2 (HEKA). Holding potential was

-80 mV. Capacitance and Ohmic leak were subtracted using a P/5 protocol. Recordings were low

pass filtered at 10 kHz and digitized at 100 kHz. Voltage clamp data were analyzed and plotted

with Igor Pro 7 or 8 (Wavemetrics). Current amplitudes at each voltage were the average from

0.19-0.20 s after voltage step. As the experiments plotted in Figure 6 Supplement 1 A were

merely to confirm functional expression of ion channels at the cell surface, series resistance

compensation was not used, and substantial cell voltage errors are predicted during these

experiments.

Kv2.1 ionic current recordings

Prior to patching, Kv2.1-CHO cells were washed in divalent-free PBS, then harvested in Versene (Gibco, Cat # 15040066). Cells were scraped and transferred to a polypropylene tube, pelleted and washed three times at 1,000 g for 2 minutes and then resuspended in the same external solution as used in the recording chamber bath. Cells were rotated in a polypropylene tube at room temperature (22-24°C) until use. Cells were then pipetted into a 50 uL recording chamber (Warner Instruments, RC-24N) prefilled with external solution and allowed to settle for 5 or more minutes. After adhering to the bottom of the glass recording chamber, cells were thoroughly rinsed with external solution using a gravity-driven perfusion system. Cells showing uniform intracellular GFP expression of intermediate intensity were selected for patching.

Voltage clamp was achieved with a patch clamp amplifier (Axon Instruments Axopatch 200B) run by Patchmaster software (HEKA). Borosilicate glass pipettes (Sutter, BF150-110-7.5HP) were pulled with blunt tips, coated with silicone elastomer (Dow Corning Sylgard 184), heat cured, and tip fire-polished to resistances less than 4MΩ. Capacitance and Ohmic leak were subtracted using a P/5 protocol. Recordings were low-pass filtered at 10kHz using the amplifier's built-in Bessel and digitized at 100kHz.

For whole cell ionic currents measurements in Kv2.1-CHO cells, the external patching solution contained (in mM): 3.5 KCl, 155 NaCl, 10 HEPES, 1.5 CaCl2, 1 MgCl2, adjusted to pH 7.4 with NaOH. The internal (pipette) solution contained (in mM): 70 KCl, 5 EGTA, 50 HEPES, 50 KF, and 35 KOH, adjusted to pH 7.4 with KOH. The osmolarity for the external solution was 315 mOsm, and 310 mOsm for the internal solution measured by a vapor presume osmometer. Following establishment of the whole-cell seal, ionic K+ current recordings were taken in the presence of a vehicle which consisted of 100 nM TTX, 10 mM glucose, and 0.1% BSA prepared in external solution. Cells were held at -100 mV with channel activation steps ranging from -80 mV to +120 mV in increments of +5 mV (100 ms) before being returned to 0 mV (100 ms) to record tail currents. The inter-sweep interval was 2 s. To determine the bioactivity of GxTX-594, Kv2.1 ionic currents were recorded once more, 5 minutes following the wash-in of bath solution also containing 100 nM GxTX-594. Wash-ins were carried out while holding at -100 mV; during

some wash-ins, the membrane potential was pulsed to 0 mV to gauge the time course of binding

and channel inhibition. To exchange solution, 100 uL was washed through the chamber and

removed distally through vacuum tubing to maintain constant bath fluid level.

Ionic current analysis

The average current in the 100 ms prior to step was used to zero-subtract the recording. Outward

current taken as the mean value between 90-100 ms of the channel activation step and used to

calculate and correct for series resistance induced voltage error. Tail current values were derived

from the mean value between 0.2-1.2 ms of the 0 mV tail current step. Tail current was

normalized by the mean activation step current from 50-80 mV and plotted against the estimated

membrane potential, which had been corrected for voltage-error and the calculated liquid

junction potential of 8.5 mV. These tail GV plots were fit with a 4th power Boltzmann (Sack,

Aldrich et al. 2004) and the fit parameters were used for statistical analysis.

Brain slice methods

Hippocampal slice culture preparation and transfection

All experimental procedures were approved by the University of California Davis Institutional

Animal Care and Use Committee and were performed in strict accordance with the Guide for the

Care and Use of Laboratory Animals of the NIH. Animals were maintained under standard light–

dark cycles and had ad libitum access to food and water. Organotypic hippocampal slice cultures

were prepared from postnatal day 5-7 rats, as described (Stoppini, Buchs et al. 1991) and

detailed in a video protocol (Opitz-Araya and Barria 2011). DIV15-30 neurons were transfected

2-6 days before imaging via biolistic gene transfer (160 psi, Helios gene gun, Bio-Rad), as

described in a detailed video protocol (Woods and Zito 2008). 10 μg of plasmid was coated to 6-

8 mg of 1.6 μm gold beads.

Two-photon excitation slice imaging

Image stacks (512 x 512 pixels, 1 mm Z-steps; 0.035 μm/pixel) were acquired using a custom

two-photon excitation microscope (Olympus LUMPLFLN 60XW/IR2 objective, 60x, 1.0 NA)

with two pulsed Ti:sapphire lasers (Mai Tai: Spectra Physics) tuned to 810 nm (for GxTX-594

imaging) and 930 nm (for GFP imaging) and controlled with ScanImage (Pologruto, Sabatini et

al. 2003). After identifying a neuron expressing Kv2.1-GFP, perfusion was stopped and GxTX-

594 was added to the static bath solution to a final concentration of 100 nM. After five minutes

incubation, perfusion was restarted, leading to wash-out of GxTX-594 from the slice bath. Red

and green photons (Chroma: 565dcxr, BG-22 glass, HQ607/45) emitted from the sample were

collected with two sets of photomultiplier tubes (Hamamatsu R3896).

Whole cell voltage clamp for brain slice imaging

Organotypic hippocampal slice cultures (DIV 6-7, not transfected) were transferred to an

imaging chamber with recirculating ACSF maintained at 30°C. To hold the slice to the bottom of

the chamber, a horseshoe-shaped piece of gold wire was used to weight the membrane holding

the slice. ACSF contained (in mM) 127 NaCl, 25 NaHCO$_3$, 25 D-glucose, 2.5 KCl, 1.25

NaH$_2$PO$_4$, 1 MgCl2, 2 CaCl$_2$, 200 nM tetrodotoxin, pH 7.3 and aerated with 95% O$_2$/ 5% CO$_2$

(~310mOsm). 4 mL of 100 nM GxTX-594 in ACSF were used in the circulating bath to allow

the toxin to reach the slice and reach the desired concentration of 100 nM throughout the

circulating bath. Images were acquired beginning 3 minutes after GxTX-594 was added.

Apparent CA1 neurons with GxTX-594 labeling in a Kv2-like pattern were selected for whole-

cell patch clamp. Voltage clamp was achieved using an Axopatch 200B amplifier (Axon

Instruments) controlled with custom software written in Matlab (MathWorks, Inc.). Patch

pipettes (5-7 MΩ) were filled with intracellular solution containing (in mM): 135 Cs-

methanesulfonate, 10 Na$_2$-phosphocreatine, 3 Na-L-ascorbate, 4 NaCl, 10 HEPES, 4 MgCl$_2$, 4

Na$_2$ATP, 0.4 NaGTP, pH 7.2. Neurons were clamped at -70 mV. Input resistance and holding

current were monitored throughout the experiment. Cells were excluded if the pipette series

resistance was higher than 25 MΩ or if the holding current exceeded -100 pA. To activate Kv2

channels, a 50 s depolarizing step from -70 mV to 0 mV was given.

Image analysis

Fluorescence images were analyzed using ImageJ 1.52n software (Schneider, Rasband et al.

2012). Regions of interest (ROIs) encompassed the entire fluorescent region of an individual cell

or neuron, unless mentioned otherwise. ROIs were drawn manually. Analysis of images was

conducted independently by multiple researchers who produced similar results, but analysis was

not conducted in a blinded nor randomized fashion. Fluorescence intensity (*F)* was background

subtracted using the mean *F* of a region that did not contain cells. In experiments with CHO cells

where the bath solution contained 9 nM GxTX-594, the apparent surface membrane of most cells

(40/47) which lack Kv2.1 protein had lower *F* than the background region that did not contain

cells (Fig 4 A, horizontal black dotted line) indicating that the background *F* was overestimated.

Based on the mean *F* from the apparent surface membrane of cells lacking Kv2.1 protein, the

background was overestimated by 24%. The signal from the surface membrane of cells stably

transfected with rat Kv2.1 (Kv2.1-CHO) was, on average, 18x higher than regions that did not

contain cells, making the error generated by our overestimation of the background about 1%. As

the error produced by this background subtraction method was relatively small, it was not

corrected. For F/F_{init} normalization, F_{init} was the mean fluorescence preceding the indicated

voltage stimuli, or the max observed intensity in dose response experiments. Further details of

specific normalization and background subtraction procedures are provided in figure legends.

Time dependence of fluorescence intensity was fit with a monoexponential decay (Eq. 1). For

colocalization analyses the Pearson coefficient was calculated using the JACoP plugin (Bolte and

Cordelieres 2006). Colocalization analyses were conducted within ROIs defining individual

cells. Plotting and curve fitting was performed with Igor Pro 7 or 8 (Wavemetrics), which

performs nonlinear least-squares fits using a Levenberg-Marquardt algorithm. Sample sizes of n

$\geq$ 3 were selected to confirm reproducibility. Sample sizes of n $\geq$ 6 were selected to power

nonparametric statistical comparisons to discern p < 0.01. Error values from individual curve

fittings are standard deviations. All other errors, including error bars, indicate standard errors.

Arithmetic means are reported for intensity measurements and correlation coefficients.

Geometric means are reported for time constants and rates. As the distributions underlying

variability in results are unknown, nonparametric statistical comparisons were conducted with

Mann-Whitney U tests and two-tailed p values reported individually if p > 0.0001.

Model calculations

Predictions of the Scheme I and II models were calculated using Microsoft Excel. A spreadsheet

containing model calculations that can be used to generate model prediction is included as a

supplement (Model Calculation Supplement).

RESULTS

GxTX-594 labels Kv2 proteins

To monitor activation of Kv2 proteins in tissue slices, we synthesized an EVAP compatible with two-photon imaging. We previously presented an EVAP that was a synthetic derivative of GxTX conjugated to a DyLight 550 fluorophore (GxTX-550) (Tilley, Eum et al. 2014). DyLight 550 has poor two-photon excitation properties, and for this study it was replaced with Alexa Fluor 594, a persulfonated Texas Red analog with a large two-photon excitation cross-section and ample spectral separation from green fluorescent protein (GFP), making it well-suited for multiplexed, two-photon excitation imaging experiments (Zito, Knott et al. 2004).

We used solid-phase peptide synthesis to generate a variant of GxTX, an amphiphilic 36-amino acid cystine knot peptide. We synthesized the same peptide used for GxTX-550, Ser13Cys GxTX, where a free thiolate side chain of cysteine 13 is predicted to extend into extracellular solution when the peptide is bound to a voltage sensor (Tilley, Eum et al. 2014). GxTX-1E folds by formation of 3 internal disulfides, and cysteine 13 was differentially protected during oxidative refolding to direct chemoselective conjugation. Following refolding and thiol deprotection, Alexa Fluor 594 C_5 maleimide was condensed with the free thiol, and Ser13Cys (Alexa Fluor 594) GxTX-1E (called "GxTX-594" throughout) was purified (Fig 1).

We performed a dose–response experiment to determine the concentration range where GxTX-594 effectively labels Kv2.1 protein. For these experiments, Kv2.1 was expressed in a CHO-K1 cell line stably transfected with rat Kv2.1 under control of a tetracycline-inducible promoter (Kv2.1-CHO) (Trapani and Korn 2003). Kv2.1 expression was induced with minocycline 2 days prior to experiments, such that all Kv2.1-CHO cells expressed Kv2.1. The K_d of our previously presented EVAP, GxTX-550, was 30.0 ± 3.9 nM which was calculated from electrophysiology at a holding potential of -100 mV (Tilley, Eum et al. 2014) and we guessed that 33 times this K_d, 1000 nM, would be a sufficient upper bound for dose-response experiments with the related GxTX-594 peptide. We performed three tenfold serial dilutions of 1000 nM GxTX-594 to generate the range of concentrations used for this dose–response and applied each concentration of GxTX-594 to Kv2.1-CHO cells for 15 minutes followed by 15 minutes of washout before the next concentration of GxTX-594 was applied (Fig 2). Due to the incomplete equilibration of

labeling at 1 and 10 nM, the true K_d value for GxTX-594 is expected to be somewhat lower, and

our measure is expected to be an overestimate of K_d. A Langmuir binding isotherm was fit to

fluorescence intensity immediately after washout of GxTX-594 from the extracellular solution

and resulted in a K_d of 26.9 ± 8.3 nM (Fig 2 C). This suggests that GxTX-594 has a similar or

higher affinity for resting voltage sensors than GxTX-550. The near saturation of the

fluorescence dose-response with 100 nM GxTX-594 and rate of fluorescence equilibration (Fig.

2 D) suggested that incubation with 100 nM GxTX-594 for 5 minutes should be sufficient for

near maximal labeling of Kv2.1-CHO cells.

We assessed whether GxTX-594 fluorescence on Kv2.1-CHO cells is due to binding Kv2

proteins. Mammals have two pore-forming Kv2 proteins, Kv2.1 and Kv2.2, both of which are

expressed in neurons throughout the brain (Trimmer 1991, Kihira, Hermanstyne et al. 2010,

Bishop, Guan et al. 2015). CHO-K1 cells were transfected with either a rat Kv2.1-GFP or Kv2.2-

GFP construct. We confirmed surface expression of these constructs by whole cell patch clamp

electrophysiology. Transfection with either Kv2.1-GFP or Kv2.2-GFP yielded delayed rectifier

K^+ currents, consistent with Kv2 channels (Fig Supplement 1 A). The CHO-K1 cell line does not

have endogenous K^+ currents (Gamper, Stockand et al. 2005). Two days post transfection,

Kv2.1-GFP or Kv2.2-GFP fluorescence intensity was non-uniform as seen previously for Kv2.1-

GFP (Tilley, Eum et al. 2014), with distinct subcellular regions of high density separated by

regions of much lower density (Fig 3 A). In neurons and other mammalian cell lines similar non-

uniformities have been referred to as "clusters," a term we adopt here to refer to high density

regions of Kv2 channels. In HEK293 cells and neurons, Kv2 clusters are sites where VAP

proteins in the endoplasmic reticulum bind Kv2 proteins in the plasma membrane to form

endoplasmic reticulum–plasma membrane junctions (Johnson, Leek et al. 2018, Kirmiz, Palacio

et al. 2018). A similar mechanism may underly the clustering of Kv2 proteins in CHO cells.

To determine the degree of association between GxTX-594 and Kv2 proteins in CHO cells we

calculated the Pearson Correlation Coefficient (PCC) between the fluorescence intensity from

GxTX-594 and Kv2-GFP proteins (Dunn, Kamocka et al. 2011). A PCC value of 0 signifies

uncorrelated pixels between two images and a PCC value of 1 signifies complete correlation

between pixels from two images. Correlation between pixels from two fluorescent recordings of

an image suggests spatial colocalization of labeled proteins. GxTX-594, and Kv2.1-GFP or

Kv2.2-GFP, had PCC that averaged 0.86 or 0.80, respectively, indicating a high degree of

colocalization of both Kv2 proteins with GxTX-594 (Fig 3 B).

We also assessed if GxTX-594 has a preference for either of the Kv2 proteins under these

labeling conditions. We calculated the ratio of GxTX-594 to GFP fluorescence intensity from

GxTX-594 and Kv2.1-GFP or Kv2.2-GFP. The ratio of GxTX-594 to GFP fluorescence intensity

was similar for Kv2.1-GFP or Kv2.2-GFP (Fig 3 C), consistent with the lack of discrimination of

GxTX between Kv2.1 and Kv2.2 (Herrington, Zhou et al. 2006).

The colocalization of GxTX-594 specifically with Kv2 proteins makes it clear that the dominant

GxTX-594 fluorescence signal from CHO cells is due to binding of Kv2 proteins. However,

GxTX accesses the membrane-embedded voltage sensors of Kv2 proteins by partitioning into the

outer leaflet of the plasma membrane bilayer (Milescu, Bosmans et al. 2009, Gupta, Zamanian et

al. 2015), and a GxTX derivative labeled with an environment sensing fluorophore is detectable

in the membrane of this same cell line under similar conditions (Fletcher-Taylor, Thapa et al.

2020). We wondered if GxTX-594 partitioning into CHO cell membranes could contribute to

fluorescence measurements conducted throughout this study. To assess GxTX-594 partitioning

into the membrane, under conditions used for voltage clamp fluorimetry later in this study, we

compared fluorescence from non-transfected CHO cells with cells transfected with Kv2-GFP

proteins. The membrane of all cells was identified using wheat germ agglutinin (WGA-405) as a

mask to manually draw a region of interest (ROI) where the subsequent analysis was performed.

We saw no indication of GxTX-594 enrichment at the plasma membrane of non-transfected cells

(Fig 4 A). GxTX-594 fluorescence from non-transfected cells had lower intensities than the

extracellular solution (Fig 4 B). We expect lower fluorescence inside of cells as GxTX-594 is

membrane impermeant and excludes the 9 nM GxTX-594 in the extracellular solution from the

cell interior. When GxTX-594 fluorescence was plotted against Kv2.1-GFP fluorescence, a

linear trend was seen with increasing Kv2.1-GFP and GxTX-594 signal (Fig 4 B). Thus GxTX-

594 fluorescence intensity from CHO cells is dependent on the presence of Kv2.1 proteins and

not membrane partitioning. These results indicate that GxTX-594 partitioning into CHO cell

membranes contributes little to the fluorescence measurements made throughout this manuscript.

GxTX-594 modulates Kv2.1 conductance

We performed electrophysiological analyses to determine if GxTX-594 retains the ability to allosterically modulate Kv2.1. GxTX is a partial inverse agonist of Kv2.1 which lowers channel open probability by stabilizing voltage sensors in a resting conformation. Consequently, more positive intracellular voltage is required to activate voltage sensors and achieve the same open probability as without GxTX (Tilley, Angueyra et al. 2019). Previously we estimated that the Kv2.1 complex with GxTX bound to the voltage sensor is at least 5400-fold more stable when the voltage sensor is in its resting conformation (Tilley, Angueyra et al. 2019). Kv2.1 requires more positive intracellular voltage to attain the same level of conductance when bound by GxTX. To characterize the efficacy of GxTX-594 in allosterically modulating Kv2.1 gating, we measured whole cell Kv2.1 ionic currents by voltage clamp in the presence and absence of GxTX-594. From these currents we analyzed the Kv2.1 conductance-voltage (GV) relation by fitting with a 4th power Boltzmann function. The voltage at which the conductance of the fitted function is 50% of maximum, V_{mid}, was +73 mV for 100 nM GxTX-594 (Fig 5). For comparison, the V_{mid} of 100 nM GxTX was +67 mV (Tilley, Angueyra et al. 2019). This small difference in V_{mid} between GxTX and GxTX-594 could potentially result from the different extracellular solutions (see methods), differences in allostery or affinities between GxTX and GxTX-594, or the inherent variability of Kv2.1 GVs between cells. The efficacy of GxTX-594 in shifting the GV indicates that it retains efficacy against Kv2.1 and suggests GxTX-594 acts by the same mechanism as GxTX.

The relationship between GxTX-594 cell surface fluorescence and Kv2.1 voltage activation

To study the relationship between voltage sensor conformational change and GxTX-594 fluorescence, we collected time-lapse fluorescence images from voltage-clamped Kv2.1-CHO cells. We analyzed fluorescence dynamics from airy disk confocal images of cells voltage-clamped in the whole-cell patch clamp configuration. For imaging experiments, Kv2.1-CHO cells were induced for 2 days to express higher densities of Kv2.1 than in Figure 5 ionic current recordings. To minimize voltage clamp error associated with the high level of whole cell Kv2.1 current, the blocker Cs^+ replaced K^+ in the pipette solution. As the dose-response experiments indicate GxTX-594 signal equilibrates within 5 min after application of 100 nM GxTX-594 (Fig

2), we adopted a protocol of applying 100 nM GxTX-594 to Kv2.1-CHO cells for 5 minutes and

then, to reduce the fluorescence in the bath solution, GxTX-594 was diluted to 9 nM before

imaging. The confocal imaging plane containing the most GxTX-594 fluorescence was the glass-

adhered surface of CHO cells. At this imaging plane we conducted time lapse imaging of

voltage-clamped CHO cells and observed a decrease in GxTX-594 fluorescence when the

membrane voltage was stepped from -80 mV to +40 mV. Upon returning the membrane voltage

to -80 mV, GxTX-594 fluorescence increased (Fig 6 A and C). During time lapse imaging of

voltage-clamped cells, we noticed that GxTX-594 fluorescence in the center of the glass-adhered

surface responded more slowly to voltage changes than the periphery. At -80 mV, at the center of

the glass-adhered cell surface, relabeling was incomplete after 500 seconds, however, at the cell

periphery relabeling neared completion within 200 seconds (Fig 6 A). To quantify this

observation, concentric circular ROIs were drawn with the smallest ROI in the center of the

glass-adhered surface. The average fluorescence intensities from each of these ROIs were

compared to each other and an ROI at the cell periphery (Fig 6 B and C). We quantified the rate

of fluorescence change ($k_{\Delta F}$) by fitting the average fluorescence intensities from each ROI with a

monoexponential function:

$F = F_{\infty} + (F_0 - F_{\infty})\, e^{-\frac{(t-t_0)}{\tau}}$ (Equation 1)

Where $\tau = 1/k_{\Delta F}$, t = time, t_0 = time at start of fit, F = fluorescence intensity, F_0 = fluorescence at

start of fit, F_{∞} = fluorescence after infinite time. In response to voltage change, the $k_{\Delta F}$ was

consistently slower at the center of the glass-adhered surface and faster towards the outer

periphery (Fig 6 D and E). While $k_{\Delta F}$ was consistently slower at the center of the glass-adhered

surface we observed that the rate of fluorescence change was more pronounced during labeling at

-80 mV than unlabeling at 40 mV. When cells were held at a membrane potential of -80 mV, $k_{\Delta F}$

at the periphery was approximately 10-fold faster than the $k_{\Delta F}$ at the center of the cell. In

comparison, when the membrane potential was held at 40 mV, $k_{\Delta F}$ at the periphery was

approximately 3-fold faster than the $k_{\Delta F}$ at the center of the cell. This difference in kinetics

suggests that the concentration of freely diffusing GxTX-594 in the restricted space between the

cell membrane and the glass surface is transiently non-uniform after voltage change. Potential

reasons for this phenomenon are considered further in the *Limitations of GxTX-594* section of the

discussion. We found that the average $k_{\Delta F}$ at -80 mV at the center of the glass-adhered surface

was comparable to the $k_{\Delta F}$ at 10 nM from the confluent cells in our dose response experiments

(Fig 2 D) at 0.0014 s^{-1} and 0.0011 s^{-1} respectively. This suggests that the Kv2 voltage sensors in

the unpatched cells for dose response experiments are the same early resting conformation as

voltage-clamped cells at -80 mV (Scholle, Dugarmaa et al. 2004, Tilley, Angueyra et al. 2019).

The gradual change in fluorescence intensity over many seconds after a voltage step is

inconsistent with a fast electrochromic effect leading to fluorescence change, as the change in

fluorescence intensity does not occur instantaneously when the membrane potential is stepped.

Additionally, the slowing of $k_{\Delta F}$ in subcellular regions further from the periphery of cells is

inconsistent with a slower electrochromic effect, as whole cell voltage clamp should render the

membrane surface isopotential.

To avoid complications resulting from the spatial variability of kinetics at the glass-adhered

surface, further voltage-dependent labeling experiments were conducted while imaging at a plane

above the coverslip, where the cell surface has unrestricted access to the bulk extracellular

solution. For consistency, we retained the protocol of initially labeling Kv2 channels with 100

nM GxTX-594 and diluting the solution to 9 nM. In retrospect it is not clear that the 100 nM step

offers any advantage when imaging at a plane above the glass-adhered surface. Voltage stimuli

were given only after fluorescence intensity stabilized at -80 mV; this was at least 9 minutes after

dilution of GxTX-594 to 9 nM. We determined the time required for GxTX-594 to reach a stable

fluorescence after dilution from 100 nM to 9 nM by time-lapse imaging during dilution (Fig

Supplement 2 A, B). The $k_{\Delta F}$ indicated that, on average, equilibration was 85% complete 9

minutes after dilution to 9 nM. After dilution to 9 nM the mean fluorescence intensity decreased

by $39.1 \pm 8.4\%$ and remained stable (Fig Supplement 2 B).

To investigate the relationship between voltage sensor conformational change and GxTX-594

fluorescence, we measured the fluorescence response of GxTX-594 when the membrane voltage

of cells prepared and imaged as described above was stepped from -80 mV to more positive

voltages that ranged from -40 mV to +80 mV (Fig 7 A). ROIs corresponding to the cell surface

were manually identified and average fluorescence intensity quantified from time-lapse

sequences. Both the amplitude and kinetics of fluorescence change from cell surface ROIs are

sensitive to voltage (Fig 7 B), similar to prior findings with GxTX-550 (Tilley, Eum et al. 2014).

The dynamic fluorescence responses of GxTX-594 to voltage changes occurred at

physiologically relevant potentials, suggesting that changes in GxTX-594 labeling intensity or

rate could occur in response to changes in cellular electrical signaling. We observed that the

voltage-dependent reduction in fluorescence equilibrates at a value above background. Even

when cells were given a +80 mV depolarizing voltage stimulus, some GxTX-594 fluorescence

remained (Fig 7 B). Most of the residual fluorescence appeared to be localized to the cell surface

membrane (Fig 7 A), and the amount varied between cells (Fig 7 C). As such surface labeling

was not present on CHO cells in the absence of Kv2 proteins (Fig 4), we consider this residual

fluorescence to originate from voltage-insensitive Kv2 proteins. These voltage-insensitive Kv2

proteins could potentially be at the cell surface and have immobilized voltage sensor which are

stuck in a resting conformation, or they could be internalized Kv2.1–GxTX-594 complexes that

remain just under the cell surface. Kv2.1 channels in HEK293 cells are internalized within 10

minutes at 37°C (Deutsch, Weigel et al. 2012, Weigel, Fox et al. 2012, Fox, Haberkorn et al.

2013, Weigel, Tamkun et al. 2013).

To compare voltage-response properties between cells, we used a normalization procedure to

analyze only the voltage-sensitive fraction of the fluorescence from each cell, which we defined

as the fluorescence that changed between -80 mV and +80 mV. The initial fluorescence at a

holding potential of -80 mV was normalized to 100% F/F_{init}, and residual fluorescence after a

+80 mV step was normalized to 0% F/F_{init} (Fig 7 D). To characterize the voltage dependence of

the Kv2.1–GxTX-594 interaction, fluorescence–voltage (FV) responses were fit with a

Boltzmann function. $f(V) = offset + A(\dfrac{1}{1+e^{-\left(\left(V-V_{\frac{1}{2}}\right)*\frac{zF}{RT}\right)}})$ Equation 2

Where V = voltage, $offset$ = the offset from zero of the Boltzmann distribution, A = the

amplitude, z = number of elementary charges, F = Faraday's constant, R = the universal gas

constant, T = temperature (held at 295 K). This fit had a half maximal voltage midpoint ($V_{1/2}$) of

-27 mV and a steepness (z) of 1.4 elementary charges (e_0) (Fig 7 D lower panel, black line). This

is similar to a fit of voltage sensor movement in Kv2.1-CHO cells, without any GxTX present:

$V_{1/2}$ = -26 mV, z = 1.6 e_0 (Tilley, Angueyra et al. 2019).

To determine the temporal response of GxTX-594 labeling to voltage change, we compared rates

of fluorescence change ($k_{\Delta F}$) at varying step potentials (Fig 7 E) in the same data set that were

analyzed in Figure 7 D. We quantified $k_{\Delta F}$ by fitting the average fluorescence from voltage-

clamped cells with a monoexponential function (Eq 1).

In response to voltage steps from a holding potential of -80 mV to more positive potentials, $k_{\Delta F}$

increased progressively as step potential was increased. After a return to -80 mV, labeling

occurred at a rate similar to unlabeling at -40 mV, indicating that the rate of relaxation to a

labeled/unlabeled equilibrium saturates below -40 mV. We noted the degree of variability in $k_{\Delta F}$

measurements became greater at more positive potentials (Fig 7 E top and bottom panels). At -80

mV there was a 2-fold range in $k_{\Delta F}$ values and a 9-fold range at +80 mV. The relatively low

variation in $k_{\Delta F}$ at -80 mV suggests that fits of the upward relaxation at -80 mV are relatively

consistent in this data set, despite variance in fluorescence intensity after rebinding (Fig 6 C and

Fig 9B). Possible reasons for cell-to-cell variability at more positive voltages are discussed

further in the *Limitations of GxTX-594* section of the Discussion. The average $k_{\Delta F}$–voltage

response was fit with a Boltzmann distribution (Eq. 2) resulting in $V_{1/2}$ = +38 ± 15 mV and z =

1.43 ± 0.35 e_0 (Fig 7 E, black line lower panel). While the precision of this fit is limited by the

variability in $k_{\Delta F}$ values, it is notable that the $V_{1/2}$ and z from the Boltzmann fit is similar to the

voltage dependence of integrated gating charge of the GxTX–Kv2.1 complex: $V_{1/2}$ = +41.3 mV,

z = 1.5 e_0 (Tilley, Angueyra et al. 2019). We later propose a mechanistic underpinning for this

correlation.

This analysis suggests that the dynamics of GxTX-594 labeling convey a surprisingly

straightforward readout of Kv2.1 voltage sensor activation: the magnitude of the response is

determined by the probability of voltage sensor activation of Kv2.1 alone and the kinetics of the

response are determined by the probability of voltage sensor activation in the Kv2.1–GxTX-594

complex.

Temperature dependence of fluorescence kinetics

As part of our efforts to identify the underpinnings of the substantial variation in GxTX-594 $k_{\Delta F}$

measurements, we assessed the temperature dependence of $k_{\Delta F}$. Given the marked temperature

sensitivity reported for Kv2.1 conductance (Yang and Zheng 2014), we wondered if this could be

due to the fluctuations in ambient temperature (27-29°C). To assess the temperature dependence

of GxTX-594 labeling, the cell bath solution was held at either 27°C or 37°C and stepped to 0

mV for a measurement of $k_{\Delta F}$. The average change in $k_{\Delta F}$ over this 10 °C difference, or Q_{10}, was

3.8-fold (Fig Supplement 3). This Q_{10} predicts only a 1.3-fold increase in $k_{\Delta F}$ can be attributed to

a 2°C variation, suggesting only a small fraction of the 6-fold variability could be attributed to

temperature fluctuations. Other potential sources of variability are considered in the *Limitations*

of GxTX-594 section of the discussion.

The relation between voltage sensor activation and GxTX-594 dynamics can be recapitulated

by rate theory modeling.

To enable translation of the intensity of fluorescence from GxTX-594 on a cell surface into a

measure of conformational change, we developed an EVAP model, a series of equations derived

from rate theory that relate cell labeling to voltage sensor activation. The framework of the

EVAP model is generalizable to fluorescent molecular probes that report conformational changes

by a change in binding affinity. In the EVAP model, the proportion of labeled vs. unlabeled Kv2

in a membrane is determined by the proportion of voltage sensors in resting vs. activated

conformations. The model assumes the innate voltage sensitivity of the Kv2 subunit is solely

responsible for controlling voltage dependence. EVAP labeling is voltage-dependent because the

binding and unbinding rates are different for resting and activated conformations of voltage

sensors.

Voltage-activation of Kv2 channels involves many conformational changes (Scholle, Dugarmaa

et al. 2004, Jara-Oseguera, Ishida et al. 2011, Tilley, Angueyra et al. 2019). However, models

which presume independent activation of a voltage sensor in each of the four Kv2.1 subunits

accurately predict many aspects of voltage activation and voltage sensor toxin binding (Lee,

Wang et al. 2003, Tilley, Angueyra et al. 2019). For simplicity, we model Kv2 proteins as having

only resting and activated conformations. With the simplifying assumptions that the

conformational change in each voltage sensor is independent, a rate theory model was developed

consisting of 4 interconnected states (Scheme I: EVAP Allosteric Expansion).

Scheme I: EVAP Allosteric Expansion

Resting labeled   Activated labeled

$k_{on,resting} \cdot [EVAP]$ $k_{off,resting}$ $k_{on,activated} \cdot [EVAP]$ $k_{off,active}$

Resting unlabeled   Activated unlabeled

Scheme II: EVAP Model

$P_{activated}/P_{resting}$ labeled

$k_{on,total} \cdot [EVAP]$ $k_{off,total}$

$P_{activated}/P_{resting}$ unlabeled

When voltage sensors change from resting to activated conformations, the binding rate of the

GxTX-594 EVAP decreases and the unbinding rate increases. When the membrane voltage is

held constant for sufficient time, the proportions of labeled and unlabeled proteins reach an

equilibrium. EVAP labeling requires seconds to equilibrate (Fig 7), whereas Kv2 channel gating

equilibrates in milliseconds (Tilley, Angueyra et al. 2019), three orders of magnitude more

quickly. These distinct time scales of equilibration suggest an approximation to model the EVAP

labeling response: voltage sensor conformations achieve equilibrium quickly such that only their

distribution at equilibrium is expected to greatly impact the kinetics of labeling and unlabeling,

allowing Scheme I to collapse into Scheme II, which depicts the structure of the EVAP Model

used for calculations.

We generated a model of dynamic GxTX-594 fluorescence using the EVAP model and empirical

measurements. In the EVAP model, at any given voltage, there is a probability that a voltage

sensor is either in its resting conformation ($P_{resting}$), or in its activated conformation ($P_{activated}$),

such that $P_{activated} = (1 - P_{resting})$. The equilibrium for voltage sensor activation is then a ratio of

activated to resting voltage sensors ($P_{activated}/P_{resting}$) in which

$$\frac{P_{activated}}{P_{resting}} \, unlabeled = e^{(V - V_{1/2,unlabeled})\frac{zF}{RT}}$$ Equation 3a

$$\frac{P_{activated}}{P_{resting}} \, labeled = e^{(V - V_{1/2,labeled})\frac{zF}{RT}}$$ Equation 3b

where $V_{1/2}$ is the voltage where $P_{activated}/P_{resting,} = 1$. In a prior study, our analysis of the

conductance–voltage relation of Kv2.1 yielded a $V_{1/2} = -32$ mV with $z = 1.5$ e_0 for the early

movement of 4 independent voltage sensors, and found that with a saturating concentration of

GxTX the $V_{1/2} = +42$ mV (Tilley, Angueyra *et al.* 2019, Fig. 1 C). These values were used for

$V_{1/2,unlabeled}$, *z, and* $V_{1/2,labeled}$ respectively (Table 1). To relate voltage sensor activation to labeling

and unlabeling, we used microscopic binding ($k_{on}[EVAP]$) and unbinding (k_{off}) rates that are

distinct for resting and activated voltage sensors. We estimated values for these rates assuming

$k_{\Delta F} = k_{on}[EVAP] + k_{off}$ (Equation 4)

and

$K_d = \frac{k_{off}}{k_{on}}$ (Equation 5)

To calculate the $k_{on,resting}$ and $k_{off,resting}$ values reported in Table 1, we used the saturating value at

negative voltages of the $k_{\Delta F}$–voltage relation (Fig 7 E), and K_d from dose-response imaging (Fig

2 D). In 9 nM GxTX-594, at greater than +40 mV, voltage-dependent unlabeling was nearly

complete, indicating that $k_{off,activated} \gg k_{on,activated}[EVAP]$. We input the saturation of $k_{\Delta F}$ at

positive voltages as $k_{off,activated}$ (Fig 7 E). The slow labeling of activated voltage sensors

confounded attempts to measure $k_{on,activated}$ directly, and we used the statistical thermodynamic

principle of microscopic reversibility (Lewis 1925) to constrain $k_{on,activated}$:

$$\frac{\frac{P_{activated}}{P_{resting}},labeled}{\frac{P_{activated}}{P_{resting}},unlabeled} = \frac{\frac{k_{off,resting}}{k_{on,resting}}}{\frac{k_{off,activated}}{k_{on,activated}}}$$ (Equation 6)

The EVAP model (Scheme II) has only a single microscopic binding rate, $k_{on,total}[EVAP]$, and

unbinding rate, $k_{off,total}$. $k_{on,total}$ is a weighted sum of both $k_{on,resting}$ and $k_{on,activated}$ from Scheme I.

The weights for $k_{on,total}$ are the relative probabilities that unlabeled voltage sensors are resting or

activated, which is determined at any static voltage by an equilibrium constant,

$\frac{P_{activated}}{P_{resting}}unlabeled$:

$$k_{on,total} = k_{on,resting}[EVAP] \cdot \frac{1}{1+\frac{P_{activated}}{P_{resting}}unlabeled} + k_{on,active}[EVAP] \cdot \frac{1}{1+\frac{1}{\frac{P_{activated}}{P_{resting}}unlabeled}}$$

(Equation 7)

Similarly, $k_{off,total}$ is determined by the unbinding rate from resting voltage sensors ($k_{off,resting}$) and

the unbinding rate from activated voltage sensors ($k_{off,activated}$), and weighted such that:

$$k_{off,total} = k_{off,resting} \cdot \frac{1}{1+\frac{P_{activated}}{P_{resting}}labeled} + k_{off,active} \cdot \frac{1}{1+\frac{1}{\frac{P_{activated}}{P_{resting}}labeled}}$$ (Equation 8)

Using $k_{on,total}$ and $k_{off,total}$ we can compute $k_{\Delta F}$ using Equation 4. To test the predictive value of the

EVAP model, we compared the predicted $k_{\Delta F}$ to empirical measurements in 9 nM GxTX-594 and

found the EVAP model (Scheme II) had a $V_{1/2}$ and z that differed by only 3.24 mV and 0.05 e_0

respectively (Fig 7 E, blue line, lower panel).

The EVAP model was also used to predict the magnitude of GxTX-594 fluorescence changes on

cell surfaces. In theory, the ratio of fluorescence at a test voltage to fluorescence at a prior

voltage (F/F_{init}) is equal to the probability that a Kv2 subunit is labeled by GxTX-594 ($P_{labeled}$):

$$\frac{F}{F_{init}} = \frac{P_{labeled}}{P_{labeled,init}}$$ (Equation 9)

The equilibrium $P_{labeled}$ at any voltage can be determined from microscopic binding rates

associated with Scheme II where:

$$P_{labeled} = \frac{1}{1+\frac{K_{d,total}}{[EVAP]}} = \frac{1}{1+\frac{k_{off,total}}{k_{on,total}\cdot[EVAP]}} \qquad \text{(Equation 10)}$$

Using this model, we calculated the fluorescence–voltage relation from $P_{labeled}$ (Equation 9) and

compared these calculated values to our experimental data. The model predicts fluorescence

response with a $V_{1/2}$ and z that differed by only -4.05 mV and 0.06 e_0 respectively (Fig 7 D, blue

line, lower panel).

The EVAP model predicts that changes in labeling report conformational changes of unlabeled Kv2 proteins.

We use the EVAP model to investigate general principles of the relation between voltage sensor

activation and labeling. In particular, we are interested in exploiting EVAP labeling to reveal

information about voltage sensor activation in the unlabeled Kv2 population, akin to how Ca^{2+}

indicators are used to reveal the concentration of free Ca^{2+}. The EVAP model was used to predict

labeling at a series of EVAP concentrations (Fig 8 A). The model does not include EVAP signal

that is insensitive to voltage (Fig 7 C). EVAP labeling is not directly sensitive to voltage but is

determined by the probability that voltage sensors are in an activated conformation (Fig 8 A). As

GxTX-594 binds resting voltage sensors much more rapidly than activated voltage sensors

($k_{on,resting} \gg k_{on,activated}$), the probability that unlabeled voltage sensors are resting determines the

labeling rate, and influences P_{bound}. The rate of fluorescence change ($k_{\Delta F}$) in response to variation

in membrane potential is determined by the probability that voltage sensors in the Kv2.1–GxTX-

594 complexes are at rest (Fig 8 B). Thus $k_{\Delta F}$ is not readily interpreted as a conformational

change in unlabeled Kv2 proteins. Since $k_{\Delta F} = k_{on}[\text{GxTX-594}] + k_{off}$, concentrations of GxTX-

594 where $k_{on}[\text{GxTX-594}] \ll k_{off}$ will cause $k_{\Delta F}$ to be determined by k_{off} and will have little

dependence on the concentration of GxTX-594. The EVAP model predicts that as GxTX-594

concentration decreases, the rate of fluorescence change $k_{\Delta F}$ asymptotically approaches the

probability that labelled channels are active (Fig 8 B). However, the EVAP model predicts that

as GxTX-594 concentration decreases, the change in labeling ($\Delta F/\Delta F_{max}$) asymptotically

approaches the probability that unlabeled voltage sensors are resting (Fig 8 A). Even at 100 nM

GxTX-594, more than 3 times the K_d for the resting conformation, the midpoint of the

Boltzmann distribution describing the change in labeling differs from the midpoint of the QV of

unlabeled Kv2.1 proteins by only 25 mV. Thus, the thermodynamics of the ensemble of the free

EVAP and labeled and unlabeled Kv2 proteins undergird a remarkable and readily interpretable

phenomenon: a decrease in GxTX-594 fluorescence indicates a decrease in the number of

unlabeled voltage sensors that are at rest. Thus, a simple qualitative interpretation can be made: a

GxTX-594 fluorescence change means unlabeled Kv2 voltage sensors have changed

conformation.

Repetitive action potential-like stimuli amplify the GxTX-594 response

Action potentials involving Kv2 currents occur on the millisecond time scale (Liu and Bean

2014, Kimm, Khaliq et al. 2015), orders of magnitude faster than the GxTX-594 response. Thus,

the slow EVAP response will integrate Kv2 conformations occurring over many seconds. The

voltage waveforms that are, arguably, most relevant to Kv2 function are repetitive action

potentials. Kv2 currents can profoundly impact the firing frequency of neurons and the ability of

neurons to sustain high frequency firing (Liu and Bean 2014, Kimm, Khaliq et al. 2015). The

particular roles of Kv2 currents in high frequency firing are underpinned by their deactivation

kinetics, which are much slower than other Kv currents (Liu and Bean 2014). Kv2.1 OFF gating

charge movement reveals that voltage sensor deactivation is even slower than pore closure

(Tilley, Angueyra et al. 2019). Thus, sufficiently high action potential frequencies can prevent

Kv2 proteins from fully deactivating before a next action potential is triggered, creating a kinetic

trap that progressively accumulates activated voltage sensors. This behavior of Kv2 proteins

predicts that a voltage stimulus of repetitive action potentials could evoke a more robust

fluorescence signal than the EVAP model, which assumes continuous equilibrium of voltage

sensors, and thus cannot kinetically trap activated conformations.

We crudely mimicked action potentials with trains of 2 ms voltage steps from -80 to +40 mV,

and observed the changes in fluorescence on GxTX-594–labeled Kv2.1-CHO cells (Fig 9 A, B).

To assess frequency response, step frequency was varied from 0.02 to 200 Hz. Unlabeling

occurred and $k_{\Delta F}$ increased with stimulus frequency (Fig 9 C, D). We compared these

fluorescence responses to those predicted by the EVAP model. Predictions of F/F_{init} and $k_{\Delta F}$

were made from the EVAP model by summing the products of time-averaged probability of

being at each voltage (P_{Vn}) and the fluorescence change predicted at that voltage (ΔF_{Vn})

$$F/F_{init} = (P_{V1} \cdot \Delta F_{V1}) + (P_{V2} \cdot \Delta F_{V2}) + \ldots + (P_{Vn} \cdot \Delta F_{Vn}) \qquad \text{(Equation 11)}$$

For voltage steps from -80 to +40 mV Eq. 11 is:

$$F/F_{init} = (P_{40mV} \cdot \Delta F_{40mV}) + (P_{-80mV} \cdot \Delta F_{-80mV})$$

When the stimulus frequency was less than 50 Hz, the EVAP model did a reasonable job of

predicting fluorescence change. At and above 50 Hz, fluorescence decreased by more than the

EVAP model predicted, even without accounting for a voltage-insensitive fraction of Kv2.1

proteins (Fig 9 C, blue line, lower panel). As discussed above, this divergence from the

equilibrium-based EVAP model is expected at frequencies where voltage steps are shorter than

Kv2 equilibration times. The time constant of activating gating current decay from Kv2.1-CHO

cells was 1.3 ms at +40 mV (Tilley, Angueyra et al. 2019), which means the majority of voltage

sensors are effectively activated during the 2 ms +40 mV steps. In contrast, the time constant of

deactivating gating current decay from Kv2.1-CHO cells was 22 ms at -80 mV (Tilley, Angueyra

et al. 2019), which means Kv2.1 is expected to become kinetically trapped in activated

conformations when stimuli to +40 mV from -80 mV are ~50 Hz or faster. Thus, the amplified

EVAP response appears consistent with voltage sensors failing to deactivate before the next

stimulus, leading to an accumulation of activated voltage sensors, and a more dramatic

fluorescence response than predicted by the EVAP model. An additional factor that could

contribute to more EVAP unlabeling than predicted by the EVAP model is the possibility that

subsaturating GxTX-594 has little impact on voltage sensor deactivation kinetics. Overall, these

dynamics indicate that the magnitude of the change in GxTX-594 fluorescence intensity will be

amplified during repetitive action potentials, a regime of electrophysiological signaling where

Kv2 currents are critical.

The EVAP model indicates that GxTX-594 response kinetics are determined by the

conformational changes of labeled Kv2 proteins. An assessment of EVAP kinetics in response to

increasing frequency of action potential-like voltages, did not reveal major deviations from the

EVAP model, in contrast to the fluorescence intensity measurements from the same cells. We

predicted EVAP kinetic responses as

$k_{\Delta F} = \left(P_{V1} \cdot k_{\Delta F, V1}\right) + \left(P_{V2} \cdot k_{\Delta F, V2}\right) + \dots + \left(P_{Vn} \cdot k_{\Delta F, Vn}\right)$ (Equation 12)

where P_{Vn} is as in Eq. 11 and $k_{\Delta F,n}$ is $k_{\Delta F}$ at that particular voltage. For voltage steps from -80 to

+40 mV Eq. 12 is:

$k_{\Delta F} = \left(P_{40mV} \cdot k_{\Delta F40mV}\right) + \left(P_{-80mV} \cdot k_{\Delta F-80mV}\right)$

The model produced a frequency dependence of $k_{\Delta F}$ that was within range of the empirical data

(Fig 9 D, blue line, lower panel).

This finding is interesting because the response of GxTX-594 labeling to rapidly varying stimuli

indicates that the intensity and kinetics of GxTX-594 fluorescence respond distinctly to different

voltage waveforms. While the change in fluorescence intensity is expected to be magnified by

high-frequency firing, the kinetics of labeling are expected to remain more true to the EVAP

model.

GxTX-594 selectively labels Kv2 proteins

An important consideration for determination of whether GxTX-594 could reveal conformational

changes of endogenous Kv2 proteins is whether the EVAP is selective for Kv2 proteins.

Electrophysiological studies have concluded that the native GxTX peptide is selective for Kv2

channels, with some off-target modulation of A-type Kv4 channels (Herrington, Zhou et al.

2006, Liu and Bean 2014, Speca, Ogata et al. 2014). However, electrophysiological testing

cannot determine whether ligands bind unless they also alter currents (Sack, Stephanopoulos et

al. 2013). Furthermore, structural differences between wild-type GxTX and the GxTX-594

variant could potentially alter selectivity among channel protein subtypes.

To test whether GxTX-594 binds other voltage-gated K$^+$ channel subtypes, we quantified surface

labeling and analyzed colocalization of GxTX-594 with a selection of GFP-labeled voltage-gated

K$^+$ channel subtypes. A series of GFP-K$^+$ channel fusions were identified that express and retain

function with a GFP tag: rat Kv4.2-GFP, rat Kv1.5-GFP, and mouse BK-GFP. Transfection of

each of these channel subtypes into CHO cells resulted in voltage-dependent outward currents,

consistent with cell surface expression of the K^+ channels (Fig Supplement 1 A). In imaging

analysis, the surface membrane was identified by labeling with a fluorescent wheat germ

agglutinin (WGA) conjugate. Transfection of all K^+ channel-GFP constructs resulted in GFP

localization at the cell surface, however the majority of the Kv4.2 and Kv1.5 appeared

intracellularly retained (Fig Supplement 1 B). Consistent with reports in other cell lines, cell

surface localization improved when Kv4.2 was co-transfected with KChIP2 (Shibata, Misonou et

al. 2003) or Kv1.5 with Kvβ2 (Shi, Nakahira et al. 1996) (Fig 10 A).

To quantify the relative labeling of the GFP-tagged K+ channels we measured the ratio of

GxTX-594 to GFP fluorescence. CHO cells expressing the GFP-tagged variants of Kv2.1,

Kv2.2, Kv4.2 + KChIP2, Kv1.5 + Kvβ2, and BK were incubated with 100 nM GxTX-594 for 5

minutes followed by washout with external solution to minimize background fluorescence (Fig

10 A). After background subtraction, the ratio of fluorescence excited at 594 nm (GxTX-594) to

fluorescence excited at 488 nm (GFP) was normalized to the mean 594 nm to 488 nm ratio for

Kv2.1-GFP. The 594:488 ratio was not distinguishable from zero for Kv4.2 + KChIP2, Kv1.5 +

Kvβ2, or BK (Fig 10 B), indicating minimal, if any, binding to these channel subtypes.

Furthermore, no colocalization was apparent between Kv4.2 + KChIP2, Kv1.5 + Kvβ2, or BK

and the residual fluorescence at GxTX-594 wavelengths (Fig 10 C). A similar set of experiments

conducted without the auxiliary subunits of Kv4.2 or Kv1.5 also gave no indication of GxTX-

594 labeling of Kv4.2 or Kv1.5 (Fig Supplement 1 B, C, D). We did not conduct a

comprehensive survey of the more than 80 known mammalian proteins containing voltage sensor

domains, and thus cannot be certain that GxTX-594 does not bind any other voltage sensors, or

other receptors on cell surfaces. However, the lack of labeling of these related voltage-gated K^+

channels indicates that GxTX-594 does not promiscuously label voltage sensors. In all tests

conducted, GxTX-594 was selective for Kv2 proteins.

To determine whether expression of Kv2 proteins embedded in tissue can be imaged with GxTX-

594, we overexpressed Kv2.1-GFP in brain slices. We examined rat brain at CA1 pyramidal

neurons of the hippocampus. We chose CA1 neurons for several reasons: they express Kv2

channels at a density typical of central neurons, the physiology of these neurons have been

intensively studied, and their electrical properties are relatively homogeneous (Misonou,

Mohapatra et al. 2005). Organotypic hippocampal slice cultures prepared from postnatal day 5-7

rats were sparsely transfected with Kv2.1-GFP, resulting in a subset of neurons displaying green

fluorescence. When imaged two to four days after transfection, GFP fluorescence was observed

in the plasma membrane surrounding neuronal cell bodies and proximal dendrites (Fig

Supplement 4 A and B). Six days or more after transfection, Kv2.1-GFP fluorescence organized

into clusters on the surface of the cell soma and proximal processes (Fig 11 A, Fig Supplement 4

C), a pattern consistent with a prior report of endogenous Kv2.1 in CA1 neurons (Misonou,

Mohapatra et al. 2005). After identifying a neuron expressing Kv2.1-GFP, solution flow into the

imaging chamber was stopped and GxTX-594 was added to the static bath solution to a final

concentration of 100 nM. After five minutes of incubation, solution flow was restarted, leading

to wash-out of excess GxTX-594 from the imaging chamber. After wash-out, GxTX-594

fluorescence remained colocalized with Kv2.1-GFP (Fig 11, Fig Supplement 4), indicating that

GxTX-594 is able to permeate through dense neural tissue and bind to Kv2 proteins on neuronal

surfaces. Pearson correlation coefficients confirmed the colocalization of GxTX-594 with Kv2.1-

GFP in multiple slices (Fig 11 C). In most images of Kv2.1-GFP expressing neurons, GxTX-594

also labeled puncta on neighboring neurons that did not express Kv2.1-GFP, but at intensities

that were roughly an order of magnitude dimmer (Fig 11 B, white arrow). The clustered GxTX-

594 fluorescence patterns on the cell body, and proximal processes of CA1 neurons was

strikingly similar to reported patterns of anti-Kv2 immunofluorescence patterns and is consistent

with GxTX-594 labeling endogenous Kv2 proteins in CA1 neurons. While we cannot exclude

the possibility that CA1 neurons have a subset of Kv2 proteins on their surface that are not

labeled by GxTX-594, we saw no indication of Kv2.1-GFP on neuronal surfaces that are not

labeled by GxTX-594.

While we observed GxTX-594 fluorescence that morphologically resembles endogenous Kv2

protein localizations, we also found that GxTX-594 occasionally labels structures not consistent

with Kv2 proteins (Fig Supplement 4 lower panel arrows). This non-Kv2 labeling was most

prevalent at the surface of the hippocampal slices and progressively decreased as the imaging

plane was moved deeper into the tissue (data not shown). Our interpretation of this phenomenon

is that GxTX-594 can accumulate in the dead tissue and debris that is present at the surface of a

hippocampal section after it is cut. For this reason, we analyzed only GxTX-594 fluorescence

with subcellular localizations consistent with Kv2 channels.

GxTX-594 labeling of neurons in brain slice responds to neuronal depolarization

To test whether GxTX-594 labeling of neurons in brain slices is consistent with labeling of

endogenous Kv2 voltage sensors, we determined whether GxTX-594 labeling responds to

voltage changes in tissue. First, we looked for Kv2-like labeling patterns on CA1 pyramidal

neurons in untransfected brain slices bathed in 100 nM GxTX-594. With two-photon excitation,

optical sections are thinner than the neuronal cell bodies, and fluorescence appeared as puncta

circumscribed by dark intracellular spaces (Fig 12 A). This was similar to the patterns of

fluorescence in Kv2.1-GFP transfected slices (Fig 11 B), and consistent with the punctate

expression pattern of Kv2.1 in CA1 pyramidal neurons seen in fixed brain slices (Misonou,

Mohapatra et al. 2005).

We tested whether the punctate fluorescence was voltage-sensitive using voltage clamp. To

ensure voltage clamp of the neuronal cell body, slices were bathed in tetrodotoxin to block Na+

channels, and Cs+ was included in the patch pipette solution to block K+ currents. In each

experiment, whole-cell configuration was achieved with a GxTX-594 labeled neuron, and

holding potential was set to -70 mV. At this point, time-lapse imaging of a two-photon excitation

optical section was initiated. Depolarization to 0 mV resulted in loss of fluorescence from a

subset of puncta at the perimeter of the voltage-clamped neuron (Fig 12 B red arrows,

Supplemental Movie). Other fluorescent puncta appeared unaltered by the 0 mV step (Fig 12 B,

white arrows). These puncta could represent Kv2 proteins on a neighboring cell, off-target

labeling by GxTX-594, or voltage-insensitive Kv2 proteins, possibly due to near-surface

internalization. To assess if the fluorescence changes were due to the voltage change or random

fluctuations during imaging, we compared fluorescence of the apparent cell membrane region to

regions more distal from the voltage-clamped cell body. To quantify this fluorescence change, an

ROI 30 pixels (1.2 μm) wide containing the apparent membrane of the cell body was compared

to other regions within each image (Fig 12 C). The region containing the membrane of the cell

body lost fluorescence during the 0 mV step (Fig 12 D, ROI 1) while neither of the regions more

distal (ROI 2, ROI 3), nor the intracellular region (ROI 0) showed a similar change. In multiple

slices, the voltage-dependent decrease of fluorescence was seen specifically in the region of the

voltage-clamped neuronal membrane (Fig 12 E). The kinetics of fluorescence response of the

voltage-clamped membrane region were similar between slices (Fig 12 F, K).

To determine whether the fluorescence response to depolarization was driven by the Kv2-like

puncta on the cell membrane, the fluorescence along a path containing the apparent cell

membrane was selected by drawing a path connecting fluorescent puncta surrounding the dark

cell body region, and averaging fluorescence in a region 10 pixels wide (0.4 μm) centered on this

path (Fig 12 G, yellow line). The fluorescence intensity along the path of the ROI revealed

distinct peaks corresponding to puncta (Fig 12 H, red trace). After stepping the neuron to 0 mV,

the intensity of fluorescence of a subset of puncta lessened (Fig 12 H, black trace, peaks

1,3,4,5,8). While the data is quite noisy, it is clear that even more reduction is observed in

individual fluorescence puncta in brain slice (Fig 12, I blue line). This suggests the voltages

sensors in these functionally-selected puncta are extensively activated. The EVAP model

predicts only a 55% reduction of fluorescence from CHO cells stepped to 0 mV in 100 nM

GxTX. However, the GxTX-594 concentration within tissue may be more dilute than in the bath

solution, and consequently the sensitive fraction calculated from the EVAP model is a lower

bound. Other puncta maintained or increased their brightness after depolarization (Fig 12 H,

black trace, peaks 2, 6, 7), but it was not certain whether these voltage-insensitive puncta

correspond to Kv2 proteins on the surface of the same voltage-clamped neuron. When the

kinetics of fluorescence intensity decay of individual voltage-sensitive puncta were fit with Eq.

1, $k_{\Delta F}$ values were similar between puncta (Fig 12 I), consistent with these spatially separated

puncta all being on the surface of the voltage-clamped neuron. The fluorescence changes

measured in these puncta were similar to those predicted by the Fig 8 model (Fig 12 I, blue line),

although the variability of these measurements was substantial. To address whether the

fluorescence change at the cell membrane was driven by decreases in regions of punctate

fluorescence, the punctate and non-punctate fluorescence intensity changes were analyzed

separately. The regions with fluorescence intensities above average for the path (Fig 12 H,

dashed line) were binned as one group. The above-average group, by definition, contained all

punctate fluorescence. When comparing the fluorescence before and during the 0 mV step, the

regions that were initially below average fluorescence maintained the same intensity ($103 \pm 8\%$);

regions that were initially above average fluorescence decreased in intensity ($70 \pm 8\%$) (Fig 12

J). This suggests that the detectable unlabeling along the cell membrane originated in the Kv2-

like puncta, not the regions of lower fluorescence intensity between them.

To determine whether the dynamics of the voltage-dependent responses of GxTX-594

fluorescence on neurons in brain slices were consistent with labeling of Kv2.1, we performed

experiments with Kv2.1-expressing CHO cells under similar conditions as brain slice: 100 nM

GxTX-594, 30°C, Cs^+-containing patch pipette solution. The rate of fluorescence changes in

Kv2.1-CHO cells were similar to neurons in brain slices, and consistent with the $k_{\Delta F}$ of 0.06 s^{-1}

predicted by Fig 8 model (Fig 12 K). However, the data underlying of $k_{\Delta F}$ measurements was

noisy, limiting our ability to detect differences.

Discussion

The molecular targeting, conformation selectivity, and spatial precision of fluorescence from

GxTX-594 enable identification of where in tissue the conformational status of Kv2 voltage

sensors becomes altered. However, the utility of GxTX-594 as an EVAP is limited by several

factors including emission intensity, variability between experiments, and inhibition of Kv2

proteins. We discuss the potential utility and limitations of the EVAP mechanism underlying

GxTX-594.

Unique capabilities of GxTX-594

The Kv2 EVAP presented here is the only imaging method we are aware of for measuring

voltage-sensitive conformational changes of a specific, endogenous protein. As GxTX binding

selectively stabilizes the fully resting conformation of Kv2.1 voltage sensors (Tilley, Angueyra

et al. 2019), GxTX-594 labeling is expected to report the fully resting conformation of the Kv2

voltage sensor, in which the first gating charge of the Kv2 S4 segment is in the gating charge

transfer center (Tao, Lee et al. 2010). Images of GxTX-594 fluorescence reveal this

conformation's occurrence, with subcellular spatial resolution. Importantly, the EVAP model we

develop allows deconvolution of the behavior of unlabeled Kv2 proteins from Kv2 proteins

inhibited by GxTX-594. This enables the subcellular locations where Kv2 voltage sensing occurs

to be seen for the first time.

Electrophysiological approaches can detect the voltage-sensitive K^+ conductance of Kv2

channels. However, the majority of Kv2 proteins on cell surface membranes do not function as

channels and are nonconducting (Benndorf, Koopmann et al. 1994, O'Connell, Loftus et al.

2010). Kv2 proteins dynamically regulate cellular physiology by nonconducting functions

(Antonucci, Lim et al. 2001, Singer-Lahat, Sheinin et al. 2007, Feinshreiber, Singer-Lahat et al.

2010, Dai, Manning Fox et al. 2012, Fox, Haberkorn et al. 2015, Johnson, Leek et al. 2018,

Kirmiz, Palacio et al. 2018, Kirmiz, Vierra et al. 2018, Kirmiz, Gillies et al. 2019, Vierra, Kirmiz

et al. 2019). The Kv2 EVAP reports on the conformation of Kv2 voltage sensors independently

from ion conductance, enabling the study of voltage sensor involvement in Kv2's nonconducting

physiological functions. We present this EVAP as a prototype molecular probe for imaging

voltage sensing by endogenous proteins, in tissue, with molecular specificity.

Here, during our initial testing of GxTX-594, we observed that the majority of Kv2 protein

detected at discrete individual clusters was voltage sensitive. While it may not sound surprising

to find that voltage-gated ion channel proteins are voltage sensitive, the voltage sensors of

surface-expressed proteins can be immobilized. For example, gating charge of the L-type Ca^{2+}

channel Cav1.2 is immobilized until it is bound by an intracellular protein (Turner, Anderson et

al. 2020). Kv2.1 channel function is extensively regulated by neurons. In rat CA1 neurons the

clustered Kv2 channels are proposed to be nonconducting (Fox, Loftus et al. 2013). Our results

show that clustered Kv2 proteins in rat CA1 neurons remain voltage-sensitive. Interestingly,

when Kv2.1 is expressed in CHO cells, a fraction of the GxTX-594 fluorescence is voltage

insensitive (Fig 7 C). This observation is consistent with voltage sensor immobilization of some

surface-expressed Kv2.1 protein, although it could be due to intracellular Kv2.1–GxTX-594

proteins that appear to be at the cell surface.

We used images of GxTX-594 fluorescence to measure the coupling between endogenous Kv2

proteins and membrane potential at specific, subcellular anatomical locations. Similarly, GxTX-

594 imaging should detect changes of voltage sensor status when Kv2 proteins become engaged

or disengaged from nonconducting functions such as formation of plasma membrane–endoplasmic reticulum junctions (Antonucci, Lim et al. 2001, Fox, Haberkorn et al. 2015, Kirmiz, Palacio et al. 2018), regulation of exocytosis (Singer-Lahat, Sheinin et al. 2007, Feinshreiber, Singer-Lahat et al. 2010), regulation of insulin secretion (Dai, Manning Fox et al. 2012) interaction with kinases, phosphatases and SUMOylases (Misonou, Mohapatra et al. 2004, Park, Mohapatra et al. 2006, Dai, Kolic et al. 2009, Cerda and Trimmer 2011, McCord and Aizenman 2013), formation of specialized subcellular calcium signaling domains (Vierra, Kirmiz et al. 2019), and interactions with astrocytic end feet (Du, Tao-Cheng et al. 1998). GxTX-594 could potentially reveal conformational changes in organs throughout the body where Kv2 proteins are expressed, which include muscle, thymus, spleen, kidney, adrenal gland, pancreas, lung, and reproductive organs (Bocksteins 2016).

Limitations of GxTX-594

There are important limitations to the GxTX-594 approach, and of the underlying EVAP mechanism generally. We discuss several limitations which are worth considering in the design of any studies.

GxTX-594 is slow

The kinetics of GxTX-594 labeling are limited to measuring changes in Kv2 activity on a time scale of tens of seconds. Thus GxTX-594 will not provide sufficient time resolution to distinguish fast electrical signaling events. However, we have found that the temporal resolution of GxTX-594 is compatible with live imaging and electrophysiology experiments. The response time of GxTX-594 is far slower than the kinetics of Kv2 conformational change, limiting measurements to the probability, averaged over time, that voltage sensors are resting or active. It is worth noting that the probability of a conformation's occurrence is a valuable measure, and the ultimate quantitation of many biophysical studies of ion channels (e.g., open probability, steady state conductance, gating charge-voltage relation).

GxTX-594 dynamics are altered in confined extracellular spaces

The kinetics of GxTX-594 dynamics varied within different regions of the same CHO cell. GxTX-594 fluorescence in the restricted space at the center of the glass-adhered surface

appeared to respond more slowly to voltage changes than the periphery, which has better access

to the bath solution (Fig 6). The location dependence of $k_{\Delta F}$ was more pronounced during GxTX-

594 labeling at -80 mV than unlabeling at +40 mV, and such a difference can be explained by the

distinct voltage dependent affinities of Kv2.1 for GxTX-594. We suspect that the more extreme

location dependence at -80 mV is due to a high density of Kv2.1 binding sites in the restricted

extracellular space between the cell membrane and glass coverslip, such that GxTX-594 is

depleted from solution by binding Kv2.1 before reaching the center of the cell. After unbinding

at +40 mV, each GxTX-594 molecule is expected to be less likely to rebind to Kv2.1 proteins

which are in an activated, low affinity conformation (Tilley, Eum et al. 2014, Tilley, Angueyra et

al. 2019).

The different GxTX-594 dynamics in a restricted space is significant because the space

extracellular to Kv2.1 channel clusters is restricted in many neurons. The Kv2 channel clusters in

the plasma membrane of hippocampal and cortical interneurons are tightly associated with

astrocytic end feet and the cleft between these two membranes is a few nanometers wide (Du,

Tao-Cheng et al. 1998). Experiments investigating Kv2 channels localized to these subcellular

regions should consider the limitations of using EVAP probes in restricted spaces; most notably

the finding that labeling dynamics are more dramatically affected than unlabeling dynamics (Fig

6 D and E).

GxTX-594 dynamics are variable

The GxTX-594 response rates and amplitudes were notably variable between Kv2.1 CHO cells.

We did not succeed in constraining this variability and offer conjecture about its origins.

Some variability results from fits of $k_{\Delta F}$ where the final value of the relaxation was poorly

determined by the data (Fig 3C, 7B, 9B and Fig Supplement 2). Fluctuations in temperature

during imaging likely had only a small effect on the fluorescence dynamics of GxTX-594;

considering the measured Q_{10} of 3.8, the $2°C$ temperature fluctuations between the experiments

in Figures 7 and 9 should produce only 1.3-fold change in $k_{\Delta F}$. We also expect that changes in

fluorescence due to photobleaching of the Alexa Flour 594 fluorophore were limited, based on

the observation that fluorescence remained unchanged in cells that were not voltage clamped but

in the same field of view as voltage clamped cells receiving voltage stimulus (Fig 9 A).

We noticed that results were more consistent between stimuli of the same cell and the variability

was greatest between cells. We suspect that cell-to-cell differences in Kv2.1 conformational

equilibria are responsible for much of the variability in GxTX-594 response. The Kv2.1

conductance-voltage relation is regulated by many cellular pathways including kinases,

phosphatases, and SUMOylases (Misonou, Mohapatra et al. 2004, Park, Mohapatra et al. 2006,

Dai, Kolic et al. 2009, Cerda and Trimmer 2011, McCord and Aizenman 2013). Large cell-to-

cell variation in Kv2.1 conductance-voltage, and gating charge-voltage relations have been

reported in CHO cells (McCrossan, Roepke et al. 2009, Tilley, Eum et al. 2014, Kang, Vanoye et

al. 2019, Tilley, Angueyra et al. 2019).

In this study, when we predicted the voltage sensor $V_{1/2}$ of Kv2.1 from electrophysiology we

observed a 6.4 mV standard deviation with a range of 19 mV, and this variance appeared to be

exacerbated by GxTX-594 (Fig 5E) (9.7 mV standard deviation and range of 36 mV). As EVAP

dynamics and the GV are both determined by voltage sensor activation, variability in the GxTX-

594 response is expected. The hypothesis that cell-to-cell variation in fluorescence dynamics is

due to the inherent variability of Kv2.1 voltage sensor activation could be more definitively

tested by identifying if a correlation exists between the $V_{1/2}$ of the QV and fluorescence–voltage

relationship from individual cells labeled with GxTX-594. While we have not attempted this, the

structure of the variance in GxTX-594 fluorescence–voltage relationships is informative. The

fluorescence–voltage relationships compiled from many cells become more variable near the

midpoint of relevant voltage sensor movements. The response amplitude, F/F_{init}, (Fig 7 D), is

determined by unlabeled voltage sensor activation, and appears most variable near the $V_{1/2}$ of the

unlabeled QV relation, -32 mV (Tilley, Angueyra et al. 2019). The response kinetics, $k_{\Delta F,}$ are

determined by activation of labeled voltage sensors and appear to become increasingly variable

at voltages higher than -20 mV (Fig 7 E bottom panel). Despite the increasing variability, the $V_{1/2}$

and z from the Boltzmann fit of the $k_{\Delta F}$–voltage relationship to measurements from many cells

was remarkably close to the same measurements in the QV of the GxTX–Kv2.1 complex with a

$V_{1/2}$ and z that differ by 3.3 mV and 0.07 e_0 respectively.

Another possibility is that variability in surface membrane composition undergirds the variability

of the GxTX-594 response. GxTX affinities are influenced by the dynamically changing

complement of lipids in the plasma membrane lipid composition. Lipids bind to the voltage

sensing domain near the GxTX binding site (Milescu, Bosmans et al. 2009, Gupta, Zamanian et

al. 2015). Sphingomyelinase D treatment, which alters the membrane composition by converting

sphingomyelin to ceramide 1-phosphate, has been shown to enhance GxTX affinity for Kv2.1 by

4-fold (Milescu, Bosmans et al. 2009). Voltage activation of Kv2.1 is also affected by

sphingomyelinase treatments. While variation in lipid composition is expected to cause variation

in GxTX-594 dynamics, we do not know the degree to which the lipid composition varies

between the Kv2.1-CHO cells in our studies.

Signal Intensity and Noise

Fluorescence signal intensity limits interpretation of EVAP labeling. We found that variability in

the intensity of GxTX-594 labeling of CA1 neurons limited the quantitative interpretation with

the EVAP model. However, the signal to noise was clearly sufficient to identify voltage sensing

of endogenous Kv2 proteins. As CA1 hippocampal neurons express Kv2 proteins at a density

typical of central neurons (Misonou, Mohapatra et al. 2004, Vacher, Mohapatra et al. 2008,

Speca, Ogata et al. 2014). Thus, we expect that GxTX-594 imaging could be implemented in

most brain regions with similar signal to noise characteristics. In other cell types, such as

neurons of the subiculum, or the inner segment of photoreceptors, Kv2 proteins are expressed at

far higher densities (Maletic-Savatic, Lenn et al. 1995, Gayet-Primo, Yaeger et al. 2018).

Improved signal to noise would be expected from such cells that express higher densities of Kv2

proteins. Kv2 proteins are also expressed by many other cell types throughout the body

(Bocksteins 2016). It remains to be seen what the signal to noise characteristics are for EVAP

Kv2 labeling in these other tissues.

GxTX-594 inhibits Kv2 proteins

GxTX-based probes inhibit the Kv2 proteins they label by stabilizing the resting conformation of

Kv2 voltage sensors. The Kv2.1–GxTX-594 complex does not open to conduct K^+ ions in the

physiological voltage range (Fig 5). Thus, GxTX-594 depletes the population of Kv2 proteins

responding normally to physiological stimuli. While the concentration of an EVAP can be

lowered to minimize interference with Kv2 activity, fluorescence intensity is diminished and

further limits the ability to detect subtle physiological changes. With a related probe, we

explored the impact of decreasing toxin concentration on fluorescence response of an EVAP

(Tilley, Eum et al. 2014). Here, we demonstrate that lower concentration and physiological

stimuli are not always required for scientifically meaningful implementation of the probe. Even

when using GxTX-594 concentrations that grossly inhibit most Kv2 proteins, the behavior of

unlabeled Kv2 proteins can be calculated using EVAP model we have developed. Of course, the

electrical feedback within cells will be altered by such protocols.

The EVAP model is oversimplified

Another limitation of the analysis developed here is that the model of Kv2 voltage sensor

conformational change is an oversimplification. The gating dynamics of Kv2 channels are more

complex than our model (Islas and Sigworth 1999, Scholle, Dugarmaa et al. 2004, Jara-

Oseguera, Ishida et al. 2011, Tilley, Angueyra et al. 2019). Under some conditions the

assumption of voltage sensor independence will limit the model's predictive power. Additionally,

the model of GxTX-594 labeling developed here assumes that voltage sensors are in continuous

equilibrium. These deviations from equilibrium likely explain the deviation of the model from

the data in response to high frequency voltage steps (Fig 9 C, D).

Conformation-selective probes reveal conformational changes of endogenous proteins

Measurements of dynamic labeling by a conformation-selective probe such as GxTX can enable

deduction of how unlabeled proteins behave. This is perhaps counterintuitive because GxTX

inhibits voltage sensor movement of the Kv2 protein it binds, and thus only bound proteins

generate optical signals. This approach is analogous to calcium imaging experiments, which have

been spectacularly informative about physiological calcium signaling (Yang and Yuste 2017),

despite the fact that no optical signals originate from the physiologically relevant free Ca^{2+}, only

from Ca^{2+} that is chelated by a dye. In all such experiments, fluorescence from Ca^{2+}-bound dyes

is deconvolved using the statistical thermodynamics of Ca^{2+} binding to calculate free Ca^{2+}

(Adams 2010). Similarly, GxTX-based probes dynamically bind to unlabeled Kv2 proteins, and

the binding rate is dependent on the probability that unlabeled voltage sensors are in a resting

conformation (Fig 8). Thus, the conformations of unlabeled Kv2 proteins influence the dynamics

of labeling with GxTX-based probes. Consequently, the dynamics of labeling reveal the

conformations of unlabeled Kv2 proteins.

Deployment of GxTX-594 to report conformational changes of endogenous proteins

demonstrates that conformation-selective ligands can be used to image occurrence of the

conformations they bind to. The same principles of action apply to any conformation-selective

labeling reagent, suggesting that probes for conformational changes of many different proteins

could be developed. Probes could conceivably be developed from the many other voltage sensor

toxins and other gating modifiers that act by a similar mechanism as GxTX, yet target the

voltage sensors of different ion channel proteins (McDonough, Mintz et al. 1997, Sack, Aldrich

et al. 2004, Catterall, Cestele et al. 2007, Swartz 2007, Schmalhofer, Calhoun et al. 2008, Peretz,

Pell et al. 2010, McCormack, Santos et al. 2013, Ahuja, Mukund et al. 2015, Dockendorff,

Gandhi et al. 2018, Zhang, Sharma et al. 2018). Conformation-selective binders have been

engineered for a variety of other proteins, and methods to quantify conformational changes from

their fluorescence are needed. For example, fluorescently-labeled conformation-selective binders

have revealed that endocytosed GPCRs continue to remain in a physiologically activated

conformation (Irannejad, Tomshine et al. 2013, Tsvetanova, Irannejad et al. 2015, Eichel and

von Zastrow 2018). A means to determine the conformational equilibria of GPCRs from

fluorescence images has not yet been developed. We suggest that the statistical thermodynamic

framework developed here could provide a starting point for more quantitative interpretation of

other conformation-selective molecular probes.

Acknowledgements

We thank Jim Trimmer (University of California, Davis) for numerous discussions, and

constructive critical reading of an early version of the manuscript. We thank Georgeann Sack

(Afferent LLC) for critical reading, editing, and feedback. This research was supported by US

National Institutes of Health grants R01NS096317, U01NS090581, R21EY026449, and

T32GM007377. P. Thapa was supported by American Heart Association postdoctoral fellowship

17POST33670698. GxTX variants were synthesized at the Molecular Foundry of the Lawrence

Berkeley National Laboratory under US Department of Energy contract DE-AC02-05CH11231.

The authors declare no competing financial interests.

Author contributions (Credit nomenclature)

Parashar Thapa: Conceptualization, Formal analysis, Investigation, Methodology, Visualization,

Writing- original draft, Writing- reviewing & editing

Robert Stewart: Conceptualization, Formal analysis, Investigation, Methodology, Visualization,

Writing- original draft, Writing- reviewing & editing

Rebecka J. Sepela: Conceptualization, Formal analysis, Investigation, Methodology,

Visualization, Writing- original draft, Writing- reviewing & editing

Oscar Vivas: Investigation, Methodology, Writing- reviewing & editing

Laxmi K. Parajuli: Investigation, Methodology, Writing- reviewing & editing

Mark Lillya: Conceptualization, Formal analysis, Investigation, Methodology, Visualization

Sebastian Fletcher-Taylor: Investigation, Methodology, Writing- reviewing & editing

Bruce E. Cohen: Conceptualization, Funding acquisition, Project administration, Supervision,

Writing- reviewing & editing

Karen Zito: Conceptualization, Funding acquisition, Investigation, Methodology, Project

administration, Supervision, Writing- original draft, Writing- reviewing & editing

Jon T. Sack: Conceptualization, Formal analysis, Funding acquisition, Investigation,

Methodology, Project administration, Supervision, Visualization, Writing- original draft,

Writing- reviewing & editing

References

Adams, S. R. (2010). "How calcium indicators work." <u>Cold Spring Harb Protoc</u> **2010**(3): pdb.top70.

Aggarwal, S. K. and R. MacKinnon (1996). "Contribution of the S4 segment to gating charge in the Shaker K+ channel." <u>Neuron</u> **16**(6): 1169-1177.

Ahuja, S., S. Mukund, L. Deng, K. Khakh, E. Chang, H. Ho, S. Shriver, C. Young, S. Lin, J. P. Johnson, Jr., P. Wu, J. Li, M. Coons, C. Tam, B. Brillantes, H. Sampang, K. Mortara, K. K. Bowman, K. R. Clark, A. Estevez, Z. Xie, H. Verschoof, M. Grimwood, C. Dehnhardt, J. C. Andrez, T. Focken, D. P. Sutherlin, B. S. Safina, M. A. Starovasnik, D. F. Ortwine, Y. Franke, C. J. Cohen, D. H. Hackos, C. M. Koth and J. Payandeh (2015). "Structural basis of Nav1.7 inhibition by an isoform-selective small-molecule antagonist." <u>Science</u> **350**(6267): aac5464.

An, W. F., M. R. Bowlby, M. Betty, J. Cao, H. P. Ling, G. Mendoza, J. W. Hinson, K. I. Mattsson, B. W. Strassle, J. S. Trimmer and K. J. Rhodes (2000). "Modulation of A-type potassium channels by a family of calcium sensors." <u>Nature</u> **403**(6769): 553-556.

Antonucci, D. E., S. T. Lim, S. Vassanelli and J. S. Trimmer (2001). "Dynamic localization and clustering of dendritic Kv2.1 voltage-dependent potassium channels in developing hippocampal neurons." <u>Neuroscience</u> **108**(1): 69-81.

Armstrong, C. M. and F. Bezanilla (1973). "Currents related to movement of the gating particles of the sodium channels." <u>Nature</u> **242**(5398): 459-461.

Benndorf, K., R. Koopmann, C. Lorra and O. Pongs (1994). "Gating and conductance properties of a human delayed rectifier K+ channel expressed in frog oocytes." <u>J Physiol</u> **477**(Pt 1): 1-14.

Bezanilla, F. (2008). "How membrane proteins sense voltage." <u>Nat Rev Mol Cell Biol</u> **9**(4): 323-332.

Bezanilla, F. (2018). "Gating currents." <u>J Gen Physiol</u> **150**(7): 911-932.

Bishop, H. I., D. Guan, E. Bocksteins, L. K. Parajuli, K. D. Murray, M. M. Cobb, H. Misonou, K. Zito, R. C. Foehring and J. S. Trimmer (2015). "Distinct Cell- and Layer-Specific Expression Patterns and Independent Regulation of Kv2 Channel Subtypes in Cortical Pyramidal Neurons." <u>J Neurosci</u> **35**(44): 14922-14942.

Bocksteins, E. (2016). "Kv5, Kv6, Kv8, and Kv9 subunits: No simple silent bystanders." <u>J Gen Physiol</u> **147**(2): 105-125.

Bolte, S. and F. P. Cordelieres (2006). "A guided tour into subcellular colocalization analysis in light microscopy." <u>J Microsc</u> **224**(Pt 3): 213-232.

Catterall, W. A., S. Cestele, V. Yarov-Yarovoy, F. H. Yu, K. Konoki and T. Scheuer (2007). "Voltage-gated ion channels and gating modifier toxins." <u>Toxicon</u> **49**(2): 124-141.

Cerda, O. and J. S. Trimmer (2011). "Activity-dependent phosphorylation of neuronal Kv2.1 potassium channels by CDK5." <u>J Biol Chem</u> **286**(33): 28738-28748.

Dai, X. Q., J. Kolic, P. Marchi, S. Sipione and P. E. Macdonald (2009). "SUMOylation regulates Kv2.1 and modulates pancreatic beta-cell excitability." <u>J Cell Sci</u> **122**(Pt 6): 775-779.

Dai, X. Q., J. E. Manning Fox, D. Chikvashvili, M. Casimir, G. Plummer, C. Hajmrle, A. F. Spigelman, T. Kin, D. Singer-Lahat, Y. Kang, A. M. Shapiro, H. Y. Gaisano, I. Lotan and P. E. Macdonald (2012). "The voltage-dependent potassium channel subunit Kv2.1 regulates insulin secretion from rodent and human islets independently of its electrical function." Diabetologia **55**(6): 1709-1720.

Deutsch, E., A. V. Weigel, E. J. Akin, P. Fox, G. Hansen, C. J. Haberkorn, R. Loftus, D. Krapf and M. M. Tamkun (2012). "Kv2.1 cell surface clusters are insertion platforms for ion channel delivery to the plasma membrane." Mol Biol Cell **23**(15): 2917-2929.

Dockendorff, C., D. M. Gandhi, I. H. Kimball, K. S. Eum, R. Rusinova, H. I. Ingolfsson, R. Kapoor, T. Peyear, M. W. Dodge, S. F. Martin, R. W. Aldrich, O. S. Andersen and J. T. Sack (2018). "Synthetic Analogues of the Snail Toxin 6-Bromo-2-mercaptotryptamine Dimer (BrMT) Reveal That Lipid Bilayer Perturbation Does Not Underlie Its Modulation of Voltage-Gated Potassium Channels." Biochemistry **57**(18): 2733-2743.

Du, J., J. H. Tao-Cheng, P. Zerfas and C. J. McBain (1998). "The K+ channel, Kv2.1, is apposed to astrocytic processes and is associated with inhibitory postsynaptic membranes in hippocampal and cortical principal neurons and inhibitory interneurons." Neuroscience **84**(1): 37-48.

Dunn, K. W., M. M. Kamocka and J. H. McDonald (2011). "A practical guide to evaluating colocalization in biological microscopy." American Journal of Physiology-Cell Physiology **300**(4): C723-C742.

Eichel, K. and M. von Zastrow (2018). "Subcellular Organization of GPCR Signaling." Trends Pharmacol Sci **39**(2): 200-208.

Feinshreiber, L., D. Singer-Lahat, R. Friedrich, U. Matti, A. Sheinin, O. Yizhar, R. Nachman, D. Chikvashvili, J. Rettig, U. Ashery and I. Lotan (2010). "Non-conducting function of the Kv2.1 channel enables it to recruit vesicles for release in neuroendocrine and nerve cells." J Cell Sci **123**(Pt 11): 1940-1947.

Fletcher-Taylor, S., P. Thapa, R. J. Sepela, R. Kaakati, V. Yarov-Yarovoy, J. T. Sack and B. E. Cohen (2020). "Distinguishing Potassium Channel Resting State Conformations in Live Cells with Environment-Sensitive Fluorescence." ACS Chem Neurosci.

Fox, P. D., C. J. Haberkorn, E. J. Akin, P. J. Seel, D. Krapf and M. M. Tamkun (2015). "Induction of stable ER-plasma-membrane junctions by Kv2.1 potassium channels." J Cell Sci **128**(11): 2096-2105.

Fox, P. D., C. J. Haberkorn, A. V. Weigel, J. L. Higgins, E. J. Akin, M. J. Kennedy, D. Krapf and M. M. Tamkun (2013). "Plasma membrane domains enriched in cortical endoplasmic reticulum function as membrane protein trafficking hubs." Mol Biol Cell **24**(17): 2703-2713.

Fox, P. D., R. J. Loftus and M. M. Tamkun (2013). "Regulation of Kv2.1 K(+) conductance by cell surface channel density." J Neurosci **33**(3): 1259-1270.

Frech, G. C., A. M. VanDongen, G. Schuster, A. M. Brown and R. H. Joho (1989). "A novel potassium channel with delayed rectifier properties isolated from rat brain by expression cloning." Nature **340**(6235): 642-645.

Gamper, N., J. D. Stockand and M. S. Shapiro (2005). "The use of Chinese hamster ovary (CHO)
cells in the study of ion channels." J Pharmacol Toxicol Methods **51**(3): 177-185.

Gayet-Primo, J., D. B. Yaeger, R. A. Khanjian and T. Puthussery (2018). "Heteromeric
KV2/KV8.2 Channels Mediate Delayed Rectifier Potassium Currents in Primate Photoreceptors."
J Neurosci **38**(14): 3414-3427.

Gordon, E., T. K. Roepke and G. W. Abbott (2006). "Endogenous KCNE subunits govern Kv2.1
K+ channel activation kinetics in Xenopus oocyte studies." Biophys J **90**(4): 1223-1231.

Gupta, K., M. Zamanian, C. Bae, M. Milescu, D. Krepkiy, D. C. Tilley, J. T. Sack, V. Yarov-
Yarovoy, J. I. Kim and K. J. Swartz (2015). "Tarantula toxins use common surfaces for interacting
with Kv and ASIC ion channels." Elife **4**: e06774.

Herrington, J., Y. P. Zhou, R. M. Bugianesi, P. M. Dulski, Y. Feng, V. A. Warren, M. M. Smith,
M. G. Kohler, V. M. Garsky, M. Sanchez, M. Wagner, K. Raphaelli, P. Banerjee, C. Ahaghotu, D.
Wunderler, B. T. Priest, J. T. Mehl, M. L. Garcia, O. B. McManus, G. J. Kaczorowski and R. S.
Slaughter (2006). "Blockers of the delayed-rectifier potassium current in pancreatic beta-cells
enhance glucose-dependent insulin secretion." Diabetes **55**(4): 1034-1042.

Irannejad, R., J. C. Tomshine, J. R. Tomshine, M. Chevalier, J. P. Mahoney, J. Steyaert, S. G.
Rasmussen, R. K. Sunahara, H. El-Samad, B. Huang and M. von Zastrow (2013). "Conformational
biosensors reveal GPCR signalling from endosomes." Nature **495**(7442): 534-538.

Islas, L. D. and F. J. Sigworth (1999). "Voltage sensitivity and gating charge in Shaker and Shab
family potassium channels." J Gen Physiol **114**(5): 723-742.

Jara-Oseguera, A., I. G. Ishida, G. E. Rangel-Yescas, N. Espinosa-Jalapa, J. A. Perez-Guzman, D.
Elias-Vinas, R. Le Lagadec, T. Rosenbaum and L. D. Islas (2011). "Uncoupling charge movement
from channel opening in voltage-gated potassium channels by ruthenium complexes." J Biol Chem
286(18): 16414-16425.

Johnson, B., A. N. Leek, L. Sole, E. E. Maverick, T. P. Levine and M. M. Tamkun (2018). "Kv2
potassium channels form endoplasmic reticulum/plasma membrane junctions via interaction with
VAPA and VAPB." Proc Natl Acad Sci U S A **115**(31): E7331-e7340.

Kaczmarek, L. K. (2006). "Non-conducting functions of voltage-gated ion channels." Nat Rev
Neurosci **7**(10): 761-771.

Kang, S. K., C. G. Vanoye, S. N. Misra, D. M. Echevarria, J. D. Calhoun, J. B. O'Connor, K. L.
Fabre, D. McKnight, L. Demmer, P. Goldenberg, L. E. Grote, I. Thiffault, C. Saunders, K. A.
Strauss, A. Torkamani, J. van der Smagt, K. van Gassen, R. P. Carson, J. Diaz, E. Leon, J. E.
Jacher, M. C. Hannibal, J. Litwin, N. R. Friedman, A. Schreiber, B. Lynch, A. Poduri, E. D. Marsh,
E. M. Goldberg, J. J. Millichap, A. L. George Jr and J. A. Kearney (2019). "Spectrum of KV2.1
Dysfunction in KCNB1-Associated Neurodevelopmental Disorders." Annals of Neurology **86**(6):
899-912.

Kihira, Y., T. O. Hermanstyne and H. Misonou (2010). "Formation of heteromeric Kv2 channels
in mammalian brain neurons." J Biol Chem **285**(20): 15048-15055.

Kimm, T., Z. M. Khaliq and B. P. Bean (2015). "Differential Regulation of Action Potential Shape and Burst-Frequency Firing by BK and Kv2 Channels in Substantia Nigra Dopaminergic Neurons." J Neurosci **35**(50): 16404-16417.

Kirmiz, M., T. E. Gillies, E. J. Dickson and J. S. Trimmer (2019). "Neuronal ER-plasma membrane junctions organized by Kv2-VAP pairing recruit Nir proteins and affect phosphoinositide homeostasis." J Biol Chem **294**(47): 17735-17757.

Kirmiz, M., S. Palacio, P. Thapa, A. N. King, J. T. Sack and J. S. Trimmer (2018). "Remodeling neuronal ER-PM junctions is a conserved nonconducting function of Kv2 plasma membrane ion channels." Mol Biol Cell **29**(20): 2410-2432.

Kirmiz, M., N. C. Vierra, S. Palacio and J. S. Trimmer (2018). "Identification of VAPA and VAPB as Kv2 Channel-Interacting Proteins Defining Endoplasmic Reticulum-Plasma Membrane Junctions in Mammalian Brain Neurons." J Neurosci **38**(35): 7562-7584.

Lee, H. C., J. M. Wang and K. J. Swartz (2003). "Interaction between extracellular Hanatoxin and the resting conformation of the voltage-sensor paddle in Kv channels." Neuron **40**(3): 527-536.

Lewis, G. N. (1925). "A New Principle of Equilibrium." Proc Natl Acad Sci U S A **11**(3): 179-183.

Li, H., W. Guo, H. Xu, R. Hood, A. T. Benedict and J. M. Nerbonne (2001). "Functional expression of a GFP-tagged Kv1.5 alpha-subunit in mouse ventricle." Am J Physiol Heart Circ Physiol **281**(5): H1955-1967.

Lin, M. Z. and M. J. Schnitzer (2016). "Genetically encoded indicators of neuronal activity." Nat Neurosci **19**(9): 1142-1153.

Liu, P. W. and B. P. Bean (2014). "Kv2 channel regulation of action potential repolarization and firing patterns in superior cervical ganglion neurons and hippocampal CA1 pyramidal neurons." J Neurosci **34**(14): 4991-5002.

Long, S. B., E. B. Campbell and R. Mackinnon (2005). "Voltage sensor of Kv1.2: structural basis of electromechanical coupling." Science **309**(5736): 903-908.

Long, S. B., X. Tao, E. B. Campbell and R. MacKinnon (2007). "Atomic structure of a voltage-dependent K+ channel in a lipid membrane-like environment." Nature **450**(7168): 376-382.

MacDonald, P. E., A. M. Salapatek and M. B. Wheeler (2003). "Temperature and redox state dependence of native Kv2.1 currents in rat pancreatic beta-cells." J Physiol **546**(Pt 3): 647-653.

Maletic-Savatic, M., N. J. Lenn and J. S. Trimmer (1995). "Differential spatiotemporal expression of K+ channel polypeptides in rat hippocampal neurons developing in situ and in vitro." J Neurosci **15**(5 Pt 2): 3840-3851.

McCord, M. C. and E. Aizenman (2013). "Convergent Ca2+ and Zn2+ signaling regulates apoptotic Kv2.1 K+ currents." Proc Natl Acad Sci U S A **110**(34): 13988-13993.

McCormack, K., S. Santos, M. L. Chapman, D. S. Krafte, B. E. Marron, C. W. West, M. J. Krambis, B. M. Antonio, S. G. Zellmer, D. Printzenhoff, K. M. Padilla, Z. Lin, P. K. Wagoner, N. A. Swain, P. A. Stupple, M. de Groot, R. P. Butt and N. A. Castle (2013). "Voltage sensor

interaction site for selective small molecule inhibitors of voltage-gated sodium channels." Proc Natl Acad Sci U S A **110**(29): E2724-2732.

McCrossan, Z. A., T. K. Roepke, A. Lewis, G. Panaghie and G. W. Abbott (2009). "Regulation of the Kv2.1 Potassium Channel by MinK and MiRP1." Journal of Membrane Biology **228**(1): 1-14.

McDonough, S. I., I. M. Mintz and B. P. Bean (1997). "Alteration of P-type calcium channel gating by the spider toxin omega-Aga-IVA." Biophys J **72**(5): 2117-2128.

Milescu, M., F. Bosmans, S. Lee, A. A. Alabi, J. I. Kim and K. J. Swartz (2009). "Interactions between lipids and voltage sensor paddles detected with tarantula toxins." Nat Struct Mol Biol **16**(10): 1080-1085.

Misonou, H., D. P. Mohapatra, M. Menegola and J. S. Trimmer (2005). "Calcium- and metabolic state-dependent modulation of the voltage-dependent Kv2.1 channel regulates neuronal excitability in response to ischemia." J Neurosci **25**(48): 11184-11193.

Misonou, H., D. P. Mohapatra, E. W. Park, V. Leung, D. Zhen, K. Misonou, A. E. Anderson and J. S. Trimmer (2004). "Regulation of ion channel localization and phosphorylation by neuronal activity." Nat Neurosci **7**(7): 711-718.

Murakoshi, H., G. Shi, R. H. Scannevin and J. S. Trimmer (1997). "Phosphorylation of the Kv2.1 K^+ Channel Alters Voltage-Dependent Activation." Molecular Pharmacology **52**(5): 821-828.

O'Connell, K. M., R. Loftus and M. M. Tamkun (2010). "Localization-dependent activity of the Kv2.1 delayed-rectifier K+ channel." Proc Natl Acad Sci U S A **107**(27): 12351-12356.

Opitz-Araya, X. and A. Barria (2011). "Organotypic hippocampal slice cultures." J Vis Exp(48).

Park, K. S., D. P. Mohapatra, H. Misonou and J. S. Trimmer (2006). "Graded regulation of the Kv2.1 potassium channel by variable phosphorylation." Science **313**(5789): 976-979.

Peltola, M. A., J. Kuja-Panula, S. E. Lauri, T. Taira and H. Rauvala (2011). "AMIGO is an auxiliary subunit of the Kv2.1 potassium channel." EMBO Rep **12**(12): 1293-1299.

Peretz, A., L. Pell, Y. Gofman, Y. Haitin, L. Shamgar, E. Patrich, P. Kornilov, O. Gourgy-Hacohen, N. Ben-Tal and B. Attali (2010). "Targeting the voltage sensor of Kv7.2 voltage-gated K+ channels with a new gating-modifier." Proc Natl Acad Sci U S A **107**(35): 15637-15642.

Plant, L. D., E. J. Dowdell, I. S. Dementieva, J. D. Marks and S. A. Goldstein (2011). "SUMO modification of cell surface Kv2.1 potassium channels regulates the activity of rat hippocampal neurons." J Gen Physiol **137**(5): 441-454.

Pologruto, T. A., B. L. Sabatini and K. Svoboda (2003). "ScanImage: flexible software for operating laser scanning microscopes." Biomed Eng Online **2**: 13.

Ramu, Y., Y. Xu and Z. Lu (2006). "Enzymatic activation of voltage-gated potassium channels." Nature **442**(7103): 696-699.

Sack, J. T., R. W. Aldrich and W. F. Gilly (2004). "A gastropod toxin selectively slows early transitions in the Shaker K channel's activation pathway." J Gen Physiol **123**(6): 685-696.

Sack, J. T., N. Stephanopoulos, D. C. Austin, M. B. Francis and J. S. Trimmer (2013). "Antibody-guided photoablation of voltage-gated potassium currents." J Gen Physiol **142**(3): 315-324.

Schmalhofer, W. A., J. Calhoun, R. Burrows, T. Bailey, M. G. Kohler, A. B. Weinglass, G. J. Kaczorowski, M. L. Garcia, M. Koltzenburg and B. T. Priest (2008). "ProTx-II, a selective inhibitor of NaV1.7 sodium channels, blocks action potential propagation in nociceptors." Mol Pharmacol **74**(5): 1476-1484.

Schneider, C. A., W. S. Rasband and K. W. Eliceiri (2012). "NIH Image to ImageJ: 25 years of image analysis." Nat Methods **9**(7): 671-675.

Schneider, M. F. and W. K. Chandler (1973). "Voltage dependent charge movement of skeletal muscle: a possible step in excitation-contraction coupling." Nature **242**(5395): 244-246.

Scholle, A., S. Dugarmaa, T. Zimmer, M. Leonhardt, R. Koopmann, B. Engeland, O. Pongs and K. Benndorf (2004). "Rate-limiting reactions determining different activation kinetics of Kv1.2 and Kv2.1 channels." J Membr Biol **198**(2): 103-112.

Seoh, S. A., D. Sigg, D. M. Papazian and F. Bezanilla (1996). "Voltage-sensing residues in the S2 and S4 segments of the Shaker K+ channel." Neuron **16**(6): 1159-1167.

Shi, G., K. Nakahira, S. Hammond, K. J. Rhodes, L. E. Schechter and J. S. Trimmer (1996). "Beta subunits promote K+ channel surface expression through effects early in biosynthesis." Neuron **16**(4): 843-852.

Shibata, R., H. Misonou, C. R. Campomanes, A. E. Anderson, L. A. Schrader, L. C. Doliveira, K. I. Carroll, J. D. Sweatt, K. J. Rhodes and J. S. Trimmer (2003). "A fundamental role for KChIPs in determining the molecular properties and trafficking of Kv4.2 potassium channels." J Biol Chem **278**(38): 36445-36454.

Singer-Lahat, D., A. Sheinin, D. Chikvashvili, S. Tsuk, D. Greitzer, R. Friedrich, L. Feinshreiber, U. Ashery, M. Benveniste, E. S. Levitan and I. Lotan (2007). "K+ channel facilitation of exocytosis by dynamic interaction with syntaxin." J Neurosci **27**(7): 1651-1658.

Speca, D. J., G. Ogata, D. Mandikian, H. I. Bishop, S. W. Wiler, K. Eum, H. J. Wenzel, E. T. Doisy, L. Matt, K. L. Campi, M. S. Golub, J. M. Nerbonne, J. W. Hell, B. C. Trainor, J. T. Sack, P. A. Schwartzkroin and J. S. Trimmer (2014). "Deletion of the Kv2.1 delayed rectifier potassium channel leads to neuronal and behavioral hyperexcitability." Genes Brain Behav **13**(4): 394-408.

Stoppini, L., P. A. Buchs and D. Muller (1991). "A simple method for organotypic cultures of nervous tissue." J Neurosci Methods **37**(2): 173-182.

Swartz, K. J. (2007). "Tarantula toxins interacting with voltage sensors in potassium channels." Toxicon **49**(2): 213-230.

Tanabe, T., K. G. Beam, J. A. Powell and S. Numa (1988). "Restoration of excitation-contraction coupling and slow calcium current in dysgenic muscle by dihydropyridine receptor complementary DNA." Nature **336**(6195): 134-139.

Tao, X., A. Lee, W. Limapichat, D. A. Dougherty and R. MacKinnon (2010). "A gating charge transfer center in voltage sensors." Science **328**(5974): 67-73.

Tilley, D. C., J. M. Angueyra, K. S. Eum, H. Kim, L. H. Chao, A. W. Peng and J. T. Sack (2019). "The tarantula toxin GxTx detains K(+) channel gating charges in their resting conformation." J Gen Physiol **151**(3): 292-315.

Tilley, D. C., K. S. Eum, S. Fletcher-Taylor, D. C. Austin, C. Dupre, L. A. Patron, R. L. Garcia, K. Lam, V. Yarov-Yarovoy, B. E. Cohen and J. T. Sack (2014). "Chemoselective tarantula toxins report voltage activation of wild-type ion channels in live cells." Proc Natl Acad Sci U S A **111**(44): E4789-4796.

Trapani, J. G. and S. J. Korn (2003). "Control of ion channel expression for patch clamp recordings using an inducible expression system in mammalian cell lines." BMC Neurosci **4**: 15.

Trimmer, J. S. (1991). "Immunological identification and characterization of a delayed rectifier K+ channel polypeptide in rat brain." Proc Natl Acad Sci U S A **88**(23): 10764-10768.

Tsvetanova, N. G., R. Irannejad and M. von Zastrow (2015). "G protein-coupled receptor (GPCR) signaling via heterotrimeric G proteins from endosomes." J Biol Chem **290**(11): 6689-6696.

Turner, M., D. E. Anderson, P. Bartels, M. Nieves-Cintron, A. M. Coleman, P. B. Henderson, K. N. M. Man, P. Y. Tseng, V. Yarov-Yarovoy, D. M. Bers, M. F. Navedo, M. C. Horne, J. B. Ames and J. W. Hell (2020). "α-Actinin-1 promotes activity of the L-type Ca(2+) channel Ca(v) 1.2." Embo j **39**(5): e102622.

Vacher, H., D. P. Mohapatra and J. S. Trimmer (2008). "Localization and targeting of voltage-dependent ion channels in mammalian central neurons." Physiol Rev **88**(4): 1407-1447.

Vierra, N. C., M. Kirmiz, D. van der List, L. F. Santana and J. S. Trimmer (2019). "Kv2.1 mediates spatial and functional coupling of L-type calcium channels and ryanodine receptors in mammalian neurons." Elife **8**.

Weigel, A. V., P. D. Fox, E. J. Akin, K. H. Ecklund, M. M. Tamkun and D. Krapf (2012). "Size of cell-surface Kv2.1 domains is governed by growth fluctuations." Biophys J **103**(8): 1727-1734.

Weigel, A. V., M. M. Tamkun and D. Krapf (2013). "Quantifying the dynamic interactions between a clathrin-coated pit and cargo molecules." Proceedings of the National Academy of Sciences **110**(48): E4591-E4600.

Woods, G. and K. Zito (2008). "Preparation of gene gun bullets and biolistic transfection of neurons in slice culture." J Vis Exp(12).

Xu, H., T. Li, A. Rohou, C. P. Arthur, F. Tzakoniati, E. Wong, A. Estevez, C. Kugel, Y. Franke, J. Chen, C. Ciferri, D. H. Hackos, C. M. Koth and J. Payandeh (2019). "Structural Basis of Nav1.7 Inhibition by a Gating-Modifier Spider Toxin." Cell **176**(5): 1238-1239.

Yang, F. and J. Zheng (2014). "High temperature sensitivity is intrinsic to voltage-gated potassium channels." Elife **3**: e03255.

Yang, W. and R. Yuste (2017). "In vivo imaging of neural activity." Nat Methods **14**(4): 349-359.

Zagotta, W. N., T. Hoshi, J. Dittman and R. W. Aldrich (1994). "Shaker potassium channel gating. II: Transitions in the activation pathway." J Gen Physiol **103**(2): 279-319.

Zhang, A. H., G. Sharma, E. A. B. Undheim, X. Jia and M. Mobli (2018). "A complicated complex: Ion channels, voltage sensing, cell membranes and peptide inhibitors." Neurosci Lett **679**: 35-47.

Zhang, G., S. Zheng, H. Liu and P. R. Chen (2015). "Illuminating biological processes through site-specific protein labeling." Chem Soc Rev **44**(11): 3405-3417.

1 Zito, K., G. Knott, G. M. Shepherd, S. Shenolikar and K. Svoboda (2004). "Induction of spine
2 growth and synapse formation by regulation of the spine actin cytoskeleton." <u>Neuron</u> **44**(2): 321-
3 334.

Figure Legends

Figure 1: *Synthesis of GxTX-594.*

A. Molecular model of GxTX-594. Scale bar is 10 angstroms. Backbone of GxTX peptide

depicted with ribbon. Cys13(maleimide-Alexa 594) depicted with CPK atoms.

B. HPLC chromatogram of Ser13Cys GxTX. Gradient described in *Materials and Methods -*

GxTX Synthesis. Ser13Cys GxTX eluted at 12.6 minutes, peak 1, which corresponds to

33% acetonitrile. MALDI-TOF MS profile of peak 1 (Inset). Rel. Int. is Relative

Intensity.

C. HPLC chromatogram of GxTX-594 conjugation reaction between Alexa Fluor 594-

maleimide and Ser13Cys GxTX. Peak 1 is Ser13Cys GxTX (Retention time: 12.8

minutes, 33% acetonitrile), peak 2 is a minor product from conjugation, and peak 3 is

GxTX-594, the major product from conjugation (Retention time: 16.4 minutes, 35%

acetonitrile). The fractions corresponding to peak 3 were combined.

D. HPLC chromatogram of the combined peak 3 fractions from panel C, a GxTX-594

preparation used in this study. 2 μL of 13.1 μM GxTX-594 diluted in 200 μL of 0.1 %

TFA was injected. MALDI-TOF MS profile of the combined peak 3 fractions (Inset).

Figure 2: *Dose-response characteristics of GxTX-594 labeling.*

A. Fluorescence from Kv2.1-CHO cells incubated in indicated concentrations of GxTX-594

for 15 minutes then washed out before imaging. Imaging plane was near the glass-

adhered cell surface. Fluorescence shown corresponds GxTX-594 (Magenta). Scale bar is

20 μm.

B. Time-lapse fluorescence intensity from the dose-response experiment shown in A.

Fluorescence was not background-subtracted. F_{max} is intensity while cells were incubated

in 1000 nM GxTX-594.

C. Relative fluorescence intensity of GxTX-594 that remains on cells immediately after

wash-out of indicated concentrations of GxTX-594 from the extracellular solution.

Symbols correspond to each of 3 experiments. Black line is the fit of a Langmuir

isotherm to obtain a dissociation constant (K_d) for GxTX-594; K_d = 26.9 nM ± 8.3.

D. Labeling or unlabeling kinetics for GxTX-594 at indicated concentrations. Symbols

correspond to each of 3 experiments. The $k_{\Delta F}$ from each point was obtained by a

monoexponential fit (Eq. 1). Error bars represent the standard deviation of. Data were fit

with Equation 4 (k_{on} = 6.372 x 10^{-5} ± 0.049 x 10^{-5} nM^{-1} s^{-1}; k_{off} = 5.929 x 10^{-4} ± 0.043 x

10^{-4} s^{-1}).

Figure 3: *GxTX-594 colocalizes with Kv2-GFP.*

A. Fluorescence from CHO cells transfected with Kv2.1-GFP (Top) or Kv2.2-GFP (Bottom)

and labeled with GxTX-594. Optical sections were near the glass-adhered cell surface.

Cells were incubated with 100 nM GxTX-594 for 5 minutes and rinsed before imaging.

Fluorescence shown corresponds to emission of GFP (Left), GxTX-594 (Middle), or an

overlay of GFP and GxTX-594 (Right). Scale bars are 20 μm.

B. Pearson correlation coefficient for GxTX-594 colocalization with Kv2.1-GFP or Kv2.2-

GFP. In panels B and C pixel intensities were background subtracted before analyses by

subtracting the average fluorescence of a background ROI that did not contain cells, from

the ROI containing the cell. All data from the same application of GxTX-594. Circles

represent measurements from individual cells.

C. Ratio of fluorescence intensity when excited at 594 nm versus 488 nm from cells

expressing Kv2.1-GFP or Kv2.2-GFP. The consistent ratio between Kv2.1 and Kv2.2

indicates that GxTX-594 labels the two proteins equivalently.

1 Figure 4: ***GxTX-594 labeling of surface membranes requires Kv2 protein.***

2 A. Fluorescence from live CHO cells transfected with Kv2.1-GFP (upper row) or Kv2.2-

3   GFP (lower row) and labeled with GxTX-594. Airy disk imaging was at a plane above

4   the glass-adhered surface. Cells were incubated with 100 nM GxTX-594 and 5 µg/mL

5   WGA-405 then diluted to 9 nM GxTX-594 and 0.45 µg/mL WGA-405 before imaging.

6   Scale bars are 20 µm. Fluorescence shown was excited by a 594 nm laser (Column 1),

7   488 nm laser (Column 2) or 405 nm laser (Column 3). Scale bars are 20 µm.

8 B. Fluorescence intensity from 594 nm excitation versus 488 nm excitation. Cell preparation

9   was transfected with Kv2.1-GFP. Each point represents one cell. Cells with obvious GFP

10   fluorescence are green points, cells without are black points. Mean background

11   fluorescence from a region that did not contain cells is indicated by dashed lines. Red line

12   represents a linear fit of cells with obvious GFP fluorescence.

Figure 5: *GxTX-594 modulates Kv2.1 conductance.*

A. Representative Kv2.1-CHO current response under whole cell voltage-clamp. Cells were

given 100 ms, 5 mV increment voltage steps ranging from -80 mV (blue) to +120 mV

(red) and then stepped to 0 mV to record tail currents. The holding potential was -100

mV.

B. Kv2.1 currents from the same cell 5 minutes after the addition of 100 nM GxTX-594.

C. Normalized conductance–voltage relationships from Kv2.1 current before application of

GxTX-594 (n = 13). Different symbols correspond to individual cells and the green

corresponds to cell in panel A.

D. Normalized conductance–voltage relationships in 100 nM GxTX-594 (n = 11).

E. Mean midpoint of each of 4 independent voltage sensors in the 4^{th} power Boltzmann fit

($V_{1/2}$) before (-31 ± 6 mV SD) and after (27 ± 10 mV SD) 100 nM GxTX-594. p <

0.0001, two-tailed T-test.

F. Mean elementary charge associated with Boltzmann fit (z) in units of elementary charge

before (1.5 ± 0.3 e^+ SD) and after (1.0 ± 0.4 e^+ SD) 100 nM GxTX-594. p = 0.0013, two-

tailed T-test.

G. Mean midpoint of conductance change in the 4^{th} power Boltzmann fit (V_{mid}) before (-

2 ± 6 mV SD) and after (73 ± 13 mV SD) 100 nM GxTX-594. p < 0.0001, two-tailed T-

test.

Figure 6: ***Extracellular access can impact GxTX-594 labeling kinetics.***

A. Time lapse airy disk images of the glass-adhered surface of a voltage-clamped Kv2.1-CHO cell in 9 nM GxTX-594. Pseudocolored images indicate maximal intensities in white. Time index is in the upper left of each panel and membrane potential is indicated in the upper right. Scale bar in lower right of last panel is 10 μm

B. Airy disk image of the glass adhered surface of a voltage-clamped Kv2.1-CHO cell in 9 nM GxTX-594. Gray lines indicate boundaries of ROIs. ROI 1, 2, and 3 are concentric circles each with a respective diameter of 1.8, 4.9, and 9.1 μm. ROI 4 was hand-drawn to contain the apparent cell surface. In all cells analyzed, ROI 1-3 were concentric circles of the same sizes, while ROI 4 varied based on cell shape. Scale bar is 10 μm.

C. Representative traces of 594 fluorescence intensity response to voltage changes. Red lines are monoexponential fits (Eq. 1): 40 mV step ROI 1 $k_{\Delta F} = 4.29 \times 10^{-2} \pm 0.26 \times 10^{-2}$ s^{-1}, ROI 2 $k_{\Delta F} = 4.39 \times 10^{-2} \pm 0.16 \times 10^{-2}$ s^{-1}, ROI 3 $k_{\Delta F} = 5.65 \times 10^{-2} \pm 0.1.5 \times 10^{-2}$ s^{-1}, ROI 4 $k_{\Delta F} = 9.69 \times 10^{-2} \pm 0.33 \times 10^{-2}$ s^{-1}. -80 mV step ROI 1 $k_{\Delta F} = 4.27 \times 10^{-4} \pm 0.11 \times 10^{-4}$ s^{-1}, ROI 2 $k_{\Delta F} = 7.999 \times 10^{-4} \pm 0.053 \times 10^{-4}$ s^{-1}, ROI 3 $k_{\Delta F} = 2.7556 \times 10^{-3} \pm 0.0074 \times 10^{-3}$ s^{-1}, and ROI 4 $k_{\Delta F} = 5.46 \times 10^{-3} \pm 0.13 \times 10^{-3}$ s^{-1}. Background for subtraction was the average intensity of a region that did not contain cells over the time course of the voltage protocol. Each trace was normalized to initial fluorescence intensity before the application of the voltage stimulus.

D. Circles indicate $k_{\Delta F}$ at +40 mV from individual cells. Circle coloring indicates data from the same cell.

E. Circles indicate $k_{\Delta F}$ at -80 mV from individual cells.

| 1 | Figure 7: ***GxTX-594 labeling responds to transmembrane voltage.*** |

2. A. Fluorescence intensities from an optical section of a voltage-clamped Kv2.1- CHO cell in
3. 9 nM GxTX-594. Pseudocolored images indicate maximal intensities in white. Rows
4. indicate voltage step taken from a holding potential of -80 mV. Columns indicate time
5. after voltage step. Scale bar is 10 μm.
6. B. GxTX-594 fluorescence during steps to indicated voltages. Smooth lines are
7. monoexponential fits (Eq. 1): -40 mV $k_{\Delta F}$ = 2.15 x 10^{-2} ± 0.22 x 10^{-2} s^{-1}: 0 mV $k_{\Delta F}$ =
8. 1.279 x 10^{-1} ± 0.023 x 10^{-1} s^{-1}, 40 mV $k_{\Delta F}$ = 2.492 x 10^{-1} ± 0.062 x 10^{-1} s^{-1} and 80 mV $k_{\Delta F}$
9. = 4.20 x 10^{-1} ± 0.11 x 10^{-1} s^{-1}. ROIs were hand-drawn around the apparent cell surface
10. membrane based on GxTX-594 fluorescence. 0% was set by subtraction of background
11. which was the average intensity of a region that did not contain cells over the time course
12. of the voltage protocol. For each trace, 100% was set from the initial fluorescence
13. intensity at -80 mV before the subsequent voltage step.
14. C. Fluorescence intensity remaining at the end of 50 s steps to +80 mV. Each circle
15. represents one cell. Background subtraction as in panel B.
16. D. Voltage-dependence of fluorescence intensity at the end of 50 s steps. For each cell,
17. 100% was set from the initial fluorescence intensity at -80 mV before the first step to
18. another voltage. Cells did not always recover to initial fluorescence intensity during
19. the -80 mV holding period between voltage steps. (Top panel) Circle coloring indicates
20. data from the same cell and lines connect points from the same cell. Gray circles
21. represent data shown in B. (Bottom panel) black bars represent the mean F/F_{init} at each
22. voltage and error bars represent the standard error of the mean. Black line is fit of a first-
23. order Boltzmann equation (Eq. 2): $V_{1/2}$ = -27.4 ± 2.5 mV, z = 1.38 ± 0.13 $e_\circ$. Green line is
24. prediction from the EVAP model at 9 nM GxTX.
25. E. Voltage dependence of fluorescence intensity kinetics ($k_{\Delta F}$). (Top panel) circle coloring is
26. the same as panel D. (Bottom panel) black bars represent the average $k_{\Delta F}$ at each voltage
27. and error bars represent the standard error of the mean. Black line is a first-order
28. Boltzmann equation fit to the $k_{\Delta F}$–voltage relation: $V_{1/2}$ = +38 ± 15 mV, z = 1.43 ± 0.35
29. $e_\circ$. Green line is prediction from the EVAP model at 9 nM GxTX. Error bars are standard
30. error of the mean.
31.
32.

Figure 8: ***Relationship of GxTX-594 labeling to probability of Kv2 voltage sensor activation.***

A. EVAP model predictions of voltage-dependence of cell surface fluorescence intensity at

different concentrations of EVAP in solution. Bottom axis represents membrane voltage.

The left axis represents the predicted fluorescence relative to the situation where all

voltage sensors are at rest and does not include EVAP signal that is insensitive to voltage.

The right axis represents the probability that voltage sensors of unlabeled Kv2.1 are in

their resting conformation.

B. EVAP model predictions of concentration- and voltage-dependence of $k_{\Delta F}$. Colors

correspond to panel A. The right axis represents the probability that voltage sensors of

labeled Kv2.1 are in their active conformation.

Figure 9: ***High frequency repetitive stimuli amplify the GxTX-594 response.***

A. Fluorescence intensity from Kv2.1- CHO cells incubated in 9 nM GxTX-594. Arrow

indicates voltage-clamped cell. Fluorescence at holding potential of -80 mV (Left), after

50 s of 200 Hz stimulus (Middle), and 100 s after the cell is returned to holding potential

of -80 mV (Right). Stimulus was a 2 ms step to +40 mV. Note that in each panel the

unclamped cell (Left cell in each panel) does not show a change in fluorescence. Scale

bar is 10 μm.

B. Representative trace with GxTX-594 unlabeling at 200 Hz. The trace was fit using a

monoexponential (Eq. 1). Fit is overlaid and is in red. The $k_{\Delta F}$ at 200 Hz was obtained

from this fit. For 200 Hz $k_{\Delta F} = 0.2327 \pm 0.0099 \times 10^{-1} \text{ s}^{-1}$. 0% F/F_{init} was set by

subtraction of the average intensity of a region that did not contain cells. 100% was set by

the initial fluorescence intensity at -80 mV.

C. Fluorescence–stimulus frequency relation. Points indicate F/F_{init} from individual cells.

(Top panel) point coloring indicates data from the same cell. (Bottom panel) black bars

represent the average F/F_{init} at each voltage and error bars represent the standard error of

the mean. Green line is the prediction of the EVAP model at a concentration of 9 nM.

D. $k_{\Delta F}$–stimulus frequency relation. Plotted as in panel C.

1	Figure 10: *GxTX-594 selectively labels Kv2 proteins.*
2	A. Fluorescence from live CHO cells transfected with Kv2.1-GFP, Kv2.2-GFP, Kv4.2-
3	GFP + KChIP2, Kv1.5-GFP + Kvβ2, or BK-GFP (Indicated by row) and labeled with
4	GxTX-594. Airy disk imaging was from a plane above the glass-adhered surface.
5	Cells were incubated with 100 nM GxTX-594 and 5 μg/mL WGA-405 then rinsed
6	before imaging. Fluorescence corresponds to emission of GFP (Column 1), GxTX-
7	594 (Column 2), WGA-405 (Column 3), or an overlay (Column 4). Scale bars are 20
8	μm.
9	B. Intensity of GxTX-594 labeling for different K^+ channel-GFP types. Fluorescence
10	intensity resulting from 594 nm excitation of GxTX-594 is divided by fluorescence
11	intensity resulting from 488 nm excitation of GFP. This value was normalized to the
12	average 594:488 ratio from GxTX-594 and Kv2.1-GFP. Circles indicate
13	measurements from individual cells. Only cells with obvious GFP expression were
14	analyzed. For analysis ROIs were drawn around the cell membrane indicated by
15	WGA-405 fluorescence. Pixel intensities were background-subtracted before analyses
16	by subtracting the average fluorescence of a background ROI that did not contain
17	cells from the ROI containing the cell; this occasionally resulted in ROIs with
18	negative intensity. Kv2.1 n = 16, Kv2.2 n = 10, Kv4.2 n = 13, Kv1.5 n = 13, BK n
19	=10; n indicates the number of individual cells analyzed during a single application of
20	GxTX-594. Bars are arithmetic mean. Significant differences were observed between
21	594:488 ratio for Kv2.1 or Kv2.2 and Kv1.5, Kv4.2, or BK by Mann-Whitney
22	($p<0.0001$).
23	C. Pearson correlation coefficients between GxTX-594 and GFP. Same cells as panel B.
24	Significant differences were observed between correlation coefficients for Kv2.1 or
25	Kv2.2 and Kv1.5, Kv4.2, or BK by Mann-Whitney ($p<0.0001$).
26	
27	

Figure 11: ***GxTX-594 labels CA1 hippocampal pyramidal neurons transfected with Kv2.1-***

GFP.

A. Two-photon excitation images of fluorescence from the soma and proximal dendrites of a

rat CA1 hippocampal pyramidal neuron in a brain slice six days after transfection with

Kv2.1-GFP (Left), labeled with 100 nM GxTX-594 (Middle) and overlay (Right). Image

represents a Z-projection of 20 optical sections. Scale bar 10 μm.

B. A single optical section of the two-photon excitation image shown in A. GxTX-594

labels both Kv2.1-GFP puncta from a transfected cell and apparent endogenous Kv2

proteins from an untransfected cell in the same cultured slice (Right, arrow).

C. Pearson correlation coefficients from CA1 hippocampal neurons two, four, and six days

after transfection with Kv2.1-GFP. Each circle represents a different neuron. Bars are

arithmetic means.

Figure 12: ***GxTX-594 puncta on hippocampal CA1 neurons are sensitive to voltage stimulus.***

    A. Two-photon excitation optical section from CA1 pyramidal neurons in a cultured hippocampal slice after incubation with 100 nM GxTX-594. Scale bar is 5 μm.

    B. 594 fluorescence from a single two-photon excitation optical section before and during depolarization of a whole cell patch-clamped neuron. Text labels of holding potential (-70 mV or 0 mV) indicate approximate position of the patch-clamp pipette. Red arrows indicate stimulus-sensitive puncta, white arrows indicate stimulus-insensitive puncta. Left panel is the average fluorescence of the three frames at -70 mV before depolarization while the right panel is the average fluorescence of three frames after holding potential was stepped to 0 mV. Scale bar is 5 μm. Supplementary Movie 1 contains time-lapse images from this experiment.

    C. ROIs used in analysis for panels D, E, and F. Same slice as panel B. ROI 1 contains the apparent plasma membrane of the cell body of the patch-clamped neuron, it was generated by drawing a path tracing the apparent plasma membrane and then expanding to an ROI containing 15 pixels on either side of the path (1.2 μm total width). ROI 2 contains the area 15-45 pixels outside the membrane path (1.2 μm total width). RO1 3 contains the area more than 45 pixels outside of the membrane path. ROI 0 contains the area more than 15 pixels inside the membrane path. Scale bar is 5 μm.

    D. Fluorescence from each ROI shown in C. Squares represent ROI 0, circles represent ROI 1, upward triangles represent ROI 2 and upside-down triangles represent ROI 3. Background was defined as the mean fluorescence of ROI 0 during the experiment. F_{init} was defined as the mean fluorescence of ROI 1 during the first six frames, after subtraction of background. Dotted lines represent the average fluorescence of the first six frames of each ROI. The voltage protocol is shown above.

    E. Change in fluorescence during a 0 mV step for different ROIs in multiple hippocampal slices. ROIs for each slice were defined by methods described in panel C. Circles represent mean fluorescence from six frames during 0 mV stimulus (F), normalized to mean fluorescence from same ROI in six frames before stimulus (F_{init}). Circle color is consistent between ROIs for each hippocampal slice, red circles are from the slice shown in panel D. Black bars are arithmetic mean ± standard error from 3 hippocampal slices.

F. Kinetics of fluorescence change from ROI 1 during a 0 mV step in multiple hippocampal

slices. ROI 1 for each slice was defined by the method described in panel C. Lines are

monoexponential fits (Eq.1). $k_{\Delta F}$ = 6.8 x 10^{-2} ± 2.6 x 10^{-2} s^{-1} (yellow), 6.8 x 10^{-2} ± 4.6 x

10^{-2} s^{-1} (red), and 4.9 x 10^{-2} ± 2.1 x 10^{-2} s^{-1} (green). The voltage protocol is shown above.

Colors of individual circles indicate the same slices as panel E.

G. Region of interest used in analysis for panels H, I, and J. Same image as panel C. The

shaded ROI was generated by drawing a path tracing the apparent plasma membrane and

then expanding to an ROI containing 5 pixels on either side of the path (0.4 μm total

width). Numbers indicate puncta that appear as peaks in panel H. Scale bar is 5 μm.

H. A plot of the fluorescence intensity along the ROI shown in panel G before (red trace)

and during (black trace) 0 mV stimulus. Numbers above peaks correspond to puncta

labeled in panel G. Red trace: mean fluorescence intensity during the three frames

immediately before the stimulus, normalized to mean intensity of entire ROI (black

dotted line), plotted against distance along path. Black trace: mean fluorescence intensity

during three frames at 0 mV, normalized by the same F_{init} value as the red trace.

I. Kinetics of fluorescence change of individual puncta from panel G. Puncta intensity are

average intensity of points extending to half maxima of each peak in panel H. Asterisks

indicate mean fluorescence intensity of puncta 1 (pink), 4 (red), and 5 (dark red). Lines

are monoexponential fits (Eq.1). $k_{\Delta F}$ = 7.1 x 10^{-2} ± 2.9 x 10^{-2} s^{-1} (pink), 6.7 x 10^{-2} ± 2.5 x

10^{-2} s^{-1} (red) and 1.2 x 10^{-1} ± 5.6 x 10^{-2} s^{-1} (dark red). Fits to other puncta had standard

deviations larger than $k_{\Delta F}$ values, and were excluded. F_{init} was defined as the mean

background subtracted fluorescence of the puncta during the six frames before stimuli.

The voltage protocol is displayed above. Blue line is prediction from Scheme I.

J. Comparison of fluorescence change of puncta and inter-puncta regions in response to 0

mV stimulus. The regions before stimulus shown in the red line of panel H that had

F/F_{init} ≥ 1 (above average) were binned separately from regions with F/F_{init} < 1. The

mean fluorescence of each region during 0 mV stimulus (panel H black trace) was

compared to the fluorescence before stimulus (panel H red trace). Circles indicate values

from three independent hippocampal slices, colors indicate same slices as panel E and F.

Black bars are arithmetic mean ± standard error.

K. Rate of fluorescence change of GxTX-594 after a 0 mV stimulus; from puncta (as in panel I), ROI 1 (as in panel F) or Kv2.1-CHO cells in 100 nM GxTX-594. Kv2.1-CHO cells were imaged at the same temperature as neurons (30 °C) using neuronal intracellular solution. CHO extracellular solution (CE) was used for Kv2.1-CHO cell experiments. Kv2.1-CHO measurements were made by airy disk confocal imaging. Black bars are mean ± SEM from data shown. Dashed blue line is $k_{\Delta F} = 6.31 \times 10^{-2}$ s^{-1} prediction of Scheme I.

Figure Supplement 1: ***GxTX-594 selectively labels Kv2 proteins on cell surfaces***

A. Exemplar whole-cell voltage clamp recordings of CHO cells expressing Kv2.1-GFP,

Kv2.2-GFP, Kv4.2-GFP, Kv1.5-GFP, or BK-GFP. Recordings shown are representative

responses to 100 ms steps from -100 mV to -40, 0, and +40 mV.

B. Confocal imaging of fluorescence from live CHO cells transfected with Kv2.1-GFP,

Kv2.2-GFP, Kv4.2-GFP Kv1.5-GFP, or BK-GFP (Indicated by row) and labeled with

GxTX-594. Confocal imaging plane was above the glass-adhered surface. Cells were

incubated with 100 nM GxTX-594 and 5 μg/mL WGA-405 and rinsed before imaging.

Fluorescence shown corresponds to emission of GFP (Column 1), Alexa Fluor 594

(Column 2), WGA-405 (Column 3), or an overlay of GFP, Alexa Fluor 594, and WGA-

405 (Column 4). Scale bars are 20 μm.

C. Fluorescence intensity resulting from 594 nm excitation of GxTX-594 is divided by

fluorescence intensity resulting from 488 nm excitation of GFP. Analysis methods as in

Figure 10 B. Kv2.1 n = 11, Kv2.2 n = 9, Kv4.2 n = 13, Kv1.5 = 13, and BK n =12. Cells

were compiled from two independent GxTX-594 applications, each circle corresponds to

a cell. Significant differences were observed between GxTX:GFP ratio for Kv2.1 or

Kv2.2 and Kv1.5, Kv4.2, or BK by Mann-Whitney ($p<0.0001$).

D. Pearson correlation coefficients between GxTX-594 and GFP. Same cells as panel C.

Analysis methods as in Figure 10 C.

Figure Supplement 2: *GxTX-594 fluorescence equilibration after dilution from 100 nM to*

9nM.

A. CHO cells transfected Kv2.1-GFP (Left) with in 100 nM GxTX-594 (Middle) and after

dilution to 9 nM GxTX-594 (Right). Regions darker than the background bath solution

are CHO cells not transfected with Kv2.1-GFP (Arrowhead). Scale bar is 20 μm.

B. Fluorescence excited at 594 nM during dilution from 100 nM GxTX-594 to 9 nM GxTX-

594. Timing of dilution is shown above the graph. Fluorescence was normalized to the

mean 594 fluorescence before dilution to 9 nM. Data are from cell labeled with arrow in

panel A. 100 nM and 9 nM images in panel A are from 0 and 12 minutes respectively.

C. Rate of GxTX-594 fluorescence decay after dilution from monoexponential fits (Eq 1).

Error bars are standard deviation of fit parameter. n = 5 cells.

Figure Supplement 3: *Variation in bath temperature does not account for variability of GxTX-*

594 kinetics.

A. Representative traces of GxTX-594 fluorescence intensity response to voltage changes at

27 °C (black) and 37 °C (gray). Smooth lines are fits of a monoexponential function (Eq.

1): 27 °C $k_{\Delta F}$ = 2.47 x 10^{-2} ± 0.39 x 10^{-2} s^{-1}; 37 °C $k_{\Delta F}$ = 7.54 x 10^{-2} ± 0.54 x 10^{-2} s^{-1}.

Background was determined by taking the average fluorescence of a region that did not

contain cells over the time course of the voltage protocol. This average was subtracted

from each experimental group. Traces were normalized to initial fluorescence intensity

before the application of the voltage stimulus.

B. $k_{\Delta F}$ at 27 °C and 37 °C. The rate of fluorescence change was significantly faster at higher

temperatures (Mann-Whitney p = 0.0005). From geometric means (Bars), a Q_{10} of 3.8

was calculated between 27 °C and 37 °C. Each circle represents one cell, n = 7 both

groups.

Figure Supplement 4: *GxTX-594 labels CA1 hippocampal pyramidal neurons transfected with*

Kv2.1-GFP.

Two-photon excitation images of rat CA1 hippocampal pyramidal neurons in brain slices as in

Figure 7 A. Kv2.1-GFP (Left), GxTX-594 (Middle) and overlay (Right). Scale bars 10 µm.

A. Pyramidal neurons two days after transfection with Kv2.1-GFP.

B. Pyramidal neurons four days after transfection with Kv2.1-GFP.

C. Pyramidal neurons six days after transfection with Kv2.1-GFP.

Table 1

Parameters used for calculations to generate Scheme I: EVAP Allosteric Expansion.

Parameter	Value
$k_{on,resting}$	$0.30\ \mu M^{-1}\ s^{-1}$
$k_{off,resting}$	$0.0081\ s^{-1}$
$k_{on,activated}$	$0.21\ \mu M^{-1}s^{-1}$
$k_{off,activated}$	$0.39\ s^{-1}$
$V_{1/2,unlabeled}$	$-32\ mV$
$V_{1/2,labeled}$	$41\ mV$
z	$1.5\ e_0$

Figure 1

Figure 2

A GxTX-594

0nM   1nM   10nM   100nM   1000nM

B

1nM 10 100 1000

100%

F / Fmax

0%

0 20 40 60 80 100
Time / min

C

100%

F / Fmax

0%

1 10 100 1000
GxTX-594 / nM

D

0.1

$k_{\Delta F}$ / s^{-1}

0.001

0 1 10 100
GxTX-594 / nM

Figure 3

Figure 4

A

B

Figure 5

Figure 6

Figure 7

Figure 8

A

B

Figure 9

Figure 10

Figure 11

Figure 12

Supplement 1

A

2 nA
50 ms

B

Kv 2.1 | GxTX-594 | Membrane | Overlay

Kv 2.2

Kv 4.2

Kv 1.5

BK

C

594:488 fluorescence

Kv2.1 Kv2.2 Kv4.2 Kv1.5 BK

D

Correlation coefficient

Kv2.1 Kv2.2 Kv4.2 Kv1.5 BK

Supplement 2

A

| Kv2.1-GFP | 100 nM GxTX-594 | 9 nM GxTX-594 |

B

100 nM ⎯⎯⎯⎯⎯⎯⎯⎯⎯ 9 nM

F / F_{init}

Time / min

C

$k_{\Delta F} / s^{-1}$

Supplement 3

Supplement 4