Optical measurement of endogenous ion channel activation in live tissue

Authors

Parashar Thapa[1‡], Robert Stewart[1‡], Rebecka J. Sepela[1], Oscar Vivas[2], Laxmi K. Parajuli[2], Mark Lillya[1], Sebastian Fletcher-Taylor[1,3], Bruce E. Cohen[3], Karen Zito[2], Jon T. Sack[1*]

[1]Department of Physiology and Membrane Biology, University of California, Davis, CA 95616
[2]Center for Neuroscience, University of California, Davis, CA 95616
[3]Molecular Foundry and Division of Molecular Biophysics & Integrated Bioimaging, Lawrence Berkeley National Laboratory, Berkeley, CA 94720

[‡]Authors contributed equally to this work
[*]Correspondence to jsack@ucdavis.edu

Abstract

A primary goal of molecular physiology is to understand how conformational changes of

proteins affect the function of cells, tissues, and organisms. Here, we describe an imaging

method for measuring the conformational changes of the voltage sensors of endogenous ion

channel proteins within live tissue. We have synthesized a fluorescent variant of a peptidyl toxin,

compatible with two-photon excitation microscopy, that targets a resting conformation of Kv2-

type voltage-gated potassium channels. Using voltage-response fluorescence, we calibrated a

statistical thermodynamic model relating probe labeling intensity to the conformations adopted

by unlabeled Kv2 proteins. Two-photon excitation imaging of rat brain slices reveals

fluorescence consistent with conformation-selective labeling of endogenous neuronal Kv2

proteins. This method of quantifying endogenous Kv2 conformational changes from

fluorescence images enables the imaging of specific protein responses to electrical activity in

tissue, without genetic modification.

Introduction

1. Voltage-gated ion channels respond to changes in voltage across a cell membrane with

2. conformational changes that initiate physiological responses. Structural biology techniques have

3. revolutionized our understanding by revealing atomistic details of these protein conformations

4. (Y. Jiang *et al.* 2003, Q.-X. Jiang *et al.* 2004, Long *et al.* 2005, Tao *et al.* 2010, Kintzer *et al.*

5. 2018, Wisedchaisri *et al.* 2019, Xu *et al.* 2019), yet the physiological relevance of each

6. conformation can be difficult to determine . Here, we describe an imaging method to detect

7. conformational changes of voltage sensors of endogenous Kv2-type voltage-gated K^+ channels

8. in tissue. Our objective in developing Kv2 imaging probes is to reveal conformational changes of

9. native proteins in live cells, with subcellular spatial resolution, which cannot be otherwise

10. measured. Conformational changes of voltage sensors have been detected with

11. electrophysiological measurements of gating currents (Armstrong & Bezanilla 1973, M. F.

12. Schneider & Chandler 1973, Bezanilla 2018) or by optical measurements from fluorophores

13. covalently attached to voltage sensors by genetic encoding (Lin & Schnitzer 2016) and/or by

14. chemical engineering (G. Zhang *et al.* 2015). Each of these techniques have enabled

15. breakthroughs in our understanding of how protein conformational changes are coupled to one

16. another (Zheng & Trudeau 2016). Experimental limitations prevent these existing techniques

17. from measuring conformational changes of voltage sensors of most endogenous proteins: gating

18. currents can only be measured when the proteins are expressed at high density in a voltage-

19. clamped membrane; engineered proteins differ from endogenous channels; and introducing point

20. mutations and conjugated fluorophores into voltage sensors alters structure and function. Here,

21. we develop a different strategy, where fluorescent probes dynamically label endogenous voltage

22. sensors to reveal conformational states.

23.

24. Members of the voltage-gated cation channel superfamily, including Kv2 channels, contain

25. specialized voltage sensor domains comprised of a bundle of four transmembrane helices termed

26. S1-S4 (Long *et al.* 2005, 2007). The S4 helix contains positively charged arginine and lysine

27. residues, gating charges, which respond to voltage changes by moving through the

28. transmembrane electric field (Aggarwal & MacKinnon 1996, Seoh *et al.* 1996, Islas & Sigworth

29. 1999). When voltage sensor domains encounter a transmembrane voltage that is more negative

30. on the inside of the cell membrane, voltage sensors are biased towards resting conformations, or

down states, in which gating charges are localized intracellularly. When voltage becomes more

positive, gating charges translate towards the extracellular side of the membrane, and voltage

sensors are progressively biased towards activated conformations, or up states, in a process of

voltage activation (Armstrong & Bezanilla 1973, Zagotta *et al.* 1994, Tao *et al.* 2010, Xu *et al.*

2019). While activation of Kv2 voltage sensor domains does not guarantee pore opening (Scholle

et al. 2004, Jara-Oseguera *et al.* 2011, Fox *et al.* 2013, Tilley *et al.* 2019), activation of voltage

sensors are required for pore opening (Islas & Sigworth 1999). Thus, measurement of whether

Kv2 voltage sensors adopt resting *versus* activated conformations determines whether a

prerequisite for opening has occurred.

To image native channel activation, we exploit the conformation-selective binding of the

tarantula peptide guangxitoxin-1E (GxTX) to measure where and when Kv2 proteins adopt

resting *versus* activated conformations in tissue. To do this, we synthesized a fluorescent

derivative of GxTX that is compatible with two-photon excitation imaging through light-

scattering tissue, GxTX-594. Heterologously overexpressed Kv2 channels in dissociated cells

have been dynamically labeled by a fluorescent GxTX conjugate, and at dilute concentrations,

the probe can report conformational changes while inhibiting only a small fraction of the

macroscopic K^+ current (Tilley et al., 2014). Here, we synthesize GxTX-594, find that it

specifically labels native Kv2 channels brain slices, and identify the conformational change that

it reports. Crucially, we also develop a method to quantify the degree of Kv2 voltage sensor

conformational change from images of GxTX-594 fluorescence. We calibrate this method to

determine the conformational equilibria of unlabeled channels using fluorescence from channels

labeled with GxTX-594. We deploy the GxTX-594 probe in brain slices and image voltage-

sensitive fluorescence changes that indicate conformational changes of endogenous neuronal

Kv2 channels.

Results

To create an optical probes of native ion channel activity in tissue, we used solid-phase peptide synthesis to generate a variant of GxTX, an amphiphilic 36-amino acid cysteine knot peptide. GxTX partitions into the outer leaflet of the plasma membrane bilayer where it selectively binds to membrane-embedded voltage sensors of Kv2 channels (Milescu *et al.* 2009, Tilley *et al.* 2014, Gupta *et al.* 2015). We synthesized Ser13Cys GxTX, where the free "spinster" thiolate side chain (Shimony *et al.* 1994) of cysteine 13 is predicted to extend into extracellular solution when the peptide is bound to a voltage sensor. GxTX-1E folds by formation of 3 internal disulfides, and cysteine 13 was differentially protected during oxidative refolding to direct chemoselective conjugation (Tilley *et al.* 2014). Following refolding and thiol deprotection, the maleimide derivative of Alexa Fluor 594, a persulfonated Texas Red analog, was condensed with the free thiol, and Ser13Cys(Alexa Fluor 594) GxTX-1E (called "GxTX-594" throughout) was purified and tested for activity against Kv2.1 (Fig. 1 and Fig 1 Supplement 1). These results demonstrate that variants of GxTX can be folded and chemoselectively ligated to reporter molecules while retaining bioactivity against ion channels. Alexa Fluor 594 exhibits a large two-photon absorption cross-section and ample spectral separation from green fluorescent protein (GFP), making it well-suited for multiplexed, two-photon excitation imaging experiments (Zito *et al.* 2004).

GxTX-594 colocalizes with Kv2.1 and Kv2.2

We first assessed whether GxTX-594 labels Kv2 channels. Mammals have two Kv2 pore-forming subunits, Kv2.1 and Kv2.2, both of which are expressed in neurons throughout the brain (Trimmer 1991, Kihira *et al.* 2010, Bishop *et al.* 2015). CHO-K1 cells were transfected with either a rat Kv2.1-GFP or Kv2.2-GFP construct. While the CHO-K1 cell line has little, if any, endogenous K^+ current (Gamper *et al.* 2005), transfection with either Kv2.1-GFP or Kv2.2-GFP yielded delayed rectifier K^+ currents (Fig 6 Supplement 1 A, top two rows). When imaged two days post transfection, Kv2.1-GFP and Kv2.2-GFP fluorescence appeared at the surface membrane of the transfected CHO cells, and the fluorescence was localized predominantly to the glass-adhered basal surface, where it adopted a clustered morphology (Fig 1 A). Clustered basal expression is a hallmark of Kv2 channel localization in neurons and other mammalian cell lines when they are adhered to glass (Shi *et al.* 1994, Antonucci *et al.* 2001). After incubation with

membrane-impermeant GxTX-594 (100 nM for 5 minutes), GxTX-594 fluorescence localized to

regions with Kv2-GFP signal (Fig 1 A). GxTX-594, and Kv2.1-GFP or Kv2.2-GFP, had Pearson

correlation coefficients that averaged 0.86 or 0.80, respectively, verifying a high degree of

colocalization of both Kv2 proteins with GxTX-594 (Fig 1 B). The ratio of GxTX-594 to GFP

fluorescence intensity was similar for Kv2.1-GFP or Kv2.2-GFP (Fig 1 C), consistent with the

lack of discrimination of GxTX between Kv2.1 and Kv2.2 (Herrington *et al.* 2006).

GxTX-594 binds Kv2.1 in the presence of its neuronal auxiliary subunit AMIGO-1

In neurons, Kv2 channels colocalize with an auxiliary subunit, AMIGO-1 (Peltola *et al.* 2011,

Bishop *et al.* 2018). AMIGO-1 is a single-pass transmembrane protein with a large extracellular

domain, an architecture similar to auxiliary subunits of voltage-gated sodium channels, Kv4

channels, or BK channels that modulate binding of peptide toxins to the channels (Xia *et al.*

1999, Gilchrist *et al.* 2013, Maffie *et al.* 2013). To determine if AMIGO-1 affects GxTX-594

labeling, Kv2.1-expressing cells were transiently transfected with an AMIGO-1-YFP construct.

For these experiments, Kv2.1 channels were expressed in a CHO-K1 cell line stably transfected

with rat Kv2.1 under control of a tetracycline-inducible promoter (Kv2.1-CHO) (Trapani & Korn

2003). Kv2.1 expression was induced with minocycline 48 hours prior to experiments, such that

all Kv2.1-CHO cells expressed Kv2.1. In this expression system, Kv2.1 channels are not tagged

with a fluorescent protein. AMIGO-1-YFP fluorescence was observed in clusters at the basal

membrane (Fig 2 A) similar to Kv2.1-GFP (Fig 1 A). GxTX-594 colocalized with AMIGO-1-

YFP (Fig 2 A). Pearson's correlation coefficient indicated colocalization between AMIGO-1-

YFP and GxTX-594, consistent with AMIGO-1 colocalizing with clustered Kv2.1 (Fig 2 B).

GxTX-594 also labeled cells that did not express AMIGO-1 (Fig 2 A). We tested whether

binding of GxTX-594 to Kv2.1 channels is different between cells which do or do not express

AMIGO-1. Increasing concentrations of GxTX-594 were applied to cells and the relationship

between GxTX-594 fluorescence intensity and dose was quantified for cells expressing AMIGO-

1-YFP (green) and cells not expressing AMIGO-1-YFP (blue) (Fig 2 C). A Langmuir binding

isotherm was fit to GxTX-594 fluorescence intensity (lines, Fig 2 D), and resulted in an

indistinguishable dissociation constant between cells expressing AMIGO-1-YFP (27 ± 14 nM)

and cells not expressing AMIGO-1-YFP (26.9 ± 8.3 nM). Thus, there is no indication that

AMIGO-1 disrupts the GxTX-594–Kv2 interaction.

GxTX-594 fluorescence dynamics report Kv2 channel voltage activation

To understand the relationship between channel gating and GxTX-594 fluorescence, we

determined how fluorescence intensity on cells expressing Kv2.1 responds to changes in

membrane voltage. For these experiments, Kv2.1-CHO cells were incubated in minocycline for

48 hours to induce Kv2.1 channel expression. Kv2.1-CHO cells were then incubated for 5

minutes in a bath solution containing 100 nM GxTX-594 before dilution with extracellular

solution to 9 nM, and imaged at least 9 minutes after dilution. Fluorescence intensity images

were generated from airy disk confocal imaging of cells voltage-clamped in the whole-cell patch

clamp configuration. Unlike images shown in Fig 1, the imaging plane for voltage-clamp

experiments was above the glass-adhered surface. This higher plane was chosen because GxTX-

594 labeling kinetics are slower in the highly restricted extracellular space where the plasma

membrane adheres to glass (Fig 3 Supplement 1). To measure voltage-dependent fluorescence

changes, cells were held at -80 mV and stepped to indicated test potentials (Fig 3 A). A region of

interest (ROI) corresponding to the cell surface was manually identified and average

fluorescence intensity quantified from time-lapse sequences. Average fluorescence intensity

from the cell surface was background-subtracted using the average fluorescence intensity from

an ROI in a region without cells. Both the amplitude and kinetics of fluorescence change from

cell surface ROIs are sensitive to voltage (Fig 3 B), similar to prior findings with GxTX-550

(Tilley *et al.* 2014). We observed that cells exhibited a baseline fluorescence intensity above

background, even when given a +80 mV depolarizing voltage stimulus (Fig 3 B). This residual

fluorescence varied between cells (Fig 3 C) and was potentially due to internalized GxTX-594

and/or autofluorescence. However, much of the residual fluorescence appeared to be localized to

the cell surface membrane (Fig 3 A). As such membrane labeling was not present on CHO cells

in the absence of Kv2 channels (Fig 6), this fluorescence could indicate a population of Kv2.1

channels that reside at the cell surface and bind GxTX-594 in a fashion that renders the Kv2.1–

GxTX complex insensitive to voltage.

To compare labeling intensity between cells, the initial fluorescence at a holding potential of -80

mV was normalized to 100% F/F_{init}, and residual fluorescence after a +80 mV step was

normalized to 0% F/F_{init} (Fig 3 D). The fluorescence–voltage response was fit with a Boltzmann

function with a half maximal voltage midpoint ($V_{1/2}$) of -27 mV and a steepness (z) of 1.4

elementary charges (e_0) (Fig 3 D, black line). This is strikingly similar to the voltage dependence

of integrated gating charge from Kv2.1-CHO cells, without any GxTX present $V_{1/2}$ = -26 mV, z =

1.6 e_0 (Tilley *et al.* 2019).

To determine the temporal response of GxTX-594 labeling to voltage change, we compared rates

of fluorescence change ($k_{\Delta F}$) to varying step potentials (Fig 3 E) in the same data set that were

analyzed in Fig 3 D. We quantified rates of fluorescence change ($k_{\Delta F}$) by fitting a

monoexponential function:

$F = F_\infty + F_0 e^{\frac{(t-t_0)}{\tau}}$ (Equation 1)

Where $\tau = 1/k_{\Delta F}$, t = time, t_0 = time at start of fit, F = fluorescence intensity, F_0 = fluorescence at

start of fit, F_∞ = fluorescence after infinite time. In response to voltage steps from a holding

potential of -80 mV to more positive potentials, $k_{\Delta F}$ increased progressively from approximately

-40 mV to +40 mV. After a return to -80 mV, labeling occurred at a rate similar to unlabeling

at -40 mV, indicating that the rate of relaxation to a labeled/unlabeled equilibrium saturates

below -40 mV. There was substantial variability in $k_{\Delta F}$ between cells (Fig 3 E). We wondered if

this could be due to fluctuations in ambient temperature (27-29°C), given the substantial

temperature sensitivity reported for Kv2.1 conductance (F. Yang & Zheng 2014). However the

temperature dependence of $k_{\Delta F}$ indicated that only 1.3-fold variation could be attributed to a 2°C

temperature variation (Fig 3 Supplement 2), suggesting that there are determinants of $k_{\Delta F}$

variability which are intrinsic to individual cells. The $k_{\Delta F}$–voltage response was fit with a

Boltzmann distribution (Eq. 2) resulting in $V_{1/2}$ = +2 mV and z = 1.1 e_0 (Fig 3 E, black line).

Notably, the dynamic fluorescence responses of GxTX-594 to voltage changes occurred at

physiologically relevant potentials, suggesting that changes in GxTX-594 labeling intensity or

rate could occur in response to changes in cellular electrical signaling.

The relation between voltage sensor activation and GxTX-594 dynamics can be recapitulated

by rate theory modeling.

To enable translation of the intensity of fluorescence from GxTX-594 on a cell surface into a

measure of channel conformational change, we developed a model relating cell labeling to

voltage sensor activation. The framework of this model is generalizable to fluorescent molecular

probes that report conformational changes by a change in binding affinity. To refer to GxTX-594

and mechanistically related probes more generically, we refer to them as Endogenous Channel

Activity Probes (ECAPs).

In our model of ECAP labeling, the ratio of labeled to unlabeled Kv2 in a membrane is

determined by confomational change from resting to activated altering the binding and unbinding

rates of the ECAP (Scheme I).

Scheme I

Scheme II

The Scheme I model assumes the innate voltage sensitivity of the Kv2 channel is solely

responsible for controlling voltage dependence. Labeling is voltage-dependent because the

binding and unbinding rates are different for resting and activated conformations of voltage

sensors. When voltage sensors change from resting to activated conformations, the binding rate

of the ECAP decreases and the unbinding rate increases. When the membrane voltage is held

constant for sufficient time, the proportions of labeled and unlabeled channels reach an

equilibrium. Scheme I assumes that voltage sensor conformational changes equilibrate much

faster than ECAP binding or unbinding, such that binding and unbinding are the rate-limiting

steps. ECAP labeling requires seconds to equilibrate (Fig 3), whereas Kv2 channel gating

equilibrates in milliseconds (Tilley *et al.* 2019), 3 orders of magnitude more quickly. This much

faster voltage sensor equilibration justifies our simplifying approximation that voltage sensor

conformations achieve equilibrium so quickly that only their distribution at equilibrium impacts

the kinetics of labeling and unlabeling. Voltage-activation of Kv2 channels involve many

conformational changes (Scholle *et al.* 2004, Jara-Oseguera *et al.* 2011, Tilley *et al.* 2019).

However, models which presume independent activation of a voltage sensor in each of the four

Kv2.1 subunits accurately predict many aspects of voltage activation and voltage sensor toxin

binding (Lee *et al.* 2003, Tilley *et al.* 2019). Thus, we found it reasonable to model voltage

sensor activation of Kv2 channels as a single, rapid transition of each voltage sensor from a

resting conformation to an activated conformation. With this model, at any given voltage, there

is a probability that a voltage sensor is either in its resting conformation ($P_{resting}$), or in its

activated conformation ($P_{activated}$), such that $P_{activated} = (1 - P_{resting})$. The equilibrium for voltage

sensor activation is then a ratio of activated to resting voltage sensors ($P_{activated}/P_{resting}$) in which

$$\frac{P_{activated}}{P_{resting}} unlabeled = e^{(V-V_{1/2,unlabeled})\frac{zF}{RT}} \qquad \text{Equation 2a}$$

$$\frac{P_{activated}}{P_{resting}} labeled = e^{(V-V_{1/2,labeled})\frac{zF}{RT}} \qquad \text{Equation 2b}$$

where $V_{1/2}$ is the voltage where $P_{activated}/P_{resting,} = 1$, z is the number of elementary charges

determining voltage dependence, F is the Faraday constant, R is the ideal gas constant, and T is

absolute temperature. In a prior study, we fit the conductance–voltage relation of Kv2.1 with a

Boltzmann distribution that yielded $V_{1/2}$ = -31.7 mV with $z = 1.5$ e_0 for the early movement of 4

independent voltage sensors, and found that with a saturating concentration of GxTX the $V_{1/2}$ =

41.3 mV (Tilley *et al.* 2019, Fig. 1 C). These values were used for $V_{1/2,unlabled}$, *z, and* $V_{1/2,labled}$

respectively (Table 1). To relate voltage sensor activation to labeling and unlabeling, we used

microscopic binding ($k_{on}[ECAP]$) and unbinding (k_{off}) rates that are distinct for resting and

activated voltage sensors. We estimated values for these rates assuming

$k_{\Delta F} = k_{on}[ECAP] + k_{off}$ (Equation 3)

and

$K_d = \dfrac{k_{off}}{k_{on}}$ (Equation 4)

To calculate the $k_{on,resting}$ and $k_{off,resting}$ values reported in Table 1, we used the saturating value at

negative voltages of the $k_{\Delta F}$–voltage relation (Fig 3 E), and K_d from dose-response imaging (Fig

2 D). In 9 nM GxTX-594, at greater than +40 mV, voltage-dependent unlabeling was nearly

complete, indicating that $k_{off,activated} \gg k_{on,activated}[ECAP]$. We input the saturation of $k_{\Delta F}$ at

positive voltages as $k_{off,activated}$ (Fig 3 E). The slow labeling of activated voltage sensors

confounded attempts to measure $k_{on,activated}$ directly, and we used the statistical thermodynamic

principle of microscopic reversibility (Lewis 1925) to constrain $k_{on,activated}$:

$\dfrac{\frac{P_{activated}}{P_{resting}},labeled}{\frac{P_{activated}}{P_{resting}},unlabeled} = \dfrac{\frac{k_{off,resting}}{k_{on,resting}}}{\frac{k_{off,activated}}{k_{on,activated}}}$ (Equation 5)

At any static voltage, the simplifying assumption that voltage sensors are in continuous

equilibrium allows Scheme I to collapse into Scheme II, which has only a single microscopic

binding rate, $k_{on,total}[ECAP]$, and unbinding rate, $k_{off,total}$. In Scheme II, $k_{on,total}$ is a weighted sum

of both $k_{on,resting}$ and $k_{on,activated}$ from Scheme I. The weights for $k_{on,total}$ are the relative

probabilities that unlabeled voltage sensors are resting or activated, which is determined at any

static voltage by an equilibrium constant, $\frac{P_{activated}}{P_{resting}} unlabeled$:

$k_{on,total} = k_{on,resting}[ECAP] \cdot \dfrac{1}{1+\frac{P_{activated}}{P_{resting}}unlabeled} + k_{on,active}[ECAP] \cdot \dfrac{1}{1+\frac{1}{\frac{P_{activated}}{P_{resting}}unlabeled}}$

(Equation 6)

Similarly, $k_{off,total}$ is determined by the unbinding rate from resting voltage sensors ($k_{off,resting}$) and

the unbinding rate from activated voltage sensors ($k_{off,activated}$), and weighted such that:

$$k_{off,total} = k_{off,resting} \cdot \frac{1}{1+\frac{P_{activated}}{P_{resting}}labeled} + k_{off,active} \cdot \frac{1}{1+\frac{1}{\frac{P_{activated}}{P_{resting}}labeled}} \qquad \text{(Equation 7)}$$

Using $k_{on,total}$ and $k_{off,total}$ we can compute $k_{\Delta F}$ using Equation 3. To test the predictive value of the Scheme II model, we compared the predicted $k_{\Delta F}$ to experimental measurements in 9 nM GxTX-594 (Fig 3 E, blue line). We found the Scheme II model reasonably reproduced the voltage dependence of Kv2.1 labeling.

The Scheme II model was also used to predict the magnitude of ECAP fluorescence changes on cell surfaces. In theory, the ratio of fluorescence at a test voltage to a prior voltage (F/F_{init}) is equal to the probability that a Kv2 subunit is labeled by the ECAP ($P_{lableled}$):

$$\frac{F}{F_{init}} = \frac{P_{labeled}}{P_{labeled,init}} \qquad \text{(Equation 8)}$$

The equilibrium $P_{labeled}$ at any voltage can be determined from microscopic binding rates associated with Scheme II where:

$$P_{labeled} = \frac{1}{1+\frac{K_{d,total}}{[ECAP]}} = \frac{1}{1+\frac{k_{off,total}}{k_{on,total}\cdot[ECAP]}} \qquad \text{(Equation 9)}$$

Using this model, we calculated the fluorescence–voltage relation from $P_{labeled}$ (Equation 8) and compared these calculated values to our experimental data. The model predicts fluorescence response within the range of the experimental data (Fig 3 D, blue line).

Rate theory modeling roughly predicts GxTX-594 labeling during dynamic voltage changes

The Scheme II model relating labeling to voltage sensor activation was developed for static voltages. However, in electrically activated cells, voltage is dynamic. In neurons, action potentials involving Kv2 currents occur on the millisecond time scale (Liu & Bean 2014, Kimm *et al.* 2015), orders of magnitude faster than the GxTX-594 response. Thus, the slow ECAP response will integrate voltage fluctuations occurring over many seconds. The Scheme II model

assumes that voltage sensors are in continuous equilibrium, and this assumption breaks down

when voltage is rapidly changing, as Kv2 voltage sensor movements with or without GxTX

require milliseconds to equilibrate (Tilley *et al.* 2019). We assessed whether the Scheme II

model retains predictive power in response action potentials by stimulating cells with brief

voltage steps that crudely mimic action potentials. We stimulated GxTX-594 labeled Kv2.1-

CHO cells with trains of 2 ms voltage steps from -80 to +40 mV, and observed a change in cell

surface fluorescence (Fig 4 A, B). To assess frequency response, step frequency was varied from

0.02 to 200 Hz. At frequencies higher than 25 Hz, detectable unlabeling occurred (Fig 4 C), and

$k_{\Delta F}$ varied with stimulus frequency (Fig 4 D). Frequency-dependent predictions of F/F_{init} and $k_{\Delta F}$

were made from the Scheme II model. These predictions were calculated assuming the summed

probability of being at every voltage determines the overall $P_{labeled}$. The general form of our

equation relating fluorescence intensity change to a stimulus that varies faster than ECAP

labeling is:

$$F/F_{init} = (P_{V1} \cdot \Delta F_{V1}) + (P_{V2} \cdot \Delta F_{V2}) + \ldots + (P_{Vn} \cdot \Delta F_{Vn}) \qquad \text{(Equation 10)}$$

where ΔF_n is ΔF at that particular voltage. For voltage steps from -80 to +40 mV Eq. 10 is:

$$F/F_{init} = (P_{40mV} \cdot \Delta F_{40mV}) + (P_{-80mV} \cdot \Delta F_{-80mV})$$

Measured changes in fluorescence intensity were 2- to 3-fold more sensitive to frequency than

responses predicted by the Eq. 10 (Fig 4 C, blue line). When the fluorescence intensity changes

were fit with a Boltzmann distribution, the fit had a similar slope but the midpoint was at a lower

stimulus frequency than the model predictions (Fig 4 C, black line). This discrepancy could be

due to Kv2.1 voltage sensors having insufficient time to equilibrate at high stimulus frequencies.

The time constant of deactivating gating current decay from Kv2.1-CHO cells was 22 ms at -80

mV (Tilley *et al.* 2019). Voltage sensors that fail to deactivate before the next stimulus could

lead to an accumulation of activated voltage sensors, and more ECAP unlabeling than predicted

by Eq. 10 which assumes continuous equilibrium.

The general form of our equation relating fluorescence intensity change to a stimulus that varies faster than ECAP labeling is:

$$k_{\Delta F} = \left(P_{V1} \cdot k_{\Delta F,V1}\right) + \left(P_{V2} \cdot k_{\Delta F,V2}\right) + \ldots + \left(P_{Vn} \cdot k_{\Delta F,Vn}\right) \qquad \text{(Equation 11)}$$

where P_{Vn} is the probability over time that the voltage has a particular value and $k_{\Delta F,n}$ is $k_{\Delta F}$ at that particular voltage. For voltage steps from -80 to +40 mV Eq. 11 is:

$$k_{\Delta F} = \left(P_{40mV} \cdot k_{\Delta F40mV}\right) + \left(P_{-80mV} \cdot k_{\Delta F-80mV}\right)$$

The model produced a frequency dependence of $k_{\Delta F}$ that was within range of the empirical data (Fig 4 D, blue line).

Overall, the response of GxTX-594 labeling to rapidly varying stimuli indicates that the fluorescence response generally follows the predictions of the Scheme II model, although GxTX-594 labeling is more sensitive to high frequency stimulation than predicted. As this very simple model recapitulated the general response of an ECAP to static and varying voltage stimuli, we further investigated its predictions of ECAP labeling responses and its utility in interpreting changes in ECAP labeling.

Changes in ECAP labeling report conformational changes of unlabeled channels.

We used the ECAP model to investigate general principles of the relation between voltage sensor activation and labeling. In particular, we are interested in exploiting ECAP labeling to reveal information about voltage sensor activation in the unlabeled channel population, akin to how Ca^{2+} indicators are used to reveal the concentration of free Ca^{2+}. The Scheme I model was used to predict labeling at a series of ECAP concentrations (Fig 5 A). While the model was calibrated to fluorescence responses at particular voltages (Fig 5 A, bottom axis), ECAP labeling is not directly sensitive to voltage, but is determined by the probability that voltage sensors are in an activated conformation (Fig 5 A, top axis). As the ECAP binds resting voltage sensors much more rapidly than activated voltage sensors ($k_{on,resting} \gg k_{on,activated}$) the probability that unlabeled

voltage sensors are resting determines the labeling rate, and influences P_{bound}. Our model predicts

that as the ECAP concentration decreased, the change in labeling ($\Delta F/\Delta F_{max}$)

asymptotically approaches the probability that unlabeled voltage sensors are resting (Fig 5 A).

Even at 100 nM GxTX-594, more than 3 times the K_d for the resting conformation, a change in

labeling corresponds to a conformational change in the voltage sensors of unlabeled channels. At

and below the K_d for the resting conformation, the change in labeling was nearly proportionate

with voltage sensor conformational change. The predicted rate of fluorescence change ($k_{\Delta F}$)

approaches a consistent response as ECAP concentration decreased below the K_d for the resting

conformation. Thus, the fluorescence response is similar at ECAP all concentrations below the

K_d for the resting conformation, when a minority of ECAP binding sites are occupied. This

indicates that if the concentration of ECAP applied to a tissue is below the K_d for the resting

conformation, ECAP dynamics are expected to be consistent throughout a tissue, even if the

ECAP concentration is lower due to difficulty penetrating tissue or is otherwise depleted.

Overall, a broad qualitative interpretation can be made from ECAP fluorescence changes: a

change in labeling indicates that voltage sensors of unlabeled channels have changed their

probability of being activated.

GxTX-594 selectively labels Kv2 channels

Armed with a model for interpretation of changes in ECAP labeling, we proceed to determine

whether GxTX-594 could reveal conformational changes of endogenous Kv2 channels. An

important consideration in interpreting fluorescence images of tissues is whether cell labeling

with the GxTX-594 ECAP could be due to receptors other than Kv2 channels.

Electrophysiological studies have concluded that the native GxTX peptide is selective for Kv2

channels, with some off-target modulation of A-type Kv4 current (Herrington *et al.* 2006, Liu &

Bean 2014, Speca *et al.* 2014). However, electrophysiological testing only indicates whether

GxTX modulates channels; electrophysiology cannot detect ligands that bind channels, but do

not alter currents (Sack *et al.* 2013). Furthermore, structural differences between wild-type

GxTX and the GxTX-594 variant could potentially alter selectivity among channel subtypes.

To test whether GxTX-594 binds other voltage-gated K^+ channel subtypes, we quantified surface

labeling and analyzed colocalization of GxTX-594 with a selection of GFP-labeled voltage-gated

K$^+$ channel subtypes. A series of GFP-K$^+$ channel fusions were identified that express and retain

function with a GFP tag: rat Kv4.2-GFP (Shibata *et al.* 2003), rat Kv1.5-GFP (Li *et al.* 2001),

and mouse BK-GFP. Transfection of each of these channel subtypes into CHO cells resulted in

voltage-dependent outward currents, consistent with cell surface expression of the K$^+$ channels

(Fig 6, Supplement 1 A). In imaging analysis, the surface membrane was identified by labeling

with a fluorescent wheat germ agglutinin (WGA) conjugate. Transfection of all K$^+$ channel-GFP

constructs resulted in GFP localization at the cell surface, however the majority of the Kv4.2 and

Kv1.5 appeared intracellularly retained (Fig 6, Supplement 1 B). Consistent with reports in other

cell lines, cell surface localization improved when of Kv4.2 was co-transfected with KChIP2

(Shibata *et al.* 2003) or Kv1.5 with Kvβ2 (Shi *et al.* 1996) (Fig 6 A). After incubation with 100

nM GxTX-594 for five minutes, cells expressing the GFP-tagged variants of Kv2.1 and 2.2

became labeled while neither Kv4.2 + KChIP2, nor Kv1.5 + Kvβ2, nor BK channels were

appreciably labeled (Fig 6 A). When the background-subtracted GxTX-594:GFP fluorescence

intensity ratio was normalized to GxTX-594:Kv2.1-GFP, this ratio was close to zero for Kv4.2 +

KChIP2, Kv1.5 + Kvβ2, or BK (Fig 6 B), suggesting minimal binding to these channel subtypes.

Furthermore, no colocalization was apparent between Kv4.2 + KChIP2, Kv1.5 + Kvβ2, or BK

and GxTX-594 (Fig 6 C). A similar set of experiments conducted without the Kv4.2 or Kv1.5

auxiliary subunits also gave no indication of GxTX-594 labeling of Kv4.2 or Kv1.5 (Fig 6,

Supplement 1 B, C, D). We did not conduct a comprehensive survey of all the more than 80

known mammalian proteins containing voltage sensor domains, and thus cannot be certain that

GxTX-594 does not bind any other voltage sensors, or other receptors on cell surfaces. However,

the lack of labeling of these related voltage-gated K$^+$ channels indicates that GxTX-594 does not

promiscuously label voltage sensors. In all tests conducted, GxTX-594 was selective for Kv2

channels.

To determine whether expression of Kv2 channels embedded in tissue can be imaged with

GxTX-594, we overexpressed Kv2.1-GFP in brain slices. We examined rat brain at CA1

pyramidal neurons of the hippocampus. CA1 neurons express Kv2 channels, the physiology of

these neurons have been intensively studied, and their electrical properties are relatively

homogeneous (Misonou *et al.* 2005). Organotypic hippocampal slice cultures prepared from

postnatal day 5-7 rats were sparsely transfected with Kv2.1-GFP, resulting in a subset of neurons

displaying green fluorescence. When imaged two to four days after transfection, GFP

fluorescence was observed in the plasma membrane surrounding neuronal cell bodies and

proximal dendrites (Fig 7 Supplement 1 A and B). Six days or more after transfection, Kv2.1-

GFP fluorescence organized into clusters on the surface of the cell soma and proximal processes

(Fig 7 A, Fig 7 Supplement 1 C), a pattern consistent with a prior report of endogenous Kv2.1 in

CA1 neurons (Misonou *et al.* 2005). After identifying a neuron expressing Kv2.1-GFP, solution

flow into the imaging chamber was stopped and GxTX-594 was added to the static bath solution

to a final concentration of 100 nM. After five minutes of incubation, solution flow was restarted,

leading to wash-out of GxTX-594 from the imaging chamber. After wash-out, GxTX-594

fluorescence remained colocalized with Kv2.1-GFP (Fig 7, Fig 7 Supplement 1), indicating that

GxTX-594 is able to permeate through dense neural tissue and bind to Kv2 channels on neuronal

surfaces. Pearson correlation coefficients confirmed the colocalization of GxTX-594 with Kv2.1-

GFP in multiple slices (Fig 7 C). In most images of Kv2.1-GFP expressing neurons, GxTX-594

labeled puncta were observed on neighboring neurons that did not express Kv2.1-GFP, but at

intensities that were roughly an order of magnitude dimmer (Fig 7 B, white arrow). The

similarity in the labeling patterns are consistent with GxTX-594 labeling endogenous Kv2

channels. The lack of GxTX-594 labeling of non-Kv2 K^+ channels, the increased labeling of

neurons where Kv2.1-GFP is expressed, and the Kv2-like labeling patterns of CA1 neurons all

indicate that GxTX-594 selectively labels Kv2 channels in living brain slices.

GxTX-594 labeling of neurons in brain slice responds to neuronal depolarization

To test whether GxTX-594 labeling of neurons in brain slices is consistent with labeling of

endogenous Kv2 voltage sensors, we determined whether GxTX-594 labeling responds to

voltage changes. First, we looked for Kv2-like labeling patterns on CA1 pyramidal neurons in

untransfected brain slices bathed in 100 nM GxTX-594. In two-photon excitation optical sections

thinner than the neuronal cell bodies, fluorescent puncta circumscribed dark intracellular spaces

(Fig 8 A). This was similar to the patterns of fluorescence in Kv2.1-GFP transfected slices (Fig 7

B), and consistent with the punctate expression pattern of Kv2.1 in CA1 pyramidal neurons seen

in fixed brain slices (Misonou *et al.* 2005).

We tested whether the punctate fluorescence was voltage-sensitive using voltage clamp. To

ensure voltage clamp of the neuronal cell body, slices were bathed in tetrodotoxin to block Na^+

channels, and Cs^+ was included in the patch pipette solution to block K^+ currents. In each

experiment, whole-cell configuration was achieved from a GxTX-594 labeled neuron, and

holding potential was set to -70 mV. At this point, time-lapse imaging of a two-photon excitation

optical section was initiated. Depolarization to 0 mV resulted in loss of fluorescence from a

subset of puncta at the perimeter of the voltage-clamped neuron (Fig 8 B red arrows,

Supplemental Movie). Other fluorescent puncta appeared unaltered by the 0 mV step (Fig 8 B,

white arrows).

To assess if the apparent stimulus-dependent fluorescence decrease noted in puncta after a

voltage step to 0 mV was due to random fluctuations during imaging, we compared fluorescence

of the apparent cell membrane region to regions more distal from the voltage-clamped cell body.

To quantify this fluorescence change, an ROI 30 pixels (1.2 µm) wide containing the apparent

membrane of the cell body was compared to other regions within each image (Fig 8 C). The

region containing the membrane of the cell body lost fluorescence during the 0 mV step (Fig 8

D, ROI 1) while neither of the regions more distal (ROI 2, ROI 3), nor the intracellular region

(ROI 0) showed a similar change. In multiple slices, the voltage-dependent decrease of

fluorescence was seen specifically in the region of the voltage-clamped neuronal membrane (Fig

8 E). The kinetics of fluorescence response of the voltage-clamped membrane region were

similar between slices (Fig 8 F).

To determine whether the fluorescence response to depolarization was driven by the Kv2-like

puncta on the cell membrane, the fluorescence along a path containing the apparent cell

membrane was selected by drawing a path connecting fluorescent puncta surrounding the dark

cell body region, and averaging fluorescence in a region 10 pixels wide (0.4 µm) centered on this

path (Fig 8 G, yellow line). The fluorescence intensity along the path of the ROI revealed distinct

peaks corresponding to puncta (Fig 8 H, red trace). After stepping the neuron to 0 mV, the

intensity of fluorescence of a subset of puncta lessened (Fig 8 H, black trace, peaks 1,3,4,5,8).

The Scheme II model predicts a 55% reduction of fluorescence, but even more reduction is

observed in individual fluorescence puncta in brain slice (Fig 8, I blue line). This would be

expected if the concentration of GxTX-594 diminished as it penetrated into the brain tissue (Fig

6 A). In addition to the depolarization-responsive fluorescence puncta, other puncta which

appeared to be on the surface of the same cell body, maintained or increased their brightness

after depolarization (Fig 8 H, black trace, peaks 2, 6, 7). These puncta could represent Kv2

proteins on a neighboring cell, internalized Kv2-GxTX-594 complexes, off-target labeling by

GxTX-594, or Kv2 channels that are unresponsive to voltage. When the kinetics of fluorescence

intensity decay of individual voltage-sensitive puncta were fit with Eq. 1, $k_{\Delta F}$ values were similar

between puncta (Fig 8 I), consistent with these spatially separated puncta all being on the surface

of the voltage-clamped neuron. The fluorescence changes measured in these puncta were similar

to those predicted by the Figure 6 model (Fig 8 I, blue line). To address whether the fluorescence

change at the cell membrane was driven by decreases in regions of punctate fluorescence, the

punctate and non-punctate fluorescence intensity changes were analyzed separately. The regions

with fluorescence intensities above average for the path (Fig 8 H, dashed line) were binned as

one group. The above-average group, by definition, contained all punctate fluorescence. When

comparing the fluorescence before and during the 0 mV step, the regions that were initially

below average maintained the same intensity ($103 \pm 8\%$); regions of above average fluorescence

decreased in the intensity ($70 \pm 8\%$) (Fig 8 J). This suggests that the detectable unlabeling along

the cell membrane originated in the Kv2-like puncta, not the regions of lower fluorescence

intensity between them.

To determine whether the dynamics of voltage-dependent response of GxTX-594 fluorescence

on neurons in brain slices were consistent with labeling of Kv2.1, we performed experiments

with Kv2.1-expressing CHO cells under similar conditions as brain slice: 100 nM GxTX-594,

30°C, Cs^+-containing patch pipette solution. The rate of fluorescence changes in Kv2.1-CHO

cells were similar to neurons in brain slices, and consistent with the $k_{\Delta F}$ of 0.06 s^{-1} predicted by

Figure 6 model (Fig 8 K). The data indicate that the voltage sensors of Kv2 channels in

hippocampal CA1 neurons have a similar response to voltage as the voltage sensors of Kv2.1

channels in CHO cells.

Discussion

We conclude that GxTX-594 dynamically labels a resting conformation of endogenous Kv2

channels in neurons. This conclusion is based on a series of findings: GxTX-594 labels Kv2

proteins (Fig 1), in the presence of its obligate neuronal auxiliary subunit AMIGO-1 (Fig 2), but

does not label related, non-Kv2, Kv proteins (Fig 6). Steps to 0 mV resulted in dynamic

unlabeling of CHO cells (Fig 3) and neurons (Fig 8), with similar kinetics (Fig 8 K). The

punctate GxTX-594 labeling of CA1 neurons is consistent with the patterns of endogenous Kv2

expression seen by immunofluorescence (Misonou *et al.* 2005, Bishop *et al.* 2015, 2018), and

Kv2.1-GFP expression (Fig 7). These results are consistent with the GxTX-594 probe labeling

endogenous Kv2 channels in rat CA1 hippocampal pyramidal neurons and unlabeling in

response to voltage activation of the channels. Thus, tracking the fluorescence of conformation-

dependent ligands such as GxTX-594 can reveal conformational changes of endogenous

proteins. Below we discuss physiological questions that could be addressed with a GxTX-based

Kv2 voltage sensor probe, the limitations of GxTX-594 as a probe, and the potential for

engineering conformation-selective imaging probes to report activity of other endogenous

proteins.

A voltage sensor probe to study Kv2 physiology

The GxTX-594 probe is an ECAP that reveals conformational changes of Kv2 voltage sensors.

Activation of Kv2 voltage sensors is required for opening of its K^+-conductive pore (Islas &

Sigworth 1999). Cells dynamically regulate the voltage sensitivity of Kv2 channel pore opening

(Murakoshi *et al.* 1997, MacDonald *et al.* 2003, Misonou *et al.* 2005, 2006, Redman *et al.* 2007,

Plant *et al.* 2011, Mandikian *et al.* 2014). It is unknown whether these regulatory processes

impact voltage sensor movement directly, or impact conformational changes that are coupled to

voltage sensor movement, such as pore opening. By specifically measuring voltage sensor

movements, the mechanisms by which pore opening is regulated can be better understood. In

studies of the mechanism of cellular regulation of Kv2.1 in HEK cells, it was concluded that

voltage sensor movements were decoupled from pore opening in the majority of channels with

punctate localization (O'Connell *et al.* 2010, Fox *et al.* 2013). This conclusion was supported by

assessing movement of voltage sensors with gating currents, which are currents arising from the

movement of voltage sensors themselves. Gating current measurements are feasible only in cell

preparations where a sufficiently high density of voltage sensors can be voltage-clamped. This density requirement has resulted in gating current measurement being most common with recombinant overexpression in heterologous systems in which a high density of protein can be trafficked to the surface membrane. Measurements of endogenous gating currents have been restricted to a limited number of cell preparations such as invertebrate giant axons (Armstrong & Bezanilla 1973) and muscle cells (M. F. Schneider & Chandler 1973). Gating currents or other measures of conformational change of endogenous voltage sensors have never, to our knowledge, been measured in a brain slice.

The GxTX-594 ECAP has revealed the first glimpse of how Kv2 voltage sensors are operating in a tissue. The kinetics of the GxTX-594 response to voltage were similar in brain slice and CHO cells, indicating that the stimulus-response relation of voltage sensor conformational change is similar in the two cell preparations. Our results indicate that the voltage sensors of endogenous Kv2 proteins clustered on the surface of CA1 pyramidal neurons in rat hippocampus are responsive to voltage. This result is perhaps expected, the result is consistent with the proposal of Tamkun and colleagues (Fox *et al.* 2013), that only a small fraction of clustered Kv2.1 channels in HEK cells conduct K^+ current, while many more channels change conformation in response to voltage change. In addition to reporting when and where Kv2 channels enter conformations that are prerequisites for channel opening, Kv2 ECAPs can reveal where nonconducting Kv2 proteins change conformation. Whether conformational changes are important for the growing list of Kv2's nonconducting functions are unknown. By reporting voltage sensor activation, the GxTX-594 probe could potentially be used to determine whether voltage sensor conformational change is crucial for the nonconducting functions of Kv2 proteins.

Limitations of using GxTX-594 to measure conformational change
The ability of conformation-selective probes to report conformational change is inherently limited. Some limitations result from the properties of the particular probe deployed, while other limitations are inherent to the mechanism of conformation-selective binding. Below, we discuss limitations of GxTX-594 and how those limitations can be minimized in the realms of temporal resolution, signal to noise, protein modulation by probes, and errors in interpretation of images.

Temporal resolution. With GxTX-594, the average probability that voltage sensors are activated

($P_{activated}$), can be measured. GxTX-594 requires seconds to respond, much slower than the

membrane voltage changes associated with action potentials. Probes with faster kinetics are

expected to enable measurements of faster patterns of channel activity. Modifying probe kinetics

becomes a question of sensor design and toxin engineering; the response properties of probes can

be altered by chemical modification of side chain residues, allowing the dynamics to be

optimized in next-generation probes. Ultimately the response time of any conformation-selective

probe will be limited by the labeling rate which is a product of the probe's concentration and its

inherent association and dissociation constants. Alternatively, ECAPs that remain bound while

producing a fluorescence change in response to their target's conformational change could

provide much faster time resolution.

Signal to noise. The fluorescence signal intensity from GxTX-594 limits spatial and temporal

resolution, estimates of $P_{activated}$, and interpretation of data. Continuing advances in microscopy

are expected to improve fundamental signal to noise of fluorescence imaging in tissue. However,

the signal to noise in all experiments is ultimately limited by the photophysics of the fluorophore

as well as photodamage to the probe and tissue. Probes with improved fluorescence

characteristics are expected to improve the ultimate signal to noise. Quantum dot nanoparticles

offer orders of magnitude improvement in signal to noise for one- and two-photon excitation

microscopy methods. We have developed synthetic methods for making Kv2 quantum dot

ECAPs from GxTX (Mann *et al.* 2018). Such probes are expected to respond to Kv2 channel

voltage sensor activation with similarly interpretable fluorescence changes, yet offer orders of

magnitude improvements in signal to noise ratio.

Protein modulation by probes. GxTX-based probes inhibit the Kv2 channels they label. Thus,

they deplete the population of channels responding normally to physiological stimuli. This is a

general problem with molecular probes: they disrupt the process they report (e.g. free calcium

indicators deplete free calcium). This probe perturbation problem can be minimized by working

with low concentrations of probe to inhibit a minimal subset of the total Kv2 population.

Channel activity can be recorded optically even when only a small fraction of Kv2 channels are

affected (Tilley *et al.* 2014). Furthermore, GxTX-594 most accurately reports conformational

change when used at sub-saturating concentrations (Fig 5). Thus, low concentrations are optimal

for both accuracy of measurement and minimizing the perturbation of physiology.

Oversimplification of Kv2 gating dynamics. The models enabling quantification of

conformational change involve approximations. The statistical thermodynamics developed here

involve ligand binding and conformational change dynamics using approximations of Eyring rate

theory (Eyring 1935a, b). Thus, models will always have limitations stemming from the

necessary approximations. The model of Kv2 voltage sensor conformational change developed

here is a further oversimplification. While the gating dynamics of Kv2 channels have never been

fully described with an explicit thermodynamic model, they are certainly more complex than our

model. Our model assumes each of the of the channel's 4 voltage sensors bind 1 GxTX, and that

the 4 voltage sensors change conformation independently of each other. However, Kv2 channels

exhibit cooperative gating when they open their pores, and pore opening impacts voltage sensors

(Islas & Sigworth 1999, Scholle, Dugarmaa, *et al.* 2004, Jara-Oseguera *et al.* 2011, Tilley *et al.*

2019). Under some conditions the assumption of voltage sensor independence may limit the

model's predictive power. Additionally, the model of GxTX-594 labeling developed here

assumes that voltage sensors are in continuous equilibrium. This simplification is expected to

result in diminished accuracy during high frequency voltage changes, such as during action

potentials. These deviations from equilibrium could explain the deviation of the model from the

data in response to high frequency voltage steps (Fig 4 C, D).

Conformation-selective probes reveal conformational change of endogenous proteins

If the limitations of conformation-selective binding are given adequate consideration,

deconvolution of fluorescence intensities can reveal conformational changes of endogenous

proteins. Importantly, measurements of dynamic labeling by a conformation-selective probe,

such as GxTX can enable deduction of how unlabeled proteins behave. This is perhaps counter-

intuitive because GxTX inhibits voltage sensor movement of the Kv2 subunit it binds, and only

the bound subunits generate optical signals (Tilley *et al.* 2014, 2019). This approach is analogous

to calcium imaging experiments, which have been spectacularly informative about physiological

calcium signaling (W. Yang & Yuste 2017), despite the fact that no optical signals originate

from the physiologically relevant free Ca^{2+}, only from Ca^{2+} that is chelated by a dye. In all such

experiments, fluorescence from Ca^{2+}-bound dyes is deconvolved using the statistical

thermodynamics of Ca^{2+} binding to calculate free Ca^{2+} (Adams 2010). Similarly, GxTX-based

probes dynamically bind to unlabeled Kv2 proteins, and the binding rate is dependent on the

probability that unlabeled voltage sensors are in a resting conformation (Fig 5). Thus, the

conformations of unlabeled Kv2 proteins influence the dynamics of labeling with GxTX-based

probes. Consequently, the dynamics of labeling reveal the conformations of unlabeled Kv2

proteins.

Deployment of GxTX-594 to report conformational changes of endogenous channels

demonstrates that conformation-selective ligands can be used to image occurrence of the

conformations they bind to. The same principles of action apply to any conformation-selective

labeling reagent, suggesting that probes for conformational changes of many different proteins

could be developed. Probes could conceivably be developed from the many other voltage sensor

toxins and other gating modifiers that act by a similar mechanism as GxTX, yet target the

voltage sensors of different ion channel proteins (McDonough *et al.* 1997, Sack *et al.* 2004,

Catterall *et al.* 2007, Swartz 2007, Schmalhofer *et al.* 2008, Peretz *et al.* 2010, McCormack *et al.*

2013, Ahuja *et al.* 2015, A. H. Zhang *et al.* 2018, Dockendorff *et al.* 2018). Conformation-

selective binders have been engineered for a variety of other proteins, yet methods to quantify

conformational changes from their fluorescence are needed. For example, fluorescently-labeled

conformation-selective binders have revealed that endocytosed GPCRs continue to remain in a

physiologically activated conformation (Irannejad *et al.* 2013, Tsvetanova *et al.* 2015, Eichel &

von Zastrow 2018). However, a means to determine the conformational equilibria of GPCRs

from fluorescence images has not yet been developed. We suggest that the statistical

thermodynamic framework developed here could provide a starting point for more quantitative

interpretation of of other conformation-selective molecular probes.

Materials and methods

GxTX-594 synthesis

The Ser13Cys GxTX peptide synthesized as described (Tilley *et al.* 2014). Methionine 35 of

GxTX was replaced by the oxidation-resistant noncanonical amino acid norleucine and serine 13

was replaced with cysteine to create a spinster thiol. Ser13Cys GxTX was labeled with a Texas

red derivative (Alexa Fluor 594 C_5 maleimide, Thermo Fisher Scientific, Catalog # 10256).

Ser13Cys GxTX lyophilisate was brought to 560 μM in 50% ACN + 1 mM Na_2EDTA. 2.4 μL

of 1M Tris (pH 6.8 with HCl), 4 μL of 10 mM of Alexa Fluor 594 C_5 maleimide in DMSO and

17.9 μL of 560 μM Ser13Cys GxTX were added for a final solution of 100 mM Tris, 1.6 mM

Alexa Fluor 594 C_5 maleimide and 0.4 mM GxTX in 24 μL of reaction solution. Reactants were

combined in a 1.5 mL low protein binding polypropylene tube (LoBind, Eppendorf, Catalog #

022431081) and mixed at 1000 RPM, 20°C for 4 hours (Thermomixer 5355 R, Eppendorf).

After incubation, the tube was centrifuged at 845 RCF for 10 minutes at room temperature. A

purple pellet was observed post-centrifugation. The supernatant was transferred to a fresh tube

and centrifuged at 845 RCF for 10 minutes. After this 2[nd] centrifugation, no visible pellet was

seen. The supernatant was injected onto a reverse-phase HPLC C_{18} column (Biobasic- 4.6mm

RP-C_{18} 5 μm, Thermo Fisher Scientific, Catalog # 2105-154630) equilibrated in 20% ACN,

0.1% TFA at 1 mL/min and eluted with a protocol holding in 20% ACN for 2 minutes,

increasing to 30% ACN over 1 minute, then increasing ACN at 0.31% per minute. HPLC

effluent was monitored by fluorescence and an absorbance array detector. 1 ml fractions were

pooled based on fluorescence (280 nm excitation, 350 nm emission) and absorbance (214 nm,

280 nm, and 594 nm). GxTX-594 peptide-fluorophore conjugate eluted at approximately 35%

ACN, and mass was confirmed by mass spectrometry using a Bruker UltraFlextreme MALDI-

TOF/TOF (Fig 1 Supplement 1). Samples for identification from HPLC eluant were mixed 1:1 in

an aqueous solution of 25% MeOH and 0.05% TFA saturated with alpha-cyano-4-hydrocinnamic

acid, pipetted onto a ground-steel plate, dried under vacuum, and ionized with 60-80% laser

power. Molecular species were detected using a reflector mode protocol and quantitated using

Bruker Daltonics flexAnalysis 3.4. Lyophilizate containing GxTX-594 conjugation product was

dissolved in CE buffer (defined below), and stored at -80 °C. GxTX-594 concentration was

determined by 280 nm absorbance using a calculated extinction coefficient of 18,900 A.U. M^{-1}

cm^{-1}.

CHO cell methods

CHO cell culture and transfection

The CHO-K1 cell line (ATCC) and a subclone transfected with a tetracycline-inducible rat

Kv2.1 construct (Kv2.1-CHO) (Trapani & Korn 2003), were cultured as described previously

(Tilley *et al.* 2014). To induce channel expression in Kv2.1-TREx-CHO cells, 1 μg/mL

minocycline (Enzo Life Sciences, Catalog # ALX-380-109-M050), prepared in 70% ethanol at 2

mg/mL, was added to the maintenance media to induce K^+ expression. Minocycline was added

40-48 hours before imaging and voltage clamp fluorometry experiments. Transfections were

achieved with Lipofectamine 2000 (Life Technologies, Catalog # 1668027). 1.1 μL of

Lipofectamine was diluted, mixed, and incubated in 110 μL of Opti-MEM (Gibco, Product

Number 31985-062, Lot Number 1917064) in a 1:100 ratio for 5 minutes at room temperature.

Concurrently, 1 μg of plasmid DNA and 110 μL of Opti-MEM were mixed in the same fashion.

DNA and Lipofectamine 2000 mixtures were combined, titurated, and let sit for 20 minutes.

Then, the transfection cocktail mixture was added to a 35 mm cell culture dish of CHO cells at

approximately 40% confluency and allowed to settle at 37°C 5% CO_2 for 4-6 hours before the

media was replaced. Rat Kv2.1-GFP (Antonucci *et al.* 2001), rat Kv2.2-GFP (Kirmiz, Vierra, *et

al.* 2018), rat Kv1.5-GFP (Li *et al.* 2001), rat Kv4.2-GFP (Shibata *et al.* 2003), mouse BK-GFP

(Trimmer Lab, UC Davis), rat Kv2.1 S586A-GFP (Kirmiz, Palacio, *et al.* 2018), rat AMIGO-

YFP (Bishop *et al.* 2015), rat KvBeta2 (Shibata *et al.* 2003) and rat KCHiP2 (An *et al.* 2000)

plasmids were all kind gifts from James Trimmer, University of California, Davis. Identities of

constructs were confirmed by sequencing from their CMV promoter.

Confocal and Airy disk imaging

Confocal images were obtained with an inverted confocal system (Zeiss LSM 880 410900-247-

075) run by ZEN black v2.1. A 63x/1.40 NA Oil DIC objective (Zeiss 420782-9900-799) was

used for most imaging experiments; a 63x/1.2 NA Water DIC objective (Zeiss 441777-9970-

000) was used for voltage clamp fluorometry experiments. GFP and YFP were excited with the

488 nm line from an argon laser (3.2 mW at installation) powered at 0.5% unless otherwise

noted. GxTX594 was excited with 594 nm helium-neon laser (0.6 mW at installation) powered at

10% unless otherwise noted. WGA-405 was excited with a 405 nm diode laser, (3.5 mW at

installation) powered at 1% unless otherwise noted. Temperature inside the microscope housing

was 27-30 °C.

In confocal imaging mode, fluorescence was collected with the microscope's 32-channel gallium

arsenide phosphide detector array arranged with a diffraction grating to measure 400-700 nm

emissions in 9.6 nm bins. Emission bands were 495-550 nm for GFP and YFP, 605-700 nm for

GxTX-594, and 420-480 nm for WGA-405. Point spread functions were calculated using ZEN

black software using emissions from 0.1 μm fluorescent microspheres, prepared on a slide

according to manufacturer's instructions (Thermo Fisher Scientific, Catalog # T7279). The point

spread functions for confocal images with the 63x/1.40 Oil DIC objective in the X-Y direction

were 228 nm (488 nm excitation) and 316 nm (594 nm excitation).

In Airy disk imaging mode, fluorescence was collected with the microscope's 32-channel

gallium arsenide phosphide detector array arranged in a concentric hexagonal pattern (Zeiss

Airyscan 410900-2058-580). After deconvolution, the point spread functions for the 63x/1.40

NA objective with 488 nm excitation was 124 nm in X-Y and 216 nm in Z; with 594 nm

excitation 168 nm in X-Y and 212 nm in Z. For the 63x/1.2 NA objective, the point spread

function with 488 nm excitation was 187 nm in X-Y and 214nm in Z; the point spread function

with 594 nm excitation was 210 nm in X-Y and 213 nm in Z.

Unless stated otherwise, cells were plated in uncoated 35 mm dishes with a 7 mm inset No. 1.5

coverslip (MatTek, Catalog # P35G-1.5-20-C). The CHO cell external (CE) solution used for

imaging and electrophysiology contained (in mM): 3.5 KCl, 155 NaCl, 10 HEPES, 1.5 $CaCl_2$, 1

$MgCl_2$, adjusted to pH 7.4 with NaOH. Measured osmolarity was 315 mOsm. When noted,

solution was supplemented with either 10 mM glucose (CEG) or 10 mM glucose and 1% BSA

(CEGB).

For time-lapse, GxTX-594 dose-response experiments, Kv2.1-CHO cells were plated onto 22 x

22 mm No. 1.5H coverglass (Deckglaser). Prior to imaging, cell maintenance media was

removed and replaced with CEGB, then coverslip was mounted on an imaging chamber (Warner

Instrument, Catalog # RC-24E) with vacuum grease. During time-lapse imaging, indicated

concentrations of GxTX-594 were applied for 10 minutes followed by manual wash-out with

CEGB solution. 15 minutes after wash-out, the next GxTX-594 concentration was added.

Solutions were added to the imaging chamber perfusion via a syringe at a flow rate of ~1 mL /

10 sec. Images were taken every 5 sec. Laser power was set to 0.5% for the 488 nm laser and

1.5% for the 594 nm laser. For colocalization experiments with GFP-tagged proteins, cells were

incubated in 100 nM GxTX-594 for 5 minutes and then washed with CEGB before imaging.

Whole cell voltage clamp for CHO cell imaging

Kv2.1-TREx-CHO cells plated in glass-bottom 35 mm dishes were washed with CEGB, placed

in the microscope, and then incubated in 100 µL of 100 nM GxTX594 for 5 minutes to label

cells. Before patch clamp, the solution was diluted with 1 mL of CEG for a working

concentration of 9 nM GxTX594 during experiments. Cells with obvious GxTX-594 surface

staining were voltage-clamped in whole-cell mode with an EPC-10 patch clamp amplifier

(HEKA) run by Patchmaster software, v2x90.2 (HEKA). The patch pipette contained a

potassium-deficient Cs^+ internal pipette solution to limit outward current and reduce voltage

error: 70 mM CsCl, 50 mM CsF, 35mM NaCl, 1 mM EGTA, 10 mM HEPES, brought to pH 7.4

with CsOH. Osmolarity was 310 mOsm. The liquid junction potential was calculated to be 3.5

mV and was not corrected. Borosilicate glass pipettes (Sutter Instruments, Catalog # BF 150-

110-10HP) were pulled with blunt tips to resistances less than 3.0 MΩ in these solutions. Cells

were held at -80 mV (unless noted otherwise) and stepped to indicated voltages. The voltage step

stimulus was maintained until any observed change in fluorescence was complete. For stimulus-

frequency dependence experiments, cells were given 2 ms steps to 40 mV at stated frequencies

(0.02, 5, 10, 20, 50, 100, 150, or 200 Hz). Images for voltage clamp fluorometry were taken in

Airy disk imaging mode with the settings described above.

Whole cell voltage clamp for CHO cell K^+ current recordings

Whole-cell voltage clamp was used to measure currents from CHO cells expressing Kv2.1-GFP,

Kv2.2-GFP, Kv1.5-GFP, Kv4.2-GFP, BK-GFP or GFP that we transfected as described above.

Cells were plated on glass bottom dishes. Cells were held at -80 mV, then 100 ms voltage steps

were delivered ranging from -80 mV to +80 mV in +5 mV increments. Pulses were repeated

every 2 seconds. The external (bath) solution contained CE solution. The internal (pipette)

solution contained (in mM): 35 KOH, 70 KCl, 50 KF, 50 HEPES, 5 EGTA adjusted to pH 7.2

with KOH. Liquid junction potential was calculated to be 7.8 mV and was not corrected.

Borosilicate glass pipettes (Sutter Instruments, Cat # BF150-110-10HP) were pulled into pipettes

with resistance less than 3 MΩ for patch clamp recording. Recordings were at room temperature

(22–24 °C). Voltage clamp was achieved with an Axon Axopatch 200B amplifier (MDS

Analytical Technologies) run by Pachmaster software, v2x90.2 (HEKA). Holding potential was

-80 mV. Capacitance and Ohmic leak were subtracted using a P/5 protocol. Recordings were low

pass filtered at 10 kHz and digitized at 100 kHz. Voltage clamp data were analyzed and plotted

with Igor Pro 7 or 8 (Wavemetrics). Current amplitudes at each voltage were the average from

0.19-0.20 s after voltage step. As the experiments plotted in Figure 6 Supplement 1 A were

merely to confirm functional expression of ion channels at the cell surface, series resistance

compensation was not used and substantial cell voltage errors are predicted during these

experiments.

Brain slice methods

Hippocampal slice culture preparation and transfection

All experimental procedures were approved by the University of California Davis Institutional

Animal Care and Use Committee and were performed in strict accordance with the Guide for the

Care and Use of Laboratory Animals of the NIH. Animals were maintained under standard light–

dark cycles and had ad libitum access to food and water. Organotypic hippocampal slice cultures

were prepared from postnatal day 5-7 rats, as described (Stoppini *et al.* 1991) and detailed in a

video protocol (Opitz-Araya & Barria 2011). DIV15-30 neurons were transfected 2-6 days

before imaging via biolistic gene transfer (160 psi, Helios gene gun, Bio-Rad), as described in a

detailed video protocol (Woods & Zito 2008). 10 µg of plasmid was coated to 6-8 mg of 1.6 µm

gold beads.

Two-photon excitation slice imaging

Image stacks (512 x 512 pixels, 1 mm Z-steps; 0.035 µm/pixel) were acquired using a custom

two-photon excitation microscope (Olympus LUMPLFLN 60XW/IR2 objective, 60x, 1.0 NA)

with two pulsed Ti:sapphire lasers (Mai Tai: Spectra Physics) tuned to 810 nm (for GxTX-594

imaging) and 930 nm (for GFP imaging) and controlled with ScanImage (Pologruto *et al.* 2003).

After identifying a neuron expressing Kv2.1-GFP, perfusion was stopped and GxTX-594 was

added to the static bath solution to a final concentration of 100 nM. After five minutes

incubation, perfusion was restarted, leading to wash-out of GxTX-594 from the slice bath. Red

and green photons (Chroma: 565dcxr, BG-22 glass, HQ607/45) emitted from the sample were

collected with two sets of photomultiplier tubes (Hamamatsu R3896).

Whole cell voltage clamp for brain slice imaging

Organotypic hippocampal slice cultures (DIV 6-7, not transfected) were transferred to an

imaging chamber with recirculating ACSF maintained at 30°C. To hold the slice to the bottom of

the chamber, a horseshoe-shaped piece of gold wire was used to weight the membrane holding

the slice. ACSF contained (in mM) 127 NaCl, 25 NaHCO$_3$, 25 D-glucose, 2.5 KCl, 1.25

NaH$_2$PO$_4$, 1 MgCl2, 2 CaCl$_2$, 200 nM tetrodotoxin, pH 7.3 and aerated with 95% O$_2$/ 5% CO$_2$

(~310mOsm). 4 mL of 100 nM GxTX-594 in ACSF were used in the circulating bath to allow

the toxin to reach the slice and reach the desired concentration of 100 nM throughout the

circulating bath. Images were acquired beginning 3 minutes after GxTX-594 was added.

Apparent CA1 neurons with GxTX-594 labeling in a Kv2-like pattern were selected for whole-

cell patch clamp. Voltage clamp was achieved using an Axopatch 200B amplifier (Axon

Instruments) controlled with custom software written in Matlab (MathWorks, Inc.). Patch

pipettes (5-7 MΩ) were filled with intracellular solution containing (in mM): 135 Cs-

methanesulfonate, 10 Na$_2$-phosphocreatine, 3 Na-L-ascorbate, 4 NaCl, 10 HEPES, 4 MgCl$_2$, 4

Na$_2$ATP, 0.4 NaGTP, pH 7.2. Neurons were clamped at -70 mV. Input resistance and holding

current were monitored throughout the experiment. Cells were excluded if the pipette series

resistance was higher than 25 MΩ or if the holding current exceeded -100 pA. To activate Kv2

channels, a 50 s depolarizing step from -70 mV to 0 mV was given.

Image analysis

Fluorescence images were analyzed using ImageJ 1.52n software (C. A. Schneider *et al.* 2012).

Regions of interest (ROIs) encompassed the entire fluorescent region of an individual cell or

neuron, unless mentioned otherwise. ROIs were drawn manually. Analysis of images was

conducted independently by multiple researchers who produced similar results, but analysis was

not conducted in a blinded nor randomized fashion. Fluorescence intensity (*F*) was background

subtracted using mean F of a region lacking cells. For F/F_{init} normalization, F_{init} was the mean

fluorescence preceding the indicated voltage stimuli, or the max observed intensity in dose

response experiments. Further details of specific normalization procedures are provided in figure

legends. Time dependence of fluorescence intensity was fit with a monoexponential decay (Eq.

1). For colocalization analyses the Pearson coefficient was calculated using the JACoP plugin

(Bolte & Cordelières 2006). Colocalization analyses were conducted within ROIs defining

individual cells. Plotting and curve fitting was performed with with Igor Pro 7 or 8

(Wavemetrics), which performs nonlinear least-squares fits using a Levenberg-Marquardt

algorithm. Error values from individual curve fittings are standard deviations. All other errors,

including error bars, indicate standard errors. Arithmetic means are reported for intensity

measurements and correlation coefficients. Geometric means are reported for time constants and

rates. Statistical comparisons were with the Mann-Whitney U test.

Model calculations

Predictions of the Scheme I and II models were calculated using Microsoft Excel. A spreadsheet

containing model calculations that can be used to generate model prediction is included as a

supplement (Model Calculation Supplement).

Acknowledgements

We thank James Trimmer (University of California, Davis) for numerous discussions, and constructive critical reading of an early version of the manuscript. We thank Georgeann Sack (Afferent, LLC) for critical reading, editing, and feedback. This research was supported by US National Institutes of Health grants R01NS096317 (B.E.C. and J.T.S.), U01NS090581, R21EY026449, and T32GM007377. P. Thapa was supported by American Heart Association postdoctoral fellowship 17POST33670698. Work at the Molecular Foundry was supported by the Director, Office of Science, Office of Basic Energy Sciences, Division of Materials Sciences and Engineering, of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. The authors declare no competing financial interests.

Author contributions (CRediT nomenclature)

Parashar Thapa: Conceptualization, Formal analysis, Investigation, Methodology, Visualization, Writing- original draft, Writing- reviewing & editing

Robert Stewart: Conceptualization, Formal analysis, Investigation, Methodology, Visualization, Writing- original draft, Writing- reviewing & editing

Rebecka J. Sepela: Conceptualization, Formal analysis, Investigation, Methodology, Visualization, Writing- original draft, Writing- reviewing & editing

Oscar Vivas: Investigation, Methodology, Writing- reviewing & editing

Laxmi K. Parajuli: Investigation, Methodology, Writing- reviewing & editing

Mark Lillya: Conceptualization, Formal analysis, Investigation, Methodology, Visualization

Sebastian Fletcher-Taylor: Investigation, Methodology, Writing- reviewing & editing

Bruce E. Cohen: Conceptualization, Funding acquisition, Project administration, Supervision, Writing- reviewing & editing

Karen Zito: Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Supervision, Writing- original draft, Writing- reviewing & editing

Jon T. Sack: Conceptualization, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Supervision, Visualization, Writing- original draft, Writing- reviewing & editing

References

Adams, S.R. (2010) How calcium indicators work. *Cold Spring Harbor Protocols*, **2010**, pdb.top70. doi:10.1101/pdb.top70

Aggarwal, S.K. & MacKinnon, R. (1996) Contribution of the S4 segment to gating charge in the Shaker K+ channel. *Neuron*, **16**, 1169–77. doi:10.1016/S0896-6273(00)80143-9

Ahuja, S., Mukund, S., Deng, L., Khakh, K., Chang, E., Ho, H., Shriver, S., Young, C., Lin, S., Johnson, J.P., Wu, P., Li, J., Coons, M., Tam, C., Brillantes, B., Sampang, H., Mortara, K., Bowman, K.K., *et al.* (2015) Structural basis of Nav1.7 inhibition by an isoform-selective small-molecule antagonist. *Science*, **350**, aac5464. doi:10.1126/science.aac5464

An, W.F., Bowlby, M.R., Betty, M., Cao, J., Ling, H.P., Mendoza, G., Hinson, J.W., Mattsson, K.I., Strassle, B.W., Trimmer, J.S. & Rhodes, K.J. (2000) Modulation of A-type potassium channels by a family of calcium sensors. *Nature*, **403**, 553–556. doi:10.1038/35000592

Antonucci, D.E., Lim, S.T., Vassanelli, S. & Trimmer, J.S. (2001) Dynamic localization and clustering of dendritic Kv2.1 voltage-dependent potassium channels in developing hippocampal neurons. *Neuroscience*, **108**, 69–81. doi:10.1016/S0306-4522(01)00476-6

Armstrong, C.M. & Bezanilla, F. (1973) Currents related to movement of the gating particles of the sodium channels. *Nature*, **242**, 459–61. doi:10.1038/242459a0

Bezanilla, F. (2018) Gating currents. *The Journal of General Physiology*, **150**, 911–932. doi:10.1085/jgp.201812090

Bishop, H.I., Cobb, M.M., Kirmiz, M., Parajuli, L.K., Mandikian, D., Philp, A.M., Melnik, M., Kuja-Panula, J., Rauvala, H., Shigemoto, R., Murray, K.D. & Trimmer, J.S. (2018) Kv2 Ion Channels Determine the Expression and Localization of the Associated AMIGO-1 Cell Adhesion Molecule in Adult Brain Neurons. *Frontiers in Molecular Neuroscience*, **11**, 1. doi:10.3389/fnmol.2018.00001

Bishop, H.I., Guan, D., Bocksteins, E., Parajuli, L.K., Murray, K.D., Cobb, M.M., Misonou, H., Zito, K., Foehring, R.C. & Trimmer, J.S. (2015) Distinct Cell- and Layer-Specific Expression Patterns and Independent Regulation of Kv2 Channel Subtypes in Cortical Pyramidal Neurons. *The Journal of Neuroscience*, **35**, 14922–42. doi:10.1523/JNEUROSCI.1897-15.2015

Bolte, S. & Cordelières, F.P. (2006) A guided tour into subcellular colocalization analysis in light microscopy. *Journal of Microscopy*, **224**, 213–32. doi:10.1111/j.1365-2818.2006.01706.x

Catterall, W.A., Cestèle, S., Yarov-Yarovoy, V., Yu, F.H., Konoki, K. & Scheuer, T. (2007) Voltage-gated ion channels and gating modifier toxins. *Toxicon*, **49**, 124–141. doi:10.1016/j.toxicon.2006.09.022

Dockendorff, C., Gandhi, D.M., Kimball, I.H., Eum, K.S., Rusinova, R., Ingólfsson, H.I., Kapoor, R., Peyear, T., Dodge, M.W., Martin, S.F., Aldrich, R.W., Andersen, O.S. & Sack, J.T. (2018) Synthetic Analogues of the Snail Toxin 6-Bromo-2-mercaptotryptamine Dimer (BrMT) Reveal That Lipid Bilayer Perturbation Does Not Underlie Its Modulation of Voltage-Gated Potassium Channels. *Biochemistry*, **57**, 2733–2743. doi:10.1021/acs.biochem.8b00292

Eichel, K. & Zastrow, M. von. (2018) Subcellular Organization of GPCR Signaling. *Trends in Pharmacological Sciences*, **39**, 200–208. doi:10.1016/j.tips.2017.11.009

Eyring, H. (1935) The Activated Complex and the Absolute Rate of Chemical Reactions. *Chemical Reviews*, **17**, 65–77, American Chemical Society. doi:10.1021/cr60056a006

Eyring, H. (1935) The Activated Complex in Chemical Reactions. *The Journal of Chemical*

Physics, **3**, 107–115, American Institute of Physics. doi:10.1063/1.1749604

Fox, P.D., Loftus, R.J. & Tamkun, M.M. (2013) Regulation of Kv2.1 K(+) conductance by cell surface channel density. *The Journal of Neuroscience*, **33**, 1259–70. doi:10.1523/JNEUROSCI.3008-12.2013

Gamper, N., Stockand, J.D. & Shapiro, M.S. (2005) The use of Chinese hamster ovary (CHO) cells in the study of ion channels. *Journal of Pharmacological and Toxicological Methods*, **51**, 177–85. doi:10.1016/j.vascn.2004.08.008

Gilchrist, J., Das, S., Petegem, F. Van & Bosmans, F. (2013) Crystallographic insights into sodium-channel modulation by the β4 subunit. *Proceedings of the National Academy of Sciences of the United States of America*, **110**, E5016-24. doi:10.1073/pnas.1314557110

Gupta, K., Zamanian, M., Bae, C., Milescu, M., Krepkiy, D., Tilley, D.C., Sack, J.T., Yarov-Yarovoy, V., Kim, J. Il & Swartz, K.J. (2015) Tarantula toxins use common surfaces for interacting with Kv and ASIC ion channels. *eLife*, **4**, e06774. doi:10.7554/eLife.06774

Herrington, J., Zhou, Y.P., Bugianesi, R.M., Dulski, P.M., Feng, Y., Warren, V.A., Smith, M.M., Kohler, M.G., Garsky, V.M., Sanchez, M., Wagner, M., Raphaelli, K., Banerjee, P., Ahaghotu, C., Wunderler, D., Priest, B.T., Mehl, J.T., Garcia, M.L., *et al.* (2006) Blockers of the Delayed-Rectifier Potassium Current in Pancreatic -Cells Enhance Glucose-Dependent Insulin Secretion. *Diabetes*, **55**, 1034–1042. doi:10.2337/diabetes.55.04.06.db05-0788

Irannejad, R., Tomshine, J.C., Tomshine, J.R., Chevalier, M., Mahoney, J.P., Steyaert, J., Rasmussen, S.G.F., Sunahara, R.K., El-Samad, H., Huang, B. & Zastrow, M. von. (2013) Conformational biosensors reveal GPCR signalling from endosomes. *Nature*, **495**, 534–8. doi:10.1038/nature12000

Islas, L.D. & Sigworth, F.J. (1999) Voltage sensitivity and gating charge in Shaker and Shab family potassium channels. *The Journal of General Physiology*, **114**, 723–42. doi:10.1085/jgp.114.5.723

Jara-Oseguera, A., Ishida, I.G., Rangel-Yescas, G.E., Espinosa-Jalapa, N., Pérez-Guzmán, J.A., Elías-Viñas, D., Lagadec, R. Le, Rosenbaum, T. & Islas, L.D. (2011) Uncoupling charge movement from channel opening in voltage-gated potassium channels by ruthenium complexes. *The Journal of Biological Chemistry*, **286**, 16414–25. doi:10.1074/jbc.M110.198010

Jiang, Q.-X., Wang, D.-N. & MacKinnon, R. (2004) Electron microscopic analysis of KvAP voltage-dependent K+ channels in an open conformation. *Nature*, **430**, 806–10. doi:10.1038/nature02735

Jiang, Y., Lee, A., Chen, J., Ruta, V., Cadene, M., Chait, B.T. & MacKinnon, R. (2003) X-ray structure of a voltage-dependent K+ channel. *Nature*, **423**, 33–41. doi:10.1038/nature01580

Kihira, Y., Hermanstyne, T.O. & Misonou, H. (2010) Formation of Heteromeric Kv2 Channels in Mammalian Brain Neurons. *Journal of Biological Chemistry*, **285**, 15048–15055. doi:10.1074/jbc.M109.074260

Kimm, T., Khaliq, Z.M. & Bean, B.P. (2015) Differential Regulation of Action Potential Shape and Burst-Frequency Firing by BK and Kv2 Channels in Substantia Nigra Dopaminergic Neurons. *The Journal of Neuroscience*, **35**, 16404–17. doi:10.1523/JNEUROSCI.5291-14.2015

Kintzer, A.F., Green, E.M., Dominik, P.K., Bridges, M., Armache, J.-P., Deneka, D., Kim, S.S., Hubbell, W., Kossiakoff, A.A., Cheng, Y. & Stroud, R.M. (2018) Structural basis for activation of voltage sensor domains in an ion channel TPC1. *Proceedings of the National*

Academy of Sciences of the United States of America, **115**, E9095–E9104. doi:10.1073/pnas.1805651115

Kirmiz, M., Palacio, S., Thapa, P., King, A.N., Sack, J.T. & Trimmer, J.S. (2018) Remodeling neuronal ER-PM junctions is a conserved nonconducting function of Kv2 plasma membrane ion channels. *Molecular Biology of the Cell*, **29**, 2410–2432. doi:10.1091/mbc.E18-05-0337

Kirmiz, M., Vierra, N.C., Palacio, S. & Trimmer, J.S. (2018) Identification of VAPA and VAPB as Kv2 Channel-Interacting Proteins Defining Endoplasmic Reticulum–Plasma Membrane Junctions in Mammalian Brain Neurons. *The Journal of Neuroscience*, **38**, 7562–7584. doi:10.1523/JNEUROSCI.0893-18.2018

Lee, H.C., Wang, J.M. & Swartz, K.J. (2003) Interaction between extracellular Hanatoxin and the resting conformation of the voltage-sensor paddle in Kv channels. *Neuron*, **40**, 527–36. doi:10.1016/S0896-6273(03)00636-6

Lewis, G.N. (1925) A New Principle of Equilibrium. *Proceedings of the National Academy of Sciences of the United States of America*, **11**, 179–83, National Academy of Sciences. doi:10.1073/pnas.11.3.179

Li, H., Guo, W., Xu, H., Hood, R., Benedict, A.T. & Nerbonne, J.M. (2001) Functional expression of a GFP-tagged Kv1.5 alpha-subunit in mouse ventricle. *American Journal of Physiology. Heart and Circulatory Physiology*, **281**, H1955-67. doi:10.1152/ajpheart.2001.281.5.H1955

Lin, M.Z. & Schnitzer, M.J. (2016) Genetically encoded indicators of neuronal activity. *Nature Neuroscience*, **19**, 1142–1153. doi:10.1038/nn.4359

Liu, P.W. & Bean, B.P. (2014) Kv2 channel regulation of action potential repolarization and firing patterns in superior cervical ganglion neurons and hippocampal CA1 pyramidal neurons. *The Journal of Neuroscience*, **34**, 4991–5002. doi:10.1523/JNEUROSCI.1925-13.2014

Long, S.B., Campbell, E.B. & Mackinnon, R. (2005) Crystal structure of a mammalian voltage-dependent Shaker family K+ channel. *Science (New York, N.Y.)*, **309**, 897–903. doi:10.1126/science.1116269

Long, S.B., Tao, X., Campbell, E.B. & MacKinnon, R. (2007) Atomic structure of a voltage-dependent K+ channel in a lipid membrane-like environment. *Nature*, **450**, 376–82. doi:10.1038/nature06265

MacDonald, P.E., Salapatek, A.M.F. & Wheeler, M.B. (2003) Temperature and redox state dependence of native Kv2.1 currents in rat pancreatic beta-cells. *The Journal of Physiology*, **546**, 647–53. doi:10.1113/jphysiol.2002.035709

Maffie, J.K., Dvoretskova, E., Bougis, P.E., Martin-Eauclaire, M.-F. & Rudy, B. (2013) Dipeptidyl-peptidase-like-proteins confer high sensitivity to the scorpion toxin AmmTX3 to Kv4-mediated A-type K+ channels. *The Journal of Physiology*, **591**, 2419–27. doi:10.1113/jphysiol.2012.248831

Mandikian, D., Bocksteins, E., Parajuli, L.K., Bishop, H.I., Cerda, O., Shigemoto, R. & Trimmer, J.S. (2014) Cell type-specific spatial and functional coupling between mammalian brain Kv2.1 K+ channels and ryanodine receptors. *The Journal of Comparative Neurology*, **522**, 3555–74. doi:10.1002/cne.23641

Mann, V.R., Powers, A.S., Tilley, D.C., Sack, J.T. & Cohen, B.E. (2018) Azide-Alkyne Click Conjugation on Quantum Dots by Selective Copper Coordination. *ACS Nano*, **12**, 4469–4477. doi:10.1021/acsnano.8b00575

McCormack, K., Santos, S., Chapman, M.L., Krafte, D.S., Marron, B.E., West, C.W., Krambis, M.J., Antonio, B.M., Zellmer, S.G., Printzenhoff, D., Padilla, K.M., Lin, Z., Wagoner, P.K., Swain, N.A., Stupple, P.A., Groot, M. de, Butt, R.P. & Castle, N.A. (2013) Voltage sensor interaction site for selective small molecule inhibitors of voltage-gated sodium channels. *Proceedings of the National Academy of Sciences of the United States of America*, **110**, E2724–E2732. doi:10.1073/pnas.1220844110

McDonough, S.I., Mintz, I.M. & Bean, B.P. (1997) Alteration of P-type calcium channel gating by the spider toxin omega-Aga-IVA. *Biophysical Journal*, **72**, 2117–28. doi:10.1016/S0006-3495(97)78854-4

Milescu, M., Bosmans, F., Lee, S., Alabi, A.A., Kim, J. Il & Swartz, K.J. (2009) Interactions between lipids and voltage sensor paddles detected with tarantula toxins. *Nature structural & molecular biology*, **16**, 1080–5. doi:10.1038/nsmb.1679

Misonou, H., Menegola, M., Mohapatra, D.P., Guy, L.K., Park, K.-S. & Trimmer, J.S. (2006) Bidirectional Activity-Dependent Regulation of Neuronal Ion Channel Phosphorylation. *Journal of Neuroscience*, **26**, 13505–13514. doi:10.1523/JNEUROSCI.3970-06.2006

Misonou, H., Mohapatra, D.P., Menegola, M. & Trimmer, J.S. (2005) Calcium- and metabolic state-dependent modulation of the voltage-dependent Kv2.1 channel regulates neuronal excitability in response to ischemia. *The Journal of Neuroscience*, **25**, 11184–93. doi:10.1523/JNEUROSCI.3370-05.2005

Murakoshi, H., Shi, G., Scannevin, R.H. & Trimmer, J.S. (1997) Phosphorylation of the Kv2.1 K+ channel alters voltage-dependent activation. *Molecular Pharmacology*, **52**, 821–8. doi:10.1124/mol.52.5.821

O'Connell, K.M.S., Loftus, R. & Tamkun, M.M. (2010) Localization-dependent activity of the Kv2.1 delayed-rectifier K+ channel. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 12351–12356. doi:10.1073/pnas.1003028107

Opitz-Araya, X. & Barria, A. (2011) Organotypic hippocampal slice cultures. *Journal of Visualized Experiments*, MyJoVE Corporation. doi:10.3791/2462

Peltola, M.A., Kuja-Panula, J., Lauri, S.E., Taira, T. & Rauvala, H. (2011) AMIGO is an auxiliary subunit of the Kv2.1 potassium channel. *EMBO Reports*, **12**, 1293–1299. doi:10.1038/embor.2011.204

Peretz, A., Pell, L., Gofman, Y., Haitin, Y., Shamgar, L., Patrich, E., Kornilov, P., Gourgy-Hacohen, O., Ben-Tal, N. & Attali, B. (2010) Targeting the voltage sensor of Kv7.2 voltage-gated K+ channels with a new gating-modifier. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 15637–42. doi:10.1073/pnas.0911294107

Plant, L.D., Dowdell, E.J., Dementieva, I.S., Marks, J.D. & Goldstein, S.A.N. (2011) SUMO modification of cell surface Kv2.1 potassium channels regulates the activity of rat hippocampal neurons. *The Journal of General Physiology*, **137**, 441–54. doi:10.1085/jgp.201110604

Pologruto, T.A., Sabatini, B.L. & Svoboda, K. (2003) ScanImage: flexible software for operating laser scanning microscopes. *Biomedical Engineering Online*, **2**, 13. doi:10.1186/1475-925X-2-13

Redman, P.T., He, K., Hartnett, K.A., Jefferson, B.S., Hu, L., Rosenberg, P.A., Levitan, E.S. & Aizenman, E. (2007) Apoptotic surge of potassium currents is mediated by p38 phosphorylation of Kv2.1. *Proceedings of the National Academy of Sciences of the United States of America*, **104**, 3568–73. doi:10.1073/pnas.0610159104

Sack, J.T., Aldrich, R.W. & Gilly, W.F. (2004) A gastropod toxin selectively slows early transitions in the Shaker K channel's activation pathway. *The Journal of General Physiology*, **123**, 685–96. doi:10.1085/jgp.200409047

Sack, J.T., Stephanopoulos, N., Austin, D.C., Francis, M.B. & Trimmer, J.S. (2013) Antibody-guided photoablation of voltage-gated potassium currents. *The Journal of General Physiology*, **142**, 315–24. doi:10.1085/jgp.201311023

Schmalhofer, W.A., Calhoun, J., Burrows, R., Bailey, T., Kohler, M.G., Weinglass, A.B., Kaczorowski, G.J., Garcia, M.L., Koltzenburg, M. & Priest, B.T. (2008) ProTx-II, a selective inhibitor of NaV1.7 sodium channels, blocks action potential propagation in nociceptors. *Molecular Pharmacology*, **74**, 1476–84. doi:10.1124/mol.108.047670

Schneider, C.A., Rasband, W.S. & Eliceiri, K.W. (2012) NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, **9**, 671–5. doi:10.1038/nmeth.2089

Schneider, M.F. & Chandler, W.K. (1973) Voltage dependent charge movement of skeletal muscle: a possible step in excitation-contraction coupling. *Nature*, **242**, 244–6. doi:10.1038/242244a0

Scholle, A., Dugarmaa, S., Zimmer, T., Leonhardt, M., Koopmann, R., Engeland, B., Pongs, O. & Benndorf, K. (2004) Rate-limiting reactions determining different activation kinetics of Kv1.2 and Kv2.1 channels. *The Journal of Membrane Biology*, **198**, 103–12. doi:10.1007/s00232-004-0664-0

Seoh, S.A., Sigg, D., Papazian, D.M. & Bezanilla, F. (1996) Voltage-sensing residues in the S2 and S4 segments of the Shaker K+ channel. *Neuron*, **16**, 1159–67. doi:10.1016/S0896-6273(00)80142-7

Shi, G., Kleinklaus, A.K., Marrion, N. V & Trimmer, J.S. (1994) Properties of Kv2.1 K+ channels expressed in transfected mammalian cells. *The Journal of Biological Chemistry*, **269**, 23204–11. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/8083226

Shi, G., Nakahira, K., Hammond, S., Rhodes, K.J., Schechter, L.E. & Trimmer, J.S. (1996) Beta subunits promote K+ channel surface expression through effects early in biosynthesis. *Neuron*, **16**, 843–52. doi:10.1016/S0896-6273(00)80104-X

Shibata, R., Misonou, H., Campomanes, C.R., Anderson, A.E., Schrader, L.A., Doliveira, L.C., Carroll, K.I., Sweatt, J.D., Rhodes, K.J. & Trimmer, J.S. (2003) A fundamental role for KChIPs in determining the molecular properties and trafficking of Kv4.2 potassium channels. *The Journal of Biological Chemistry*, **278**, 36445–54. doi:10.1074/jbc.M306142200

Shimony, E., Sun, T., Kolmakova-Partensky, L. & Miller, C. (1994) Engineering a uniquely reactive thiol into a cysteine-rich peptide. *Protein engineering*, **7**, 503–7. doi:10.1093/protein/7.4.503

Speca, D.J., Ogata, G., Mandikian, D., Bishop, H.I., Wiler, S.W., Eum, K., Wenzel, H.J., Doisy, E.T., Matt, L., Campi, K.L., Golub, M.S., Nerbonne, J.M., Hell, J.W., Trainor, B.C., Sack, J.T., Schwartzkroin, P.A. & Trimmer, J.S. (2014) Deletion of the Kv2.1 delayed rectifier potassium channel leads to neuronal and behavioral hyperexcitability. *Genes, Brain, and Behavior*, **13**, 394–408. doi:10.1111/gbb.12120

Stoppini, L., Buchs, P.A. & Muller, D. (1991) A simple method for organotypic cultures of nervous tissue. *Journal of Neuroscience Methods*, **37**, 173–82. doi:10.1016/0165-0270(91)90128-M

Swartz, K.J. (2007) Tarantula toxins interacting with voltage sensors in potassium channels. *Toxicon*, **49**, 213–30. doi:10.1016/j.toxicon.2006.09.024

Tao, X., Lee, A., Limapichat, W., Dougherty, D.A. & MacKinnon, R. (2010) A Gating Charge Transfer Center in Voltage Sensors. *Science*, **328**, 67–73. doi:10.1126/science.1185954

Tilley, D.C., Angueyra, J.M., Eum, K.S., Kim, H., Chao, L.H., Peng, A.W. & Sack, J.T. (2019) The tarantula toxin GxTx detains K+ channel gating charges in their resting conformation. *The Journal of General Physiology*, **151**, 292–315. doi:10.1085/jgp.201812213

Tilley, D.C., Eum, K.S., Fletcher-Taylor, S., Austin, D.C., Dupre, C., Patron, L.A., Garcia, R.L., Lam, K., Yarov-Yarovoy, V., Cohen, B.E. & Sack, J.T. (2014) Chemoselective tarantula toxins report voltage activation of wild-type ion channels in live cells. *Proceedings of the National Academy of Sciences of the United States of America*, **111**, E4789-96. doi:10.1073/pnas.1406876111

Trapani, J.G. & Korn, S.J. (2003) Control of ion channel expression for patch clamp recordings using an inducible expression system in mammalian cell lines. *BMC Neuroscience*, **4**, 15. doi:10.1186/1471-2202-4-15

Trimmer, J.S. (1991) Immunological identification and characterization of a delayed rectifier K+ channel polypeptide in rat brain. *Proceedings of the National Academy of Sciences of the United States of America*, **88**, 10764–8. doi:10.1073/pnas.88.23.10764

Tsvetanova, N.G., Irannejad, R. & Zastrow, M. von. (2015) G Protein-coupled Receptor (GPCR) Signaling via Heterotrimeric G Proteins from Endosomes. *Journal of Biological Chemistry*, **290**, 6689–6696. doi:10.1074/jbc.R114.617951

Wisedchaisri, G., Tonggu, L., McCord, E., Gamal El-Din, T.M., Wang, L., Zheng, N. & Catterall, W.A. (2019) Resting-State Structure and Gating Mechanism of a Voltage-Gated Sodium Channel. *Cell*, **178**, 993-1003.e12. doi:10.1016/j.cell.2019.06.031

Woods, G. & Zito, K. (2008) Preparation of gene gun bullets and biolistic transfection of neurons in slice culture. *Journal of Visualized Experiments*, pii:675. doi:10.3791/675

Xia, X.M., Ding, J.P. & Lingle, C.J. (1999) Molecular basis for the inactivation of Ca2+- and voltage-dependent BK channels in adrenal chromaffin cells and rat insulinoma tumor cells. *The Journal of Neuroscience*, **19**, 5255–64. doi:10.1523/jneurosci.19-13-05255.1999

Xu, H., Li, T., Rohou, A., Arthur, C.P., Tzakoniati, F., Wong, E., Estevez, A., Kugel, C., Franke, Y., Chen, J., Ciferri, C., Hackos, D.H., Koth, C.M. & Payandeh, J. (2019) Structural Basis of Nav1.7 Inhibition by a Gating-Modifier Spider Toxin. *Cell*, **176**, 1238–1239. doi:10.1016/j.cell.2019.01.047

Yang, F. & Zheng, J. (2014) High temperature sensitivity is intrinsic to voltage-gated potassium channels. *eLife*, **3**, e03255. doi:10.7554/eLife.03255

Yang, W. & Yuste, R. (2017) In vivo imaging of neural activity. *Nature Methods*, **14**, 349–359. doi:10.1038/nmeth.4230

Zagotta, W.N., Hoshi, T., Dittman, J. & Aldrich, R.W. (1994) Shaker potassium channel gating. II: Transitions in the activation pathway. *The Journal of General Physiology*, **103**, 279–319. doi:10.1085/jgp.103.2.279

Zhang, A.H., Sharma, G., Undheim, E.A.B., Jia, X. & Mobli, M. (2018) A complicated complex: Ion channels, voltage sensing, cell membranes and peptide inhibitors. *Neuroscience Letters*, **679**, 35–47. doi:10.1016/j.neulet.2018.04.030

Zhang, G., Zheng, S., Liu, H. & Chen, P.R. (2015) Illuminating biological processes through site-specific protein labeling. *Chemical Society Reviews*, **44**, 3405–3417. doi:10.1039/C4CS00393D

Zheng, J. & Trudeau, M.C. (2016) *Handbook of Ion Channels*. (J. Zheng & M. C. Trudeau, Eds.), CRC Press.

Zito, K., Knott, G., Shepherd, G.M.G., Shenolikar, S. & Svoboda, K. (2004) Induction of Spine Growth and Synapse Formation by Regulation of the Spine Actin Cytoskeleton. *Neuron*, **44**, 321–334. doi:10.1016/j.neuron.2004.09.022

Figure Legends

Figure 1: *GxTX-594 colocalizes with Kv2-GFP channels.*

A. Fluorescence from CHO cells transfected with Kv2.1-GFP (Top) or Kv2.2-GFP (Bottom) and labeled with GxTX-594. Optical sections were near the glass-adhered cell surface. Cells were incubated with 100 nM GxTX-594 for 5 minutes and rinsed before imaging. Fluorescence shown corresponds to emission of GFP (Left), Alexa Fluor 594 (Middle), or an overlay of GFP and Alexa Fluor 594 channels (Right). Scale bars are 20 µm.

B. Pearson correlation coefficient for GxTX-594 colocalization with Kv2.1-GFP or Kv2.2-GFP.

C. Ratio of magenta to green intensity between GxTX-594 and Kv2-GFP. Circles indicate measurements from individual cells. In panels B and C, pixel intensities were background subtracted before analyses by subtracting the average fluorescence of a background ROI that did not contain cells, from the ROI containing the cell. All data from the same application of GxTX-594. Bars are arithmetic mean.

Figure 1 Supplement 1: *Synthesis of GxTX-594.*

A. Molecular model of GxTX-594. Scale bar is 10 angstroms.

B. HPLC chromatogram of Ser13Cys GxTX. Gradient described in *Methods* Ser13Cys GxTX eluted at 12.6 minutes, peak 1, which corresponds to 33% acetonitrile. MALDI-TOF MS profile of peak 1 (Inset). Rel. Int. is Relative Intensity.

C. HPLC chromatogram of GxTX-594 conjugation reaction between Alexa Fluor 594-maleimide and Ser13Cys GxTX. Peak 1 is Ser13Cys GxTX (Retention time: 12.8 minutes, 33% acetonitrile), peak 2 is a minor product from conjugation, and peak 3 is GxTX-594, the major product from conjugation (Retention time: 16.4 minutes, 35% acetonitrile). The fractions corresponding to peak 3 were combined.

D. HPLC chromatogram of the combined peak 3 fractions from panel C, a GxTX-594 preparation used in this study. 2 μL of 13.1 μM GxTX-594 diluted in 200 μL of 0.1 % TFA was injected. MALDI-TOF MS profile of the combined peak 3 fractions (Inset).

Figure 2: *GxTX594 binds Kv2.1 in the presence of its auxiliary subunit AMIGO-1.*

A. Fluorescence from Kv2.1-CHO cells, some of which have been transfected with AMIGO-1-YFP. Imaging plane was near the glass-adhered cell surface. Cells were incubated in indicated concentrations of GxTX-594 for 15 minutes then washed out before imaging. Fluorescence shown corresponds to YFP (Green) and Alexa Fluor 594 (Magenta). Scale bar is 20 µm. Green line surrounds a representative AMIGO-1 positive ROI. Blue line surrounds an AMIGO-1 negative ROI.

B. Pearson correlation coefficient between AMIGO-1-YFP and GxTX-594 within AMIGO positive ROIs. Coefficients from 1000 nM dose of GxTX-594. n = 8 ROIs total over 3 independent experiments. Bar is arithmetic mean.

C. Time-lapse GxTX-594 fluorescence intensity from the dose-response experiment shown in A. The green trace represents the AMIGO-1-YFP positive ROI, the blue trace represents the AMIGO-1-YFP negative ROI. Fluorescence was not background-subtracted. F_{max} is intensity while cells were incubated in 1000 nM GxTX-594.

D. Changes in fluorescence were analyzed at each concentration from multiple dose response experiments. Hollow markers correspond to 3 independent experiments. Green represents AMIGO-1-YFP positive ROIs, blue represents AMIGO-1-YFP negative ROIs. Lines are fits of a Langmuir isotherm (Hill equation with slope held at 1) to obtain a dissociation constant (K_d) for GxTX-594; AMIGO-1-YFP positive K_d = 26.7 nM ± 14.3; AMIGO-1-YFP negative K_d = 26.9 nM ± 8.3.

Figure 3: ***GxTX-594 labeling responds to transmembrane voltage.***

A. Fluorescence intensities from an optical section of a voltage-clamped Kv2.1- CHO cell in 9 nM GxTX-594. Pseudocolored images indicate maximal intensities in white. Rows indicate voltage step taken from a holding potential of -80 mV. Columns indicate time after voltage step. Scale bar is 10 μm.

B. GxTX-594 fluorescence during steps to indicated voltages. Smooth lines are fits of monoexponential (Eq. 1): -40 mV $k_{\Delta F}$ = 2.2 x 10^{-2} ± 2.2 x 10^{-3} s^{-1}: 0 mV $k_{\Delta F}$ = 1.3 x 10^{-1} ± 2.3 x 10^{-3} s^{-1}, 40 mV $k_{\Delta F}$ = 2.6 x 10^{-1} ± 6.2 x 10^{-3} s^{-1} and 80 mV $k_{\Delta F}$ = 5.3 x 10^{-1} ± 1.1 x 10^{-2} s^{-1}. ROIs were hand-drawn around the apparent cell surface membrane based on GxTX-594 fluorescence. Background for subtraction was the average intensity of a region that did not contain cells over the time course of the voltage protocol. Each trace was normalized to initial fluorescence intensity before the application of the voltage stimulus.

C. Fluorescence intensity remaining at the end of 50 s steps to +80 mV. Each circle represents one cell. Background subtraction and normalization as in panel B.

D. Voltage-dependence of fluorescence intensity at the end of 50 s steps. Data normalized to 100% for -80 mV and 0% for +80 mV. Circle coloring indicates data from the same cell. Gray circles represent data shown in B. Black line is fit of a first-order Boltzmann equation as in Tilley et al., 2019: $V_{1/2}$ = -27.4 ± 2.5 mV, z = -1.38 ± 0.13 e_0. Blue line is prediction from Scheme I at 9 nM GxTX. Bars are arithmetic mean ± standard error.

E. Voltage dependence of fluorescence intensity kinetics at end of 50 s steps. Circle coloring is the same as panel D. Black line is a first-order Boltzmann equation fit to the logarithm of $k_{\Delta F}$: $V_{1/2}$ = 2.3 ± 7.9 mV, z = -1.14 ± 0.38 e_0. Blue line is prediction from Scheme I at 9 nM GxTX. Bars are geometric mean ± standard error.

Figure 3 Supplement 1: *Spatial dependence of channel labeling kinetics at the glass-adhered surface.*

We conducted experiments to determine the region of voltage-clamped cells from which to quantify dynamic fluorescence. The confocal imaging plane containing the most GxTX-594 fluorescence was the glass-adhered basal surface. We noticed that GxTX-594 fluorescence in the center of the basal surface appeared to respond more slowly to voltage changes than the periphery. To quantify this observation, concentric ROIs were drawn in the center of the basal surface of the cell and their average fluorescence intensities were compared to each other and an ROI at the cell periphery. In response to voltage change, the kinetics of fluorescence change were slower at the center of the basal surface and faster towards the outer periphery. We quantified rates of fluorescence change ($k_{\Delta F}$) by fitting a monoexponential function (Eq 1). $k_{\Delta F}$ progressively slowed as the ROI contracted from the perimeter of the cell. This difference in kinetics suggests that the concentration of freely diffusing GxTX-594 in the restricted space between the cell membrane and the glass surface is transiently non-uniform after voltage change. The location dependence of $k_{\Delta F}$ was more pronounced during GxTX-594 labeling at -80 mV than unlabeling at +40 mV. We suspect that the more extreme location dependence at -80 mV is due to a high density of Kv2.1 binding sites in the restricted extracellular space between the cell membrane and glass coverslip, such that GxTX-594 is depleted from solution by binding Kv2.1 before reaching the center of the cell. After unbinding at +40 mV, each GxTX-594 molecule is expected to be less likely to rebind to Kv2.1 proteins as it diffuses out from under the cell, because Kv2.1 voltage sensors are in an activated, low affinity conformation (Tilley *et al.* 2014, 2019). This expected difference in GxTX-594 buffering by Kv2.1 could explain the difference in location dependence of $k_{\Delta F}$ during labeling at -80 mV and unlabeling at +40 mV. To avoid complications resulting from the spatial variability of kinetics at the basal surface, further voltage-dependent labeling experiments were conducted while imaging at a plane above the coverslip, where the cell surface has unrestricted access to the extracellular solution.

A. Time lapse airy disk images of the glass-adhered surface of a voltage-clamped Kv2.1-CHO cell in 9 nM GxTX-594. Time index is in the upper left of each panel and membrane potential is indicated in the upper right. Scale bar in lower right of last panel is 10 μm

B. Airy disk image of the glass adhered surface of a voltage-clamped Kv2.1-CHO cell in 9 nM GxTX-594. Grey lines indicate boundaries of ROIs. ROI 1, 2, and 3 are concentric circles each with a respective diameter of 1.8, 4.9, and 9.1 μm. ROI 4 was hand-drawn to contain the apparent cell surface. In all cells analyzed, ROI 1-3 were concentric circles of the same sizes, while ROI 4 varied based on cell shape. Scale bar is 10 μm.

C. Representative traces of GxTX-594 fluorescence intensity response to voltage changes. Red lines are monoexponential fits (Eq. 1): 40 mV step ROI 1 $k_{\Delta F} = 4.3 \times 10^{-2} \pm 2.6 \times 10^{-3}$ s^{-1}, ROI 2 $k_{\Delta F} = 4.4 \times 10^{-2} \pm 1.6 \times 10^{-3}$ s^{-1}, ROI 3 $k_{\Delta F} = 5.7 \times 10^{-2} \pm 1.5 \times 10^{-3}$ s^{-1}, ROI 4 $k_{\Delta F} = 9.7 \times 10^{-2} \pm 3.3 \times 10^{-3}$ s^{-1}. -80 mV step ROI 1 $k_{\Delta F} = 4.3 \times 10^{-4} \pm 1.1 \times 10^{-5}$ s^{-1}, ROI 2 $k_{\Delta F} = 7.9 \times 10^{-4} \pm 5.3 \times 10^{-6}$ s^{-1}, ROI 3 $k_{\Delta F} = 2.8 \times 10^{-3} \pm 7.4 \times 10^{-6}$ s^{-1}, and ROI 4 $k_{\Delta F} = 5.5 \times 10^{-3} \pm 1.3 \times 10^{-4}$ s^{-1}. Background for subtraction was the average intensity of a region that did not contain cells over the time course of the voltage protocol. Each trace was normalized to initial fluorescence intensity before the application of the voltage stimulus.

D. Circles indicate $k_{\Delta F}$ at 40 mV from individual cells. Circle coloring indicates data from the same cell.

E. Circles indicate $k_{\Delta F}$ at -80 mV from individual cells. Circle coloring indicates data from the same cell.

Figure 3 Supplement 2: *Temperature dependence of GxTX-594 on Kv2.1 CHO cells.*

To assess the temperature dependence of GxTX-594 labeling, the cell bath solution was held at either 27°C or 37°C and stepped to 0 mV for a measurement of $k_{\Delta F}$. The fold change in $k_{\Delta F}$ over this 10 °C difference, or Q_{10}, was 3.8-fold. This Q_{10} predicts a 1.3-fold increase in $k_{\Delta F}$ between 27 and 29°C, the temperature range of expreiments reported in Fig 3 E. This 1.3-fold increase was less than cell-to-cell variability observed.

A. Representative traces of GxTX-594 fluorescence intensity response to voltage changes at 27 °C (black) and 37 °C (gray). Smooth lines are fits of a monoexponential function (Eq. 1): 27 °C $k_{\Delta F} = 2.8$ x $10^{-2} \pm 3.9$ x 10^{-3} s^{-1}; 37 °C $k_{\Delta F} = 7.5$ x $10^{-2} \pm 5.4$ x 10^{-3} s^{-1}. Background was determined by taking the average fluorescence of a region that did not contain cells over the time course of the voltage protocol. This average was subtracted from each experimental group. Traces were normalized to initial fluorescence intensity before the application of the voltage stimulus.

B. $k_{\Delta F}$ at 27 °C and 37 °C. The rate of fluorescence change was significantly faster at higher temperatures (Mann-Whitney p = 0.0005). From geometric means (Bars), a Q_{10} of 3.8 was calculated between 27 °C and 37 °C. Each circle represents one cell, n = 7 both groups.

Figure 4: *Frequency dependence of GxTX-594 response*

A. Fluorescence intensity on Kv2.1-expressing CHO cells incubated in 9 nM GxTX-594. GxTX-594 fluorescence before (Left), after 50 s of 200 Hz stimulus (Middle), and 100 s after the cell is returned to resting potential of -80 mV (Right). Note that in each panel the unpatched cell (Left cell in each panel) does not show a change in fluorescence. Scale bar is 10 μm. Patched cell is indicated by a white arrow.

B. Representative trace of GxTX-594 unlabeling at 200 Hz. The trace was fit using a monoexponential (Eq. 1). Fit is overlaid and is in red. The $k_{\Delta F}$ at 200 Hz was obtained from this fit. For 200 Hz $k_{\Delta F}$ = 2.23 x 10^{-1} ± 9.9 x 10^{-3} s^{-1}. ROIs were hand drawn capturing the cell membrane based on GxTX-594 fluorescence such that only the cell membrane was selected without selecting the dark intracellular region. Background was determined by taking the average fluorescence of a region that did not contain cells over the time course of the protocol. This average was subtracted from each experimental group. F_{init} is the initial fluorescence intensity before the application of the frequency stimulus.

C. Decrease in fluorescence with increasing stimulus frequency. Circles indicate F/F_{init} from individual cells when stepped to +40 mV at indicated frequencies from a holding potential of -80 mV. Data normalized to 100% for -80 mV before frequency pulse. Circle coloring indicates data from the same cell. Black line is fit of a first-order Boltzmann equation as in Tilley et al., 2019: $V_{1/2}$ = -0.35 ± 0.03 mV, z = 51.23 ± 5.19 e_o. Blue line represents decreases in fluorescence at increasing voltages derived from the model shown in Scheme I at a concentration of 9 nM. Bars are arithmetic mean ± standard error.

D. Increases in $k_{\Delta F}$ with increasing stimulus frequency. Rates were obtained from fitting decreases in fluorescence with a monoexponential (Eq. 1) as shown in B. Blue line represents the rates of fluorescence change at increasing frequencies derived from the model shown in Scheme I at a concentration of 9 nM. Bars are geometric mean ± standard error.

Figure 5: ***Relationship of GxTX-594 labeling to probability of channel activation.***

A. Scheme I model predictions of concentration- and voltage-dependence of cell surface fluorescence intensity. Bottom axis represents membrane voltage while the top axis represents the ratio of activated to resting voltage sensors for unlabeled Kv2.1. Dashed black line represents the probability that voltage sensors of unlabeled Kv2.1 are in their resting conformation.

B. Scheme I model predictions of concentration- and voltage-dependence of $k_{\Delta F}$. Colors correspond to panel A.

Figure 6: *GxTX-594 is selective for Kv2 channels.*

A. Fluorescence from live CHO cells transfected with Kv2.1-GFP, Kv2.2-GFP, Kv4.2-GFP + KChIP2, Kv1.5-GFP + Kvβ2, or BK-GFP (Indicated by row) and labeled with GxTX-594. Airy disk imaging was from a plane above the glass-adhered surface. Cells were incubated with 100 nM GxTX-594 and 5 μg/mL WGA-405 then rinsed before imaging. Fluorescence shown corresponds to emission of GFP (Column 1), Alexa Fluor 594 (Column 2), WGA-405 (Column 3), or an overlay of GFP, Alexa Fluor 594, and WGA-405 channels (Column 4). Scale bars are 20 μm.

B. GxTX-594:GFP fluorescence intensity ratios for different K^+ channel-GFP types. F_{init} is the average GxTX-594:Kv2.1-GFP ratio. Circles indicate measurements from individual cells. Only cells with obvious GFP expression were analyzed. For analysis ROIs were drawn around the cell membrane indicated by WGA-405 fluorescence. Pixel intensities were background-subtracted before analyses by subtracting the average fluorescence of a background ROI that did not contain cells from the ROI containing the cell; this occasionally resulted in ROIs with negative intensity. Kv2.1 n = 16, Kv2.2 n = 10, Kv4.2 n = 13, Kv1.5 n = 13, BK n =10; n indicates the number of individual cells analyzed during a single application of GxTX-594. Bars are arithmetic mean. Significant differences were observed between GxTX:GFP ratio for Kv2.1 or Kv2.2 and Kv1.5, Kv4.2, or BK by Mann-Whitney (p<0.0001).

C. Pearson correlation coefficients between GxTX-594 and GFP. Same cells as panel B. Significant differences were observed between correlation coefficients for Kv2.1 or Kv2.2 and Kv1.5, Kv4.2, or BK by Mann-Whitney (p<0.0001).

Figure 6 Supplement 1: *GFP tagged K$^+$ channels exhibit expected outward currents.*

A. Exemplar whole-cell voltage clamp recordings of CHO cells expressing Kv2.1-GFP, Kv2.2-GFP, Kv4.2-GFP, Kv1.5-GFP, or BK-GFP. Recordings shown are representative responses to 100 ms steps from -100 mV to -40, 0 and +40 mV.

B. Fluorescence from live CHO cells transfected with Kv2.1-GFP, Kv2.2-GFP, Kv4.2-GFP Kv1.5-GFP, or BK-GFP (Indicated by row) and labeled with GxTX-594. Confocal imaging plane was above the glass-adhered surface. Cells were incubated with 100 nM GxTX-594 and 5 µg/mL WGA-405 and rinsed before imaging. Fluorescence shown corresponds to emission of GFP (Column 1), Alexa Fluor 594 (Column 2), WGA-405 (Column 3), or an overlay of GFP, Alexa Fluor 594, and WGA-405 channels (Column 4). Scale bars are 20 µm.

C. Background-subtracted GxTX-594:GFP ratio for different GFP and K$^+$ channel subtypes normalized to the average GxTX-594:Kv2.1-GFP ratio. Analysis same as in Figure 6 B. Kv2.1 n = 11, Kv2.2 n = 9, Kv4.2 n = 13, Kv1.5 = 13, and BK n =12. Cells were compiled from two independent GxTX-594 applications, each circle corresponds to a cell. Significant differences were observed between GxTX:GFP ratio for Kv2.1 or Kv2.2 and Kv1.5, Kv4.2, or BK by Mann-Whitney (p<0.0001).

D. Pearson correlation coefficients between GxTX-594 and GFP. Unless otherwise noted analysis was same as in Figure 6 C. Same cells as panel C.

Figure 7: **GxTX-594 labels CA1 hippocampal pyramidal neurons transfected with Kv2.1-GFP.**

A. Two-photon excitation images of fluorescence from the soma and proximal dendrites of a rat CA1 hippocampal pyramidal neuron in a brain slice six days after transfection with Kv2.1-GFP (Left), labeled with 100 nM GxTX-594 (Middle) and overlay (Right). Image represents a z-projection of 20 optical sections. Scale bar 10 μm.

B. A single optical section of the two-photon excitation image shown in A. GxTX-594 labels both Kv2.1-GFP puncta from a transfected cell and apparent endogenous Kv2 channels from an untransfected cell in the same cultured slice (Right, arrow).

C. Pearson correlation coefficients from CA1 hippocampal neurons two, four, and six days after transfection with Kv2.1-GFP. Each circle represents a different neuron. Bars are arithmetic means.

Figure 7 Supplement 1: **GxTX-594 labels CA1 hippocampal pyramidal neurons transfected with Kv2.1-GFP.**

> Two-photon excitation images rat CA1 hippocampal pyramidal neurons in brain slices as in Figure 7 A. Kv2.1-GFP (Left), GxTX-594 (Middle) and overlay (Right). Scale bars 10 μm.

A. Two days after transfection with Kv2.1-GFP.

B. Four days after transfection with Kv2.1-GFP.

C. Six days after transfection with Kv2.1-GFP.

Figure 8: **GxTX-594 puncta on hippocampal CA1 neurons are sensitive to voltage stimulus**

A. Single two-photon excitation optical section of fluorescence from CA1 pyramidal neurons in a cultured hippocampal slice after incubation with 100 nM GxTX-594. Scale bar is 5 μm.

B. GxTX-594 fluorescence from a single two-photon excitation optical section before and during depolarization of a whole cell patch-clamped neuron. Text labels of holding potential (-70 mV or 0 mV) indicate approximate position of the patch-clamp pipette. Red arrows indicate stimulus-sensitive puncta, white arrows indicate stimulus-insensitive puncta. Left panel is the average fluorescence of the three frames at -70 mV before depolarization while the right panel is the average fluorescence of three frames after holding potential was stepped to 0 mV. Scale bar is 5 μm. Supplementary Movie 1 contains time-lapse images from this experiment.

C. ROIs used in analysis for panels D, E, and F. Same slice as panel B. ROI 1 contains the apparent plasma membrane of the cell body of the patch-clamped neuron, it was generated by drawing a path tracing the apparent plasma membrane and then expanding to an ROI containing 15 pixels on either side of the path (1.2 μm total width). ROI 2 contains the area 15-45 pixels outside the membrane path (1.2 μm total width). RO1 3 contains the area more than 45 pixels outside of the membrane path. ROI 0 contains the area more than 15 pixels inside the membrane path. Scale bar is 5 μm.

D. Fluorescence from each ROI shown in C. Squares represent ROI 0, circles represent ROI 1, upward triangles represent ROI 2 and upside-down triangles represent ROI 3. Background was defined as the mean fluorescence of ROI 0 during the experiment. F_{init} was defined as the mean fluorescence of ROI 1 during the first six frames, after subtraction of background. Dotted lines represent the average fluorescence of the first six frames of each ROI. The voltage protocol is shown above.

E. Change in fluorescence during a 0 mV step for different ROIs in multiple hippocampal slices. ROIs for each slice were defined by methods described in panel C. Circles represent mean fluorescence from six frames during 0 mV stimulus (F), normalized to mean fluorescence from same ROI in six frames before stimulus (F_{init}). Circle color is consistent between ROIs for each hippocampal slice, red circles are from the slice shown in panel D. Black bars are arithmetic mean ± standard error from 3 hippocampal slices.

F. Kinetics of fluorescence change from ROI 1 during a 0 mV step in multiple hippocampal slices. ROI 1 for each slice was defined by the method described in panel C. Lines are monoexponential fits (Eq.1). $k_{\Delta F} = 6.8 \times 10^{-2} \pm 2.6 \times 10^{-2}$ s^{-1} (yellow), $6.8 \times 10^{-2} \pm 4.6 \times 10^{-2}$ s^{-1} (red), and $4.9 \times 10^{-2} \pm 2.1 \times 10^{-2}$ s^{-1} (green). The voltage protocol is shown above. Colors of individual circles indicate the same slices as panel E.

G. Region of interest used in analysis for panels H, I, and J. Same image as panel C. The shaded ROI was generated by drawing a path tracing the apparent plasma membrane and then expanding to an ROI containing 5 pixels on either side of the path (0.4 μm total width). Numbers indicate puncta that appear as peaks in panel H. Scale bar is 5 μm.

H. A plot of the fluorescence intensity along the ROI shown in panel G before (red trace) and during (black trace) 0 mV stimulus. Numbers above peaks correspond to puncta labeled in panel G. Red trace: mean fluorescence intensity during the three frames immediately before the stimulus, normalized to mean intensity of entire ROI (black dotted line), plotted against distance along path. Black trace: mean fluorescence intensity during three frames at 0 mV, normalized by the same F_{init} value as the red trace.

I. Kinetics of fluorescence change of individual puncta from panel G. Puncta intensity are average intensity of points extending to half maxima of each peak in panel H. Asterisks indicate mean fluorescence intensity of puncta 1 (pink), 4 (red), and 5 (dark red). Lines are monoexponential fits (Eq.1). $k_{\Delta F} = 7.1 \times 10^{-2} \pm 2.9 \times 10^{-2}$ s^{-1} (pink), $6.7 \times 10^{-2} \pm 2.5 \times 10^{-2}$ s^{-1} (red) and $1.2 \times 10^{-1} \pm 5.6 \times 10^{-2}$ s^{-1} (dark red). Fits to other puncta had standard deviations larger than $k_{\Delta F}$ values, and were excluded. F_{init} was defined as the mean background subtracted fluorescence of the puncta during the six frames before stimuli. The voltage protocol is displayed above. Blue line is prediction from Scheme I.

J. Comparison of fluorescence change of puncta and inter-puncta regions in response to 0 mV stimulus. The regions before stimulus shown in the red line of panel H that had $F/F_{init} \geq 1$ (above average) were binned separately from regions with $F/F_{init} < 1$. The mean fluorescence of each region during 0 mV stimulus (panel H black trace) was compared to the fluorescence before stimulus (panel H red trace). Circles indicate values from three independent hippocampal slices, colors indicate same slices as panel E and F. Black bars are arithmetic mean ± standard error.

K. Rate of fluorescence change of GxTX-594 after a 0 mV stimulus; from puncta (as in panel I), ROI 1 (as in panel F) or Kv2.1-CHO cells in 100 nM GxTX-594. Kv2.1-CHO cells were imaged at the same temperature as neurons (30 °C) using neuronal intracellular solution. Normal CHO extracellular solution (CE) was used for Kv2.1-CHO cell experiments. Kv2.1-CHO measurements were made by airy disk confocal imaging. Black bars are mean ± SEM from data shown. Dashed blue line is $k_{\Delta F} = 6.31 \times 10^{-2}$ s^{-1} prediction of Scheme I.

Table 1

Parameters used for calculations with Scheme I Model

Parameter	Value
$k_{on,resting}$	$0.30\ \mu M^{-1}\ s^{-1}$
$k_{off,resting}$	$0.0081\ s^{-1}$
$k_{on,activated}$	$0.21\ \mu M^{-1}s^{-1}$
$k_{off,activated}$	$0.39\ s^{-1}$
$V_{1/2,unlabeled}$	$-32\ mV$
$V_{1/2,labeled}$	$41\ mV$
z	$1.5\ e_0$

Figure 1

Figure 1 Supplement 1

Figure 2

A AMIGO-1

GxTX-594

AMIGO-1 (blue label in 1000nM panel)
+ AMIGO-1 (green label in 1000nM panel)

0nM   1nM   10nM   100nM   1000nM

B

Correlation coefficient

1.0
0.8
0.6
0.4
0.2
0.0

C

1nM 10 100 1000

F / Fmax

100%

- AMIGO-1
+ AMIGO-1

0%

0 20 40 60 80 100
Time / min

D

F / Fmax

100%

- AMIGO-1
+ AMIGO-1

0%

1 10 100 1000
GxTX-594 / nM

Figure 3

Figure 3 Supplement 1

A

0 s — -80mV | 50 s — 40mV | 130 s — -80mV | 540 s — -80mV

B

C

40mV
-80mV

F / F_{init}

100%

40%

0 100 200 300 400 500

Time / sec

D 40mV unlabeling

$k_{\Delta F} / s^{-1}$

1 2 3 4
Center ⟶ Periphery

E -80mV labeling

$k_{\Delta F} / s^{-1}$

1 2 3 4
Center ⟶ Periphery

Figure 3 Supplement 2

Figure 4

Figure 5

Figure 6

A

B

C

Figure 7

Figure 7 Supplement 1

Figure 8