Adverse effects of *Bacillus thuringiensis* bioinsecticide on non-target *Drosophila* species

Aurélie **Babin**, Marie-Paule **Nawrot-Esposito**, Armel **Gallet**, Jean-Luc **Gatti**[$*], Marylène **Poirié**[$]

Université Côte d'Azur, INRA, CNRS, ISA, France

*Corresponding author

Institut Sophia Agrobiotech, INRA, 400 route des chappes, 06903 Sophia Antipolis, France.

E-mail : jean-luc.gatti@inra.fr

[$]**Last co-authors**

Running title: Unintentional impact of *Bt* on *Drosophila* species

Keywords: *Drosophila*, *Bacillus thuringiensis*, biopesticide, toxicity, non-intentional effects,

development, fitness, longevity, offspring number, ecotoxicology

Word count (main text): 6 584

Figures: 7 figures and 1 table

Supplementary information

Abstract

Biopesticides based on *Bacillus thuringiensis kurstaki* (*Btk*) and *israelensis* (*Bti*) spores and toxins are widely used to control insect pests, increasing environmental risks to non-target biodiversity. Here, we tested for potential effects of larval ingestion of *Bt* commercial formulations on *Drosophila* species. Doses equivalent to those recommended for field application ($\leqslant 10^6$ CFU/g of fly medium) had no effect whereas *Btk* doses 10 to 100-fold higher (10^7-10^8 CFU/g) altered the development (decreased emergence due to larval mortality and increased development time), and moderately influenced adult fitness-related traits. At the highest *Btk* and *Bti* dose (10^9 CFU/g), all larvae died before pupation. The impact of *Btk* formulations resulted from the spores/cleaved toxins synergy, but also additives. While recommended doses had no effect on non-target *Drosophila* species, the accumulation of *Bt* bioinsecticides in the environment could have adverse side-effects on the populations of these species and therefore their associated communities.

Introduction

The world's population is expected to reach more than 9 billion people by 2050 (United Nations, 2015), increasing the demand for agricultural resources in the future. Increasing agricultural production requires improved management of pests, especially insects that cause more than 30% of losses (Pimentel and Burgess, 2014). Nowadays, their management still largely relies on conventional chemical insecticides. However, their use and efficiency have been considerably reduced due to the emergence of pests' resistance, development of secondary pests, adverse side-effects on non-target species (natural enemies of pests, pollinators) (Devine and Furlong, 2007; Sanchis and Bourguet, 2008), and more generally the impacts on biodiversity and human health (e.g. neurological disorders, functional impairment of reproduction, cancers) (WHO Report, 2007; Baldi et al. 2013; Gilden et al. 2016; Rizzati et al. 2016). Developed as an alternative, biopesticides are considered more specific and safer for the environment and human health. Today, they still represent less than 5% of the pesticide market, the large majority being microbial insecticide formulations based on viable spores and toxins of *Bacillus thuringiensis* (*Bt*) (over 400 registered formulations) (Sanchis and Bourguet, 2008; Lacey et al. 2015).

Bt is a Gram-positive endospore-forming bacterium that synthesizes a wide range of toxins with different chemical structures, modes of action and biological targets. The most abundant and studied are Cry δ-endotoxins encoded by genes located on large plasmids and produced as parasporal crystalline inclusions during the stationary growth phase (Crickmore 2017, Adang et al. 2014). *Bt* produces other insecticidal toxins, the Cyt (cytolytic δ-endotoxins) and Vip (secreted Vegetative Insecticidal Proteins) that synergize their effects with Cry toxins, virulence factors such as β-exotoxins (or thuringiensin), a secreted nucleotide toxic for almost all tested life forms thus prohibited in commercial formulations (WHO Report, 1999), and anti-fungal factors (Bravo et al. 2017; Rabinovitch et al. 2017). *Bt* subspecies and strains can differ in their plasmid number and in the synthesized toxins cocktail responsible for their biological activity, which was used to delineate potential target insects (Palma et al. 2014). For instance, *Bt* subsp. *kurstaki* (*Btk*) harbours 11

different plasmids and produces the 5 Cry toxins, Cry1Aa, Cry1Ab, Cry1Ac, Cry2Aa and Cry2Ab

(Ben-Dov et al. 1997; Adang et al. 2014), while *Bt* subsp. israelensis (*Bti*) produces a combination

of Cry4Aa, Cry4Ba, Cry10Aa, and Cry11Aa (Ben-Dov et al. 1999; Berry et al. 2002). The different

toxin cocktails produced by some *Bt* subspecies can also be harmful to non-insect organisms such

as nematodes and protozoa (Palma et al. 2014).

The formulations based on spores and toxin crystals of *Btk* and *Bti* are the most sprayed in

organic farming and natural areas (e.g. forests, swamps) to deal with larvae of Lepidopteran pests

and Dipteran larvae of mosquitoes and black flies, respectively. It is generally accepted that once

ingested by insect larvae, the toxin crystals are dissolved by the midgut alkaline pH, releasing ~130

kDa pro-toxins that are then processed by digestive proteases into smaller, soluble, active toxin

fragments of ~ 60-70 kDa (Bravo et al. 2007; Wei et al. 2016). *Active toxins* bind to specific

receptors of midgut epithelial cells, eliciting pores formation in the cell membrane, cell lysis and

gut epithelium disorganization (Schnepf et al. 1998; Bravo et al. 2011). This allows gut bacteria,

including *Bt*, to colonize the hemocoel, and leads to rapid septicaemia and death (Obata et al. 2009;

Caccia et al. 2016).

The increasing use of *Bt* has recently raised concern about its potential impact on non-target

species. Numerous impact studies of field application rates and acute intoxications have concluded

that *Bt* is safe or has a limited impact on non-target vertebrates and invertebrates, and associated

species communities (Glare and O'Callaghan, 2000). Yet, there is growing evidence of direct and

indirect cross-effects of *Bt* formulations and toxins across insect species and orders, or even across

phyla, suggesting that *Bt* targeting is only partly specific (van Frankenhuyzen 2017; Venter and

Bøhn, 2016). In addition, data showed that almost all of the *Btk* was still present on the leaves

surface 72 hours after spraying (Bizzarri and Bishop, 2008), its amount returning close to

environmental levels only 28 days after treatment (Raymond et al. 2010). Finally, *Bt* spores can

survive in the soil and different supports for months and even years after application (Hendriksen et

al. 2002; Duchet et al. 2014; Hung et al. 2016a, b; Enger et al. 2018). *Bt* formulations contain also

numerous compounds to protect spores and crystals and aggregate them into a wettable form, surfactants to facilitate spraying and dispersion on plants, and phagostimulants (Couch, 2000; Brar et al. 2006). Nevertheless, spores and toxins are somewhat sensitive to biotic and abiotic conditions (e.g. UV, pH, rainfall), which requires frequent applications to achieve the required pest control level (Brar et al. 2006). All this can lead to *Bt* accumulation in the environment, thus raising the rarely addressed issue of potential side-effects of chronic exposure (*i.e.* continuous and increasing exposure for an extended period) of non-target species to doses unexpectedly above the recommended application rates.

Diptera are worldwide distributed insects, most of which are not targets for *Bt* and its toxins. This is the case of the genus *Drosophila*, represented by ~ 1500 described species (Bächli, 1999-2008), including the model organism *D. melanogaster*. In the field, most of these flies feed and reproduce mainly on ripening or rotting fruits and are therefore present in areas treated with *Bt* such as orchards, vineyards and gardening areas. Unable to disperse between food patches, early stages of *Drosophila* larvae eat intensively and grow exponentially (Tennessen and Thumel, 2011), and may thus ingest high doses of *Bt* bioinsecticides. Surprisingly, few studies have focused on *Drosophila* species (Benz and Perron, 1967; Saadoun et al. 2001; Khyami-Horani 2002; Obeidat 2008; Obeidat et al. 2012; Cossentine et al. 2016; Biganski et al. 2017; Haller et al. 2017) and most of them showed susceptibility of these species to *Btk*. However, definitive conclusions were difficult to draw since most of these studies used mainly late 3rd instar larvae preparing for pupation, *i.e.* when they feed much less than younger larvae, and the tested *Bt* preparations possibly contain highly toxic β-exotoxins, especially in the case of field isolates.

Here, we have tested the dose-dependent chronic side-effects of different commercial formulations of *Btk* (devoid of β-exotoxins) and, to a lesser extent of *Bti*, on the wild-type *D. melanogaster* Canton S, with a focus on developmental traits (developmental time, emergence rate). The spore-forming Gram-positive *Bacillus subtilis* and the *Btk* strain (4D22), devoid of Cry toxin genes and thus of crystals, were used as non-pathogenic controls. We also analysed two fitness-

related traits of adult flies (male and female longevity, offspring number) after entire development

in presence of *Btk* formulation. Finally, we extended these experiments to several *D. melanogaster*

strains and *Drosophila* species to explore the potential implications in terms of competition and

associated communities.

Results

***Btk* formulations adversely impact the development of *D. melanogaster*.**

The wild-type Canton S strain of *D. melanogaster* was used to evaluate the dose-dependent effect

of the commercial *Btk* formulation Delfin[®] A on the emergence rate (ER, proportion of emerged

flies from the initial egg pool) and developmental time (DT, mean number of days from egg to adult

emergence). Eggs were transferred on a standard low-protein/high-sugar fly medium containing

Delfin[®] A at doses ranging from 5×10^5 CFU/g of medium (mean equivalent of the maximum

recommended doses for field application; see Methods and Supplementary information S1) to 10^9

CFU/g (~ 1,000 times the recommended dose). To check for specific effects of *Btk* formulations

and the respective role of *Btk* spores and Cry toxins, we tested the same dose range of the

commercial *Bti* formulation Vectobac[®] targeting mosquitoes that contains different Cry toxins

(Bravo et al. 2011), of the Cry-free strain *Btk* 4D22, and of the *Drosophila* non-pathogenic spore-

forming *Bacillus subtilis*.

Developmental traits (ER and DT) of exposed and non-exposed control flies were similar at

doses up to 10^7 CFU/g of Delfin[®] A (Fig. 1a-b; Table 1). At higher doses, both ER and DT were

affected in a dose-dependent manner: ER was reduced by 17% at 5×10^7 CFU/g (although not

statistically significant), up to 100% at 10^9 CFU/g, dose at which no individual reached the pupal

stage. The lethal dose 50 (LD50) was estimated between 5×10^7 and 10^8 CFU/g (Fig. 1a). DT was

increased of about 0.5 day at 5×10^7 CFU/g (+4% compared to controls), up to 1.5 days (+14%) at

10^8 CFU/g (Fig. 1b; Table 1). The sex-ratio at emergence (SR, proportion of males) was strongly

biased towards males at the highest dose at which complete development occurred (10^8 CFU/g),

with 58% more males compared to controls (Supplementary information S2). Because addition of

Btk formulation could modify parameters of the fly medium and thus contribute to these effects, we

checked the pH of the dose-responses medium. The presence or dose of the formulation had no

effect (Supplementary information S4).

We observed no change in ER using the same dose range of the *Btk* Cry-free strain 4D22 (Fig.

1a, 1e; Table 1) and the non-pathogenic *Bacillus subtilis* (Fig. 1a, Table 1). Addition of *Bti*

Vectobac$^®$ did not affect ER up to 10^8 CFU/g but reduced it by 89% at 10^9 CFU/g (~2,000 times the

highest recommended dose for field application; Fig. 1a; Table 1; Supplementary information S1).

DT varied with the dose of *Btk* 4D22, mainly due to differences between doses other than the

control. DT increased by ~1.5 days at the highest dose of Vectobac$^®$ (Fig. 1b; Table 1) and showed

a similar trend with *B. subtilis* ($p = 0.06$; Fig. 1b; Table 1). None of these three treatments

influenced dramatically the SR, the slight decrease in male proportion for most of the Vectobac$^®$

doses being due to the higher average sex-ratio for the control dose compared to those for the two

other treatments (Supplementary information S2).

To test whether these effects are generic to *Btk* formulations, the fly development was

evaluated on two other formulations, Delfin$^®$ B (same brand) and Scutello DF (brand Dipel$^®$), at the

critical doses 10^8 and 10^9 CFU/g. As Delfin$^®$ A, these formulations contain spores and Cry toxins

such as Cry-1A as pro-toxins of ~130 kDa, activated toxins of ~60-70 kDa, but also as smaller

fragments (Wei et al. 2016; Fig. 1e, red asterisks). ER remained unchanged at 10^8 CFU/g whereas

no individual reached pupation at 10^9 CFU/g on Delfin$^®$ B and very few individuals reached the

adult stage on Scutello DF$^®$, DT being increased by more than 2 days (Fig. 1c-d; Table 1). No

significant bias in SR was observed for either formulation (Supplementary information S2).

Adverse effects of *Btk* formulation strongly impact the early development.

Larval stages were tested for their susceptibility to *Btk* formulation in two independent and

complementary dose-response tests of survival on Delfin$^®$ A, at doses ranging from 10^5 to 10^9

CFU/g of high protein/sugar free medium. We focused on the 1^{st} and 2^{nd} larval instars, during

which growth is exponential (Tennessen and Thummel 2011), so that larvae are most heavily

exposed to the bioinsecticide. In the first test, the cumulative survival was measured by counting

late 1st and 2nd instar larvae alive which have been exposed to Delfin[®] A from the egg stage.

Larval survival was not influenced at 10^7 CFU/g, whereas it decreased for both larval instars above

that dose to reach up to 37% mortality at 10^9 CFU/g (Fig. 2a). Reduced survival tended to occur at a

lower dose when cumulative survival was measured later in the development, *i.e.* 10^9 for late 1^{st}

instar larvae and 10^8 CFU/g for 2^{nd} instar larvae (Fig. 2a; Table 1). For both instars, larvae surviving

10^9 CFU/g were noticeably smaller and less active than those surviving lower doses. In emergence

assays with planned exposure from the egg to the adult stage, none of these individuals reached the

pupal stage (see results above). In the second test, larval survival was measured after early 1^{st} and

2^{nd} instar larvae had been exposed for 24 hours to Delfin[®] A. Survival of 1^{st} instar larvae decreased

by 36% on 10^9 CFU/g whereas that of 2^{nd} instar larvae did not change (Fig. 2b, Table 1).

Developmental exposure to *Btk* formulation does not strongly influence fitness-related traits

in adults.

Long-term consequences on flies of exposure to *Btk* formulation throughout the development were

evaluated on two fitness-related traits, longevity and total offspring number**.** Traits were measured

on a *Btk*-free low-protein/high-sugar medium after individuals had completed their development on

the same fly medium but in presence of selected doses of Delfin[®] A: $5×10^6$ CFU/g, which had no

impact on development, and $5×10^7$ and 10^8 CFU/g, which caused moderate and strong

developmental alterations, respectively (see Fig. 1a).

Adult longevity was analysed in two independent experiments on groups of 15 females and 15

males held together. Despite large variation between experiments (Table 1), the longevity of adults

reared on $5×10^6$ CFU/g of Delfin[®] A was similar to that of non-exposed controls (Fig. 3). Males

and females which developed on the two higher doses showed a moderate longevity benefit, higher

in females for 10^8 CFU/g (Fig. 3a-b, d-e; Table 1). Males generally survived better than females

(Table 1) but their longevity benefit of developing on 10^8 CFU/g was only observed in the second

experiment (Fig. 3b, e).

The female offspring number - the sum of offspring produced by the 15 females of each fly

group during the longevity experiment - varied depending on both the experiment and the Delfin® A

dose (Table 1). In the 1^{st} experiment, adults from larvae reared on 10^8 CFU/g had fewer offspring

compared to control adults and to adults developed on the other doses whereas the total offspring

number varied regardless of the *Btk* dose in the 2^{nd} experiment (Fig. 3c, f, Table 1).

Btk-formulation dose-dependent alterations of development are not specific to the _D._

melanogaster strain.

Dose-dependent effects of *Btk* formulation on the development were tested on three additional *D.*

melanogaster strains: the wild-type Nasrallah (strain 1333), the wild-type Sefra population reared in

the laboratory for 4 years, and the double mutant YW1118. The emergence rates (ER) and

developmental times (DT) were measured on a high-protein/sugar-free medium (rearing medium of

these strains) mixed with Delfin® A doses ranging from 10^5 to 10^9 CFU/g. To allow the comparison

with previous results with Canton S flies on low-protein/high sugar fly medium, Canton S was also

reared and tested on the high-protein/sugar-free medium along with the other strains.

None of the fly strains was impacted at doses up to 10^7 CFU/g, whereas ER was strongly reduced

and DT was increased at higher doses for all the strains (Fig. 4a-b, Table 1), with no individual

reaching the pupal stage at 10^9 CFU/g (LD50 between 10^8 and 10^9 CFU/g). At 10^8 CFU/g, the

magnitude of effects on Canton S flies was lower than that observed on the low-protein/high-sugar

medium. At this dose, ER varied between strains, the largest reduction being observed for Sefra

(Table 1). We observed no dose-dependent bias in SR (Supplementary information S3).

Btk formulation also affects other _Drosophila_ species.

The ER and DT were analysed for seven other *Drosophila* species from different phylogenetic clades at doses of Delfin® A from 10^5 to 10^9 CFU/g of high-protein/sugar-free medium (rearing medium of all the species). Tested species were *D. simulans* (*D. melanogaster* sister species), the African *D. yakuba*, *D. subobscura*, *D. immigrans, D. hydei*, and the invasive *D. suzukii*, all belonging to the *Drosophila* subgenus, and *D. busckii* from the *Dorsilopha* subgenus. For all the species, doses up to 10^6 CFU/g of Delfin® A had no effect on ER and DT whereas all individuals failed to reach the pupal stage and no fly emerged at 10^9 CFU/g (Fig. 5-6). Amplitudes of development alterations at 10^7 and 10^8 CFU/g varied between species (Fig. 5-6; Table 1). All species were affected at 10^8 CFU/g as was *D. melanogaster* (see Fig. 4a for comparison). *D. simulans* and *D. busckii* had unchanged ER, but DT was slightly increased for *D. simulans* (although slightly reduced at 10^7 CFU/g; similar results with a Japanese strain, data not shown) and strongly increased for *D. busckii* (by 20%, *i.e.* ~ 4 days) (Fig. 5-6, Table 1). *D. yakuba* ER and DT were similar to those of *D. melanogaster*, with an LD50 around 10^8 CFU/g and a moderate DT increase of ~ 1 day (Fig. 5-6, Table 1; similar results with a strain from Sweden, data not shown). The ER of *D. hydei* and *D. subobscura* were very low at 10^8 CFU/g (LD50 below this dose), with a high DT (Fig. 5-6; Table 1), while *D. immigrans* did not survive. No *D. suzukii* individual emerged at 10^8 CFU/g and development was already moderately impacted at 10^7 CFU/g (Fig. 5-6). No dose-dependent bias in SR was detected for either species (Supplementary information S5).

Development alterations may result from a synergy between formulation components.

Bt spores and toxins represent about 30% of the weight of commercial formulations, with up to about 10% of insecticidal protein toxins, mainly Cry pro-toxins and activated toxins (see Fig. 1e) (Koch et al. 2015). The remaining weight consists of various compounds such as residues of culture medium and various additives including surfactant, anti-foaming agents, etc. (Glare and O'Callaghan, 2000; Brar et al. 2006). Since other compounds of formulations can be more harmful in some cases than the active ingredient (Bradberry et al. 2004), we explored the role of small

diffusible molecular weight components of Delfin® A in the alterations of ER and DT of *D. melanogaster* Canton S. For that, we mixed a 10 kDa dialyzed suspension of Delfin® A at 10^7, 10^8, and 10^9 CFU/g with low-protein/high-sugar medium. ER and DT were unaffected by the presence of the dialyzed suspension from the 10^7 CFU/g dose, whereas no individual reached the adult stage (no pupation) with the suspension from the 10^9 CFU/g dose (Fig. 7a; Table 1). At 10^8 CFU/g, ER was not modified but DT increased by ~ 1 day, only in one of the two experiments, partially reproducing the changes observed without dialysis (Fig. 7a-b; see also Fig. 1a-b, Table 1; 3 independent experiments for ER, 2 independent experiments for DT).

Cry1A profiles of dialyzed Delfin® A suspensions, like those of the non-dialyzed ones, comprised 130-kDa pro-toxins and 60-70 kDa activated toxins, but also showed toxin degradation as evidenced by additional smaller fragments of activated toxins (Fig. 7c). The respective roles of *Btk* toxin fragments and spores in the alterations of *D. melanogaster* development were further explored through experiments of dialysis followed by successive centrifugations to eliminate most of the spores and toxin crystals. Despite variation between experiments, ER was strongly affected only in one of the three experiments while DT was always significantly increased when flies were reared in presence of centrifuged supernatants that contained a limited range of Cry 1A toxin fragments (Supplementary information S6).

Discussion

The increasing use of bioinsecticides based on *Bacillus thuringiensis* (*Bt*) raised concern about potential side-effects on non-target biodiversity because of their partial specific targeting (de Souza Machado et al. 2017; van Frankenhuyzen 2017; Venter and Bohn 2016), persistence in the environment (Duchet et al. 2014; Hung et al. 2016a, b), and requirement of repeated spraying to reach the desired pest control level (Brar et al. 2006). Especially, side-effects of chronic exposure on non-target biodiversity, including insects present on treated areas, remain under-evaluated. Here, we tested the side-effects of ingestion of *Bt* formulations (mainly *Bt kurstaki* (*Btk*) but also *Bt*

israelensis (*Bti*)) throughout the entire development of several non-target species of *Drosophila* flies which are naturally present in treated areas. While formulation doses up to those recommended for field sprayings ($\leq 10^6$ CFU/g of medium) had no effect on *Drosophila* development, mortality and/or developmental delay occurred from doses only 10 times and 50 times higher than the maximum recommended dose of the main tested *Btk* formulation for *D. suzukii* (10^7 CFU/g) and the *D. melanogaster* strains (5×10^7 CFU/g), respectively. Besides, all the tested species except *D. simulans* were strongly affected at 10^8 CFU/g, and no (or extremely limited) fly development occurred at the highest tested dose (10^9 CFU/g), equivalent to 1000 times the maximum recommended dose but below common acute intoxication doses (WHO Report, 2007). Recommended doses are single-spraying doses on a homogeneous and dry zone without covering areas. In the field, both repeated spraying of stabilized formulation and rainfall washouts can increase *Bt* spores and toxins presence in both space and time. While the highest dose tested here would hardly be reached in the field, the minimal doses at which flies development was impacted may be readily obtained. Furthermore, the minimal quantity of *Bt* formulation inducing developmental alterations may be even lower since a single *Drosophila* larva is unlikely to process 1g of medium given its size and feeding rate. Our data also evidence a window of susceptibility to *Btk* during the larval development, ingestion during the 1st larval instar being responsible for a large part of the observed detrimental effects on the development.

When testing for generic effects of *Bt* formulations, slightly different results were observed with two other *Btk* formulations and a formulation of *Bti*: there was no effect on *D. melanogaster* development at the doses up to 10^8 CFU/g but a strong detrimental effect at the highest dose tested, 10^9 CFU/g. All the *Btk* formulations, based on two different bacterial strains (see Methods), had similar profiles of Cry1A protoxins and activated toxins, but they differed in their efficient spore contents, formulation type, and likely additives, which may account for the observed variation in the half-lethal dose. The *Bti* formulation, widely used against Dipteran Nematoceran insects (e.g. mosquitoes, black flies; Becker 2000), impacted *D. melanogaster* development only at the highest

dose tested. These impacts of *Bt* formulations on *D. melanogaster* development are consistent with growing evidence suggesting a partly specific targeting of *Bt* (van Frankenhuyzen 2013; Venter and Bøhn, 2016). Until recently, it has generally been accepted that the mode of action of *Bt* after ingestion by insects relies on key steps of specific binding of proteolyzed *Bt* toxins to receptors of midgut epithelial cells, defining targets for each *Bt* subspecies (Bravo et al. 2007, 2011; Palma et al. 2014). Several primary and secondary types of toxin receptors, including cadherin-like proteins, aminopeptidases, GPI-anchored alkaline phosphatases (Adang et al. 2014), and more recently the ATP dependent binding cassette reporter C2 (Stevens et al. 2017), have been identified in Lepidoptera and Dipteran mosquitoes. Focusing on the action of *Btk* targeting Lepidoptera, no Lepidoptera Cry receptor orthologues were found in *Drosophila* (Stevens et al. 2017), supporting the idea that these flies would not be affected by the spraying of *Btk* formulation. However, the existence of orthologues of other types of Cry receptors in *Drosophila* flies remains unknown. In addition, the substantial amounts of active Cry1A toxin fragments in *Btk* formulations could compensate for the possible lack of solubilization of protoxin crystals in the fly midgut and proteolytic activation of toxins by fly gut proteases, both required for Cry activity in insect larvae (Bravo et al. 2007). Other toxins synthesized by *Btk* and present in the formulations could also play a role in the observed cross-order activity as some, such as Cry2A, have an insecticidal effect on both Lepidoptera and Diptera (George and Crickmore 2012).

The lack of effect of ingestion of *Bacillus subtilis* or *Btk* Cry-free 4D22 on the development of *D. melanogaster* excludes that developmental alterations result from severe disruption of digestion and nutrient uptake/competition in the presence of high spore/bacteria loads in the larval gut throughout development. It supports the idea of a synergistic action of *Btk* spores and Cry toxins, consistent with the models of *Bt* action on insect larvae in which toxins first breach the gut epithelium, allowing the gut content, including *Bt* spores, to colonize the hemocoel (Bravo et al. 2007; Obata et al. 2009; Bravo et al. 2011; Caccia et al. 2016). The partially reproduced mortality rate and delayed development in dialysis experiments further indicate that low diffusible molecular

weight compounds in *Btk* formulations (e.g., culture media residues, salts, additives) may contribute to these developmental alterations. This is supported by the lack of impact on *D. melanogaster* development of the ingestion of spores and Cry toxins of a *Btk* 4D1 strain (or HD1, one of the strains used in commercial *Btk* formulations, but here produced in the laboratory) used without additives, even at the highest dose 10^9 CFU/g (additional information S7; Fig. S7a, b). The *Btk* 4D1 culture contained few active Cry toxins and smaller toxin fragments, in contrast to commercial *Btk* formulations (Fig. S7c), supporting the possible contribution of these toxin fragments to the cross-order activity of *Btk* formulations on *Drosophila*.

As observed for *D. suzukii* exposed to laboratory-produced *Btk* cultures (Cossentine et al. 2016), mortality of *D. melanogaster* during development on *Btk* formulation already occurred early in development. First and second instars larvae are probably highly exposed due to their high feeding rate and their exponential growth (Santos et al. 1997). As the observed larval mortality was only about 40% at the highest dose (10^9 CFU/g), while none of the individuals reached the pupal stage, the remaining mortality likely occurred during the third larval stage, maybe due to delayed action of *Btk* spores and toxins. Interestingly, alterations of the development (mortality and delayed emergence) mimicked those typically generated by nutritional stress conditions in insect larvae (Nepoux et al. 2010; Vantaux et al. 2016). Accordingly, the developmental alterations were partially rescued on a protein rich fly medium, probably through compensatory protein intake, as in other arthropod species (Moret and Schmid-Hempel 2000; Kutzer and Armitage 2016; Vantaux et al. 2016). In addition, the sex ratio of flies was strongly biased towards males after development on the dose of *Btk* formulation affecting fly emergence (10^8 CFU/g) and under low protein conditions. This highlights the importance of nutritional conditions in *Btk* impacts on development, with sex-specific differences in larval susceptibility to environmental stressors, here the accumulation of *Btk* formulation, under protein restriction conditions as previously reported in *D. melanogaster* (Andersen et al. 2010).

The development on sublethal doses of *Btk* formulation did not dramatically affect the

longevity of *D. melanogaster* adults and the offspring number throughout life. Developmental exposure to *Btk* doses that slightly and strongly reduced the likelihood of reaching the adult stage even gave males and females a dose-dependent longevity benefit, in addition to the male higher longevity observed in mixed-sex populations (Khazaeli and Curtsinger, 2000), and slightly increased the offspring number (although not significantly). Surviving the exposure to *Btk* formulation throughout the development has likely selected for fitter individuals. This is similar to the increased longevity of adult insects that have survived developmental nutritional stress (Rion and Kawecki, 2007, Burger et al. 2010), or are resistant to environmental stressors (Khazaeli and Curtsinger 2000).

The origin of *Drosophila* (species and population/strain) influenced the magnitude of the impacts of the *Btk* formulation on the development. Within the *D. melanogaster* species, all strains tested were susceptible to the *Btk* formulation with both mortality and delayed development at the same dose, but with variation in the effect magnitude. This suggests potential population-specific differences in susceptibility to *Btk* formulation accumulation in the environment, and hence potential spatial and temporal heterogeneity of *Btk* spraying impacts for each *Drosophila* species. At the fly community level, differences in susceptibility to *Btk* formulation, in terms of effect magnitude and type of developmental alteration (mortality and/or developmental delay) occurred between *Drosophila* species, regardless of their phylogenetic distances. In the *Drosophila* subgenus, *D. simulans* was less susceptible than its sister species *D. melanogaster*, whereas the African *D. yakuba* experienced similar impacts on the development as *D. melanogaster*. The three species *D. immigrans*, *D. subobscura* and *D. hydei* were similarly more susceptible than *D. melanogaster*, but with slight differences in effect magnitudes. The phylogenetically distant *D. busckii* (*Dorsilopha* subgenus) was the least affected of all the species tested in terms of developmental mortality, but its development was strongly delayed. The five species *D. melanogaster*, *D. simulans*, *D. hydei*, *D. immigrans*, and *D. busckii* belong to the guild of cosmopolitan domestic *Drosophila* species, *D. subobscura* is sub-cosmopolitan species, and *D.*

busckii is an opportunistic frugivorous species (Atkinson and Shorrocks 1977). All these species coexist frequently and compete on the same discrete and ephemeral rotting fruit patches, with seasonal variations in the composition of the fly community (Shorrocks 1991; Benado and Brncic 1994; Nunney 1996; Mitsui and Kimura 2000). Differences in species susceptibility to accumulation of *Btk* formulation could modify larval competition conditions and lead to additional local and temporal variations in *Drosophila* communities' composition. The potential side-effects of *Bt* sprays on non-target *Drosophila* communities would be hardly predictable as they depend on spatial patterns of *Bt* accumulation.

The *Btk* formulation clearly impacted the development of the invasive *D. suzukii*, as recently reported by Cossentine et al. (2016), this species being the most susceptible here with effects already detectable at 10 times the recommended spraying dose. Compared with the other seven species that live on rotten fruits, *D. suzukii* poses a threat to fruit production because it feeds and lays eggs on healthy ripening fruits (Walsh et al. 2011; Delbac et al. 2014; Poyet et al. 2014) and hence colonizes orchards and vineyards earlier during the fruit season. The greater susceptibility of *D. suzukii* to the accumulation of *Btk* formulation in the environment might mitigate the potential ecological burden of its invasion for local communities of *Drosophila* frugivorous species in orchards. Alternatively, as *D. suzukii* attacks on fruits can accelerate their decomposition by microorganisms, its higher susceptibility to *Btk* could reduce the number of fruits made suitable for other *Drosophila* species.

In conclusion, we show here that repeated spraying and accumulation of *Btk* formulation can potentially impact non-target insect communities, and the magnitude of this impact could depend on the formulation used and the insect species. Recent studies have reported similar adverse side-effects due to repeated spraying of the *Bti* formulation, directly on non-target organisms (e.g. Duguma et al. 2015), and indirectly on predators via food webs (e.g. Poulin et al. 2010). These studies and the data presented here highlight that pest control with *Bt* bioinsecticides should be done with caution in the field to avoid, or at least limit, potential negative impacts on non-target

biodiversity and species communities within ecosystems. At last, *D. melanogaster*, a model species

in many research fields, could also serve as a study model to assess the toxicity of *Bt* on non-target

species, and identify the mechanisms underlying these side-effects.

Methods

Commercial formulations, Bacillus *productions and Colony Forming Unit measurement*

The tested commercial brands of *Bacillus thuringiensis kurstaki* (*Btk*; serotype 3a, b, c; Zeigler,

1999) were Delfin® A and B (strain SA11; wettable granules, Valent BioSciences, AMM 9200482,

32,000 UI/mg) and Scutello DF (a Dipel® sub-brand; strain ABTS-351; wettable granules, Biobest®,

AMM 2010513, 540g/kg). The commercial brand of *Bacillus thuringiensis israelensis* (*Bti*; strain

HD-14; serotype 14; Ziegler, 1999) was VectoBac® WG (wettable granules, Bayer, AMM 2020029,

3000 UTI/mg). For each formulation, the number of viable spores (expressed as Colony Forming

Units (CFU) per mg of granules) was estimated using serial dilutions of a suspension on LB agar

plates and counting of bacterial colonies after overnight incubation at 30°C. CFU estimations were

5×10^7 CFU/mg for *Btk* Delfin® A; 2.5×10^7 CFU/mg for *Btk* Delfin® B; 2.2×10^7 CFU/mg for *Btk*

Scutello DF; 6×10^7 CFU/mg for *Bti* VectoBac®. No change in CFU estimations occurred during the

time frame of the experiments. Manufacturer-recommended doses for Delfin® range from 0.15 to

1.5 kg/ha depending on the crop type. Based on our CFU estimations, this corresponds to

recommended doses of 7.5×10^4 to 7.5×10^5 CFU/cm^2 of Delfin® A, and 3.75×10^4 to 3.75×10^5

CFU/cm^2 of Delfin® B for each spraying in the field. For Scutello DF, recommended doses range

from 0.1 to 1 kg/ha, equivalent to 2.2×10^4 to 2.2×10^5 CFU/cm^2. Vectobac® WG is used at 0.125 to

1 kg/ha, equivalent to 7.5×10^4 to 6×10^5 CFU/cm^2.

The acrystillipherous (Cry toxin-free) *Btk* 4D22 strain (depleted for the toxin-encoding

plasmids; Gonzalez et al. 1982) obtained from the Bacillus Genetic Stock Center (http://bgsc.org;

Columbus USA), and a *Drosophila* non-pathogenic *Bacillus subtilis* (gift from Dr. E. Bremer,

University of Marburg, Germany; A. Brun-Barale, pers. comm.) were grown at 30°C in the

sporulation-specific medium PGSM (Bactopeptone® 7.5 g, KH_2PO_4 3.4 g, K_2HPO_4 4.35 g, glucose 7.5 g, PGSM salts 5 mL, $CaCl2$ 0.25 M, distilled water qsp 1L, pH 7.2; PGSM salts: $MgSO_4.7H_2O$, $MnSO_4.H_2O$, $ZnSO_4.7H_2O$, $FeSO_4.7H_2O$) for about 14 days for sporulation to occur. Following elimination of vegetative cells (1h at 70 °C), spore pellets were collected after centrifugation (4,500 rpm, 20 min, 4 °C), washed with sterile water, and lyophilized. CFU numbers were counted for each preparation as described above.

Fly stocks

The four tested strains of *Drosophila melanogaster* (phylogenetic subgroup: melanogaster) were the standard wild-type Canton S (Bloomington Drosophila Centre) used as a reference strain, the wild-type Nasrallah strain from Tunisia (strain 1333, Gif-sur-Yvette), the double mutant standard strain YW1118 (white and yellow mutations; gift from Dr. B. Charroux, IBD, Marseille-Luminy)*,* and a recently field-collected strain (caught in Southern France in 2013) that we named "Sefra". For *Drosophila* species comparison, we included 6 species of the *Drosophila* subgenus, *D. simulans* (strain 1132; phylogenetic subgroup: melanogaster), *D. yakuba* (strain 1880; phylogenetic subgroup: melanogaster), *D. hydei* (phylogenetic subgroup: hydei) and *D. suzukii* (phylogenetic subgroup: immigrans) (both kindly provided by Dr. R. Allemand, LBBE, University Lyon 1), *D. immigrans* (phylogenetic subgroup: immigrans), *D. subobscura* (phylogenetic subgroup: obscura), and one species of the *Dorsilopha* subgenus, *D. busckii* (all three species collected in South-East of France in Spring 2015).

All strains and species were maintained at controlled densities (150-200 eggs/40 ml of fly medium) under standard laboratory conditions (25°C or 20°C for recently collected species, 60 % relative humidity, 12:12 light/dark cycle), on a high-protein/sugar-free fly medium (10 % cornmeal, 10 % yeast, 0 % sugar). The *D. melanogaster* Canton S strain was also reared on a standard low-protein/sugar-free fly medium (8 % cornmeal, 2 % yeast, 2.5 % sugar) to test for the influence of the medium composition on *Btk* exposure effects.

Intoxication method and dose-response assay

Commercial formulations and laboratory spore productions were suspended and diluted in buffer to perform dose-response assays with doses from 10^5 to 10^9 CFU/g of fly medium. All doses were prepared in 100 µl and homogenized thoroughly with the fly medium (100µl/g). *Drosophila* eggs and larvae were collected from stock vials at the suitable developmental stage and transferred carefully to the intoxication vials and dishes, then maintained under standard laboratory conditions until a) the emergence of adults, or, in the larvae survival tests, b) until a given developmental stage was reached from the egg, and c) for 24h. Control groups of individuals were transferred on fly medium homogenized with the same volume of buffer.

Development-related traits and larval survival

To evaluate emergence rates and developmental times upon intoxication throughout the entire development, precise numbers of eggs from mass oviposition were transferred to intoxication vials containing fly medium mixed with doses of *Bt* formulations or bacteria productions and let to develop under standard laboratory conditions until the fly emergence. Eggs without chorion and transparent eggs were discarded. The initial number of eggs was adjusted depending on the species biology and the vial size: 20 eggs for 2 g of fly medium in small vials (Ø 3.3 cm, surface ~8.5 cm^2, 0.24 g/cm^2) for tests with *D. melanogaster* Canton S, 50 eggs for 6 g of fly medium for comparison of *D. melanogaster* strains and *Drosophila* species in wider vials (Ø 4.6 cm, surface ~16 cm^2, 0.37 g/cm^2) except for *D. hydei*, *D. suzukii* and *D. immigrans* for which 30 eggs were transferred on 6 g of fly medium. Numbers and sex of emerging flies were recorded once a day until the day the pupae of the next generation should form. From these data, the emergence rate (proportion of emerged flies from the initial eggs; ER), the mean developmental time (mean number of days for completion of development; DT), and the sex-ratio (proportion of male flies; SR) were calculated for each intoxication vial.

For the larval survival tests, 20 eggs or larvae from a 4-hour mass oviposition at the indicated

developmental stage, were transferred to small dishes containing 1 g of fly medium (Ø 3 cm,

surface ~7 cm^2) homogenized with increasing doses of Delfin® A. Surviving larvae were counted at

the indicated developmental stage, or after 24-hour intoxication, to calculate the proportion of

surviving larvae. For the test from the egg, eggs which did not hatch were not included in the

counting. As a control measurement, we measured the pH of the fly medium in the presence of the

dose range of *Bt* formulations (see Supplementary Information S4).

Adult fitness-related traits

For the longevity and offspring number tests, males and females emerged from several rearing vials

for each dose of Delfin® A were pooled when aged 2 days. Groups of 15 males and 15 females were

transferred into vials with fresh fly medium without formulation. Fly medium was renewed every 3-

4 days. After each fly transfer to fresh food, discarded maintenance vials were incubated under

standard laboratory conditions for the offspring to develop. Mortality and sex of dead flies were

recorded daily until the last fly died. Offspring numbers were counted from the first emergence

until pupae of the next generation appeared. The tests were repeated twice. Due to the variation in

the duration of the two longevity experiments, offspring numbers of each vial were summed to

obtain a total offspring number per dose of Delfin® A for each experiment.

Dialysis and Cry toxin analysis

A suspension of 2×10^{10} CFU of Delfin® A was dialyzed against PBS (KH_2PO_4 1.06 mM,

$Na_2HPO_4(2H_2O)$ 3mM, NaCl 154 mM, qsp distilled water, pH 7.2), at 250 rpm, 4°C overnight,

using an 8-10 kDa MW cut-off membrane (ZelluTrans, Roth®). The CFUs of the dialyzed

suspension and the effects on ER and DT were analysed as described above. The dialyzed

suspension was also subject to a 12.5 % SDS-PAGE and compared to the non-dialyzed suspension

after silver staining. The presence of Cry1A pro-toxins, activated toxins and toxin fragments was

probed by Western-blot using an in-house anti-Cry1A rabbit polyclonal antibody.

Data analysis

Data on development traits (emergence rate ER and developmental time DT), sex-ratio (SR),

survival of larval stages and offspring number were analysed with mixed effect models including

the dose of *Btk* formulation/spore production, the *D. melanogaster* strain, the *Drosophila* species or

the developmental stage as fixed effects, and replicate (plus the experiment when necessary) as

random effects (for ER data, data were analysed with bias-corrected models with replicate as fixed

effect to allow pairwise comparisons; similar results obtained with models including replicate as

random effect). ER, SR and survival of larval stages were analysed with generalized linear models,

with binomial distribution and logit link. DT and offspring number were analysed with linear

models. DT were transformed into developmental rates (1/developmental time) to fulfil the

assumptions of the analysis of variance (homoscedasticity and residuals normality). Adult longevity

data were analysed with proportional hazard Cox regression models including fly sex and dose of

Btk formulation as fixed effects, and replicates as a random effect. For all the data sets, the main

fixed effects and their interactions were tested with log-likelihood ratio tests. *Post hoc* pairwise

comparisons were made for pairs of *D. melanogaster* strains, formulation/spore treatments, and

between the control dose and the other doses. All the analyses were performed in R (R

Development Core Team, 2008) using the packages lme4 (Bates et al. 2015), brglm (Kosmidis

2017), multcomp (Horton et al. 2008), survival (Terry et al. 2000) and coxme (Terry and Therneau,

2015).

References

Adang, M.J., Crickmore, N., Jurat-Fuentes, J.L. 2014. Diversity of *Bacillus thuringiensis* crystal

toxins and mechanism of action. In: Dhadialla, T.S. and Gill, S.S., *Advances in Insect Physiology*

vol. 47, Insect midgut and insecticidal proteins. Oxford Academic Press, pp. 39–87.

Andersen, L.H., Kristensen, T.N., Loeschcke, V., Toft, S., Mayntz, D. 2010. Protein and
carbohydrate composition of larval food affects tolerance to thermal stress and desiccation in
adult *Drosophila melanogaster*. *Journal of Insect Physiology*, 56: 336-340.

Atkinson, W., Shorrocks, B. 1977. Breeding site specificity in the domestic species of *Drosophila*.
Oecologia, 29(3): 223-232.

Bächli, G. 1999-2008. TaxoDros: the database on the taxonomy of Drosophilidae. Available at:
http://www.taxodros.uzh.ch.

Baldi, I., Cordier, S., Coumoul, X., Elbaz, A. 2013. Pesticides, effets sur la santé. Expertise
collective. INSERM. Available at: http://ipubli-inserm.inist.fr/handle/10608/1.

Bates, D., Maechler, M., Bolker, B., Walker, S. 2015. Fitting linear mixed-effects models using
lme4. *Journal of Statistical Software*, 67(1): 1-48.

Becker, N. 2000. Bacterial control of vector-mosquitoes and black flies. In: Charles, J.-F.,
Delécluse, A., Nielsen-Roux, C. (ed.). Entomopathogenic bacteria: from laboratory to field
application. Kluwer Academic Publishers. p. 383-398.

Ben-Dov, E., Zaritsky, A., Dahan, E., Barak, Z., Sinai, R., Manasherob, R., Khamraev, A.,
Troitskaya, E., Dubitsky, A., Berezina, N., Margalith, Y. 1997. Extended screening by PCR for
seven cry-group genes from field-collected strains of *Bacillus thuringiensis*. *Applied and
Environmental Microbiology*, 63(12): 4883–4890.

Ben-Dov, E., Nissan, G., Pelleg, N., Manasherob, R., Boussiba, S., Zaritsky, A. 1999. Refined,
circular restriction map of the *Bacillus thuringiensis* subsp. *israelensis* plasmid carrying the
mosquito larvicidal genes. *Plasmid*, 42(3): 186-191.

Benado, M., Brncic, D. 1994. An eight-year phenological study of a local drosophilid community in
Central Chile. *Journal of Zoological Systematics and Evolutionary Research*, 32: 51-63.

Benz, G., Perron, J.M. 1967. The toxic action of *Bacillus thuringiensis* "exotoxin" on *Drosophila
reared in yeast-containing and yeast-free media. *Experientia*, 23(10): 871–872.

Berry, C., O'Neil, S., Ben-Dov, E., Jones, A.F., Murphy, L., Quail, M.A., Holden, M.T.G., Harris,

D., Zaritsky, A., Parkhill, J. 2002. Complete sequence and organization of pBtoxis, the toxin-coding plasmid of *Bacillus thuringiensis* subsp. *israelensis*. *Applied and Environmental Microbiology*, 68(10): 5082-5095.

Biganski, S., Jehle, J.A., Kleepies, R.G. 2017. *Bacillus thuringiensis* serovar *israelensis* has no effect on *Drosophila suzukii* Matsumura. *Journal of Applied Entomology*, 142: 33-36.

Bizzarri, M.F., Bishop, A.H. 2008. The ecology of *Bacillus thuringiensis* on the phylloplane: colonization from soil, plasmid transfer, and interaction with larvae of *Pieris brassicae*. *Microbial Ecology*, 56(1): 133-139.

Bradberry, S.M., Proudfoot, A.T., Vale, J.A. 2004. Glyphosate poisoning. *Toxicological Reviews*, 23(3): 159-167. Brar, S.K., Verma, M., Tyagi, R.D., Valéro, J.R. 2006. Recent advances in downstream processing and formulations of *Bacillus thuringiensis* based biopesticides. *Process Biochemistry*, 41(2): 323–342.

Bravo, A., Gill, S.S., Soberon, M. 2007. Mode of action of *Bacillus thuringiensis* Cry and Cyt toxins and their potential for insect control. *Toxicon*, 49: 423-435.

Bravo, A., Likitvivatanavong, S., Gill, S.S., Soberon, M. 2011. *Bacillus thuringiensis*: a story of a successful bioinsecticide. *Insect Biochemistry and Molecular Biology*, 41(7): 423–431.

Bravo, A., Pacheco, S., Gomez, I., Garcia-Gomez B., Onofre, J., Soberon, M. 2017. Insecticidal proteins from *Bacillus thuringiensis* and their mechanism of action. In: Fiuza, L.M., Polanczyk, R.A., Crickmore, N. (eds). *Bacillus thuringiensis* and *Lysinibacillus sphaericus*, characterization and use in the field of biocontrol. pp. 53-66.

Burger, J.M.S., Buechel, S.D., Kawecki, T.J. 2010. Dietary restriction affects lifespan but not cognitive aging in *Drosophila melanogaster*. *Aging Cell*, 9: 327-335.

Caccia, S., Di Lelio, I., La Storia, A., Marinelli, A., Varricchio, P., Franzetti, E., Banyuls, N., Tettamanti, G., Casartelli, M., Giordana, B., Ferre, J., Gigliotti, S., Ercolini, D., Pennacchio, F. 2016. Midgut microbiota and host immunocompetence underlie *Bacillus thuringiensis* killing mechanism. *Proceedings of the National Academy of Sciences of the United States of America*,

113(34): 9486-9491.

Cossentine, J., Robertson, M., Xu, D. 2016. Biological activity of *Bacillus thuringiensis* in

Drosophila suzukii (Diptera: Drosophilidae). *Journal of Economic Entomology*, 0(0): 1-8.

Couch, T.L., 2000. Industrial fermentation and formulation of entomopathogenic bacteria. In:

Charles, J.-F., Delecluse, A., Nielsen-LeRoux, C. (Eds.), Entomopathogenic Bacteria: From

Laboratory to Field Application. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp.

297–316.

Crickmore, N. 2017. *Bacillus thuringiensis* toxin classification. In: Fiuza, L., Polanczyk, R.,

Crickmore, N. (eds) *Bacillus thuringiensis* and *Lysinibacillus sphaericus*. Springer.

de Souza Machado, A.A., Zarfl, C., Rehse, S., Kloas, W. 2017. Low-dose effects: nonmonotonic

responses for the toxicity of a *Bacillus thuringiensis* biocide to *Daphnia magna*. *Environmental*

Science & Technology, 51: 1679-1686.

Delbac, L., Rusch, A., Rouzes, R., Ravidat, M.-L., Launes, S., Thiéry, D. 2014. *Drosophila suzukii*

est-elle une menace pour la vigne? *Phytoma*, 679 : 16-21.

Devine, G.J., Furlong, M.J. 2007. Insecticide use: contexts and ecological consequences.

Agriculture and Human Values, 24(3): 281–306.

Duchet, C., Tetreau, G., Marie, A., Rey, D., Besnard, G., Perrin, Y., Paris, M., David, J.-P.,

Lagneau, C., Desprès, L. 2014. Persistence and recycling of bioinsecticidal *Bacillus*

thuringiensis subsp. *israelensis* spores in contrasting environments: evidence from field

monitoring and laboratory experiments. *Microbial Ecology*, 67(3): 576–586.

Duguma, D., Hall, M.W., Rugman-Jones, P., Stouthamer, R., Neufeld, J.D., Walton, W.E. 2015.

Microbial communities and nutrient dynamics in experimental microcosms are altered after the

application of a high dose of *Bti*. *Journal of Applied Ecology*, 52: 763-773.

Enger, K.S., Mitchell, J., Murali, B., Bridsell, D.N., Keim, P., Gurian, P.L., Wagner, D.M. 2018.

Evaluating the long-term persistence of *Bacillus* spores on common surfaces. *Microbial*

Biotechnology, 56: 3073, 12.

George, Z., Crickmore, N. 2012. *Bacillus thuringiensis* applications in agriculture. In: E.

Sansinenea (ed.). *Bacillus thuringiensis* biotechnology. p. 392.

Gilden, R.C., Huffling, K., Sattler, B. 2010. Pesticides and health risks. *Journal of Obstetric,*

Gynecologic & Neonatal Nursing, 39(1): 103–110.

Glare, T.R., O'Callaghan, M. 2000. *Bacillus thuringiensis*: Biology, ecology and safety. John Wiley

& Sons, UK. pp. 350.

Gonzales, J.M.Jr, Brown, B.J., Carlton, B.C. 1982. Transfer of *Bacillus thuringiensis* plasmids

coding for δ-endotoxin among strains of *B. thuringiensis* and *B. cereus*. *Proceedings of the*

National Academy of Sciences of the United States of America, 79: 6951-6955.

Haller, S., Romeis, J.X.R., Meissle, M., 2017. Effects of purified or plant-produced Cry proteins on

Drosophila melanogaster (Diptera: Drosophilidae) larvae. *Scientific Reports*, 1-11.

Hendriksen, N.B., Hansen, B.M. 2002. Long-term survival and germination of *Bacillus*

thuringiensis var. *kurstaki* in a field trial. *Canadian Journal of Microbiology*, 48(3): 256–261.

Horton, T., Bretz, F., Westfall, P. 2008. Simultaneous inference in general parametric models.

Biometrical Journal, 50(3): 346-363.

Hung, T.P., Truong, L.V., Binh, N.D., Frutos, R., Quiquampoix, H., Staunton, S. 2016. Persistence

of detectable insecticidal proteins from *Bacillus thuringiensis* (Cry) and toxicity after adsorption

on contrasting soils. *Environmental Pollution*, 208: 318–325.

Hung, T.P., Truong, L.V., Binh, N.D., Frutos, R., Quiquampoix, H., Staunton, S. 2016. Fate of

insecticidal *Bacillus thuringiensis* Cry protein in soil: differences between purified toxin and

biopesticide formulation. *Pest Management Science*, 72: 2247–2253.

Khazaeli, A.A., Curtsinger, J.W. 2000. Genetic analysis of extended lifespan in *Drosophila*

melanogaster III. On the relationship between artificially selected and wild stocks. *Genetica*, 109:

245-253.

Khyami-Horani, H. 2002. Toxicity of *Bacillus thuringiensis* and *B. sphaericus* to laboratory

populations of *Drosophila melanogaster* (Diptera: Drosophilidae). *Journal Basic Microbiology*,

42(2): 105-110.

Koch, M.S., Ward, J.M., Levine, S.L., Baum, J.A., Vicini, J.L., Hammond, B.G. 2015. The food

and environmental safety of *Bt* crops. *Frontiers in Plant Science*, 6: 283.

Kosmidis I (2017). *brglm: Bias Reduction in Binary-Response Generalized Linear Models*. R

package version 0.6.1, http://www.ucl.ac.uk/~ucakiko/software.html.

Kutzer, M.A., Armitage, S.A.O. 2016. The effect of diet and time after bacterial infection on

fecundity, resistance, and tolerance in *Drosophila melanogaster*. *Ecology and Evolution*, 6(13):

4229-4242.

Lacey, L.A., Grzywacz, D., Shapiro-Ilan, D.I., Frutos, R., Brownbridge, M., Goettel, M.S. 2015.

Insect pathogens as biological control agents: Back to the future. *Journal of Invertebrate*

Pathology, 132(C): 1–41.

Mitsui, H., Kimura, M.T. 2000. Coexistence of drosophilid flies: aggregation, patch size diversity

and parasitism. *Ecological Research*, 15: 93-100.

Moret, Y., Schmid-Hempel, P. 2000. Survival for immunity: the price of immune system activation

for bumblebee workers. *Science*, 290(5494): 1166-1168.

Nepoux, V., Haag, C.R., Kawecki, T.J. 2010. Effects of inbreeding on aversive learning in

Drosophila. *Journal of Evolutionary Biology*, 23: 2333-2345.

Nunney, L. 1996. The colonization of oranges by the cosmopolitan *Drosophila*. *Oecologia*, 108:

552-561.

Obata, F., Kitami, M., Inoue, Y., Atsumi, S., Yoshizawa, Y., Sato, R. 2009. Analysis of the region

for receptor binding and triggering of oligomerization on *Bacillus thuringiensis* Cry1Aa toxin.

FEBS Journal, 276: 5949-5959.

Obeidat, M. 2008. Toxicity of local *Bacillus thuringiensis* isolates against *Drosophila melanogaster*.

World Journal of Agricultural Sciences, 4(2): 161-167.

Obeidat, M., Khymani-Horani, H. and Al-Momani, F. 2012. Toxicity of *Bacillus thuringiensis* β-

exotoxins and δ-endotoxins to *Drosophila melanogaster*, *Ephestia kuhniella* and human

erythrocytes. *African Journal of Biotechnology*, 11(46):10504-10512.

Palma, L., Muñoz, D., Berry, C., Murillo, J., Caballero, P. 2014. *Bacillus thuringiensis* toxins: an overview of their biocidal activity. *Toxins*, 6(12): 3296–3325.

Pimentel, D., Burgess, M. 2014. Environmental and economic costs of the application of pesticides primarily in the United States. In: Integrated Pest Management. Dordrecht: Springer Netherlands. pp. 47–71.

Poulin, B., Lefebvre, G., Paz, L. 2010. Red flag for green spray: adverse trophic effects of *Bti* on breeding birds. *Journal of Applied Ecology*, 47: 884-889.

Poyet, M., Eslin, P., Héraude, M., Le Roux, V., Prévost, G., Givert, P., Chabrerie, O. 2014. Invasive host for invasive pest: when the Asiatic cherry fly (*Drosophila suzukii*) meets the American black cherry (*Prunus serotine*) in Europe. *Agricultural and Forest Entomology*, 1-9, DOI: 10.1111/afe012052.

R Development Core Team 2008. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL http://www.R-project.org.

Rabinovitch, L., Vivoni, A.M., Machado, V., Knaak, N., Berlitz, D.L., Polanczyk, R.A., Fiuza, L.M. 2017. *Bacillus thuringiensis* characterization: morphology, physiology, biochemistry, pathotype, cellular, and molecular aspects. In: Fiuza, L.M., Polanczyk, R.A., Crickmore, N. (eds). *Bacillus thuringiensis* and *Lysinibacillus sphaericus*, characterization and use in the field of biocontrol. pp. 1-18.

Raymond, B., Wyres, K.L., Sheppard, S.K., Ellis, R.J., Bonsall, M.B. 2010. Environmental factors determining the epidemiology and population genetic structure of the *Bacillus cereus* group in the field. *PloS Pathogens*, 6(5): e1000905.

Rion, S., Kawecki, T.J. 2007. Evolutionary biology of starvation resistance: what we have learned from *Drosophila*. *Journal of Evolutionary Biology*, 20(5): 1655-1664.

Rizzati, V., Briand, O., Guillou, H., Gamet-Payrastre, L. 2016. Effects of pesticide mixtures in

human and animal models: an update of the recent literature. *Chemico-Biological Interactions*,

254: 231–246.

Saadoun, I., Al-Moman, F., Obeidat, M., Meqdam, M., Elbetieha, A. 2001. Assessment of toxic

potential of local Jordanian *Bacillus thuringiensis* strains on *Drosophila melanogaster* and *Culex*

sp (Diptera). *Journal of Applied Microbiology*, 90: 866-872.

Sanchis, V., Bourguet, D. 2008. *Bacillus thuringiensis*: applications in agriculture and insect

resistance management. A review. *Agronomy for Sustainable Development*, 28(1): 11–20.

Santos, M., Borash, D.J., Joshi, A., Bounlutay, N., Mueller, L.D. 1997. Density-dependent natural

selection in *Drosophila*: evolution of growth rate and body size. *Evolution*, 51(2): 420-432.

Schnepf, E., Crickmore, N., Van Rie, J., Lereclus, D., Baum, J., Feitelson, J., Zeigler, D.R., Dean,

D.H. 1998. *Bacillus thuringiensis* and its pesticidal crystal proteins. *Microbiology and Molecular*

Biology Reviews, 62(3): 775–806.

Shorrocks, B. 1991. Competition on a divided and ephemeral resource: a cage experiment.

Biological Journal of the Linnean Society, 43: 211-220.

Stevens, T., Song, S., Bruning, J.B., Choo, A., Baxter, S.W. 2017. Expressing a moth abcc2 gene in

transgenic *Drosophila* causes susceptibility to Bt Cry1Ac without requiring a cadherin-like

protein receptor. *Insect Biochemistry and Molecular Biology*, 80: 61-70.

Tennessen, J.M., Thummel, C.S. 2011. Coordinating growth and maturation - Insights from

Drosophila. *Current Biology*, 21(18): R750–757.

Therneau, T.M. 2015. coxme: Mixed Effects Cox Models. R package version 2.2-5.

https://CRAN.R-project.org/package=coxme

Therneau, T.M., Grambsch, P.M. 2000. Modeling survival data: extending the Cox model. Springer,

New York, ISBN 0-387-98784-3.

United Nations, Department of Economic and Social Affairs, Population Division 2015. World

Population Prospects 2015 – Data Booklet (ST/ESA/ SER.A/377).

van Frankenhuyzen K. 2013. Cross-order and cross-phylum activity of *Bacillus thuringiensis*

pesticidal proteins. Journal of Invertebrate Pathology, 114(1): 76–85.

van Frankenhuyzen, K. 2017. Specificity and cross-order activity of *Bacillus thuringiensis*

pesticidal proteins. In: Fiuza, L.M., Polanczyk, R.A., Crickmore, N. (eds). *Bacillus thuringiensis*

and *Lysinibacillus sphaericus*, characterization and use in the field of biocontrol. pp. 127-172.

Vantaux, A., Ouattarra, I., Lefèvre, T., Dabiré, K.R. 2016. Effects of larvicidal and larval nutritional

stresses on *Anopheles gambiae* development, survival and competence for *Plasmodium*

falciparum. *Parasites & Vectors*, 9: 226.

Venter, H.J., Bøhn, T. 2016. Interactions between Bt crops and aquatic ecosystems: A review.

Environmental Toxicology and Chemistry, 35(12): 2891–2902.

Walsh, D.B., Bolda, M.P., Goodhue, R.E., Dreves, A.J., Lee, J., Bruck, D.J., Walton, V.M., O'Neal,

S.D., Zalom, F.G. 2011. *Drosophila suzukii* (Diptera: Drosophilidae): invasive pest of ripening

soft fruit expanding its geographic range and damage potential. *Journal of Integrated Pest*

Management, 2(1): DOI: 10.1603/IPM10010.

Wei, J., Liang, G., Wang, B., Zhong, F., Chen, L., Khaing, M.M., Zhang, J., Guo, Y., Wu, K.,

Tabashnik, B.E. 2016. Activation of Bt protoxin Cry1Ac in resistant and susceptible cotton

bollworm. *PLoS ONE*, 11(6): e0156560.

WHO report 1999. Guideline specification for bacterial larvicides for public health use. WHO

document WHO/CDS/CPC/WHOPES/99.2. World Health Organization, Geneva.

WHO report 2007. WHO specifications and evaluations for public health pesticides: *Bacillus*

thuringiensis subspecies israelensis strain AM65-52. World Health Organization, Geneva.

Zeigler, D.R. 1999. *Bacillus* genetic stock center catalog of strains, 7[th] edition. Part 2: *Bacillus*

thuringiensis and *Bacillus cereus*. pp. 58.

Acknowledgements

We thank Xiao Han, Jingru Li and Abir Oueslati for help with preliminary experiments, L.

Kremmer, C. Rebuf and O. Magliano for providing and rearing flies and help in preparing fly

medium, A. Brun-Barale for the *Bacillus subtilis* spores, D. Pauron for preparation of Cry1A toxin,

Hugo Mathé-Hubert for advice on statistical analyses, and M. Amichot for helpful discussions. The

Cry1A antibody was produced in collaboration with the INRA-PFIE platform (Nouzilly, France).

Financial Supports

This work was supported by the French National Agency for Research (ANR-13-CESA-0003-001

ImBio), the European Union's Seventh Framework Program for research, technological

development and demonstration under grant agreement No. 613678 (DROPSA), the "Investments

for the Future" LABEX SIGNALIFE (ANR-11-LABX-0028), and the INRA Plant Health

Department.

Author contributions

AB, MP NE, AG, JLG and MP designed the experiments. AB performed the experiments with

contributions of MPNE. AB performed the statistical analyses. AB, JLG, and MP wrote the

manuscript with contributions from all the authors.

Additional information

Supplementary information

Competing financial interest: The authors declare no competing financial interests.

Table 1. Results of statistical analyses to assess the effect of the dose of formulation/spore

production and its interaction with the treatment, the larval instar, the experiment, the sex, the fly

strain and the fly species when appropriate. See figures for *post hoc* comparisons of the doses with

the control dose.

Source of variation/Data	χ^2 / Deviance	d.f.	P value
Development on *Btk* Delfin® A, *Btk* 4D22, *Bti* Vectobac®, *Bacillus subtilis*			
Emergence rate			
Dose × Treatment	285.7	20	**< 0.0001**
Dose for each treatment:			
- Delfin® A	237.5	6	**< 0.0001**
- 4D22	7.0	7	0.40
- Vectobac®	165.8	5	**< 0.0001**
- *B. subtilis*	1.9	6	0.93
Developmental time			
Dose × Treatment	220.8	19	**< 0.0001**
Dose for each treatment:			
- Delfin® A	68.8	6	**< 0.0001**
- 4D22	16.08	7	**0.024**
- Vectobac®	37.5	6	**< 0.0001**
- *B. subtilis*	13.5	7	0.060
Development on *Btk* Delfin® B and Scutello DF (dose effect)			
Emergence rate			
- Delfin® B	151.2	2	**< 0.0001**
- Scutello DF	105.1	2	**< 0.0001**
Developmental time			
- Delfin® B	2.5	1	0.12
- Scutello DF	30.9	2	**< 0.0001**
Role of formulation components in the development alterations (dialysis)			
Dose effect			
Emergence rate	459.8	3	**< 0.0001**
Developmental time	13.7	2	**0.0011**
Survival of larval stages on Delfin® A			
Cumulative survival			
Dose × Larval instar	16.2	5	**0.0063**
Dose for each instar:			
- late 1st instar	87.4	5	**< 0.0001**
- late 2nd instar	25.7	5	**0.0001**
24-hour survival			
Dose × Larval instar	15.9	5	**0.007**
Dose for each instar:			
- late 1st instar	55.9	5	**< 0.0001**
- late 2nd instar	3.76	5	0.58
Adult fitness-related traits after development on Delfin® A			
Longevity			
Experiment	20.1	1	**< 0.0001**
- *1st experiment:*			
Dose	12.3	3	**0.0065**
Sex	35.0	1	**< 0.0001**
(e^β coefficient males *vs* females ± se: 0.55 ± 0.16)			
Dose × Sex	20.4	3	**0.00014**
Sexes analyzed separately			
- females	12.0	3	**0.0073**
(e^β coefficients *vs* control ± se: 5×10⁶: 1.05 ± 0.17, 5×10⁷: 0.71 ± 0.16, 10⁸: 0.60 ± 0.21)			
- males	20.4	3	**0.00014**
(e^β coefficients *vs* control ± se: 5×10⁶: 0.80 ± 0.16, 5×10⁷: 0.66 ± 0.16, 10⁸: 1.53 ± 0.18)			

760
761

Source of variation/Data	χ^2 / Deviance	d.f.	P value
Adult fitness-related traits after development on Delfin® A			
- 2ⁿᵈ experiment:			
Dose	16.5	3	**0.00090**
Sex	31.5	1	**< 0.0001**
(e^β coefficient males *vs* females ± se: 0.45 ± 0.22)			
Dose × Sex	0.69	3	0.88
Sexes analyzed separately			
- females	13.2	3	**0.0043**
(e^β coefficients doses *vs* control ± se: 5×10⁶: 0.92 ± 0.22, 5×10⁷: 0.63 ± 0.21, 10⁸: 0.51 ± 0.21)			
- males	7.01	3	**0.072**
(e^β coefficients doses *vs* control ± se: 5×10⁶: 1.02 ± 0.22, 5×10⁷: 0.70 ± 0.22, 10⁸: 0.64 ± 0.22)			
Total numbers of offspring			
Dose × Experiment	28.1	3	**< 0.0001**
Dose for each experiment:			
- 1ˢᵗ experiment	26.3	3	**< 0.0001**
- 2ⁿᵈ experiment	4.1	3	0.25
Development of other strains of *D. melanogaster* on Delfin® A (including Canton S)			
Emergence rate			
Dose × Fly strain	105.5	15	**< 0.0001**
Dose for each fly strain:			
- Canton S	588.6	5	**< 0.0001**
- Nasrallah	745.3	5	**< 0.0001**
- Sefra	900.7	5	**< 0.0001**
- YW1118	636.9	5	**< 0.0001**
Developmental time			
Dose × Fly strain	9.3	12	0.68
Dose for each fly strain:			
- Canton S	40.3	4	**< 0.0001**
- Nasrallah	18.0	4	**0.0012**
- Sefra	27.2	4	**< 0.0001**
- YW1118	28.9	4	**<0.0001**
Development of other *Drosophila* species on Delfin® A			
Emergence rate			
Dose × Fly species	538.2	30	**< 0.0001**
Dose for each species:			
- *D. simulans*	461.0	5	**< 0.0001**
- *D. yakuba*	750.7	5	**< 0.0001**
- *D. hydei*	596.8	5	**< 0.0001**
- *D. immigrans*	726.3	5	**< 0.0001**
- *D. subobscura*	729.6	5	**< 0.0001**
- *D. suzukii*	725.0	5	**< 0.0001**
- *D. busckii*	586.0	5	**< 0.0001**
Developmental time			
Dose × Fly species	59.9	22	**< 0.0001**
Dose for each species:			
- *D. simulans*	25.9	4	**< 0.0001**
- *D. yakuba*	34.7	4	**< 0.0001**
- *D. hydei*	11.5	4	**0.022**
- *D. immigrans*	6.01	3	0.11
- *D. subobscura*	68.8	4	**< 0.0001**
- *D. suzukii*	11.7	3	**0.0085**
- *D. busckii*	58.8	4	**< 0.0001**

 Figure legends

 Figure 1. Development of *D. melanogaster* Canton S flies on *Btk* and *Bti* commercial

 formulations. (**a**) Emergence rate (mean ± sem) and (**b**) developmental time (mean ± sem) of 20

 initial eggs on increasing doses of *Btk* Delfin$^{®}$ A (red triangles), the Cry-free *Btk* 4D22 (open

 lozenges), the mosquito-targeting *Bti* Vectobac$^{®}$ (green squares), and the non-pathogenic *Bacillus*

 subtilis (light grey circles). For Vectobac$^{®}$ and *B. subtilis*, N = 4-7 per dose; for Delfin$^{®}$ A and *Btk*

 4D22, N = 9-12 for the control, N = 3 for 5.10^{5} and 10^{9}, N = 4-9 for 10^{6}, N = 7-14 from 5.10^{6} to 10^{8}.

 (**c**) Emergence rate (mean ± sem) and (**d**) developmental time (mean ± sem) on increasing doses of

 the two *Btk* formulations Delfin$^{®}$ B (dark red circles) and Scutello DF (orange squares). N = 4

 replicates of 20 eggs per dose and formulation, except for controls and 10^{8} CFU/g of Delfin$^{®}$ B (9-

 10 replicates of 20 eggs). Results of *post hoc* comparisons of each dose to the control: · $0.05<p<0.1$;

 * $0.01<p<0.05$; ** $0.001<p<0.01$; *** $p<0.001$. (**e**) Immunoblotting with an anti-Cry1A polyclonal

 antibody on proteins from a suspension of laboratory-produced spores of Cry-free *Btk* 4D22, the

 three *Btk* formulations Delfin$^{®}$ A, B, Scutello DF, and a suspension of laboratory-produced Cry1A

 toxins. Red asterisks indicate the Cry protoxins (~130 kDa) and the activated fragments (~60 kDa

 and ~70 kDa).

 Figure 2. Survival of *D. melanogaster* Canton S larval stages on increasing doses of *Btk*

 Delfin$^{®}$ A. (**a**) Proportion of surviving larvae (mean ± sem) upon *Btk* exposure from the egg to late

 1^{st} instar (open lozenges) and late 2^{nd} instar (black triangles). (**b**) Proportion of surviving larvae

 (mean ± sem) upon 24-hour *Btk* exposure of early 1^{st} instar larvae (open lozenges) and 2^{nd} instar

 larvae (black triangles). N = 5-7 replicates of 20 individuals per dose. Results of *post hoc*

 comparisons of each dose with the control: * $0.01<p<0.05$; ** $0.001<p<0.01$; *** $p<0.001$.

 Figure 3. Fitness-related traits of adults (longevity and total offspring number) after

 development on *Btk* Delfin$^{®}$ A. (**a**, **d**) Female longevity (mean survival fraction over time ± sem),

(**b**, **e**) Male longevity (mean ± sem), and (**c**, **f**) total offspring number (mean ± sem), measured on

individuals that developed without *Btk* (blue items) and on $5×10^6$ CFU/g of *Btk* Delfin® A (green

items), $5×10^7$ CFU/g (red items), and 10^8 CFU/g (dark red items). Data from 2 experiments (a-c,

experiment 1; d-f, experiment 2). For each trait, N = 3-5 replicates of 15 males and 15 females per

dose in experiment 1, N = 3 replicates of 15 males and 15 females in experiment 2. Results of *post*

hoc comparisons of each dose with the control: * $0.01<p<0.05$; ** $0.001<p<0.01$; *** $p<0.001$.

Figure 4. Development of four *D. melanogaster* strains on increasing doses of *Btk* Delfin® A. (a)

Emergence rate (mean ± sem), (**b**) Developmental time (mean ± sem) of the strains Canton S (blue

lozenges), Nasrallah (yellow triangles), Sefra (green squares), and YW1118 (red circles). N = 4

groups of 50 eggs per dose and fly strain for each trait. Results of *post hoc* comparisons of each

dose to the control: · $0.05<p<0.1$; * $0.01<p<0.05$; ** $0.001<p<0.01$; *** $p<0.001$.

Figure 5. Emergence rate of seven *Drosophila* species on increasing doses of *Btk* Delfin® A.

Mean emergence rate (± sem). N = 4 replicates of 50 eggs per dose for *D. simulans*, *D. yakuba*, *D.*

subobscura, and *D. busckii*, N = 4 replicates of 30 eggs per dose for *D. hydei*, *D. suzukii*, and *D.*

immigrans. Results of *post hoc* comparisons of each dose with the control: · $0.05<p<0.1$; *

$0.01<p<0.05$; ** $0.001<p<0.01$; *** $p<0.001$.

Figure 6. Developmental time of seven *Drosophila* species on increasing doses of *Btk* Delfin® A.

Mean developmental time (± sem). N = 4 replicates of 50 eggs per dose for *D. simulans*, *D. yakuba*,

D. subobscura, and *D. busckii*, N = 4 replicates of 30 eggs per dose for *D. hydei*, *D. suzukii*, and *D.*

immigrans. Results of *post hoc* comparisons of each dose with the control: * $0.01<p<0.05$; **

$0.001<p<0.01$; *** $p<0.001$.

Figure 7. Evaluation of the role of small molecular weight components of *Btk* Delfin® A (dialysis; membrane cut-off: 8-10 kDa) in the altered development of *D. melanogaster* Canton S. (**a**) Emergence rate (mean ± sem) and (**b**) developmental time (mean ± sem) on increasing doses of dialyzed Delfin® A. $N = 3$ experiments of 4 replicates with 20 eggs per dose for the emergence rate, $N = 2$ experiments of 4 replicates per dose for the developmental time. Results of *post hoc* comparisons of each dose with the control: · $0.05 < p < 0.1$; * $0.01 < p < 0.05$; ** $0.001 < p < 0.01$; *** $p < 0.001$. (**c**) Anti-Cry1A probed immunoblot of non-dialyzed (ND) and dialyzed (D) suspensions showing the decrease in the amount of ~ 130 kDa protoxins and the increase in that of ~ 60/70 kDa activated toxins after dialysis.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7