Differential side-effects of *Bacillus thuringiensis* bioinsecticide on non-target *Drosophila* flies

Aurélie **Babin**, Marie-Paule **Nawrot-Esposito**, Armel **Gallet**, Jean-Luc **Gatti**[$*], Marylène **Poirié**[$]

Université Côte d'Azur, INRAE, CNRS, ISA, France

*Corresponding author

Institut Sophia Agrobiotech, INRAE, 400 route des chappes, 06903 Sophia Antipolis, France.

E-mail : jean-luc.gatti@inra.fr

[$]**Co-last authors**

Running title: *Bt* bioinsecticide and *Drosophila* flies

Keywords: *Drosophila*, *D. melanogaster*, *D. suzukii*, *Bacillus thuringiensis*, biopesticide, non-

intentional effects, development, longevity, fitness

Word count: main text: 5 284, abstract: 207

References: 90

Figures: 7 figures and 1 table

Supplementary information

Abstract

Biopesticides based on *Bacillus thuringiensis* (*Bt*) spores and toxins are alternate pest management solutions widely used to control insect pests. Their increasing use could lead to accumulation in the environment, hence leading to chronic exposure of non-target organisms. Here, we tested for potential non-intentional side-effects of chronic exposure to *Bt* biopesticide on larvae of non-target *Drosophila* species present in *Bt*-treated areas. Doses up to those recommended for field application ($\leqslant 10^6$ CFU/g of fly medium) had no effect on the fly development, whereas doses 10 to 100-fold higher (10^7-10^8 CFU/g) increased developmental time and decreased adult emergence rates in a dose-dependent manner and with varying effect amplitudes for all the species and strains tested. For all them, all larvae died before pupation at the highest dose tested (10^9 CFU/g). Focusing on *D. melanogaster*, delayed development and reduced emergence resulted from stage-dependent larval mortality, and fitness-related traits of adult flies emerging from surviving *Bt* biopesticide exposure were moderately increased. The effects of *Bt* biopesticide seemed to result from the spores/cleaved toxins synergy, and possibly additives. While recommended doses had no effect on non-target *Drosophila* species, misuse or local accumulation of *Bt* bioinsecticides in the environment could have non-intentional side-effects on fly populations with potential implications for their associated communities.

Introduction

The world's population is expected to reach more than 9 billion people by 2050,[1] increasing the demand for agricultural resources. Thisl requires to improve pest management, especially insects that cause more than 30% of agricultural losses.[2] Nowadays, their management largely relies on conventional chemical insecticides. However, their use and efficiency have been considerably reduced due to the emergence of pests' resistance, development of secondary pests, adverse side-effects on non-target species (natural enemies of pests, pollinators),[3,4] and more generally the impacts on biodiversity and human health (e.g. neurological disorders, functional impairment of reproduction, cancers).[5-8] Developed as a more specific and safer alternative, biopesticides represent less than 5% of the pesticide market, the large majority being microbial insecticide formulations based on viable spores and toxins of the bacterium *Bacillus thuringiensis* (*Bt*) (over 400 registered formulations).[4,9]

Bt is a Gram-positive endospore-forming bacterium that synthesizes a wide range of toxins with different chemical structures, modes of action and biological targets. The most abundant and studied are Cry δ-endotoxins encoded by genes located on large plasmids and produced as parasporal crystalline inclusions during the stationary growth phase.[10,11] *Bt* produces other insecticidal toxins, the Cyt (cytolytic δ-endotoxins) and Vip (secreted Vegetative Insecticidal Proteins) that synergize their effects with Cry toxins, virulence factors such as β-exotoxins (or thuringiensin), a secreted nucleotide toxic for almost all tested life forms thus prohibited in commercial formulations,[12] and anti-fungal factors.[13,14] *Bt* subspecies and strains can differ in their plasmid number and in the synthesized toxins cocktail responsible for their biological activity, which was used to delineate potential target insects.[15] For instance, *Bt* subsp. *kurstaki* (*Btk*) produces the 5 Cry toxins, Cry1Aa, Cry1Ab, Cry1Ac, Cry2Aa and Cry2Ab,[10,16] while *Bt* subsp. israelensis (*Bti*) produces a combination of Cry4Aa, Cry4Ba, Cry10Aa, and Cry11Aa,[17,18] both strains being commercially used. The different toxin cocktails produced by some *Bt* subspecies can also be detrimental to non-insect organisms such as nematodes, protozoa, and even molluscs.[15]

The bioinsecticide formulations based on spores and toxin crystals of *Btk* and *Bti* are the most

sprayed in organic and conventional farming, and natural areas (e.g. forests, swamps) to deal with

larvae of Lepidopteran pests and Dipteran larvae of mosquitoes and black flies, respectively. It is

generally accepted that once ingested by insect larvae, the toxin crystals are dissolved by the midgut

alkaline pH, releasing ~130 kDa pro-toxins that are then processed by digestive proteases into

smaller, soluble, active toxin fragments of ~ 60-70 kDa.[19,20] Active toxins bind to specific

receptors of midgut epithelial cells, eliciting pores formation in the cell membrane, cell lysis and

gut epithelium disorganization.[21,22] This allows gut bacteria, including *Bt*, to colonize the

hemocoel, and leads to rapid septicaemia and death.[23,24]

Numerous impact studies of field application rates and acute intoxications have shown that *Bt*

bioinsecticides are safe or have limited impact on non-target vertebrates and invertebrates, and

associated species communities.[25,26] However, the increasing use of biopesticides based on *Bt*

spores and toxins has recently raised concern[27] and led to the assessment of their potential effects

on non-target species, such as auxiliary insects of biological control,[28], pollinators,[29] and species

and species communities which simply share their habitat with *Bt*-targeted insect pests.[30-32] Yet,

there is growing evidence of direct and indirect cross-effects of *Bt* bioinsecticides and toxins across

insect species and orders, or even across phyla, suggesting that *Bt* targeting is only partly

specific.[33,34] Data showed that almost all of the applied *Btk* dose was still present on the leaves

surface 72 hours after spraying,[35] its amount returning close to environmental levels only 28 days

after treatment.[36] Finally, *Bt* spores can survive in the soil and on different supports for months and

even years after application.[37-41] *Bt* formulations contain also numerous compounds to protect

spores and crystals and aggregate them into a wettable form, surfactants to facilitate spraying and

dispersion on plants, and phagostimulants.[42,43] Nevertheless, since toxin crystals, and to a much

lesser extent spores,[44] are somewhat sensitive to abiotic conditions (e.g. UV, pH, rainfall), repeated

sprayings with a minimum delay of 3 to 8 days is often recommended over the period of pest

occurrence to achieve the required pest control level[43,45] (http://www.certiseurope.fr,

http://www.certisusa.com). All these can potentially lead to *Bt* accumulation in the environment,

thus raising the rarely addressed issue of potential side-effects of chronic exposure (*i.e.* continuous

and increasing exposure dose for an extended period) of non-target species to doses unexpectedly

above the recommended spraying doses.

Diptera are worldwide distributed insects, most of which are not targets for *Bt* and its toxins.

This is the case of the genus *Drosophila*, represented by ~ 1 500 described species,[46] including the

model organism *D. melanogaster*. In the field, most of these flies feed and reproduce mainly on

ripening or rotting/fermenting fruits and are therefore naturally present in areas treated with *Bt* such

as orchards, vineyards and gardening areas. Unable to disperse between food patches, early

developmental stages of *Drosophila* eat intensively and grow exponentially,[47] and may thus ingest

high doses of *Bt* bioinsecticides that have accumulated during the treatment periods. Surprisingly,

despite the presence of many *Drosophila* species in *Bt*-treated areas, their role in the decomposition

of organic matter, and the ease of study of some species, only a few studies have focused on these

flies. However, most of them suggested susceptibility to *Btk*, but they used mainly late 3rd instar

larvae preparing for pupation, which do not feed much, and used *Bt* preparations, especially field

isolates, that possibly contained highly toxic β-exotoxins, which are not authorized in commercial

Bt formulations.[48-55] So far, no study addressed the effects of chronic exposure to *Bt* formulations

containing spores and toxin crystals but no β-exotoxins, on developing stages of these Dipterans

that are present in *Bt*-treated areas.

Here, we have tested the chronic side-effects of different commercial formulations of *Btk*

(devoid of β-exotoxins) and, to a lesser extent of *Bti*, with doses starting from the maximum

recommended spraying dose up to 1 000 times this dose (i.e. below acute intoxication doses used in

most studies). We mainly focused on developmental traits (developmental time, emergence rate),

firstly using the wild-type *D. melanogaster* CantonS. The spore-forming Gram-positive *Bacillus*

subtilis and the *Btk* strain (4D22), devoid of Cry toxin genes and thus of crystals, were used as non-

pathogenic controls. We also analysed two fitness-related traits of adult flies (male and female

longevity, offspring number) after entire development in presence of *Btk* formulation. To test for

effects specific to the fly genetic background, developmental traits upon exposure to *Btk*

formulation were measured on four other *D. melanogaster* strains. Finally, we extended further

these development experiments to seven other *Drosophila* species. We have chosen six species in

addition to *D. melanogaster*, including cosmopolitan species, which can co-occur in the field[56-60]

and are usually present in *Bt*-treated areas, and the invasive *D. suzukii* that can now co-exist with

the six species in the areas it has recently invaded. This aims at providing a first-step in the

exploration of potential implications of chronic exposure to *Btk* formulation in terms of species

competition and species community composition and dynamics.

Material and methods

Commercial formulations, Bacillus *productions and Colony Forming Unit measurement*

The tested commercial brands of *Bacillus thuringiensis kurstaki* (*Btk*; serotype 3a, b, c[61]) were

Delfin® A and B (strain SA-11; wettable granules, Valent BioSciences, AMM 9200482, 32,000

UI/mg) and Scutello DF (a Dipel® sub-brand; strain ABTS-351; wettable granules, Biobest®, AMM

2010513, 540g/kg). The commercial brand of *Bacillus thuringiensis israelensis* (*Bti*; strain HD-14;

serotype 14[61]) was VectoBac® WG (wettable granules, Bayer, AMM 2020029, 3000 UTI/mg). For

each formulation, the number of viable spores (expressed as Colony Forming Units (CFU) per mg

of granules) was estimated using serial dilutions of a suspension on LB agar plates and counting of

bacterial colonies after overnight incubation at 30°C. CFU estimations were 5×10^7 CFU/mg for *Btk*

Delfin® A; 2.5×10^7 CFU/mg for *Btk* Delfin® B; 2.2×10^7 CFU/mg for *Btk* Scutello DF; 6×10^7

CFU/mg for *Bti* VectoBac®. Of note, our CFU estimations fell in those appended on the commercial

packaging, between 10^{13} and 5×10^{13} CFU/kg (32 000 UI/mg; http://www.certiseurope.fr). No

change in CFU estimations occurred during the time frame of the experiments. Manufacturer-

recommended doses for Delfin® range from 0.15 to 1.5 kg/ha depending on the crop type. Based on

our CFU estimations, this corresponds to recommended doses of 7.5×10^4 to 7.5×10^5 CFU/cm² of

Delfin[®] A, and 3.75×10^4 to 3.75×10^5 CFU/cm^2 of Delfin[®] B for each spraying in the field. For

Scutello DF, recommended doses range from 0.1 to 1 kg/ha, which are equivalent to 2.2×10^4 to

2.2×10^5 CFU/cm^2. Vectobac[®] WG is used at 0.125 to 1 kg/ha, equivalent to 7.5×10^4 to 6×10^5

CFU/cm^2.

The acrystillipherous (Cry toxin-free) *Btk* 4D22 strain (depleted for the toxin-encoding

plasmids[62]) obtained from the Bacillus Genetic Stock Center (http://bgsc.org; Columbus USA),

and a *Drosophila* non-pathogenic *Bacillus subtilis* (gift from Dr. E. Bremer, University of Marburg,

Germany; A. Brun-Barale, pers. comm.) were grown at 30°C in the sporulation-specific medium

PGSM (Bactopeptone[®] 7.5 g, KH$_2$PO$_4$ 3.4 g, K$_2$HPO$_4$ 4.35 g, glucose 7.5 g, PGSM salts 5 mL,

CaCl2 0.25 M, distilled water qsp 1L, pH 7.2; PGSM salts: MgSO$_4$.7H$_2$O, MnSO$_4$.H$_2$O,

ZnSO$_4$.7H$_2$O, FeSO$_4$.7H$_2$O) for about 14 days for sporulation to occur. Following elimination of

vegetative cells (1h at 70 °C), spore pellets were collected after centrifugation (4,500 rpm, 20 min,

4 °C), washed with sterile water, and lyophilized. CFU numbers were counted for each preparation

as described above.

Fly stocks

The four tested strains of *Drosophila melanogaster* (phylogenetic subgroup: melanogaster) were the

standard wild-type Canton S (Bloomington Drosophila Centre) used as a reference strain, the wild-

type Nasrallah strain from Tunisia (strain 1333, Gif-sur-Yvette), the double mutant standard strain

YW1118 (white and yellow mutations; gift from Dr. B. Charroux, IBD, Marseille-Luminy)*,* and a

recently field-collected strain (caught in Southern France in 2013) that we named "Sefra". For

Drosophila species comparison, we included 6 species of the *Drosophila* subgenus, *D. simulans*

(strain 1132; phylogenetic subgroup: melanogaster), *D. yakuba* (strain 1880; phylogenetic subgroup:

melanogaster), *D. hydei* (phylogenetic subgroup: hydei) and *D. suzukii* (phylogenetic subgroup:

immigrans) (both kindly provided by Dr. R. Allemand, LBBE, University Lyon 1), *D. immigrans*

(phylogenetic subgroup: immigrans), *D. subobscura* (phylogenetic subgroup: obscura), and one

species of the *Dorsilopha* subgenus, *D. busckii* (all three species collected in South-East of France

in Spring 2015).

All strains and species were maintained at controlled densities (150-200 eggs/40 ml of fly

medium) under standard laboratory conditions (25°C or 20°C for recently collected species, 60 %

relative humidity, 12:12 light/dark cycle), on a high-protein/sugar-free fly medium (10 % cornmeal,

10 % yeast, 0 % sugar). The *D. melanogaster* Canton S strain was also reared on a standard low-

protein/sugar-free fly medium (8 % cornmeal, 2 % yeast, 2.5 % sugar) to test for the influence of

the medium composition on *Btk* exposure effects.

Intoxication method and dose-response assay

Commercial formulations and laboratory spore productions were suspended and diluted in buffer to

perform dose-response assays with doses from 10^5 to 10^9 CFU/g of fly medium. All doses were

prepared in 100 µl and homogenized thoroughly with the fly medium (100µl/g). *Drosophila* eggs

and larvae were collected from stock vials at the suitable developmental stage and transferred

carefully to the intoxication vials and dishes, then maintained under standard laboratory conditions

until a) the emergence of adults, or, in the larvae survival tests, b) until a given developmental stage

was reached from the egg, and c) for 24h. Control groups of individuals were transferred on fly

medium homogenized with the same volume of buffer.

Development-related traits and larval survival

To evaluate emergence rates and developmental times upon intoxication throughout the entire

development, precise numbers of eggs from mass oviposition were transferred to intoxication vials

containing fly medium mixed with doses of *Bt* formulations or bacteria productions and let to

develop under standard laboratory conditions until the fly emergence. Eggs without chorion and

transparent eggs were discarded. The initial number of eggs was adjusted depending on the species

biology and the vial size: 20 eggs for 2 g of fly medium in small vials (Ø 3.3 cm, surface ~8.5 cm^2,

0.24 g/cm^2) for tests with *D. melanogaster* Canton S, 50 eggs for 6 g of fly medium for comparison

of *D. melanogaster* strains and *Drosophila* species in wider vials (Ø 4.6 cm, surface ~16 cm^2, 0.37

g/cm^2) except for *D. hydei*, *D. suzukii* and *D. immigrans* for which 30 eggs were transferred on 6 g

of fly medium. Numbers and sex of emerging flies were recorded once a day until the day the pupae

of the next generation should form. From these data, the emergence rate (proportion of emerged

flies from the initial eggs; ER), the mean developmental time (mean number of days for completion

of development; DT), and the sex-ratio (proportion of male flies; SR) were calculated for each

intoxication vial.

For the larval survival tests, 20 eggs or larvae from a 4-hour mass oviposition at the indicated

developmental stage, were transferred to small dishes containing 1 g of fly medium (Ø 3 cm,

surface ~7 cm^2) homogenized with increasing doses of Delfin® A. Surviving larvae were counted at

the indicated developmental stage, or after 24-hour intoxication, to calculate the proportion of

surviving larvae. For the test from the egg, eggs which did not hatch were not included in the

counting. As a control measurement, we measured the pH of the fly medium in the presence of the

dose range of *Bt* formulations (see Supplementary Information S4).

Adult fitness-related traits

For the longevity and offspring number tests, males and females emerged from several rearing vials

for each dose of Delfin® A were pooled when aged 2 days. Groups of 15 males and 15 females were

transferred into vials with fresh fly medium without formulation. Fly medium was renewed every 3-

4 days. After each fly transfer to fresh food, discarded maintenance vials were incubated under

standard laboratory conditions for the offspring to develop. Mortality and sex of dead flies were

recorded daily until the last fly died. Offspring numbers were counted from the first emergence

until pupae of the next generation appeared. The tests were repeated twice. Due to the variation in

the duration of the two longevity experiments, offspring numbers of each vial were summed to

obtain a total offspring number per dose of Delfin® A for each experiment.

Dialysis and Cry toxin analysis

A suspension of 2×10^{10} CFU of Delfin® A was dialyzed against PBS (KH_2PO_4 1.06 mM, $Na_2HPO_4(2H_2O)$ 3mM, NaCl 154 mM, qsp distilled water, pH 7.2), at 250 rpm, 4°C overnight, using an 8-10 kDa MW cut-off membrane (ZelluTrans, Roth®). The CFUs of the dialyzed suspension and the effects on emergence rate (ER) and developmental time (DT) were analysed as described above. The dialyzed suspension was also subject to a 12.5 % SDS-PAGE and compared to the non-dialyzed suspension after silver staining. The presence of Cry1A pro-toxins, activated toxins and toxin fragments was probed by Western-blot using an in-house anti-Cry1A rabbit polyclonal antibody.

Data analysis

Data on development traits (emergence rate ER and developmental time DT), sex-ratio (SR), survival of larval stages and offspring number were analysed with mixed effect models including the dose of *Btk* formulation/spore production, the *D. melanogaster* strain, the *Drosophila* species or the developmental stage as fixed effects, and replicate (plus the experiment when necessary) as random effects (for ER data, data were analysed with bias-corrected models with replicate as fixed effect to allow pairwise comparisons; similar results obtained with models including replicate as random effect). ER, SR and survival of larval stages were analysed with generalized linear models, with binomial distribution and logit link. DT and offspring number were analysed with linear models. DT were transformed into developmental rates (1/developmental time) to fulfil the assumptions of the analysis of variance (homoscedasticity and residuals normality). Adult longevity data were analysed with proportional hazard Cox regression models including fly sex and dose of *Btk* formulation as fixed effects, and replicates as a random effect. For all the data sets, the main fixed effects and their interactions were tested with log-likelihood ratio tests. *Post hoc* pairwise comparisons were made for pairs of *D. melanogaster* strains, formulation/spore treatments, and

between the control dose and the other doses. All the analyses were performed in R[63] using the

packages lme4,[64] brglm,[65] multcomp,[66], survival,[67] and coxme.[68]

Results

***Btk* formulations adversely impact the development of *D. melanogaster*.**

The wild-type Canton S strain of *D. melanogaster* was used to evaluate the dose-dependent effect

of the commercial *Btk* formulation Delfin® A on the emergence rate (ER, proportion of emerged

flies from the initial egg pool) and developmental time (DT, mean number of days from egg to adult

emergence). Eggs were transferred on a standard low-protein/high-sugar fly medium containing

Delfin® A at doses ranging from 5×10^5 CFU/g of medium (mean equivalent of the maximum

recommended doses for one field application; see Methods and Supplementary information S1) to

10^9 CFU/g (~ 1,000 times the recommended dose). To check for specific effects of *Btk* formulations

and the respective role of *Btk* spores and Cry toxins, we tested the same dose range of the

commercial *Bti* formulation Vectobac® targeting mosquitoes that contains different Cry toxins,[22]

of the Cry-free strain *Btk* 4D22, and of the *Drosophila* non-pathogenic spore-forming *Bacillus*

subtilis.

Developmental traits (ER and DT) of exposed and non-exposed control flies were similar at

doses up to 10^7 CFU/g of Delfin® A (Fig. 1a-b; Table 1). At higher doses, both ER and DT were

affected in a dose-dependent manner: ER was reduced by 17% at 5×10^7 CFU/g (although not

statistically significant), up to 100% at 10^9 CFU/g, dose at which no individual reached the pupal

stage. The lethal dose 50 (LD50) was estimated between 5×10^7 and 10^8 CFU/g (Fig. 1a). DT was

increased of about 0.5 day at 5×10^7 CFU/g (+4% compared to controls), up to 1.5 days (+14%) at

10^8 CFU/g (Fig. 1b; Table 1). The sex-ratio at emergence (SR, proportion of males) was strongly

biased towards males at the highest dose at which complete development occurred (10^8 CFU/g),

with 58% more males compared to controls (Supplementary information S2). Because addition of

Btk formulation could modify parameters of the fly medium and thus contribute to these effects, we

checked the pH of the medium: the presence of formulation and its dose had no effect on it

(Supplementary information S4).

We observed no change in ER using the same dose range of the *Btk* Cry-free strain 4D22 (Fig.

1a, 1e; Table 1) and the non-pathogenic *Bacillus subtilis* (Fig. 1a, Table 1). Addition of *Bti*

Vectobac[®] did not affect ER up to 10^8 CFU/g but reduced it by 89% at 10^9 CFU/g (~2,000 times the

highest recommended dose for field application; Fig. 1a; Table 1; Supplementary information S1).

DT varied with the dose of *Btk* 4D22, the differences being mainly between doses but not with the

control. DT increased by ~1.5 days at the highest dose of Vectobac[®] (Fig. 1b; Table 1) and showed

a similar trend with *B. subtilis* (P = 0.06; Fig. 1b; Table 1). None of these three treatments

influenced dramatically the SR, the slight decrease in male proportion for most of the Vectobac[®]

doses being due to the higher average sex-ratio for the control dose compared to those for the two

other treatments (Supplementary information S2).

To test whether these effects are generic to *Btk* formulations, the fly development was

evaluated on two other formulations, Delfin[®] B (same brand) and Scutello DF (brand Dipel[®]), at the

critical doses 10^8 and 10^9 CFU/g. As Delfin[®] A, these formulations contain spores and Cry toxins

such as Cry-1A as pro-toxins of ~130 kDa, activated toxins of ~60-70 kDa, but also as smaller

fragments[20] (Fig. 1e, red asterisks). ER remained unchanged at 10^8 CFU/g whereas no individual

reached pupation at 10^9 CFU/g on Delfin[®] B and very few individuals reached the adult stage on

Scutello DF[®], DT being increased by more than 2 days (Fig. 1c-d; Table 1). No significant bias in

SR was observed for either formulation (Supplementary information S2).

Adverse effects of *Btk* formulation strongly impact the early development.

Larval stages were assessed for their susceptibility to *Btk* formulation in two independent and

complementary dose-response tests of survival to Delfin[®] A, at doses ranging from 10^5 to 10^9

CFU/g of high protein/sugar free medium (this medium is used to rear fly species which are

difficult to rear in the lab (see below) and is less limiting for the development of early larval stages).

We focused on the 1^{st} and 2^{nd} larval instars, during which growth is exponential,[47] so that larvae

are most heavily exposed to the bioinsecticide. In the first test, the cumulative survival was

measured by counting alive late 1st and 2nd instar larvae which have been exposed to Delfin® A

from the egg stage. Larval survival was not influenced at 10^7 CFU/g, whereas it decreased for both

larval instars above that dose to reach up to 37% mortality at 10^9 CFU/g (Fig. 2a). Reduced survival

tended to occur at a lower dose when cumulative survival was measured later in the development,

i.e. 10^9 for late 1^{st} instar larvae and 10^8 CFU/g for 2^{nd} instar larvae (Fig. 2a; Table 1). For both

instars, larvae surviving 10^9 CFU/g were noticeably smaller and less active than those surviving

lower doses. In emergence assays with planned exposure from the egg to the adult stage, none of

these individuals reached the pupal stage (see results above). In the second test, larval survival was

measured after early 1^{st} and 2^{nd} instar larvae had been exposed for 24 hours to Delfin® A. Survival

of 1^{st} instar larvae decreased by 36% on 10^9 CFU/g whereas that of 2^{nd} instar larvae did not change

(Fig. 2b, Table 1).

Developmental exposure to *Btk* formulation does not strongly influence fitness-related traits

in adults.

Long-term consequences on flies of exposure to *Btk* formulation throughout the development were

evaluated on two fitness-related traits, longevity and total offspring number. Traits were measured

on a *Btk*-free low-protein/high-sugar medium after individuals had completed their development on

the same fly medium but in presence of selected doses of Delfin® A: 5×10^6 CFU/g, which had no

impact on development, and 5×10^7 and 10^8 CFU/g, which caused moderate and strong development

alterations, respectively (see Fig. 1a).

Adult longevity was analysed in two independent sets of experimental replicates on groups of

15 females and 15 males held together. Despite large variation between the two sets of experimental

replicates (Table 1), the longevity of adults reared on 5×10^6 CFU/g of Delfin® A was similar to that

of non-exposed controls (Fig. 3). Males and females which developed on the two higher doses

showed a moderate longevity benefit, higher in females for 10^8 CFU/g (Fig. 3a-b, d-e; Table 1).

Males generally survived better than females (Table 1) but their longevity benefit of developing on

10^8 CFU/g was only observed in the second experiment (Fig. 3b, e).

The female offspring number - the sum of offspring produced by the 15 females of each fly

group during the longevity experiment - varied depending on both the experiment and the Delfin$^®$ A

dose (Table 1). In the 1st experiment, adults from larvae reared on 10^8 CFU/g had fewer offspring

compared to control adults and to adults developed on the other doses whereas the total offspring

number varied regardless of the *Btk* dose in the 2nd experiment (Fig. 3c, f, Table 1).

Btk*-formulation dose-dependent alterations of development are not specific to the *D.

***melanogaster* Canton S strain.**

Dose-dependent effects of *Btk* formulation on the development were tested on three additional *D.*

melanogaster strains: the wild-type Nasrallah (strain 1333), the wild-type Sefra population reared in

the laboratory for 4 years, and the double mutant YW1118. The emergence rates (ER) and

developmental times (DT) were measured on a high-protein/sugar-free medium (rearing medium of

these strains) mixed with Delfin$^®$ A doses ranging from 10^5 to 10^9 CFU/g. To allow the comparison

with previous results with Canton S flies on low-protein/high sugar fly medium, Canton S was also

reared and tested on the high-protein/sugar-free medium along with the other strains.

None of the fly strains was impacted at doses up to 10^7 CFU/g, whereas ER was strongly

reduced and DT was increased at higher doses for all the strains (Fig. 4a-b, Table 1), with no

individual reaching the pupal stage at 10^9 CFU/g (LD50 between 10^8 and 10^9 CFU/g). At 10^8

CFU/g, the magnitude of effects on Canton S flies was lower than that observed on the low-

protein/high-sugar medium. At this dose, ER varied between strains, the largest reduction being

observed for Sefra (Table 1). We observed no dose-dependent bias in SR (Supplementary

information S3).

Btk formulation also affects other _Drosophila_ species, including the invasive pest _D. suzukii_.

The ER and DT were analysed for seven other _Drosophila_ species from different phylogenetic clades at doses of Delfin® A from 10^5 to 10^9 CFU/g of high-protein/sugar-free medium (rearing medium of all the species). Tested species were _D. simulans_ (_D. melanogaster_ sister species), the African _D. yakuba_, _D. subobscura_, _D. immigrans_, _D. hydei_, and the invasive _D. suzukii_, all belonging to the _Drosophila_ subgenus, and _D. busckii_ from the _Dorsilopha_ subgenus. For all the species, doses up to 10^6 CFU/g of Delfin® A had no effect on ER and DT whereas all individuals failed to reach the pupal stage and no fly emerged at 10^9 CFU/g (Fig. 5-6). Amplitudes of development alterations at 10^7 and 10^8 CFU/g varied between species (Fig. 5-6; Table 1). All species were affected at 10^8 CFU/g as was _D. melanogaster_ (see Fig. 4a for comparison). _D. simulans_ and _D. busckii_ had unchanged ER, but DT was slightly increased for _D. simulans_ (although slightly reduced at 10^7 CFU/g; similar results with a Japanese strain, data not shown) and strongly increased for _D. busckii_ (by 20%, _i.e._ ~ 4 days) (Fig. 5-6, Table 1). _D. yakuba_ ER and DT were similar to those of _D. melanogaster_, with an LD50 around 10^8 CFU/g and a moderate DT increase of ~ 1 day (Fig. 5-6, Table 1; similar results with a strain from Sweden, data not shown). The ER of _D. hydei_ and _D. subobscura_ were very low at 10^8 CFU/g (LD50 below this dose), with a high DT (Fig. 5-6; Table 1), while _D. immigrans_ did not survive. No _D. suzukii_ individual emerged at 10^8 CFU/g and development was already moderately impacted at 10^7 CFU/g (Fig. 5-6). No dose-dependent bias in SR was detected for either species (Supplementary information S5).

Development alterations may result from a synergy between formulation components.

_Bt_ spores and toxins can represent more than half the weight of commercial formulations (85% for Delfin®, http://www.certisusa.com), with up to about 10% of insecticidal protein toxins within this fraction, mainly Cry pro-toxins and activated toxins[69] (see Fig. 1e). The remaining weight consists of various compounds such as residues of culture medium and various additives including surfactant, anti-foaming agents, etc…[25,43] It has been shown that, for some products, additives can

be more harmful in some cases than the active ingredient,[70] we explored the role of small

diffusible molecular weight components of Delfin® A in the alterations of ER and DT of *D.*

melanogaster Canton S. For that, we mixed a 10 kDa dialyzed suspension of Delfin® A at 10^7, 10^8,

and 10^9 CFU/g with low-protein/high-sugar medium. ER and DT were unaffected by the presence

of the dialyzed suspension at the 10^7 CFU/g dose, whereas no individual reached the adult stage (no

pupation) with the suspension at the 10^9 CFU/g dose (Fig. 7a; Table 1). At 10^8 CFU/g, ER was not

modified but DT increased by ~ 1 day, only in one of the two experiments, partially reproducing the

changes observed without dialysis (Fig. 7a-b; see also Fig. 1a-b, Table 1; 3 independent

experiments for ER, 2 independent experiments for DT).

Cry1A profiles of dialyzed Delfin® A suspensions, like those of the non-dialyzed ones,

comprised 130-kDa pro-toxins and 60-70 kDa activated toxins, but also showed toxin degradation

as evidenced by additional smaller fragments of activated toxins (Fig. 7c). The respective roles of

Btk toxin fragments and spores in the alterations of *D. melanogaster* development were further

explored through experiments of dialysis such as those previously described, followed by

successive centrifugations to eliminate most of the spores and toxin crystals. Despite variation

between experiments, ER was strongly affected only in one of the three experiments while DT was

always significantly increased when flies were reared in presence of centrifuged supernatants that

contained a limited range of Cry 1A toxin fragments (Supplementary information S6).

Discussion

The increasing use of bioinsecticides based on *Bacillus thuringiensis* (*Bt*) raises concern about their

potential non-intentional side-effects on non-target organisms, and biodiversity in general, due to

their partially specific targeting,[33,34,71] persistence in the environment,[35,36,38,40,41,44,72] and the

requirement of and recommendations for repeated spraying to reach the desired pest control

level.[43] Especially, side-effects of chronic exposure on non-target organisms, including insects

present on treated areas, remain under-evaluated. Here, we have tested the side-effects of ingestion

of *Bt* bioinsecticide formulations (mainly made of *Bt kurstakii* strains (*Btk*) but also of *Bt israelensis*

(*Bti*)) throughout the entire development of several non-target species of *Drosophila* flies which are

naturally present in treated areas. While formulation doses up to those recommended for each field

sprayings ($\leq 10^6$ CFU/g of medium) had no effect on *Drosophila* development, mortality and/or

developmental delay occurred markedly from doses only 10 times and 50 times higher than the

maximum recommended dose of the main tested *Btk* formulation for *D. suzukii* (10^7 CFU/g) and the

D. melanogaster strains (5×10^7 CFU/g), respectively. Development alterations were already strong

at these doses, suggesting the occurrence of alterations already at lower doses, i.e. between 10^6 and

10^7 CFU/g for *D. suzukii*, and between 10^7 and 5×10^7 CFU/g for *D. melanogaster* strains. Accurate

analyses would be needed to verify this possibility. Besides, all the tested species, except *D.*

simulans, were strongly affected at 10^8 CFU/g, and no (or extremely limited) fly development

occurred at the highest tested dose (10^9 CFU/g), equivalent to 1000 times the maximum

recommended dose but far below common acute intoxication doses used in many studies.[5]

Recommended doses are those for each spraying on a homogeneous and dry zone without covering

areas. In the field, recommended repeated sprayings of stabilized formulation and rainfall washouts

following spraying may increase *Bt* spores and toxins presence in both space and time. While the

highest dose tested here (10^9 CFU/g) would be hardly reached in the field as those for acute

intoxications, the minimal doses at which the fly development was markedly impacted and the

lower ones from which changes in development appeared may be readily obtained. Furthermore, the

minimal quantity of *Bt* formulation inducing development alterations may be even lower since a

single *Drosophila* larva is unlikely to process 1g of medium given its size and feeding rate. Our data

also evidenced a window of *Btk* susceptibility during larval development, with the ingestion during

the 1st larval instar accounting for a large part of the observed detrimental effects on the

development (see below in the discussion).

When testing for non-intentional generic effects of *Bt* formulations, similar patterns of

development alterations were observed but shifted to higher doses with two other *Btk* formulations

and a formulation of *Bti*: there was no effect on *D. melanogaster* development at the doses up to 10^8

CFU/g but a strong detrimental effect at the highest dose tested, 10^9 CFU/g. The three *Btk*

formulations tested, based on two different bacterial strains (see Methods), had similar profiles of

Cry1A protoxins and activated toxins, but they differed in their efficient spore contents, formulation

type, and likely additives, which may account for the observed variation in the half-lethal dose. The

Bti formulation, widely used against Dipteran Nematocera insects (e.g. mosquitoes, black flies),[73]

impacted *D. melanogaster* development only at the highest dose tested. The impacts of *Btk*

formulations on *D. melanogaster* development are consistent with growing evidence suggesting a

partly specific targeting of *Bt*.[33,74] Until recently, it has been generally accepted that the mode of

action of *Bt* after ingestion by insects relies on key steps of specific binding of proteolyzed *Bt* toxins

to receptors of midgut epithelial cells, defining targets for each *Bt* subspecies.[15,19,22] Several

primary and secondary types of toxin receptors, including cadherin-like proteins, aminopeptidases,

GPI-anchored alkaline phosphatases,[10] and more recently the ATP dependent binding cassette

reporter C2,[75] have been identified in Lepidoptera and Diptera mosquitoes. Focusing on the action

of *Btk* targeting Lepidoptera, no Lepidoptera cadherin-like Cry receptor orthologues were found in

Drosophila,[75] supporting the idea that these flies would not be affected by the spraying of *Btk*

formulation. The existence of other types of Cry receptors in *Drosophila* flies would explain the

observed developmental impacts but remains to be investigated. In addition, the substantial amounts

of active Cry1A toxin fragments in *Btk* formulations could compensate for the possible lack of

solubilization of protoxin crystals in the fly midgut and proteolytic activation of toxins by fly gut

proteases, both required for Cry activity in insect larvae.[19] Other toxins synthesized by *Btk* and

present in the formulations could also play a role in the observed cross-order activity as some, such

as Cry2A, have an insecticidal effect on both Lepidoptera and Diptera.[76]

The lack of effect of ingestion of *Bacillus subtilis* or *Btk* Cry-free 4D22 on the development

of *D. melanogaster* excludes that development alterations result from severe disruption of digestion

and nutrient uptake/competition in the presence of high spore/bacteria loads in the larval gut

throughout development. It supports the idea of a synergistic action of *Btk* spores and Cry toxins,

consistent with the models of *Bt* action on insect larvae in which toxins first breach the gut

epithelium, allowing the gut content, including *Bt* spores, to colonize the hemocoel.[19,22-24] The

partially reproduced mortality rate and delayed development in dialysis experiments may further

indicate that low diffusible molecular weight compounds in *Btk* formulations (e.g., culture media

residues, salts, additives) may contribute to these development alterations. This is supported by the

lack of impact on *D. melanogaster* development of the ingestion of laboratory-produced spores and

Cry toxins of a *Btk* 4D1 strain (or HD1, a reference strain used as a control strain here, not used in

commercial formulations) used without additives, even at the highest dose 10^9 CFU/g (additional

information S7; Fig. S7a, b). The *Btk* 4D1 culture contained few active Cry toxins and smaller toxin

fragments, in contrast to commercial *Btk* formulations (Fig. S7e), supporting the possible

contribution of these toxin fragments to the cross-order activity of *Btk* formulations on *Drosophila*.

As previously reported for *D. suzukii* exposed to laboratory-produced *Btk* cultures,[53]

mortality of *D. melanogaster* during development on *Btk* formulation occurred early in

development. First and second instars larvae are likely highly exposed due to their high feeding rate

and their exponential growth.[77] As the observed larval mortality was only about 40% at the highest

dose (10^9 CFU/g) (Figure 2), while none of the individuals reached the pupal stage at this dose

(Figure 4), the remaining mortality likely occurred during the third larval stage, maybe due to a

delayed action of *Btk* spores and toxins. Interestingly, alterations of the development (mortality and

delayed emergence) mimicked those typically generated by nutritional stress conditions in insect

larvae.[78,79] Accordingly, the development alterations were partially rescued on a protein rich fly

medium, probably through compensatory protein intake, as in other arthropod species.[79-81] In

addition, the sex ratio of flies was strongly biased towards males after development on the dose of

Btk formulation affecting fly emergence (10^8 CFU/g) and under low protein conditions. This

highlights the importance of nutritional conditions in *Btk* impacts on *Drosophila* development, with

sex-specific differences in larval susceptibility to environmental stressors, here the accumulation of

Btk formulation, under protein restriction conditions as reported previously in *D. melanogaster*.[82]

The development on sublethal doses of *Btk* formulation did not dramatically affect the

longevity of *D. melanogaster* adults and the lifetime offspring number. Developmental exposure to

Btk doses that slightly and strongly reduced the likelihood of reaching the adult stage even gave to

males and females a dose-dependent longevity benefit, in addition to the male higher longevity

observed in mixed-sex populations,[83] and slightly increased the offspring number (although not

significantly). Surviving the exposure to *Btk* formulation throughout the development has likely

selected for fitter individuals. This is similar to the increased longevity of adult insects that have

survived developmental nutritional stress,[84,85] or are resistant to environmental stressors.[83]

The origin of *Drosophila* (species and population/strain) influenced the magnitude of the

development alterations induced by *Btk* formulation. For *D. melanogaster*, all the strains tested

were susceptible to *Btk* formulation with both mortality and delayed development at the same dose,

but with variation in the effect magnitudes. This suggests potential population-specific differences

in susceptibility to *Btk* formulation accumulation in the environment, and hence potential spatial

and temporal heterogeneity of *Btk* spraying impacts for each *Drosophila* species. Among the other

seven species tested, differences in susceptibility to *Btk* formulation, in terms of effect magnitude

and type of development alteration (mortality and/or developmental delay), occurred between

Drosophila species, regardless of their phylogenetic distances. For the *Drosophila* subgenus, *D.*

simulans was less susceptible than its sister species *D. melanogaster*, whereas the African *D.*

yakuba experienced similar impacts on the development as *D. melanogaster*. The three species *D.*

immigrans, *D. subobscura* and *D. hydei* were similarly more susceptible than *D. melanogaster*, but

with slight differences in effect magnitudes. The phylogenetically distant *D. busckii* (*Dorsilopha*

subgenus) was the least affected of all the species tested in terms of developmental mortality, but its

development was strongly delayed. The five species *D. melanogaster*, *D. simulans*, *D. hydei*, *D.*

immigrans, and *D. busckii* belong to the guild of cosmopolitan domestic *Drosophila* species, *D.*

subobscura is sub-cosmopolitan species, and *D. busckii* is an opportunistic frugivorous species.[86]

All these species coexist frequently and compete on the same discrete and ephemeral rotting fruit

patches, with seasonal variations in the composition of the fly community.[56-59] At the community

level, differences in the species susceptibility to accumulation of *Btk* formulation could modify

larval competition conditions and lead to additional local and temporal variations in *Drosophila*

communities' composition. The potential side-effects of *Bt* sprays on non-target *Drosophila*

communities would be hardly predictable as they would depend on spatial patterns of *Bt*

accumulation. A formal mesocosm study of *Drosophila* community dynamics under exposure to

Btk formulation, at least under semi-field conditions, would allow assessing the consequences of *Bt*

accumulation on species competition and community composition.

As for the other species, the presence of *Btk* formulation impacts the development of the

invasive *D. suzukii*, as recently reported by Cossentine et al.,[53] this species being the most

susceptible here with effects already clearly detectable at 10 times the recommended spraying dose.

Compared with the other seven species that live on rotten fruits, *D. suzukii* poses a threat to fruit

production because it feeds and lays eggs in healthy ripening soft fruits[87-89] and hence colonizes

orchards and vineyards earlier during the fruit season. Despite its oviposition mode with a saw-like

ovipositor inserted inside ripening soft fruits,[87] the exposure of *D. suzukii* larvae may be quite

similar to that of larvae of other *Drosophila* species laying on the surface of fermenting fruits or

rotting fruit flesh. Indeed, the saw-like ovipositor likely carries *Bt* spores and toxins from the

surface of treated fruits when piercing the fruit skin, and additional *Bt* may then enter the fruit

through the drilled holes. From an ecological point of view, the greater susceptibility of *D. suzukii*

to the accumulation of *Btk* formulation in the environment might mitigate the potential ecological

burden of its invasion for local communities of *Drosophila* frugivorous species in orchards.

Alternatively, as *D. suzukii* attacks on fruits can accelerate their decomposition by microorganisms,

its higher susceptibility to *Btk* could reduce the number of fruits made suitable for other *Drosophila*

species.

In conclusion, our results showed that at recommended *Bt* doses, no visible effects on

Drosophila was observed, but repeated sprayings and subsequent accumulation of *Btk* biopesticide

above these doses can potentially impact these non-target insects. The magnitude of this impact

possibly depends on both the formulation used and the insect species. Although our study was

carried out under controlled laboratory conditions which may dramatically differ from natural

conditions encountered in the field (temperature, pH, humidity, food availability, photoperiod,

predator/parasite/pathogen pressures, etc…), standard laboratory strains and flies derived from wild

populations recently collected exhibited similar patterns of development alterations, suggesting our

results may not be specific to laboratory-influenced genetic backgrounds. Recent studies have

reported similar adverse side-effects due to repeated sprayings of the *Bti* formulation, directly on

non-target organisms,[31] and indirectly on predators via food webs.[90] These studies and the data

presented here highlight that pest control with *Bt* bioinsecticides should be done with caution in the

field to avoid, or at least limit, potential non-intentional side-effects on non-target organisms and

hence on biodiversity. At last, *D. melanogaster*, a model species in many research fields, could also

serve as a study model to identify the mechanisms underlying these side-effects and/or the potential

emergence of resistance to these biopesticides.

Acknowledgements

We thank Xiao Han, Jingru Li and Abir Oueslati for help with preliminary experiments, L.

Kremmer, C. Rebuf and O. Magliano for providing and rearing flies and help in preparing fly

medium, A. Brun-Barale for the *Bacillus subtilis* spores, D. Pauron for preparation of Cry1A toxin,

Hugo Mathé-Hubert for advice on statistical analyses, and M. Amichot for helpful discussions. The

Cry1A antibody was produced in collaboration with the INRA-PFIE platform (Nouzilly, France).

Financial Supports

This work was supported by the French National Agency for Research (ANR-13-CESA-0003-001

ImBio), the European Union's Seventh Framework Program for research, technological

development and demonstration under grant agreement No. 613678 (DROPSA), the "Investments

for the Future" LABEX SIGNALIFE (ANR-11-LABX-0028), the INRA Plant Health Department

(to MPNE and JLG), the CNRS (to AG), and the University Nice Côte d'Azur (to MP).

Author contributions

AB, MPNE, AG, JLG and MP designed the experiments. AB performed the experiments with

contributions of MPNE. AB performed the statistical analyses. AB, JLG, and MP wrote the

manuscript with contributions from all the authors.

Additional information:

Supplementary information

Competing interest: The authors declare no competing interests.

References

1. United Nations, Department of Economic and Social Affairs, Population Division 2015. World

Population Prospects 2015 – Data Booklet (ST/ESA/ SER.A/377).

2. Pimentel, D., Burgess, M. 2014. Environmental and economic costs of the application of

pesticides primarily in the United States. In: Integrated Pest Management. Dordrecht: Springer

Netherlands. pp. 47–71.

3. Devine, G.J., Furlong, M.J. 2007. Insecticide use: contexts and ecological consequences.

Agriculture and Human Values, 24(3): 281–306.

4. Sanchis, V., Bourguet, D. 2008. *Bacillus thuringiensis*: applications in agriculture and insect

resistance management. A review. *Agronomy for Sustainable Development*, 28(1): 11–20.

5. WHO report 2007. WHO specifications and evaluations for public health pesticides: *Bacillus*

thuringiensis subspecies israelensis strain AM65-52. World Health Organization, Geneva.

6. Baldi, I., Cordier, S., Coumoul, X., Elbaz, A. 2013. Pesticides, effets sur la santé. Expertise

collective. INSERM. Available at: http://ipubli-inserm.inist.fr/handle/10608/1.

7. Gilden, R.C., Huffling, K., Sattler, B. 2010. Pesticides and health risks. *Journal of Obstetric,*

Gynecologic & Neonatal Nursing, 39(1): 103–110.

8. Rizzati, V., Briand, O., Guillou, H., Gamet-Payrastre, L. 2016. Effects of pesticide mixtures in

human and animal models: an update of the recent literature. *Chemico-Biological Interactions*,

254: 231–246.

9. Lacey, L.A., Grzywacz, D., Shapiro-Ilan, D.I., Frutos, R., Brownbridge, M., Goettel, M.S. 2015.

Insect pathogens as biological control agents: Back to the future. *Journal of Invertebrate*

Pathology, 132(C): 1–41.

10. Adang, M.J., Crickmore, N., Jurat-Fuentes, J.L. 2014. Diversity of *Bacillus thuringiensis* crystal

toxins and mechanism of action. In: Dhadialla, T.S. and Gill, S.S., *Advances in Insect Physiology*

vol. 47, Insect midgut and insecticidal proteins. Oxford Academic Press, pp. 39–87.

11. Crickmore, N. 2017. *Bacillus thuringiensis* toxin classification. In: Fiuza, L., Polanczyk, R.,

Crickmore, N. (eds) *Bacillus thuringiensis* and *Lysinibacillus sphaericus*. Springer.

12. WHO report 1999. Guideline specification for bacterial larvicides for public health use. WHO

document WHO/CDS/CPC/WHOPES/99.2. World Health Organization, Geneva.

13. Bravo, A., Pacheco, S., Gomez, I., Garcia-Gomez B., Onofre, J., Soberon, M. 2017. Insecticidal

proteins from *Bacillus thuringiensis* and their mechanism of action. In: Fiuza, L.M., Polanczyk,

R.A., Crickmore, N. (eds). *Bacillus thuringiensis* and *Lysinibacillus sphaericus*, characterization

and use in the field of biocontrol. pp. 53-66.

14. Rabinovitch, L., Vivoni, A.M., Machado, V., Knaak, N., Berlitz, D.L., Polanczyk, R.A., Fiuza,

L.M. 2017. *Bacillus thuringiensis* characterization: morphology, physiology, biochemistry,

pathotype, cellular, and molecular aspects. In: Fiuza, L.M., Polanczyk, R.A., Crickmore, N. (eds).

Bacillus thuringiensis and *Lysinibacillus sphaericus*, characterization and use in the field of

biocontrol. pp. 1-18.

15. Palma, L., Muñoz, D., Berry, C., Murillo, J., Caballero, P. 2014. *Bacillus thuringiensis* toxins:

an overview of their biocidal activity. *Toxins*, 6(12): 3296–3325.

16. Ben-Dov, E., Zaritsky, A., Dahan, E., Barak, Z., Sinai, R., Manasherob, R., Khamraev, A.,

Troitskaya, E., Dubitsky, A., Berezina, N., Margalith, Y. 1997. Extended screening by PCR for

seven cry-group genes from field-collected strains of *Bacillus thuringiensis*. *Applied and*

Environmental Microbiology, 63(12): 4883–4890.

17. Ben-Dov, E., Nissan, G., Pelleg, N., Manasherob, R., Boussiba, S., Zaritsky, A. 1999. Refined,

circular restriction map of the *Bacillus thuringiensis* subsp. *israelensis* plasmid carrying the

mosquito larvicidal genes. *Plasmid*, 42(3): 186-191.

18. Berry, C., O'Neil, S., Ben-Dov, E., Jones, A.F., Murphy, L., Quail, M.A., Holden, M.T.G.,

Harris, D., Zaritsky, A., Parkhill, J. 2002. Complete sequence and organization of pBtoxis, the

toxin-coding plasmid of *Bacillus thuringiensis* subsp. *israelensis*. *Applied and Environmental*

Microbiology, 68(10): 5082-5095.

19. Bravo, A., Gill, S.S., Soberon, M. 2007. Mode of action of *Bacillus thuringiensis* Cry and Cyt

toxins and their potential for insect control. *Toxicon*, 49: 423-435.

20. Wei, J., Liang, G., Wang, B., Zhong, F., Chen, L., Khaing, M.M., Zhang, J., Guo, Y., Wu, K.,

Tabashnik, B.E. 2016. Activation of Bt protoxin Cry1Ac in resistant and susceptible cotton

bollworm. *PLoS ONE*, 11(6): e0156560.

21. Schnepf, E., Crickmore, N., Van Rie, J., Lereclus, D., Baum, J., Feitelson, J., Zeigler, D.R.,

Dean, D.H. 1998. *Bacillus thuringiensis* and its pesticidal crystal proteins. *Microbiology and*

Molecular Biology Reviews, 62(3): 775–806.

22. Bravo, A., Likitvivatanavong, S., Gill, S.S., Soberon, M. 2011. *Bacillus thuringiensis*: a story of

a successful bioinsecticide. *Insect Biochemistry and Molecular Biology*, 41(7): 423–431.

23. Obata, F., Kitami, M., Inoue, Y., Atsumi, S., Yoshizawa, Y., Sato, R. 2009. Analysis of the

region for receptor binding and triggering of oligomerization on *Bacillus thuringiensis* Cry1Aa

toxin. *FEBS Journal*, 276: 5949-5959.

24. Caccia, S., Di Lelio, I., La Storia, A., Marinelli, A., Varricchio, P., Franzetti, E., Banyuls, N., Tettamanti, G., Casartelli, M., Giordana, B., Ferre, J., Gigliotti, S., Ercolini, D., Pennacchio, F. 2016. Midgut microbiota and host immunocompetence underlie *Bacillus thuringiensis* killing mechanism. *Proceedings of the National Academy of Sciences of the United States of America*, 113(34): 9486-9491.

25. Glare, T.R., O'Callaghan, M. 2000. *Bacillus thuringiensis*: Biology, ecology and safety. John Wiley & Sons, UK. pp. 350.

26. Rubio-Infante, N., Moreno-Fierros, L. 2016. An overview of the safety and biological effects of *Bacillus thuringiensis* Cry toxins in mammals. *Journal of Applied Toxicology*, 36: 630-648.

27. EFSA Panel on Biological Hazards (BIOHAZ) 2016. Risks for public health related to the presence of *Bacillus cereus* and other *Bacillus* spp. including *Bacillus thuringiensis* in foodstuffs. *EFSA Journal*, doi: 10.2903/j.efsa.2016.4524.

28. Amichot, M., Curty, C., Benguettat-Magliano, O., Gallet, A., Wajnberg, E. 2016. Side effects of *Bacillus thuringiensis* var. *kurstaki* on the hymenopterous parasitic wasp *Trichogramma chilonis*. *Environmenal Science and Pollution Research*, 23: 3097-3103.

29. Renzi, M.T., Amichot, M., Pauron, D., Tchamitchian, S., Brunet, J.-L., Kretzschmar, A., Maini, S., Belzunces, L.P. 2016. Chronic toxicity and physiological changes induced in the honey bee by the exposure to fipronil and *Bacillus thuringiensis* spores alone or combined. *Ecotoxicology and Environmental Safety*, 127: 205-213.

30. Caquet, T., Roucaute, M., Le Goff, P., Lagadic, L. 2011. Effects of repeated field applications of two formulations of *Bacillus thuringiensis* var. *israelensis* on non-target saltmarsh invertebrates in Atlantic coastal wetlands. *Ecotoxicology and Environmental Safety*, 74: 1122-1130.

31. Duguma, D., Hall, M.W., Rugman-Jones, P., Stouthamer, R., Neufeld, J.D., Walton, W.E. 2015. Microbial communities and nutrient dynamics in experimental microcosms are altered after the application of a high dose of *Bti*. *Journal of Applied Ecology*, 52: 763-773.

32. Lagadic, L., Schäfer, R.B., Roucaute, M., Szöcs, E., Chouin S., de Maupeou, J., Duchet, C.,

Franquet, E., Le Hunsec, B., Bertrand, C., Fayolle, S., Francès, B., Rozier, Y., Foussadier, R.,

Santoni, J.-B., Lagneau, C. 2016. No association between the use of Bti for mosquito control and

the dynamics of non-target aquatic invertebrates in French coastal and continental wetlands.

Science of the Total Environment, 553: 486-494.

33. Venter, H.J., Bøhn, T. 2016. Interactions between Bt crops and aquatic ecosystems: A review.

Environmental Toxicology and Chemistry, 35(12): 2891–2902

34. van Frankenhuyzen, K. 2017. Specificity and cross-order activity of *Bacillus thuringiensis*

pesticidal proteins. In: Fiuza, L.M., Polanczyk, R.A., Crickmore, N. (eds). *Bacillus thuringiensis*

and *Lysinibacillus sphaericus*, characterization and use in the field of biocontrol. pp. 127-172.

35. Bizzarri, M.F., Bishop, A.H. 2008. The ecology of *Bacillus thuringiensis* on the phylloplane:

colonization from soil, plasmid transfer, and interaction with larvae of *Pieris brassicae*.

Microbial Ecology, 56(1): 133-139.

36. Raymond, B., Wyres, K.L., Sheppard, S.K., Ellis, R.J., Bonsall, M.B. 2010. Environmental

factors determining the epidemiology and population genetic structure of the *Bacillus cereus*

group in the field. *PloS Pathogens*, 6(5): e1000905.

37. Hendriksen, N.B., Hansen, B.M. 2002. Long-term survival and germination of *Bacillus*

thuringiensis var. *kurstaki* in a field trial. *Canadian Journal of Microbiology*, 48(3): 256–261.

38. Duchet, C., Tetreau, G., Marie, A., Rey, D., Besnard, G., Perrin, Y., Paris, M., David, J.-P.,

Lagneau, C., Desprès, L. 2014. Persistence and recycling of bioinsecticidal *Bacillus*

thuringiensis subsp. *israelensis* spores in contrasting environments: evidence from field

monitoring and laboratory experiments. *Microbial Ecology*, 67(3): 576–586.

39. Hung, T.P., Truong, L.V., Binh, N.D., Frutos, R., Quiquampoix, H., Staunton, S. 2016a.

Persistence of detectable insecticidal proteins from *Bacillus thuringiensis* (Cry) and toxicity after

adsorption on contrasting soils. *Environmental Pollution*, 208: 318–325.

40. Hung, T.P., Truong, L.V., Binh, N.D., Frutos, R., Quiquampoix, H., Staunton, S. 2016b. Fate of

insecticidal *Bacillus thuringiensis* Cry protein in soil: differences between purified toxin and

biopesticide formulation. *Pest Management Science*, 72: 2247–2253.

41. Enger, K.S., Mitchell, J., Murali, B., Bridsell, D.N., Keim, P., Gurian, P.L., Wagner, D.M. 2018.

Evaluating the long-term persistence of *Bacillus* spores on common surfaces. *Microbial*

Biotechnology, 56: 3073, 12.

42. Couch, T.L., 2000. Industrial fermentation and formulation of entomopathogenic bacteria. In:

Charles, J.-F., Delecluse, A., Nielsen-LeRoux, C. (Eds.), Entomopathogenic Bacteria: From

Laboratory to Field Application. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp.

297–316.43. Brar, S.K., Verma, M., Tyagi, R.D., Valéro, J.R. 2006. Recent advances in

downstream processing and formulations of *Bacillus thuringiensis* based biopesticides. *Process*

Biochemistry, 41(2): 323–342.

44. Setlow, P. 2014. Spore resistance properties. *Microbiology Spectrum*, 2(5): TBS-0003-2012.

45. European Food Safety Authority 2012. Conclusion on the peer review of the pesticide risk

assessment of the active substance *Bacillus thuringiensis* subsp. *Kurstaki* (*strains ABTS 351, PB*

54, SA 11, SA 12, EG 2348). *EFSA Journal*, 10(2): 2540.

46. Bächli, G. 1999-2008. TaxoDros: the database on the taxonomy of Drosophilidae. Available at:

http://www.taxodros.uzh.ch.

47. Tennessen, J.M., Thummel, C.S. 2011. Coordinating growth and maturation - Insights from

Drosophila. *Current Biology*, 21(18): R750–757.

48. Benz, G., Perron, J.M. 1967. The toxic action of *Bacillus thuringiensis* "exotoxin" on

Drosophila reared in yeast-containing and yeast-free media. *Experientia*, 23(10): 871–872.

49. Saadoun, I., Al-Moman, F., Obeidat, M., Meqdam, M., Elbetieha, A. 2001. Assessment of toxic

potential of local Jordanian *Bacillus thuringiensis* strains on *Drosophila melanogaster* and *Culex*

sp (Diptera). *Journal of Applied Microbiology*, 90: 866-872.

50. Khyami-Horani, H. 2002. Toxicity of *Bacillus thuringiensis* and *B. sphaericus* to laboratory

populations of *Drosophila melanogaster* (Diptera: Drosophilidae). *Journal Basic Microbiology*,

42(2): 105-110.

51. Obeidat, M. 2008. Toxicity of local *Bacillus thuringiensis* isolates against *Drosophila*

melanogaster. *World Journal of Agricultural Sciences*, 4(2): 161-167.

52. Obeidat, M., Khymani-Horani, H. and Al-Momani, F. 2012. Toxicity of *Bacillus thuringiensis*

β-exotoxins and δ-endotoxins to *Drosophila melanogaster*, *Ephestia kuhniella* and human

erythrocytes. *African Journal of Biotechnology*, 11(46):10504-10512.

53. Cossentine, J., Robertson, M., Xu, D. 2016. Biological activity of *Bacillus thuringiensis* in

Drosophila suzukii (Diptera: Drosophilidae). *Journal of Economic Entomology*, 0(0): 1-8.

54. Biganski, S., Jehle, J.A., Kleepies, R.G. 2017. *Bacillus thuringiensis* serovar *israelensis* has no

effect on *Drosophila suzukii* Matsumura. *Journal of Applied Entomology*, 142: 33-36.

55. Haller, S., Romeis, J.X.R., Meissle, M., 2017. Effects of purified or plant-produced Cry proteins

on *Drosophila melanogaster* (Diptera: Drosophilidae) larvae. *Scientific Reports*, 1-11.

56. Shorrocks, B. 1991. Competition on a divided and ephemeral resource: a cage experiment.

Biological Journal of the Linnean Society, 43: 211-220.

57. Benado, M., Brncic, D. 1994. An eight-year phenological study of a local drosophilid

community in Central Chile. *Journal of Zoological Systematics and Evolutionary Research*, 32:

51-63.

58. Nunney, L. 1996. The colonization of oranges by the cosmopolitan *Drosophila*. *Oecologia*, 108:

552-561.

59. Mitsui, H., Kimura, M.T. 2000. Coexistence of drosophilid flies: aggregation, patch size

diversity and parasitism. *Ecological Research*, 15: 93-100.

60. Withers, P., Allemand, R. 2012. Les drosophiles de la region Rhône-Alpes (Diptera,

Drosophilidae). *Bulletin de la Société entomologique de France*, 117(4) : 473-482.

61. Zeigler, D.R. 1999. *Bacillus* genetic stock center catalog of strains, 7[th] edition. Part 2: *Bacillus*

thuringiensis and *Bacillus cereus*. pp. 58.

62. Gonzales, J.M.Jr, Brown, B.J., Carlton, B.C. 1982. Transfer of *Bacillus thuringiensis* plasmids

coding for δ-endotoxin among strains of *B. thuringiensis* and *B. cereus*. *Proceedings of the*

National Academy of Sciences of the United States of America, 79: 6951-6955.

63. R Development Core Team 2008. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL

http://www.R-project.org.

64. Bates, D., Maechler, M., Bolker, B., Walker, S. 2015. Fitting linear mixed-effects models using

lme4. *Journal of Statistical Software*, 67(1): 1-48.

65. Kosmidis I (2017). *brglm: Bias Reduction in Binary-Response Generalized Linear Models*. R

package version 0.6.1, http://www.ucl.ac.uk/~ucakiko/software.html.

66. Horton, T., Bretz, F., Westfall, P. 2008. Simultaneous inference in general parametric models.

Biometrical Journal, 50(3): 346-363.

67. Therneau, T.M., Grambsch, P.M. 2000. Modeling survival data: extending the Cox model.

Springer, New York, ISBN 0-387-98784-3.

68. Therneau, T.M. 2015. coxme: Mixed Effects Cox Models. R package version 2.2-5.

https://CRAN.R-project.org/package=coxme

69. Koch, M.S., Ward, J.M., Levine, S.L., Baum, J.A., Vicini, J.L., Hammond, B.G. 2015. The food

and environmental safety of *Bt* crops. *Frontiers in Plant Science*, 6: 283.

70. Bradberry, S.M., Proudfoot, A.T., Vale, J.A. 2004. Glyphosate poisoning. *Toxicological*

Reviews, 23(3): 159-167.

71. de Souza Machado, A.A., Zarfl, C., Rehse, S., Kloas, W. 2017. Low-dose effects:

nonmonotonic responses for the toxicity of a *Bacillus thuringiensis* biocide to *Daphnia magna*.

Environmental Science & Technology, 51: 1679-1686.

72. Vettori, C., Paffetti, D., Saxena, D., Stotzky, G;, Giannini, R. 2003. Persistence of toxins and

cells of *Bacillus thuringiensis* subsp. *kurstaki* introduced in sprays to Sardinia soils. *Soil Biology*

and Biochemistry, 35(12): 1635-1642.

73. Becker, N. 2000. Bacterial control of vector-mosquitoes and black flies. In: Charles, J.-F.,

Delécluse, A., Nielsen-Roux, C. (ed.). Entomopathogenic bacteria: from laboratory to field

application. Kluwer Academic Publishers. p. 383-398.

74. van Frankenhuyzen K. 2013. Cross-order and cross-phylum activity of *Bacillus thuringiensis*

pesticidal proteins. Journal of Invertebrate Pathology, 114(1): 76–85.

75. Stevens, T., Song, S., Bruning, J.B., Choo, A., Baxter, S.W. 2017. Expressing a moth abcc2

gene in transgenic *Drosophila* causes susceptibility to Bt Cry1Ac without requiring a cadherin-

like protein receptor. *Insect Biochemistry and Molecular Biology*, 80: 61-70.

76. George, Z., Crickmore, N. 2012. *Bacillus thuringiensis* applications in agriculture. In: E.

Sansinenea (ed.). *Bacillus thuringiensis* biotechnology. p. 392.

77. Santos, M., Borash, D.J., Joshi, A., Bounlutay, N., Mueller, L.D. 1997. Density-dependent

natural selection in *Drosophila*: evolution of growth rate and body size. *Evolution*, 51(2): 420-

432.

78. Nepoux, V., Haag, C.R., Kawecki, T.J. 2010. Effects of inbreeding on aversive learning in

Drosophila. *Journal of Evolutionary Biology*, 23: 2333-2345.

79. Vantaux, A., Ouattarra, I., Lefèvre, T., Dabiré, K.R. 2016. Effects of larvicidal and larval

nutritional stresses on *Anopheles gambiae* development, survival and competence for

Plasmodium falciparum. *Parasites & Vectors*, 9: 226.

80. Moret, Y., Schmid-Hempel, P. 2000. Survival for immunity: the price of immune system

activation for bumblebee workers. *Science*, 290(5494): 1166-1168.

81. Kutzer, M.A., Armitage, S.A.O. 2016. The effect of diet and time after bacterial infection on

fecundity, resistance, and tolerance in *Drosophila melanogaster*. *Ecology and Evolution*, 6(13):

4229-4242.

82. Andersen, L.H., Kristensen, T.N., Loeschcke, V., Toft, S., Mayntz, D. 2010. Protein and

carbohydrate composition of larval food affects tolerance to thermal stress and desiccation in

adult *Drosophila melanogaster*. *Journal of Insect Physiology*, 56: 336-340.

83. Khazaeli, A.A., Curtsinger, J.W. 2000. Genetic analysis of extended lifespan in *Drosophila*

melanogaster III. On the relationship between artificially selected and wild stocks. *Genetica*, 109:

245-253.

84. Rion, S., Kawecki, T.J. 2007. Evolutionary biology of starvation resistance: what we have

learned from *Drosophila*. *Journal of Evolutionary Biology*, 20(5): 1655-1664.

85. Burger, J.M.S., Buechel, S.D., Kawecki, T.J. 2010. Dietary restriction affects lifespan but not

cognitive aging in *Drosophila melanogaster*. *Aging Cell*, 9: 327-335.

86. Atkinson, W., Shorrocks, B. 1977. Breeding site specificity in the domestic species of

Drosophila. *Oecologia*, 29(3): 223-232.

87. Walsh, D.B., Bolda, M.P., Goodhue, R.E., Dreves, A.J., Lee, J., Bruck, D.J., Walton, V.M.,

O'Neal, S.D., Zalom, F.G. 2011. *Drosophila suzukii* (Diptera: Drosophilidae): invasive pest of

ripening soft fruit expanding its geographic range and damage potential. *Journal of Integrated*

Pest Management, 2(1): DOI: 10.1603/IPM10010.

88. Delbac, L., Rusch, A., Rouzes, R., Ravidat, M.-L., Launes, S., Thiéry, D. 2014. *Drosophila*

suzukii est-elle une menace pour la vigne? *Phytoma*, 679 : 16-21.

89. Poyet, M., Eslin, P., Héraude, M., Le Roux, V., Prévost, G., Givert, P., Chabrerie, O. 2014.

Invasive host for invasive pest: when the Asiatic cherry fly (*Drosophila suzukii*) meets the

American black cherry (*Prunus serotine*) in Europe. *Agricultural and Forest Entomology*, 1-9,

DOI: 10.1111/afe012052.

90. Poulin, B., Lefebvre, G., Paz, L. 2010. Red flag for green spray: adverse trophic effects of *Bti*

on breeding birds. *Journal of Applied Ecology*, 47: 884-889.

Table 1. Results of statistical analyses to assess the effect of the dose of formulation/spore

production and its interaction with the treatment, the larval instar, the experiment, the sex, the fly

strain and the fly species when appropriate. See figures for *post hoc* comparisons of the doses with

the control dose.

Source of variation/Data	χ^2 / Deviance	d.f.	P value
Development on *Btk* Delfin® A, *Btk* 4D22, *Bti* Vectobac®, *Bacillus subtilis*			
Emergence rate			
Dose × Treatment	285.7	20	**< 0.0001**
Dose for each treatment:			
- Delfin® A	237.5	6	**< 0.0001**
- 4D22	7.0	7	0.40
- Vectobac®	165.8	5	**< 0.0001**
- *B. subtilis*	1.9	6	0.93
Developmental time			
Dose × Treatment	220.8	19	**< 0.0001**
Dose for each treatment:			
- Delfin® A	68.8	6	**< 0.0001**
- 4D22	16.08	7	**0.024**
- Vectobac®	37.5	6	**< 0.0001**
- *B. subtilis*	13.5	7	0.060
Development on *Btk* Delfin® B and Scutello DF (dose effect)			
Emergence rate			
- Delfin® B	151.2	2	**< 0.0001**
- Scutello DF	105.1	2	**< 0.0001**
Developmental time			
- Delfin® B	2.5	1	0.12
- Scutello DF	30.9	2	**< 0.0001**
Role of formulation components in the development alterations (dialysis)			
Dose effect			
Emergence rate	459.8	3	**< 0.0001**
Developmental time	13.7	2	**0.0011**
Survival of larval stages on Delfin® A			
Cumulative survival			
Dose × Larval instar	16.2	5	**0.0063**
Dose for each instar:			
- late 1st instar	87.4	5	**< 0.0001**
- late 2nd instar	25.7	5	**0.0001**
24-hour survival			
Dose × Larval instar	15.9	5	**0.007**
Dose for each instar:			
- late 1st instar	55.9	5	**< 0.0001**
- late 2nd instar	3.76	5	0.58
Adult fitness-related traits after development on Delfin® A			
Longevity			
Experiment	20.1	1	**< 0.0001**
- *1st experiment:*			
Dose	12.3	3	**0.0065**
Sex	35.0	1	**< 0.0001**
(e^β coefficient males *vs* females ± se: 0.55 ± 0.16)			
Dose × Sex	20.4	3	**0.00014**
Sexes analyzed separately			
- females	12.0	3	**0.0073**
(e^β coefficients *vs* control ± se: 5×10^6: 1.05 ± 0.17, 5×10^7: 0.71 ± 0.16, 10^8: 0.60 ± 0.21)			
- males	20.4	3	**0.00014**
(e^β coefficients *vs* control ± se: 5×10^6: 0.80 ± 0.16, 5×10^7: 0.66 ± 0.16, 10^8: 1.53 ± 0.18)			

818
819

Source of variation/Data	χ^2 / Deviance	d.f.	P value
Adult fitness-related traits after development on Delfin® A			
- 2nd experiment:			
Dose	16.5	3	**0.00090**
Sex	31.5	1	**< 0.0001**
(e^β coefficient males *vs* females ± se: 0.45 ± 0.22)			
Dose × Sex	0.69	3	0.88
Sexes analyzed separately			
- females	13.2	3	**0.0043**
(e^β coefficients doses *vs* control ± se: 5×10^6: 0.92 ± 0.22, 5×10^7: 0.63 ± 0.21, 10^8: 0.51 ± 0.21)			
- males	7.01	3	**0.072**
(e^β coefficients doses *vs* control ± se: 5×10^6: 1.02 ± 0.22, 5×10^7: 0.70 ± 0.22, 10^8: 0.64 ± 0.22)			
Total numbers of offspring			
Dose × Experiment	28.1	3	**< 0.0001**
Dose for each experiment:			
- 1st experiment	26.3	3	**< 0.0001**
- 2nd experiment	4.1	3	0.25
Development of other strains of *D. melanogaster* on Delfin® A (including Canton S)			
Emergence rate			
Dose × Fly strain	105.5	15	**< 0.0001**
Dose for each fly strain:			
- Canton S	588.6	5	**< 0.0001**
- Nasrallah	745.3	5	**< 0.0001**
- Sefra	900.7	5	**< 0.0001**
- YW1118	636.9	5	**< 0.0001**
Developmental time			
Dose × Fly strain	9.3	12	0.68
Dose for each fly strain:			
- Canton S	40.3	4	**< 0.0001**
- Nasrallah	18.0	4	**0.0012**
- Sefra	27.2	4	**< 0.0001**
- YW1118	28.9	4	**<0.0001**
Development of other *Drosophila* species on Delfin® A			
Emergence rate			
Dose × Fly species	538.2	30	**< 0.0001**
Dose for each species:			
- *D. simulans*	461.0	5	**< 0.0001**
- *D. yakuba*	750.7	5	**< 0.0001**
- *D. hydei*	596.8	5	**< 0.0001**
- *D. immigrans*	726.3	5	**< 0.0001**
- *D. subobscura*	729.6	5	**< 0.0001**
- *D. suzukii*	725.0	5	**< 0.0001**
- *D. busckii*	586.0	5	**< 0.0001**
Developmental time			
Dose × Fly species	59.9	22	**< 0.0001**
Dose for each species:			
- *D. simulans*	25.9	4	**< 0.0001**
- *D. yakuba*	34.7	4	**< 0.0001**
- *D. hydei*	11.5	4	**0.022**
- *D. immigrans*	6.01	3	0.11
- *D. subobscura*	68.8	4	**< 0.0001**
- *D. suzukii*	11.7	3	**0.0085**
- *D. busckii*	58.8	4	**< 0.0001**

Figure legends

Figure 1. Development of *D. melanogaster* Canton S flies on *Btk* and *Bti* commercial formulations. (**a**) Emergence rate (mean ± s.e.m.) and (**b**) developmental time (mean ± s.e.m.) of 20 initial eggs on increasing doses of *Btk* Delfin® A (red triangles), the Cry-free *Btk* 4D22 (open lozenges), the mosquito-targeting *Bti* Vectobac® (green squares), and the non-pathogenic *Bacillus subtilis* (light grey circles). For Vectobac® and *B. subtilis*, $N = 4\text{-}7$ per dose; for Delfin® A and *Btk* 4D22, $N = 9\text{-}12$ for the control, $N = 3$ for 5.10^5 and 10^9, $N = 4\text{-}9$ for 10^6, $N = 7\text{-}14$ from 5.10^6 to 10^8. (**c**) Emergence rate (mean ± s.e.m.) and (**d**) developmental time (mean ± s.e.m.) on increasing doses of the two *Btk* formulations Delfin® B (dark red circles) and Scutello DF (orange squares). $N = 4$ replicates of 20 eggs per dose and formulation, except for controls and 10^8 CFU/g of Delfin® B (9-10 replicates of 20 eggs). Results of *post hoc* comparisons of each dose to the control: $^{.}$ $0.05 < P < 0.1$; * $0.01 < P < 0.05$; ** $0.001 < P < 0.01$; *** $P < 0.001$. (**e**) Immunoblotting with an anti-Cry1A polyclonal antibody on proteins from a suspension of laboratory-produced spores of Cry-free *Btk* 4D22, the three *Btk* formulations Delfin® A, B, Scutello DF, and a suspension of laboratory-produced Cry1A toxins. Red asterisks indicate the Cry protoxins (~130 kDa) and the activated fragments (~60 kDa and ~70 kDa).

Figure 2. Survival of *D. melanogaster* Canton S larval stages on increasing doses of *Btk* Delfin® A. (**a**) Proportion of surviving larvae (mean ± s.e.m.) upon *Btk* exposure from the egg to late 1st instar (open lozenges) and late 2nd instar (black triangles). (**b**) Proportion of surviving larvae (mean ± s.e.m.) upon 24-hour *Btk* exposure of early 1st instar larvae (open lozenges) and 2nd instar larvae (black triangles). $N = 5\text{-}7$ replicates of 20 individuals per dose. Results of *post hoc* comparisons of each dose with the control: * $0.01 < P < 0.05$; ** $0.001 < P < 0.01$; *** $P < 0.001$.

Figure 3. Fitness-related traits of adults (longevity and total offspring number) after

development on *Btk* Delfin® A. (**a**, **d**) Female longevity (mean survival fraction over time ± s.e.m.),

(**b**, **e**) Male longevity (mean ± s.e.m.), and (**c**, **f**) total offspring number (mean ± s.e.m.), measured

on individuals that developed without *Btk* (blue items) and on 5×10^6 CFU/g of *Btk* Delfin® A (green

items), 5×10^7 CFU/g (red items), and 10^8 CFU/g (dark red items). Data from 2 experiments (a-c,

experiment 1; d-f, experiment 2). For each trait, N = 3-5 replicates of 15 males and 15 females per

dose in experiment 1, N = 3 replicates of 15 males and 15 females in experiment 2. Results of *post*

hoc comparisons of each dose with the control: * $0.01 < P < 0.05$; ** $0.001 < P < 0.01$; *** $P < 0.001$.

Figure 4. Development of four *D. melanogaster* strains on increasing doses of *Btk* Delfin® A. (**a**)

Emergence rate (mean ± s.e.m.), (**b**) Developmental time (mean ± s.e.m.) of the strains Canton S

(blue lozenges), Nasrallah (yellow triangles), Sefra (green squares), and YW1118 (red circles). N =

4 groups of 50 eggs per dose and fly strain for each trait. Results of *post hoc* comparisons of each

dose to the control: · $0.05 < P < 0.1$; * $0.01 < P < 0.05$; ** $0.001 < P < 0.01$; *** $P < 0.001$.

Figure 5. Emergence rate of seven *Drosophila* species on increasing doses of *Btk* Delfin® A.

Mean emergence rate (± s.e.m.). N = 4 replicates of 50 eggs per dose for *D. simulans*, *D. yakuba*, *D.*

subobscura, and *D. busckii*, N = 4 replicates of 30 eggs per dose for *D. hydei*, *D. suzukii*, and *D.*

immigrans. Results of *post hoc* comparisons of each dose with the control: · $0.05 < P < 0.1$; *

$0.01 < P < 0.05$; ** $0.001 < P < 0.01$; *** $P < 0.001$.

Figure 6. Developmental time of seven *Drosophila* species on increasing doses of *Btk* Delfin® A.

Mean developmental time (± s.e.m.). N = 4 replicates of 50 eggs per dose for *D. simulans*, *D.*

yakuba, *D. subobscura*, and *D. busckii*, N = 4 replicates of 30 eggs per dose for *D. hydei*, *D. suzukii*,

and *D. immigrans*. Results of *post hoc* comparisons of each dose with the control: * $0.01 < P < 0.05$;

** $0.001 < P < 0.01$; *** $P < 0.001$.

Figure 7. Evaluation of the role of small molecular weight components of *Btk* Delfin[®] A

(dialysis; membrane cut-off: 8-10 kDa) in the altered development of *D. melanogaster* Canton

S. (a) Emergence rate (mean ± s.e.m.) and (**b**) developmental time (mean ± s.e.m.) on increasing

doses of dialyzed Delfin[®] A. $N = 3$ experiments of 4 replicates with 20 eggs per dose for the

emergence rate, $N = 2$ experiments of 4 replicates per dose for the developmental time. Results of

post hoc comparisons of each dose with the control: · $0.05 < P < 0.1$; * $0.01 < P < 0.05$; **

$0.001 < P < 0.01$; *** $P < 0.001$. (**c**) Anti-Cry1A probed immunoblot of non-dialyzed (ND) and

dialyzed (D) suspensions showing the decrease in the amount of ~ 130 kDa protoxins and the

increase in that of ~ 60/70 kDa activated toxins after dialysis.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Emergence rate (y-axis) vs Dose (CFU/g) (x-axis) for seven Drosophila species: *Drosophila simulans*, *Drosophila yakuba*, *Drosophila immigrans*, *Drosophila subobscura*, *Drosophila hydei*, *Drosophila suzukii*, *Drosophila busckii*.

Figure 6

Figure 7