A primer genetic toolkit for exploring mitochondrial biology and disease using zebrafish

Ankit Sabharwal[1], Jarryd M. Campbell[1,2], Zachary WareJoncas[1], Mark Wishman[1], Hirotaka Ata[1,2], Wiebin Liu[1.3], Noriko Ichino[1], Jake D. Bergren[1], Mark D. Urban[1], Rhianna Urban[1], Tanya L. Poshusta[1], Yonghe Ding[1,3], Xiaolei Xu[1,3], Karl J. Clark[1,2], Stephen C. Ekker[1,2]*

1. Department of Biochemistry and Molecular Biology, Mayo Clinic, Rochester, MN 55905, USA

2. Center for Clinical and Translational Sciences, Mayo Clinic, Rochester, MN 55905, USA

3. Division of Cardiovascular Diseases, Mayo Clinic, Rochester, MN 55905, USA

*Corresponding author: Ekker.Stephen@mayo.edu

Keywords:

Abstract

Mitochondria are a dynamic eukaryotic innovation that play diverse roles in biology and disease. The mitochondrial genome is remarkably conserved in all vertebrates, encoding the same 37 gene set and overall genomic structure ranging from 16,596 base pairs (bp) in the teleost zebrafish (*Danio rerio*) to 16,569 bp in humans. Mitochondrial disorders are amongst the most prevalent inherited diseases affecting roughly 1 in every 5000 individuals. Currently, few effective treatments exist for those with mitochondrial ailments, representing a major unmet patient need. Mitochondrial dysfunction is also implicated to be a common component of a wide variety of other human illnesses ranging from neurodegenerative disorders like Huntington's disease and Parkinson's disease to autoimmune illnesses such as multiple sclerosis and rheumatoid arthritis. The electron transport chain (ETC) component of mitochondria is critical for mitochondrial biology and defects can lead to many mitochondrial disease symptoms. Here we present a publicly available collection of genetic mutants created in highly conserved, nuclear-encoded mitochondrial genes in *Danio rerio*. The zebrafish system represents a potentially powerful new opportunity for the study of mitochondrial biology and disease due to the large number of orthologous genes shared with humans and the many advanced features of this model system from genetics to imaging. This collection includes 22 mutant lines in 18 different genes created by locus-specific gene editing to induce frameshift or splice acceptor mutations leading to predicted protein truncation during translation. Also included are 6 lines created by the random insertion of the gene-breaking transposon (GBT) protein trap cassette. All of these targeted mutant alleles truncate conserved domains of genes critical to the proper function of the ETC or genes that have been implicated in human mitochondrial disease. This collection is designed to accelerate the use of zebrafish to study of many different aspects of

37 mitochondrial function with the goal of widening our understanding of their role in biology and

38 human disease.

Introduction

Mitochondria are semi-autonomous organelles critical for eukaryotic cell function. The mitochondrial endosymbiotic genesis origin hypothesis proposes its evolution from an alpha proteobacterial ancestor, *Rickettsia prowazekii* [1,2] that were harnessed by a eukaryotic cell as the host billions of years ago [3]. The proteobacterium became a symbiote of the host cell, bringing with it a system for more efficient generation of cellular energy in the form of ATP.

During the course of evolution, the genetic material of the mitochondria underwent reductive expansion and was transferred in a retrograde fashion to the nuclear genome. The retrograde movement of genes from the mitochondria to its eukaryotic host paved the way for the mitochondria to specialize as energy production organelles rather than consuming energy repetitively replicating its own multi-copy genome. The mitochondrial-encoded genetic material at present is a vestige of the original proteobacterial genome [4–6] meaning despite having DNA of their own, mitochondria rely heavily on the nuclear genes for most of their functions.

Mitochondria have critical functions in metabolism, organ homeostasis, apoptosis and aging. They also play important but still largely mysterious roles in human pathology, as demonstrated by the enormous biological variation and diverse disorders in patients with mitochondrial disease [7–10]. Nearly every organ system can be compromised, but with highly variable and complex physiological and biochemical outcomes. Imaging and basic science of mitochondria showcase how this highly dynamic organelle responds differentially to extrinsic and intrinsic biological signals. However, understanding how mitochondria function in normal biology, and how human mitochondrial DNA variations contribute to health and disease, has been hampered by a lack of effective approaches to manipulate the powerhouse of the cell.

The vertebrate mitochondrial chromosome is circular and includes 37 genes, 13 encoding for

protein subunits of the electron transport chain, 22 coding for transfer RNAs, and 2 encoding

ribosomal RNAs (**Figure 1**) [11,12]. The mitochondrial gene order, strand specific nucleotide

bias and codon usage is highly conserved [13]. However, mtDNA encoded genes lack introns

and utilize a divergent genetic code than their nuclear counterparts [14,15]. For instance, AUA

codon codes for methionine as per mitochondrial genetic code, whereas the same sequence codes

for isoleucine in the nuclear genetic code. Similarly, nuclear stop codon UGA is read as the

tryptophan amino acid by the mitochondrial codon cypher.

Mitochondria are unique cellular compartments with different DNA and RNA repair and editing

rules, hampering attempts at directly manipulating these nucleic acid components. For example,

DNA nucleases that introduce double stranded breaks and subsequent repair in nuclear DNA

induce the degradation of mtDNA [16–18]. Indeed, none of the canonical DNA repair pathways

found in the nucleus have been shown to be active in mitochondria [18–20]. Finally, no system

has demonstrated the ability to deliver exogenous DNA or RNA to mitochondria, restricting the

tools available for mtDNA editing [21,22]. All of these factors are distinct from the nuclear

genome, making mtDNA a far less accessible genome for traditional gene editing methods and

reagents.

The diverse functions that mitochondria are capable of, including oxidative phosphorylation,

would not be possible with the small subset of 13 proteins encoded in mtDNA. Discoveries led

by high throughput proteomic approaches have enhanced our knowledge of the mitochondrial

proteome [23–27]. There are approximately 1158 nuclear-encoded proteins that localize to the

mammalian mitochondria, exerting a dual genetic control via its nuclear counterpart. Nuclear-

encoded mitochondrial proteins on the basis of their mitochondrial function can be broadly

classified into different categories including oxidative phosphorylation, energy production,

membrane dynamics, genome maintenance, and ion/metabolite homeostasis.

Unlike the nuclear genome, where most cells have only two copies, each cell can harbor

thousands of copies of mtDNA, depending on environmental needs, making mitochondrial

genome engineering a population genetics challenge. When every mtDNA molecule is identical

with non-mutagenic variants, the cell is in a genetic state, called homoplasmy, and the

mitochondria functions normally. In people with mtDNA-based mitochondrial disease, mtDNA

genomes harboring pathogenic mutations typically co-exist with the healthy genomes in a state

called heteroplasmy. Homoplasmy may also arise when the cell harbors only mutant mtDNA

molecules, resulting in severe clinical manifestations. [28]. The ratio of pathogenic-harboring to

nonpathogenic-harboring mtDNA is critical to mitochondrial disease onset. A threshold exists

for most mitochondrial disease, at which, the healthy mtDNA genomes can no longer

compensate for the pathogenic genomes, resulting in disease onset leading to breakdown of

oxidative phosphorylation and other mitochondrial function(s) [29].

Mitochondrial disorders are a heterogeneous group of clinical manifestations resulting from

either inherited or spontaneous mutations in mtDNA or nDNA leading to altered structure or

function of the proteins or RNA that reside in mitochondria [6,10,30,31]. The pathophysiology

of mitochondrial disorders involves functions of both the nuclear and mitochondrial genomes,

conferring additional complexity to the manifestation of the syndromes. The prevalence of

mitochondrial disorders is about 1 in 5000 live births; however, the prevalence is often

influenced by the presence of founder mutations and consanguinity in populations [31–33].

These conditions range in severity dependent on the specific mutation and its prevalence and

localization within the patient. In addition to diseases specifically progenerated by mitochondrial

defects, mitochondrial dysfunction has also been implicated as a cofactor or by-product of many

other conditions; including cancer, immune diseases, developmental delays and

neurodegenerative condition such as Alzheimer's [34–38].

Mitochondrial dysfunction tends to affect primarily high energy systems and can therefore have

devastating effects on a wide range of body systems including brain function, liver function,

vision, hearing, immune function, and all muscle types [31,39]. The identification and

characterization of novel mitochondrial genes in human genetic disorders have also enhanced our

knowledge of mitochondrial function [40–44]. However, despite their known prevalence and

severity, mitochondrial diseases remain understudied as compared to other genetic conditions,

and the options for treatment are limited.

The conserved roles of mitochondria have been traditionally uncovered using accessible model

systems such as the yeast *S.cerevisiae* [45–48] and invertebrates including *C.elegans* [49–52]

and *D. melanogaster* [53–56]. Vertebrate mitochondria, however, are known to have additional

innovations and encode unique subunits not present in invertebrate models [5,45,57–67]. Most

exploration to date of vertebrate mitochondrial biology has been conducted either in cell culture

or in mouse models. However, cells in a dish are missing important environmental cellular and

organismal context, whereas mouse mitochondrial experimental work can be hindered by limited

population size and invasive imaging techniques.

We provide here an initial genetic toolkit for modeling mitochondrial biology and disease in

Danio rerio (zebrafish) in an effort to help catalyze the use of this invaluable model organism in

mitochondrial research. The zebrafish is a tropical freshwater teleost that has proven to be an

invaluable resource for the study of human disease and genetics [68–70]. Principal among the

many research amenable characteristics of zebrafish are their high genetic orthology to humans,

high fecundity, their optical transparency (in the embryonic stage) that lends itself to facile

imaging, and the ease of reagent delivery through microinjection into the single celled embryo,

which is a full millimeter across before the first mitotic division. Zebrafish also serve as a

potentially powerful vertebrate model organism to study human mitochondrial disorders because

of conserved mitochondrial genome and mitochondrial genetic machinery. Zebrafish and human

mitochondrial chromosomes display ~65% sequence identity at the nucleotide level, and share

the same codon usage, strand specific nucleotide bias and gene order [12] **(Figure 1)**.

Over the recent years, zebrafish research has helped shed light on mitochondrial biology and

further developed our understanding of the mechanisms of mitochondrial-associated pathology

[71–76]. Zebrafish have also been successfully used as a model system to study mitochondrial

targeting drugs with implications in development and cardiovascular function [71,77,78]. These

drugs can simply be dissolved in the water housing the larvae and offer an advantage of large

sample size with minimal volumes of drug administered. Drug-screening studies have aided in

understanding the pathogenesis of various diseases and helped to identify targets for treatment

[77].

To help engender an expansion of zebrafish deployment in mitochondrial research, we present a

panel or "starter-pack" toolkit of genetic mutants made in nuclear-encoded mitochondrial genes

in zebrafish. This mutant panel, which we have named the Marriot Mitochondrial Collection

(MMC), consists of 28 zebrafish lines with mutations in 23 different mitochondrial genes. The

mutant collection focusses primarily on genes that encode for components of the energy

generating electron transport chain with at least one mutant in each complex of the ETC. Other

mutants consist of assembly factors, protein chaperones that manage mitochondrial membrane

traffic, and genes related to mitochondrial replication. These mutants were made either by the

use of targeted endonuclease [79], or curated from our research group's library of randomly

generated insertional mutants [80]. We hope that this collection, in addition to being intrinsically

useful, will also help serve as a primer to the modeling of mitochondrial biology and disease in

zebrafish.

Methods

Zebrafish Handling

All animal work was conducted under Mayo Clinic's institutional animal welfare approvals

(IACUC number: A34513-13-R16).

Identification of zebrafish orthologs having putative mitochondrial function

A previous study combining discovery and subtractive proteomics with computational,

microscopy identified 1098 mouse genes that could encode for proteins residing in mitochondria

[24]. They further identified 1013 human orthologs for these genes, providing an initial

inventory of the genes coding for proteins resident in mitochondria. Using literature assessment

and HUGO database curation approaches, we identified 97 proteins that are involved in the

biogenesis and assembly of electron transport chain in mitochondria. Using zebrafish orthologs

of human genes from ZFIN (Zebrafish Information Network), 93 zebrafish mitochondrial

orthologs were identified. These orthologs were systemically annotated with respect to clinical

phenotype by extensive mining from PubMed based published case reports and OMIM database

(Supplementary Table 1).

Mutant Generation

Mutant lines were created by one of two methods, either through the targeted use of

Transcription Activator Like Effector Nucleases (TALENs) [79] or by screening Gene Breaking

Transposon (GBT) lines for integrations into mitochondrial genes [80]. Reagents were delivered

in both methods by the microinjection of either TALEN pairs or GBT transposon and Tol2

transposase into single cell zebrafish embryos.

TALEN Design/Assembly/Delivery

The TALEN mutants in this collection were originally generated as part of a previously

published study [79]. In brief, TALEN pairs were designed using the Mojohand software

platform [81] (www.talendesign.org) to target highly conserved, and therefore likely functionally

important, areas of nuclear-encoded mitochondrial genes. TALEN RVDs were then cloned into

pT3Ts-GoldyTALEN (TALEN vector with a T3 transcriptional promoter for in vitro

transcription) using the FusX rapid TALEN assembly system [79], which uses the RVD

definitions: HD=C, NN=G, NI=A, NG=T. Following assembly, mRNA was synthesized in vitro

using the mMessage Machine T3 kit (Ambion) and extracted by a phenol-chloroform extraction

as prescribed in the mMessage Machine manual. The extracted mRNA was then delivered into

single cell zebrafish embryos at 100pg doses (50pg per TALEN arm) by microinjection.

TALEN Mutant Screening

Following microinjection, genomic DNA was extracted from F0 larvae three days post

fertilization (dpf) by sodium hydroxide extraction. DNA for eight individual larvae was analyzed

for NHEJ activity at the TALEN target site by Restriction Fragment Length Polymorphism

(RFLP) analysis. Groups with high reported NHEJ activity by RFLP were raised to adulthood

and outcrossed to create an F1 generation. Suspected NHEJ mutants were further verified by

Sanger sequencing. F1 larvae demonstrating NHEJ mutations confirmed by both RFLP and

Sanger sequencing were raised to adulthood. Fin biopsies were performed on these adults and

DNA was extracted by sodium hydroxide extraction.

199 Gene Breaking Transposon System

The Gene Breaking Transposon System of protein trap system and a complete repository of

protocols for the creation and screening of GBT mutant lines has been described [80]. In short,

protein trap transposons were delivered in combination with mRNA for Tol2 transposase (25pg

each) into single cell zebrafish embryos by microinjection. Embryos were screened for GFP

fluorescence at 3-4dpf and classified broadly into three classes [80]. Class three embryos, those

with whole body GFP expression, were raised to adulthood and outcrossed to non-transgenic

lines to create an F1 generation. mRFP-expressing F1 embryos were sorted by expression

pattern, assigned a GBT number, and raised to adulthood. These adult fish were then outcrossed

to non-transgenic lines to create an F2 generation upon which all subsequent propagation, testing

and imaging was conducted.

To determine genes tagged by the protein trap system, rapid amplification of cDNA ends

(RACE) was performed as described [82] with minor updates to primer sequences. cDNA was

generated using a transposon-specific primer (5R-mRFP-P0) against 250ng of total mRNA in the

reverse transcription reaction. PCR was then performed using the following gene-specific

primers: 5R-mRFP-P1 and 5R-mRFP-P2. The resulting products were TA cloned for further

amplification and then sequence verified for in-frame mRFP fusions by Sanger sequencing.

In some cases, inverse PCR was also conducted to RACE PCR as described [80]. Whole

genomic DNA was extracted from individual F2 embryos using a sodium hydroxide extraction

and 800ng was digested in a combination reaction using AvrII, NheI, SpeI, and XbaI restriction

enzymes. Approximately 200ng of the product of this digestion was self-ligated and used as a

template for PCR using the following nested and primary primers. 5' side: 5R-mRFP-P1 and 5R-

mRFP-P2 paired with INV-OPT-P1 and INV-OPT-P2, respectively. 3' side: 5R-GFP-P1 and 5R-

GFP-P2 with Tol2-ITR(L)-O1 and Tol2-ITR(L)-O3, respectively. The products of the final

nested reaction were gel-extracted, cloned, and sequenced. Prospective in frame mRFP fusions

were further verified by comparing suspected protein trap allele to GBT expression pattern and

by PCR against DNA or cDNA from mRFP carrier siblings versus non-carrier siblings.

Protein trap lines as verified by these methods were catalogued for specific genomic fusions

using the National Center for Biotechnology Information's (NCBI) Homologene Database.

Human orthologs for tagged genes were further identified using blastX searches against the

human genome. GBT lines with protein trap fusions to nuclear encoded mitochondrial genes

with human orthologues have been included in this collection.

***In silico* analysis to determine protein homology and alteration of DNA sequence by**

TALENs and GBT in zebrafish mutants

Human amino acid sequences were compared to both the zebrafish wild type sequence and each

specific zebrafish mutant sequence associated with each allele. Amino acid sequence information

for a particular gene was gathered for humans from https://www.uniprot.org/ along with the wild

type amino acid sequence for the zebrafish. The first/most common isoform for each entry was

used for this analysis. Any added information regarding functional domains or regions was also

gathered. A protein-protein BLAST was conducted using https://blast.ncbi.nlm.nih.gov/Blast.cgi

to compare the sequences of the human and wild type zebrafish amino acid sequences. BLAST

search settings used included the blastp algorithm. Regions of low and high homology were then

mapped out for the sequences.

For zebrafish mutant to human comparisons, the cDNA or DNA sequences were gathered via

sequencing and put through the translate tool at https://web.expasy.org/translate/. The standard

genetic code was used for conversion. Stop codons arising from the frameshifted DNA sequence

were found in all mutants. The BLAST analysis conducted on the mutants was taken from the

starting methionine to the first encountered stop codon. Regions of low and high homology were

then mapped for each sequence.

Access to all reported reagents – zebrafish and sequences

All listed tools are immediately available through the Mayo Clinic Zebrafish Facility, and all fish

lines will be available via ZIRC. Sequences needed for genotyping and related metadata are

currently on zfishbook [83].

RESULTS

Generation of zebrafish mutant collection

We generated zebrafish mutants for a wide selection of nuclear-encoded mitochondrial genes by

employing two genetic engineering approaches, gene breaking transposons and TALENs. The

collection of zebrafish mutants is referred as the Marriot Mitochondrial Collection (MMC) and

comprises of 28 mutants for 23 nuclear-encoded mitochondrial proteins **(Figure 2)**. Out of these,

22 were created by TALEN indel mutagenesis and 6 were created by gene breaking trap

mutagenesis. The mutants include proteins from all known functional pathways involved in

mitochondrial homeostasis and ATP generation **(Figure 2)**.

Broadly, the pathways can be classified as subunits of oxidative phosphorylation complexes,

chaperones for assembly of oxidative phosphorylation proteins, maintenance proteins for the

mitochondrial genome (replication, mtRNA folding and translation), calcium homeostasis and

mitochondrial protein import. All of these genes have human orthologs, nearly all with known

mutations in which lead to severe clinical manifestations such as Leigh syndrome,

cardiomyopathy, progressive external ophthalmoplegia, and oxidative phosphorylation

deficiency (**Table 1**).

Table 1: List of nuclear-encoded genes for which zebrafish mutants were generated as part

of MMC collection: The current MMC resource list summarizes information on human gene,

zebrafish ortholog, mouse ortholog, relevant clinical phenotypes and diseases, protein function,

and OMIM ID. Green = Created by TALEN Indel Mutagenesis, Orange = Curated from GBT

protein trap lines. Allele designators are highlighted in bold for respective zebrafish paralog.

(Chr- Chromosome; OMIM ID: Online Mendelian Inheritance in Man ID; NA-not available).

Approved symbol	Approved name	Human chr	Zebrafish orthologs	Zebrafish chr	Allele designator	Mouse orthologs	Mouse chr	Clinical phenotype observed in humans	Disease	Biological function	OMIM ID
NDUFAB1	NADH:ubiquinone oxidoreductase subunit AB1	16	*ndufab1a* ***ndufab1b***	1 3	*mn0135	Ndufab1b	7	-na-	-na-	Oxidative phosphorylation enzymes - Complex I	603836
NDUFAF6	NADH:ubiquinone oxidoreductase complex assembly factor 6	8	*ndufaf6*	16	mn0117	Ndufaf6	4	Focal right-hand seizures, ataxia, lactic acidosis, exercise intolerance, weakness and muscle tension	Leigh syndrome, mitochondrial complex I deficiency	Oxidative phosphorylation complex assembly - Complex I	612392
NDUFS4	NADH:ubiquinone oxidoreductase subunit S4	5	*ndufs4*	5	mn0118	Ndufs4	13	Mitochondrial complex I deficiency	Mitochondrial complex-I deficiency	Oxidative phosphorylation enzymes - Complex I	602694
SDHA	Succinate dehydrogenase complex flavoprotein subunit A	5	*sdha*	19	mn0121	Sdha	13	Dyspnea, cardiomegaly, cardiomyopathy, nystagmus, hypotonia, gastrointestinal stromal tumors, paragangliomas, pheochromocytoma, psychomotor regression and severe hyperandrogenism	Mitochondrial complex-II deficiency; Cardiomyopathy, Leigh syndrome; Paraganglioma	Oxidative phosphorylation enzymes - Complex II	600857

Approved symbol	Approved name	Human chr	Zebrafish orthologs	Zebrafish chr	Allele designator	Mouse orthologs	Mouse chr	Clinical phenotype observed in humans	Disease	Biological function	OMIM ID
BCS1L	BCS1 homolog, ubiquinol-cytochrome C reductase complex chaperone	2	bcs1l	9	mn0102	Bcs1l	1	Growth retardation, aminoaciduria, cholestasis, iron overload, lactic acidosis, early death, hypotonia, encephalopathy, microcephaly, psychomotor regression, hypoglycemia and hepatic failure	Björnstad syndrome: GRACILE syndrome; Leigh syndrome; Mitochondrial complex III deficiency, nuclear type 1	Oxidative phosphorylation complex assembly - Complex III	603647
UQCRQ	ubiquinol-cytochrome c reductase complex III subunit VII	5	uqcrq	14	mn0128 mn0129	Uqcrq	11	Severe psychomotor retardation and extrapyramidal signs, dystonia, athetosis, ataxia, mild axial hypotonia and marked global dementia	Mitochondrial complex III deficiency	Mitochondrial complex III: ubiquinol-cytochrome c reductase complex subunits	612080
COX10	Cytochrome c oxidase assembly factor heme A: farnesyltransferase COX10	17	cox10	12	mn0107 mn0108	Cox10	11	Muscle weakness, hypotonia, ataxia, ptosis, pyramidal syndrome, status epilepticus, lactic acidosis, hypertrophic cardiomyopathy, hypoglycemia and metabolic acidosis	Leigh syndrome due to mitochondrial COX IV deficiency; Mitochondrial complex IV deficiency	Oxidative phosphorylation complex assembly - Complex IV	602125
COX4I2	Cytochrome C Oxidase Subunit 4I2	20	cox4i2b	23	mn0104	Cox4i2	2	Exocrine pancreatic insufficiency, dyserythropoietic anemia, and calvarial hyperostosis	Exocrine pancreatic insufficiency, dyserythropoietic anemia, and calvarial hyperostosis	Oxidative phosphorylation complex enzymes - Complex IV	607976
SURF1	SURF1, Cytochrome C Oxidase Assembly Factor	9	surf1	5	mn0123	Surf1	2	Childhood onset neuropathy, lactic acidosis, mild intellectual disability ataxia, facial dysmorphism, encephalopathy, hypotonia, cerebellar ataxia, deafness, ophthalmoplegia, growth retardation and nystagmus	Charcot-Marie-Tooth disease, type 4K; Leigh syndrome, due to COX IV deficiency	Involved in biogenesis of cytochrome c oxidase complex	185620
ATP5F1E	ATP synthase F1 subunit epsilon	20	atp5f1e	6	mn0101	Atp5f1e	2	Neonatal-onset lactic acidosis, 3-methylglutaconic aciduria, mild mental retardation and peripheral neuropathy	Mitochondrial complex (ATP synthase) deficiency	Oxidative phosphorylation complex assembly - ATP synthase	606153
TMEM70	Transmembrane protein 70	8	tmem70	2	mn0126 mn0127	Tmem70	1	Lactic acidosis, encephalopathy, histiocytoid cardiomyopathy, microcephaly, hypotonia, facial dysmorphism and 3-methylglutaconic aciduria psychomotor delay and hyperammonemia	Mitochondrial complex V (ATP synthase) deficiency	Biogenesis of mitochondrial ATP synthase	612418

Approved symbol	Approved name	Human chr	Zebrafish orthologs	Zebrafish chr	Allele designator	Mouse orthologs	Mouse chr	Clinical phenotype observed in humans	Disease	Biological function	OMIM ID
COQ2	Coenzyme Q2, Polyprenyltransfe--rase	4	coq2	5	mn0106	Coq2	5	Multiple system atrophy, unsteadiness of gait, nystagmus, gait ataxia, dysarthria, speech difficulty, dysmetria, lactic acidosis, urinary dysfunction and nystagmus	Coenzyme Q10 deficiency, primary; Multiple system atrophy	Biosynthesis of CoQ, coenzyme in mitochondrial respiratory chain	609825
PDSS2	Decaprenyl diphosphate synthase subunit 2	6	pdss2	13	mn0120	Pdss2	10	Hypotonia, seizures, cortical blindness, lactic acidosis, encephalopathy and nephrotic syndrome	Coenzyme Q10 deficiency	Involved in biosynthesis of coenzyme Q	610564
MICU1	Mitochondrial calcium uptake 1	10	micu1	13	mn0132	Micu1	10	Proximal muscle weakness and learning disabilities	Myopathy with extrapyramidal signs	Regulator of mitochondrial calcium uptake	605084
SMDT1	Single-pass membrane protein with aspartate rich tail	22	smdt1a *smdt1b	3 1	*mn0122	Smdt1	15	-na-	-na-	Core regulatory subunit of calcium channel in mitochondrial inner membrane	615588
MCU	Mitochondrial calcium uniporter	10	mcu	13	mn0111	Mcu	10	-na-	-na-	Calcium transporter and mediates calcium uptake in mitochondria	614197
TK2	Thymidine kinase 2, mitochondrial	16	tk2	7	mn0125	Tk2	8	Infantile onset fetal encephalomyopathy, failure to thrive, muscle weakness, mild facial weakness, myopathy, axonal neuropathy, brain atrophy, hypertrophic cardiomyopathy, regression of motor development, psychomotor arrest, dysphonia, dysphagia, ptosis, impaired gait, bilateral ptosis, hypotonia, lactic acidosis and epilepsy	Progressive external ophthalmoplegia; Mitochondrial DNA depletion syndrome 2 (myopathic type)	Required for mitochondrial DNA synthesis	188250
MTFMT	Mitochondrial methionyl-tRNA formyltransferase	15	mtfmt	7	mn0113 mn0114	Mtfmt	9	Psychomotor developmental delay, renal dysplasia, mild facial dysarthria and ataxia	Combined oxidative phosphorylation deficiency	Catalyzes the formylation of methionyl-tRNA	611766
MCU	Mitochondrial calcium uniporter	10	mcu	13	mn0599gt	Mcu	10	-na-	-na-	Calcium transporter and mediates calcium uptake in mitochondria	614197
LRPPRC	Leucine rich pentatricopeptide repeat containing	2	lrpprc	13	mn0235gt	Lrpprc	17	Delayed psychomotor development, mental retardation, mild dysmorphic facial features, hypotonia, ataxia,development of lesions in the brainstem and basal ganglia, seizures, dysphagia and hypertrophic cardiomyopathy	Leigh syndrome, French-Canadian type	Involved in translation of mitochondrial encoded cox subunits and mediation of folding of mitochondrial transcriptome	607544

Approved symbol	Approved name	Human chr	Zebrafish orthologs	Zebrafish chr	Allele designator	Mouse orthologs	Mouse chr	Clinical phenotype observed in humans	Disease	Biological function	OMIM ID
MRPS18B	Mitochondrial ribosomal protein S18B	6	mrps18b	19	mn0425gt	Mrps18b	17	-na-	-na-	Part of small 28S subunit of mitoribosome	611982
TIMM50	Translocase of inner mitochondrial membrane 50	19	timm50	15	mn0906gt	Timm50	7	Severe intellectual disability, seizure and 3-methylglutaconic aciduria	3-methylglutaconic aciduria, type IX; Mitochondrial complex V deficiency	Subunit of TIM23 inner mitochondrial membrane complex and recognizes mitochondrial targeting signal or pre-sequence	607381
OGDH	Oxoglutarate dehydrogenase	7	ogdha *ogdhb	8 10	*mn0281gt	Ogdh	11	Colorectal cancer	Alpha-ketoglutarate dehydrogenase deficiency	Catalyzes the conversion of 2-oxoglutarate to succinyl-CoA and CO_2	613022
IDH2	Isocitrate dehydrogenase (NADP (+)) 2, mitochondrial	15	idh2	18	mn0268gt	Idh2	7	Acute myeloid leukemia and abnormal production of D-2-hydroxyglutaric acid	D-2-hydroxyglutaric aciduria 2	Catalyzes the conversion of isocitrate to 2-oxoglutarate	147650

TALEN- and GBT-mediated targeting of nuclear-encoded mitochondrial genes

Amino acid analysis using protein-protein BLAST functions between the wild type human, wild

type zebrafish, and mutant zebrafish sequences showed consistent results between the three

conditions. The comparison of wild type human and wild type zebrafish sequences showed high

areas of homology following the mitochondrial targeting domain in almost every gene, with 80-

100% similarity in catalytic or active domains of the transcripts. Analysis of the mutant zebrafish

and human wild type comparison showed a range of difference with predicted frameshift

mutations leading to truncation of the protein very early in the transcript (**Figure 3A-X**). Of the

alleles created by TALEN mutagenesis in the MMC collection, all but one (*micu1*) showed a

predicted frameshift mutation leading to a truncation event immediately following the NHEJ-

mediated insertion or deletion. The GBT mutants showed high levels of homology prior to the

transposon integration site followed by a nearly complete loss of normal transcript levels

following splicing into the GBT cassette.

Discussion

In a multicellular organism, each cell is able to carry out its functions due to well-orchestrated cross-talk between the nuclear and mitochondrial genomes. Many mitochondrial functions like energy production, genome maintenance, ion/metabolite homeostasis, membrane dynamics and transport of biomolecules can be attributed to approximately 1158 known nuclear proteins residing in the mitochondria [24,27]. Due to advancements in proteomic technologies in recent years, there has been a surge in documentation of many characterized and uncharacterized mitochondrial proteins. However, correlation between mitochondrial localization of these proteins and their physiological significance in disease progression remains largely unexplored. Out of the 1158 nuclear encoded proteins, only 245 have functional evidence in mitochondrial clinical manifestations [39]. For the remaining 913 proteins, mitochondrial involvement in disease progression has yet to be demonstrated. The obstacles in mitochondrial genetic research, and thus delays in finding effective treatments, are primarily due to the limited tools available to mitochondrial researchers, specifically the small number of available model systems and animals.

Most mitochondrial research thus far has made use of model systems that each present their own unique challenge to accurate study of human disease. Yeast, one of the most common laboratory eukaryotes, have been incredibly useful in mitochondrial research, but unlike humans and other animals they lack complex I [45]. Complex 1 deficiency is the most frequent among mitochondrial disorders caused by mutations in 28 out of 48 genes contributing to its assembly

and biogenesis [84]. Also, complex 1 is responsible for the most ROS generation, which acts as a

cue in various signaling pathways [85].

Human cell culture has been valuable in exploring mitochondrial disease models, principally

through cytoplasmic hybrids that are made by replacing the mitochondria of an immortalized cell

line with mutated patient mitochondria [86–89]. However, while these cells are a more faithful

representation of human mitochondrial activity than yeast, they have different barriers to

accuracy. First, immortalized cells, and cultured cells in general, often have modified metabolic

pathways compared to cells *in vivo*. Second, as is common to all cell culture research, cells in a

dish cannot accurately represent the complex interactions and systems biology inherent to a

complete organism.

In this study, we aim to support the establishment of the zebrafish as a complementary animal

model for understanding the role of nuclear-encoded mitochondrial proteins in biology and the

pathophysiology of mitochondrial disorders. We propose that zebrafish can be an excellent

model organism to study mitochondrial biology primarily because of their conserved genome,

codon bias and synteny. Other advantages include their amenability to genetic manipulation and

optical clarity, which facilitates direct observation. New gene editing techniques such as

TALENs and CRISPRs have aided in the development of humanized disease models in

organisms such as zebrafish, mice, and even pigs [90–97].

The MMC collection described here encompasses mutations in 23 different nuclear-encoded

genes related to mitochondrial function enabling a diverse study of the important roles the

mitochondria play in cellular biology. The collection includes mutants in all five complexes of

the electron transport chain as well as many other crucial pathways related to protein transport,

metabolite synthesis, mtDNA replication and expression, and calcium homeostasis. The first set

of mutants were created using custom gene editing to exons coding for critical functional

domains of the protein product. These domains were targeted because they shared high levels of

homology to their human ortholog. The second group of mutants consisted of fish that had been

injected with a protein trap transposon system that truncates the expected protein products as

well as sorting and imaging through fluorescent expression patterns *in vivo*. Included here, we

have curated small group of mutants where the GBT integrated into a mitochondrial gene of

interest.

In recent years zebrafish have been used to understand the pathophysiology of human

mitochondrial disorders such as cardiovascular, multisystemic, neurological, and erythropoiesis,

to name a few. Studies for loss of function have been carried out to model left ventricular non-

compaction, a genetically heterogeneous cardiomyopathy caused by mutations in nicotinamide

nucleotide transhydrogenase (*NNT*) gene [98,99]. Transiently suppressing the levels of *nnt* gene

did mirror the symptoms, such as early ventricular malformation and contractility defects,

observed in human patients [99]. Transient suppression of *ndufb11* protein in zebrafish results in

cardiac anomalies, confirming the role of *NDUFB11* in histiocytoid cardiomyopathy

pathogenesis [41].

Parkinson disease (PD) is another example where *parkin* and *pink* knockdown larvae display

mitochondrial fragmentation accompanied by accumulation of oxidative species, mimicking the

phenotype observed in PD patients. Loss of dendritic arborization and peripheral neurons were

also observed in the knockdown animals [72,100–103]. The zebrafish model of Dravet syndrome

shows altered metabolism associated with the pathophysiology [104]. Downregulation of the

sodium channel, voltage-gated, type I like, alpha b (*scnl1ab*) in zebrafish leads to mitochondrial

structural abnormalities accompanied by accumulation of oxidative species [104–106].

Sideroblastic anemia, caused by mutations in mitoferrin (*mfrn1*) gene, has been successfully

modelled in zebrafish [107]. The elucidation of its role in zebrafish as iron importer does shed

some light on iron related defects in progression of anemia, accompanied by mitochondrial

dysfunction. Zebrafish erythrocytes are nucleated and possess organelles such as mitochondria

[108] in their cytoplasm, making them an ideal model organism to study ribosomopathies and

hematological malignancies [109–114]. Mitochondria in erythrocytes may offer valuable insight

into novel biological pathways resident in mitochondria.

Transient infantile liver failure and mitochondrial deafness, caused by mutations in tRNA 5-

methylaminomethyl-2-thiouridylate methyltransferase (*TRMU*) gene [115] have been

phenocopied in zebrafish [116]. Zebrafish *trmu* mutants exhibit abnormal mitochondrial

morphology, deficient oxidative phosphorylation, defects in hair cells and decreases in the steady

state levels of mt-tRNAs, resembling the observations made in patients suffering from this

syndrome [116].

Taking cues from these studies, many of the genes selected here were prioritized on the basis of

the mutations and clinical phenotype reported in patients. Detailed phenotypic analyses will be

described in subsequent studies. The zebrafish mutants encode for genes that have been involved

in myriad of clinical phenotypes such as encephalopathy, nephrotic syndrome, intellectual

disability, psychomotor developmental delay, respiratory chain deficiency, lactic acidosis, iron

overload, microcephaly, hypertrophic cardiomyopathy, etc. The superficial investigations of the

tissue-specific phenotypes associated with, for example, housekeeping nuclear encoded proteins,

have to be refined with organelle level investigations. The repository of mutants encoding for

proteins involved in respiratory chain biogenesis and assembly offers the potential to understand

the moonlighting role of these proteins. Mutations in these proteins are associated with organ-

specific phenotypes such as neurological, paraganglioma, cardiovascular etc. [41,117]. Extensive

energy demand by these tissues is one possible explanation for the progression of these

manifestations; however, the role of these proteins in organ-specific cellular niches or

homeostasis is also a possibility. These models can help to decipher the role of these proteins in

the mitochondrial interactome, when studied *in vivo*. This underpins the utility of this MMC

collection in deciphering the role of mitochondrial proteins in tissue-specific biological

pathways.

Zebrafish are also amenable to small molecule mediated pharmacological modelling of

mitochondrial phenotypes. To test hypoxia as a potential protective therapy in mitochondrial

disorders, Von Hippel-Lindau (*vhl*) null mutant zebrafish, when treated with antimycin mediated

mitochondrial insult, exhibited improved survival. In addition, FG-4592 was found to improve

survival in response to respiratory chain inhibition, possibly due to increase in hypoxia response

[118]. A series of ETC complex specific pharmacological inhibitors such as rotenone (complex

I), azide (complex IV), oligomycin (complex V) and chloramphenicol (mitochondrial protein

translation) have been used to model respiratory chain dysfunction in zebrafish [78,119,120].

Zebrafish larvae display a series of phenotypes, such as developmental arrest, when treated with

rotenone. Treatment with azide induced decreased heart rate, loss of motor function, inability to

respond to tactical stimulation, neurological damage and mortality. Organ specific manifestations

such as neurological and behavioral dysfunction have been reported in 6-7 days post fertilization

(dpf) zebrafish larvae upon administration of a titred drug concentration [120]. These

observations tilt the scales in favor of zebrafish for modeling mitochondrial clinical

manifestations. Gaining insights from these studies, our compendium of mutants provides an

excellent resource to test a large number of biological and chemical modulators as potential

therapies for mitochondrial disorders for which treatment remains elusive.

Zebrafish offer many unique advantages for *in vivo* imaging experiments, as compared to their

mammalian counterparts. A classical study corroborating this advantage is *in vivo* imaging of

mitochondrial transport in a single axon, as demonstrated by Yang and his colleagues [121].

Seok and his colleagues generated a transgenic zebrafish line, expressing green fluorescent

protein fused to mitochondrial localization sequence from cytochrome c oxidase [71].

Mitochondrial function is often measured by various parameters such as estimation of membrane

potential, mitochondrial superoxide species and energy production. Superoxide activity and

membrane potential in zebrafish has been measured by employing the use of cell permeable

chemical probes such as MitoSox [122] and Dihydrorhodamine 123 (DH123) [73], respectively.

Zebrafish transgenic lines expressing genetically encoded calcium and oxidation indicators, have

also served as an excellent model to measure calcium homeostasis and oxidative status *in vivo* in

models of mechanosensory hair cell damage and death [123]. Constructs such as Mitotimer allow

the measurement of mitochondrial turnover, transport, and changes in the redox history of

mitochondria during organogenesis events in zebrafish embryos [124]. This transgenic model

enables a sweeping picture of the mitochondrial network, helping in studying cellular processes

such as mitophagy and apoptosis.

Zebrafish have also provided novel mechanistic insights to interrogate the mitochondrial

dynamics, to be called as "*in-vivo* life cycle of mitochondria" in healthy and diseased conditions

[124]. One such example is that of Mitofish [74], that recapitulates mitochondrial network

biogenesis, unraveling the role of the organelle in cell-type specific niches of different organ

systems. Mitofish is a transgenic zebrafish line that fluorescently labels the mitochondria in the

neurons, enabling non-invasive *in vivo* observation. Advancement in adaptive optics and lattice

light sheet microscopy have empowered researchers to look at vibrant and colorful images of

organelles in zebrafish. These technologies have been applied in investigating the organellar

dynamics in brain during early development and in the eye of an adult zebrafish [125]. These

MMMC mutants provide an excellent platform to study mitochondrial remodeling and

homeostasis in the progression of various clinical disorders by mapping the mitochondrial wiring

of the cell.

These zebrafish mutants have been cryopreserved for the purpose of sharing the lines as a

resource with the field for expanding the study of mitochondrial genetics and biology. In

addition, our template for the creation of mitochondrial mutants in zebrafish should enable the

creation of higher animal models of specific variants of interest. For the purpose of accelerating

this research, all lines in this project have had their information deposited with ZIRC, and all of

the lines have been made available via zfishbook (www.zfishbook.com) [83].

The goal of this study is to disseminate the use of zebrafish as a *noblesse oblige* in the field of

mitochondrial biology and medicine, paving way for the development of novel insight for

diagnostic and therapeutic strategies. Ultimately, using this collection of mutants we hope to

unravel a small part of the mystery that shrouds one of the most crucial organelles in the

cell. We hope that this MMMC mutant collection and the primer we provide for adding to it,

will help to usher in a new mitochondrial research using the zebrafish.

Acknowledgements

The authors thank Dr. Eric Schon and Dr. Vamsi Mootha for helping select the gene set targeted

in this collection, and the following who contributed to the generation of these new zebrafish

lines including Alexandra Cook, Roberto Lopez Cervera, and the Mayo Clinic Zebrafish Facility

staff. Gabriel Martinez Galvez helped with paper illustration design. This work was sponsored

by the Mayo Foundation, a gift by the Marriott Foundation and by NIH grants DK84567,

GM63904 and HG006431 to SCE.

Author Contributions

450 The paper was written by AS, MW, ZWJ and SCE. Experiments were executed by JMC, MW,

451 HA, NI, MDU, JDB, RMH, WL, YD and TLP with experimental guidance from XX, KJC and

452 SCE. Data analysis was completed by JMC, NI, JDB, MDU, RMH, MW, WL, YD and AS.

References

1.   Andersson SG, Zomorodipour A, Andersson JO, Sicheritz-Ponten T, Alsmark UC,

Podowski RM, et al. The genome sequence of Rickettsia prowazekii and the origin of

mitochondria. Nature. England; 1998;396: 133–140. doi:10.1038/24094

2.   Muller M, Martin W. The genome of Rickettsia prowazekii and some thoughts on the

origin of mitochondria and hydrogenosomes. Bioessays. United States; 1999;21: 377–381.

doi:10.1002/(SICI)1521-1878(199905)21:5<377::AID-BIES4>3.0.CO;2-W

3.   Gray MW, Burger G, Lang BF. Mitochondrial evolution. Science. United States;

1999;283: 1476–1481.

4.   Thorsness PE, Weber ER. Escape and migration of nucleic acids between chloroplasts,

mitochondria, and the nucleus. Int Rev Cytol. United States; 1996;165: 207–234.

5.   Lynch M, Koskella B, Schaack S. Mutation pressure and the evolution of organelle

genomic   architecture.   Science.   United   States;   2006;311:   1727–1730.

doi:10.1126/science.1118884

6.   Vafai SB, Mootha VK. Mitochondrial disorders as windows into an ancient organelle.

468   Nature. England; 2012;491: 374–383. doi:10.1038/nature11707

469   7.   Elson JL, Samuels DC, Turnbull DM, Chinnery PF. Random intracellular drift explains

470   the clonal expansion of mitochondrial DNA mutations with age. Am J Hum Genet. United

471   States; 2001;68: 802–806. doi:10.1086/318801

472   8.   DiMauro S, Hirano M. Mitochondrial DNA Deletion Syndromes. In: Adam MP, Ardinger

473   HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, et al., editors. Seattle (WA); 1993.

474   9.   Wallace DC. Mitochondrial DNA mutations in disease and aging. Environ Mol Mutagen.

475   United States; 2010;51: 440–450. doi:10.1002/em.20586

476   10.   Elson JL, Smith PM, Greaves LC, Lightowlers RN, Chrzanowska-Lightowlers ZMA,

477   Taylor RW, et al. The presence of highly disruptive 16S rRNA mutations in clinical

478   samples indicates a wider role for mutations of the mitochondrial ribosome in human

479   disease. Mitochondrion. Netherlands; 2015;25: 17–27. doi:10.1016/j.mito.2015.08.004

480   11.   Andrews RM, Kubacka I, Chinnery PF, Lightowlers RN, Turnbull DM, Howell N.

481   Reanalysis and revision of the Cambridge reference sequence for human mitochondrial

482   DNA. Nature genetics. United States; 1999. p. 147. doi:10.1038/13779

483   12.   Broughton RE, Milam JE, Roe BA. The complete sequence of the zebrafish (Danio rerio)

484   mitochondrial genome and evolutionary patterns in vertebrate mitochondrial DNA.

485   Genome Res. United States; 2001;11: 1958–1967. doi:10.1101/gr.156801

486   13.   Wolstenholme DR. Animal mitochondrial DNA: structure and evolution. Int Rev Cytol.

487   United States; 1992;141: 173–216.

488   14.   Osawa S, Jukes TH, Watanabe K, Muto A. Recent evidence for evolution of the genetic
489   code. Microbiol Rev. United States; 1992;56: 229–264.

490   15.   Richter R, Pajak A, Dennerlein S, Rozanska A, Lightowlers RN, Chrzanowska-
491   Lightowlers ZMA. Translation termination in human mitochondrial ribosomes. Biochem
492   Soc Trans. England; 2010;38: 1523–1526. doi:10.1042/BST0381523

493   16.   Bayona-Bafaluy MP, Blits B, Battersby BJ, Shoubridge EA, Moraes CT. Rapid directional
494   shift of mitochondrial DNA heteroplasmy in animal tissues by a mitochondrially targeted
495   restriction endonuclease. Proc Natl Acad Sci U S A. United States; 2005;102: 14392–
496   14397. doi:10.1073/pnas.0502896102

497   17.   Srivastava S, Moraes CT. Double-strand breaks of mouse muscle mtDNA promote large
498   deletions similar to multiple mtDNA deletions in humans. Hum Mol Genet. England;
499   2005;14: 893–902. doi:10.1093/hmg/ddi082

500   18.   Nissanka N, Minczuk M, Moraes CT. Mechanisms of Mitochondrial DNA Deletion
501   Formation. Trends Genet. England; 2019; doi:10.1016/j.tig.2019.01.001

502   19.   Alexeyev M, Shokolenko I, Wilson G, LeDoux S. The maintenance of mitochondrial
503   DNA integrity--critical analysis and update. Cold Spring Harb Perspect Biol. United
504   States; 2013;5: a012641. doi:10.1101/cshperspect.a012641

505   20.   Zinovkina LA. Mechanisms of Mitochondrial DNA Repair in Mammals. Biochemistry
506   (Mosc). United States; 2018;83: 233–249. doi:10.1134/S0006297918030045

21.   Patananan AN, Wu T-H, Chiou P-Y, Teitell MA. Modifying the Mitochondrial Genome.

Cell Metab. United States; 2016;23: 785–796. doi:10.1016/j.cmet.2016.04.004

22. Gammage PA, Moraes CT, Minczuk M. Mitochondrial Genome Engineering: The

Revolution May Not Be CRISPR-Ized. Trends Genet. England; 2018;34: 101–110.

doi:10.1016/j.tig.2017.11.001

23. Lopez MF, Kristal BS, Chernokalskaya E, Lazarev A, Shestopalov AI, Bogdanova A, et

al. High-throughput profiling of the mitochondrial proteome using affinity fractionation

and automation. Electrophoresis. Germany; 2000;21: 3427–3440. doi:10.1002/1522-

2683(20001001)21:16<3427::AID-ELPS3427>3.0.CO;2-L

24. Pagliarini DJ, Calvo SE, Chang B, Sheth SA, Vafai SB, Ong S-E, et al. A mitochondrial

protein compendium elucidates complex I disease biology. Cell. United States; 2008;134:

112–123. doi:10.1016/j.cell.2008.06.016

25. Elstner M, Andreoli C, Klopstock T, Meitinger T, Prokisch H. The mitochondrial

proteome database: MitoP2. Methods Enzymol. United States; 2009;457: 3–20.

doi:10.1016/S0076-6879(09)05001-0

26. Smith AC, Robinson AJ. MitoMiner, an integrated database for the storage and analysis of

mitochondrial proteomics data. Mol Cell Proteomics. United States; 2009;8: 1324–1337.

doi:10.1074/mcp.M800373-MCP200

27. Calvo SE, Clauser KR, Mootha VK. MitoCarta2.0: an updated inventory of mammalian

mitochondrial proteins. Nucleic Acids Res. England; 2016;44: D1251-7.

doi:10.1093/nar/gkv1003

28. Taylor RW, Turnbull DM. Mitochondrial DNA mutations in human disease. Nat Rev

Genet. England; 2005;6: 389–402. doi:10.1038/nrg1606

29.   Rossignol R, Faustin B, Rocher C, Malgat M, Mazat J-P, Letellier T. Mitochondrial

threshold effects. Biochem J. England; 2003;370: 751–762. doi:10.1042/BJ20021594

30.   Chinnery PF. Mitochondrial Disorders Overview. In: Adam MP, Ardinger HH, Pagon RA,

Wallace SE, Bean LJH, Stephens K, et al., editors. Seattle (WA); 1993.

31.   Gorman GS, Chinnery PF, DiMauro S, Hirano M, Koga Y, McFarland R, et al.

Mitochondrial diseases. Nature reviews. Disease primers. England; 2016. p. 16080.

doi:10.1038/nrdp.2016.80

32.   Skladal D, Halliday J, Thorburn DR. Minimum birth prevalence of mitochondrial

respiratory chain disorders in children. Brain. England; 2003;126: 1905–1912.

doi:10.1093/brain/awg170

33.   Schaefer AM, McFarland R, Blakely EL, He L, Whittaker RG, Taylor RW, et al.

Prevalence of mitochondrial DNA disease in adults. Ann Neurol. United States; 2008;63:

35–39. doi:10.1002/ana.21217

34.   Wallace DC, Singh G, Lott MT, Hodge JA, Schurr TG, Lezza AM, et al. Mitochondrial

DNA mutation associated with Leber's hereditary optic neuropathy. Science. United

States; 1988;242: 1427–1430.

35.   Swerdlow RH, Wooten GF. A novel deafness/dystonia peptide gene mutation that causes

dystonia in female carriers of Mohr-Tranebjaerg syndrome. Ann Neurol. United States;

2001;50: 537–540.

36.   Garone C, Tadesse S, Hirano M. Clinical and genetic spectrum of mitochondrial

neurogastrointestinal encephalomyopathy. Brain. England; 2011;134: 3326–3332.

doi:10.1093/brain/awr245

37. Urrutia PJ, Mena NP, Nunez MT. The interplay between iron accumulation, mitochondrial

dysfunction, and inflammation during the execution step of neurodegenerative disorders.

Front Pharmacol. Switzerland; 2014;5: 38. doi:10.3389/fphar.2014.00038

38. Singh RK, Saini S, Verma D, Kalaiarasan P, Bamezai RNK. Mitochondrial ND5 mutation

mediated elevated ROS regulates apoptotic pathway epigenetically in a P53 dependent

manner for generating pro-cancerous phenotypes. Mitochondrion. Netherlands; 2017;35:

35–43. doi:10.1016/j.mito.2017.05.001

39. Craven L, Alston CL, Taylor RW, Turnbull DM. Recent Advances in Mitochondrial

Disease. Annu Rev Genomics Hum Genet. United States; 2017;18: 257–275.

doi:10.1146/annurev-genom-091416-035426

40. Ghezzi D, Saada A, D'Adamo P, Fernandez-Vizarra E, Gasparini P, Tiranti V, et al.

FASTKD2 nonsense mutation in an infantile mitochondrial encephalomyopathy

associated with cytochrome c oxidase deficiency. Am J Hum Genet. United States;

2008;83: 415–423. doi:10.1016/j.ajhg.2008.08.009

41. Shehata BM, Cundiff CA, Lee K, Sabharwal A, Lalwani MK, Davis AK, et al. Exome

sequencing of patients with histiocytoid cardiomyopathy reveals a de novo NDUFB11

mutation that plays a role in the pathogenesis of histiocytoid cardiomyopathy. Am J Med

Genet A. United States; 2015;167A: 2114–2121. doi:10.1002/ajmg.a.37138

42. Wesolowska M, Gorman GS, Alston CL, Pajak A, Pyle A, He L, et al. Adult Onset Leigh

Syndrome in the Intensive Care Setting: A Novel Presentation of a C12orf65 Related

Mitochondrial Disease. J Neuromuscul Dis. Netherlands; 2015;2: 409–419. doi:10.3233/JND-150121

43. Floyd BJ, Wilkerson EM, Veling MT, Minogue CE, Xia C, Beebe ET, et al. Mitochondrial Protein Interaction Mapping Identifies Regulators of Respiratory Chain Function. Mol Cell. United States; 2016;63: 621–632. doi:10.1016/j.molcel.2016.06.033

44. Kohda M, Tokuzawa Y, Kishita Y, Nyuzuki H, Moriyama Y, Mizuno Y, et al. A Comprehensive Genomic Analysis Reveals the Genetic Landscape of Mitochondrial Respiratory Chain Complex Deficiencies. PLoS Genet. United States; 2016;12: e1005679. doi:10.1371/journal.pgen.1005679

45. Boumans H, Grivell LA, Berden JA. The respiratory chain in yeast behaves as a single functional unit. J Biol Chem. United States; 1998;273: 4872–4877.

46. Schwimmer C, Rak M, Lefebvre-Legendre L, Duvezin-Caubet S, Plane G, di Rago J-P. Yeast models of human mitochondrial diseases: from molecular mechanisms to drug screening. Biotechnol J. Germany; 2006;1: 270–281. doi:10.1002/biot.200500053

47. Altmann K, Durr M, Westermann B. Saccharomyces cerevisiae as a model organism to study mitochondrial biology: general considerations and basic procedures. Methods Mol Biol. United States; 2007;372: 81–90. doi:10.1007/978-1-59745-365-3_6

48. Denoth Lippuner A, Julou T, Barral Y. Budding yeast as a model organism to study the effects of age. FEMS Microbiol Rev. England; 2014;38: 300–325. doi:10.1111/1574-6976.12060

49. Addo MG, Cossard R, Pichard D, Obiri-Danso K, Rotig A, Delahodde A. Caenorhabditis

elegans, a pluricellular model organism to screen new genes involved in mitochondrial

genome maintenance. Biochim Biophys Acta. Netherlands; 2010;1802: 765–773.

doi:10.1016/j.bbadis.2010.05.007

50.   Harrington AJ, Hamamichi S, Caldwell GA, Caldwell KA. C. elegans as a model

organism to investigate molecular pathways involved with Parkinson's disease. Dev Dyn.

United States; 2010;239: 1282–1295. doi:10.1002/dvdy.22231

51.   Moreno-Arriola E, Cardenas-Rodriguez N, Coballase-Urrutia E, Pedraza-Chaverri J,

Carmona-Aparicio L, Ortega-Cuellar D. Caenorhabditis elegans: A useful model for

studying metabolic disorders in which oxidative stress is a contributing factor. Oxid Med

Cell Longev. United States; 2014;2014: 705253. doi:10.1155/2014/705253

52.   Tissenbaum HA. Using C. elegans for aging research. Invertebr Reprod Dev. England;

2015;59: 59–63. doi:10.1080/07924259.2014.940470

53.   Sanchez-Martinez A, Luo N, Clemente P, Adan C, Hernandez-Sierra R, Ochoa P, et al.

Modeling human mitochondrial diseases in flies. Biochim Biophys Acta. Netherlands;

2006;1757: 1190–1198. doi:10.1016/j.bbabio.2006.05.008

54.   Fernandez-Moreno MA, Farr CL, Kaguni LS, Garesse R. Drosophila melanogaster as a

model system to study mitochondrial biology. Methods Mol Biol. United States;

2007;372: 33–49. doi:10.1007/978-1-59745-365-3_3

55.   Guo M. Drosophila as a model to study mitochondrial dysfunction in Parkinson's disease.

Cold Spring Harb Perspect Med. United States; 2012;2. doi:10.1101/cshperspect.a009944

56.   Sen A, Cox RT. Fly Models of Human Diseases: Drosophila as a Model for

Understanding Human Mitochondrial Mutations and Disease. Curr Top Dev Biol. United

States; 2017;121: 1–27. doi:10.1016/bs.ctdb.2016.07.001

57.   Lightowlers RN, Rozanska A, Chrzanowska-Lightowlers ZM. Mitochondrial protein

synthesis: figuring the fundamentals, complexities and complications, of mammalian

mitochondrial   translation.   FEBS   Lett.   England;   2014;588:   2496–2503.

doi:10.1016/j.febslet.2014.05.054

58.   Pett W, Lavrov D V. Cytonuclear Interactions in the Evolution of Animal Mitochondrial

tRNA   Metabolism.   Genome   Biol   Evol.   England;   2015;7:   2089–2101.

doi:10.1093/gbe/evv124

59.   Lavrov D V, Pett W. Animal Mitochondrial DNA as We Do Not Know It: mt-Genome

Organization and Evolution in Nonbilaterian Lineages. Genome Biol Evol. England;

2016;8: 2896–2913. doi:10.1093/gbe/evw195

60.   Brown WM, George MJ, Wilson AC. Rapid evolution of animal mitochondrial DNA.

Proc Natl Acad Sci U S A. United States; 1979;76: 1967–1971.

61.   Moyes CD, Suarez RK, Hochachka PW, Ballantyne JS. A comparison of fuel preferences

of mitochondria from vertebrates and invertebrates. Can J Zool. NRC Research Press;

1990;68: 1337–1349. doi:10.1139/z90-201

62.   Grossman LI, Wildman DE, Schmidt TR, Goodman M. Accelerated evolution of the

electron transport chain in anthropoid primates. Trends Genet. England; 2004;20: 578–

585. doi:10.1016/j.tig.2004.09.002

63.   Gershoni M, Fuchs A, Shani N, Fridman Y, Corral-Debrinski M, Aharoni A, et al.

Coevolution predicts direct interactions between mtDNA-encoded and nDNA-encoded

subunits of oxidative phosphorylation complex i. J Mol Biol. England; 2010;404: 158–

171. doi:10.1016/j.jmb.2010.09.029

64.   Haen KM, Pett W, Lavrov D V. Parallel loss of nuclear-encoded mitochondrial

aminoacyl-tRNA synthetases and mtDNA-encoded tRNAs in Cnidaria. Mol Biol Evol.

United States; 2010;27: 2216–2219. doi:10.1093/molbev/msq112

65.   Rea SL, Graham BH, Nakamaru-Ogiso E, Kar A, Falk MJ. Bacteria, yeast, worms, and

flies: exploiting simple model organisms to investigate human mitochondrial diseases.

Dev Disabil Res Rev. United States; 2010;16: 200–218. doi:10.1002/ddrr.114

66.   Kuzmenko A V, Levitskii SA, Vinogradova EN, Atkinson GC, Hauryliuk V, Zenkin N, et

al. Protein biosynthesis in mitochondria. Biochemistry (Mosc). United States; 2013;78:

855–866. doi:10.1134/S0006297913080014

67.   Zhao J, Lendahl U, Nister M. Regulation of mitochondrial dynamics: convergences and

divergences between yeast and vertebrates. Cell Mol Life Sci. Switzerland; 2013;70:

951–976. doi:10.1007/s00018-012-1066-6

68.   Lieschke GJ, Currie PD. Animal models of human disease: zebrafish swim into view. Nat

Rev Genet. England; 2007;8: 353–367. doi:10.1038/nrg2091

69.   Lin C-Y, Chiang C-Y, Tsai H-J. Zebrafish and Medaka: new model organisms for modern

biomedical research. J Biomed Sci. England; 2016;23: 19. doi:10.1186/s12929-016-0236-

5

70.   Blanco-Sanchez B, Clement A, Phillips JB, Westerfield M. Zebrafish models of human

eye and inner ear diseases. Methods Cell Biol. United States; 2017;138: 415–467.

doi:10.1016/bs.mcb.2016.10.006

71.   Kim MJ, Kang KH, Kim C-H, Choi S-Y. Real-time imaging of mitochondria in transgenic

659   zebrafish expressing mitochondrially targeted GFP. Biotechniques. England; 2008;45:

660   331–334. doi:10.2144/000112909

661   72.   Flinn L, Mortiboys H, Volkmann K, Koster RW, Ingham PW, Bandmann O. Complex I

662   deficiency and dopaminergic neuronal cell loss in parkin-deficient zebrafish (Danio rerio).

663   Brain. England; 2009;132: 1613–1623. doi:10.1093/brain/awp108

664   73.   Song Y, Selak MA, Watson CT, Coutts C, Scherer PC, Panzer JA, et al. Mechanisms

665   underlying metabolic and neural defects in zebrafish and human multiple acyl-CoA

666   dehydrogenase deficiency (MADD). PLoS One. United States; 2009;4: e8329.

667   doi:10.1371/journal.pone.0008329

668   74.   Plucinska G, Paquet D, Hruscha A, Godinho L, Haass C, Schmid B, et al. In vivo imaging

669   of disease-related mitochondrial dynamics in a vertebrate model system. J Neurosci.

670   United States; 2012;32: 16203–16212. doi:10.1523/JNEUROSCI.1327-12.2012

671   75.   Steele SL, Prykhozhij S V, Berman JN. Zebrafish as a model system for mitochondrial

672   biology   and   diseases.   Transl   Res.   United   States;   2014;163:   79–98.

673   doi:10.1016/j.trsl.2013.08.008

674   76.   Byrnes J, Ganetzky R, Lightfoot R, Tzeng M, Nakamaru-Ogiso E, Seiler C, et al.

675   Pharmacologic modeling of primary mitochondrial respiratory chain dysfunction in

676   zebrafish. Neurochem Int. England; 2017; doi:10.1016/j.neuint.2017.07.008

77.   Pinho BR, Santos MM, Fonseca-Silva A, Valentao P, Andrade PB, Oliveira JMA. How

mitochondrial dysfunction affects zebrafish development and cardiovascular function: an

in vivo model for testing mitochondria-targeted drugs. Br J Pharmacol. England;

2013;169: 1072–1090. doi:10.1111/bph.12186

78.   Polyak E, Ostrovsky J, Peng M, Dingley SD, Tsukikawa M, Kwon YJ, et al. N-

acetylcysteine and vitamin E rescue animal longevity and cellular oxidative stress in pre-

clinical models of mitochondrial complex I disease. Mol Genet Metab. United States;

2018;123: 449–462. doi:10.1016/j.ymgme.2018.02.013

79.   Ma AC, McNulty MS, Poshusta TL, Campbell JM, Martinez-Galvez G, Argue DP, et al.

FusX: A Rapid One-Step Transcription Activator-Like Effector Assembly System for

Genome   Science.   Hum   Gene   Ther.   United   States;   2016;27:   451–463.

doi:10.1089/hum.2015.172

80.   Clark KJ, Balciunas D, Pogoda H-M, Ding Y, Westcot SE, Bedell VM, et al. In vivo

protein trapping produces a functional expression codex of the vertebrate proteome. Nat

Methods. United States; 2011;8: 506–515. doi:10.1038/nmeth.1606

81.   Neff KL, Argue DP, Ma AC, Lee HB, Clark KJ, Ekker SC. Mojo Hand, a TALEN design

tool for genome editing applications. BMC Bioinformatics. England; 2013;14: 1.

doi:10.1186/1471-2105-14-1

82.   Clark KJ, Geurts AM, Bell JB, Hackett PB. Transposon vectors for gene-trap insertional

mutagenesis   in   vertebrates.   Genesis.   United   States;   2004;39:   225–233.

doi:10.1002/gene.20049

83.   Clark KJ, Argue DP, Petzold AM, Ekker SC. zfishbook: connecting you to a world of

zebrafish revertible mutants. Nucleic Acids Res. England; 2012;40: D907-11.

doi:10.1093/nar/gkr957

84. Sharma LK, Lu J, Bai Y. Mitochondrial respiratory complex I: structure, function and

implication in human diseases. Curr Med Chem. United Arab Emirates; 2009;16: 1266–

1277.

85. Liu Y, Fiskum G, Schubert D. Generation of reactive oxygen species by the mitochondrial

electron transport chain. J Neurochem. England; 2002;80: 780–787.

86. King MP, Attardi G. Human cells lacking mtDNA: repopulation with exogenous

mitochondria by complementation. Science. United States; 1989;246: 500–503.

87. Koga Y, Davidson M, Schon EA, King MP. Analysis of cybrids harboring MELAS

mutations in the mitochondrial tRNA(Leu(UUR)) gene. Muscle Nerve Suppl. United

States; 1995;3: S119-23.

88. Vithayathil SA, Ma Y, Kaipparettu BA. Transmitochondrial cybrids: tools for functional

studies of mutant mitochondria. Methods Mol Biol. United States; 2012;837: 219–230.

doi:10.1007/978-1-61779-504-6_15

89. Wilkins HM, Carl SM, Swerdlow RH. Cytoplasmic hybrid (cybrid) cell lines as a practical

model for mitochondriopathies. Redox Biol. Netherlands; 2014;2: 619–631.

doi:10.1016/j.redox.2014.03.006

90. Bedell VM, Wang Y, Campbell JM, Poshusta TL, Starker CG, Krug RG 2nd, et al. In vivo

genome editing using a high-efficiency TALEN system. Nature. England; 2012;491: 114–

118. doi:10.1038/nature11537

91.   Hwang WY, Fu Y, Reyon D, Maeder ML, Tsai SQ, Sander JD, et al. Efficient genome

editing in zebrafish using a CRISPR-Cas system. Nat Biotechnol. United States; 2013;31:

227–229. doi:10.1038/nbt.2501

92.   Jiang W, Zhou H, Bi H, Fromm M, Yang B, Weeks DP. Demonstration of

CRISPR/Cas9/sgRNA-mediated targeted gene modification in Arabidopsis, tobacco,

sorghum and rice. Nucleic Acids Res. England; 2013;41: e188. doi:10.1093/nar/gkt780

93.   Fujihara Y, Ikawa M. CRISPR/Cas9-based genome editing in mice by single plasmid

injection. Methods Enzymol. United States; 2014;546: 319–336. doi:10.1016/B978-0-12-

801185-0.00015-5

94.   Hsu PD, Lander ES, Zhang F. Development and applications of CRISPR-Cas9 for

genome   engineering.   Cell.   United   States;   2014;157:   1262–1278.

doi:10.1016/j.cell.2014.05.010

95.   Zhang F, Wen Y, Guo X. CRISPR/Cas9 for genome editing: progress, implications and

challenges. Hum Mol Genet. England; 2014;23: R40-6. doi:10.1093/hmg/ddu125

96.   Hashimoto M, Bacman SR, Peralta S, Falk MJ, Chomyn A, Chan DC, et al. MitoTALEN:

A General Approach to Reduce Mutant mtDNA Loads and Restore Oxidative

Phosphorylation Function in Mitochondrial Diseases. Mol Ther. United States; 2015;23:

1592–1599. doi:10.1038/mt.2015.126

97.   Niu D, Wei H-J, Lin L, George H, Wang T, Lee I-H, et al. Inactivation of porcine

endogenous retrovirus in pigs using CRISPR-Cas9. Science. United States; 2017;357:

1303–1307. doi:10.1126/science.aan4187

98.   Meimaridou E, Kowalczyk J, Guasti L, Hughes CR, Wagner F, Frommolt P, et al.

Mutations in NNT encoding nicotinamide nucleotide transhydrogenase cause familial

glucocorticoid deficiency. Nat Genet. United States; 2012;44: 740–742.

doi:10.1038/ng.2299

99.   Bainbridge MN, Davis EE, Choi W-Y, Dickson A, Martinez HR, Wang M, et al. Loss of

Function Mutations in NNT Are Associated With Left Ventricular Noncompaction. Circ

Cardiovasc Genet. United States; 2015;8: 544–552.

doi:10.1161/CIRCGENETICS.115.001026

100.   Anichtchik O, Diekmann H, Fleming A, Roach A, Goldsmith P, Rubinsztein DC. Loss of

PINK1 function affects development and results in neurodegeneration in zebrafish. J

Neurosci. United States; 2008;28: 8199–8207. doi:10.1523/JNEUROSCI.0979-08.2008

101.   Fett ME, Pilsl A, Paquet D, van Bebber F, Haass C, Tatzelt J, et al. Parkin is protective

against proteotoxic stress in a transgenic zebrafish model. PLoS One. United States;

2010;5: e11783. doi:10.1371/journal.pone.0011783

102.   Sallinen V, Kolehmainen J, Priyadarshini M, Toleikyte G, Chen Y-C, Panula P.

Dopaminergic cell damage and vulnerability to MPTP in Pink1 knockdown zebrafish.

Neurobiol Dis. United States; 2010;40: 93–101. doi:10.1016/j.nbd.2010.06.001

103.   Xi Y, Ryan J, Noble S, Yu M, Yilbas AE, Ekker M. Impaired dopaminergic neuron

development and locomotor function in zebrafish with loss of pink1 function. Eur J

Neurosci. France; 2010;31: 623–633. doi:10.1111/j.1460-9568.2010.07091.x

104.   Kumar MG, Rowley S, Fulton R, Dinday MT, Baraban SC, Patel M. Altered Glycolysis

and Mitochondrial Respiration in a Zebrafish Model of Dravet Syndrome. eNeuro. United

States; 2016;3. doi:10.1523/ENEURO.0008-16.2016

105. Baraban SC, Dinday MT, Hortopan GA. Drug screening in Scn1a zebrafish mutant
identifies clemizole as a potential Dravet syndrome treatment. Nat Commun. England;
2013;4: 2410. doi:10.1038/ncomms3410

106. Grone BP, Qu T, Baraban SC. Behavioral Comorbidities and Drug Treatments in a
Zebrafish scn1lab Model of Dravet Syndrome. eNeuro. United States; 2017;4.
doi:10.1523/ENEURO.0066-17.2017

107. Guernsey DL, Jiang H, Campagna DR, Evans SC, Ferguson M, Kellogg MD, et al.
Mutations in mitochondrial carrier family gene SLC25A38 cause nonsyndromic
autosomal recessive congenital sideroblastic anemia. Nat Genet. United States; 2009;41:
651–653. doi:10.1038/ng.359

108. Bronnimann D, Annese T, Gorr TA, Djonov V. Splitting of circulating red blood cells as
an in vivo mechanism of erythrocyte maturation in developing zebrafish, chick and mouse
embryos. J Exp Biol. England; 2018;221. doi:10.1242/jeb.184564

109. Berman J, Hsu K, Look AT. Zebrafish as a model organism for blood diseases. Br J
Haematol. England; 2003;123: 568–576.

110. Danilova N, Sakamoto KM, Lin S. Ribosomal protein S19 deficiency in zebrafish leads to
developmental abnormalities and defective erythropoiesis through activation of p53
protein family. Blood. United States; 2008;112: 5228–5237. doi:10.1182/blood-2008-01-
132290

111. Butler MG, Gore A V, Weinstein BM. Zebrafish as a model for hemorrhagic stroke.

Methods Cell Biol. United States; 2011;105: 137–161. doi:10.1016/B978-0-12-381320-
6.00006-0

112. Kulkeaw K, Sugiyama D. Zebrafish erythropoiesis and the utility of fish as models of
anemia. Stem Cell Res Ther. England; 2012;3: 55. doi:10.1186/scrt146

113. Kretz CA, Weyand AC, Shavit JA. Modeling Disorders of Blood Coagulation in the
Zebrafish. Curr Pathobiol Rep. United States; 2015;3: 155–161. doi:10.1007/s40139-015-
0081-3

114. Avagyan S, Zon LI. Fish to Learn: Insights into Blood Development and Blood Disorders
from Zebrafish Hematopoiesis. Hum Gene Ther. United States; 2016;27: 287–294.
doi:10.1089/hum.2016.024

115. Prezant TR, Agapian J V, Bohlman MC, Bu X, Oztas S, Qiu WQ, et al. Mitochondrial
ribosomal RNA mutation associated with both antibiotic-induced and non-syndromic
deafness. Nat Genet. United States; 1993;4: 289–294. doi:10.1038/ng0793-289

116. Zhang Q, Zhang L, Chen D, He X, Yao S, Zhang Z, et al. Deletion of Mtu1 (Trmu) in
zebrafish revealed the essential role of tRNA modification in mitochondrial biogenesis
and   hearing   function.   Nucleic   Acids   Res.   England;   2018;46:   10930–10945.
doi:10.1093/nar/gky758

117. DiMauro S, Schon EA. Mitochondrial respiratory-chain diseases. N Engl J Med. United
States; 2003;348: 2656–2668. doi:10.1056/NEJMra022567

118. Jain IH, Zazzeron L, Goli R, Alexa K, Schatzman-Bone S, Dhillon H, et al. Hypoxia as a
therapy   for   mitochondrial   disease.   Science.   United   States;   2016;352:   54–61.

doi:10.1126/science.aad9642

119.   Bretaud S, Lee S, Guo S. Sensitivity of zebrafish to environmental toxins implicated in

Parkinson's disease. Neurotoxicol Teratol. United States; 2004;26: 857–864.

doi:10.1016/j.ntt.2004.06.014

120.   Byrnes J, Ganetzky R, Lightfoot R, Tzeng M, Nakamaru-Ogiso E, Seiler C, et al.

Pharmacologic modeling of primary mitochondrial respiratory chain dysfunction in

zebrafish. Neurochem Int. England; 2018;117: 23–34. doi:10.1016/j.neuint.2017.07.008

121.   Xu Y, Chen M, Hu B, Huang R, Hu B. In vivo Imaging of Mitochondrial Transport in

Single-Axon Regeneration of Zebrafish Mauthner Cells. Front Cell Neurosci. Switzerland;

2017;11: 4. doi:10.3389/fncel.2017.00004

122.   Dowling JJ, Arbogast S, Hur J, Nelson DD, McEvoy A, Waugh T, et al. Oxidative stress

and successful antioxidant treatment in models of RYR1-related myopathy. Brain.

England; 2012;135: 1115–1127. doi:10.1093/brain/aws036

123.   Pickett SB, Thomas ED, Sebe JY, Linbo T, Esterberg R, Hailey DW, et al. Cumulative

mitochondrial activity correlates with ototoxin susceptibility in zebrafish mechanosensory

hair cells. Elife. England; 2018;7. doi:10.7554/eLife.38062

124.   Mandal A, Pinter K, Drerup CM. Analyzing Neuronal Mitochondria in vivo Using

Fluorescent Reporters in Zebrafish. Front cell Dev Biol. Switzerland; 2018;6: 144.

doi:10.3389/fcell.2018.00144

125.   Liu T-L, Upadhyayula S, Milkie DE, Singh V, Wang K, Swinburne IA, et al. Observing

the cell in its native state: Imaging subcellular dynamics in multicellular organisms.

Science. United States; 2018;360. doi:10.1126/science.aaq1392

 Figures Legends:

Figure 1: Circular representation of the zebrafish (NC_002333.2) and human

mitochondrial genomes (NC_012920.1): Both genomes share the same synteny, number of

genes and are nearly identical in size.

Figure 2: Zebrafish Marriot mitochondrial mutant collection: Schematic representation of

various mitochondrial resident pathways for which zebrafish mutants were generated. The

nuclear-encoded mitochondrial proteins have been illustrated according to the function they are

involved in mitochondrial maintenance and homeostasis. Mutants generated by gene editing are

depicted as grey, whereas those generated by the GBT system are depicted in pink.

Figure 3: A depiction of homology of the mutants created for proteins involved in different

mitochondrial resident pathways: A-C: Complex 1 of mitochondrial respiratory chain; A:

NADH: ubiquinone oxidoreductase subunit AB1 b (Ndufab1b); B: NADH dehydrogenase

association factor 6 (Ndufaf6); C: NADH dehydrogenase iron sulfur protein 4 (Ndufs4); **D:**

Complex 2 of mitochondrial respiratory chain, Succinate dehydrogenase subunit A (Sdha),

subunit; **E-F: Complex 3 of mitochondrial respiratory chain;** E: Mitochondrial chaperone

protein BCS1 (Bcs1l); F: Ubiquinol-cytochrome c reductase subunit VIII (Uqcrq); **G-I:**

Complex 4 of mitochondrial respiratory chain; G: Protoheme IX farnesyltransferase (Cox10);

H: Cytochrome c oxidase IV subunit 1 (Cox4i2b); I: Surfeit locus protein 1 (Surf1); **J-K:**

Complex 5 of mitochondrial respiratory chain; J: ATP synthase subunit epsilon (Atp5e); K:

Transmembrane protein 70 (Tmem70); **L-M: Biosynthesis of coenzyme Q;** L: 4-

hydroxybenzoate polyprenyltransferase (Coq2) M: Decaprenyl-disphosphate synthase subunit 2

(Pdss2); **N-Q: Mitochondrial calcium homeostasis;** N: Mitochondrial calcium uptake protein 1

(Micu1); O: Calcium uniporter protein (Mcu) generated by TALEN and P; GBT; Q: Essential

MCU regulator a (Smdt1a); **R-U: Mitochondrial genome maintenance;** R: Thymidine kinase 2

(Tk2); S: Leucine-rich pentatricopeptide repeat-containing (Lrpprc); T: Mitochondrial

methionyl-tRNA formyltransferase (Mtfmt); U: 28s Ribosomal protein s18b, mitochondrial

(Mrps18b); **V: Mitochondrial protein import;** Mitochondrial import inner membrane

translocase subunit TIM50 (Timm50); **W-X: Mitochondrial metabolite synthesis;** W: 2-

Oxoglutarate dehydrogenase b (Ogdhb); X: Isocitrate dehydrogenase (Idh2).

Figures:

Figure 1:

Figure 2:

Figure 3:

Amino Acid Homology 0-40% ☐ 40-60% ☐ 60-80% ☐ 80-100% ■

M Decaprenyl-disphosphate synthase subunit 2 (Pdss2)
Human: 1 ── 377
Zebrafish Wild Type: 1 ─28─ 399
pdss2^mn0120 1 ── ↓ TALEN Cut Site 185

N Mitochondrial calcium uptake protein 1 (Micu1)
Human: 1 ── 476
Zebrafish Wild Type: 1 ─103─ 488
micu1^mn0132 1 ── ↓ TALEN cut site / 3 amino acid substitution 489

O Essential MCU regulator (Smdt1a)
Human: 1 ── 107
Zebrafish Wild Type: 1 ─69─ 124
smdt1^mn0122 1 ── ↓ TALEN cut site 20

P Mitochondrial calcium uniporter (Mcu)
Human: 1 ── 351
Zebrafish Wild Type: 1 ─76─ 376
mcu^mn0111 1 ── ↓ TALEN cut site *Exon 2 deleted 39 / 40 ── 338

Q Mitochondrial calcium uniporter (Mcu)
Human: 1 ── 351
Zebrafish Wild Type: 1 ─76─ 376
mcu^mn0599GT: 1 ── ↓ GBT insertion site 97

R Thymidine kinase 2 (Tk2)
Human: 1 ── * Active site 265
Zebrafish Wild Type: 1 ─58─ 268
tk2^mn0125 1 ── ↓ TALEN cut site 222

S Leucine-rich pentatricopeptide repeat-containing (Lrpprc)
Human: 1 ── 1394
Zebrafish Wild Type: 1 ── 1461
lrpprc^mn0235GT 1 ── ↓ GBT insertion site 817

T Mitochondrial methionyl-tRNA formyltransferase (Mtfmt)
Human: 1 ── 389
Zebrafish Wild Type: 1 ─50─ 390
mtfmt^mn0114 1 ── ↓ TALEN cut site 233
mtfmt^mn0113 1 ── ↓ TALEN cut site 234

U 28s ribosomal protein s18b (Mrps18b)
Human: 1 ── 258
Zebrafish Wild Type: 1 ── 242
mrps18b^mn0425GT 1 ── ↓ GBT insertion site 75

V Mitochondrial import inner membrane translocase subunit TIM50 (Timm50)
Human: 1 ── 353
Zebrafish Wild Type: 1 ─96─ 387
timm50^mn0906GT 1 ── ↓ GBT Insertion Site 139

W 2-oxoglutarate dehydrogenase (Ogdhb)
Human: 1 ── 1023
Zebrafish Wild Type: 1 ── 1023
ogdh^mn0281GT 1 ── ↓ GBT Insertion Site 174

X Isocitrate Dehydrogenase (Idh2)
Human: 1 ── 451
Zebrafish Wild Type: 1 ── 449
idh2^mn0268GT 1 ── ↓ GBT Insertion Site 320

49