Evolutionary rates are correlated between cockroach symbiont

and mitochondrial genomes

Daej A. Arab[1], Thomas Bourguignon[1,2,3], Zongqing Wang[4], Simon Y. W. Ho[1], & Nathan Lo[1]

[1]School of Life and Environmental Sciences, University of Sydney, Sydney, Australia

[2]Okinawa Institute of Science and Technology Graduate University, Tancha, Onna-son,

Okinawa, Japan

[3]Faculty of Forestry and Wood Sciences, Czech University of Life Sciences, Prague, Czech

Republic

[4]College of Plant Protection, Southwest University, Chongqing, China

Authors for correspondence:

Daej A. Arab

e-mail: daej.arab@sydney.edu.au

Nathan Lo

e-mail: nathan.lo@sydney.edu.au

Keywords: host-symbiont interaction, *Blattabacterium cuenoti*, phylogeny, molecular

evolution, substitution rate, cockroach.

Abstract

Bacterial endosymbionts evolve under strong host-driven selection. Factors influencing host
evolution might affect symbionts in similar ways, potentially leading to correlations between
the molecular evolutionary rates of hosts and symbionts. Although there is evidence of rate
correlations between mitochondrial and nuclear genes, similar investigations of hosts and
symbionts are lacking. Here we demonstrate a correlation in molecular rates between the
genomes of an endosymbiont (*Blattabacterium cuenoti*) and the mitochondrial genomes of
their hosts (cockroaches). We used partial genome data for multiple strains of *B. cuenoti* to
compare phylogenetic relationships and evolutionary rates for 55 cockroach/symbiont
pairs. The phylogenies inferred for *B. cuenoti* and the mitochondrial genomes of their hosts
were largely congruent, as expected from their identical maternal and cytoplasmic mode of
inheritance. We found a correlation between evolutionary rates of the two genomes, based on
comparisons of root-to-tip distances and on comparisons of the branch lengths of
phylogenetically independent species pairs. Our results underscore the profound effects that
long-term symbiosis can have on the biology of each symbiotic partner.

1. Introduction

Rates of molecular evolution are governed by a multitude of factors and vary significantly among species [1, 2]. In the case of symbiotic organisms, such rates may be influenced by the biology of their symbiotic partner, in addition to their own. This is particularly the case for strictly vertically transmitted, obligate intracellular symbionts (hereafter 'symbionts'), which have a highly intimate relationship with their hosts [3]. For example, a small host effective population size will potentially lead to increased fixation of slightly deleterious mutations within both host and symbiont genomes, owing to the reduced efficacy of selection.

When the phylogenies of host and symbiont taxa are compared, simultaneous changes in evolutionary rate between host-symbiont pairs might be evident in their branch lengths. Some studies have found a correlation in evolutionary rates between nuclear and mitochondrial genes in sharks [4], herons [5], and turtles [6], suggesting that host biology affects substitution rates in nuclear and cytoplasmic genomes in similar ways. In insects, nuclear genes that interact directly with mitochondrial proteins have shown rate correlations with mitochondrial genes [7].

Potential correlations in evolutionary rates between hosts and bacterial symbionts remain untested. Evidence for correlated levels of synonymous substitutions was found in a study of one nuclear gene and two mitochondrial genes from *Camponotus* ants and three genes from their *Blochmannia* symbionts [8]. However, the study did not determine whether this correlation was driven by rates of evolution, time since divergence, or both. Numbers of substitutions tend to be low for closely related pairs of hosts and their corresponding symbionts, and high for more divergent pairs, leading to a correlation with time that does not necessarily reflect correlation in evolutionary rates.

Blattabacterium cuenoti (hereafter *Blattabacterium*) is an intracellular bacterial symbiont that has been in an obligatory intracellular and mutualistic relationship with cockroaches for over 200 million years [9, 10]. Found in highly specialized cells in the fat bodies of cockroaches, *Blattabacterium* is required for host fitness and fertility, and is transovarially transmitted from the mother to the progeny [11, 12]. Genome-wide analyses of the symbiont have confirmed its role in host nitrogen metabolism and the synthesis of essential amino acids [13, 14]. The genomes of 21 *Blattabacterium* strains sequenced to date are highly reduced compared with those of their free-living relatives, ranging in size from 590 to 645 kb [15, 16]. They contain genes encoding enzymes for DNA replication and repair, with some exceptions (*holA*, *holB*, and *mutH*) [13, 16, 17]. The extent to which host

nuclear proteins are involved in the cell biology of *Blattabacterium*, and particularly DNA

replication, is not well understood.

We recently performed a study of cockroach evolution and biogeography using

mitochondrial genomes [9]. During this process, we obtained partial genomic information for

several *Blattabacterium* strains. These data provide the opportunity to test for correlation of

molecular evolutionary rates between *Blattabacterium* and host-cockroach mitochondrial

DNA. Here we infer phylogenetic trees for 55 *Blattabacterium* strains on the basis of 104

genes and compare branch lengths and rates of evolution for host-symbiont pairs across the

phylogeny. We find evidence of markedly increased rates of evolution in some

Blattabacterium lineages, which appear to be matched by increased rates of evolution in

mitochondrial DNA of host lineages.

2. Materials and methods

A list of samples and collection data for each cockroach examined is provided in table S1

(see electronic supplementary material, ESM). For the majority of taxa examined in this

study, we obtained *Blattabacterium* sequence data from genomic libraries originally used in a

previous study of cockroach mitochondrial genomes carried out by our laboratories [9]. In

some cases, new genomic data were obtained from fat bodies of individual cockroaches (see

ESM for further details). We obtained 104 genes of 55 *Blattabacterium* strains from these

data and aligned them with orthologues from seven outgroup taxa from Flavobacteriales

(details provided in ESM).

Genomic data were assembled and annotated, and then aligned and tested for

saturation. After the exclusion of 3rd codon sites in each data set, total lengths for the

mitochondrial and *Blattabacterium* alignments were 11,051 bp and 71,458, respectively. The

former was partitioned into four subsets (1st codon sites, 2nd codon sites, rRNAs, and

tRNAs), and the latter into two subsets according to codon positions. Trees were inferred

using maximum likelihood in RAxML v8.2 [18], using 1000 bootstrap replicates to estimate

node support. We examined congruence between host and symbiont phylogenies using the

distance-based ParaFit [19] in R 3.5.1 [20]. Root-to-tip distances from the RAxML analyses

for each host and symbiont pair were subjected to Pearson correlation analysis. Branch-length

differences between hosts and symbionts were compared for 27 phylogenetically independent

species pairs across the topology (see figure S1 in ESM). These were calculated using a fixed

topology (derived from the *Blattabacterium* analysis described above) for each of the

following three data sets: 1) 1st+2nd codon sites of protein-coding genes; 2) translated amino

acid sequences; 3) 1st+2nd codon positions of protein-coding genes plus the inclusion of

rRNAs and tRNAs in the case of the mitochondrial data set. Further details on phylogenetic

methods are provided in the ESM.

107 3. Results

In all analyses, there was strong support for the monophyly of each cockroach family with

the exception of Ectobiidae (figure 1). The topologies inferred from the host and symbiont

data sets were congruent ($p = 0.001$). In only two cases was a disagreement found to be

supported by >85% bootstrap support in both trees (the sister group of

Lamproblattidae+Anaplectidae; the sister group to *Carbrunneria paraxami+Beybienkoa*

kurandensis).

We found a correlation between root-to-tip distances for protein-coding genes from

hosts and their symbionts ($R = 0.75$, figure 2*a*). Similar results were found when rRNAs and

tRNAs were included in the host data set ($R = 0.73$, figure S4*a*). The highest rates of

evolution in the host and symbiont data sets (on the basis of branch lengths; see figure 1)

were in members of an ectobiid clade containing *Allacta* sp., *Amazonina* sp., *Balta* sp.,

Chorisoserrata sp., and *Euphyllodromia* sp., and a separate clade containing the

Anaplectidae. After excluding these taxa, evolutionary rates remained correlated, although to

a lesser degree ($R = 0.35$, figure 2*b*). The sharing of branches between taxa in the estimation

of root-to-tip distances renders the data in these plots phylogenetically non-independent and

precludes statistical analysis.

A comparison of branch lengths among phylogenetically independent pairs of host

and symbiont taxa based on protein-coding genes revealed a significant correlation between

their rates of evolution ($R = 0.40$, $p = 0.039$; figure 2*c*, figure S1). Equivalent analyses of

branch lengths inferred from amino acid data also revealed a significant rate correlation

between host and symbiont ($R = 0.43$, $p = 0.023$; figure S3*a*). However, there was no rate

correlation between host and symbiont following inclusion of rRNAs and tRNAs in the host

mitochondrial data set ($R = 0.27$, $p = 0.17$; figure 2*d*). Analyses involving standardization of

branch-length differences yielded significant rate correlations for the protein-coding gene and

amino acid data sets ($R = 0.43–0.48$, $p = 0.011–0.023$; see figures S2, S3), and mixed results

in the case of the inclusion of rRNAs and tRNAs in the host mitochondrial data set ($R =$
0.34–0.40, $p = 0.041$–0.085; see figure S4).

4. Discussion

We have detected a correlation in molecular evolutionary rates between *Blattabacterium* and
host mitochondrial genomes, using two different methods of analysis. To our knowledge, this
is the first evidence of such a correlation in a host-symbiont relationship. Previous studies
found a correlation in evolutionary rates between mitochondrial and nuclear genes in various
animal groups [4-7].

Similar forces acting on the underlying mutation rates of both host and symbiont
genomes could translate into a relationship between their substitution rates. This could
potentially occur if symbiont DNA replication depends on the host's DNA replication and
repair machinery [21]. A better understanding of the level of integration of host-encoded
proteins in the metabolism of *Blattabacterium* will allow further exploration of this issue. A
study of insect genomic data found a correlation in rates between mitochondrial genes and
nuclear genes that encode proteins targeted to mitochondria [7]. However, since
Blattabacterium is not known to be directly associated with mitochondria, interactions
between symbiont and organellar proteins are unlikely to explain the correlations in rates that
we have found here.

Short host generation times could potentially lead to elevated evolutionary rates in host
and symbiont [22], assuming that increased rates of symbiont replication are associated with
host reproduction, as is found in *Blochmannia* symbionts of ants [23]. Variations in metabolic
rate and effective population size between host taxa could also explain the rate correlations
that we have observed. Unfortunately, with the exception of a few pest and other species,
generation time, metabolic rates, and effective population sizes are poorly understood in
cockroaches. This precludes an examination of their influence on evolutionary rates in host
and symbiont.

The addition of mitochondrial rRNA and tRNA data weakened the correlations found
in the branch-length comparisons of species pairs. The reasons for this are unclear but they
might be associated with the conserved nature of tRNAs and the stem regions of rRNAs, or
highly variable loop regions in the latter.

Blattabacterium is a vertically transmitted, obligate intracellular mutualistic symbiont,

whose phylogeny is expected to mirror that of its hosts. This is especially the case for

phylogenies inferred from mitochondrial DNA, since mitochondria are linked with

Blattabacterium through vertical transfer to offspring through the egg cytoplasm. As has been

found in previous studies [24–26], we observed a high level of agreement between the

topologies inferred from cockroach mitochondrial and *Blattabacterium* genome data sets. In

some cases, however, disagreements were observed between well-supported relationships.

The variability in rates that we observed between some lineages, and/or the highly increased

rate of mitochondrial DNA compared with *Blattabacterium* DNA, could be responsible for

these disagreements. Owing to long periods of co-evolution and co-cladogenesis between

cockroaches and *Blattabacterium* [9, 24], potential movement of strains between hosts (for

example, via parasitoids) is not expected to result in the establishment of new symbioses,

especially between hosts that diverged millions of years ago.

In conclusion, our results highlight the profound effects that long-term symbiosis can

have on the biology of each symbiotic partner. The rate of evolution is a fundamental

characteristic of any species; our study indicates that it can become closely linked between

organisms as a result of symbiosis. Further studies are required to determine whether the

correlation that we have found here also applies to the nuclear genome of the host. Future

investigations of generation time, metabolic rate, and effective population sizes in

cockroaches and *Blattabacterium* will allow testing of their potential influence on

evolutionary rates.

Authors' contributions. N.L., D.A.A., T.B., and S.Y.W.H. designed the study. Z.W. and T.

B. collected and provided specimens; D.A.A., T.B. generated sequence data; D.A.A., T.B,

N.L., and S.Y.W.H. performed data analysis and interpreted results; N.L. and D.A.A. wrote

the manuscript, with contributions from T.B. and S.Y.W.H. All authors approved the final

version of the manuscript and agree to be accountable for all aspects of the work.

Funding. D.A.A. was supported by an International Postgraduate Research Stipend from the

Australian Government. S.Y.W.H. and N.L. were supported by Future Fellowships from the

Australian Research Council. T.B. was supported by the Japan Society for the Promotion of

Science KAKENHI 18K14767, and by an EVA4.0 grant (No.

CZ.02.1.01/0.0/0.0/16_019/0000803) from the OP RDE. Z.W. acknowledges funding from

the National Natural Sciences Foundation of China (31872271).

Competing interests. There are no competing interests.

Acknowledgements. We thank Qian Tang, Frantisek Juna, James Walker, and David Rentz

for providing specimens, and Charles Foster for assistance with data curation.

References

1.   Bromham L, Penny D. 2003 The modern molecular clock. *Nat. Rev. Genet.* **4**, 216–224.
2.   Ho SYW, Lo N. 2013 The insect molecular clock. *Aust. J. Entomol.* **52**, 101–105.
3.   Douglas AE. 2010 *The Symbiotic Habit*. Princeton, NJ: Princeton University Press.
4.   Martin AP. 1999 Substitution rates of organelle and nuclear genes in sharks:
implicating metabolic rate (again). *Mol. Biol. Evol.* **16**, 996–1002.
5.   Sheldon FH, Jones CE, McCracken KG. 2000 Relative patterns and rates of evolution
in heron nuclear and mitochondrial DNA. *Mol. Biol. Evol.* **17**, 437–450.
6.   Lourenço JM, Glémin S, Chiari Y, Galtier N. 2013 The determinants of the molecular
substitution process in turtles. *J. Evol. Biol.* **26**, 38–50.
7.   Yan Z, Ye G, Werren JH. 2019 Evolutionary rate correlation between mitochondrial-
encoded and mitochondria-associated nuclear-encoded proteins in insects. *Mol. Biol.*
Evol. **36**, 1022–1036.
8.   Degnan PH, Lazarus AB, Brock CD, Wernegreen JJ. 2004 Host–symbiont stability and
fast evolutionary rates in an ant–bacterium association: cospeciation of *Camponotus*
species and their endosymbionts, *Candidatus blochmannia*. *Syst. Biol.* **53**, 95–110.
9.   Bourguignon T *et al.* 2018 Transoceanic dispersal and plate tectonics shaped global
cockroach distributions: evidence from mitochondrial phylogenomics. *Mol. Biol. Evol.*
35, 970–983.
10. Evangelista DA *et al.* 2019 An integrative phylogenomic approach illuminates the
evolutionary history of cockroaches and termites (Blattodea). *Proc. R. Soc. B* **286**,
20182076.
11. Cornwell PB. 1968 *The Cockroach*. London: Hutchinson.
12. Sacchi L, Corona S, Grigolo A, Laudani U, Selmi MG, Bigliardi E. 1996 The fate of
the endocytobionts of *Blattella germanica* (Blattaria: Blattellidae) and *Periplaneta*
americana (Blattaria: Blattidae) during embryo development. *Ital. J. Zool.* **63**, 1–11.
13. López-Sánchez MJ, Neef A, Peretó J, Patiño-Navarrete R, Pignatelli M, Latorre A,
Moya A. 2009 Evolutionary convergence and nitrogen metabolism in *Blattabacterium*
strain Bge, primary endosymbiont of the cockroach *Blattella germanica. PLOS Genet.*
5, e1000721.
14. Sabree ZL, Huang CY, Arakawa G, Tokuda G, Lo N, Watanabe H, Moran NA. 2012
Genome shrinkage and loss of nutrient-providing potential in the obligate symbiont of

the primitive termite *Mastotermes darwiniensis*. *Appl. Environ. Microbiol.* **78**, 204–
210.

15. Kinjo Y *et al.* 2018 Parallel and gradual genome erosion in the *Blattabacterium*
endosymbionts of *Mastotermes darwiniensis* and *Cryptocercus* wood roaches. *Genome*
Biol. Evol. **10**, 1622–1630.

16. Vicente CSL, Mondal SI, Akter A, Ozawa S, Kikuchi T, Hasegawa K. 2018 Genome
analysis of new *Blattabacterium* spp., obligatory endosymbionts of *Periplaneta*
fuliginosa and *P. japonica*. *PLOS ONE* **13**, e0200512.

17. Kambhampati S, Alleman A, Park Y. 2013 Complete genome sequence of the
endosymbiont *Blattabacterium* from the cockroach *Nauphoeta cinerea* (Blattodea:
Blaberidae). *Genomics* **102**, 479–483.

18. Stamatakis A. 2014 RAxML version 8: a tool for phylogenetic analysis and post-
analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313.

19. Legendre P, Desdevises Y, Bazin E. 2002 A statistical test for host–parasite
coevolution. *Syst. Biol.* **51**, 217–234.

20. R Core Team. 2018 R: A language and environment for statistical computing. Vienna,
Austria: R Foundation for Statistical Computing.

21. Moran NA, Bennett GM. 2014 The tiniest tiny genomes. *Annu. Rev. Microbiol.* **68**,
195–215.

22. Bromham L. 2009 Why do species vary in their rate of molecular evolution? *Biol. Lett.*
5, 401–404.

23. Wolschin F, Hölldobler B, Gross R, Zientz E. 2004 Replication of the endosymbiotic
bacterium *Blochmannia floridanus* is correlated with the developmental and
reproductive stages of its ant host. *Appl. Environ. Microbiol.* **70**, 4096–4102.

24. Lo N, Bandi C, Watanabe H, Nalepa C, Beninati T. 2003 Evidence for cocladogenesis
between diverse dictyopteran lineages and their intracellular endosymbionts. *Mol. Biol.*
Evol. **20**, 907–913.

25. Clark JW, Hossain S, Burnside CA, Kambhampati S. 2001 Coevolution between a
cockroach and its bacterial endosymbiont: a biogeographical perspective. *Proc. R. Soc.*
B **268**, 393–398.

26. Garrick RC, Sabree ZL, Jahnes BC, Oliver JC. 2017 Strong spatial-genetic congruence
between a wood-feeding cockroach and its bacterial endosymbiont, across a
topographically complex landscape. *J. Biogeogr.* **44**, 1500–1511.

274
275

Figure captions

Figure 1. Congruence between (*a*) phylogenetic tree of host cockroaches inferred using maximum likelihood from whole mitochondrial genomes, and (*b*) phylogenetic tree of *Blattabacterium* inferred using maximum likelihood from 104 protein-coding genes (3rd codon sites excluded from both data sets). Shaded circles at nodes indicate bootstrap values (black = 100%, grey = 85–99%). Nodes without black or grey circles have bootstrap values <85%. Red outlines on circles indicate disagreement between the phylogenies. Branches are coloured according to their membership of different cockroach families.

Figure 2. Comparison of evolutionary rates of *Blattabacterium* symbionts and their host cockroaches. (*a*) Correlation of root-to-tip distances in phylogenies of *Blattabacterium* and cockroaches, inferred using maximum-likelihood analysis of protein-coding genes from each data set, with 3rd codon sites excluded. (*b*) Correlation of root-to-tip differences following the removal of five rapidly evolving ectobiid taxa (*Amazonina* sp., *Chorisoserrata* sp., *Allacta* sp., *Balta* sp., and *Euphyllodromia* sp.) and two anaplectids. Colours represent data from representatives of different cockroach families, as shown in the colour key. Grey circles represent internal branches. (*c*) Correlation of log-transformed branch-length differences between phylogenetically independent pairs of host and symbiont taxa, based on protein-coding genes only*,* and (*d*) with the addition of rRNAs and tRNAs to the host mitochondrial data set.

(*a*) **Cockroach mtDNA**

(*b*) *Blattabacterium*

Lamproblattidae — *Lamproblatta* sp.
Anaplectidae — *Anaplecta calosoma* / *Anaplecta omei*
Blattidae — *Periplaneta americana*, *Shelfordella lateralis*, *Protagonista lugubris*, *Blatta orientalis*, *Deropeltis paulinoi*, *Melanozosteria* sp., *Platyzosteria* sp., *Cosmozosteria* sp., *Methana* sp., *Eurycotis decipiens*
Tryonicidae — *Tryonicus* sp.
Cryptocercidae — *Cryptocercus hirtus*, *Cryptocercus punctulatus*
Isoptera — *Mastotermes darwiniensis*
Corydiidae — *Eupolyphaga sinensis*, *Therea regularis*
Ectobiidae — *Ectoneura hanitschi*, *Ectobius* sp., *Phyllodromica* sp., *Amazonina* sp., *Chorisoserrata* sp., *Euphyllodromia* sp., *Allacta* sp., *Balta* sp., *Anallacta methanoides*, *Megaloblatta* sp., *Blattella germanica*, *Parcoblatta virginica*, *Paratemnopteryx couloniana*, *Carbrunneria paramaxi*, *Beybienkoa kurandanensis*, *Ischnoptera deropeltiformis*
Blaberidae — *Epilampra maya*, *Galiblatta cribrosa*, *Paranauphoeta circumdata*, *Opisthoplatia* sp., *Rhabdoblatta* sp., *Neolaxta mackerrasae*, *Laxta* sp., *Panesthia angustipennis*, *Panesthia* sp., *Macropanesthia rhinoceros*, *Pycnoscelus femapterus*, *Panchlora nivea*, *Gyna capucina*, *Blaptica dubia*, *Blaberus giganteus*, *Eublaberus distanti*, *Gromphadorhina grandidieri*, *Elliptorhina davidi*, *Aeluropoda insignis*, *Nauphoeta cinerea*

0.05 subs/site   0.05 subs/site

299   Figure 1

300
301 Figure 2
302