Socio-spatial heterogeneity in participation in mass dog vaccination campaigns, Arequipa, Peru

Ricardo Castillo-Neyra*, Amparo M. Toledo, Claudia Arevalo-Nieto, Hannelore MacDonald, Micaela DelaPuente, Cesar Naquira-Velarde, Valerie A. Paz-Soldan, Alison M. Buttenheim, Michael Z. Levy

* Corresponding author

E-mail: cricardo@pennmedicine.upenn.edu (RCN)

Short title: Socio-spatial heterogeneity in mass dog vaccination

Keywords:

Community participation, Disease outbreaks, Dog diseases, Immunization programs, Mass vaccination, One Health, Rabies, Vaccination coverage, Zoonoses.

Abstract

To control and prevent rabies in Latin America, mass dog vaccination campaigns (MDVC) are implemented mainly through fixed-location vaccination points: owners have to bring their dogs to the vaccination points where they receive the vaccination free of charge. Dog rabies is still endemic in some Latin-American countries and high overall

dog vaccination coverage and coverage evenness are desired attributes of MDVC to halt

rabies virus transmission. In Arequipa, Peru, we conducted a door-to-door post-campaign

survey on >6,000 houses to assess the placement of vaccination points on these two

attributes. We found that the odds of participating in the campaign decreased by 16% for

every 100 m from the owner's house to the nearest vaccination point (p=0.041) after

controlling for potential covariates. We found social determinants associated with

participating in the MDVC: for each child under 5 in the household the odds of

participating in the MDVC decreased by 13% (p=0.032), and for every ten more years

living in the area, the odds of participating in the MDVC increased by 9% (p<0.001),

after controlling for distance and other covariates. We also found significant spatial

clustering of unvaccinated dogs over 500m from the vaccination points, which created

pockets of unvaccinated dogs that may sustain rabies virus transmission.

Understanding the barriers to dog owners' participation in community-based dog-

vaccination programs will be crucial to implement effective zoonotic disease preventive

activities. Spatial and social elements of urbanization play an important role in coverage

of MDVCs and should be considered during their planning and evaluation.

1. Introduction

The city of Arequipa is in the midst of a dog rabies outbreak. The introduction of rabies

virus into the city has been ascribed to the transport of rabid dogs from the rabies-

endemic state of Puno during human migration [1-3], and it is likely that the persistence

of transmission is due to low coverage in the annual city-wide dog vaccination campaigns

[3]. Following the detection of the outbreak in Arequipa city in 2015, the Ministry of

Health of Peru (MOH) initiated additional vaccination campaigns in the city with varying

intensity [4]. These additional efforts have not quelled the epidemic: more than 150 rabid

dogs have been detected as of 2019.

Epidemics of dog rabies are ongoing in major urban centers across Latin America and

worldwide [1,5-8]. Since bites from rabies-infected dogs cause 99% of human rabies

deaths [9], the control and elimination of dog-mediated human rabies is based on a

population-wide animal-centered strategy: mass dog rabies vaccination [8,10,11]. Dog

vaccination has dramatically decreased the global burden of human rabies since 1955

[5,12,13] and specifically in the Americas, national programs centered around mass dog

vaccination have achieved enormous advances [8,14,15], reducing the incidence of dog

rabies by 98% since 1983 [14]. In most rabies-affected countries, the official health

organizations (e.g. MOH) organize annual MDVC that are held in outdoor settings. These

campaigns are usually free of charge and voluntary [8,16] and campaign promotion

varies greatly in format, content and intensity [1,17,18]. There are three non-mutually

exclusive strategies to implement MDVC: fixed vaccination posts, mobile teams setting

up temporary mobile post or conducting 'street vaccination', and door-to-door

vaccination. For the fixed-point strategy the vaccinators wait for the dog owners to bring

the dogs to a unique place. For the mobile team strategy, the vaccinators move from one

location to another, waiting for the dog owners to bring the dogs to these moving

locations. For the door-to-door strategy, vaccinators knock on doors asking for dogs to

vaccinate. Locations of the fixed vaccination sites are typically determined by a

combination of convenience and prominence of the location (e.g. the entrance to a health

post) [11]. In Peru, routes for door-to-door and mobile team approaches may or may not

be decided in advance, and teams may move during the course of the day looking for

high-demand locations.

The fixed-point strategy has been extensively used in Latin America and Africa, even

though it has frequently failed to reach the appropriate coverage [1,8,19-23]. The main

reasons for its extensive application is that fixed-point vaccination is easier to implement

and less costly than other strategies [18]. However, in many cases fixed-point is

combined with other strategies. The use of multiple strategies becomes more common

when initial activities are unsuccessful [19,23,25]. However, high dog owner

participation in MDVC and other dog-centered health campaigns (e.g. de-worming dogs

to prevent human hydatid disease) has proven difficult to achieve in many areas [23,24].

It is necessary to understand barriers to community-based control strategies targeting dog

populations in dog rabies-affected countries where coverage does not reach the minimum

70% recommended by the World Health Organization to attain herd immunity [11], much

less the 80% recommended by the Pan-American Health Organization for the region [25].

In cities, the social and spatial aspects of urbanization can facilitate the emergence of dog

rabies and complicate its control [26-28]. In Arequipa, the locations of rabid dogs have

been associated with urban structures [26], and dog owners from areas with different

levels of urbanization have reported distinct factors associated with vaccinating their

dogs against rabies [1]. The changing urban landscape and social processes in rapidly

growing cities have been associated with uptake of health-related services [29-32] and

may be related to the low dog vaccination coverage in Arequipa. The objectives of the

present study were to quantitatively assess the effect of distance to a vaccination point on

dog owner participation in mass dog vaccination campaigns in an urban setting and to

evaluate the effect of such distance on overall vaccination coverage and evenness.

2. Methods

2.1. Ethics statement

Ethical approval was obtained from Universidad Peruana Cayetano Heredia (approval

number: 65369), Tulane University (approval number: 14–606720), and University of

Pennsylvania (approval number: 823736).

2.2. Study setting

The study was conducted in the Alto Selva Alegre (ASA) (human population for 2015:

82,412; density: 11,902 people/km^2), one of the 14 districts of the city of Arequipa.

Arequipa, Peru's second largest city, is home to 969,000 people and is situated at ~2,300

meters above sea level. The first detection of a rabid dog in the city of Arequipa occurred

in March 2015 in ASA. By June 2016, when our data were collected, 43 rabid dogs had

been detected in 8 districts, but it is assumed that number represents a small fraction of

the total number of cases [26]. The city of Arequipa comprises communities spanning

different stages of urbanization and different migration histories, from old established

neighborhoods, to young neighborhoods, to recent invasions [33]. Within this gradient of

development, young neighborhoods and recent invasions are often located on the

periphery of the city (peri-urban area) and the older localities are nearer to the center

(urban area) [33]. Compared to the urban area, peri-urban areas generally have lower

socioeconomic status, fewer community resources, more security problems, and often

more rugged and uneven terrain (Figure 1). As new neighborhoods mature into

established neighborhoods with wealthier residents, homes are improved with better

quality construction material and permanent utility connections, and connectivity with the

rest of the city increases with better sidewalks, roads, and transportation access. ASA

transects the city, running from the center to the periphery, and the district continues to

grow towards the outskirts of the city. In our study, participants represented either the

urban or peri-urban areas of the city of Arequipa. We included 21 urban neighborhoods

founded many decades ago, and 9 peri-urban neighborhoods that originated around 2000

or later.

In ASA, the MOH conducted a mass dog vaccination campaign in June 2016. A detailed

description of the mass dog vaccination campaigns can be found elsewhere [1], and the

dog population estimator used in Arequipa can be found in [26].

126
127
128 **Figure 1**. Study communities represent landscape heterogeneity. A: Urban area. B: Peri-
129 urban.

2.3.   Data collection

In collaboration with the MOH, we georeferenced every stationary and mobile

vaccination team during the three weekends when the campaign was implemented in

ASA. Due to volunteer assistance, some dogs were vaccinated in weekdays during the

campaign; we did not georeference the location of those volunteering vaccinators, mainly

because their schedule was haphazard and unpredictable. We started door-to-door

surveys immediately after the vaccination campaign.

The door-to-door survey was designed based on the rabies literature and based on our

qualitative studies on local communities, where we found specific barriers at the

household level [1]. We collected household variables (e.g. number of household

members; number of children under 5 years old), dog owner or interviewee variables (e.g.

gender; educational attainment), and dog variables (e.g. vaccination status; age). All

houses in the study localities were geocoded and the survey data were linked to the

household coordinates. We estimated the Euclidean distance between households and the

closest vaccination point (fixed, mobile, or either).

2.4.   Statistical Analysis

We estimated the total vaccination coverage in the study area and compared the coverage

in urban vs. peri-urban localities with chi-squared test. We estimated the human to dog

ratio and bootstrapped it 10,000 times to estimate its confidence interval. To evaluate the

baseline characteristics of households and dog owners by participation in the MDVC, we

defined an ordinal outcome for houses with dogs: *no participation* (no dog vaccinated in

the house), *partial participation* (some, but not all dogs in the house, vaccinated), and *full*

participation in the MDVC (all dogs in the house vaccinated). We used a chi-square test

to compare categorical variables with 10 or more observations per group, Fisher's exact

test for categorical variables with fewer than 10 observations in any subgroup, and

Mann–Whitney U test for age of the dog owner or interviewee, which did not follow a

normal distribution and was trunked at 18 years. We compared the individual

characteristics of vaccinated and unvaccinated dogs with chi-square for categorical

variables and with Mann–Whitney U test for dog's age.

Our main objective was to assess the effect of distance to the vaccination point on

participation in the MDVC. For distance to the vaccination point, we used the Euclidean

distance from the dog's house to the closest vaccination point, either fixed or mobile. For

participation in the MDVC we used the ordinal values described above: *no participation*,

partial participation, and *full participation.* We compared proportional odds logistic

regression (POLR), non-proportional odds logistic regression, and multinomial

regression. The ordinal models were superior to the multinomial regression, and the

proportional odds assumption holds for most of the covariates. Given that the categories

of participation are inherently ordinal and that providing one point estimate per covariate

is more interpretable, the POLR model was favored. To adjust the model, based on the

recent literature, we chose this set potential of covariates: having a dog leash at home,

number of children under 5 years old at home, time living in the area, rabies status of the

last place they lived in before living in the study area, number of dogs at the house, age

and gender of the dog owner, and educational attainment. We considered transformed

distance to capture non-linear effects and interactions between distance and having a

leash. The fit of the alternative models to the data was compared with Akaike's

Information Criteria (AIC). We also attempted building a hierarchical model to take into

account the spatial autocorrelation within locality. However, given that within each

locality there was at most one vaccination point, the variable distance from the house to

the vaccination point would be unidentifiable under such hierarchical model. The final

POLR model fitted with the R package MASS (REF) was:

$$ln\left\{\frac{Odds(Y \leq k)}{Odds(Y > k)}\right\} = b_0 + b_1.distance + b_2.leash + b_3.children + b_4.\underset{in\ the\ area}{time\ living} + b_5.\underset{previous\ residence}{rabies\ status\ of} + \varepsilon$$

Where k takes the values 0 (no participation), 1 (partial participation), and 2 (full

participation). All statistical tests were 2-sided, and significance level was 0.05.

We tested the spatial pattern of vaccinated and unvaccinated dogs in relation to

vaccination tents for clustering using the bivariate cross K-function. This function

estimates spatial dependence between two types of points (i.e. unvaccinated dogs and

vaccination points) by measuring the expected number of points of type i within a given

distance to a point of type j divided by the overall density of the points of type i. We used

the Kcross function in the R package spatstat [34] to estimate deviations between the K

function estimated for our data and the theoretical K function corresponding to a

completely random Poisson point process for vaccinated dogs to tents and unvaccinated

dogs to tents. Deviations greater than the theoretical K function indicate that the mean

point count is higher than expected under complete spatial randomness (CSR) and thus

some degree of clustering is present between the two event types at the indicated

distance. Similarly, deviations less than the theoretical K function indicate that the mean

point count is lower than expected under CSR which therefore indicates that some degree

of dispersion is present between the two event types at the indicated distance.

In order to investigate the effect of geolocation on the odds of canine vaccination we

fitted Generalized Additive Models (GAMs) to our data using the R package MapGAM

[39]. GAMs are an extension of linear regression models in which both parametric and

non-parametric terms are used to estimate the outcome of interest. We used a two-

dimensional locally weighted smooth (LOESS) of latitude and longitude for our non-

parametric term. The LOESS smoother fits each data point by weighting it towards

nearby points, where weighting is based on the distance to the point being fitted. The

percentage of data points in the region that will be used to predict a particular point is

referred to as the span. The optimal span size used for smoothing was determined by

minimizing Akaike's Information Criterion. We mapped the odds ratio for each point on

pre-specified grids of each locality (from polygon data) and next tested the null

hypothesis that the odds of each points' vaccination status did not depend on geolocation

using permutation tests. For each test, the paired latitude and longitude coordinates were

randomly permuted but vaccination status was held fixed. 1000 permutations were run

for each locality and contour lines encircle areas with significantly increased or decreased

vaccination odds as indicated by point wise p-values computed from the permutation

ranks. All models and figures were created with R [35].

3. Results

The vaccination coverage of the MOH MDVC was only 58.1%, and it was low in both urban and peri-urban localities (58.0% vs. 58.6% respectively, chi2=0.086, p=0.769). Only 3.4% of dogs were (reportedly) vaccinated in private clinics, bringing our estimated total coverage to 61.5%. Participation in our survey was higher in the urban area (88.8%) compared to the peri-urban area (61.6%) (mean= 82.0%, chi2=6458.5, p<0.001). The total number of dogs in the surveyed houses was 5,292 and the human to dog ratio was 3.78:1 (95% CI: 3.69:1 - 3.89:1). In total, 65.3% of surveyed houses had dogs, but this number was higher in peri-urban areas (70.0% compared to 64.6% in urban areas, chi2=6.529, p=0.011). For 76.9% of vaccinated dogs in the area, the person who took them to the MDVC is the person who responded the survey. In our study area the urbanization process involves new localities being founded and settlers moving in. Accordingly, we found that the time of residency (or the year people moved in to this area) was clustered at the locality level. However, we found that people living in or founding a locality do not necessarily share the same place of origin or previous residence.

When we compared the distance from each house to the closest fixed vaccination point, to the closest mobile vaccination point, and to either the closest fixed or mobile vaccination point, we found a clear gradient with higher average distances for non-participant houses to houses that partially participated to houses that fully participated (Table 1). The proportion of households with children under 5 was higher in households that did not participate of the campaign (36%) compared to houses that participated fully

or partially of the campaign (31% and 29% respectively). The proportion of houses with

a dog leash increases from those who did not participate, to those who participated

partially, to those that fully participated of the MDVC. Houses that did not vaccinate their

dogs reported not knowing about the campaign nine times more than houses that

vaccinated all or some of their dogs. There were some differences in MDVC participation

by migration history: people who have lived for longer in ASA tend to report higher

participation in the MDVC compared to those who have lived fewer years in ASA. Also,

there is a slight difference in participation in the MDVC depending on rabies status of

previous residence, with more people participating in the campaign, partially or fully, if

they were migrants from a rabies endemic area (Table 1). Other variables, such as

educational attainment, the proportion of female dog owners or interviewees, and the

proportion of households in urban localities were all similar in those households that

participated fully or partially in the campaign compared to those households that did not

participate of the campaign (Table 1). Having multiple dogs is a prerequisite to be in the

partial participation group; therefore, houses with partial participation had on average

more dogs, but the number of dogs per house was very similar in houses that participated

fully compared to those that did not participate in the campaign (Table 1). Those who did

not participate in the MDVC reported more frequently not knowing about the campaign

before it happened (Table 1), but many of them reported learning about the campaign the

same day it occurred. In S1_Table, we report the media channels through which they

learned about the campaign either after it occurred or the same day it occurred.

Table 1. Characteristics of the dog owner population in the study area by participation in
the MDVC, Arequipa City, Peru, 2016.
266

Interviewee/Owner Variables	Vaccinated no dog (n=1,017)	Vaccinated some dogs (n=176)	Vaccinated all dogs (n=1,434)	p
Distance to any closest vaccination point – Mean (SD)	125m (96)	117m (114)	103m (83)	<0.001 [a]
Distance to closest fixed vaccination point – Mean (SD)	501m (479)	490m (420)	461m (422)	0.119 [a]
Distance to closest mobile vaccination point – Mean (SD)	137m (101)	127m (119)	114m (93)	<0.001 [a]
Number of dogs in houses with dogs – Mean (SD)	1.79 (1.10)	2.96 (1.24)	1.80 (1.05)	<0.001 [a]
Houses with children under 5	36.4%	28.9%	30.8%	0.019 [b]
Age of dog owner/interviewee – Median (IQR)	39 years (28 - 52)	40 years (29 - 50)	42 years (30 - 53)	0.024 [c]
Time living in ASA if migrant – Mean (SD)	18.1 years (13.7)	21.2 years (13.9)	22.1 years (14.8)	<0.001 [a]
Area before living in ASA				0.036 [b]
Rabies epidemic (Arequipa)	90.7%	94.1%	89.8%	
Rabies endemic (Puno)	1.6%	2.4%	3.3%	
Rabies free (rest of country)	7.7%	3.6%	6.9%	
Educational attainment				0.509 [b]
Illiterate	0.6%	0.5%	0.6%	
Primary school	11.5%	11.4%	13.6%	
High school	43.2%	48.3%	41.1%	
Technical academy	21.2%	15.3%	20.6%	
University	22.3%	23.3%	23.3%	
Preferred not to respond	1.2%	1.1%	0.7%	
Owns at least one dog leash	51.0%	54.5%	60.1%	<0.001 [b]
Female interviewees/owners	64.6%	73.3%	64.4%	0.032 [b]
Did not know about the campaign before it happened	15.4%	2.3%	1.3%	<0.001 [b]
Live in urban locality (vs. peri-urban)	78.8%	78.1%	79.4%	0.002 [b]

p values estimated with [a] one-way ANOVA; [b] Chi square test; and [c] Kruskal Wallis test.

Compared to vaccinated dogs, unvaccinated dogs were older, they were more likely

females, they had more free access to the street, there was no leash for them at home, and

they were less likely to be walked. It seems that multipurpose dogs (dogs reported as

guard and company dogs) were more likely to be vaccinated (Table 2). It does not seem

that dogs from a specific source have higher chances of being vaccinated. If any, dogs

received as gifts are more likely to be vaccinated and dogs born at home or

adopted/picked on the street are less likely to be vaccinated (Table 2). Being considered

purebred or being spayed/neutered was not associated with dog vaccination status (Table

2).

Table 2. Dog characteristics by vaccination status in the MDVC (dogs vaccinated
privately not included).
280

Dog Variables	Unvaccinated dog (n=2,019)	Vaccinated dog (n=2,800)	p
Age of dogs – Mean (SD)	24 months (12 – 60)	20 months (10 – 54)	0.039 [a]
Female dogs	42.1%	36.8%	<0.001 [b]
Free access to the street	20.6%	25.7%	0.002 [b]
Leash for the dog	33.8%	45.3%	<0.001 [b]
Dogs that are walked, either with or without leash.	55.9%	66.9%	<0.001 [b]
Dog function			<0.001 [b]
Company	52.0%	49.0%	
Protection	23.4%	20.9%	
Company & Protection	24.6%	29.6%	
Breeder/business	0.0%	0.5%	
Dog source			<0.001 [b]
Gift	51.2%	56.5%	
Bought at market/store	17.4%	18.3%	
Born at home	15.6%	12.3%	
Adopted/Picked on the street	12.3%	10.8%	
Bought from friend/neighbor	2.6%	2.0%	
Do not know	1.0%	0.0%	
Purebred as reported by interviewee	25.7%	24.9%	0.589 [b]
Spayed/Neutered	3.5%	2.6%	0.157 [b]

p values estimated with [a] Student's t-test; and [b] Chi square test.

We found that distance to the vaccination site is strongly associated with participation of

the MDVC. The odds of not participating in the MDVC were 16% lower for someone

who lived 100 meters farther from the vaccination point after adjusting for other

covariates and this difference was statistically significant (Table 3). Those having a dog

leash at home had 35% higher odds of participation of the MDVC, either fully or

partially, compared to those who did not have a dog leash after adjusting for other

covariates. The odds of participating in the MDVC, either fully or partially, were 13%

lower for each additional child under 5 years at home after adjusting for other covariates.

The time living in the area was also associated with participation in the MDVC; the odds

of participating in the campaign were 9% higher when we compared people who had

lived 10 years longer in the area with those who moved to the area more recently.

Another component of migration history, the previous residence region, was also

associated with participating in the MDVC: those whose previous residence was a rabies-

free region or was any district within Arequipa were 23% to 32% less likely to participate

of the campaign compared to those whose previous residence was a rabies-endemic

region, after adjusting for other covariates (Table 3). Demographic variables such as

owner's/interviewee's age, gender and educational attainment, and other house-level

variables, such as number of dogs at the house, were dropped during model selection

because they neither improved the fit of the model nor were statistically associated with

participating in the MDVC.

Table 3. Factors associated with participating in the Mass Dog Vaccination Campaign,
Arequipa, Peru.
306

Variable	OR	95% CI	p
Distance to closest vaccination point (100m)	0.84	0.76 – 0.98	0.041
Having a leash at home	1.35	1.10 – 1.61	0.003
Number of children U5Y at home	0.87	0.78 – 0.97	0.032
Time living in ASA (10 more years)	1.09	1.06 – 1.11	<0.001
Rabies status of previous residence			
Endemic for rabies	Ref.		
Epidemic for rabies	0.77	0.63 – 0.93	0.006
Free of rabies	0.68	0.57 – 0.82	<0.001

Odds ratios estimated with multiple proportional odds logistic regressions.

The significant effect of distance to the vaccination point on the odds of participating in

the MDVC has consequences for the distribution of unvaccinated dogs in the area. We

observed spatial clustering of unvaccinated owned dogs as a function of the distance from

the house to the vaccination point; that is, unvaccinated dogs are closer to each other than

expected by chance. These pockets of unvaccinated dogs occur at 500 meters from the

vaccination point or further (Figure 2). We also analyzed the spatial odds of participating

in the MDVC, that is, the effect of their specific geolocation with regards to the location

of the vaccination point. For areas served by fixed-point vaccination, there was a clear

smooth spatial effect with higher odds of participating for houses closer to the

vaccination point and a decreasing gradient farther away from the vaccination point. The

spatial effect of the fixed-point vaccination strategy creates two clearly defined zones: a

large zone with statistically significantly high odds of participation in the MDVC and

another large zone with statistically significantly low odds of participation in the MDVC

(Figure 3). For areas served by mobile teams, there were more spots of significant low

and high odds of participation in the MDVC and these spots were spread in the study area

without a clear association between the spots and the locations where the vaccination

teams stopped to wait for dogs or to vaccinate dogs (Figure 3). The spatial odds shown in

maps A and B in Figure 3 are relative to themselves. When the odds found in the area

served by mobile teams were adjusted to the scale of the area served by fixed teams, the

odds of participating in most of the area are similar to the odds of participating for

someone who lives mid-distance from the fixed point in map A.

332

333
334

335   **Figure 2.** Clustering of unvaccinated dogs as a function of distance from the vaccination
336   point.
337

339
340   **Figure 3.** Spatial odds ratios for participating in the MDVC in a locality served by a
341 fixed vaccination point (cross in A), and by a mobile team (in B, X's represent locations
342 where the mobile team stopped to wait for dogs or to vaccinate dogs). The insert in B
343 shows the spatial odds for that area after adjusting the odds scale to the scale in A.
344 Contour lines represent areas where the spatial odds were statistically significant.
345

4. Discussion

In Arequipa, the social and spatial aspects of urbanization facilitate the emergence of dog

rabies and complicate its control. In 21 urban and 9 peri-urban localities in Arequipa,

Peru, we found low vaccination coverage and coverage that was spatially uneven. We

found a strong effect from a more proximal explanation for low participation in the

MDVC: distance to the vaccination point. The unadjusted data show a clear negative

gradient with higher levels of participation in the MDVC at shorter distances to the

vaccination point. After accounting for other important individual- and household-level

variables, distance to the vaccination point remains an important factor associated to

participating to the campaign. The effect of distance on participation in the MDVC also

has effects on the spatial coverage of vaccination. We found areas with statistically

significant lower odds of dogs being vaccinated, and the LOESS smoother maps

correlated well with maps of vaccination coverage. Therefore, it seems that both fixed

point and mobile team canine vaccination approaches produced spatially heterogeneous

vaccination coverage. However, we found that vaccination coverage was more "patchy"

coverage in localities served by mobile vaccination teams. This combination of mobile

and fixed points was used also in 2015, but the same localities are not always served with

the same approach (e.g. a locality that was served with mobile teams in 2015 could be

served with fixed point vaccination in 2016). As others have reported [36], there is

potential that in some localities owners in 2016 expected that the MDVC would be

brought to their doors and did not plan or intend to bring the dogs to the fixed points in

their areas. Spatially heterogeneous vaccination coverage is undesirable for dog rabies

control and elimination. Townsend et al. [37] found that such patchy coverage can

"profoundly damage prospects of elimination […] by creating pockets where rabies could

persist", and modeled patchy coverage within 1 km^2 cell grids. The low coverage

'patches' in our study had smaller areas than 1 km^2, thus the potential for a threat to

elimination efforts may be different or non-existent. However, in these densely populated

areas it is unknown if these 'patches' are large enough to sustain rabies transmission in

the city.

Many studies have explored logistical, informational, social and structural barriers for

dog rabies vaccination experienced by owners in rabies-affected areas

[1,17,18,22,23,36,38-48]. Two of the most common reasons identified are difficulty

handling the dogs [1,22,23,44,45,47,48] and lack of time [1,23,44,45,47,48]. These two

logistical barriers are correlated with a less studied element: the distance to the

vaccination campaigns. Distance to health services has been fairly well studied in terms

of availability of and access to health care and impact on health, especially for maternal

health, treatment and prevention of chronic diseases and treatment adherence for

infectious diseases [49-51], but not as much for preventing infectious diseases. Some

rural studies mention distance as a potential factor for low dog vaccination coverage

[22,52] and two studies directly evaluated the effect of distance on overall rural villages

vaccination coverage [45] and the effect of distance on attendance to the MDVC in Sub-

Saharan Africa [18]. In the sub-Saharan Africa study, researchers found that distance in

dispersed communities have an impact on MDVC attendance [18]. Given the hilly

landscape with rare direct paths between houses and vaccination points, they estimated

the shortest-path distance for their analysis to take into consideration the long and

tortuous routes dog owners had to follow to visit the vaccination sites. In our study area,

with non-dispersed urbanized highly-populated localities and high density of street

intersections that increase walkability, distance is still an important proximal explanation

for low participation in the vaccination campaigns.

Our study area consisted of urban and peri-urban localities. Surprisingly, there were no

clear differences in participation in the MDVC between these two groups. However, the

distribution of other proximal rabies-related characteristics is different between them (e.g.

more free-roaming dogs in peri-urban areas, more neutered/spayed dogs in urban areas).

There are other social determinants that provide more distal explanations for participation

in the MDVC. In previous focus groups conducted by our team, young females reported

that having a baby at home could prevent them for participating in the campaign [1].

Similarly, we found that households with children under 5 years old were less likely to

vaccinate their dogs compared to houses without children, and each additional child

under 5 reduced the odds of vaccinating the dogs in the house. This knowledge suggests

that redirecting the attention to the dog to protect the household children from rabies

could increase participation in the MDVC for this group of households. There was also a

clear difference in participation between those who lived in a dog rabies-endemic area

before living in Arequipa. A possible explanation for that difference is higher awareness

among that group. In our focus groups, we found low awareness and low perception of

severity among residents of Arequipa. Interestingly, another component of migration

history was also associated with participation in the MDVC: time living in the area. This

phenomenon has been observed for the utilization of other health services in different

settings and populations [53-55]. Migration, settlement and adaptation are processes that

take time and are necessary for the uptake of health services [56-59], and could be

influencing the participation in the MDVC. Importantly, in the peri-urban localities there

are more households with children under 5, more recent migrants, and more people

whose previous residence was in a rabies-affected region.

Our study has a number of limitations. Other studies have focused attention on dog-level

variables (e.g. sex, age, function) that might be associated to participation in MDVC [18].

We did not analyze these; rather we focus on owner and community characteristics that

can be utilized by the health authorities (who rarely have the opportunity to collect

detailed house-by-house information) to increase participation in MDVC. Vaccination

status and access to the street were reported by the owner/interviewee. Given the bad

publicity in the local media about owned free roaming unvaccinated dogs and the

authorities' threats to fine 'irresponsible' dog owners, there is potential for social

desirability bias to inflate our estimate of vaccination coverage and deflate that of the

proportion of dogs that have access to the street. We did not ask the interviewees to show

the vaccination certificate they receive at the MDVC since many of them do not save

these certificates. We used Euclidean distances, which are only a proxy for the real

distance traveled by individuals to a vaccination post.

Dog rabies virus transmission continues in Arequipa, putting at risk millions of people in

the city and the surrounding departments. Dog-focused public health strategies are not

limited to rabies: deworming dogs to prevent human echinococcosis [60], use of

insecticide treatment [61] or vaccines [62] on dogs to prevent human Leishmaniasis or

Chagas disease, are just a few examples. These programs, if they are to be successful,

require high coverage and evenness in their implementation. The same approaches to

reach the appropriate levels of community participation that might have worked in the

1980s are not working today. Understanding the barriers for dog owners' participation in

community-based programs will be crucial to implement effective zoonotic disease

preventive activities. Distance to health services and the heterogeneous social

composition of growing cities have to be taken into consideration when designing field

programs to protect against zoonotic diseases.

Supporting information

S1_Resumen en Español (Abstract in foreign language).

S1_Table. MDVC communication channels interviewees were exposed by participation

level.

How they learned about the campaign	Vaccinate none dog (n=858)	Vaccinated some dogs (n=170)	Vaccinated all dogs (n=1,397)	p
Megaphone	56.9%	64.5%	66.5%	<0.001 [a]
TV	36.4%	33.1%	32.1%	0.842 [a]
Radio	33.9%	23.3%	30.4%	0.122 [a]
Poster/Banner	7.8%	9.9%	7.4%	0.337 [a]
Relative/friend/neighbor	4.8%	8.7%	5.7%	0.027 [a]
Newspaper	1.3%	0.6%	1.1%	0.798 [a]
Flier	1.3%	1.2%	0.7%	0.497 [b]
At municipality	0.6%	0.6%	0.6%	--
Social media	0.1%	0.6%	0.7%	0.066 [b]
Community meeting	0.1%	0.0%	0.0%	0.454 [b]

455   p-values estimated with [a] Chi square test and [b] Fisher exact test.
456

5. Acknowledgments

458   We thank all the survey participants for agreeing to be part of our study and sharing their

459   time with us. We acknowledge the work of the members of the Zoonotic Disease

460   Research Laboratory for their contribution to the implementation of this study, especially

461   Katty Borrini and Jorge Apaza. This work was partially supported by a grant from NIH-

462   NICHD 5R01HD075869.

6. References

1. Castillo-Neyra R, Brown J, Borrini K, Arevalo C, Levy MZ, Buttenheim A, et al. Barriers to dog rabies vaccination during an urban rabies outbreak: Qualitative findings from Arequipa, Peru. PLoS neglected tropical diseases. 2017;11: e0005460.

2. Castillo-Neyra R, Levy MZ, Náquira C. Efecto del sacrificio de perros vagabundos en el control de la rabia canina. Rev perú med exp salud publica. 2016;33: 772. doi:10.17843/rpmesp.2016.334.2564

3. Ministerio de Salud del Peru - Direccion General de Epidemiologia. Alerta Epidemiológica [Epidemiologic Alert]. Lima; 2015.

4. El Presidente de la República. Decreto Supremo que declara en Emergencia Sanitaria por el plazo de noventa (90) días calendario, a la provincia de Arequipa del departamento de Arequipa. 2016;: 582224–582229.

5. Aréchiga Ceballos N, Karunaratna D, Aguilar Setién A. Control of canine rabies in developing countries: key features and animal welfare implications. Revue scientifique et technique (International Office of Epizootics). 2014;33: 311–321.

6. Wallace RM, Reses H, Franka R, Dilius P, Fenelon N, Orciari L, et al. Establishment of a Canine Rabies Burden in Haiti through the Implementation of a Novel Surveillance Program. PLoS neglected tropical diseases. 2015;9: e0004245.

7. Widdowson M-A, Morales GJ, Chaves S, McGrane J. Epidemiology of urban canine rabies, Santa Cruz, Bolivia, 1972-1997. Emerging infectious diseases. 2002;8: 458–461.

8. Vigilato MAN, Clavijo A, Knobl T, Silva HMT, Cosivi O, Schneider MC, et al. Progress towards eliminating canine rabies: policies and perspectives from Latin America and the Caribbean. Philosophical transactions of the Royal Society of London Series B, Biological sciences. 2013;368: 20120143.

9. Fooks AR, Banyard AC, Horton DL, Johnson N, McElhinney LM, Jackson AC. Current status of rabies and prospects for elimination. The Lancet. 2014;384: 1389–1399.

10. Hampson K, Coudeville L, Lembo T, Sambo M, Kieffer A, Attlan M, et al. Estimating the global burden of endemic canine rabies. PLoS neglected tropical diseases. 2015;9: e0003709.

11. World Health Organisation. WHO Expert Consultation on Rabies. Second report. World Health Organization technical report series. 2013;: 1–139– back cover.

498   12.   Franka R, Smith TG, Dyer JL, Wu X, Niezgoda M, Rupprecht CE. Current and
499   future tools for global canine rabies elimination. Antiviral research. 2013;100:
500   220–225.

501   13.   Lackay SN, Kuang Y, Fu ZF. Rabies in small animals. The Veterinary clinics of
502   North America Small animal practice. 2008;38: 851–859.

503   14.   Clavijo A, Del Rio Vilas VJ, Mayen FL, Yadon ZE, Beloto AJ, Vigilato MAN, et
504   al. Gains and future road map for the elimination of dog-transmitted rabies in the
505   Americas. The American journal of tropical medicine and hygiene. 2013;89:
506   1040–1042.

507   15.   Vigilato MAN, Cosivi O, Knobl T, Clavijo A, Silva HMT. Rabies Update for
508   Latin America and the Caribbean. Emerging infectious diseases. 2013;19: 678–
509   679.

510   16.   Davlin SL, Vonville HM. Canine rabies vaccination and domestic dog population
511   characteristics in the developing world: a systematic review. Vaccine. 2012;30:
512   3492–3502.

513   17.   Cleaton JM, Wallace RM, Crowdis K, Gibson A, Monroe B, Ludder F, et al.
514   Impact of community-delivered SMS alerts on dog-owner participation during a
515   mass rabies vaccination campaign, Haiti 2017. Vaccine. 2018;36: 2321–2325.

516   18.   Mazeri S, Gibson AD, Meunier N, Bronsvoort BMD, Handel IG, Mellanby RJ, et
517   al. Barriers of attendance to dog rabies static point vaccination clinics in Blantyre,
518   Malawi. PLoS neglected tropical diseases. 2018;12: e0006159.

519   19.   Cleaveland S, Kaare M, Tiringa P, Mlengeya T, Barrat J. A dog rabies vaccination
520   campaign in rural Africa: impact on the incidence of dog rabies and human dog-
521   bite injuries. Vaccine. 2003;21: 1965–1973.

522   20.   Jibat T, Hogeveen H, Mourits MCM. Review on dog rabies vaccination coverage
523   in Africa: a question of dog accessibility or cost recovery? PLoS neglected tropical
524   diseases. 2015;9: e0003447.

525   21.   Gsell AS, Knobel DL, Kazwala RR, Vounatsou P, Zinsstag J. Domestic dog
526   demographic structure and dynamics relevant to rabies control planning in urban
527   areas in Africa: the case of Iringa, Tanzania. BMC veterinary research. 2012;8:
528   236.

529   22.   Kaare M, Lembo T, Hampson K, Ernest E, Estes A, Mentzel C, et al. Rabies
530   control in rural Africa: Evaluating strategies for effective domestic dog
531   vaccination. Vaccine. 2009;27: 152–160.

532   23.   Muthiani Y, Traoré A, Mauti S, Zinsstag J, Hattendorf J. Low coverage of central
533   point vaccination against dog rabies in Bamako, Mali. Preventive veterinary
534   medicine. 2015;120: 203–209.

535   24.   Perry BD, Kyendo TM, Mbugua SW, Price IE, Varma S. Increasing rabies
536   vaccination coverage in urban dog populations of high human population density
537   suburbs: a case study in Nairobi, Kenya. Preventive veterinary medicine. 1995;22:
538   137–142.

539   25.   Schneider MC, Belotto A, Adé MP, Hendrickx S, Leanes LF, Rodrigues MJ de F,
540   et al. Current status of human rabies transmitted by dogs in Latin America.
541   Cadernos de saúde pública. 2007;23: 2049–2063.

542   26.   Castillo-Neyra R, Zegarra E, Monroy Y, Bernedo R, Cornejo-Rosello I, Paz-
543   Soldan V, et al. Spatial Association of Canine Rabies Outbreak and Ecological
544   Urban Corridors, Arequipa, Peru. Tropical Medicine and Infectious Disease.
545   2017;2: 38.

546   27.   Burgos-Cáceres S. Canine Rabies: A Looming Threat to Public Health. Animals :
547   an open access journal from MDPI. 2011;1: 326–342.

548   28.   World Health Organization. WHO Expert Consultation on Rabies. Geneva; 2005.

549   29.   Zhang L, Li X, Yang H, Mao R, Zhao Q. Health status and health-seeking
550   behaviour between interprovincial and intraprovincial rural-to-urban young
551   migrants in Nanjing China. Asia-Pacific Population Journal. 2011;26: 39–54.

552   30.   Li Y. Understanding health constraints among rural-to-urban migrants in China.
553   Qualitative health research. 2013;23: 1459–1469.

554   31.   Benyoussef A, Cutler JL, Levine A, Mansourian P, Phan-Tan T, Baylet R, et al.
555   Health effects of rural-urban migration in developing countries--Senegal. Social
556   science & medicine (1982). 1974;8: 243–262.

557   32.   Carrillo-Larco RM, Bernabé-Ortiz A, Pillay TD, Gilman RH, Sanchez JF, Poterico
558   JA, et al. Obesity risk in rural, urban and rural-to-urban migrants: prospective
559   results of the PERU MIGRANT study. International journal of obesity (2005).
560   2015.

561   33.   Levy MZ, Barbu CM, Castillo-Neyra R, Quispe Machaca VR, Ancca Juárez J,
562   Escalante-Mejia P, et al. Urbanization, land tenure security and vector-borne
563   Chagas disease. Proceedings Biological sciences / The Royal Society. 2014;281:
564   20141003.

565   34.   Baddeley A, Turner R. spatstat: An RPackage for Analyzing Spatial Point Patterns.
566   Journal of Statistical Software. 2005;12.

567   35.   Team. R: A Language and Environment for Statistical Computing. 2013;: 1–3604.

568   36.   Durr S, Mindekem R, Kaninga Y, Doumagoum Moto D, Meltzer MI, Vounatsou
569   P, et al. Effectiveness of dog rabies vaccination programmes: comparison of

owner-charged and free vaccination campaigns. Epidemiology and Infection.
2009;137: 1558.

37. Townsend SE, Sumantra IP, Pudjiatmoko, Bagus GN, Brum E, Cleaveland S, et al.
Designing programs for eliminating canine rabies from islands: Bali, Indonesia as
a case study. PLoS neglected tropical diseases. 2013;7: e2372.

38. Widyastuti MDW, Bardosh KL, Sunandar   , Basri C, Basuno E, Jatikusumah A,
et al. On dogs, people, and a rabies epidemic: results from a sociocultural study in
Bali, Indonesia. Infectious Diseases of Poverty. 2015;4: 60.

39. Bardosh K, Sambo M, Sikana L, Hampson K, Welburn SC. Eliminating rabies in
Tanzania? Local understandings and responses to mass dog vaccination in
Kilombero and Ulanga districts. PLoS neglected tropical diseases. 2014;8: e2935.

40. Serebe SG, Tadesse KA. Study on community knowledge, attitude and practice of
rabies in and nearby Gondar town, North West Ethiopia. Journal of Public ….
2014.

41. Kabeta T, Deresa B, Tigre W, Ward MP, Mor SM. Knowledge, Attitudes and
Practices of Animal Bite Victims Attending an Anti-rabies Health Center in Jimma
Town, Ethiopia. PLoS neglected tropical diseases. 2015;9: e0003867.

42. De Ramos M, Bravo LC. Knowledge, attitudes, and practices of community
regarding animals bite and rabies. Paediatric Infectious Disease Society of the ….
2004.

43. Singh US, Choudhary SK. Knowledge, attitude, behavior and practice study on
dog-bites and its management in the context of prevention of rabies in a rural
community of Gujarat. Indian Journal of Community Medicine. 2005.

44. Kayali U, Mindekem R, Yemadji N. Coverage of pilot parenteral vaccination
campaign against canine rabies in N'Djamena, Chad. Bulletin of the World ….
2003.

45. Minyoo AB, Steinmetz M, Czupryna A. Incentives Increase Participation in Mass
Dog Rabies Vaccination Clinics and Methods of Coverage Estimation Are
Assessed to Be Accurate. PLoS Negl Trop …. 2015.

46. Lembo T, Hampson K, Haydon DT, Craft M, Dobson A, Dushoff J, et al.
Exploring reservoir dynamics: a case study of rabies in the Serengeti ecosystem.
The Journal of Applied Ecology. 2008;45: 1246–1257.

47. Lapiz SMD, Miranda MEG, Garcia RG, Daguro LI, Paman MD, Madrinan FP, et
al. Implementation of an intersectoral program to eliminate human and canine
rabies: the Bohol Rabies Prevention and Elimination Project. PLoS neglected
tropical diseases. 2012;6: e1891.

48.   Ferguson EA, Hampson K, Cleaveland S, Consunji R, Deray R, Friar J, et al.
607   Heterogeneity in the spread and control of infectious disease: consequences for the
608   elimination of canine rabies. Scientific reports. 2015;5: 18232.

609   49.   Jacobs B, Ir P, Bigdeli M, Annear PL, Van Damme W. Addressing access barriers
610   to health services: an analytical framework for selecting appropriate interventions
611   in low-income Asian countries. Health Policy and Planning. 2012;27: 288–300.

612   50.   Bissonnette L, Wilson K, Bell S, Shah TI. Neighbourhoods and potential access to
613   health care: The role of spatial and aspatial factors. Health & Place. 2012;18: 841–
614   853.

615   51.   Hiscock R, Pearce J, Blakely T, Witten K. Is Neighborhood Access to Health Care
616   Provision Associated with Individual-Level Utilization and Satisfaction? Health
617   Services Research. 2008;43: 2183–2200.

618   52.   Cleaveland S. The epidemiology of rabies and canine distemper in the Serengeti,
619   Tanzania.

620   53.   Zhang L, Li X, Yang H, Mao R, Zhao Q. Health status and health-seeking
621   behaviour between interprovincial and intraprovincial rural-to-urban young
622   migrants in Nanjing China. Asia-Pacific Population Journal. 2011;26: 39–54.

623   54.   Phillips DR. Spatial variations in attendance at general practitioner services. Social
624   Science & Medicine. 1979;13D: 169–181.

625   55.   Sanz B, Regidor E, Galindo S, Pascual C, Lostao L, Díaz JM, et al. Pattern of
626   health services use by immigrants from different regions of the world residing in
627   Spain. International Journal of Public Health. SP Birkhäuser Verlag Basel;
628   2011;56: 567–576. doi:10.1007/s00038-011-0237-9

629   56.   Benyoussef A, Cutler JL, Levine A, Mansourian P, Phan-Tan T, Baylet R, et al.
630   Health effects of rural-urban migration in developing countries--Senegal. Social
631   science & medicine (1982). 1974;8: 243–262.

632   57.   Carrillo-Larco RM, Bernabé-Ortiz A, Pillay TD, Gilman RH, Sanchez JF, Poterico
633   JA, et al. Obesity risk in rural, urban and rural-to-urban migrants: prospective
634   results of the PERU MIGRANT study. International journal of obesity (2005).
635   2015.

636   58.   Berry JW. Acculturation and Adaptation in a New Society. International
637   Migration. John Wiley & Sons, Ltd (10.1111); 1992;30: 69–85.
638   doi:10.1111/j.1468-2435.1992.tb00776.x

639   59.   Hull D. Migration, adaptation, and illness: a review. Soc Sci Med Med Psychol
640   Med Sociol. 1979;13A: 25–36.

60.   Fu Q, Han X-M, Wang L-Y, Zhaxi S-M, Ma X, Wang Y-S, et al. Effect of control
pattern with emphasis on canine deworming in a pilot of echinococcosis in highly
endemic area, southern Qinghai Plateau. Zhongguo ji sheng chong xue yu ji sheng
chong bing za zhi = Chinese journal of parasitology & parasitic diseases. 2011;29:
293–295.

61.   Loza A, Talaga A, Herbas G, Canaviri RJ, Cahuasiri T, Luck L, et al. Systemic
insecticide treatment of the canine reservoir of Trypanosoma cruzi induces high
levels of lethality in Triatoma infestans, a principal vector of Chagas disease.
Parasites & Vectors. 2017;10: 344.

62.   Aparicio-Burgos JE, Zepeda-Escobar JA, de Oca-Jiménez RM, Estrada-Franco JG,
Barbabosa-Pliego A, Ochoa-García L, et al. Immune protection against
Trypanosoma cruzi induced by TcVac4 in a canine model. PLoS neglected tropical
diseases. 2015;9: e0003625.

Effect of distance from dog owner's house to the closest vaccination point on clustering of unvaccinated dogs

Distance from Vaccination Point in Meters