Socio-spatial heterogeneity in participation in mass dog rabies vaccination campaigns, Arequipa, Peru

Ricardo Castillo-Neyra[1,2]*, Amparo M. Toledo[2], Claudia Arevalo-Nieto[2], Hannelore MacDonald[3], Micaela De la Puente-León[2], Cesar Naquira-Velarde[2], Valerie A. Paz-Soldan[2,4], Alison M. Buttenheim[2,5], Michael Z. Levy[1,2]

[1] Department of Biostatistics, Epidemiology and Informatics, Perelman School of Medicine, University of Pennsylvania Perelman School of Medicine, Philadelphia, Pennsylvania, USA
[2] Zoonotic Disease Research Lab, One Health Unit, School of Public Health and Administration, Universidad Peruana Cayetano Heredia, Arequipa, Peru
[3] Department of Biology, School of Arts and Sciences, University of Pennsylvania, Philadelphia, Pennsylvania, USA
[4] Global Community Health and Behavioral Sciences, Tulane School of Public Health and Tropical Medicine, New Orleans, Louisiana, USA
[5] Department of Family and Community Health, School of Nursing, University of Pennsylvania, Philadelphia, Pennsylvania, USA

* Corresponding author

E-mail: cricardo@pennmedicine.upenn.edu

Short title: Socio-spatial heterogeneity in mass dog vaccination campaign

Keywords:

Community participation, Disease outbreaks, Dog diseases, Immunization programs, Mass vaccination, One Health, Rabies, Vaccination coverage, Zoonoses.

Abstract

To control and prevent rabies in Latin America, mass dog vaccination campaigns (MDVC) are implemented mainly through fixed-location vaccination points: owners have to bring their dogs to the vaccination points where they receive the vaccination free of charge. Dog rabies is still endemic in some Latin-American countries and high overall dog vaccination coverage and coverage evenness are desired attributes of MDVC to halt rabies virus transmission. In Arequipa, Peru, we conducted a door-to-door post-campaign survey on >6,000 houses to assess the placement of vaccination points on these two attributes. We found that the odds of participating in the campaign decreased by 16% for every 100 m from the owner's house to the nearest vaccination point (p=0.041) after controlling for potential covariates. We found social determinants associated with participating in the MDVC: for each child under 5 in the household, the odds of participating in the MDVC decreased by 13% (p=0.032), and for each decade less lived in the area, the odds of participating in the MDVC decreased by 8% (p<0.001), after controlling for distance and other covariates. We also found significant spatial clustering of unvaccinated dogs over 500 m from the vaccination points, which created pockets of unvaccinated dogs that may sustain rabies virus transmission. Understanding the barriers to dog owners' participation in community-based dog-vaccination programs will be crucial to implementing effective zoonotic disease preventive activities. Spatial and social elements of urbanization play an important role in coverage of MDVCs and should be considered during their planning and evaluation.

Author summary

In Peru and other dog rabies-affected countries, mass dog vaccination campaigns (MDVC) are implemented primarily through fixed-location vaccination points: owners have to bring their dogs to the vaccination points where they receive the vaccination. To stop rabies virus transmission, a high and even dog vaccination coverage is desired. In Arequipa, Peru, following a MDVC, we conducted a door-to-door survey of >6,000 houses to assess how the placement of vaccination points affected coverage of the campaign. When comparing dog owners with similar characteristics, we found that the odds of participating in the MDVC was reduced by 16% for every 100 m distance from the nearest vaccination point. Some social conditions were also associated with participating in the MDVC: for each child under 5 in the household, odds of participating in the MDVC decreased by 13%, and for each decade less lived in the area, the odds of participating in the MDVC decreased by 8%. Distance to the vaccination point and variation in social conditions across the city play important roles in achieving coverage of MDVCs and should be considered during campaign planning and evaluation.

1. Introduction

The city of Arequipa is in the midst of a sustained dog rabies outbreak. The introduction of rabies virus into the city has been ascribed to the unintentional transport of rabid dogs from the rabies-endemic state of Puno during human migration [1-3], and the persistence of transmission is likely due to low coverage in the annual city-wide dog vaccination campaigns [3]. Following the detection of the outbreak in Arequipa city in 2015, the

Ministry of Health of Peru (MOH) initiated additional vaccination campaigns in the city

with varying intensity [4]. These additional efforts have not quelled the epidemic: more

than 160 rabid dogs have been detected as of 2019.

Epidemics of dog rabies are ongoing in major urban centers across Latin America and

worldwide [1,5-8]. Since bites from rabies-infected dogs cause 99% of human rabies

deaths [9], the control and elimination of dog-mediated human rabies relies on a

population-wide animal-centered strategy: mass dog rabies vaccination [8,10,11]. Dog

vaccination has dramatically decreased the global burden of human rabies since 1955

[5,12,13]; in the Americas, national programs centered around mass dog vaccination have

achieved enormous advances [8,14,15], reducing the incidence of dog rabies by 98%

since 1983 [14]. In most rabies-affected countries, government health entities (e.g. MOH)

organize annual MDVC that are held in outdoor settings. These campaigns are usually

free of charge and voluntary [8,16] and campaign promotion varies greatly in format,

content and intensity [1,17,18].

There are three non-mutually exclusive strategies used in Peru to implement MDVC:

fixed vaccination posts, mobile teams setting up a temporary mobile post or conducting

'street vaccination', and door-to-door vaccination [19]. For the fixed-point strategy, the

vaccinators wait for the dog owners to bring the dogs to a unique place. For the mobile

team strategy, the vaccinators move from one location to another during the day,

vaccinating dogs along their way and spending short periods (i.e. under an hour) in a

location before moving on, waiting for the dog owners to bring the dogs to these moving

locations. For the door-to-door strategy, vaccinators knock on doors asking to vaccinate

dogs in the household. Locations of the fixed vaccination sites are typically determined

by a combination of convenience and prominence of the location (e.g. the entrance to a

health post, a well-known park) [11]. In Peru, routes for door-to-door and mobile team

approaches may or may not be decided in advance, and teams may move during the

course of the day looking for dogs along their routes.

The fixed-point strategy has been extensively used in Latin America and Africa, even

though it has frequently failed to attain coverage targets [1,8,20-24]. The main reasons

for its extensive application are that fixed-point vaccination is easier to implement and

less costly than other strategies [18]. In many cases fixed-point is combined with other

strategies, particularly when initial activities are unsuccessful [19,23,25]. However, high

dog owner participation in MDVC and other dog-centered health campaigns (e.g. de-

worming dogs to prevent human hydatid disease) has proven difficult to achieve in many

areas [24,25]. It is necessary to understand barriers to community-based control strategies

targeting dog populations in dog rabies-affected countries where coverage does not reach

the minimum 70% recommended by the World Health Organization to attain herd

immunity [11], much less the 80% recommended by the Pan-American Health

Organization for the region [26].

In cities, the social and spatial aspects of urbanization can facilitate the emergence of dog

rabies and complicate its control [27,28]. In Arequipa, the locations of rabid dogs have

been associated with urban structures [27], and dog owners from areas with different

levels of urbanization have reported distinct correlates of vaccinating their dogs against

rabies [1]. The changing urban landscape and social processes in rapidly-growing cities

have been associated with uptake of health-related services [29-32] and may be related to

the low dog vaccination coverage in Arequipa. The objectives of the present study were

to quantitatively assess the association between distance to a vaccination point and dog

owner participation in mass dog vaccination campaigns in an urban setting, and to

evaluate the effect of such distance on overall vaccination coverage and evenness.

2. Methods

2.1. Ethics statement

128 Ethical approval was obtained from Universidad Peruana Cayetano Heredia (approval

129 number: 65369), Tulane University (approval number: 14–606720), and University of

130 Pennsylvania (approval number: 823736). All human subjects in this study were adults.

2.2. Study setting

The study was conducted in Alto Selva Alegre (ASA) (human population for 2015:

82,412; density: 11,902 people/km^2), one of the 14 districts of the city of Arequipa.

Arequipa, Peru's second largest city, is home to 969,000 people and is situated at ~2,300

meters above sea level. The first detection of a rabid dog in the city of Arequipa occurred

in March 2015 in ASA. By June 2016, when our data were collected, 43 rabid dogs had

been detected in 8 districts, but it is assumed that number represents a small fraction of

the total number of cases [27]. The city of Arequipa comprises communities spanning

different stages of urbanization and different migration histories, from old established

neighborhoods, to young neighborhoods, to recent invasions [33]. Within this gradient of

development, young neighborhoods and recent invasions are often located on the

periphery of the city (peri-urban area) and the older localities are nearer to the center

(urban area) [33]. Compared to the urban area, peri-urban areas generally have lower

socioeconomic status, fewer community resources, more security problems, and often

more rugged and uneven terrain (Figure 1). As new neighborhoods mature into

established neighborhoods with wealthier residents, homes are improved with better

quality construction material and permanent utility connections, and connectivity with the

rest of the city increases with better sidewalks, roads, and transportation access. ASA

transects the city, running from the center to the periphery, and the district continues to

grow towards the outskirts of the city. In our study, participants represented either the

urban or peri-urban areas of the city of Arequipa. We included 21 urban neighborhoods

founded many decades ago, and 9 peri-urban neighborhoods that originated around 2000

or later.

In ASA, the MOH conducted a mass dog vaccination campaign in June 2016. A detailed

description of the mass dog vaccination campaigns can be found elsewhere [1], and the

dog population estimator used in Arequipa can be found in [27].

159
160
161 **Figure 1**. Study communities display landscape heterogeneity. A: Urban area. B: Peri-
162 urban.
163

2.3. Data collection

In collaboration with the MOH, we georeferenced every stationary and mobile

vaccination team during the three weekends when the campaign was implemented in

ASA. Due to volunteer assistance, some dogs were vaccinated in weekdays during the

campaign; we did not georeference the location of those volunteering vaccinators, mainly

because their schedule was haphazard and unpredictable. We started door-to-door

surveys immediately after the vaccination campaign, visiting every household in the

study area and consenting and surveying only one adult (≥18) per household.

The door-to-door survey was designed based on the rabies literature and based on our

qualitative studies of local communities, in which we found specific household-level

barriers to vaccination [1]. We collected household variables (e.g. number of household

members; number of children under 5 years old), dog owner or interviewee variables (e.g.

gender; educational attainment), and dog variables (e.g. vaccination status; age). All

houses in the study localities were geocoded and the survey data were linked to the

household coordinates. We estimated the Euclidean distance between households and the

closest vaccination point (fixed, mobile, or either).

2.4. Statistical Analysis

We estimated the total vaccination coverage in the study area and compared the coverage

in urban vs. peri-urban localities with a chi-squared test. We estimated the human to dog

ratio and bootstrapped it 10,000 times to estimate its confidence interval. To evaluate the

baseline characteristics of households and dog owners by participation in the MDVC, we

defined an ordinal outcome for houses with dogs: *no participation* (no dog vaccinated in

the house), *partial participation* (some, but not all dogs in the house, vaccinated), and *full*

participation in the MDVC (all dogs in the house vaccinated). We used a chi-square test

to compare categorical variables with 10 or more observations per group, Fisher's exact

test for categorical variables with fewer than 10 observations in any subgroup, and

Mann–Whitney U test for age of the dog owner or interviewee, which did not follow a

normal distribution and was truncated at 18 years. We compared the individual

characteristics of vaccinated and unvaccinated dogs with chi-square for categorical

variables and with Mann–Whitney U test for dog's age.

Our main objective was to assess the association between distance to the vaccination

point and participation in the MDVC. For distance to the vaccination point, we used the

Euclidean distance from the dog's house to the closest vaccination point, either fixed or

mobile. For participation in the MDVC we used the ordinal values described above: *no*

participation, *partial participation*, and *full participation.* We compared proportional

odds logistic regression (POLR), non-proportional odds logistic regression, and

multinomial regression. The ordinal models were superior to the multinomial regression,

and the proportional odds assumption holds for most of the covariates. Given that the

categories of participation are inherently ordinal and that providing a single point

estimate per covariate is more interpretable, the POLR model was favored. Based on the

recent literature and our local studies [1,34,35], the following covariates were used for

model building: having a dog leash at home, number of children under 5 years old at

home, time living in the area, rabies status of the last place they lived in before living in

the study area, number of dogs at the house, age and gender of the dog owner, and

educational attainment. We considered transformed distance to capture non-linear effects

and interactions between distance and having a leash. The fit of the alternative models to

the data was compared with Akaike's Information Criteria (AIC). We also attempted to

build a hierarchical model to take into account the spatial autocorrelation within locality.

However, given that within each locality there was at most one vaccination point, the

variable distance from the house to the vaccination point would be unidentifiable under

such hierarchical model. The final POLR model fitted with the R package MASS [36]

was:

$$ln\left\{\frac{Odds\,(Y \le k)}{Odds\,(Y > k)}\right\} = b_0 + b_1.\,distance + b_2.\,leash + b_3.\,children + b_4.\,\substack{recency\ in \\ the\ area} + b_5.\,\substack{rabies\ status\ of \\ previous\ residence} + \varepsilon$$

where k takes the values 0 (no participation), 1 (partial participation), and 2 (full

participation). All statistical tests were 2-sided, and significance level was 0.05.

We tested the spatial pattern of vaccinated and unvaccinated dogs in relation to

vaccination tents for clustering using the bivariate cross K-function. This function

estimates spatial dependence between two types of points (i.e. unvaccinated dogs and

vaccination points) by measuring the expected number of points of type i within a given

distance to a point of type j divided by the overall density of the points of type i. We used

the Kcross function in the R package spatstat [37] to estimate deviations between the K

function estimated for our data and the theoretical K function corresponding to a

completely random Poisson point process for vaccinated dogs to tents and unvaccinated

dogs to tents. Deviations greater than the theoretical K function indicate that the mean

point count is higher than expected under complete spatial randomness (CSR) and thus

some degree of clustering is present between the two event types at the indicated

distance. Similarly, deviations less than the theoretical K function indicate that the mean

point count is lower than expected under CSR which therefore indicates that some degree

of dispersion is present between the two event types at the indicated distance.

In order to investigate the association between geolocation and the odds of canine

vaccination we fitted Generalized Additive Models (GAMs) to our data using the R

package MapGAM [38]. GAMs are an extension of linear regression models in which

both parametric and non-parametric terms are used to estimate the outcome of interest.

We used a two-dimensional locally weighted smooth (LOESS) of latitude and longitude

for our non-parametric term. The LOESS smoother fits each data point by weighting it

towards nearby points, where weighting is based on the distance to the point being fitted.

The percentage of data points in the region that will be used to predict a particular point

is referred to as the span. The optimal span size used for smoothing was determined by

minimizing the AIC. We mapped the odds ratio for each point on pre-specified grids of

each locality (from polygon data) and next tested the null hypothesis that the odds of each

points' vaccination status did not depend on geolocation using permutation tests. For

each test, the paired latitude and longitude coordinates were randomly permuted but

vaccination status was held fixed. 1000 permutations were run for each locality and

contour lines encircle areas with significantly increased or decreased vaccination odds as

indicated by point wise p-values computed from the permutation ranks. All models and

figures were created with R [39].

3. Results

Based on our survey, the estimated vaccination coverage of the MOH MDVC was only

58.1%, and it was low in both urban and peri-urban localities (58.0% vs. 58.6%

respectively, chi2=0.086, p=0.769). Only 3.4% of dogs were (reportedly) vaccinated in

private clinics, bringing our estimated total coverage to 61.5%. Participation in our

survey was higher in the urban area (88.8%) compared to the peri-urban area (61.6%)

(mean= 82.0%, chi2=6458.5, p<0.001). The total number of dogs in the surveyed houses

was 5,292 and the human to dog ratio was 3.78:1 (95% CI: 3.69:1 - 3.89:1). In total,

65.3% of surveyed houses had dogs, but this number was higher in peri-urban areas

(70.0% compared to 64.6% in urban areas, chi2=6.529, p=0.011). For 76.9% of

vaccinated dogs in the area, the person who took them to the MDVC is the person who

responded to the survey. In our study area the urbanization process involves new

localities being founded and settlers moving in. Accordingly, we found that the time of

residency (or the year people moved into this area) was clustered at the locality level.

However, we found that people living in or founding a locality do not necessarily share

the same place of origin or previous residence.

When we compared the distance from each house to the closest fixed vaccination point, to the closest mobile vaccination point, and to either the closest fixed or mobile vaccination point, we found a clear gradient in distance from vaccination point from non-participant houses (farther) to houses that partially participated (closer) to houses that fully participated (closest) (Table 1). The proportion of households with children under 5 was higher in households that did not participate in the campaign (36%) compared to houses that participated fully or partially in the campaign (31% and 29% respectively). The proportion of houses with a dog leash increases from those who did not participate, to those who participated partially, to those that fully participated of the MDVC. There were some differences in MDVC participation by migration history: people who have lived longer in ASA tend to report higher participation in the MDVC compared to those who have lived fewer years in ASA. Also, there is a slight difference in participation in the MDVC depending on rabies status of previous residence, with more people participating in the campaign, partially or fully, if they were migrants from a rabies endemic area (Table 1). Other variables, such as educational attainment, the proportion of female dog owners or interviewees, and the proportion of households in urban localities were all similar in those households that participated fully or partially in the campaign compared to those households that did not participate of the campaign (Table 1). Having multiple dogs is a prerequisite to be in the partial participation group; therefore, houses with partial participation had on average more dogs, but the number of dogs per house was very similar in houses that participated fully compared to those that did not participate in the campaign (Table 1). Those who did not participate in the MDVC

292   reported more frequently not knowing about the campaign before it happened (Table 1),

293   but many of them reported learning about the campaign the same day it occurred. In

294   S1_Table, we report the media channels through which they learned about the campaign

295   either after it occurred or the same day it occurred.

297   Table 1. Characteristics of the dog owner population in the study area by participation in
298   the MDVC, Arequipa City, Peru, 2016.

299

Interviewee/Owner Variables	Vaccinated no dog (n=1,017)	Vaccinated some dogs (n=176)	Vaccinated all dogs (n=1,434)	p
Distance to any closest vaccination point – Mean (SD)	125m (96)	117m (114)	103m (83)	<0.001 [a]
Distance to closest fixed vaccination point – Mean (SD)	501m (479)	490m (420)	461m (422)	0.119 [a]
Distance to closest mobile vaccination point – Mean (SD)	137m (101)	127m (119)	114m (93)	<0.001 [a]
Number of dogs in houses with dogs – Mean (SD)	1.79 (1.10)	2.96 (1.24)	1.80 (1.05)	<0.001 [a]
Houses with children under 5	36.4%	28.9%	30.8%	0.019 [b]
Age of dog owner/interviewee – Median (IQR)	39 years (28 - 52)	40 years (29 - 50)	42 years (30 - 53)	0.024 [c]
Time living in ASA if migrant – Mean (SD)	18.1 years (13.7)	21.2 years (13.9)	22.1 years (14.8)	<0.001 [a]
Area before living in ASA				0.036 [b]
Rabies epidemic (Arequipa)	90.7%	94.1%	89.8%	
Rabies endemic (Puno)	1.6%	2.4%	3.3%	
Rabies free (rest of country)	7.7%	3.6%	6.9%	
Educational attainment				0.509 [b]
Illiterate	0.6%	0.5%	0.6%	
Primary school	11.5%	11.4%	13.6%	
High school	43.2%	48.3%	41.1%	
Technical academy	21.2%	15.3%	20.6%	
University	22.3%	23.3%	23.3%	
Preferred not to respond	1.2%	1.1%	0.7%	
Owns at least one dog leash	51.0%	54.5%	60.1%	<0.001 [b]
Female interviewees/owners	64.6%	73.3%	64.4%	0.032 [b]
Did not know about the campaign before it happened	15.4%	2.3%	1.3%	<0.001 [b]
Live in urban locality (vs. peri-urban)	78.8%	78.1%	79.4%	0.002 [b]

300   *p* values estimated with [a] one-way ANOVA; [b] Chi square test; and [c] Kruskal Wallis test.

301

302   Compared to vaccinated dogs, unvaccinated dogs were older, were more likely female,

303   had more free access to the street, had owners with no leash for them at home, and were

304   less likely to be walked. Multipurpose dogs (dogs reported as guard and company dogs)

were more likely to be vaccinated (Table 2). Based on the stated source of the dog, those

received as gifts are more likely to be vaccinated and dogs born at home or

adopted/picked on the street are less likely to be vaccinated (Table 2). Being considered

purebred or being spayed/neutered was not associated with dog vaccination status (Table

2).

Table 2. Dog characteristics by vaccination status in the MDVC (dogs vaccinated
privately not included).
312

Dog Variables	Unvaccinated dog (n=2,019)	Vaccinated dog (n=2,800)	p
Age of dogs – Mean (SD)	24 months (12 – 60)	20 months (10 – 54)	0.039 [a]
Female dogs	42.1%	36.8%	<0.001 [b]
Free access to the street	20.6%	25.7%	0.002 [b]
Leash for the dog	33.8%	45.3%	<0.001 [b]
Dogs that are walked, either with or without leash.	55.9%	66.9%	<0.001 [b]
Dog function			<0.001 [b]
Company	52.0%	49.0%	
Protection	23.4%	20.9%	
Company & Protection	24.6%	29.6%	
Breeder/business	0.0%	0.5%	
Dog source			<0.001 [b]
Gift	51.2%	56.5%	
Bought at market/store	17.4%	18.3%	
Born at home	15.6%	12.3%	
Adopted/Picked on the street	12.3%	10.8%	
Bought from friend/neighbor	2.6%	2.0%	
Do not know	1.0%	0.0%	
Purebred as reported by interviewee	25.7%	24.9%	0.589 [b]
Spayed/Neutered	3.5%	2.6%	0.157 [b]

p values estimated with [a] Student's t-test; and [b] Chi square test.

In the multivariable regression analysis, we found that distance to the vaccination site is

strongly associated with participation of the MDVC. The odds of not participating in the

MDVC were 16% lower for someone who lived 100 meters farther from the vaccination

point after adjusting for other covariates and this difference was statistically significant

(Table 3). Those having a dog leash at home had 35% higher odds of participation of the

MDVC, either fully or partially, compared to those who did not have a dog leash after

adjusting for other covariates. The odds of participating in the MDVC, either fully or

partially, were 13% lower for each additional child under 5 years at home, after adjusting

for other covariates. Migration history was associated with participating in the MDVC;

participation was lower in those who migrated more recently to the study area (8% lower

odds of participation for each decade less lived in the area, after adjusting for other

covariates). Another component of migration history, the previous residence region, was

also associated with participating in the MDVC: those whose previous residence was a

rabies-free region or was any district within Arequipa were 23% to 32% less likely to

participate of the campaign compared to those whose previous residence was a rabies-

endemic region, after adjusting for other covariates (Table 3). Demographic variables

such as owner's/interviewee's age, gender and educational attainment, and other

household-level variables, such as number of dogs at the house, were dropped during

model selection because they neither improved the fit of the model nor were statistically

associated with participating in the MDVC.

Table 3. Factors associated with participating in the Mass Dog Vaccination Campaign,
Arequipa City, Peru, 2016.
338

Variable	OR	95% CI	p
Distance to closest vaccination point (100m)	0.84	0.76 – 0.98	0.041
Having a leash at home	1.35	1.10 – 1.61	0.003
Number of children U5Y at home	0.87	0.78 – 0.97	0.032
Recency in ASA (1 fewer decade)	0.92	0.90 – 0.94	<0.001
Rabies status of previous residence			
Endemic for rabies	Ref.		
Epidemic for rabies	0.77	0.63 – 0.93	0.006
Free of rabies	0.68	0.57 – 0.82	<0.001

Odds ratios estimated with multiple proportional odds logistic regressions.

The significant association between distance to the vaccination point and odds of

participating in the MDVC has consequences for the distribution of unvaccinated dogs in

the area. We observed spatial clustering of unvaccinated owned dogs as a function of the

distance from the house to the vaccination point. These pockets of unvaccinated dogs

closer to each other than expected by chance occur at 500 meters from the vaccination

point or further (Figure 2). We also analyzed the spatial odds of participating in the

MDVC, that is, the association between their specific geolocation and the vaccination

point. For areas served by fixed-point vaccination, there was a clear smooth spatial effect

with higher odds of participating for houses closer to the vaccination point and a

decreasing gradient farther away from the vaccination point. The spatial effect of the

fixed-point vaccination strategy creates two clearly defined zones: a large zone with

statistically significantly high odds of participation in the MDVC and another large zone

with statistically significantly low odds of participation in the MDVC (Figure 3). For

areas served by mobile teams, there were more spots of significant low and high odds of

participation in the MDVC and these spots were spread in the study area without a clear

association between the spots and the locations where the vaccination teams stopped to

wait for dogs or to vaccinate dogs (Figure 3).

Figure 2. Clustering of unvaccinated dogs as a function of distance from the vaccination
361   point.
362

Figure 3. Spatial odds ratios for participating in the MDVC in a locality served by a fixed vaccination point (cross in A), and by a mobile team (in B, X's represent locations where the mobile team stopped to wait for dogs or to vaccinate dogs). Created with MapGAM package [38] in R [39].

4. Discussion

In Arequipa Peru, the social and spatial aspects of urbanization facilitate the emergence of dog rabies and complicate its control. In 21 urban and 9 peri-urban localities in Arequipa, Peru, we found low vaccination coverage and coverage that was spatially uneven. We found a strong effect of a potential proximal determinant of participation in the MDVC: distance to the vaccination point. The unadjusted data show a clear negative gradient with higher levels of participation in the MDVC at shorter distances to the vaccination point. After accounting for other important individual- and household-level

variables, distance to the vaccination point remains an important factor associated with

participating in the campaign. The association between distance to and participation in

the MDVC also impacts the spatial evenness of vaccination. We found areas with

statistically significant lower odds of dogs being vaccinated, and the LOESS smoother

maps correlated well with maps of vaccination coverage. Therefore, it seems that both

fixed point and mobile team canine vaccination approaches produced spatially

heterogeneous vaccination coverage. However, we found that vaccination coverage was

more "patchy" in localities served by mobile vaccination teams. This combination of

mobile and fixed points was used also in 2015, but the same localities are not always

served with the same approach (e.g. a locality that was served with mobile teams in 2015

could be served with fixed point vaccination in 2016). As others have reported [40], there

is potential that in some localities owners in 2016 expected that the MDVC would be

brought to their doors and did not plan or intend to bring the dogs to the fixed points in

their areas. Spatially heterogeneous vaccination coverage is undesirable for dog rabies

control and elimination. Townsend et al. [41] found that such patchy coverage can

"profoundly damage prospects of elimination […] by creating pockets where rabies could

persist", and modeled patchy coverage within 1 km^2 cell grids. The low coverage

'patches' in our study had smaller areas than 1 km^2, thus the potential for a threat to

elimination efforts may be different or non-existent. However, in these densely populated

areas it is unknown if these 'patches' are large enough to sustain rabies transmission in

the city.

Many studies have explored logistical, informational, social and structural barriers for

dog rabies vaccination experienced by owners in rabies-affected areas

[1,17,18,23,24,40,42-52]. Two of the most common reasons identified are difficulty

handling the dogs [1,23,24,48,49,51,52] and lack of time [1,24,48,49,51,52]. These two

logistical barriers are correlated with a less studied element: the distance to the

vaccination campaigns. Distance to health services has been fairly well studied in terms

of availability of and access to health care and impact on health, especially for maternal

health, treatment and prevention of chronic diseases and treatment adherence for

infectious diseases [53-55], but not as much for preventing infectious diseases. Some

rural studies mention distance as a potential factor for low dog vaccination coverage

[23,56] and two studies directly evaluated the association between distance and overall

rural villages vaccination coverage [49] and the association between distance and

attendance at the MDVC in Sub-Saharan Africa [18]. In the sub-Saharan Africa study,

researchers found that distance in dispersed communities have an impact on MDVC

attendance [18]. Given the hilly landscape with rare direct paths between houses and

vaccination points, they estimated the shortest-path distance for their analysis to take into

consideration the long and tortuous routes dog owners had to follow to visit the

vaccination sites. In our study area, with non-dispersed urbanized highly-populated

localities and high density of street intersections that increase walkability, distance is still

an important proximal explanation for low participation in the vaccination campaigns.

Our study area consisted of urban and peri-urban localities. Surprisingly, there were no

clear differences in participation in the MDVC between these two groups. However, the

distribution of other proximal rabies-related characteristics is different between them (e.g.

more free-roaming dogs in peri-urban areas, more neutered/spayed dogs in urban areas).

There are other social determinants that provide more distal explanations for participation

in the MDVC. In previous focus groups conducted by our team, young females reported

that having a baby at home could prevent them from participating in the campaign [1].

Similarly, we found that households with children under 5 years old were less likely to

vaccinate their dogs compared to houses without children, and each additional child

under 5 reduced the odds of vaccinating the dogs in the house. This insight suggests an

opportunity to increase participation in the MDCV by framing the decision to vaccinate

as an action taken to protect children in the household from rabies. There was also a clear

difference in participation between those who lived in a dog rabies-endemic area before

living in Arequipa. A possible explanation for that difference is higher awareness among

that group. In our focus groups, we found low awareness and low perception of severity

among residents of Arequipa [1]. Interestingly, another component of migration history

was also associated with participation in the MDVC: time living in the area. This

phenomenon has been observed for the utilization of other health services in different

settings and populations [53-55]. Migration, settlement and adaptation are processes that

take time and are necessary for the uptake of health services [56-59], and could be

influencing the participation in the MDVC. Importantly, in the peri-urban localities there

are more households with children under 5, more recent migrants, and more people

whose previous residence was in a rabies-affected region.

Our study has a number of limitations. Other studies have focused attention on dog-level

variables (e.g. sex, age, function) that might be associated to participation in MDVC [18].

We did not analyze these; rather we focus on owner and community characteristics that

can be utilized by the health authorities (who rarely have the opportunity to collect

detailed house-by-house information) to increase participation in MDVC. Vaccination

status and access to the street were reported by the owner/interviewee. Given the bad

publicity in the local media about owned free roaming unvaccinated dogs and the

authorities' threats to fine 'irresponsible' dog owners, there is potential for social

desirability bias to inflate our estimate of vaccination coverage and deflate that of the

proportion of dogs that have access to the street. We did not ask the interviewees to show

the vaccination certificate they receive at the MDVC since many of them do not save

these certificates. We used Euclidean distances, which are only a proxy for the real

distance traveled by individuals to a vaccination post.

Dog rabies virus transmission continues in Arequipa, putting at risk millions of people in

the city and the surrounding departments. Dog-focused public health strategies are not

limited to rabies: deworming dogs to prevent human echinococcosis [57], use of

insecticide treatment [58] or vaccines [59] on dogs to prevent human Leishmaniasis or

Chagas disease, are just a few examples. These programs, if they are to be successful,

require high coverage and evenness in their implementation. The same approaches to

reach the appropriate levels of community participation that might have worked in the

1980s are not working today. Understanding the barriers for dog owners' participation in

community-based programs will be crucial to implement effective zoonotic disease

preventive activities. Distance to health services and the heterogeneous social

composition of growing cities have to be examined when designing field programs to

protect against zoonotic diseases.

5. Acknowledgments

We thank all the survey participants for agreeing to be part of our study and sharing their

time with us. We acknowledge the work of the members of the Zoonotic Disease

Research Laboratory for their contribution to the implementation of this study, especially

Katty Borrini and Jorge Apaza.

6. References

1. Castillo-Neyra R, Brown J, Borrini K, Arevalo C, Levy MZ, Buttenheim A, et al. Barriers to dog rabies vaccination during an urban rabies outbreak: Qualitative findings from Arequipa, Peru. Recuenco S, editor. PLoS Negl Trop Dis. 2017;11: e0005460. doi:10.1371/journal.pntd.0005460

2. Castillo-Neyra R, Levy MZ, Náquira C. [Effect of free-roaming dogs culling on the control of canine rabies]. Rev perú med exp salud publica. 2016;33: 772–779. doi:10.17843/rpmesp.2016.334.2564

3. Ministerio de Salud del Peru - Direccion General de Epidemiologia. Alerta Epidemiológica [Epidemiologic Alert]. Lima; 2015. Report No.: AE-DEVE N. 003-2015.

4. El Presidente de la República. Decreto Supremo que declara en Emergencia Sanitaria por el plazo de noventa (90) días calendario, a la provincia de Arequipa del departamento de Arequipa [Internet]. Apr 2, 2016 pp. 582224–582229. Available: busquedas.elperuano.pe/normaslegales/decreto-supremo-que-declara-en-emergencia-sanitaria-por-el-p-decreto-supremo-n-015-2016-sa-1363166-5/

5. Aréchiga Ceballos N, Karunaratna D, Aguilar Setién A. Control of canine rabies in developing countries: key features and animal welfare implications. Rev - Off Int Epizoot. 2014;33: 311–321.

6. Wallace RM, Reses H, Franka R, Dilius P, Fenelon N, Orciari L, et al. Establishment of a Canine Rabies Burden in Haiti through the Implementation of a Novel Surveillance Program. PLoS Negl Trop Dis. 2015;9: e0004245. doi:10.1371/journal.pntd.0004245

7. Widdowson M-A, Morales GJ, Chaves S, McGrane J. Epidemiology of urban canine rabies, Santa Cruz, Bolivia, 1972-1997. Emerging Infect Dis. 2002;8: 458–461. doi:10.3201/eid0805.010302

503   8.   Vigilato MAN, Clavijo A, Knobl T, Silva HMT, Cosivi O, Schneider MC, et al.
504   Progress towards eliminating canine rabies: policies and perspectives from Latin
505   America and the Caribbean. Philos Trans R Soc Lond, B, Biol Sci. 2013;368:
506   20120143. doi:10.1098/rstb.2012.0143

507   9.   Fooks AR, Banyard AC, Horton DL, Johnson N, McElhinney LM, Jackson AC.
508   Current status of rabies and prospects for elimination. The Lancet. 2014;384:
509   1389–1399. doi:10.1016/S0140-6736(13)62707-5

510   10.   Hampson K, Coudeville L, Lembo T, Sambo M, Kieffer A, Attlan M, et al.
511   Estimating the global burden of endemic canine rabies. PLoS Negl Trop Dis.
512   2015;9: e0003709. doi:10.1371/journal.pntd.0003709

513   11.   World Health Organisation. WHO Expert Consultation on Rabies. Second report.
514   World Health Organ Tech Rep Ser. 2013;: 1–139– back cover.

515   12.   Franka R, Smith TG, Dyer JL, Wu X, Niezgoda M, Rupprecht CE. Current and
516   future tools for global canine rabies elimination. Antiviral Res. 2013;100: 220–
517   225. doi:10.1016/j.antiviral.2013.07.004

518   13.   Lackay SN, Kuang Y, Fu ZF. Rabies in small animals. The Veterinary clinics of
519   North America Small animal practice. 2008;38: 851–859.

520   14.   Clavijo A, Del Rio Vilas VJ, Mayen FL, Yadon ZE, Beloto AJ, Vigilato MAN, et
521   al. Gains and future road map for the elimination of dog-transmitted rabies in the
522   Americas. Am J Trop Med Hyg. The American Society of Tropical Medicine and
523   Hygiene; 2013;89: 1040–1042. doi:10.4269/ajtmh.13-0229

524   15.   Vigilato MAN, Cosivi O, Knobl T, Clavijo A, Silva HMT. Rabies Update for
525   Latin America and the Caribbean. Emerging Infect Dis. 2013;19: 678–679.
526   doi:10.3201/eid1904.121482

527   16.   Davlin SL, Vonville HM. Canine rabies vaccination and domestic dog population
528   characteristics in the developing world: a systematic review. Vaccine. 2012;30:
529   3492–3502. doi:10.1016/j.vaccine.2012.03.069

530   17.   Cleaton JM, Wallace RM, Crowdis K, Gibson A, Monroe B, Ludder F, et al.
531   Impact of community-delivered SMS alerts on dog-owner participation during a
532   mass rabies vaccination campaign, Haiti 2017. Vaccine. 2018;36: 2321–2325.
533   doi:10.1016/j.vaccine.2018.03.017

534   18.   Mazeri S, Gibson AD, Meunier N, Bronsvoort BMD, Handel IG, Mellanby RJ, et
535   al. Barriers of attendance to dog rabies static point vaccination clinics in Blantyre,
536   Malawi. Rodriguez-Barraquer I, editor. PLoS Negl Trop Dis. 2018;12: e0006159.
537   doi:10.1371/journal.pntd.0006159

538   19.   World Health Organisation. WHO Expert Consultation on Rabies. World Health
539   Organization technical report series. Geneva; 2018 pp. 1–139.

540   20.   Cleaveland S, Kaare M, Tiringa P, Mlengeya T, Barrat J. A dog rabies vaccination
541   campaign in rural Africa: impact on the incidence of dog rabies and human dog-
542   bite injuries. Vaccine. 2003;21: 1965–1973.

543   21.   Jibat T, Hogeveen H, Mourits MCM. Review on dog rabies vaccination coverage
544   in Africa: a question of dog accessibility or cost recovery? Rupprecht CE, editor.
545   PLoS Negl Trop Dis. 2015;9: e0003447. doi:10.1371/journal.pntd.0003447

546   22.   Gsell AS, Knobel DL, Kazwala RR, Vounatsou P, Zinsstag J. Domestic dog
547   demographic structure and dynamics relevant to rabies control planning in urban
548   areas in Africa: the case of Iringa, Tanzania. BMC Vet Res. 2012;8: 236.
549   doi:10.1186/1746-6148-8-236

550   23.   Kaare M, Lembo T, Hampson K, Ernest E, Estes A, Mentzel C, et al. Rabies
551   control in rural Africa: Evaluating strategies for effective domestic dog
552   vaccination. Vaccine. 2009;27: 152–160. doi:10.1016/j.vaccine.2008.09.054

553   24.   Muthiani Y, Traoré A, Mauti S, Zinsstag J, Hattendorf J. Low coverage of central
554   point vaccination against dog rabies in Bamako, Mali. Prev Vet Med. 2015;120:
555   203–209. doi:10.1016/j.prevetmed.2015.04.007

556   25.   Perry BD, Kyendo TM, Mbugua SW, Price IE, Varma S. Increasing rabies
557   vaccination coverage in urban dog populations of high human population density
558   suburbs: a case study in Nairobi, Kenya. Prev Vet Med. 1995;22: 137–142.
559   doi:10.1016/0167-5877(94)00407-A

560   26.   Schneider MC, Belotto A, Adé MP, Hendrickx S, Leanes LF, Rodrigues MJ de F,
561   et al. Current status of human rabies transmitted by dogs in Latin America. Cad
562   Saude Publica. 2007;23: 2049–2063.

563   27.   Castillo-Neyra R, Zegarra E, Monroy Y, Bernedo R, Cornejo-Rosello I, Paz-
564   Soldan V, et al. Spatial Association of Canine Rabies Outbreak and Ecological
565   Urban Corridors, Arequipa, Peru. Tropical Medicine and Infectious Disease.
566   2017;2: 38. doi:10.3390/tropicalmed2030038

567   28.   Burgos-Cáceres S. Canine Rabies: A Looming Threat to Public Health. Animals
568   (Basel). 2011;1: 326–342. doi:10.3390/ani1040326

569   29.   Zhang L, Li X, Yang H, Mao R, Zhao Q. Health status and health-seeking
570   behaviour between interprovincial and intraprovincial rural-to-urban young
571   migrants in Nanjing China. Asia-Pacific Population Journal. 2011;26: 39–54.

572   30.   Li Y. Understanding health constraints among rural-to-urban migrants in China.
573   Qual Health Res. 2013;23: 1459–1469. doi:10.1177/1049732313507500

574   31.   Benyoussef A, Cutler JL, Levine A, Mansourian P, Phan-Tan T, Baylet R, et al.
575   Health effects of rural-urban migration in developing countries--Senegal. Soc Sci
576   Med. 1974;8: 243–262.

32.   Carrillo-Larco RM, Bernabé-Ortiz A, Pillay TD, Gilman RH, Sanchez JF, Poterico
JA, et al. Obesity risk in rural, urban and rural-to-urban migrants: prospective
results of the PERU MIGRANT study. Int J Obes (Lond). 2015.
doi:10.1038/ijo.2015.140

33.   Urbanization, land tenure security and vector-borne Chagas disease.

34.   Taylor LH, Wallace RM, Balaram D, Lindenmayer JM, Eckery DC, Mutonono-
Watkiss B, et al. The Role of Dog Population Management in Rabies Elimination-
A Review of Current Approaches and Future Opportunities. Front Vet Sci.
Frontiers; 2017;4: 109. doi:10.3389/fvets.2017.00109

35.   Sambo M, Lembo T, Cleaveland S, Ferguson HM, Sikana L, Simon C, et al.
Knowledge, Attitudes and Practices (KAP) about Rabies Prevention and Control:
A Community Survey in Tanzania. Rupprecht CE, editor. PLoS Negl Trop Dis.
2014;8: e3310. doi:10.1371/journal.pntd.0003310

36.   Venables WN, Ripley BD. Modern Applied Statistics with S. 4 ed. New York:
Springer; 2002.

37.   Baddeley A, Turner R. spatstat: An RPackage for Analyzing Spatial Point Patterns.
Journal of Statistical Software. 2005;12. doi:10.18637/jss.v012.i06

38.   Bai L, Bartell S, Bliss R, Vieira V. MapGAM: Mapping Smoothed Effect
Estimates from Individual-Level Data [Internet]. 1st ed. Available:
https://CRAN.R-project.org/package=MapGAM

39.   Team. R: A Language and Environment for Statistical Computing [Internet].
Vienna, Austria: R Foundation for Statistical Computing; 2013. pp. 1–3604.
Available: http://www.R-project.org/

40.   Durr S, Mindekem R, Kaninga Y, Doumagoum Moto D, Meltzer MI, Vounatsou
P, et al. Effectiveness of dog rabies vaccination programmes: comparison of
owner-charged and free vaccination campaigns. Epidemiol Infect. 2009;137:
1558–1567. doi:10.1017/S0950268809002386

41.   Townsend SE, Sumantra IP, Pudjiatmoko, Bagus GN, Brum E, Cleaveland S, et al.
Designing programs for eliminating canine rabies from islands: Bali, Indonesia as
a case study. Rupprecht CE, editor. PLoS Negl Trop Dis. 2013;7: e2372.
doi:10.1371/journal.pntd.0002372

42.   Widyastuti MDW, Bardosh KL, Sunandar   , Basri C, Basuno E, Jatikusumah A,
et al. On dogs, people, and a rabies epidemic: results from a sociocultural study in
Bali, Indonesia. Infectious Diseases of Poverty. 2015;4: 60. doi:10.1186/s40249-
015-0061-1

43.   Bardosh K, Sambo M, Sikana L, Hampson K, Welburn SC. Eliminating rabies in
Tanzania? Local understandings and responses to mass dog vaccination in
Kilombero and Ulanga districts. Rupprecht CE, editor. PLoS Negl Trop Dis.
2014;8: e2935. doi:10.1371/journal.pntd.0002935

44.   Serebe SG, Tadesse KA. Study on community knowledge, attitude and practice of
rabies in and nearby Gondar town, North West Ethiopia. Journal of Public ….
2014.

45.   Kabeta T, Deresa B, Tigre W, Ward MP, Mor SM. Knowledge, Attitudes and
Practices of Animal Bite Victims Attending an Anti-rabies Health Center in Jimma
Town, Ethiopia. Zinsstag J, editor. PLoS Negl Trop Dis. 2015;9: e0003867.
doi:10.1371/journal.pntd.0003867

46.   De Ramos M, Bravo LC. Knowledge, attitudes, and practices of community
regarding animals bite and rabies. Paediatric Infectious Disease Society of the ….
2004.

47.   Singh US, Choudhary SK. Knowledge, attitude, behavior and practice study on
dog-bites and its management in the context of prevention of rabies in a rural
community of Gujarat. Indian Journal of Community Medicine. 2005.

48.   Kayali U, Mindekem R, Yemadji N. Coverage of pilot parenteral vaccination
campaign against canine rabies in N'Djamena, Chad. Bulletin of the World ….
2003.

49.   Minyoo AB, Steinmetz M, Czupryna A. Incentives Increase Participation in Mass
Dog Rabies Vaccination Clinics and Methods of Coverage Estimation Are
Assessed to Be Accurate. PLoS Negl Trop …. 2015.

50.   Lembo T, Hampson K, Haydon DT, Craft M, Dobson A, Dushoff J, et al.
Exploring reservoir dynamics: a case study of rabies in the Serengeti ecosystem.
The Journal of Applied Ecology. 2008;45: 1246–1257. doi:10.1111/j.1365-
2664.2008.01468.x

51.   Lapiz SMD, Miranda MEG, Garcia RG, Daguro LI, Paman MD, Madrinan FP, et
al. Implementation of an intersectoral program to eliminate human and canine
rabies: the Bohol Rabies Prevention and Elimination Project. Zinsstag J, editor.
PLoS Negl Trop Dis. 2012;6: e1891. doi:10.1371/journal.pntd.0001891

52.   Ferguson EA, Hampson K, Cleaveland S, Consunji R, Deray R, Friar J, et al.
Heterogeneity in the spread and control of infectious disease: consequences for the
elimination of canine rabies. Sci Rep. 2015;5: 18232. doi:10.1038/srep18232

53.   Jacobs B, Ir P, Bigdeli M, Annear PL, Van Damme W. Addressing access barriers
to health services: an analytical framework for selecting appropriate interventions
in low-income Asian countries. Health Policy and Planning. 2012;27: 288–300.
doi:10.1093/heapol/czr038

54.   Bissonnette L, Wilson K, Bell S, Shah TI. Neighbourhoods and potential access to
651   health care: The role of spatial and aspatial factors. Health & Place. 2012;18: 841–
652   853. doi:10.1016/j.healthplace.2012.03.007

653   55.   Hiscock R, Pearce J, Blakely T, Witten K. Is Neighborhood Access to Health Care
654   Provision Associated with Individual-Level Utilization and Satisfaction? Health
655   Services Research. 2008;43: 2183–2200. doi:10.1111/j.1475-6773.2008.00877.x

656   56.   Cleaveland S. The epidemiology of rabies and canine distemper in the Serengeti,
657   Tanzania. doi:https://doi.org/10.17037/PUBS.00682291

658   57.   Fu Q, Han X-M, Wang L-Y, Zhaxi S-M, Ma X, Wang Y-S, et al. Effect of control
659   pattern with emphasis on canine deworming in a pilot of echinococcosis in highly
660   endemic area, southern Qinghai Plateau. Zhongguo Ji Sheng Chong Xue Yu Ji
661   Sheng Chong Bing Za Zhi. 2011;29: 293–295.

662   58.   Loza A, Talaga A, Herbas G, Canaviri RJ, Cahuasiri T, Luck L, et al. Systemic
663   insecticide treatment of the canine reservoir of Trypanosoma cruzi induces high
664   levels of lethality in Triatoma infestans, a principal vector of Chagas disease.
665   Parasites & Vectors. BioMed Central; 2017;10: 344. doi:10.1186/s13071-017-
666   2278-2

667   59.   Aparicio-Burgos JE, Zepeda-Escobar JA, de Oca-Jiménez RM, Estrada-Franco JG,
668   Barbabosa-Pliego A, Ochoa-García L, et al. Immune protection against
669   Trypanosoma cruzi induced by TcVac4 in a canine model. Santiago HDC, editor.
670   PLoS Negl Trop Dis. 2015;9: e0003625. doi:10.1371/journal.pntd.0003625

7. Supporting Information Legends

675   S1 Alternative Language Abstract. Resumen en español.

677   S1 Table. Reported exposure to MDVC communication channels by participation level.

Effect of distance from dog owner's house to the closest vaccination point on clustering of unvaccinated dogs