Marked compositional changes in harvestmen assemblages in Amazonian forest islands induced by a mega dam

Ana Lúcia Tourinho[1*], Maíra Benchimol[2], Willians Porto[3], Carlos A. Peres[4] & Danielle Storck-Tonon[5]

1. Universidade Federal do Mato Grosso, Instituto de Ciências Naturais, Humanas e Sociais, Núcleo de Estudos da Biodiversidade da Amazônia Matogrossense (NEBAM), Av. Alexandre Ferronato, 1200, Setor Industrial, CEP: 78577-267, Sinop, MT, Brazil.

2. Laboratório de Ecologia Aplicada à Conservação, Universidade Estadual de Santa Cruz, Rodovia Jorge Amado km 16, CEP: 45662-900 Ilhéus, BA, Brazil.

3. División de Aracnología, Museo Argentino de Ciencias Naturales–CONICET, Av. Ángel Gallardo 470, C1405DJR Buenos Aires, Argentina.

4. School Environmental Sciences, University of East Anglia, Norwich NR47TJ, UK

5. Programa de Pós-Graduação em Ambiente e Sistemas de Produção Agrícola, Universidade do Estado de Mato Grosso, Rod. MT 358, km 7 - Jardim Aeroporto. CEP: 78300-000, Tangará da Serra, MT, Brazil.

*Corresponding author: Ana Lúcia Tourinho, <amtourinho@gmail.com>

Running Title: Mega dam drives changes in harvestmen assemblages

Abstract

Mega hydroelectric dams have become one of the main drivers of habitat loss in tropical forests, converting large tracts of pristine forests into isolated forest islands. Understanding how biodiversity cope with landscape modification in these archipelagic landscapes is of paramount importance to assess the environmental consequences of dam infrastructure and propose mitigation actions for biodiversity conservation. In this context, harvestmen (Opiliones, Arachnida) comprise a good indicator taxon of habitat quality, given their high sensitivity to desiccation and microclimatic change. Here, we investigate the effects of landscape change induced by a mega hydropower dam on forest harvestmen species richness, abundance and composition within the Balbina Hydroelectric Dam, Central Brazilian Amazon. Using both nocturnal search and diurnal beating tray surveys, we sampled 20 islands and five mainland continuous forests, relating our biological response variables to local, patch and landscape scale metrics. We captured 783 harvestmen representing 31 species, 22 genera and 9 families considering all sampling sites, with species composition differing across islands and mainland sites. Although species richness was unaffected by any local, patch and landscape variables, species composition and abundance were differentially affected by a set of predictor variables at different scales. Forest cover and fallen woody stems were significant predictors of species composition patterns, whereas vegetation density, forest cover, island area, abundance of palm trees, and fallen woody stems best explained harvestmen abundance. We recommend retaining large forest habitat patches surrounded by a great amount of forest cover to prevent the rapid decay in tree communities and minimize forest disturbance effects in large hydroelectric dams.

Introduction

Hydroelectric dams have been posed as a major new villain for Amazonian rainforest and freshwater ecosystems, severely inducing the widespread habitat loss and wholesale local species extinctions (Benchimol & Peres, 2015a; Bobrowiec & Tavares, 2017, Lees et al., 2016). The Amazon basin accounts for 18% of global scale river discharge (Dai & Trenberth, 2002), so it is not surprising that this region has rapidly become a major target for hydropower development; in addition to 191 existing dams, 246 additional dams may be built across the whole Amazon Basin in the next few decades (Lees et al. 2016). Indeed, more than 1,300,00 ha of forest has already been flooded by operational dams in the Brazilian Amazon alone (ECOA, 2018).

Several studies have shown the pervasive social and environmental impacts of large dams. From local communities (Esselman & Opperman, 2010, Fearnside, 2016) to fluvial hydrology (Nilsson et al., 2005), large dams affect indigenous communities by flooding their territories, and increasing greenhouse gas emissions (Almeida et al., 2013). They also drive local extinctions of both aquatic (Alho, 2011; Liermann et al., 2012; Palmeirim et al., 2014) and terrestrial fauna and flora (Terborgh et al., 1997; Benchimol & Venticinque, 2014, Benchimol & Peres, 2015a,b,c). However, few studies have assessed the effects of large dams on terrestrial invertebrates, despite their immense importance for forest cycling and ecosystem services (Storck-Tonon & Peres, 2017, Carvalho et al., 2012). For instance, a study has shown that most orchid bee species are driven to local extinction after 26 years of insularization within a mega dam in central Brazilian Amazonia, with potential effects on pollination (Storck-Tonon & Peres, 2017).

Several groups of arthropods can be described as effective bioindicators, providing valuable information on environmental quality at both narrow and large scales on the basis of very brief surveys, enabling their efficient inclusion in environmental licensing studies prior to development projects (Maleque et al., 2006, Melo et al., 2015). Additionally, arthropods play an important role in many key processes influencing ecosystem cycling of carbon and other nutrients, and modulating the amount and quality of resources entering the detrital foodweb (Yang & Gratton, 2014). These processes may occur via direct or indirect pathways mediated through species interactions via plants and soil microbes, integrating community-level interactions with ecosystem processes. Arachnids are mostly generalist predators, and are often more dependent on the physical structure of microhabitats than on prey availability (Villanueva-Bonilla et al., 2017). Although most species can inhabit humid habitats, harvestmen species (Opiliones, Arachnida) are physiologically dependent of humid environments, due to their high susceptibility to desiccation, nocturnal activity and low vagility (Pinto-da-Rocha et al., 2007).

Despite the significant ecological importance of harvestmen (Fig. 2), serving as excellent predictors of environmental quality given their rapid responses to habitat change and fragmentation (Bragagnolo et al., 2007), this charismatic taxonomic group remains poorly investigated in tropical forest landscapes. Additionally, patterns of diversity in harvestmen assemblages are highly correlated with vegetation structure at

meso and local scales, and are especially responsive to the presence of large old trees and palms in Amazonian forests (Colmenares et al., 2016; Tourinho et al., 2014, 2017, Porto et al., 2016). This occurs because palms increase habitat complexity at ground level, as a consequence of the fallen litter trapped within their leaves (Vasconcelos *et al*., 1990, Franken et al., 2013). Therefore, both palm and non-palm trees are directly responsible for the quality and depth of the soil leaf-litter layer, which is a key microhabitat component affecting assemblages of small-bodied harvestmen species and other invertebrates (Tarli et al., 2014, Colmenares et al., 2016, Porto et al. 2016).

To date, only one study, conducted in forest patches embedded within heterogeneous vegetation matrices in the Brazilian Atlantic forest, has assessed the effects of forest fragmentation on harvestmen within a tropical forest landscape (Bragagnolo, et al. 2007). In this study, harvestmen assemblages were more clearly structured by forest quality and size than other arthropod taxa, suggesting that they are good community-wide indicators of intermediate-level disturbances and can be used as a robust model to monitor tropical forest disturbance. However, we still lack information on how this key taxonomic group is affected by habitat insularization, with archipelagic ecosystems formed in the aftermath of rising floodwaters induced by large dams providing an unique natural experiment in this context.

Here, we assessed the degree to which forest fragmentation induced by a mega hydroelectric dam affects harvestmen assemblages in the Brazilian Amazon. We conducted sampling surveys at five mainland continuous forest sites and 20 widely distributed forest islands differing in size, shape and degree of isolation within the ~443,772-hectares Balbina Hydroelectric Dam. Specifically, we investigate (1) which are the major factors structuring harvestmen assemblages in these insular forest environments; and (2) the potential effect of habitat structure on harvestmen assemblages, and how they respond to the habitat insularization process.

Materials and Methods

Study site

We conducted this research at the Balbina Hydroelectric Reservoir landscape - BHR (1°48'S; 59°29'W), located in the state of Amazonas, central Brazilian Amazonia. The

Balbina Dam is a large man-made reservoir that was formed in 1987 by the damming of the Uatumã river, aiming to supply power to the growing capital city of Manaus. An area of 312,900 ha was inundated, resulting in the creation of 3,546 islands ranging in size from <1 to 4,878 ha. Most islands are covered by primary sub-montane dense forests, and are surrounded by open-water and dead trees that rise above the water level; the mean water column depth range from 7.4 m (in most of the reservoir) to 30 m (in the former river channel; Eletronorte, 1997). Mean annual temperature is 28°C and the mean annual precipitation is 2376 mm. A large tract of pristine continuous forest sites surrounds the Balbina archipelagic landscape, and share similar floristic composition, fauna, climate and topography (Benchimol & Peres, 2015a,c). In 1990, part of the archipelago and an extensive tract of neighbouring pristine forest became protected by the creation of the ~940,358-ha Uatumã Biological Reserve, the largest Biological Reserve in Brazil. Although islands are unaffected by logging and hunting pressure, several islands experienced ephemeral understorey fires during the El Niño drought of late-1997 (Benchimol & Peres, 2015a).

Study Design

We selected 20 islands using two cloudless georeferenced Landsat ETM+ scenes. Additionally, we surveyed five continuous ('pseudo-control') forest sites located in the southernmost part of the Uatumã Reserve, within a permanent study grid of the PPBio (Brazilian Biodiversity Research Program, Fig. 1). Islands were selected on the basis of their size, shape, isolation and spatial distribution, keeping a minimum distance of 1 km from one another (Fig. 1). Likewise, each continuous forest site in the mainland was also spaced by 1 km from each other. Island area ranged from 11.8 to 218.5 ha (mean ± SD = 91.5 ± 69.7 ha), and ranged widely in isolation distances [to the nearest mainland continuous forest] from 10 m to 9,000 m (mean ± SD = 3,220 ± 2,648 m).

Harvestmen sampling

Between 2006 and 2008, we systematically sampled harvestmen assemblages (Fig. 2) using four 300-m^2 transects (30 m × 10 m) at each forest site, separated by 30 m, totalling 1,200 m^2 of transect area per site (see Tourinho et al., 2014, Porto et al., 2016 and Tourinho et al., 2018). To avoid edge effects on islands, transects were established at least 250 m from the nearest forest edge. Each plot was surveyed using both a

nocturnal search method (Sørensen et al., 2002; Pinto-da-Rocha and Bonaldo, 2006; Resende et al., 2012) and diurnal beating tray surveys. Nocturnal searchers were performed by one collector for 1 hour, during which all harvestmen (and other arachnids) encountered on the forest floor to a height of 2 m, on tree trunks and other vegetation were captured. Beating tray surveys were conducted during daylight hours, by one collector for 1 h, during which shrubs up to 3 m in height were struck 20 times with a wooden stick. We used a 1-m^2 white fabric wooden frame positioned underneath the shrubs to capture any fallen harvestmen.

All individuals collected were subsequently identified at the lowest taxonomic level by experts on harvestmen taxonomy (Ana Lúcia Tourinho, Pío Colmenares and Willians Porto). For this, we examined the external morphology of individual specimens under a stereomicroscope, which were compared with original descriptions provided in the literature (Pinto-da-Rocha, 1994, 1996, 1997, 2004; Kury, 2003), type specimens, and photographs of type specimens. We excluded from the dataset nymphs and females, which have ambiguous morphology, because they cannot be assigned to any genus or species level. In the case of species groups with a very conservative external morphology and/or poorly understood taxonomy (e.g. Cosmetidae, Sclerosomatidae, Zalmoxidae) we also examined their male genitalia to improve species delimitation, following Acosta et al. (2007). All specimens collected were preserved in 70% ethanol and deposited in the Invertebrate Collection of the Instituto Nacional de Pesquisas da Amazônia, Manaus, Amazonas, Brazil, under curator, Dr Célio Magalhães.

Landscape and habitat variables

We used RapidEye$^©$ high-resolution (5-m pixel) imagery for the entire Balbina archipelago to quantify landscape structure variables at the patch and landscape-scales, associated with each of surveyed forest site. At the patch scale, we obtained the total island area in hectares ($\log_{10}$ x; 'area'); the isolation distance from each island to the nearest mainland ('$D_{mainland}$'); and the island shape ('shape') by dividing the total island area by the total island perimeter. At the landscape scale, we estimated the total amount of forest cover ('cover') within a specified radius, and modified the proximity index of McGarigal et al., (2012) by considering the total size of any land mass within the buffer, rather than excluding land areas outside the buffer for patches contained within the

buffer ('prox'). We considered multiple buffers (250m, 500m and 1000m) outside the perimeter of each focal island and mainland forest sites for landscape-scale analyses.

We also obtained variables at local scale, including the litter depth ('litdepth'), the density of trees ('dtrees'), the total number of palm trees ('palms') and the number of fallen woody stems ('fallen'). 'Fallen' considered down wood and does not include arborescent palm left bracts all of which were used here as proxies of vegetation habitat structure available to harvestmen (Pio et al. 2016). For this, we mapped all non-palm and palm trees with a diameter at breast height (DBH) > 30 cm within each 300-m^2 transect. We also mapped all fallen stems within each plot. At every 5 m along the long axis of the transects, a measurement of litter depth was taken. These measurements consisted of forcing a wooden stick 0.5 cm in diameter into the litter until it reached the soil and noting the height (in cm) above ground to the top layer of litter.

Data analysis

We firstly compared the coefficient of determination (R^2) from regression analysis of harvestmen species richness for each forest cover radius (i.e., 250 m, 500 m, 1000 m) to determine which radial buffer best explained diversity patterns. The highest R^2 values were obtained at 1000 m radius (see Table S1), so we used this buffer size for all subsequently analyses. We thus performed a Pearson correlation matrix between all landscape and habitat variables, excluding variables that were highly correlated ($r \geq 0.50$) from the analysis, resulting in five habitat and landscape variables maintained in the models: area, litter depth, number of palm trees, number of fallen woody stems, and forest cover.

To assess differences in harvestmen assemblage structure among the habitat types (islands and mainland) we used permutational multivariate analysis of variance (PERMANOVA). To analyse changes in species composition between all surveyed sites, we performed multiple regression on distance matrices (MRM) analysis. MRM is similar to partial Mantel tests and was used to examine the correlation between the dependent distance matrix and the independent explanatory distance matrices. We used the abundance matrix of each species obtained within the transects to run a Principal Coordinates Analysis (PCoA) and reduce the dimensionality of the database to two or more axes, as this allowed easier visualization and interpretation of the data collected,

resulting in the creation of principal coordinates that represents the axes which explain the highest amount of variation. We ordinated the transects by species composition using the Bray-Curtis dissimilarity index and used the first three PCoA axes as predictors of species community. Following the theory of island biogeography (MacArthur & Wilson 1967), we first examined the potential effects of island size ($\log(x+1)$) and isolation ($D_{mainland}$) on harvestmen species richness, abundance and composition using linear regression. For the latter, we reduced the data dimensionality using NMDS ordinations based on both the Jaccard (presence/absence) and Bray-Curtis (abundance) similarity values, and used the first axis as a predictor of species composition.

We further performed Generalized Linear Models (GLMs) to examine the relationship between harvestmen species richness, abundance and composition ($NMDS_1$) of islands and the explanatory variables. Multicollinearity among these variables was tested using Variation Inflation Factors (VIF) (Domman et al. 2013) using the 'HH' package (Heiberger, 2016), deleting the least moderately redundant or collinear variables (VIF > 6). We used Poisson, Gaussian and quasi-Poisson error structures for GLMs of species richness, composition, and abundance, respectively. We ran all GLMs using the 'MuMIn' package (Barton, 2018) and retained all models that differed by $\Delta AIC \leq 2.00$ (Burnham & Anderson 2002). All analyses were performed within the R 3.3.1 environment (R Core Team 2016).

Results

A total of 783 harvestmen representing 31 species, 22 genera and 9 families were collected across all sampling sites throughout the BHR (Table 2). These ranged along the size spectrum from the small-bodied Fissiphalliidae, Zalmoxidae and Stygnomatidae (2.00 ± 2.50 mm) to the large-bodied Cranaidae, Stygnidae species of *Protimesius* (4.95 ± 10.85 mm) (Fig. 2). The number of species co-occurring within any given island or continuous forest site ranged from 1 to 20 (mean $\pm$ SD = 6.25 ± 4.14) species. The families containing the largest numbers of species recorded were Stygnidae (9), Sclerosomatidae (6), Manaosbiidae (4), and Cosmetidae (4) (Table 2), whereas the most abundant families were Stygnidae (209) and Manaosbiidae (202). Nine species were collected exclusively on islands, 14 were exclusively collected at mainland sites, and

nine in both islands and mainland sites (Fig. 5, Table 2). The families with largest number of species on islands were Stygnidae (7) and Zalmoxidae (3), whereas the Sclerosomatidae (6) and Stygnidae (5) were the most speciose families in the mainland. All Sclerosomatidae (194 individuals, 6 species) were exclusively restricted to mainland sites (Fig. 5).

The species composition of harvestmen (Fig. 3) differed between islands and mainland continuous forest sites (PERMANOVA: F= 4.76, P =0.01). Three PCoA axes represent the species composition and they collectively explained 57% of overall variation in the data (PCoA axis 1: 25%, axis 2: 18%, axis 3: 15%). Species composition similarity, as represented by the second PCoA axis, was affected by geographic distance across all trap sites (MRM r = 0.14, p = 0.009), whereas the other two axes were unrelated to a distance effect. When we consider only islands in the MRM model, none of compositional metrics were affected by distance (MRM r = 0.67, p = 0.35), and the first three PCoA axes explained 61% of overall variation (PCoA axis 1: 30%, PCoA axis 2: 18%, PCoA axis 3: 15%).

Considering only islands, there was no significant species-area relationship (SAR) on the basis of a semi-log model (R^2_{adj} = 0.000, P = 0.471). Island isolation distance to the nearest mainland area also failed to explain harvestmen species richness (R^2_{adj} = 0.000, P = 0.835). Similar patterns were also observed for species composition, with neither island size (R^2_{adj} = 0.001, P = 0.326) nor island isolation (R^2_{adj}= 0.000, P = 0.422) significantly affecting the first NMDS axis. Island size and isolation also failed to affect patterns of overall harvestmen abundance within sites (Area - R^2_{adj} = 0.000, P = 0.423; Isolation - R^2_{adj} = 0.000, P = 0.363).

When all explanatory variables were also incorporated into the models, GLMs showed that landscape-scale forest cover and fallen woody stems were the main predictors of local species composition. Considering species richness, however, none of the local habitat nor landscape variables were significant predictors in the 'best' model. In contrast, vegetation density, forest cover, island size, abundance of palm trees, and the amount of fallen woody stems were all significant predictors of harvestmen abundance (Table 1).

Discussion

As far as we are aware, this is the first study to examine harvestmen responses to habitat insularization induced by a large dam within tropical forests. Islands harboured fewer species than mainland continuous forest sites, and exhibited a simplified species composition. Contrary to our expectations, island area and isolation did not predict patterns of species richness, as predicted by the equilibrium theory of island biogeography (MacArthur and Wilson, 1967). However, a set of local, patch and landscape variables were included in the best models predicting species composition and abundance. Due to the detrimental impacts of hydroelectric dams on forest biota, we provide additional evidence that these infrastructure projects are far from the 'green energy' sources they are often considered to be (Gibson et al., 2017), and propose conservation actions to enhance the long term persistence of harvestmen assemblages and other arthropods within reservoir areas.

Main drivers of harvestmen assemblage structure

Contrary to our expectations, none of the local, patch and landscape variables were included in the best models explaining harvestmen species richness on islands. In Amazonian upland forest sites, microhabitat structure and availability per unit area explained patterns of species composition but also failed to explain species richness in harvestmen assemblages (Colmenares et al., 2016). In the Balbina archipelagic landscape, island size is a strong positive predictor of species richness in other taxonomic groups ─ including birds (Aurélio-Silva et al., 2016), small mammals (Palmeirim et al. 2018) and large mammals (Benchimol & Peres, 2015a). Although species-area relationship is one of the few ironclad rules in ecology (Mathews et al. 2016), the absence of this relationship can be attributed to other local habitat features. However, there were marked shifts in species composition, and we detected a significant species turnover between islands and continuous forest sites, indicating pervasive changes of harvestmen species assemblages on islands after 20 years of isolation. Both forest cover and fallen woody stems were the main joint drivers structuring the composition of harvestmen species on islands. Given that the presence of large canopy trees and palm tree architecture provide more available habitat, these variables have been considered as a good proxy of harvestmen diversity in studies at upland Amazonian forests (Colmenares et al., 2016, Porto et al., 2016, Tourinho et al., 2014).

Additionally, habitat quantity and quality have been equally proposed as robust predictors of harvestmen species composition in a fragmented Atlantic forest landscape (Bragagnolo et al., 2007), providing more evidence of the importance of habitat quality in maintaining tropical harvestmen assemblages. Fallen woody stems are often considered as one of the most likely microhabitats to contain harvestmen, especially aggregations during diurnal periods (Pinto-da-Rocha et al. 2007), yet the associations between opilione assemblages and this particular microhabitat have not been properly quantified in tropical forests. In fact, higher density of fallen woody stems may increase habitats favoured by harvestmen, providing ideal microclimatic conditions of humidity and temperature required for their development (Mestre and Pinto-da-Rocha, 2004; Curtis and Machado, 2007; Proud et al., 2012). Notably, this was an important variable structuring both harvestmen species composition and abundance across the Balbina islands. A positive association between harvestmen assemblages and fallen logs have not always been detected in undisturbed tropical forests, but this microhabitat type is often associated with specific harvestmen species in Amazonian forests (Colmenares et al., 2016).

The amount of landscape-scale forest cover was another variable retained in the best model influencing havestmen composition on Balbina islands. This variable was also included in the best model affecting patterns of abundance. The role of total amount of surrounding suitable habitat area is widely recognized in determining local patterns of diversity (Andrén 1994), and has been considered as a good proxy of habitat loss (Fahrig, 2003, 2013). Indeed, several studies in the Atlantic forest has showed the paramount importance of forest cover in predicting abundance patterns of insects (Morante-Filho et al., 2016) and vertebrates (Lira et al., 2012; Morante-Filho et al., 2015)

The amount of surrounding forest cover, number of palm and tree density, fallen woody stems and island size explained harvestmen abundance patterns at the Balbina islands. Thus, larger islands harbouring higher forest basal area, particularly arborescent palms, are necessary to sustain larger populations of harvestmen species on islands. In fact, tree microhabitats harboured the most diverse and abundant harvestmen assemblages, and was the only variable positively associated with harvestmen abundances in a Central Amazonian forest site distant ~190 km from Balbina (Colmenares et al., 2016).

Overall harvestmen abundance declined in response to local, patch and landscape-scale variables on surveyed islands. For instance, tree density was a strong predictor of harvestmen abundance patterns. Arboreal trees are highly sensitive to forest fragmentation, with many trees species becoming rapidly extirpated from small forest remnants (Silva & Tabarelli 2000; Laurance et al. 2006; Tabarelli, Peres & Melo 2012). This effect is potentially stronger on islands created by hydroelectric dams, with a rapid decay in the floristic diversity detected on small Balbina islands (Benchimol & Peres, 2015b). Fallen woody stems on small Balbina islands – where tree habitat diversity has been severely reduced – may therefore function as important alternative microhabitats for harvestmen assemblages. Therefore, most islands created by the Balbina dam are likely unfavourable for the long-term persistence of viable harvestmen populations, which may lead to local extinctions, creating a set of species-poor islands. On the other hand, species associated with fallen woody stems may benefit from tree mortality induced by high rates of windfalls on small islands (Benchimol & Peres 2015b), thereby becoming highly successful species in this archipelago (two Manaosbiidae and two *Stygnus*-Stygnidae species).

Although the influence of habitat quantity and structure over the abundance and spatial distribution of harvestmen has not been sufficiently explored, harvestmen have a close relationship with vegetation structure and microclimate which create environmental gradients affecting their species composition. Islands containing higher densities of mature trees and palms harbour distinct assemblages and provide the appropriate microclimatic conditions to retain harvestmen species (Colmenares et al., 2016). Harvestmen and other arachnids, such as spiders, are associated with tree bole surface area, particularly those providing the appropriate bark properties, which are important niches for resting, foraging (Southwood, 1978; Pinzón & Spence, 2010), and appropriate substrates for daytime camouflage and web attachment (Messas et al., 2014).

Species-specific responses to landscape insularization

According to several other studies, the diversity of harvestmen recorded in the islands was low (Pinto-da-Rocha & Bonaldo, 2006; Bonaldo et al., 2009; Colmenares et al., 2016; Tourinho et al., 2014, Porto et al., 2016; Tourinho et al. 2017). The small number of species detected per islands also reinforces diversity impoverishment in forest fragments, as suggested for the Atlantic Forest (Bragagnolo et al., 2007). Indeed,

55% of the sampled islands exhibited only one to five species, 45% harboured more than five, and from that just 15% (3 islands) presented 7, 8 and 10 species recorded. Differently, mainland sites retained 11-20 species, exhibiting a similar pattern obtained in other Amazonian sites (Tourinho et al. 2011, 2014, 2017, Porto et al. 2016, Bonaldo et. al, 2009). The intense sampling effort applied in this study - 20 islands (4 replicates each) - is also an indicative of islands impoverishment (160 hours sampling). As an example, species richness recovered in only 6 control in mainland areas, with a much inferior effort (48 h) rendered 22 species which may be considered comparable to other areas in the Amazon, and new additions in numbers of sites sampled would certainly increase the number of species sampled as well.

Interestingly, the number of exclusive species retained in islands (9) was not low as expected, but still lower than the number of exclusive species recorded in mainland forest sites, even with lower sampling effort (13). However, looking into these results we note that exclusive species sampled in islands are minute harvestmen species and/or specimens, except one. They are members of Agoristenidae, tiny Stygnidae genera, but mostly members of families of Zalmoxoidea and Samooidea that inhabit leaf litter layers and other cryptic environments in the Amazon forest, for which little collecting effort had been invested in this study. Given that we sampled few individuals occasionally, mostly singletons or doubletons, the lack of registers in mainland sites can also be an artefact, so we would expect to enhance the records of these specimens sifting leaf litter in mainland sites, as happened before in other Amazonian sites (Bonaldo et al. 2009, Tourinho et al. 2014).

Conversely, Cosmetidae and Sclerosomatidae species seem to be highly responsive to forest alteration and fragmentation. We did not record any of the six mainland Sclerosomatidae species in islands, and from the four species of Cosmetidae, we sampled just 7 specimens of *E. duapunctata* in only 5 islands, contrasting with 47 specimens sampled across all 5 mainland sites. Those results are quite uncommon because *E. duapunctata* followed by other species of the same genus and at least one or two species of Gagrellinae (Sclerosomatidae) are the most common, widespread and abundant species (Tourinho et al. 2014, 2017, Porto et al., 2016). Additionally, Cosmetidae, Sclerosomatidae are the top most diverse and abundant species/families in virtually all investigated sites in the Amazon (Bonaldo et al., 2009, Tourinho et al.,

2017, Tourinho et al., 2014, Porto et. al., 2016) and our results in mainland followed this pattern as expected in undisturbed areas.

The majority of harvestmen species are highly endemic and have poor disperser skills, tending to move a few dozen of meters during their lives (Mestre & Pinto-da-Rocha, 2004, Pinto-da-Rocha et al., 2007). Exceptions are some Eupnoi species, such as species of Sclerosomatidae, which exhibited higher vagility. Species of harvestmen are generally predators, and most of the species of Sclerosomatidae observed in the Amazon were predating on small insects (e.g. flies and beetles, Tourinho person. comm.), being more susceptible to habitat and other environmental structure and size than other trophic groups (Srivastava et al. 2008, Villanueva-Bonilla et al. 2017). Possibly they possess a closer relationship with the structure and architecture of some specific species of trees, palms and/or other plants in the Amazon forests, and because of their physiology and strong associations with plant microhabitat, it might be probably difficult for harvestmen species from forested environments in the Amazon to succeed in colonizing highly disturbed island areas in Balbina lake.

Conclusions

Arthropods play vital roles in various ecosystem functions, and are widely considered as good indicators of overall biodiversity and forest integrity (Maleque et al., 2006). Among the Arachnida, harvestmen comprise the third most diverse order (Shultz & Pinto-da-Rocha, 2007), exhibiting high sensitivity to habitat reduction and edge effects (Bragagnolo et al. 2007). Our study shows that forest loss significantly affected the species composition of harvestmen assemblages across islands created by a major hydroelectric dam, following 20 years of isolation. Although total species richness did not respond to local, patch and landscape variables, the assemblage structure of harvestmen was severely affected by landscape-scale forest amount and the rapid decay in tree diversity habitats in most small forest islands in this reservoir. Given the pivotal importance of microhabitat quality for harvestmen (Bragagnolo et al., 2007), we expect that a reduction in environmental heterogeneity is directly impacting several harvestmen species. Observed shifts in community structure can potentially lead to an impoverished forest functioning and integrity, especially because changes in tree diversity in insular forests at Balbina are expected to be aggravated in the future (Benchimol & Peres

2015b). We therefore provide additional evidence of the detrimental ecological consequences of dam impoundments on biodiversity, emphasising that dams are far from the often hailed green sources of energy (Gibson et al., 2017). For existing dams, we recommend that well-preserved and large forest islands, surrounded by large amounts of forest cover, should be protected in the long-term. This can buffer the effects of natural forest disturbance and preclude rapid decay in tree communities, contributing therefore to maintain high diversity of several faunal groups.

Acknowledgments

Adriano Kury provided key assistance with harvestmen type specimens, Pio Colmenares identified Laniatores specimens and tabulated species data; their support have made this work possible. Regiane Saturnino kindly made available local environmental data collected for her master thesis during arachnids sampling at the Balbina islands. This study was supported by Beca Program, from International Institute of Education of Brazil Programa Beca #B/2005/02/BMP/14, Reserva Biológica do Uatumã and a grant from the Coordination of Improvement of Higher Education Personnel - CAPES to ALT.

References

Acosta L.E., Pérez-González A., Tourinho A.L. 2007. Methods for taxonomic study. Harvestmen: The Biology of Opiliones. (ed. by Pinto-da-Rocha R., Machado G., Giribet G.), pp. 494–510. Harvard University Press, Cambridge, USA.

Alho, C.J.R., 2011. Environmental effects of hydropower reservoirs on wild mammals and freshwater turtles in Amazonia: a review. Oecol Australis. 15(3), 593–604.

Almeida, C.A., Valeriano, D.M.; Escada, M.I.S.; Rennó, C.D. 2010. Estimativa de área de vegetação secundária na Amazônia Legal brasileira. Acta Amazonica. 40. 289–302.

Andren, H., 1994. Effects of habitat fragmentation on birds and mammals in landscapes with different proportions of suitable habitat: a review. Oikos. 71(3), 355–366. DOI: 10.2307/3545823

Aurélio-Silva, M., Anciães, M., Henriques, L.M.P., Benchimol, M., Peres, C.A., 2016. Patterns of local extinction in an Amazonian archipelagic avifauna following 25 years of insularization. Biological Conservation. 199, 101–109. https://doi.org/10.1016/j.biocon.2016.03.016

Barton, K. 2018. MuMIn: Multi-Model Inference. R package version 1.40.4.

Benchimol, M., Venticinque, E.M. 2014. Responses of Primates to Landscape Change in Amazonian Land☐bridge islands—a Multi☐scale Analysis. Biotropica, 56(4), 470–478. https://doi.org/10.1111/btp.12122

Benchimol, M., Peres, C.A. 2015a. Widespread forest vertebrate extinctions induced by a mega hydroelectric dam in lowland Amazonia. PLoS ONE. 10, e0129818. https://doi.org/10.1371/journal.pone.0129818

Benchimol, M., Peres, C.A. 2015b. Edge-mediated compositional and functional decay of tree assemblages in Amazonian forest islands after 26 years of isolation. Journal of Ecology, 103, 408–420.

Benchimol, M., Peres, C.A. 2015c. Predicting local extinctions of Amazonian vertebrates in forest islands created by a mega dam. Biological Conservation, 187, 61–72.

Bobrowiec, P.E.D., Tavares, V.C. 2017. Establishing baseline biodiversity data prior to hydroelectric dam construction to monitoring impacts to bats in the Brazilian Amazon. PLoS One, 12(9), e0183036. 10.1371/journal.pone.0183036

Bonaldo, A.B., Carvalho, L.S., Pinto-da-Rocha, R., Tourinho, A., Miglio, L.T., Candiani, D.F., Lo-Man-Hung, N.F., Abrahim, N., Rodrigues, B.V.B, Brescovit, A.D., Saturnino, R.F., Bastos, N.C., Dias, S.C., Silva, B.J.F., Pereira-Filho, J.M.B., Rheims, C.A., Lucas, S.M., Polotow, D., Ruiz, G.R.S., Indicatti, R.P. 2009. Inventário e história natural dos aracnídeos da Floresta Nacional de Caxiuanã. In: Lisboa, P.L.B. (Ed.), Caxiuanã: Desafios Para a Conservação De Uma Floresta Nacional Na Amazônia. Museu Paraense Emílio Goeldi, Belém, pp. 545–588.

Bragagnolo, C., Nogueira, A.A., Pinto-da-Rocha, R., Pardini, R., 2007. Harvestmen in an Atlantic forest fragmented landscape: evaluating assemblage response to habitat quality and quantity. Biological Conservation, 139, 389–400.

Burnham, K.P., Anderson, D.R. 2002. Model Selection and Multimodel Inference: a Practical Information-theoretical Approach, 2nd edn. New York: Springer.

Carvalho, L.S., Gomes, J.O., Neckel-Oliveira, S., Lo-Man-Hung, N.F., 2012, Microhabitat use and intraspecific associations in the whip spider *Heterophrynus longicornis* (Arachnida: Amblypygi) in forest fragments formed by the Tucuruí Dam lake, Pará, Brazil. Journal of Natural History, 46, 1263–1272. https://doi.org/10.1080/00222933.2012.654515

Colmenares P.A, Baccaro, F.B., Tourinho, A.L. 2016. Assessing the Relationship of Vegetation Structure and Harvestmen Assemblage in an Amazonian Upland Forest. bioRxiv. doi:10.1101/078220

Curtis, D.J. 2007. Methods and Techniques of Study. In Harvestmen: The Biology of Opiliones.(ed. by Pinto-da-Rocha, R., Machado, G. & Giribet, G.). Harvard University Press, Cambridge, pp. 489–494.

Curtis, D.J. & Machado, G. 2007. Ecology. In Harvestmen: The Biology of Opiliones.(ed. by Pinto-da-Rocha, R., Machado, G. & Giribet, G.). Harvard University Press, Cambridge, pp. 280–308.

Dai, A., Trenberth, K. E., 2002. Estimates of Freshwater Discharge from Continents: Latitudinal and Seasonal Variations, J. Hydrometeorol., 3(6), 660–687, doi:10.1175/1525-7541(2002)003<0660:EOFDFC>2.0.CO;2

ECOA (Ecologia em Ação). 2018. International Rivers and Fundacion Proteger, Dams in Amazonia. Available online: <http://www.dams-info.org/en> (accessed 03.02.2018).

Eletronorte, Ibama. 1997. Plano de Manejo Fase I Reserva Biológica Uatumã. http://www.icmbio.gov.br/portal/images/stories/docs-planos-de-manejo/rebio_uatuma_pm.pdf

Esselman, P.C., Opperman, J.J. 2010. Overcoming information limitations for the prescription of an environmental flow regime for a central american river. Ecology and Society, 15(1), 6.

ESRI. 2016. ArcGIS Desktop: Release 10. Environmental Systems Research Institute, Redlands, CA.

Fahrig, L., 2003. Effects of habitat fragmentation on biodiversity. Annu. Rev. Ecol. Evol. Syst. 34, 487–515.

Fahrig, L., 2013. Rethinking patch size and isolation effects: the habitat amount hypothesis. J. Biogeogr. 40, 1649–1663

Fearnside, P.M. 2016. Environmental and Social Impacts of Hydroelectric Dams in Brazilian Amazonia: Implications for the Aluminum Industry. World Development, 77, 48–65. https://doi.org/10.1016/j.worlddev.2015.08.015

Franken, E.P., Baccaro, F., Gasnier, T.R., 2013. Is there a refuge for ants in litter accumulated at the base of *Attalea attaleoides* (Barb. Rodr.) Wess. Boer (Arecaceae)? Entomotropica 28, 27–37.

Gibson, L., Wilman, E.N., & Laurance, W.F. 2017. How Green is 'Green'Energy?. Trends in Ecology & Evolution, 32(12), 922–935 https://doi.org/10.1016/j.tree.2017.09.007

Heiberger, R. M. (2016). HH: Statistical analysis and data display: Heiberger and Holland.–R package ver. 3.1-31.

Kury, A.B. 2003. Annotated catalogue of the Laniatores of the New World (Arachnida, Opiliones). Revista Ibérica de Aracnología, 1, 1–337.

Laurance, W. F., Nascimento, H.E.M., Laurance, S.G., Andrade, A.C.,Fearnside, P.M., Ribeiro, J. E. L., Capretz, R. L. 2006. Rain forest fragmentation and the proliferation of successional trees. Ecology, 87(2), 469–82. https://doi.org/10.1890/05-0064

Lees, A.C., Peres, C.A., Fearnside, P.M., Schneider, M., & Zuanon, J.A. 2016. Hydropower and the future of Amazonian biodiversity. Biodiversity and conservation, 25(3), 451–466. https://doi.org/10.1007/s10531-016-1072-3

Liermann C.R., Nilsson C., Robertson J., NG, R.Y. 2012. Implications of dam obstruction for global freshwater fish diversity. Bioscience. 62, 539–548.

Lira, P.K., Ewers, R.M., Banks⸆Leite, C., Pardini, R. and Metzger, J.P., 2012. Evaluating the legacy of landscape history: extinction debt and species credit in bird and small mammal assemblages in the Brazilian Atlantic Forest. Journal of Applied Ecology, 49(6), 1325-1333.

MacArthur, R. H. & Wilson, E. O.1967. The Theory of Island Biogeography. Princeton Univ. Press, Princeton.

Maleque, M.A., Ishii, H.T., & Maeto, K. 2006. The use of arthropods as indicators of ecosystem integrity in forest management. Journal of Forestry, 104(3), 113–117.

Matthews, T. J., Guilhaumon, F., Triantis, K.A., Borregaard, M.K., & Whittaker, R.J. 2016. On the form of species-area relationships in habitat islands and true

islands. Global Ecology and Biogeography, 25(7), 847–858. https://doi.org/10.1111/geb.12269

McGarigal, K., Cushman, S.A. & Ene, E. (2012) F FRAGSTATS v4: Spatial Pattern Analysis Program for Categorical and Continuous Maps. Computer software program produced by the authors at the University of Massachusetts, Amherst. Available from http://www.umass.edu/landeco/research/fragstats/fragstats.html (accessed January 2014).

Melo, T.S., Peres, M.C.L, Andrade, A.R.S., Oliveira, M.L.T., Tinoco, M.S. 2015. Artrópodes Terrestres no Licenciamento Ambiental: Um Modelo Para Utilização Por Empresas De Consultoria. Latin American Journal of Business Management. 6(3), 126–144.

Messas, Y., Souza, H., Gonzaga, M., Vasconcellos-Neto, J., 2014. Spatial distribution and substrate selection by the orb-weaver spider Eustala perfida Mello-Leitão, 1947 (Araneae: Araneidae). J. Nat. Hist. 48, 2645–2660.

Mestre, L.A.M., Pinto-da-Rocha, R. 2004. Population biology of the harvestmen *Ilhaia cuspidata* (Opiliones; Gonyleptidae) in an Araucaria Forest fragment (Curitiba - Paraná, Brazil). The Journal of Arachnology, 32(2), 208–220.

Morante⬚Filho, J.C., Arroyo⬚Rodríguez, V., Lohbeck, M., Tscharntke, T. and Faria, D., 2016. Tropical forest loss and its multitrophic effects on insect herbivory. Ecology, 97(12), 3315-3325.

Morante-Filho, J.C., Faria, D., Mariano-Neto, E. and Rhodes, J., 2015. Birds in anthropogenic landscapes: the responses of ecological groups to forest loss in the Brazilian Atlantic Forest. PLoS One, 10(6), p.e0128923.

Nilsson, C., Reidy, C.A., Dynesius, M., Revenga, C. 2005. Fragmentation and Flow Regulation of the World's Large River Systems. Science, 308, 405–408. 10.1126/science.1107887

Palmeirim A.F., Peres C.A., Rosas F.C.W. 2014. Giant otter population responses to habitat expansion and degradation induced by a mega hydroelectric dam. Biol Conserv. 174, 30–38. https://doi.org/10.1016/j.biocon.2014.03.015

Palmeirim, A.F., Benchimol, M., Vieira, M.V., Peres, C.A. 2018. Small mammal responses to Amazonian forest islands are modulated by their forest dependence. Oecologia, 187, 191–204. https://doi.org/10.1007/s00442-018-4114-6

Pinto-da-Rocha, R. 1994. A new species of Santinezia from central Amazonia, with systematic comments on the genus (Arachnida, Opiliones, Cranaidae). Andrias, 13, 29–32.

Pinto-da-Rocha, R. 1995. Redescription of Stenostygnus Pusio Simon and synonymy of Caribbiantinae with Stenostygninae (Opiliones:Laniatores, Biantidae). The Journal of Arachnology, 23, 194–198.

Pinto-da-Rocha, R. 1997. Systematic review of the Neotropical family Stygnidae (Opiliones, Laniatores, Gonyleptoidea). Arquivos de Zoologia, 33, 163–342.

Pinto-da-Rocha, R. 2004. A new species of Fissiphalliidae from Brazilian Amazon Rain Forest (Arachnida: Opiliones). Zootaxa, 6, 1–6.

Pinto-da-Rocha, R. & Bonaldo, A.B. 2006. A structured inventory of harvestmen (Arachnida , Opiliones) at Juruti River plateau , State of Pará , Brazil. Revista Ibérica de Aracnología, 13, 155–162.

Pinto-da-Rocha, R., Machado, G, Giribet, G. 2007. Harvestmen: the Biology of Opiliones. 2007. Harvard University Press. Cambridge, Massachusetts. 597 pp.

Pinzón, J., Spence, J., 2010. Bark-dwelling spider assemblages (Araneae) in the boreal forest: dominance, diversity, composition and life-histories. J. Insect Conservat. 14, 439–458.

Porto, W., Pequeno, P.A.L, Tourinho, A.L. 2016. When less means more: Reduction of both effort and survey methods boosts efficiency and diversity of harvestmen in a tropical forest. Ecological Indicators 69, 771–779. https://doi.org/10.1016/j.ecolind.2016.05.014

Proud, D.N., Felgenhauer, B.E., Townsend, V.R., Osula, D.O., Gilmore, W.O., Napier, Z.L., Van Zandt, P.A. 2012. Diversity and Habitat Use of Neotropical Harvestmen (Arachnida: Opiliones) in a Costa Rican Rainforest. ISRN Zoology, 2012, 1–16.

Resende, L.P.A., Pinto-da-Rocha, R. & Bragagnolo, C. 2012. Diversity of harvestmen (Arachnida , Opiliones) in Parque da Onça Parda , southeastern Brazil. Ilheringia, 102, 99–105.

Shultz, J. W. & Pinto-da-Rocha, R. 2007. Morphology and functional anatomy. In: Pinto-da-Rocha, R.; Machado, G. & Giribet, G. eds. Harvestmen: The Biology of Opiliones. Cambridge, Harvard University Press. p.14-61.

Silva, J.M.C., Tabarelli, M. 2000. Tree species impoverishment and the future flora of the Atlantic forest of northeast Brazil, Nature, 404(6773), 72–74. DOI: 10.1038/35003563

Sørensen, L.L., Coddington, J., Scharff, N., 2002. Inventorying and estimating subcanopy spider diversity using semiquantitative sampling methods in an afromontane forest. Environ. Entomol. 31(2), 319–330. http://www.bioone.org/doi/abs/10.1603/0046-225X-31.2.319.

Southwood, L., 1978. Ecological Methods: with Particular Reference to the Study of Insect Populations. Chapman and Hall, London, U.K.

Srivastava, D.S., Trzcinski, M.K., Richardson, B.A., Gilbert, B. 2008. Why Are Predators More Sensitive to Habitat Size than Their Prey? Insights from Bromeliad Insect Food Webs. the american naturalist. The American Naturalist, 172 (6), 761–771.

Storck-Tonon, D., Peres, C.A. 2017. Forest patch isolation drives local extinctions of Amazonian orchid bees in a 26 years old archipelago. Biological Conservation. 214, 270–277. https://doi.org/10.1016/j.biocon.2017.07.018

Tabarelli, M., Peres, C.A., & Melo, P.L. 2012. The 'few winners and many losers' paradigm revisited: Emerging prospects for tropical forest biodiversity. Biological Conservation, 155, 136–140.

Tarli, V.D., Pequeno, P.A.C.L., Franklin, E., Oliveira, J. & Guilherme, D.R. 2014. Multiple environmental controls on cockroach assemblage structure in a tropical rain forest. Biotropica. 46, 598–607. https://doi.org/10.1111/btp.12138

Terborgh, J., Lopez, L., Tello, J., Yu, D., Bruni, A.R. 1997. Transitory states in relaxing ecosystems of land bridge islands. In W. F. Laurance, and R. O. Bierregaard Jr (Eds.). Tropical forest remnants – Ecology, management and conservation of fragmented communities, pp. 256– 274. University of Chicago Press, Chicago, Illinois.

Tourinho, A.L., Saturnino, R., Dias, S.C., Guimarães, S., Assimen, S. & Colmenares, P.A. 2011. Aranhas e opiliões da Reserva de Desenvolvimento Sustentável do Tupé, Amazonas - Brasil. In BioTupé: Meio Físico, Diversidade Biológica e Sociocultural do Baixo Rio Negro, Amazônia Central(ed. by Scudeller, V.V.). Manaus, pp. 271–289.

Tourinho, A.L., Lança, L.S., Baccaro, F.B., Dias, S.C., 2014. Complementarity among sampling methods for harvestman assemblages. Pedobiologia, 57(1), 37–45. https://doi.org/10.1016/j.pedobi.2013.09.007

Tourinho, L.S., Dias, S.C., Lo Man Hung, N.F., Bonaldo, A.B., Pinto-da-Rocha, R., Baccaro, F.B. 2017. Optimizing survey methods for spiders and harvestmen assemblages in an Amazonian upland forest. Pedobiologia, 67, 35–44. https://doi.org/10.1016/j.pedobi.2017.11.002

Vasconcelos, H.L. 1990. Effect of litter collection by understory palms on the associated macroinvertebrate fauna in Central Amazonia. Pedobiologia, 34, 157–160.

Villanueva-Bonilla, G.A., Salomão, A.T, Vasconcellos-Neto, J. 2017. Trunk structural traits explain habitat use of a tree-dwelling spider (Selenopidae) in a tropical forest. Acta Oecologica, 85, 108–115. https://doi.org/10.1016/j.actao.2017.10.004

Yang, L.H., Gratton, C. 2014. Insect as drivers of ecosystem processes. Insect Science, 2, 26–32. https://doi.org/10.1016/j.cois.2014.06.004

Caption of Figures

Figure 1. Spatial distribution of the 20 sampled islands (red) and 5 continuous forest sites (black circles) within the Balbina Hydroelectric Reservoir of central Brazilian Amazonia.

Figure 2. Harvestmen species registered at the Balbina Hydroelectric Reservoir - BHR: a) Female of *Phareicranaus manauara* (Cranaidae) caring for egg clutches; b) Male of *Saramacia lucaseae* (Manaosbiidae), c) Male of *Eucynortella duapunctata* (Cosmetidae); d) Male of *Protimesius longipalpis* (Stygnidae); e) Male of *Fissiphallius martensi* (Fissiphalliidae); f) Male of *Avima matintaperera* (Agoristenidaae); g) Female of *Stygnus pectinipes* (Stygnidae); h) Female of *Discocyrtus carvalhoi* (Gonyleptidae).

Figure 3. Nonmetric multidimensional scaling (NMDS) ordination using a Jaccard similarity matrix of harvestman composition in islands (gray circles) and mainland forest (green circles) sampled across the Balbina archipelago. Circles are sized in proportion to the area of sampled sites.

Figure 4. Relationships between the first non-metric multidimensional scaling (NMDS) axis representing the harvestmen species composition and forest cover within a 1000 m buffer. Circles proportional to the log area.

Figure 5. Site records of 31 harvestman species sampled across 25 trapping sites distributed throughout the Balbina Hydroelectric Reservoir. Species occupancies within island and mainland forest sites are represented by gray and green rectangles, respectively.

FIGURE 1

FIGURE.2

FIGURE 3

FIGURE 4

Table 1. Description of the explanatory variables used in the analyses.

Variable name	Codenames	Scale	Description
Litter depth	litdepth	Local	
Density of trees	$D_{mainland}$	Local	
Number of palm trees	palms	Local	
Number of fallen stems	fallen	Local	
Island size	area	Patch	Total island area of each focal island. In $\log_{10}$ x hectares.
Isolation	$D_{mainland}$	Patch	Shortest linear distance from each island to the nearest mainland. In meters.
Forest cover	cover	Landscape	Forest cover amount within a buffer. Buffer considered in the final analysis: 500 m. In percentage.

			Sum of all island areas divided by the
Proximity	prox	Landscape	squared edge-to-edge distances from
			each focal island to all islands within
			a specified buffer. Instead of
			considering the area of each island
			within the buffer (as in McGarigal *et*
			al. 2012), we considered the total
			area of each island. We assigned a
			value one order of magnitude greater
			than our largest island for every
			mainland continuous forest site
			included within a focal island buffer
			area. Buffer considered in the final
			analysis: 500 m. In meters.

Table 1. Results of the Generalized Linear Models for the effects of island area ($\log_{10}$), litter depth, fallen woody stems, palm trees, forest cover (buffer 1000m) and vegetation density. Only the best models are shown (ΔAICc <2).

Response variables	Best Models	AICc	ΔAICc	Akaike weights
	NULL	88.7	0.00	0.116
Species Richness				

	Litter depht	89.4	0.77	0.070
Species composition	Forest cover	23.8	0.00	0.244
	Forest cover + fallen woody stems	25.4	1.60	0.109
Species abundance	Vegetation density + Forest cover + Area+ Palm trees + Fallen woody stems	220.7	0.00	0.253
	Forest cover + Area+ Palm trees+ Fallen woody stems	220.8	0.10	0.241
	Vegetation density + forest cover + area + palm trees	221.3	0.62	0.186

Table 2. Number of individuals of each harvestmen species recorded at the Balbina Hydroelectric Reservoir

Species	Mainland	Island	Total
AGORISTENIDAE			
Avima matintaperera (Pinto-da-Rocha, 1996)		8	8
COSMETIDAE			
Cosmetidae1 sp.1	42		42

Cosmetidae1 sp.2	13		13
Eucynortella duapunctata Goodnight & Goodnight, 1943	41	7	48
Eucynortella sp.1	44		44
FISSIPHALLIIDAE			
Fissiphallius martensi		4	4
Fissiphallius sp.1		4	4
Fissiphallius sp.2	2		2
GONYLEPTIDAE			
Discocyrtus carvalhoi Mello-Leitão, 1940	5	2	7
MANAOSBIIDAE			
Manaosbia sp.1	17	120	137
Manaosbiidae1 sp.1	4	58	62
Manaosbiidae2 sp.1	2		2
Saramacia lucasae Jim & Soares, 1991	1		1
SCLEROSOMATIDAE			
Sclerosomatidae1 sp.1	48		48
Sclerosomatidae1 sp.2	58		58
Sclerosomatidae2 sp.1	79		79
Sclerosomatidae2 sp.2	2		2
Sclerosomatidae3 sp.1	4		4
Sclerosomatidae4 sp.1	3		3
STYGNIDAE		3	3
Auranus sp1			
Auranus xerxes Colmenares, Porto & Tourinho	1	16	17
Imeri sp1		2	2

Protimesius longipalpis Roewer, 1943	23		23
Stygnidae1 sp.1	12	88	100
Stygnus armatus Perty, 1833	14	18	32
Stygnus pectinipes (Roewer, 1943)		3	3
Stygnus sp.1	9	20	29
STYGNOMMATIDAE			
Stygnomma sp1		1	1
ZALMOXIDAE			
Zalmoxidae1 sp.1		1	1
Zalmoxidae2 sp.1	2	1	3
Zalmoxidae3 sp.1		1	1
Total	426	357	783

Table S1. Number of individuals and species of harvestmen (Arachnida: Opiliones) sampled across 20 islands and 5 continuous forest sites across the Balbina Hydroelectric reservoir, Central Brazilian Amazonia.

Site name	Area (ha)	UTM (X)	UTM (Y)	Number of species	Number of individuals
Angelin	67.44	240659	9801730	4	9
Cachoeira	11.84	236641	9793981	5	20
Cipoal	213.66	190871	9810812	5	17
Copa	19.20	229635	9798056	4	13

Folharal	102.30	192046	9806910	6	17
Galo de briga	190.84	197368	9807612	3	9
Gemeas	111.09	203037	9814880	6	38
Grande	95.61	204026	9806251	6	48
Igarapé	76.20	236013	9805808	4	10
Josue	146.00	205982	9801462	6	24
Manoel gato	194.59	223583	9802328	6	18
Massaranduba	17.33	201171	9802527	7	25
Mucura	22.57	229798	9800932	1	1
Neto	41.59	237861	9796638	4	5
Palhal 1	20.78	237245	9800112	2	5
Palhal 2	25.81	227557	9801956	6	16
Pontal	126.01	201747	9797850	8	27
Santa Luzia	218.53	228363	9808143	3	4
Serrinha	50.09	231033	9792680	2	2
U	79.83	204181	9799600	10	49
Mainland 1	∞	251524	9801136	14	89
Mainland 2	∞	250493	9801920	11	55
Mainland 3	∞	247494	9800939	15	67
Mainland 4	∞	251567	9798861	20	168
Mainland 5	∞	251564	9797900	11	47
TOTAL				31	783