A proposed microcosm for landscape ecology – beyond the binary to the patch-

mosaic model

Yolanda F. Wiersma[1], Rachel D. Wigle[1], R. Troy McMullin[2]

[1]Department of Biology, Memorial University of Newfoundland, St. John's, NL, A1B 3X9,

Canada.

[2]Canadian Museum of Nature, Research and Collections, Station D, PO Box 3443, Ottawa, ON,

K1P 6P4, Canada.

*Author for correspondence: ywiersma@mun.ca; ORCID: 0000-0003-4604-9240

Abstract

Background. Microcosms such as pitcher plants, or patches of mosses on a rock surface, have

been used worldwide to allow for manipulative experiments that test hypotheses for patterns

observed at larger extents, such as dispersal or community assemblage. Such microcosms can

also be applied to questions in landscape ecology, but are limited by their binary (patch/non-

patch structure). Here we examine a more realistic model landscape system that shares the

patch-mosaic structure common to kilometres-extent landscapes. This system of lichen thalli on

tree trunks has been shown to have consistent spatial patterns across replicate microcosms, but

only when sampling within a limited area. To be relevant for experimentation across scales, it is

necessary to determine whether previously observed patterns are consistent when sampling

across a broader region and when using different tree species. Here, we test for consistent

landscape patch pattern in both maco- and micro-lichens across 21 balsam fir (*Abies balsamea*)

and yellow birch (*Betula alleghaniensis*) trees.

Methods. We measured spatial pattern of lichen thalli along the trunks of two species of trees,

at two spatial resolutions; trees within a single stand (~100 x 50 m) and trees dispersed across a

larger region (500 km²). We used a "lichen ladder" comprised of 5 10 x 10 cm sampling blocks to

quantify number of species and individuals in a 50 cm section of the tree trunk. We tested for

similar patterns along the trunk and between the north and south sides at both sampling

intensities using perMANOVA.

Results. We find that lichen patches on tree trunks can function as replicate microscoms for

landscape ecology. Patterns of thalli along the trunks of trees and between the north and south

aspects of the trunk are statistically significantly consistent, although there is variation between

tree species, and groups of lichens included. Our microcosm could be used as a model system for

landscape ecology research; but researchers should test for consistent patterns first.

Introduction

Research in ecology has always been challenged with the trade-off between experimental rigour

and real-world applicability (Schindler 1998; Kohler 2002). The tenets of good experimental

design (control, randomization and replication) are done most easily through manipulative

experiments in a lab setting, while observational studies allow for natural processes to remain

present, at the expense of the experimental rigour (Kohler 2002). A widely-used solution to

bridge this divide is through the use of mesocosms (Stewart et al. 2013) or microcosms

(Srivastava et al. 2004). *Mesocosms* and *microcosms* (see Box 1 for glossary) share the property

of being contained systems, usually small enough to allow for adequate replication for statistical

power and that can be experimentally manipulated. The distinction between the two is that

mescocosms are artificially constructed while microscoms (the focus of this paper) occur

naturally (Box 1). Commonly used microcosms include pitcher plants (e.g., Mouquet et al.

2008), bromeliads (e.g., Talaga et al. 2015), patches of mosses (e.g., Gonzalez et al. 1998) and

rocky tidepools (e.g., Schiesari et al. 2018). These are sometimes termed "*microecosystems*" in

both lab (Drake 1993) and field (Gilbert et al. 1998; Lau et al. 2018) contexts; however in this

paper we will take natural microecoysystems as equivalent to microcosms (Box 1). In a review,

Srivastava et al. (2004) suggest that natural microcosms are potentially highly powerful tools

that can advance ecological research questions but caution that the utility of microcosms is

limited to the types of questions applied (Srivastava et al. 2004). In addition, Srivastava et al.

(2004) and others (Zuk and Travisano 2018) emphasize that microcosms should be viewed as

complementary tools to theoretical models, lab experiments and field observations.

Ecological research using microcosms has mainly focused on questions in community

and population ecology and has successfully used them as a tool to examine topics such as

dispersal (e.g., Grainger and Gilbert 2016), predator-prey dynamics (e.g., Busse et al. 2019) and

community assemblage (e.g., Pereira et al. 2018). Although the microcosm as a whole is often

referred to as a "landscape" (Drake 1993; Gilbert et al. 1998), the experimental systems

described above are all simple binary landscapes of habitat and non-habitat patches (i.e., pitcher

plant leaves and the surrounding air, tide pools and the surrounding rocks). Even though

experimental manipulation of moss patches (Gilbert et al. 1998; Gonzalez et al. 1998; Gonzalez

and Chaneton 2002) contained three species of bryophytes, these different species were treated

as functionally equivalent patches in the experiments. Such simplification can be appropriate,

depending on the research question; however a binary patch/non-patch landscape is an over-

simplification of the way landscape ecologists view the world (Box 1), and may explain why

microcosms have mainly been used for questions in population and community ecology and not

yet expanded widely to landscape ecology (but see Jenerette and Shen 2012).

Landscape ecologists commonly view their object of study using the "patch-mosaic"

model, that is, that *landscapes* are comprised of a diversity of *patch* types, each of which has its

own intrinsic properties of size, shape and cover, but which interacts with adjacent patterns to

form a *landscape mosaic* (Box 1; Wu 2013). Thus, while the binary microcosms of moss patches,

bromeliads or tide pools have been shown to be good model systems for some processes that

occur at larger extents (e.g., species dispersal and assembly on oceanic islands as predicted by

island biogeography theory), they may not function well as model systems for more complex

landscapes.

There have not been many developments of microcosms for landscape ecology

experiments, although experimental mesocosms have been proposed as a tool (e.g., Wiens and

Milne 1989). Model systems that have been proposed for landscape ecology include soil

biocrusts (Bowker et al. 2014) and lichen-covered tree trunks (Wiersma and McMullin 2018).

Bowker et al. (2014) point out that soil biocrusts are most like scaled-down landscapes in that

they have similar attributes to kilometers-extent landscapes, such as patch heterogeneity.

Moreover, their small size makes them tractable for manipulative experiments with good

potential for experimental replicates, and they can also be collected and transferred into

controlled environments (e.g., greenhouses) for increased experimental rigour. At the same

time, they are interesting ecological systems independent of their use as model systems, which is

an often-overlooked property of models (Zuk and Travisano 2018). Wiersma and McMullin

(2018) proposed a similar system of heterogeneous patches of cryptogams as Bowker et al.

(2014), but growing along the boles of trees as replicate landscapes. Although Bowker et al.'s

(2014) system and Wiersma and McMullin's (2018) share similar types of organisms and patch

pattern, Wiersma and McMullin's (2018) has the additional feature of being explicitly

hierarchical (Fig. 1) with the disadvantage relative to Bowker et al.'s (2014) of not being

transferrable to a lab setting.

Hierarchy is a key concept in landscape ecology (Urban et al. 1987; O'Neill et al. 1989;

Kotliar and Wiens 1990) and it has been suggested that by harnessing principles of hierarchy

and scaling (Urban et al. 1987; Urban 2005; Bowker et al. 2014), that microlandscape patterns

and processes (a level at which experimentation is more feasible) can be extrapolated to larger

extents (a level at which many management challenges occur). Wiersma and McMullin (2018)

demonstrated that 2 sets of 12 balsam fir (*Abies balsamea*) trees within a very small area (< 100

m apart) share similar patch patterns of macro-lichens along the boles, and proposed that these

trees could be considered replicate (micro)landscapes. While this system holds promise as a

potential new type of microcosm for landscape ecology, the limited focus of a single species of

tree in a restricted geographic area limits the broader applicability of their proposed model

system. Here, we use additional data – from two species of trees and across a much wider extent

(500 km²) and with a broader suite of lichens, to scrutinize Wiersma and McMullin's (2018)

hypothesis that patches of lichens along a tree bole are a model system for landscapes. If the

patch-patterns along the boles of trees holds true in this expanded study, then we posit that

similar systems in other regions might also function as complex landscape microcosms that

could be harnessed for experimentation.

Methods

Study area

We carried out sampling in the Avalon Forest Ecoregion on the island of Newfoundland,

Canada. The Avalon Forest is the smallest (500 km²) ecoregion in the province of Newfoundland

and Labrador, characterized by high humidity and precipitation, cool summers, and mild

winters (South 1983). The forested areas are on rolling hills called ribbed moraines, landscape

features created by glaciers (Hättestrand and Kleman 1999). Interspersed between the moraines

are open, sphagnum-dominated wetlands (South 1983). Forests are dominated by balsam fir

(*Abies balsamea*) with black spruce (*Picea mariana*) in wet areas and occasional yellow birch

(*Betula alleghaniensis*) stands on north facing slopes (South 1983).

Field work was carried out under Permit Nos. 2015/16-14 and 2017/18-10 issued by the

Department of Environment and Conservation, Government of Newfoundland and Labrador.

 Experimental Design

We visited 22 sites across the region (Fig. 2) and at 21 of these, selected two trees, one balsam fir

and one yellow birch that were similar in diameter and within 25 m of each other. The

remaining site (indicated as "Halls Gullies" on Fig. 2) was the original location intensively

sampled (24 trees total) by Wiersma and McMullin (2018). All sampling locations were balsam-

fir dominated stands, mostly occurring on moraines. On each tree, we sampled on the north and

south sides of the bole using a 10 cm x 50 cm "lichen ladder", divided into five 10 x 10 cm

"blocks", placed from 1.1 m to 1.6 m up the bole (Fig. 3). We identified and counted all lichen

species within each 10 cm block. Species that could not be identified in the field were collected

for identification using standard processes, including microscopy, chemical spot tests (Brodo et

al. 2001), and thin-layer chromatography (Culberson and Kristinsson 1970). We inventoried

both macro- (those with a more three-dimensional growth from, and growing on the substrate,

which includes foliose and fruticose growth forms) and micro-lichens (those growing within the

substrate, i.e., crustose growth forms) at the 21 new sites; Wiersma and McMullin (2018) limited

their study to field-identifiable macro-lichens. Thus, we analyzed the data from the 21 dispersed

sites with macro-lichens only to compare to the previous study; we also repeated the analysis for

more dispersed trees using data on both macro- and micro-lichens to assess whether the

landscape patterns observed by Wiersma and McMullin (2018) held when examining a broader

suite of lichen species and when looking across a larger sampling area.

Statistical analysis

We used a perMANOVA analysis (Anderson 2001) to assess whether the pattern of lichen

patches along the bole was consistent across all trees when stratifying for aspect, and whether

the patterns between the north and south sides were consistent when stratifying by height up

the bole. We separately analysed the data from the 21 balsam fir and the 21 yellow birch trees in

the expanded study area and did not combine the data from the 21 balsam fir in the wider

ecoregion with the data from the 24 balsam fir in the single stand because of differences in

geographic sampling intensity. We carried out all statistical analysis using R (version 1.0.136; R

Core Team 2016) with the package vegan (Oksanen et al. 2016).

3. Results

The 24 balsam fir trees in the single stand showed a significantly consistent lichen patch pattern

between the north and south sides along the 50 cm portion of the bole, when stratifying by tree

(perMANOVA R^2 = 0.00552, p = 0.05) but not when we controlled for position along the bole

(perMANOVA R^2 = 0.00552, p = 0.223). There was also a significant pattern for position along

the bole when controlling for the tree (perMANOVA R^2 = 0.001727, p = 0.024) but not for

position along the bole when we stratified for aspect (perMANOVA R^2 = 0.01727, p = 0.412).

The perMANOVA results for the 21 more spatially dispersed balsam fir trees also showed

a significant pattern for macro-lichens between the north and south sides of the tree, when

stratifying by tree (perMANOVA R^2 = 0.00988, p = 0.009) but not when controlling for position

along the bole (perMANOVA R^2 = 0.0098, p = 0.081). Unlike for the trees in the single stand,

there was no significant pattern for position along the bole when controlling for the tree

(perMANOVA R^2 = 0.00476, p = 0.157) nor for position along the bole when stratified for aspect

(perMANOVA R^2 = 0.00476, p = 0.444). When examining a different tree species, yellow birch,

there was no significantly consistent pattern of lichen between the north and south sides of the

bole, either when stratifying by tree (perMANOVA R^2 = 0.00147, p = 0.772) or by position along

the bole (perMANOVA R^2 = 0.00147, p = 0.85). Nor was there any significant pattern along the

bole of the yellow birch when stratifying by tree (perMANOVA R^2 = 0.00143, p = 0.721) or when

controlling for aspect (perMANOVA R^2 = 0.00143, p = 0.811). Overall patterns for macro-

lichens are summarized in Table 1.

When we looked at both macro- and micro-lichens, the patterns were different. Balsam

fir did not show any significant pattern (aspect stratified by tree perMANOVA R^2 = 0.00497 p =

0.051; aspect stratified by position along bole perMANOVA R^2 = 0.00497, p = 0.439; position

along bole stratified by tree perMANOVA R^2 = 0.00245, p = 0.471; position along bole stratified

by aspect perMANOVA R^2 = 0.00245, p = 0.843). In contrast, there was a significant pattern for

aspect for yellow birch, both when stratifying by tree (perMANOVA R^2 = 0.01528, p = 0.001)

and by position along the bole (perMANOVA R^2 = 0.01528, p = 0.002). However, there was not

a significant pattern for location up the bole for yellow birch, neither when stratifying by tree

(perMANOVA R^2 = 0.00373, p = 0.297) nor by aspect (perMANOVA R^2 = 0.00373, p = 0.647).

Table 2 summarizes the overall patterns for macro- and micro-lichens combined.

Discussion

The consistent patterns of patches of lichens growing along tree boles, even when trees are

widely dispersed (and presumably exposed to different micro-climate and stand-level

constraints) suggest that these microecosystems can be considered a model system for

landscape ecology. Macro-lichen patch pattern on the north vs. south sides of the bole of balsam

fir trees that were widely dispersed followed that of the more aggregated trees. Although the

pattern along the gradient of the bole was non-significant for the dispersed trees, we believe this

is due to the fact that the microtransect we used here was only 50 cm in length. Wiersma and

McMullin (2018) detected consistent patch patterns along a 1 metre gradient of a tree bole, and

(Young et al. 2018) documented gradients in both epiphyte diversity and microfauna along

gradients running 25 m up Douglas fir (*Pseudotsuga menziesii*) trees. Landscape patterns and

gradients in kilometer-extent landscapes are only detected when measured at an appropriate

scale, and likely 50 cm along the bole was insufficient to robustly detect any gradient in the 21

dispersed trees. We did, however, detect a statistically significantly consistent pattern along a 50

cm gradient of the bole in the 24 more clumped trees (where Wiersma and McMullin (2018)

detected an even stronger pattern along a 1 m gradient), which increases our confidence that a

pattern does exist in the more dispersed trees, we were simply unable to detect it.

While macro-lichens growing on the balsam fir trees appear to have consistent patch

patterns, and thus suggest these as microcosms amenable to experimentation, this proposed

system should be approached cautiously. Macro-lichens along the bole of yellow birch trees did

not show any consistent pattern. However, when we included both macro- and micro-lichens,

yellow birch did show consistent patch patterns on the north vs. south sides of the boles.

Conversely, patterns on balsam fir disappeared when both macro- and micro-lichens were

included. Thus, researchers wishing to adopt systems of cryptogams growing along tree boles

should first assess which community (i.e., just macro-lichens, or macro- and micro-lichens

combined), growing on which tree species, has a consistent spatial pattern and hence can

function as a landscape microcosm.

Despite that not all lichens growing on both species of trees showed a consistent spatial

pattern, we believe this microcosm of lichens along a tree bole holds promise for research. As

with other natural microcosms, this system allows for experimental replicates to test different

kinds of ecological questions. The complexity of multiple patch types and patch patterns allows

for different kinds of questions than those afforded by the binary patch/non-patch landscapes of

most microcosm work to date. Some questions that could be addressed include the spatial

distribution of microfauna (e.g., Bokhorst et al. 2015; Trekels et al. 2017; Young et al. 2018) or

more complex "island biogeography" questions (e.g., Southwood and Kennedy 1983). Srivastava

et al. (2004) suggested four future research questions that could be addressed with natural

microcosms; the patch patterns of lichen-covered tree trunks would be especially relevant to

question about how species characteristics and interactions drive patterns at spatial scales

(Srivastava et al. 2004). The hierarchical nature of this proposed model system will also yield

insights into how ecological processes change with system size (Srivastava et al. 2004). Research

on meta-communities is an exciting and expanded area (Logue et al. 2011), but often lacks a

spatially explicit approach. The addition of a microcosm with a spatially explicit pattern adds a

novel tool to both the community and landscape ecologist's tool box.

Potential limitations to this system are that, unlike the soil crusts proposed as a model

system for community and landscape ecology by Bowker et al. (2014), trees cannot be

transported to a lab for experimentation under controlled conditions. Moreover, the slow

growth of lichens may make the response time to certain manipulations (e.g., experimental

fragmentation) too slow to be practical for most studies. However, some lichens are relatively

fast growing (Phillips 1969; Faegri 1980; Bidussi and Gauslaa 2015) and thus by carefully

choosing manipulations, this system has the potential to contribute a novel type of microcosm

that replicates the patterns and processes observed (but not amenable to experimentation) in

kilometers-extent landscapes.

Our finding here, that spatial patterns in a natural microcosm are reasonably consistent

for trees that are widely dispersed, and that the landscape patch-mosaic model appears to be

hierarchical, has advantages for other kinds of research questions. Previously, Wiersma and

McMullin (2018) exploited these microcosms in a single, spatially-constrained stand to test

whether particular features of these micro-landscapes influenced presence or absence of a rare

species. In this case, controlling for stand-level constraints was helpful to eliminate other larger-

extent factors that might influence rare species persistence. However, the ability to replicate

microcosms across a broader region allows for natural-experiments to see how landscapes

respond to meso-scale conditions such as distance to different types of habitat (e.g., open bogs),

or broader gradients in elevation and climate. It also allows for manipulative experiments at

larger extents that might mimic real-world processes, for example, looking at responses to

different disturbance levels.

Conclusion

Thinking of the microcosm of epiphytes growing along a tree trunk as a model system for

landscapes opens up a whole suite of research opportunities not available in the binary

microcosms that have been the norm to date. The replicated patch-pattern allows for these

microcosms to be treated as micro-landscapes and thus have replicate landscape units to test

spatially-explicit questions; replication that is not possible with kilometers-extent landscapes.

Such a system is not amenable, or even necessary, for all types of research questions;

parsiomony should dictate when a binary landscape is sufficient for hypothesis testing. At the

same time, our system is interesting in and of itself; something that Zuk and Travisano (2018)

emphasize should not be overlooked with model systems. Micro-fauna and flora are often

incredibly diverse, and not always well-understood. Thus, we encourage further research on

these fascinating microcosms which occur in forests worldwide.

Acknowledgements

Thanks are due to T. Sato, P. Lauriault, J. Godfrey, M. Fahmy, E. Kissler, T. Padgett, G.

Wachinger, M.Wilkes and C. Ziter for assistance with field work, and to E. Lopez and M. Fahmy

for data entry. Salmonier Nature Park graciously supplied housing to RTM for the 2015 field

work.

References

Anderson MJ (2001) A new method for non-parametric multivariate analysis of variance.

Austral Ecol 26:32–46. doi: 10.1046/j.1442-9993.2001.01070.x

Bidussi M, Gauslaa Y. 2015. Relative growth rates and secondary compounds in epiphytic

lichens along canopy height gradients in forest gaps and meadows in inland British

Columbia. Botany 93: 123–131. doi: 10.1139/cjb-2014-0214

Bokhorst S, Asplund J, Kardol P, Wardle DA (2015) Lichen physiological traits and growth

forms affect communities of associated invertebrates. Ecology 96:2394–2407. doi:

285 10.1890/14-1030.1

286 Bowker MA, Maestre FT, Eldridge D, et al (2014) Biological soil crusts (biocrusts) as a model

287  system in community, landscape and ecosystem ecology. Biodivers Conserv 23:1619–1637.

288  doi: 10.1007/s10531-014-0658-x

289 Brodo IM, Sharnoff SD, Sharnoff S (2001) No Title. Lichens North Am

290 Busse A, Schoreisz JJ, Petermann JS (2019) Predators and priority effects suggested as potential

291  drivers of microfauna communities in a community transplantation experiment along an

292  elevational gradient. Oikos. doi: 10.1111/oik.05894

293 Culberson CF, Kristinsson H-D (1970) A standardized method for the identification of lichen

294  products. J Chromatogr A 46:85–93. doi: 10.1016/S0021-9673(00)83967-9

295 Drake JA (1993) The construction and assembly of an ecological landscape. J Anim Ecol 62:117–

296  130. doi: 10.2307/5487

297 Faegri K. 1980. Growth in an *Ochrolechia androgyna* thallus 1961-1979. Lichenologist 12: 248–

298  250. doi:10.1017/S0024282980000217

299 Gilbert F, Gonzalez A, Evans-Freke I (1998) Corridors maintain species richness in the

300  fragmented landscapes of a microecosystem. Proc R Soc B Biol Sci 265:577–582. doi:

301  10.1098/rspb.1998.0333

302 Gonzalez A, Chaneton EJ (2002) Heterotroph species extinction, abundance and biomass

303  dynamics in an experimentally fragmented microecosystem. J Anim Ecol 71:594–602. doi:

304  10.1046/j.1365-2656.2002.00625.x

305 Gonzalez A, Lawton JH, Gilbert FS, et al (1998) Metapopulation dynamics, abundance, and

306  distribution in a microecosystem. Science (80-) 281:2045–2047. doi:

307  10.1126/science.281.5385.2045

308 Grainger TN, Gilbert B (2016) Dispersal and diversity in experimental metacommunities:

309  linking theory and practice. Oikos 125:1213–1223. doi: 10.1111/oik.03018

310 Hättestrand C, Kleman J (1999) Ribbed moraine formation. Quat Sci Rev 18(1):43-61

Jenerette GD, Shen W (2012) Experimental landscape ecology. Landsc Ecol 27:1237–1248. doi:

10.1007/s10980-012-9797-1

Kohler RE (2002) Landscapes and labscapes: Exploring the lab-field border in Biology.

University of Chicago Press.

Kotliar NB, Wiens JA (1990) Multiple scales of patchiness and patch structure: a hierarchical

framework for the study of heterogeneity. Oikos 59:253–260. doi: 10.2307/3545542

Lau MK, Baiser B, Northrop A, et al (2018) Regime shifts and hysteresis in the pitcher-plant

microecosystem. Ecol Modell 382:1–8. doi: 10.1016/j.ecolmodel.2018.04.016

Logue JB, Mouquet N, Peter H, Hillebrand H (2011) Empirical approaches to metacommunities:

A review and comparison with theory. Trends Ecol Evol 26:482–491. doi:

10.1016/j.tree.2011.04.009

Mouquet N, Daufresne T, Gray SM, Miller TE (2008) Modelling the relationship between a

pitcher plant (Sarracenia purpurea) and its phytotelma community: Mutualism or

parasitism? Funct Ecol 22:728–737. doi: 10.1111/j.1365-2435.2008.01421.x

O'Neill R V, Johnson AR, King AW (1989) A hierarchical framework for the analysis of scale.

Landsc Ecol 3:193–205. doi: 10.1007/BF00131538

Oksanen J, Blanchet GF, Kindt R, et al No Title. Wagner, H (2016) Vegan Community Ecol

Packag R Packag version 23-4

Pereira CL, Araújo MB, Matias MG (2018) Interplay between productivity and regional species

pool determines community assembly in aquatic microcosms. Aquat Sci 80:. doi:

10.1007/s00027-018-0596-x

Phillips HC. 1969. Annual growth rates of three species of foliose lichens determined

photographically. Bull Torrey Bot Club. 96: 202–206. doi: 10.2307/2483333

Schiesari L, Sgambatti Monteiro A, Ilha P, et al (2018) The ecology of a system of natural

mesocosms: Rock pools in the Atlantic Forest. Freshw Biol 63:1077–1087. doi:

10.1111/fwb.13118

Schindler DW (1998) Replication versus realism: The need for ecosystem-scale experiments.

Ecosystems 1:323–334. doi: 10.1007/s100219900026

South G (1983) Biogeography and ecology of the Island of Newfoundland, 1st edn. Dr. W. Junk

Publishers, The Hague

Southwood TRE, Kennedy CEJ (1983) Trees as islands. Oikos 41:359–371. doi:

10.2307/3544094

Srivastava DS, Kolasa J, Bengtsson J, et al (2004) Are natural microcosms useful model systems

for ecology? Trends Ecol Evol 19:379–384. doi: 10.1016/j.tree.2004.04.010

Stewart RIA, Dossena M, Bohan DA, et al (2013) Mesocosm Experiments as a Tool for

Ecological Climate-Change Research. Adv. Ecol. Res. 48:71–181

Talaga S, Dézerald O, Carteron A, et al (2015) Tank bromeliads as natural microcosms: A

facultative association with ants influences the aquatic invertebrate community structure.

Comptes Rendus - Biol 338:696–700. doi: 10.1016/j.crvi.2015.05.006

Trekels H, Driesen M, Vanschoenwinkel B (2017) How do patch quality and spatial context

affect invertebrate communities in a natural moss microlandscape? Acta Oecologica

85:126–135. doi: 10.1016/j.actao.2017.10.003

Urban DL (2005) Modeling ecological processes across scales. Ecology 86:1996–2006. doi:

10.1890/04-0918

Urban DL, O'Neill R V, Shugart HH (1987) Landscape ecology. Bioscience 37:119–127

Wiens JA, Milne BT (1989) Scaling of "landscapes" in landscape ecology, or, landscape ecology

from a beetle's perspective. Landsc Ecol 3:87–96. doi: 10.1007/BF00131172

Wiersma YF, McMullin RT (2018) Is it common to be rare on the landscape? A test using a novel

model system. Landsc Ecol 33:. doi: 10.1007/s10980-017-0599-3

Wu J (2013) Landscape Ecology. In: Ecological systems. pp 179–200

Young AR, Miller JED, Villella J, et al (2018) Epiphyte type and sampling height impact

mesofauna communities in Douglas-fir trees. PeerJ 2018:. doi: 10.7717/peerj.5699

Zuk M, Travisano M (2018) Models on the runway: How do we make replicas of the world? Am

Nat 192:1–9. doi: 10.1086/697508

Box 1. Glossary of Terms

mesocosm: a contained system (e.g., aquarium) that is designed to mimic a natural, larger ecosystem for the purposes of experimental manipulation.

microcosm: a small, contained natural ecosystem, such as insects in a pitcher plant, or a tide pool community.

microecosystem: a small ecosystem that is constrained/bounded by specific structural or environmental factors (e.g., pH, substrate).

patch: the fundamental unit in a landscape; a patch is considered to be homogenous within, relative to the landscape and have its own intrinsic properties (e.g., size, shape).

landscape: an areas that is spatially heterogeneous in at least one factor of interest.

landscape mosaic: a landscape comprised of two or more patches (ecosystems) that interact to create a pattern at a larger extent.

hierarchy: a conceptual framework for natural systems; lower levels of organization (finer grain/smaller extent) are constrained by higher levels, yet at the same time, lower levels influence higher level patterns. For example, gaps are nested within forest stands; stands are nested within tracts and tracts within a larger ecosystem which is nested within a biome.

Figure Captions

Figure 1. A schematic of the hierarchical structure of the proposed model system for landscapes. At the smallest scale (panel a) the landscape is centimeters in extent and is comprised of patches of lichens on the bole for trees. Each tree can be considered a replicate microlandscape. Within a single forest stand (panel b), the landscape is metres in extent. Some stands are homogenous in terms of tree species, age/size, but across the stand there will be fine-grained heterogeneity in microtopography and microclimate. These stands are then patches in a kilometers-extent landscape (panel c); some stands of similar composition may be replicated across the landscape but interspersed with other patch types (e.g., non-forested bog, meadow, anthropogenic patches).

Figure 2. Location of sampling sites in the Avalon Forest Ecoregion on the island of Newfoundland, Canada. Inset map shows the location of the Avalon Forest Ecoregion (black polygon). Black stars on the main map are sampling locations for the 21 sites where we sampled both balsam fir and yellow birch. The location labelled "Halls Gullies" designates the stand within which 24 balsam fir were sampled.

Figure 3. A lichen ladder, which was used to sample lichen diversity on tree boles. Dimensions of each square are 10 x 10 cm; the ladder is 50 cm in length and was placed with the top rung at 1.6 m from the ground.

390 **Table 1**. **Summary of perMANOVA analysis for trees within the Avalon Forest Ecoregion**.

391 Date is from two sampling schema; one for 24 balsam fir in a single stand; and for 21 more

392 widely dispersed sites with one balsam fir and one yellow birch at each site. The analysis is for

393 consistency in lichen patterns on the tree boles for macro-lichens only along a 50 cm section of

394 the bole on the north- and south-facing sides.

Variable	Balsam fir in a single stand ($n = 24$)	Balsam fir dispersed ($n = 21$)	Yellow birch dispersed ($n = 21$)
Aspect controlling for tree	significant	significant	non-significant
Aspect controlling for location up tree	non-significant	non-significant	non-significant
Location up tree controlling for tree	significant	non-significant	non-significant
Location up tree controlling for aspect	non-significant	non-significant	non-significant

397 **Table 2. Summary of perMANOVA analysis for 21 widely dispersed sites across the Avalon**

398 **Forest Ecoregion.** Data are shown for balsam fir and yellow birch; there was one of each at each

399 site. The analysis is for consistency in lichen patterns along 50 cm of the tree boles along the

400 north and south-facing sides, for macro-lichens and mirco-lichens combined.

Variable	Balsam fir dispersed ($n = 21$)	Yellow birch dispersed ($n = 21$)
Aspect controlling for tree	non-significant	significant
Aspect controlling for location up tree	non-significant	significant
Location up tree controlling for tree	non-significant	non-significant
Location up tree controlling for aspect	non-significant	non-significant

A

⊢ cm ⊣

B

⊢ m ⊣

C

⊢ km ⊣

Trans-Canada Highway

Halls Gullies

Hwy 91

Manitoba River

Newfoundland

Sites
Water bodies

0   2.5   5   10
Kilometers