Attention updates the perceived position of moving objects

Ryohei Nakayama[1][2]*, Alex O. Holcombe[1]

[1] School of Psychology, The University of Sydney, NSW 2006, Australia

[2] Center for Information and Neural Networks (CiNet), National Institute of Information

and Communications Technology, 1-4 Yamadaoka, Suita City, Osaka, 565-0871, Japan

*Correspondence: ryouhei.nakayama@gmail.com

Abstract

The information used by conscious perception may be somewhat different from that

which drives certain actions. In support of this notion, recent studies reported that

although internal grating motion can accumulate over seconds into a large illusory

position shift, this position shift is not reflected in saccade targeting (action). Another

possibility however is that rather than saccades and other actions having privileged

access to the correct position, the attention shift thought to precede saccades resets the

accumulated position shift to zero. Here we found that the accumulation of illusory

position shift can be reset by transients near the moving object and also by the observer

pressing a button at the time the object is perceived to reach a particular position. This

creates a striking impression of the object jumping back to near its actual position. These

results suggest that both stimulus-driven attention and attention associated with control

of action may update the perceived position of moving objects and mediate the

previously reported dissociation between conscious perception and saccades.

1 Results

Visual motion signals can strongly affect the perception of position. For example, texture
motion within a stationary object shifts that object's position in the direction of the
texture motion [1,2]. Interestingly, if such an object moves by itself in a trajectory
orthogonal to the texture motion, the illusory position shift accumulates over seconds,
giving a sensation of trajectory oriented between the directions of texture and object
motions [3,4]. This is referred to as the curveball or double-drift illusion. Lisi & Cavanagh
[5] reported that when participants are asked to move their eyes (saccade) to a
double-drifting object, their eyes land in approximately the veridical position rather than
the position perceived. Lisi & Cavanagh concluded that the information used by
conscious perception is somewhat different from that which drives certain actions
(saccades). The present study instead suggests that the attention shifts thought to
precede saccades resets the accumulated position shift to zero, explaining why saccades
are made to the veridical object location, and highlighting a new role for attention.

We find that both luminance and isoluminant transients near the moving object ("flash"
and "color change" experiments respectively) and an observer's button press
subjectively timed to the object reaching a particular position
("button-press" experiment) can reset the accumulation of illusory position shift,
creating an impression of the object jumping back to its actual position. What do these
manipulations have in common? Arguably, a shift of attention toward a particular
location in the moving object's trajectory.

In each trial of our flash experiment, a pair of circular objects moved diagonally down on
a screen and horizontally outward from a fixation point in the left and right upper visual
fields respectively (Figure 1A) while internal grating patterns parallel to the object
trajectories were either stationary within the circular envelopes (single-drifting objects)
or drifted inward (double-drifting objects). During the single-drift trials and half of the
double-drift trials, white squares were flashed on both sides of each of the two moving
objects for one movie frame, which occurred at a random time between 0.7-1.1 sec after
stimulus motion onset. The moving stimuli disappeared 1.8 sec after onset. Participants
maintained fixation throughout (monitored by an eye tracker) and drew the object
trajectories they perceived with a computer mouse. All trials containing saccades in a
period from 0.7 to 1.5 sec were removed from subsequent analyses.

Figure 1B shows the trajectories reported for double-drifting objects (left column),

double-drifting objects accompanied by flashes (middle column), and single-drifting

objects accompanied by flashes (right column). Pink-shaded bands indicate intervals

wherein flashes were presented. It can be seen that the internal grating motion of the

majority of the perceived trajectories of the double-drifting objects biased the

trajectories perceived toward fixation, especially in the first half of the trajectory. The

appearance of the flashes (middle column) frequently results in a sharp trajectory

change, sometimes described by participants as a jump, toward the true position of the

object. On the other hand, single-drifting objects keep moving straight despite the flash

presentation. In summary, the luminance flash frequently resets the accumulation of

illusory position shift caused by double drifts.

The variance of the trajectories drawn for different trials and participants is significant.

To quantitatively and reproducibly assess the proportion of trials in which a jump was

indicated in the trajectory drawn, we classified them as "jump" and "no jump" trials with

a likelihood-ratio test [6]: a statistical comparison of models with versus without a

horizontal jump in a linear or curved course of stimulus trajectories. Specifically, a

quadratic curve ($x = ay^2 + by + c$; x and y denote horizontal and vertical positions; a, b,

and c are free parameters) both with and without two additional free parameters to

create a horizontal jump (corresponding to the timing and the size of the jump) was

fitted to each trajectory, by maximum likelihood assuming that the model's errors are

normally distributed.

$$Likelihood = \prod_{i=1}^{n} \exp\left(\frac{-(x_i - \mu_i)^2}{2\sigma^2}\right) \qquad (1)$$

Here, μ and σ denote a model prediction of horizontal position (x) and standard

deviation of horizontal positions, respectively. The fits of the curves were very good – on

average with the jump/no-jump model r^2 = 0.970/0.962 (SE = 0.014/0.019) for

double-drifting objects, 0.958/0.900 (SE = 0.012/0.030) for same objects with flashes,

and 0.992/0.986 (SE = 0.002/0.004) for single-drifting objects with flashes (also see

Figure S1 panels for the best-fit curves). The likelihood-ratio test was used to decide

whether the jump model fit significantly better than the no-jump model. This test used

the maximum likelihood for the jump and no-jump models and the following test statistic

(*T)*, which follows a Chi-squared distribution (degrees of freedom = 1; the timing

parameter operates fully depending on the presence/absence of the size parameter).

$$T = 2log\left(\frac{Likelihood\ with\ jump}{Likelihood\ without\ jump}\right) \qquad (2)$$

Perceived (sketched) trajectories were classified as "jump" paths if T values exceeded

3.84 (corresponding to $p<.05$), which are drawn in bold in Figure 1B. Figure 1C shows

the proportion of the "jump" paths in each condition. Jumps were much more common

for double-drifting objects with flashes compared to the other two conditions (one-way

ANOVA $F_{(2,8)}$ = 48.92, p < 0.0001; between double-drifting objects with and without

flashes $t_{(8)}$ = 8.85, p < 0.0001; between double- and single-drifting objects with flashes

$t_{(8)}$ = 8.25, p < 0.0001; between double-drifting objects without flashes and

single-drifting objects with flashes $t_{(8)}$ = 0.60, p = 0.56). For a potentially corroborating

analysis, instead of using the T value to classify the trials and comparing their proportion,

we compared T values between conditions directly. Circles in Figure 1C plot average T

values. Similar to the classification-based analysis, the T value is significantly larger for

double-drifting stimuli with flashes (one-way ANOVA $F_{(2,8)}$ = 16.79, p = 0.001; $t_{(8)}$ =

4.98, p = 0.001; $t_{(8)}$ = 5.05, p = 0.001; $t_{(8)}$ = 0.07, p = 0.95 in comparisons of the same

order as above). The perceived jumps occurred around the time of the flashes (relative

to the flashes' time, -0.05 ± 0.22 sec, 95% CI), based on the curve fits after scaling the

vertical distance of the sketched trajectory to that of the actual trajectory.

These results indicate that presentation of the luminance flashes reset the accumulated

illusory position shift, causing a jump. Why do the flashes cause a jump? It might

conceivably be mediated by the first-order motion energy of the flashes [7,8], the fact

that flashes can suppress certain percepts [9], or from the participant's attention being

grabbed by the sudden appearance of the flashes. If it does result from stimulation of

motion or luminance transient detectors, perceptual jumps may not occur without a

strong luminance transient. To investigate this, in a subsequent experiment the moving

objects' color was abruptly changed halfway through the trajectory after assessing

equiluminance points to avoid or minimize a luminance component (Figure 2A). Such

chromatic changes can attract attention [10]. As detailed below, this isoluminant

chromatic change also resulted in jumps, which suggests it is a shift of attention that

triggers the perceptual jump, not the luminance change.

Figure 1. Schematic stimulus display and results of the flash experiment. (A) Two circular objects

moved diagonally for 1.8 sec with/without orthogonal motion of internal grating patterns:

double-drifting (see Movie S1 for no flashes, Movie S2 for flashes) or single-drifting objects (see

Movie S3). Black arrows indicate the physical motion directions of envelope and grating for the

double-drifting objects. White squares were flashed for one screen frame between 0.7-1.1 sec

after the motion began, except there was no flash in half of the double-drift trials. Orange lines

(not shown during experiment) indicate possible perceived trajectories of double-drifting objects

with (solid) or without (dashed) flashes, respectively. (B) Object trajectories sketched by

participants. Rows correspond to participants and columns correspond to stimulus conditions.

Center of screen is (0,0) in X and Y coordinates. Display colors correspond to trial numbers and

bold curves denote paths classified as jumps by maximum likelihood. Black diagonal lines indicate

physical trajectories and pink-shaded bands indicate spatial locations of flash presentations. (C)

Bars show average proportion of perceived trajectories classified as jumps. Circles plot average

likelihood ratio or T (right-side ordinate). Error bars represent $\pm$ 1 SE across participants.

In this chromatic change experiment, reddish and greenish objects were presented on a

yellowish background adjusted to equiluminance for each participant (see Methods).

During the single-drift trials and half of the double-drift trials, at a random time 0.7 to

1.1 sec into the trial, the red and green colors were swapped. Otherwise, stimulus

display, drawing task, and participants were all same as the previous experiment.

Trajectory data were analyzed in the same way as the previous analyses. The jump/no

jump models again fit well, r^2 = 0.966/0.956 (SE = 0.021/0.023) for double-drifting

objects, 0.971/0.935 (SE = 0.009/0.021) for same objects with color changes, and

0.996/0.995 (SE = 0.001/0.001) for single-drifting objects with color changes (also see

Figure S2 panels for the best-fit curves).

The paths classified as jumps by the likelihood-ratio test are drawn in bold in Figure 2B.

Trajectory jumps are mainly observed for double-drifting objects accompanied by color

changes, except for Participant 5 who strangely shows more jumps for double-drifting

objects without color changes than with them. Note that because the three types of

stimulus trials are randomly interleaved, participants will have some expectation of a

color change, which might result in some anticipatory attention shifts and trigger jumps

in some non-color-change trials, if attention shifts cause jumps.

The proportion of trials in each condition that yielded jumps is shown by Figure 2C.

Jumps occurred more often for double-drifting objects with color changes than for the

other two conditions (one-way ANOVA $F_{(2,8)}$ = 7.15, p = 0.02; between double-drifting

objects with and without color changes $t_{(8)} = 2.65$, $p = 0.03$; between double- and single-drifting objects with color changes $t_{(8)} = 3.66$, $p = 0.006$; between double-drifting objects without color changes and single-drifting objects with color changes $t_{(8)} = 1.01$, $p = 0.34$). Average T values (circles) were larger for double-drifting objects with color changes than without (one-way ANOVA $F_{(2,8)} = 5.21$, $p = 0.04$; $t_{(8)} = 2.31$, $p = 0.0498$; $t_{(8)} = 3.11$, $p = 0.01$; $t_{(8)} = 0.80$, $p = 0.45$ in comparisons of the same order as above). The curve fits estimate that the jumps occurred, relative to the color changes, at 0.07 ± 0.33 sec (95% CI).

These results indicate that a sudden chromatic change in moving objects can reset the accumulation of illusory position shift caused by double drifts, despite the lack of luminance change, which reduces input to motion detectors and flash suppression processes.

The next experiment assessed whether it was truly attentional engagement with the objects, rather than something more specific to stimulus transients, that causes jumps. Participants pressed a button at the moment that moving objects arrived at a certain position. Stationary landmarks flanking that position were presented at the beginning of the trial but then removed prior to the presentation of moving objects, to eliminate the possibility of any sensory interaction. We expected that participants would particularly direct attention to the objects around the time that they should respond to them.

2　Figure 2. Schematic stimulus display and results of the color change experiment. (A) Reddish and

3　greenish objects, with mean luminance matched to the yellowish background, moved diagonally.

4　In the color change condition, they abruptly changed their color with each other. Other stimulus

5　configurations were identical to those in the flash experiment. Arrows with numbers (not shown in

the experiment) indicate the predicted perceived trajectories of double-drifting objects (see Movie

S4), double-drifting objects with color change (see Movie S5), and single-drifting objects with

color change (see Movie S6). (B) Object trajectories sketched by participants. Black diagonal lines

indicate physical trajectories and pink-shaded bands indicate region in which the color change

happened. (C) Bars plot average proportion of "jump" paths (left-side ordinate). Circles plot

average likelihood ratio or T (right-side ordinate). Error bars represent ± 1 SE across

participants.

At the start of each trial, white squares were presented for 1.6 sec on both sides of the

future object trajectories. After a 0.4-sec blank fixation period, the moving objects

appeared. Participants were instructed to press a button with the right index finger when

the moving objects passed between the former locations of the squares, while keeping

their eyes on the fixation point. Other aspects were the same as the first experiment.

Trajectory data were analyzed in the same way as the previous analyses, the fits for the

jump/no-jump model are r^2 = 0.980/0.955 (SE = 0.005/0.011) for double-drifting

objects, 0.997/0.996 (SE = 0.001/0.001) for single-drifting objects (also see Figure S3

panels for the best-fit curves).

The paths classified as jumps by the likelihood-ratio test are drawn in bold (Figure 3A).

All paths classified as jumps were in the double-drifting objects condition. Bars in Figure

3B show average rates of the "jump" paths. One-way ANOVA reveals that it is

significantly higher for double-drifting objects compared to single-drifting objects, $F_{(1,4)}$

= 25.02, p = 0.008. Circles in Figure 3B plot average T values. Similar to this

classification-based analysis, one-way ANOVA indicates the T is larger for double-drifting

objects, $F_{(1,4)}$ = 15.12, p = 0.02. The jumps occurred around the time of the button

press (-0.03 ± 0.26 sec, 95% CI, relative to when the participants pressed the button),

consistent with an attention shift around that time.

These results indicate that the button-press task frequently does reset the accumulation

of illusory position shift caused by double drifts. This also shows that stimulus transients

are not necessary for the occurrence of trajectory jumps. Putting together all the results,

we suggest that a shift of attention to (close to) the moving objects resets the

accumulation of illusory position shift caused by double drifts and causes a jump from

the accumulated illusory position to closer to the real position.

Figure 3. Results of the button-press experiment. (A) Object trajectories sketched by participants.

Black diagonal lines indicate physical trajectories and pink-shaded bands indicate spatial locations

of initial landmark presentations. (B) Bars plot average proportion of "jump" paths (left-side

ordinate). Circles plot average likelihood ratio or T (right-side ordinate). Error bars represent ± 1

SE across participants.

Discussion

We have found that an accumulated illusory position shift is often reset to near zero by

a luminance or chromatic transient in a moving object. This reset can also be triggered

by a task involving sensorimotor reactions to a particular position of a moving object.

These results are consistent with the proposition that stimulus-driven [11] and

top-down attention [12] can reset the perceived position of a moving object.

Lisi & Cavanagh (2015) found that saccades to a double-drifting object were unaffected by the illusion. They suggested that this constituted the first clear and large difference between perceived position and saccade targeting, providing critical evidence for the long-running debate over how perception and action differ in their use of sensory signals. However, the present study suggests that the reason saccades are made to the correct position can be explained by the associated allocation of attention [13] rather than a dissociation between perception and action. This would also explain similar jump percepts caused by a gaze shift to a double-drifting object as described in the first report of the curveball illusion [3].

The present study hints that rather than saccades and other actions having privileged access to the correct position, perceptual and oculomotor systems rely on the same position information [14], at least in the case of this illusion. Several previous studies have investigated stationary objects with internally-drifting texture, which elicits a small position shift and relatively little accumulation [1,2,15]. One study [16], while showing that the size of the shift varies with the stimulus used to compare position to, specifically found that it is *smaller* if compared to flashed lines than if judged by actions, specifically pointing and saccades. We suggest that this result may reflect the flashed lines resetting MIPS close to zero, while the actions are guided by some information after the attention shift, during which MIPS may accumulate [15].

While an eyetracker was used to exclude trials with saccades, smaller eye movements, termed micro-saccades, may be a concern as abrupt stimulus changes induce micro-saccades [17]. However, as can be seen in Movie S7, the curveball illusion still occurs when a fixation target in motion is pursued with the eyes. Smooth pursuit regularly contains small saccades to catch up to a moving target tracked but such small jitters in retinal image seem not to reset the accumulation of position shift.

The objects in the present experiments were likely attentionally tracked by the participants throughout most trials, since the participants' task was to report their entire trajectories. Diluting attentional tracking resources among four such objects [18,19] appears to have little effect on the curveball illusion [20]. But the transients that accompany sudden visual changes have long been suggested to engage attention to a greater extent than typically occurs with top-down attention (e.g., [21]), and may even engage qualitatively different attentional functions than does top-down attention

[22,23], such as resetting the phase of ongoing oscillations [24]. However, because we found similar effects here with the button-press task that involved no transients, we suspect that a greater intensity of top-down attention causes the reset.

Why would a greater intensity of attention reset the position perceived during the curveball illusion? Attention to an object can lead to activation of the primary visual cortex neurons stimulated by that object [25], and with early cortical cells' greater spatial precision, it is adaptive to use them for position decoding. Presumably greater attention to an object activates early retinotopic cells to a greater degree. Primary visual cortex representations are shifted in the direction of object motion [26], so use of these representations should not completely eliminate the curveball illusion. However, because the amount of shift in retinotopic areas is proportional to the size of the receptive fields [26], the use of early visual cortical areas should greatly reduce the size of the illusion relative to the use of later visual cortical areas. Thus an increase in attention can cause a perceptual jump of the object towards its actual location.

Methods

Participants

Four naive participants and one of the authors (RN) participated in the experiments. All participants had normal/corrected-to-normal vision. All experiments were conducted in accordance with the Declaration of Helsinki (2003) and were approved by the ethics committee of the University of Sydney. All participants provided written informed consent.

Apparatus

Images were displayed on a gamma-corrected 22-inch CRT monitor (Mitsubishi Diamond Pro 2070SB; 1280×1024 pixel) with a frame rate of 75 Hz. The CRT pixel resolution was 1.8 min/pixel at a viewing distance of 57 cm and mean luminance was 45.8 cd/m^2. CIE chromaticity was (0.61, 0.35) in R, (0.29, 0.62) in G, and (0.15, 0.07) in B channels, respectively. Eye movements of the right eye were monitored at a sampling rate of 1000 Hz (EyeLink 1000 version 4.56; SR Research Ltd, Mississauga, Ontario, Canada).

Stimuli

Visual stimuli were a pair of sine-wave grating patterns (0.8 cycle/deg) each within a

circular envelope (3.6 deg in diameter) such that grating edges would smoothly fade out

into the gray background according to a quarter-cycle cosine 0.3 deg gradient. A fixation

point (0.2 deg in diameter; 91.6 cd/m^2) was placed 3.0 deg below the center of screen.

The envelopes (objects) moved at 6.7 deg/sec diagonally (diagonally down and outward

at an angle of 30 degrees from straight down) in left and right upper visual fields

respectively. Their initial position was 15.5 deg away from the fixation. The orientation

of each grating was parallel to the object trajectory, its luminance contrast was 0.5, and

its initial spatial phase was randomized. The grating of the double-drifting objects drifted

inward at 2.2 deg/sec. For the single-drifting object, it was stationary relative to the

envelope.

<u>Procedure</u>

All experiments were conducted in a dark room. In each trial, participants maintained

gaze on the fixation point throughout the 1.8-sec presentation of the moving objects and

then, after the objects disappeared, traced the object trajectories by manually dragging

a computer mouse. Participants had the option to re-draw their response as many times

as they liked before confirming their response. Ten trials (except for Participant 3, who

participated in twenty trials) per stimulus condition were randomly interlaced in each

experiment.

For the flash experiment, white squares (0.7 x 0.7 deg) were presented for 13.3 msec

(one movie frame) on both sides of each of the two moving objects with horizontal

offsets of 5.4 deg from them, which was randomly timed between 0.7 and 1.1 sec after

the start of each trial.

For the color change experiment, instead of gray-scale grating patterns, reddish and

greenish grating patterns were used against a yellowish background. The B channel was

not used. The luminance contrast of the grating was 0.1 and the chromatic contrast

between the reddish and greenish patterns was maximized. One of the objects was

initially green, and the other initially red. The colors were initially randomly assigned to

the left and right objects and then swapped at a random time between 0.7 and 1.1 sec

after the start of each trial. Before the experiment started, the mean luminance of the

grating was subjectively matched to the background luminance by means of flicker

photometry [27] – participants minimized the subjective flicker by adjusting the green

intensity of the objects, positioned at the centers of the trajectories, alternating between the red and green colors at a rate of 8 Hz.

For the button response experiment, before the moving objects presentation, the same white squares as the flash experiment were presented for 1.6 sec then removed, followed by a blank fixation period of 0.4 sec. During the moving objects presentation, participants were told to press a button with the right-hand index finger when the objects would have passed between the squares if they had still been present. An occluder was positioned so that participants could not see their hand.

Analysis of eye movements and button presses

We removed trials that contained blinks or saccades (eye movements over 30 deg/sec or 8000 deg/sec^2) in a period from 0.7 to 1.5 sec after the start of each trial. On average, 81% (SD = 15%) of trials in the flash experiment, 87% (SD = 14%) of trials in the color change experiment, and 87% (SD = 17%) of trials in the button-press experiment were used in subsequent analyses. In the button-press experiment, a further 23% (SD = 19%) trials in which participants responded at a time that deviated from the physical alignment time by over 0.5 sec were removed.

Acknowledgments

This study was carried out when RN was a JSPS research fellow, a visiting researcher at the University of Sydney, and a visiting researcher at CiNet, NICT.

Author Contributions

RN and AOH designed the study. RN performed experiments and analyzed data. RN and AOH wrote the manuscript.

Declaration of Interests

The authors declare no competing interests.

References

1. Valois, R. De, and Valois, K. De (1991). Vernier acuity with stationary moving Gabors. Vision Res. *31*, 1619–1926.

2. Whitney, D. (2002). The influence of visual motion on perceived position. Trends Cogn. Sci. *6*, 211–216.

3. Shapiro, A., Lu, Z.L., Huang, C.B., Knight, E., and Ennis, R. (2010). Transitions between central and peripheral vision create spatial/temporal distortions: A hypothesis concerning the perceived break of the curveball. PLoS One *5*, e13296.

4. Tse, P.U., and Hsieh, P.J. (2006). The infinite regress illusion reveals faulty integration of local and global motion signals. Vision Res. *46*, 3881–3885.

5. Lisi, M., and Cavanagh, P. (2015). Dissociation between the perceptual and saccadic localization of moving objects. Curr. Biol. *25*, 2535–2540.

6. Neyman, J., and Pearson, E.S. (1933). On the Problem of the Most Efficient Tests of Statistical Hypotheses. Philos. Trans. R. Soc. A Math. Phys. Eng. Sci. *231*, 289–337.

7. Adelson, E.H., and Bergen, J.R. (1985). Spatiotemporal energy models for the perception of motion. J. Opt. Soc. Am. A. *2*, 284–299.

8. Watson, A.B. (1986). Temporal Sensitivity. In Handbook of Perception and Human Performance, Vol. 1: Sensory Processes and Perception, K. R. Boff, L. Kaufman, and J. P. Thomas, eds. (New York: Wiley), pp. 1–43.

9. Wolfe, J.M. (1984). Reversing ocular dominance and suppression in a single flash. Vision Res. *24*, 471–478.

10. Snowden, R.J. (2002). Visual attention to color: Parvocellular guidance of attentional resources? Psychol. Sci. *13*, 180–184.

11. Hollingworth, A., Simons, D.J., and Franconeri, S.L. (2010). New Objects Do Not Capture Attention without a Sensory Transient. Attention, Perception, Psychophys. *72*, 1298–1310.

12. Baluch, F., and Itti, L. (2011). Mechanisms of top-down attention. Trends Neurosci. *34*, 210–224.

13. Corbetta, M. (1998). Frontoparietal cortical networks for directing attention and the eye to visual locations: Identical, independent, or overlapping neural systems? Proc. Natl. Acad. Sci. U.S.A. *95*, 831–838.

14. Van Heusden, E., Rolfs, M., Cavanagh, P., and Hogendoorn, H. (2018). Motion extrapolation for eye movements predicts perceived motion-induced position shifts. J. Neurosci. *38*, 8243–8250.

15. Nishida, S., and Johnston, A. (1999). Influence of motion signals on the perceived position of spatial pattern. Nature *397*, 610–612.

16. Kerzel, D., and Gegenfurtner, K.R. (2005). Motion-induced illusory displacement reexamined: Differences between perception and action? Exp. Brain Res. *162*, 191–201.

17. Engbert, R. (2006). Microsaccades : a microcosm for research on oculomotor control, attention, and visual perception. Prog. Brain Res. *154*, 177–192.

18. Holcombe, A.O., Chen, W., and Howe, P.D.L. (2014). Object tracking : Absence of long-range spatial interference supports resource theories. J. Vis. *14*, 1–21.

19. Alvarez, G.A., and Franconeri, S.L. (2007). How many objects can you track?: Evidence for a resource-limited attentive tracking mechanism. J. Vis. *7*, 1–10.

20. Haladjian, H.H., Lisi, M., and Cavanagh, P. (2018). Motion and position shifts induced by the double-drift stimulus are unaffected by attentional load. Attention, Perception, Psychophys. *80*, 884–893.

21. Nakayama, K., and Mackeben, M. (1989). Sustained and transient components of focal visual attention. Vision Res. *29*, 1631–1647.

22. Nishida, S., and Johnston, A. (2002). Marker correspondence, not processing latency, determines temporal binding of visual attributes. Curr. Biol. *12*, 359–68.

23. Holcombe, A.O. (2009). Temporal binding favours the early phase of colour changes, but not of motion changes, yielding the colour–motion asynchrony illusion. Vis. cogn. *17*, 232–253.

24. Wood, D.K., Gu, C., Corneil, B.D., Gribble, P.L., and Goodale, M.A. (2015). Transient visual responses reset the phase of low-frequency oscillations in the skeletomotor periphery. Eur. J. Neurosci. *42*, 1919–1932.

1   25.   Roelfsema, P.R., Lamme, V.A.F., and Spekreijse, H. (1998). Object-based
2   attention in the primary visual cortex of the macaque monkey. Nature *395*, 376–
3   381.

4   26.   Harvey, B.M., and Dumoulin, S.O. (2016). NeuroImage Visual motion transforms
5   visual space representations similarly throughout the human visual hierarchy.
6   Neuroimage *127*, 173–185.

7   27.   Anstis, S.M., and Cavanagh, P. (1983). A Minimum Motion Technique for Judging
8   Equiluminance. Colour Vis. Psychophys. Physiol., 155–166.
9
10

Supporting Information

Figure S1. Best fits of object trajectories sketched by participants in the flash experiment. Rows

correspond to participants and columns correspond to stimulus conditions. Center of screen is

(0,0) in X and Y coordinates. Display colors correspond to trial numbers and bold curves denote

paths classified as jumps by maximum likelihood. Black diagonal lines indicate physical trajectories

and pink-shaded bands indicate spatial locations of flash presentations.

2 Figure S2. Best fits of object trajectories sketched by participants in the color change experiment.

3 Rows correspond to participants and columns correspond to stimulus conditions. Center of screen

4 is (0,0) in X and Y coordinates. Display colors correspond to trial numbers and bold curves denote

5 paths classified as jumps by maximum likelihood. Black diagonal lines indicate physical trajectories

6 and pink-shaded bands indicate region in which the color change happened.

Figure S3. Best fits of object trajectories sketched by participants in the button-press experiment.

Rows correspond to participants and columns correspond to stimulus conditions. Center of screen

is (0,0) in X and Y coordinates. Display colors correspond to trial numbers and bold curves denote

paths classified as jumps by maximum likelihood. Black diagonal lines indicate physical trajectories

and pink-shaded bands indicate spatial locations of initial landmark presentations.

Movie S1. Double-drifting objects. Texture motion accumulates over seconds into a large illusory position shift of objects, whose diagonal trajectories are perceived as straight down or slightly slanted in periphery (curveball illusion).

Movie S2. Double-drifting objects accompanied by flashes. Accumulated position shift is reset to near zero by luminance flashes, causing jump percepts in object trajectories.

Movie S3. Single-drifting objects accompanied by flashes. Diagonal trajectories are perceived mostly as they are regardless of luminance flashes (control condition).

Movie S4. Chromatic double-drifting objects. Texture motion accumulates over seconds into a large illusory position shift of objects, whose diagonal trajectories are perceived as straight down or somewhat slanted in periphery (curveball illusion).

Movie S5. Chromatic double-drifting objects accompanied by color changes. Accumulated position shift is reset to near zero by color changes (adjusted to equiluminance in the experiment), causing jump percepts in object trajectories.

Movie S6. Chromatic single-drifting objects accompanied by color changes. Diagonal trajectories are perceived mostly as they are regardless of color changes (control condition).

Movie S7. Double-drifting objects with a smooth-pursuing target. Curveball illusion occurs during smooth pursuing with the eyes a target in horizontal motion.