RapidAIM: A culture- and metaproteomics-based Rapid Assay of Individual Microbiome responses to drugs

Authors: Leyuan Li[1], Zhibin Ning[1], Xu Zhang[1], Janice Mayne[1], Kai Cheng[1], Alain Stintzi*[1], Daniel Figeys*[1,2,3]

Affiliations:

[1] Department of Biochemistry, Microbiology and Immunology, Ottawa Institute of Systems Biology, Faculty of Medicine, University of Ottawa, Ottawa, Canada.
[2] Department of Chemistry and Biomolecular Sciences, University of Ottawa, Ottawa, Canada.
[3] Canadian Institute for Advanced Research, Toronto, Canada.
*To whom correspondence should be addressed: Emails: dfigeys@uottawa.ca (DF), and astintzi@uottawa.ca (AS).

Abstract:

Background: Human-targeted drugs may exert off-target effects on the gut microbiota. However, our understanding of such effects is limited due to a lack of rapid and scalable assay to comprehensively assess microbiome responses to drugs. Drugs can drastically change the overall microbiome abundance, microbial composition and functions of a gut microbiome. Although we could comprehensively observe these microbiome responses using a series of tests, for the purpose of a drug screening, it is important to decrease the number of analytical tools used.

Results: Here, we developed an approach to screen compounds against individual microbiomes *in vitro* using metaproteomics adapted for both absolute bacterial abundances and functional profiling of the microbiome. Our approach was evaluated by testing 43 compounds (including four antibiotics) against five individual microbiomes. The method generated technically highly reproducible readouts, including changes of overall microbiome abundance, microbiome composition and functional pathways. Results show that besides the antibiotics, compounds berberine and ibuprofen inhibited the accumulation of biomass during *in vitro* growth of the microbiome. By comparing genus and species level-biomass contributions, selective antibacterial-like activities were found with 36 of the 39 non-antibiotic compounds. Seven of our compounds led to a global alteration of the metaproteome, with apparent compound-specific patterns of functional responses. The taxonomic distributions of responded proteins varied among drugs, i.e. different drugs affect functions of different members of the microbiome. We also showed that bacterial function can shift in response to drugs without a change in the abundance of the bacteria.

Conclusions: Current drug-microbiome interaction studies largely focus on relative microbiome composition and microbial drug metabolism. In contrast, our workflow enables multiple insights into microbiome absolute abundance and functional responses to drugs using metaproteomics as the one-stop screening tool. The workflow is robust, reproducible and quantitative, and is scalable for personalized high-throughput drug screening applications.

Keywords: Gut microbiome, drug response, *in vitro* culturing, metaproteomics, absolute abundance, functional profile

Background

Human-targeted drugs are primarily developed for their effects on the host, and little is known on their effects on the microbiome. Microbiome response to drugs could contribute to off-target drug effect [1]. In addition, the gut microbiome has been linked to gastroenterological, neurologic, respiratory, metabolic, hepatic, and cardiovascular diseases [2]. Therefore, targeting the microbiome could lead to novel therapies [3]. Although the effects of some drugs and compounds on the microbiome have been reported [4], many drug-microbiome interactions are unknown. This is due in part to the extremely high numbers of marketed drugs [5] and compounds in development [6] together with the lack of assays that can rapidly and comprehensively assess the effects of compounds on individual microbiomes.

Different *in vitro* approaches have been employed to study drug-microbiome interactions. One strategy involves long term stabilization of the microbiome, as shown in various intestinal microbiome simulators based on continuous flow [7-9]. This approach typically requires a long culture period to stabilize the microbiome (15-20 days), and notable shifts in taxonomic compositions compared with the inoculum have been shown [7, 10]. Moreover, the size and complexity of these culturing systems limit the number of individual microbiomes and drugs that can be examined [9], and thus may not be suitable for high-throughput drug screening purpose. Another strategy is to culture individual bacteria strains isolated from microbiomes. A recent study examined the effects of approved drugs on the biomass of forty individually-cultured bacterial strains in a high-throughput manner [11]. This approach highlighted the importance of biomass in identifying antibacterial-like effects. However, it did not take into account the complexity of a microbial community that could lead to different microbial responses.

Approaches such as optical density measurement [11], flow cytometry [12] and quantitative real-

time PCR [13] can be used to compare microbiome biomass. However, these approaches lack

insights into drug impact on microbial composition and functions, which are highly related to

healthy and disease states. Although we could comprehensively observe microbiome responses

by combining multiple tools, for the purpose of initial drug screening, it is important to minimize

the number of analytical tools used. There has been no report of an *in vitro* gut microbiome-

based drug screening approach that could assess both biomass responses and functional

alterations in one test.

The development of meta-omics approaches has allowed rapid and deep measurement of

microbiome compositions and functional activities. Genetic approaches such as 16S rDNA and

shotgun metagenomics have been regarded as the "gold standard" in microbiome analysis,

providing relative quantifications of microbiome membership composition and functional

capabilities [14, 15]. However, different microbial members can differ by several orders of

magnitude in biomass [16]. Moreover, there is little insight on which microbial traits actually

contribute to the functional activities of the microbiome, as functions predicted from 16S rDNA

or metagenomics analyses are not necessarily expressed. Studies have shown that gene copy

numbers are not representative of protein levels [17]. In addition, RNA expression have limited

correlation to the actual protein abundance [18]. In contrast, mass spectrometry (MS)-based

metaproteomics technology allows for deep insight into proteome-level information of the

microbiome [19, 20], providing quantified protein abundances that estimate the functional

activities of the microbiome. Proteins not only provide the biological activities to the

microbiome, but also build up a large amount of biomass in microbial cells. Hence, the

metaproteomic readouts can also be used to assess the microbiome biomass and analyze

community structure [21]. It has been validated that metaproteomics is a good estimator of

biomass contributions of microbiome members [22]. Despite its coverage could not compare to

that of the genomic sequencing-based technologies, metaproteomics could confidently quantify

proteins of the bacterial species that constitute>90% of the total biomass [23], making it

sufficient for a fast-pass drug screening application.

Here we report an approach named Rapid Assay of an Individual's Microbiome (RapidAIM)

facing gut microbiome-targeted drug screening, and evaluated the applicability of

metaproteomics for insights of microbiome responses to drugs. Briefly, in RapidAIM, individual

microbiomes are cultured in a previously optimized culture system for 24 hours, and the samples

are then analyzed using a metaproteomics-based analytical approach. A high-throughput equal-

volume based protein extraction and digestion workflow was applied to enable absolute biomass

assessment along with the functional profiling. To demonstrate the feasibility and performance

of the RapidAIM assay, we carried out a proof-of-concept study involving 43 compounds and

five individual gut microbiomes. Microbiome responses including changes in biomass, taxon-

specific biomass contributions, taxon-specific functional activities, and detailed responses of

interested enzymatic pathways can be obtained following the assay.

Results

Development and evaluation of RapidAIM

RapidAIM consists of an optimized microbiome culturing method, an equal-volume based

protein extraction and digestion workflow and a metaproteomic analysis pipeline (**Figure 1a**).

Briefly, fresh human stool samples are inoculated in 96-well deep-well plates and cultured with

drugs for 24 hours. We have previously optimized the culture model and validated that it

maintains the composition and taxon-specific functional activities of individual gut microbiomes

in 96-well plates [24]. After 24 hours, the cultured microbiomes are prepared for metaproteomic

analysis using a microplate-based metaproteomic sample processing workflow (**Supplementary**

Figure S1) adapted from our single-tube protocol [25]. The microplate-based workflow consists

of bacterial cell purification, cell lysis under ultra-sonication in 8M urea buffer, in-solution

tryptic digestion, and desalting. We validated each step of this workflow and found no significant

differences in identification efficiency between 96-well plate processing and single-tube

processing (**Supplementary Figure S1**). To compare total biomass, taxon-specific biomass and

pathway contributions between samples in a high-throughput assay format, we applied an equal

sample volume strategy to our recently developed metaproteomics techniques [20, 26, 27]. To

validate the absolute quantification of microbiome abundance by comparing total peptide

intensity, an equal volume of samples from a microbiome dilution series (simulating different

levels of drug effects) was taken for tryptic digestion and LC-MS/MS analysis. Summed peptide

intensity in each sample showed good linearity ($R^2 = 0.991$, **Figure 1b**) with a standard

colorimetric protein assay, showing that the total peptide intensity is a good indicator for

microbiome biomass levels. Since drugs could cause drastically change in microbiome

abundance, we then evaluated whether biomass differences between wells could cause bias in

identified functional and taxonomic compositions. We confirmed that the level of total biomass

didn't bias the composition of functional profiles (**Figure 1c**), protein groups (**Supplementary**

Figure S2a), and taxonomic abundances (**Supplementary Figure S2b**).

RapidAIM: Proof-of-concept study

We conducted a proof-of-concept (POC) study on the use of RapidAIM to characterize drug-

microbiome interactions. We selected 43 compounds that have been previously suggested to

impact, interact with, or be metabolized by the gut microbiome (**Supplementary Table S1**).

Thirty-seven of these compounds are FDA-approved drugs; four are antibiotics, and the others

include nonsteroidal anti-inflammatory drugs (NSAIDs), anti-diabetic drugs, aminosalicylate,

and statins, etc. Each compound, at a dose corresponding to the assumed maximal fecal

concentration of its daily dose, was added to five wells of 96-well plates containing 1 ml culture

medium in each well. The drug solvent, dimethyl sulfoxide (DMSO), was used as the negative

control. Then, each of the five wells for each compound was inoculated with a different fecal

microbiome from healthy human volunteers. Following 24 hours of culturing, the samples were

processed through the microplate-based workflow **(Figure S1)** and were subjected to a 90 min

gradient-based rapid LC-MS/MS analysis. Using our automated metaproteomic data analysis

software MetaLab [27], 101,995 peptide sequences corresponding to 24,631 protein groups were

quantified across all samples with a false discovery rate (FDR) threshold of 1% (**Figure 1e**). The

average MS/MS identification rate was 32.4 ±8.8% (mean ±SD); an average of 15,017 ±3,654

unique peptides and 6,684 ±998 protein groups were identified per sample. To provide a global

overview of the microbiome responses, a PCA was performed based on label-free quantification

(LFQ) intensities of protein groups **(Figure 1f)**. As expected, the samples clustered based on the

original microbiome source and not based on drug treatment. Within each individual microbiome

group, a number of drug-treated samples clustered closely to their control while several drug-

treated samples clearly separated from the non-treated control.

We next evaluated the robustness and reproducibility of the method by culturing one microbiome

with drugs in technical triplicates. Cultured triplicates yielded high Pearson's *r* for LFQ protein

group intensities **(Figure 1d)**. Hierarchical clustering based on Pearson's *r* of LFQ protein group

intensities between samples showed that with the exception of several compounds which

clustered closely with DMSO, cultured triplicates were clustered together **(Supplementary**

Figure S3a). Moreover, total biomass, functional enzymes, and species biomass contributions

were highly reproducible between triplicates as shown in **Supplementary Figure S3b-d**.

 Figure 1. Rapid Assay of Individual Microbiome (RapidAIM) workflow and performance. (**a**)

Experimental, analytical, and bioinformatics components of the RapidAIM workflow. Each individual's

gut microbiome samples are cultured with the test compounds in a 96-well deep-well plate at 37℃ in strict

anaerobic conditions for 24 hours followed by high-throughput sample preparation and rapid LC-MS/MS

analysis. Peptide and protein identification and quantification, taxonomic profiling, and functional

annotation were performed using the automated MetaLab software [27]. (**b**) A series of six dilutions

(dilution gradients: GRD1~6) of a same microbiome sample was tested in triplicate through the equal-

volume digestion and equal-volume MS loading protocol; the summed peptide intensity was compared to a

set of protein concentration standards provided with DC protein concentration assay and showed good

linearity (center points and error bars represent mean $\pm$ SD). (**c**) Stacked bars of clusters of orthologous

groups (COG) category levels across the six concentrations showing no bias at the functional

quantifications. (**d**) Analysis of three technical replicates in RapidAIM showing high Pearson's correlation.

(**e**) Numbers of MS/MS submitted, peptide sequence and protein group identifications in the POC dataset.

(**f**) PCA based on LFQ intensities of protein groups for all POC samples.

Effects of compounds on microbiome abundance and composition

We examined the effect of the 43 compounds on the overall abundance (biomass) of each

individual microbiome by comparing the total peptide intensity (**Figure 2a**). As expected, the

antibiotics greatly reduced total microbial biomass in most individual microbiomes (with one

exception of increased microbiome abundance in response to rifaximin, further examination is

shown in **Supplementary Figure S4**). Closely clustered with these antibiotics, berberine and

ibuprofen also inhibited the biomass of all individual microbiomes.

We next explored the effects of drugs on the microbiome composition based on bacterial

biomass contributions. To evaluate the overall shift of the microbiome, Bray-Curtis distance [28,

29] between drug-treated and DMSO control microbiome indicated that fructooligosaccharide

(FOS), rifaximin, berberine, diclofenac, ciprofloxacin, metronidazole and isoniazid significantly

shifted the microbiome (pairwise Wilcoxon test, FDR-adjusted $p < 0.05$; **Figure 2c**).

Our metaproteomic dataset allowed us to further examine the response of bacterial absolute

abundance by comparing summed peptide intensities of each taxon (**Figure 2b**). As expected,

the broad-spectrum antibiotics rifaximin, ciprofloxacin and metronidazole significantly inhibited

the absolute abundance of a high number of bacterial genera (Wilcoxon test, $p < 0.05$). Non-

antibiotic compounds, such as berberine, FOS, pravastatin, ibuprofen, diclofencac, flucycosine,

and indomethacin also showed significant decrease in the abundances of over ten genera. In

addition, selective antibacterial activities were found with 35 out of the other 39 compounds at

the genus level. Interestingly, it is clear in **Figure 2b** that while several genera were inhibited,

the absolute intensity of *Bifidobacterium* was significantly increased by fructose-

oligosaccharides (FOS). Compared to the absolute abundance, the relative abundance provided a

different insight into microbiome composition changes (**Supplementary Figure S5).** For

example, as opposed to the finding that ciprofloxacin and metronidazole inhibited the biomass of

most genera (Figure 2b), they significantly increased the relative abundance of genera

Bifidobacterium, *Ruminococcus*, *Butyrivibrio*, *Paenibacillus*, etc. Several genera including

Bifidobacterium, *Collinsella*, *Fusobacterium*, *Butyrivibrio* and *Leuconostoc* were significantly

increased in their relative abundances by FOS. Interestingly, members of the Actinobacteria

phyla, including *Eggerthella*, *Gordonibacter*, *Slackia*, and *Adlercreutzia* were more susceptible

to drugs compared to most other genera. Moreover, at the species level, we found that 36 of the

43 compounds significantly affected the biomass of at least one bacterial species (one-sided

Wilcoxon rank sum test, FDR-adjusted $p < 0.05$; **Supplementary Table S2**). To this end,

RapidAIM allowed for the assessment of changes in both absolute and relative abundances of

microbes in response to the compounds.

Figure 2. Response of microbiome abundance and composition to compounds. (**a**) Biomass responses of individuals' microbiomes to compounds relative to DMSO control. Ratio of peptide intensity between compound and DMSO control samples was calculated for each individual microbiome. (**b**) Log₂ fold-change of absolute abundance at the genus level in response to each drug compared with the DMSO control. Genera that existed in ≥80% of the volunteers are shown. Star (*) indicate significantly changed bacterial abundance by Wilcoxon test, $p < 0.05$. (**c**) Bray-Curtis distance of genus-level composition between drug-treated microbiomes and the corresponding DMSO control samples. Heatmap colors are generated with average of log2-fold changes among the five individual samples. Statistical significance was

calculated by pairwise Wilcoxon test (FDR-adjusted $p < 0.05$). Box spans interquartile range (25th to 75th

percentile) and line within box denotes median. For full compound names, see abbreviation list in

Supplementary Table S1.

Gut microbiome functions in response to compounds

The Bray-Curtis distance of protein group profiles showed that all the four antibiotics, as well as

FOS, berberine and diclofenac significantly altered the microbiome functions (**Figure 3a**). These

functional alterations likely stemmed from changes in taxonomic composition as revealed by the

genus-level Bray-Curtis distance analysis (**Figure 2c**). We next analyzed the protein group

intensities by partial least square discriminant analysis (PLS-DA) to determine whether

metaproteomic profiles could be used to discriminate between the DMSO-control and each of the

drug-treated microbiomes. In agreement with the Bray-Curtis analysis results, PLS-DA

interpretation identified drug-specific metaproteomic patterns associated with seven of our

compounds, including the four antibiotics, FOS, berberine and diclofenac (**Supplementary**

Figure S6). Hence, hereafter we named these seven compounds as class I compounds, whereas

others were named class II compounds. To gain a better understanding of the global effects of

class I compounds on the gut microbiome, we applied an unsupervised non-linear dimensionality

reduction algorithm, t-distributed stochastic neighbor embedding [30], to visualize this subgroup

of metaproteomic data based on protein group abundances (**Figure 3b**). Class I compounds led

to a global alteration of the metaproteome, with apparent compound-specific patterns.

We next examined the drug impacts on the abundance of functional proteins according to

clusters of orthologous groups (COG) of proteins. We identified 535 COGs significantly

decreased by at least one drug treatment; 15 of these COGs were decreased by ≥ 10 compounds

(**Supplementary Figure S7**). Diclofenac and FOS were the only two compounds that

significantly increased COGs (55 and 81 COGs, respectively). Enrichment analysis based on

these significantly altered COGs shows that COG categories found to be enriched were

responsive to 13 of our compounds (**Figure 3c**), six of those were class I compounds.

Interestingly, the non-antibiotic NSAID diclofenac increased the abundance of several COG

categories (**Figure 3c**). By mapping these significantly increased proteins from these COG

categories against the string database, we found that these altered proteins are functionally

interconnected (**Supplementary Figure S8**). Interestingly, one of the proteins that were highly

connected in the string network, COG0176 – transaldolase (1.76 $\pm$ 0.21 fold-change), is involved

in the biosynthesis of ansamycins, bacterial secondary metabolites that have antibiotic activities

[31].

Figure 3. Effect of compounds on metaproteomic profiles of the microbiome. (**a**) Bray-Curtis

distance of protein groups between drug-treated microbiomes and the corresponding DMSO

control samples. Statistical significance was calculated by pairwise Wilcoxon test (FDR-adjusted

$p < 0.05$). (**b**) Unsupervised dimensionality reduction analysis suggesting three different classes of

compound effects. (**c**) Enrichment analysis of all significantly different COGs in the POC dataset.

Significantly altered COGs with a p-value cutoff of 0.05 are shown.

Taxon-specific functional responses to class I compounds

We next performed a taxonomic analysis of the functional responses to diclofenac, FOS,

ciprofloxacin, and berberine, which represent four different types of compounds (NSAID,

oligosaccharide, antibiotics, anti-diabetes) in the class I. Protein groups with VIP scores >1

(thereafter defined as differential proteins) were extracted from each model, and were annotated

with their taxonomic and COG information. The taxonomic distributions of the differential

proteins varied among drugs (**Figure 4a)**. Moreover, mapping of the differential proteins to

phyla-specific pathways revealed phyla-specific responses, as shown for berberine in

Supplementary Figure S9. In agreement with **Figure 4a,** a higher proportion of down-regulated

than up-regulated pathways were identified in Firmicutes and Actinobacteria, while the opposite

pattern was observed in Bacteroidetes, Proteobacteria and Verrucomicrobia. In some cases, the

phylum-specific responses included up-regulation and down-regulation of different proteins

within the same pathway (black lines, **Supplementary Figure S9**). For example, we observed

this pattern in fatty acid, carbohydrate, and nucleotide metabolism pathways in Firmicutes.

Genus-level analysis revealed genus-specific responses to berberine (**Figure 4b**). In most genera,

the genus-specific responses correlated with the overall abundance of the corresponding genus

(**Figure 4b**, right panel). Nevertheless, some genera showed functional shifts in response to

berberine without changes in overall abundance. For example, *Bifidobacteria*, *Roseburia*,

Eubacterium, *Clostridium*, *Ruminococcus*, *Blautia*, and *Subdoligranulum* exhibited down-

regulation of proteins in various COG categories but no changes in biomass were observed.

Figure 4. Global functional effects of berberine, ciprofloxacin, FOS and diclofenac (**a**)
Taxon-function distribution of protein groups responding to berberine, FOS, ciprofloxacin and
diclofenac. Responding protein groups were selected by PLS-DA based on ComBat-corrected
data. The semicircle diameter represents the number of PLS-DA VIP>1 protein groups
corresponding to each phyla-COG category pair. (**b**) Genus-level shifts in functional activities in
response to berberine and the alterations in biomass of the corresponding genera. Functional shifts
(differential protein groups) were identified by PLS-DA. For each genus, the percentages of the

total numbers of up- and down-regulated protein groups corresponding to each COG category are

shown. Shifts in bacterial biomass in the five microbiomes are shown in box plots with the boxes

spanning interquartile range (25th to 75th percentile), and the vertical lines denoting the median

for each genus.

Enzymatic pathways in response to class I compounds

Next we examined the ability of RapidAIM in observing detailed enzymatic pathways of interest.

As examples, we show the effects of FOS and ciprofloxacin at the enzymatic pathway level.

Protein groups to were annotated to KEGG (Kyoto Encyclopedia of Genes and Genomes)

enzymes and were mapped against the KEGG pathway database. **Figure 5a** shows that FOS

increased enzymes responsible for fructan and sucrose uptake, as well as enzymes for conversion

of D-fructose into D-fructose-1-phosphate, D-mannose-6-phosephate and β-D-fructose-6-

phosphate. FOS also affected enzymes involved in the interconversion between glutamine,

glutamate and GABA (molecules involved in gut-brain communication). In addition, enzymes

involved in sulphide accumulation were affected, including decrease of dissimilatory sulfite

reductase (EC 1.8.99.5) and increase of cysteine synthase (EC 2.5.1.47).

Ciprofloxacin significantly altered the levels of enzymes involved in glycolysis/glycogenesis and

pentose phosphate pathways **(Figure 5b)**. The majority of enzymes involved in glycolysis were

significantly increased by ciprofloxacin. Ciprofloxacin down-regulated enzymes (ECs

1.1.1.49/1.1.1.363, 3.1.1.31, 1.1.1.44/1.1.1.343) involved in synthesis of ribulose-5-phosphate,

which can be isomerized to ribose 5-phosphate for nucleotide biosynthesis [32]. Moreover, the

levels of antioxidant enzymes superoxide dismutase (SOD) and catalase (CAT) were increased,

suggesting that ciprofloxacin induces oxidative stress in gut bacteria.

311
Figure 5. **Response of enzymatic pathways to drug treatment.** (a) Effect of FOS treatment on

enzymes involved in fructose and mannose metabolism, GABA production and sulfide

metabolism pathways. (b) effect of ciprofloxacin treatment on enzymes involved in the

glycolysis/gluconeogenesis and pentose phosphate pathway. The five blocks of each enzyme

represent the five individual microbiomes. Colors in the blocks represent differences between

normalized KEGG enzyme intensities with drug treatment versus DMSO (log2-transformation of

the original intensity followed by a quotient normalization (x/mean)).

Gut microbiome functions altered by class II compounds

Class II compounds, in contrast to class I compounds, did not cause a global shift in the five

individual microbiomes (an example is given by indomethacin, **Figure 6a**). However, **Figure 6a**

as well as the Bray-Curtis analyses (**Figure 2c** and **3a**) suggest that there were could be

individual variabilities in the extent of drug response. We show that if analyzed on an individual

sample basis, significant individualized functional effects could be revealed (**Figure 6b and c**),

suggesting high sensitivity of the RapidAIM assay in its application to personalized drug

screenings. For example, we identified 303 significantly altered protein groups in cultured

replicates of a single indomethacin-treated microbiome (V1). Taxon-function coupled

enrichment analysis showed that down-regulated functions were highly enriched in the genus

Bacteroides, while up-regulated functions were mostly enriched in the order Enterobacterales

(**Figure 6d and e**). The up-regulated functions of Enterobacterales included COG0459

chaperonin GroEL (HSP60 family) and COG0234 co-chaperonin GroES (HSP10) (**Figure 6e**).

Figure 6. Individual functional responses to indomethacin. (**a**) When visualizing several individual microbiome responses with PCA (based on LFQ intensities of protein groups), inter-individual variability can be greater than drug-induced functional shifts. (**b**) PCA clearly differentiated the response of microbiome V1 treated in triplicates using RapidAIM. (**c**) 303 significant protein group responders were found by *t*-test (FDR-adjusted *p*<0.05). (**d**) Taxon enrichment analysis based on the differential protein groups, (*p*-adjusted=0.05). (**e**) Taxon-function coupled enrichment analysis of up-regulated protein groups.

Discussion

In the present study, we developed an approach named Rapid Assay of Individual Microbiome (RapidAIM) to evaluate the effects of xenobiotics on individual microbiomes. The range of xenobiotics that reach the intestine and may interact with the gut microbiome is massive and expanding. These xenobiotics include antibiotics and other pharmaceuticals, phytochemicals, polysaccharides, food additives and many other compounds. With the exception of antibiotics,

we remain surprisingly ignorant on the extent to which these compounds affect the functions of

the gut microbiome. This understanding was limited by lacking an efficient and scalable

approach that could maximally obtain insights into microbiome responses while minimizing the

number of analytical tools being used.

Here we describe an approach which enables the exploration of drug-microbiome interactions

using an optimized *in vitro* culturing model and a metaproteomic approach. We have achieved

the maintenance of the representativeness of the initial individual microbiome [24]. Besides, for

an *in vitro* culturing simulating the *in vivo* microbiome, it is important to note that the population

of gut bacteria in the human body is highly dynamic. It has been estimated that there are

~$0.9 \cdot 10^{11}$ bacteria/g wet stool and a total of ~$3.8 \cdot 10^{13}$ bacteria in the colon. Approximately 200 g

wet daily stool would be excreted [33], leading to a dramatic decrease of the bacterial number in

the gut; on the other hand, new bacterial biomass starts growing on nutrients passing through the

gut. Current technologies examining the effect of xenobiotic stimulation are usually based on

microbiome stabilized after over two weeks of culturing. However, at the stable phase of

microbiome growth, the ecosystem has reached its carrying capacity (stable population size),

limiting possible observations such as drug effect on the biomass. In our studies, we have

previously validated that our composition of gut microbiome is well-maintained along the

growth curve [24], so we were able to observe drug responses of growing gut microbiomes by

adding the compounds at the initial inoculation stage. Subsequently, combined with our

quantitative metaproteomics approach based on equal-sample volume digestion, we were able to

observe the drug responses of overall microbiome abundance and taxon-specific biomass

contributions.

Here we address the importance of absolute quantification of microbiome biomass contributions.

Knowing the change of total microbiome biomass would be helpful to assess the antibiotic-like

activity of a compound. As our results clearly showed that the tested antibiotics inhibited the

accumulation of microbiome biomass, we found that non-antibiotic compounds ibuprofen and

berberine also showed inhibitory effects. Ibuprofen has been frequently used as a safe

medication. A study based on relative abundance discussed that ibuprofen had less aggressive

effects on the gut microbiome compared to some other NSAIDs [34]. However, in our study,

ibuprofen significantly inhibited the overall microbiome biomass through suppressing common

gut commensals such as *Bacteroides*, *Clostridium*, *Dorea*, *Eggerthella*, *Akkermansia*, etc. In

terms of berberine, previous studies suggested that berberine has positive effect on beneficial gut
microbes, e.g. selectively enriched a few putative short-chain fatty acid producing bacteria [35],
and increased the relative abundance of *Akkermansia* spp [36]. However, under the background
of an overall inhibition, an enrichment of a taxon (increase in relative abundance) does not
necessarily relate to its outgrowth. As another example, our result showed that ciprofloxacin,
metronidazole and FOS significantly enriched *Bifidobacterium* (**Supplementary Figure S5**).
The fact that *Bifidobacterium* has a certain resistance to ciprofloxacin and metronidazole [37]
could contribute to its higher adaptability over other genera. However, we didn't find evidence of
increase in its absolute abundance in presence of ciprofloxacin and metronidazole with
RapidAIM. On the contrary, FOS, which could be utilized as the carbon source of
Bifidobacterium [38, 39], significant increased both absolute and relative abundances of this
genus in our study.

We showed that the RapidAIM assay yielded insights into functional responses at multiple
levels. Using PLS-DA, we found that berberine, FOS, metronidazole, isoniazid, ciprofloxacin,
diclofenac, and rifaximin consistently shifted the metaproteome of the individual gut
microbiomes. By annotating the altered proteins at taxonomy, function and pathway levels, we
revealed the actions of the different drugs on the microbiome. For example, FOS treatment
elevated enzymes involved in fructan and sucrose uptake, as well as enzymes involved in the
interconversion among glutamine, glutamate and GABA, which are associated with microbiome
communication via the gut-brain axis [40]. In agreement, a study has shown that FOS
administration increased GABA receptor genes in mice, and further exhibited both
antidepressant and anxiolytic effects [41]. FOS also decreased proteins involved in sulfide
generation, suggesting decreased sulfide accumulation in the microbiome. This observation is in
agreement with *in vivo* studies showing that FOS treatment decreased the concentration of fecal
H_2S [42-44]. Ciprofloxacin treatment increased enzymes SOD and CAT, which was in
agreement with several reports indicating that ciprofloxacin triggers oxidative stress in several
bacteria [45-47]. With berberine treatment, we showed that taxon-specific functional shifts can
occur either with or without a change in the taxon's biomass. This observation highlights the
strength of our workflow which enables quantitative metaproteomic profiling of the microbiome.
Indeed, current classical sequencing-based approaches (16S rDNA or metagenomics
sequencing), which generate relative abundances, would not detect these types of changes.

Finally, we showed that although a compound may not show global impacts across the five
tested microbiomes, it could result in significant alterations on a single microbiome basis. The
example given by Indomethacin showed that the order Enterobacterales were enriched with
increased chaperonin GroEL (HSP60 family) and co-chaperonin GroES (HSP10) (**Figure 6e**),
which have been implicated in infection and diseases pathology [48].

Our workflow still exhibits certain limitations. In particular, MS analysis is a time-consuming
process. To this end, a fast-pass screening process such as tandem mass tags (TMT)[49, 50]
could be used to multiplex multiple microbiome samples in one MS analysis. Furthermore, our
workflow only measures the direct effects of compounds on the microbiome. In its current
implementation, it does not take into account the host effect on the microbiome and/or the effects
of drug metabolites produced by the host. Future efforts could be aimed at incorporating co-
culture of host cells/tissue and gut bacteria [51-53] into a high-throughput drug screening process
for achieving more comprehensive insights on host-drug-microbiome interaction.
Metaproteomics is a tool that is orthogonal to other omics technologies [17], hence for the need
of deeper investigations, RapidAIM could also be coupled with techniques such as
metagenomics or metabolomics for a multiple dimension view of the microbiome interaction
with drugs.

Conclusion

To date, the field of drug-microbiome interactions largely focuses on relative microbiome
composition and microbial drug metabolism, with a limited understanding of the effects of
pharmaceuticals on the absolute abundance and the function of the gut microbiome. A better
understanding of these interactions is essential given that the drug effects on the microbiome
biomass and functions may have important health consequences. Our workflow enabled the
insights into both absolute abundances and functional responses of the gut microbiome to drugs
using metaproteomics as the single analytical tool. We have shown that our workflow is robust,
reproducible and quantitative, and is easily adaptable for high-throughput drug screening
applications.

Methods

Stool sample preparation

The Research Ethics Board protocol (# 20160585-01H) for stool sample collection was approved by the Ottawa Health Science Network Research Ethics Board at the Ottawa Hospital. Stool samples were obtained from 5 healthy volunteers (age range 27 - 36 years; 3 males and 2 females). Exclusion criteria were: IBS, IBD, or diabetes diagnosis; antibiotic use or gastroenteritis episode in the last 3 months; use of pro-/pre-biotic, laxative, or anti-diarrheal drugs in the last month; or pregnancy. All volunteers were provided with a stool collection kit, which included a 50 ml Falcon tube containing 15 ml of sterile phosphate-buffered saline (PBS) pre-reduced with 0.1% (w/v) L-cysteine hydrochloride, a 2.5 ml sterile sampling spoon (Bel-Art, United States), plastic wrap, gloves and disposal bags. Briefly, each volunteer placed the plastic wrap over a toilet to prevent the stool from contacting water, collected ~3 g of stool with the sampling spoon, and dropped the spoon into the prepared 50 ml tube. The sample was immediately weighed by a researcher and transferred into an anaerobic workstation (5% H_2, 5% CO_2, and 90% N_2 at 37 ℃), where the tube was uncapped to remove O_2 before homogenization with a vortex mixer. Then the homogenate was filtered using sterile gauzes to remove large particles and obtain the microbiome inoculum.

Culturing of microbiomes and drug treatments

Each microbiome inoculum was immediately inoculated at a concentration of 2% (w/v) into a 96-well deep well plate containing 1 ml culture medium and a compound dissolved in 5 μl DMSO (or 5 μl DMSO as the control) in each well. The culture medium contained 2.0 g L^{-1} peptone water, 2.0 g L^{-1} yeast extract, 0.5 g L^{-1} L-cysteine hydrochloride, 2 mL L^{-1} Tween 80, 5 mg L^{-1} hemin, 10 μL L^{-1} vitamin K1, 1.0 g L^{-1} NaCl, 0.4 g L^{-1} K_2HPO_4, 0.4 g L^{-1} KH_2PO_4, 0.1 g L^{-1} $MgSO_4 \cdot 7H_2O$, 0.1 g L^{-1} $CaCl_2 \cdot 2H_2O$, 4.0 g L^{-1} $NaHCO_3$, 4.0 g L^{-1} porcine gastric mucin (cat# M1778, Sigma-Aldrich), 0.25 g L^{-1} sodium cholate and 0.25 g L^{-1} sodium chenodeoxycholate. The culture medium was sterile and had been pre-reduced overnight in an anaerobic workstation. Concentration of each compound was determined based on the assumption that maximal oral dosage of the drug distributed in 200 g average weight of the colon contents. However, several compounds (i.e. cimetidine, ciprofloxacin, flucytosine, mesalamine, metformin, metronidazole, naproxen-sodium, paracetamol, rifaximin, sodium butyrate, and sulfalazine) exceeded solubility in the given volume of DMSO (5 μl). After confirming that these compounds still showed effect

after a $10\times$ dilution (as can be seen from hierarchical clustering in **Supplementary Figure S3**),

the concentrations corresponding to the 1/10 highest oral dosages were used for these compounds.

Detailed catalogue number and concentration of each compound is listed in **Supplementary Table**

S1. After inoculation, the 96-well deep well plate was covered with a sterile silicone gel mat with

a vent hole for each well made by a sterile syringe needle. Then, the plate was shaken at 500 rpm

with a digital shaker (MS3, IKA, Germany) at 37 ℃ for 24 hours in the anaerobic chamber.

Metaproteomic sample processing and LC-MS/MS analysis

The sample processing was based on a previously reported metaproteomic sample processing

workflow[25], we adapted it for microplates (**Supplementary Figure S1**). Briefly, after

culturing for 24 hours, each 96-well plate was transferred out of the anaerobic station and was

immediately centrifuged at 300 g for 5 min to remove debris. The supernatants were transferred

into new 96-well deep well plates for another two rounds of debris removal at 300 g. The

supernatants were then transferred to a new plate and centrifuged at 2,272 g for 1 hour to pellet

the microbiome. The supernatant was removed and the pelleted bacterial cells were washed three

times with cold PBS in the same 96-deep well plate, pelleting the cells after each wash by a

2,272 g centrifugation for 1 hour. The 96-well plate containing harvested bacterial cells was then

stored overnight at -80°C before bacterial cell lysis and protein extraction. The lysis buffer was

freshly prepared, containing 8 M urea in 100 mM Tris-HCl buffer (pH = 8.0), plus Roche

PhosSTOP™ and Roche cOmplete™ Mini tablets. Microbial cell pellets were then re-suspended

in 150 μl lysis buffer and lysed on ice using a sonicator (Q125 Qsonica, USA) with an 8-tip-horn

probe. 100% amplitude was used (i.e. 15.6 Watts per sample), and four cycles of 30 s

ultrasonication and 30 s cooling down were performed. Protein concentrations of the DMSO

control samples were measured in triplicate using a detergent compatible (DC) assay (Bio-Rad,

USA). Then, a volume equivalent to the average volume of 50 μg of protein in the DMSO

control samples was acquired from each sample and placed into a new 96-deep well plate. The

samples were reduced and alkylated with 10 mM dithiothreitol (DTT) and 20 mM iodoacetamide

(IAA), followed by a $10\times$ dilution using 100 mM Tris-HCl (pH = 8.0) and tryptic digestion

under 37 ℃ for 18 hours using 1 μg of trypsin per well (Worthington Biochemical Corp.,

Lakewood, NJ). Digested peptides were desalted using a panel of lab-made 96-channel filter tips

generated by inserting 96 20 μl filter tips into a 96-well cover mat and stacking each filter tip

with 5 mg of 10-μm C18 column beads. After being washed twice with 0.1% formic acid (v/v),

tryptic peptides were then eluted with 80% acetonitrile (v/v)/0.1% formic acid (v/v).

After freeze-drying, each sample was re-dissolved in 100 μl 0.1% formic acid (v/v), and 2 μl of

the solution (corresponding to 1 μg of proteins in the DMSO control) was loaded for LC-MS/MS

analysis in a randomized order. An Agilent 1100 Capillary LC system (Agilent Technologies,

San Jose, CA) and a Q Exactive mass spectrometer (ThermoFisher Scientific Inc.) were used.

Peptides were separated with a tip column (75 μm inner diameter $\times$ 50 cm) packed with reverse

phase beads (1.9 μm/120 Å ReproSil-Pur C18 resin, Dr. Maisch GmbH, Ammerbuch, Germany)

following a 90 min gradient from 5 to 30% (v/v) acetonitrile at a 200 nL/min flow rate. 0.1%

(v/v) formic acid in water was used as solvent A, and 0.1% FA in 80% acetonitrile was used as

solvent B. The MS scan was performed from 300 to 1800 m/z, followed by data-dependent

MS/MS scan of the 12 most intense ions, a dynamic exclusion repeat count of two, and repeat

exclusion duration of 30 s.

Assessment of the equal-volume strategy

Six dilutions of a single microbiome sample were prepared in triplicate wells and an equal

volume was taken from each sample for tryptic digestion and LC-MS/MS analysis.

Metaproteomic sample processing and analysis followed the same procedures stated above, and

total peptide intensity was calculated. A DC protein concentration assay was also performed with

each sample. Linearity between total protein concentration and total peptide intensity quantified

by LC-MS/MS was then compared.

Metaproteomics data analysis

1) Metaproteomic database search:

Protein/peptide identification and quantification, taxonomic assignment and functional

annotations were done using the MetaLab software (version 1.1.0)[27]. MetaLab is a software

that automates an iterative database search strategy, i.e. MetaPro-IQ [26]. The search was based

on a human gut microbial gene catalog containing 9,878,647 sequences from

http://meta.genomics.cn/. A spectral clustering strategy were used for database construction from

all raw files, then the peptide and protein lists were generated by applying strict filtering based

on a FDR of 0.01, and quantitative information of proteins were obtained with the maxLFQ

algorithm on MaxQuant (version 1.5.3.30). Carbamidomethyl (C) was set as a fixed modification

and oxidation (M) and N-terminal acetylation (Protein N-term) were set as variable

modifications. Instrument resolution was set as "High-High".

2）Microbiome biomass analyses:

Total microbiome biomass was estimated for each sample by summing peptide intensities.

Taxonomic identification was achieved by assigning peptide sequences with lineage of lowest

common ancestor (LCA). The "peptide to taxonomy" database (pep2tax database) was selected

for mapping identified peptides to the taxonomic lineages [27]. Bacteria, eukaryota, viruses, and

archaea were included in the LCA calculation. Taxonomic biomass was quantified by summing

the intensities of the peptides corresponding to each taxon. A Bray-Curtis dissimilarity-based

approach [28] was applied for evaluating the variation of genus-level biomass contributions

between drug-treated and DMSO control groups. Calculation of the Bray-Curtis distance was

performed using the R package "vegan"[29].

3） Functional analysis:

The quantified protein groups were first filtered according to the criteria that the protein appears

in >80% of the microbiomes with at least one drug treatment. Then LFQ protein group intensities

of the filtered file was $\log_2$-transformed and normalized through quotient transformation

(x/mean) using the R package clusterSim. Then, LFQ protein group intensities were processed

by a ComBat process [54, 55] using iMetalab.ca [56] to remove possible batch effects between

individual microbiomes. Using the ComBat-corrected data, an unsupervised non-linear

dimensionality reduction algorithm, t-distributed stochastic neighbor embedding (t-SNE)[30]

was then applied to visualize similarities between samples using the R package Rtsne. Parameter

for the function Rtsne() were, perplexity=10, max_iter = 1200 (number of iterations), other

parameters were set as default. The R function geom_polygon implemented in ggplot2 was used

to visualize the t-SNE results.

Functional annotations of protein groups, including COG and KEGG information, were obtained

in the MetaLab software. In addition, KEGG ortholog (KO) annotation of protein FASTA

sequences was conducted using GhostKOALA (https://www.kegg.jp/ghostkoala/)[57]. $\log_2$

fold-change of each drug-treated sample relative to the corresponding DMSO control was

calculated using the abundances of proteins annotated to COG categories and COGs. Functional

enrichment analysis was performed using the enrichment module on iMetalab.ca through

inputting the list of COG functional proteins. Adjusted *p*-value cutoff was set at 0.05 for the

enrichment analysis.

Statistical analysis

We examined data distribution on all levels of data, and results indicated non-normal

distributions of the dataset (examples shown in **Supplementary Figures S10 and 11**). Hence, a

non-parametric statistical hypothesis test, the Wilcoxon rank sum test, was applied in statistical

analyses. For multiple comparisons, *p*-values were adjusted using the Benjamini-Hochberg false

discovery rate (FDR) procedure[58]. For multivariate analysis, partial least-squares discriminant

analyses (PLS-DA) based on ComBat-corrected protein group intensities were performed using

MetaboAnalyst (http://www.metaboanalyst.ca/)[59]. PLS-DA model were evaluated by cross-

validation of R^2 and Q^2.

Data visualizations

Box plots, violin plots, hierarchical clustering, 3D scatter plots, heatmaps, PCA, and t-SNE were

visualized using R packages ggplot2, gridExtra, scatterplot3d, and pheatmap. Pathway maps

were visualized using iPATH 3 (https://pathways.embl.de/)[60] and Pathview Web

(https://pathview.uncc.edu/)[61]. Stacked column bars and functional enrichments were

visualized on iMetaLab.ca.

List of abbreviations

COG	Clusters of orthologous groups
DMSO	Dimethyl sulfoxide
FDR	False-discovery rate
FOS	Fructooligosaccharide
GABA	Gamma-Aminobutyric Acid
KEGG	Kyoto Encyclopedia of Genes and Genomes
LC-MS/MS	Liquid chromatography–tandem mass spectrometry
LFQ	Label-free quantification
NSAIDs	Nonsteroidal anti-inflammatory drugs
PCA	Principle component analysis
PLS-DA	Partial least squares discriminant analysis
POC	Proof-of-concept
VIP	Variable importance in projection

Declarations

Ethics approval and consent to participate

The Research Ethics Board protocol (# 20160585-01H) for stool sample collection was approved
by the Ottawa Health Science Network Research Ethics Board at the Ottawa Hospital. All
volunteers signed consent forms.

Consent for publication

Not applicable.

Competing interests

D.F. and A.S. have co-founded Biotagenics and MedBiome, clinical microbiomics companies. All other authors declare no competing interests.

Funding

This work was supported by the Government of Canada through Genome Canada and the Ontario Genomics Institute (OGI-114 and OGI-149), CIHR grant (ECD-144627), the Natural Sciences and Engineering Research Council of Canada (NSERC, grant no. 210034), and the Ontario Ministry of Economic Development and Innovation (REG1-4450).

Author contributions

D.F., A.S. and L.L. designed the study. J.M. coordinated the volunteers and collected the samples. L.L., X.Z. and Z.N. developed the workflow and performed the experiments. L.L., Z.N., X.Z. and K.C. analyzed and visualized the data. L.L, D.F. and A.S. wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgments

The authors wish to thank Dr. Kendra Hodgkinson for editing the manuscript.

References:

1. Zimmermann M, Zimmermann-Kogadeeva M, Wegmann R, Goodman AL: Separating host and microbiome contributions to drug pharmacokinetics and toxicity. Science 2019; 363:eaat9931.

2. Lynch SV, Pedersen O: The human intestinal microbiome in health and disease. N Engl J Med 2016; 375:2369-79.

3. Jia W, Li H, Zhao L, Nicholson JK: Gut microbiota: a potential new territory for drug targeting. Nat Rev Drug Discov 2008; 7:123.

4. Le Bastard Q, Al-Ghalith GA, Grégoire M, Chapelet G, Javaudin F, Dailly E, Batard E, Knights D, Montassier E: Systematic review: human gut dysbiosis induced by non-antibiotic prescription medications. Aliment Pharmacol Ther 2017; 47:332-45.

5. Urquhart L: FDA new drug approvals in Q2 2018. Nat Rev Drug Discov 2018; 17:536.

6. Harvey AL, Edrada-Ebel R, Quinn RJ: The re-emergence of natural products for drug discovery in the genomics era. Nat Rev Drug Discov 2015; 14:111.

7. McDonald JAK, Schroeter K, Fuentes S, Heikamp-deJong I, Khursigara CM, de Vos WM, Allen-Vercoe E: Evaluation of microbial community reproducibility, stability and composition in a human distal gut chemostat model. J Microbiol Methods 2013; 95:167-74.

8. Van den Abbeele P, Belzer C, Goossens M, Kleerebezem M, De Vos WM, Thas O, De Weirdt R, Kerckhof F-M, Van de Wiele T: Butyrate-producing *Clostridium cluster* XIVa species specifically colonize mucins in an *in vitro* gut model. ISME J 2012; 7:949.

9. Van de Wiele T, Van den Abbeele P, Ossieur W, Possemiers S, Marzorati M: The Simulator of the Human Intestinal Microbial Ecosystem (SHIME®). In *The Impact of Food Bioactives on Health: in vitro and ex vivo models.* Edited by Verhoeckx K, Cotter P, López-Expósito I, Kleiveland C, Lea T, Mackie A, Requena T, Swiatecka D, Wichers H. Cham: Springer International Publishing; 2015: 305-17

10. Auchtung JM, Robinson CD, Britton RA: Cultivation of stable, reproducible microbial communities from different fecal donors using minibioreactor arrays (MBRAs). Microbiome 2015; 3:42.

11. Maier L, Pruteanu M, Kuhn M, Zeller G, Telzerow A, Anderson EE, Brochado AR, Fernandez KC, Dose H, Mori H, et al: Extensive impact of non-antibiotic drugs on human gut bacteria. Nature 2018; 555:623.

12.   Props R, Kerckhof F-M, Rubbens P, De Vrieze J, Hernandez Sanabria E, Waegeman W, Monsieurs P,
Hammes F, Boon N: Absolute quantification of microbial taxon abundances. ISME J 2017; 11:584-87.

13.   Jian C, Luukkonen P, Yki-Järvinen H, Salonen A, Korpela K: Quantitative PCR provides a simple and
accessible method for quantitative microbiome profiling. bioRxiv 2018:478685.

14.   Ranjan R, Rani A, Metwally A, McGee HS, Perkins DL: Analysis of the microbiome: advantages of whole
genome shotgun versus 16S amplicon sequencing. Biochem Biophys Res Commun 2016; 469:967-77.

15.   Bashiardes S, Zilberman-Schapira G, Elinav E: Use of metatranscriptomics in microbiome research.
Bioinform Biol Insights 2016; 10:19-25.

16.   Milo R: What is the total number of protein molecules per cell volume? A call to rethink some published
values. BioEssays 2013; 35:1050-55.

17.   Mills RH, Vázquez-Baeza Y, Zhu Q, Jiang L, Gaffney J, Humphrey G, Smarr L, Knight R, Gonzalez DJ:
Evaluating Metagenomic Prediction of the Metaproteome in a 4.5-Year Study of a Patient with Crohn's
Disease. mSystems 2019; 4:e00337-18.

18.   Liu Y, Beyer A, Aebersold R: On the Dependency of Cellular Protein Levels on mRNA Abundance. Cell
2016; 165:535-50.

19.   Zhang X, Deeke SA, Ning Z, Starr AE, Butcher J, Li J, Mayne J, Cheng K, Liao B, Li L, et al:
Metaproteomics reveals associations between microbiome and intestinal extracellular vesicle proteins in
pediatric inflammatory bowel disease. Nat Comm 2018; 9:2873.

20.   Zhang X, Chen W, Ning Z, Mayne J, Mack D, Stintzi A, Tian R, Figeys D: Deep metaproteomics approach
for the study of human microbiomes. Anal Chem 2017; 89:9407-15.

21.   Kleiner M: Metaproteomics: Much More than Measuring Gene Expression in Microbial Communities.
mSystems 2019; 4:e00115-19.

22.   Kleiner M, Thorson E, Sharp CE, Dong X, Liu D, Li C, Strous M: Assessing species biomass contributions
in microbial communities via metaproteomics. Nat Comm 2017; 8:1558.

23.   Zhang X, Figeys D: Perspective and Guidelines for Metaproteomics in Microbiome Studies. J Proteome
Res 2019.

24.   Li L, Abou-Samra E, Ning Z, Zhang X, Mayne J, Wang J, Cheng K, Walker K, Stintzi A, Figeys D: An *in
vitro* model maintaining taxon-specific functional activities of the gut microbiome. bioRxiv 2019:616656.

25.   Zhang X, Li L, Mayne J, Ning Z, Stintzi A, Figeys D: Assessing the impact of protein extraction methods
for human gut metaproteomics. J Proteom 2018; 180:120-27.

26.   Zhang X, Ning Z, Mayne J, Moore JI, Li J, Butcher J, Deeke SA, Chen R, Chiang C-K, Wen M, et al:
MetaPro-IQ: a universal metaproteomic approach to studying human and mouse gut microbiota.
Microbiome 2016; 4:31.

27.   Cheng K, Ning Z, Zhang X, Li L, Liao B, Mayne J, Stintzi A, Figeys D: MetaLab: an automated pipeline
for metaproteomic data analysis. Microbiome 2017; 5:157.

28.   Consortium THMP, Huttenhower C, Gevers D, Knight R, Abubucker S, Badger JH, Chinwalla AT, Creasy
HH, Earl AM, FitzGerald MG, et al: Structure, function and diversity of the healthy human microbiome.
Nature 2012; 486:207.

29.   vegan: Community ecology package. R package version 2.5-1. https://CRAN.R-
project.org/package=vegan

30.   Maaten LJPvd, Hinton GE: Visualizing high-dimensional data using t-SNE. J Mach Learn Res 2008;
9:2579-605.

31.   Rinehart KL, Shield LS: Chemistry of the ansamycin antibiotics. In *Fortschritte der Chemie Organischer
Naturstoffe / Progress in the Chemistry of Organic Natural Products.* Edited by Cimino G, Coates RM, De
Stefano S, Fontana A, Hemmerich P, Minale L, Rinehart KL, Shield LS, Sodano G, Toniolo C*, et al*.
Vienna: Springer Vienna; 1976: 231-307

32.   Mayes PA, Bender DA: The pentose phosphate pathway & other pathways of hexose metabolism. In
Harper's illustrated biochemistry. 26 edition. Edited by Murray RK, Granner DK, Mayes PA, Rodwell
VW. New York: Lange Medical Books/McGraw-Hill; 2003: 163-72

33.   Brodribb AJ, Groves C: Effect of bran particle size on stool weight. Gut 1978; 19:60.

34.   Rogers MAM, Aronoff DM: The influence of non-steroidal anti-inflammatory drugs on the gut
microbiome. Clin Microbiol Infect 2016; 22:178.e1-78.e9.

35.   Zhang X, Zhao Y, Zhang M, Pang X, Xu J, Kang C, Li M, Zhang C, Zhang Z, Zhang Y, et al: Structural
680   changes of gut microbiota during berberine-mediated prevention of obesity and insulin resistance in high-
681   fat diet-fed rats. PLOS ONE 2012; 7:e42529.

682   36.   Zhu L, Zhang D, Zhu H, Zhu J, Weng S, Dong L, Liu T, Hu Y, Shen X: Berberine treatment increases
683   Akkermansia in the gut and improves high-fat diet-induced atherosclerosis in Apoe−/− mice.
684   Atherosclerosis 2018; 268:117-26.

685   37.   Charteris WP, Kelly PM, Morelli L, Collins JK: Antibiotic susceptibility of potentially probiotic
686   Bifidobacterium isolates from the human gastrointestinal tract. Lett Appl Microbiol 1998; 26:333-37.

687   38.   Rossi M, Corradini C, Amaretti A, Nicolini M, Pompei A, Zanoni S, Matteuzzi D: Fermentation of
688   fructooligosaccharides and inulin by bifidobacteria: a comparative study of pure and fecal cultures. Appl
689   Environ Microbiol 2005; 71:6150-58.

690   39.   Marx SP, Winkler S, Hartmeier W: Metabolization of β-(2,6)-linked fructose-oligosaccharides by different
691   bifidobacteria. FEMS Microbiol Lett 2000; 182:163-69.

692   40.   Cryan JF, Dinan TG: Mind-altering microorganisms: the impact of the gut microbiota on brain and
693   behaviour. Nat Rev Neurosci 2012; 13:701.

694   41.   Burokas A, Arboleya S, Moloney RD, Peterson VL, Murphy K, Clarke G, Stanton C, Dinan TG, Cryan JF:
695   Targeting the microbiota-gut-brain axis: Prebiotics have anxiolytic and antidepressant-like effects and
696   reverse the impact of chronic stress in mice. Biol Psychiatry 2017; 82:472-87.

697   42.   Kawaguchi M, Tashiro Y, Adachi T, Tamura Z: Changes in intestinal condition, fecal microflora and
698   composition of rectal gas after administration of fructooligosaccharide and lactulose at different doses.
699   Bifidobacteria and Microflora 1993; 12:57-67.

700   43.   Swanson KS, Grieshop CM, Flickinger EA, Bauer LL, Chow J, Wolf BW, Garleb KA, Fahey JGC:
701   Fructooligosaccharides and *Lactobacillus acidophilus* modify gut microbial populations, total tract nutrient
702   digestibilities and fecal protein catabolite concentrations in healthy adult dogs. J Nutr 2002; 132:3721-31.

703   44.   Lei XJ, Cheong JY, Park JH, Kim IH: Supplementation of protease, alone and in combination with
704   fructooligosaccharide to low protein diet for finishing pigs. Animal Sci J 2017; 88:1987-93.

705   45.   Albesa I, Becerra MC, Battán PC, Páez PL: Oxidative stress involved in the antibacterial action of different
706   antibiotics. Biochem Biophys Res Commun 2004; 317:605-09.

707   46.   Becerra MC, Albesa I: Oxidative stress induced by ciprofloxacin in *Staphylococcus aureus*. Biochem
708   Biophys Res Commun 2002; 297:1003-07.

709   47.   Becerra MC, Páez PL, Laróvere LE, Albesa I: Lipids and DNA oxidation in *Staphylococcus aureus* as a
710   consequence of oxidative stress generated by ciprofloxacin. Mol Cell Biochem 2006; 285:29-34.

711   48.   Ranford JC, Henderson B: Chaperonins in disease: mechanisms, models, and treatments. Mol Pathol 2002;
712   55:209-13.

713   49.   Guirro M, Costa A, Gual-Grau A, Mayneris-Perxachs J, Torrell H, Herrero P, Canela N, Arola L: Multi-
714   omics approach to elucidate the gut microbiota activity: Metaproteomics and metagenomics connection.
715   Electrophoresis 2018; 39:1692-701.

716   50.   Thompson A, Schäfer J, Kuhn K, Kienle S, Schwarz J, Schmidt G, Neumann T, Hamon C: Tandem mass
717   tags: a novel quantification strategy for comparative analysis of complex protein mixtures by MS/MS.
718   Anal Chem 2003; 75:1895-904.

719   51.   Shah P, Fritz JV, Glaab E, Desai MS, Greenhalgh K, Frachet A, Niegowska M, Estes M, Jäger C, Seguin-
720   Devaux C, et al: A microfluidics-based in vitro model of the gastrointestinal human–microbe interface. Nat
721   Comm 2016; 7:11535.

722   52.   Kim HJ, Lee J, Choi J-H, Bahinski A, Ingber DE: Co-culture of living microbiome with microengineered
723   human intestinal villi in a gut-on-a-chip microfluidic device. J Vis Exp 2016:54344.

724   53.   Jalili-Firoozinezhad S, Gazzaniga FS, Calamari EL, Camacho DM, Fadel CW, Bein A, Swenor B, Nestor
725   B, Cronce MJ, Tovaglieri A, et al: A complex human gut microbiome cultured in an anaerobic intestine-on-
726   a-chip. Nat Biomed Eng 2019.

727   54.   Chen C, Grennan K, Badner J, Zhang D, Gershon E, Jin L, Liu C: Removing batch effects in analysis of
728   expression microarray data: an evaluation of six batch adjustment methods. PLOS ONE 2011; 6:e17238.

729   55.   Nyamundanda G, Poudel P, Patil Y, Sadanandam A: A novel statistical method to diagnose, quantify and
730   correct batch effects in genomic studies. Sci Rep 2017; 7:10849.

56.   Liao B, Ning Z, Cheng K, Zhang X, Li L, Mayne J, Figeys D: iMetaLab 1.0: a web platform for
metaproteomics data analysis. Bioinformatics 2018:bty466-bty66.

57.   Kanehisa M, Sato Y, Morishima K: BlastKOALA and GhostKOALA: KEGG tools for functional
characterization of genome and metagenome sequences. J Mol Bio 2016; 428:726-31.

58.   Benjamini Y, Hochberg Y: Controlling the false discovery rate: a practical and powerful approach to
multiple testing. J Royal Stat Soc 1995; 57:289-300.

59.   Chong J, Soufan O, Li C, Caraus I, Li S, Bourque G, Wishart DS, Xia J: MetaboAnalyst 4.0: towards more
transparent and integrative metabolomics analysis. Nucleic Acids Res 2018; 46:W486-W94.

60.   Yamada T, Letunic I, Okuda S, Kanehisa M, Bork P: iPath2.0: interactive pathway explorer. Nucleic Acids
Res 2011; 39:W412-W15.

61.   Luo W, Pant G, Bhavnasi YK, Blanchard JSG, Brouwer C: Pathview Web: user friendly pathway
visualization and data integration. Nucleic Acids Res 2017; 45:W501-W08.

RapidAIM: A culture- and metaproteomics-based Rapid Assay of Individual Microbiome responses to drugs

Leyuan Li[1], Zhibin Ning[1], Xu Zhang[1], Janice Mayne[1], Kai Cheng[1], Alain Stintzi*[1], Daniel Figeys*[1,2,3]

[1] Department of Biochemistry, Microbiology and Immunology, Ottawa Institute of Systems Biology, Faculty of Medicine, University of Ottawa, Ottawa, Canada

[2] Department of Chemistry and Biomolecular Sciences, University of Ottawa, Ottawa, Canada

[3] Canadian Institute for Advanced Research, Toronto, Canada

* Corresponding Authors, Email: dfigeys@uottawa.ca (DF), and astintzi@uottawa.ca (AS)

Supplementary figures

Supplementary Figure S1

1. Culturing in 96-deepwell
2. Bacterial cell purification
3. Ultrasonic cell disruption
4. In-solution digestion
5. LC-MS/MS analysis

1

Peptides identified — NS — 1 ml / 2 ml

Protein groups identified — NS — 1 ml / 2 ml

Cultured volume reduced from 2 ml (tube-based) to 1 ml (96-deepwell-based) → No significant difference in metaproteomic identification between 1 ml- and 2 ml- based protein extractions

2

Supernatant

3 cycles
300 x g, 10 min

Supernatant

2,270 x g, 1 hr

3 cycles
wash with PBS

14,000 x g, 20 min

Limitation of centrifugation speed when using a 96-well plate

OD600 vs Concentration
- EP tubes
- 96 wells

No significant difference in bacterial concentration between the two cell purification workflows

3 **4**

Ultrasonication
200 ul 8M urea
30 s x 4 cycles

Debris removal
16,000 x g centrifugation?

DC assay
50 µg proteins for digestion

Digestion
2 µl DTT, 56 °C, 30 min
2 µl IAA, RT dark, 40 min
1 µg Trypsin, 37 °C 18 hr

Desalting
C18 beads
+
96-channel filter tips

Limitation of centrifugation speed when using a 96-well plate

5

LC-MS/MS analysis — Equal-volume based sample loading

Peptides identified — NS — Centrifugation / No centrifugation

Protein groups identified — NS — Centrifugation / No centrifugation

Centrifugation / No centrifugation heatmap

- 6-phosphofructokinase
- ABC-type glycerol-3-phosphate transport system
- Acetylornithine/ succinyldiaminopimelate/ putrescine aminotransferase
- Glutamyl- or glutaminyl- tRNA synthetase
- Ribosomal proteins
- Sugar kinase of the NBD/HSP70 family
- TRAP-type uncharacterized transport system

Whole-workflow validation

Peptides identified — NS — Tube / 96 Wells

Protein groups identified — NS — Tube / 96 wells

Figure S1. Establishment and step-by-step validation of the microplate-based metaproteomic sample
preparation workflow of the RapidAIM assay.
After culturing in a 96-well deepwell plate **[1]**, the bacterial cells were washed with PBS **[2]**, then disrupted with
four cycles of 30 s ultrasonication **[3]**. Protein concentration was measured in the DMSO control sample using a DC
protein assay. This concentration was used to calculate a volume equivalent to 50 μg proteins in the control sample.
This volume was taken from every sample and digested with trypsin **[4]**, then desalted with a panel of 96 filter tips
packed with C18 beads and equal volumes of each sample were analyzed by LC-MS/MS **[5]**.
Due to a few differences in the metaproteomic procedure compared with tube-based protocols, we examined the
effect of protocol differences on sample quality step-by-step. In previously published protocols, samples were
cultured in 2 ml culture tubes[1,2]. In step **[1]**, culturing samples in a 96-well plate reduced the sample size to 1 ml. No
significant difference was shown by *t*-test in peptide and protein group identification numbers between protein
extractions from 1 ml and 2 ml samples. For step **[2]**, the 96-deepwell plates limited centrifugation speed to 2,270 g
(versus 14,000 g in the original protocol), so we extended the centrifugation time from 20 min to 1 hour and tested
whether bacterial concentration was affected by the altered centrifugation process[1,3]. Concentration of purified
bacterial cells were compared by OD_{600} after being re-suspended in 1 ml, 2 ml, 4 ml, 8 ml, and 16 ml PBS. No
significant difference in OD_{600} reads was observed between the two cell purification protocols. In step **[3]**, the
original protocol used a high-speed centrifugation (16,000 g) to remove cell debris after the ultrasonication. Due to
the limitation of centrifugation speed when using a microplate, we compared metaproteomic profiles of the sample
when digested with or without cell debris removal. No significant differences were found in the number of protein
identifications. In terms of differentially identified proteins, we found that the samples without cell debris removal
resulted in more identifications of cell-membrane proteins such as the ABC-type transport systems, as well as
cytoskeleton-related proteins, such as the translation elongation factor EF-Tu[4,5], 6-phosphofructokinase[6], and
ribosomes[7]. Therefore, we infer that eliminating the centrifugation process could reduce the removal of cytoskeleton
and cell membrane proteins. For step **[4]**, no validation was necessary as the same type of filter tips were used in
both protocols. Finally, we performed a whole-workflow comparison of the metaproteomic outcomes between
traditional tube-based and 96-well-based processes. The microplate-based metaproteomic workflow showed no
statistically significant difference in peptide and protein identification compared with the tube-based workflow.
804

Supplementary Figure S2

a

b

Figure S2. Assessment of the equal-volume digestion and LC-MS/MS analysis strategy.

(**a**) In triplicates, six dilutions of a single microbiome sample were subjected to this equal-volume based analysis. The LFQ intensities of protein groups showed Pearson's correlation coefficient $r > 0.95$ between most dilutions, but a lower r was seen in the samples with the lowest concentration. (**b**) Comparison of taxonomic biomass contributions on different levels (summed peptide intensity assigned to different taxa) among diluted groups suggested very low level of bias. GRD1-6 are six different dilution gradients (protein concentration shown in Figure 1b), and t1-t3 are technical replicates of the same conditions.

Supplementary Figure S3

Figure S3. Reproducibility of RapidAIM assay on different levels.

(**a**) Clustering using protein group composition information showed that triplicates of drug treatments were closely clustered. A hierarchical tree was generated based on Pearson's correlation coefficient. The cluster corresponding to the gray box contained DMSO control samples; samples in this cluster indicated drugs that had very weak effects on the microbiome. (**b**) Box chart suggesting that biomass effects by drugs are highly reproducible. (**c**) Examples showing reproducible functional responses to drugs at the enzyme level. COC0031 and COG2873 are cysteine synthase and O-acetylhomoserine/O-acetylserine sulfhydrylase (pyridoxal phosphate-dependent), respectively; both enzymes are involved in the conversion of sulfide to L-cysteine[8]. COG2221 is dissimilatory sulfite reductase (desulfoviridin), alpha and beta subunits; this enzyme reduces sulfite to sulfide[9]. (**d**) Examples showing reproducible taxonomic responses to drugs at the species level. *F. prausnitzii* is a ubiquitous bacterium of the intestinal microbiota[10]; *B. worthia* is a taurine-degrading bacterium which can reduce sulfite to sulfide by dissimilatory sulfite reductase[9]. COG2221 and *B. worthia* show a correlation in their response to different drugs. Box spans interquartile range (25th to 75th percentile), and line within box denotes median.

Supplementary Figure S4

Individual microbiome sample V1:

DMSO RFXM

Figure S4. Case study on microbiome V1's response to rifaximin (RFXM). We have shown that the antibiotic rifaximin did not show overall biomass inhibition on microbiome sample V1 (Figure 2a and Supplementary Figure S3b). We examined whether the expressions of antibiotic resistance proteins were affected. Briefly, the LC-MS/MS raw files were specifically searched against the Structured Antibiotic Resistance Genes (SARG) database 48 which highlighted 118 antibiotic resistance protein groups across the dataset. (**a**) We found an increase in relative abundance of total antibiotic resistance proteins in microbiome V1 in response to rifaximin. (**b and c**) Particularly, antibiotic resistance peptide sequences unique to Proteobacteria and Firmicutes were increased. (**d**) Moreover, despite no significant change in total microbiome biomass in V1, a significant 6.5-fold increase in the relative biomass of Proteobacteria in the whole microbial community was observed. (**e**) Non-parametric test resulted in seven significantly increased antibiotic resistance protein groups (FDR-adjusted p value<0.05). These protein groups belonged predominantly to Proteobacteria (5 out of 7). Increase of Proteobacteria is associated with dysbiosis in gut microbiota 49. These together suggested a potential risk of rifaximin administration in individual V1. (*p* values were based on two-sided Wilcoxon test; box spans interquartile range (25th to 75th percentile), and line within box denotes median.

Supplementary Figure S5

Figure S5. Log₂ fold-change of relative abundance at the genus level in response to each drug compared with the DMSO control. Genera that existed in ≥80% of the volunteers are shown. Star (*) indicate significantly changed bacterial abundance by Wilcoxon test, $p <0.05..$

 Supplementary Figure S6

860 **Figure S6. Score plots and cross-validations of seven PLS-DA models**

861 PLS-DA models of microbiomes responses to each compound were established using MetaboAnalyst 4.0. PLS-DA

862 model qualities were assessed through cross-validation, and accuracy, R^2 and Q^2 were given for each model. Seven

863 compounds were found with valid PLS-DA models distinguishing the effect of the compound from the DMSO

864 control.

Supplementary Figure S7

Figure S7. Log₂ fold-change of functions at the COG protein level

On the COG functional protein level, 535 COGs were significantly decreased by at least one drug treatment. The 15 COGs that were affected by ≥ 10 compounds are shown in this figure. Statistical significance was evaluated by one-sided Wilcoxon rank sum test, FDR-adjusted p-values: *, $p < 0.05$; **, $p < 0.01$.

878   **Supplementary Figure S8**

879
880   **Figure S8. String interaction of COG functional proteins significantly stimulated by diclofenac**

 Supplementary Figure S9

Figure S9. Phylum-specific functional responses to Berberine

Protein groups with VIP scores of >1 were annotated to phyla and COGs. Up- and down-regulated (red and green

lines) COGs corresponding to different phyla were illustrated on a metabolic pathway map using iPath 3. Pathway

maps for each phylum were combined, and overlapped pathways were shown in black lines. Our data suggest that

different phyla responded differently at a functional pathway level.

Supplementary Figure S10

892
893 **Figure S10. Randomly selected LFQ intensities of protein groups showing heavy tailed distribution on the Q-**
894 **Q plots.**

895
896

Supplementary Figure S11

Figure S11. Randomly selected log₂-fold changes of COGs showing heavy tailed distribution on the Q-Q plots.

References

1.   Li, L.*, et al.* Evaluating *in vitro* culture medium of gut microbiome with orthogonal experimental design and a metaproteomics approach. *Journal of Proteome Research* **17**, 154-163 (2018).

2.   Zhang, X.*, et al. In vitro* metabolic labeling of intestinal microbiota for quantitative metaproteomics. *Analytical Chemistry* **88**, 6120-6125 (2016).

3.   Zhang, X.*, et al.* Assessing the impact of protein extraction methods for human gut metaproteomics. *Journal of Proteomics* **180**, 120-127 (2018).

4.   Defeu Soufo, H.J., Reimold, C., Breddermann, H., Mannherz, H.G. & Graumann, P.L. Translation elongation factor EF-Tu modulates filament formation of actin-like MreB protein *in vitro*. *Journal of Molecular Biology* **427**, 1715-1727 (2015).

5.   Mayer, F. Cytoskeletal elements in bacteria *Mycoplasma pneumoniae*, *Thermoanaerobacterium* sp., and *Escherichia coli* as revealed by electron microscopy. *Journal of Molecular Microbiology and Biotechnology* **11**, 228-243 (2006).

6.   Vértessy, B.G., Orosz, F., Kovács, J. & Ovádi, J. Alternative binding of two sequential glycolytic enzymes to microtubules: molecular studies in the phosphofructokinase/aldolase/microtubule system. *Journal of Biological Chemistry* **272**, 25542-25546 (1997).

7.   Mayer, F. Cytoskeletons in prokaryotes. *Cell Biology International* **27**, 429-438 (2013).

8.   Lithgow, J.K., Hayhurst, E.J., Cohen, G., Aharonowitz, Y. & Foster, S.J. Role of a cysteine synthase in *Staphylococcus aureus*. *Journal of Bacteriology* **186**, 1579 (2004).

922   9.   Laue, H., Friedrich, M., Ruff, J. & Cook, A.M. Dissimilatory sulfite reductase (desulfoviridin) of the
923   taurine-degrading, non-sulfate-reducing bacterium *Bilophila wadsworthia* RZATAU contains a fused
924   DsrB-DsrD subunit. *Journal of Bacteriology* **183**, 1727 (2001).
925   10.   Miquel, S*., et al. Faecalibacterium prausnitzii* and human intestinal health. *Current Opinion in*
926   *Microbiology* **16**, 255-261 (2013).
927
928
929
930