Title**:**

Insulin-stimulated glucose uptake partly relies on p21-activated kinase (PAK)-2, but not

PAK1, in mouse skeletal muscle

Lisbeth L. V. Møller[1], Merna Jaurji[1], Rasmus Kjøbsted[1], Giselle A. Joseph[2], Agnete B. Madsen[1],

Jonas R. Knudsen[1], Annemarie Lundsgaard[1], Nicoline R. Andersen[1], Peter Schjerling[3,4], Thomas E.

Jensen[1], Robert S. Krauss[2], Erik A. Richter[1*], and Lykke Sylow[1,*,#]

[1] Section of Molecular Physiology, Department of Nutrition, Exercise and Sports, Faculty of Science, University of

Copenhagen, DK-2100 Copenhagen, Denmark

[2] Department of Cell, Developmental, and Regenerative Biology, Icahn School of Medicine at Mount Sinai, New York,

NY 10029, USA

[3] Institute of Sports Medicine, Department of Orthopaedic Surgery M, Bispebjerg Hospital, DK-2400 Copenhagen,

Denmark

[4] Center for Healthy Aging, Faculty of Health Sciences, University of Copenhagen, DK-2100 Copenhagen, Denmark

[*] *Authors contributed equally*

[#]*Corresponding author and person to who reprint requests should be addressed:*

Lykke Sylow. E-mail: Lshansen@nexs.ku.dk. Tel.: +4520955250

Section of Molecular Physiology, Department of Nutrition, Exercise and Sports, August Krogh

Building, 13 Universitetsparken, DK-2100 Copenhagen, Denmark

Short title: Group I PAKs and insulin-stimulated skeletal muscle glucose uptake

Abstract

Objective: Skeletal muscle glucose uptake is essential for maintaining whole-body glucose homeostasis and accounts for the majority of glucose disposal in response to insulin. The group I p21-activated kinase (PAK) isoforms PAK1 and PAK2 are activated in response to insulin in skeletal muscle. Interestingly, PAK1/2 signalling is impaired in insulin-resistant mouse and human skeletal muscle and PAK1 has been suggested to be required for insulin-stimulated GLUT4 translocation. However, the relative contribution of PAK1 and PAK2 to insulin-stimulated glucose uptake in mature skeletal muscle is unresolved. The aim of the present investigation was to determine the requirement for PAK1 and PAK2 in whole-body glucose homeostasis and insulin-stimulated glucose uptake in skeletal muscle.

Methods: Glucose uptake was measured in isolated skeletal muscle incubated with a pharmacological inhibitor (IPA-3) of group I PAKs and in muscle from whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO and double whole-body PAK1 and muscle-specific PAK2 knockout mice.

Results: The whole-body respiratory exchange ratio was largely unaffected by lack of PAK1 and/or PAK2. Whole-body glucose tolerance was mildly impaired in PAK2 mKO, but not PAK1 KO mice. IPA-3 partially reduced (-20%) insulin-stimulated glucose uptake in mouse soleus muscle. In contrast to a previous study of GLUT4 translocation in PAK1 KO mice, PAK1 KO muscles displayed normal insulin-stimulated glucose uptake *in vivo* and in isolated muscle. On the contrary, glucose uptake was slightly reduced in response to insulin in glycolytic extensor digitorum longus muscle lacking PAK2, alone (-18%) or in combination with PAK1 KO (-12%).

Conclusions: Insulin-stimulated glucose uptake partly relies on PAK2, but not PAK1, in mouse skeletal muscle. Thus, the present study challenges that group I PAKs, and especially PAK1, are

major regulators of whole-body glucose homeostasis and insulin-stimulated glucose uptake in

skeletal muscle.

Keywords

Skeletal muscle; Insulin; Glucose uptake; GLUT4 translocation; p21-activated kinases; Metabolism

Abbreviations

2DG, 2-Deoxyglucose; AUC, area under the curve; BSA, bovine serum albumin; BW, body weight;

dKO, double knockout; EDL, extensor digitorum longus; FM, fat mass; GLUT4, glucose

transporter 4; GTT, glucose tolerance test; HFD, high-fat diet; HOMA-IR, homeostatic model

assessment of insulin resistance; ITT, insulin tolerance test; i.p., intraperitoneal; KO, knockout; L6-

GLUT4myc, rat L6 skeletal muscle cells overexpressing myc-tagged GLUT4; LBM, lean body

mass; mKO, muscle-specific knockout; NOX, NADPH oxidase; PAK, p21-activated kinase; PI3K,

phosphoinositide 3-kinase; RER, respiratory exchange ratio; r.o., retro-orbital; VO2, oxygen uptake

1. Introduction

Skeletal muscles account for the majority of insulin-mediated whole-body glucose disposal [1,2] and muscle insulin resistance is an early defect in the pathophysiology of peripheral insulin resistance and type 2 diabetes mellitus [1,3]. As diabetes globally is approaching epidemic proportions, it is important to understand the mechanisms regulating glucose uptake by skeletal muscle.

Insulin stimulates glucose uptake in skeletal muscle by activation of a signalling cascade that leads to the translocation of glucose transporter (GLUT)4-containing vesicles to the sarcolemma and transverse tubuli [4]. This signalling cascade has been proposed to include activation of p21-activated kinase 1 (PAK1) downstream of PI3K [5–7]. PAKs are serine/threonine kinases and involved in numerous signalling networks regulating essential cellular activities, including cell proliferation, differentiation, apoptosis, and cytoskeleton dynamics [8–11]. Group I PAKs (PAK1-3) are downstream targets of the Rho GTPases Cdc42 and Rac1 [12]. Previous studies suggest that only PAK1 and PAK2 are expressed in skeletal muscle, whereas PAK3 mRNA and protein expression is below the detection limit [6,13,14]. In muscle cells and mouse skeletal muscle, PAK1 is proposed to be required for GLUT4 translocation in response to insulin stimulation [6,7], likely downstream of Rac1 [15–20]. Thus, together with Akt, proposed to regulate GLUT4 translocation via phosphorylation of the Rab GTPase-activating protein TBC1D4 [21–23], Rac1-PAK1 activation is suggested to be necessary for insulin-stimulated GLUT4 translocation.

Upon activation of PAK1 and PAK2, conformational changes allow autophosphorylation of T423 and T402, respectively, thereby relieving the autoinhibition of PAK1 and PAK2 [12,24,25]. In vastus lateralis muscle from subjects with obesity and type 2 diabetes, phosphorylation of PAK1/2 at T423/402 in response to insulin was 50% reduced compared to healthy controls [18]. Likewise,

insulin-stimulated pPAK1/2 T423/402 was diminished in palmitate-treated insulin-resistant L6

myotubes, even though upstream of PAK1/2, insulin-stimulated Rac1-GTP binding (i.e. activation)

was not impaired [26]. Together, these studies [18,26] associate dysregulated activity of PAK1 and

PAK2 with insulin resistance. In addition, a pharmacological inhibitor of group I PAKs, IPA-3

abolished insulin-stimulated GLUT4 translocation and glucose uptake in L6 myoblasts and

myotubes overexpressing myc-tagged GLUT4 (L6-GLUT4myc), respectively [6]. This indicates

that group I PAKs are required for insulin-stimulated glucose uptake. The effect of IPA-3 has

largely been ascribed to inhibition of PAK1, as whole-body genetic ablation of PAK1 in mice

impaired glucose tolerance [7,27] and blocked insulin-stimulated GLUT4 translocation in skeletal

muscle [7]. Further supporting PAK1 being the major PAK isoform regulating GLUT4

translocation, insulin-stimulated GLUT4 translocation was unaffected by a 75% knockdown of

PAK2 in L6-GLUT4myc myoblasts [6]. The suggested downstream mechanisms whereby PAK1

regulates GLUT4 translocation include simultaneous cofilin-mediated actin depolymerization and

N-WASP-cortactin-mediated actin polymerization [6,28,29].

Although such studies implicate group I PAKs, and in particular PAK1, in the regulation of glucose

homeostasis and GLUT4 translocation in skeletal muscle, the relative role of PAK1 and PAK2 in

insulin-stimulated glucose uptake remains to be identified in mature skeletal muscle. Therefore, we

performed a systematic series of pharmacologic and genetic experiments to analyze the involvement

of group I PAKs in the regulation of insulin-stimulated glucose uptake in mouse skeletal muscle.

We hypothesized that group I PAKs, and in particular PAK1, would be necessary for glucose

uptake in response to insulin. Contradicting our hypothesis, our results revealed that insulin-

stimulated glucose uptake partly relies on PAK2 in glycolytic mouse muscle, while PAK1 is

dispensable for whole-body glucose homeostasis and insulin-stimulated muscle glucose uptake.

2. Methods

2.1. Animals

All animal experiments complied with the European Convention for the protection of vertebrate animals used for experimental and other scientific purposes (No. 123, Strasbourg, France, 1985; EU Directive 2010/63/EU for animal experiments) and were approved by the Danish Animal Experimental Inspectorate. All mice were maintained on a 12:12-hour light-dark cycle and housed at 22°C (with allowed fluctuation of ±2°C) with nesting material. The mice were group-housed. Female C57BL/6J mice (Taconic, Denmark) were used for the inhibitor incubation study. The mice received a standard rodent chow diet (Altromin no. 1324; Brogaarden, Denmark) and water ad libitum.

2.1.1. Whole-body PAK1$^{-/-}$ mice

For a complete overview of the different cohorts of genetically modified mice, see supplementary Table S1. Whole-body PAK1$^{-/-}$ mice on a C57BL/6J background were generated as previously described [30]. The mice were obtained by heterozygous crossing. PAK1$^{-/-}$ mice (referred to as PAK1 KO) and paired littermate PAK1$^{+/+}$ mice (referred to as controls) were used for experiments. Female and male mice were used for measurements of body composition, glucose tolerance and insulin-stimulated glucose uptake in isolated muscle. The mice were 12-24 weeks of age at the time of tissue dissection and measurement of glucose uptake. Number of mice in each group: Control, *n = 6/7* (female/male); PAK1 KO, *n = 4/8*. Mice received standard rodent chow diet and water ad libitum. For measurement of *in vivo* insulin-stimulated glucose uptake in chow- and 60E% high-fat diet (HFD; no. D12492; Brogaarden, Denmark)-fed PAK1 KO mice, mice were assigned to a chow or HFD group. Chow-fed mice were 10-24 weeks of age at the time of glucose uptake measurements and tissue dissection. Number of mice in each group: Control-Chow, *n = 14/8* (female/male); PAK1 KO-Chow, *n = 6/4*. For mice receiving HFD, the diet intervention started at

6-16 weeks of age and lasted for 21 weeks. HFD-fed mice were used for body composition, glucose

tolerance and *in vivo* glucose uptake and were 27-37 weeks of age at the time of glucose uptake

measurements and tissue dissection. Number of mice in each group: Control-HFD, *n = 7/7*

(female/male); PAK1 KO-HFD, *n = 11/5*. Energy intake was measured over a period of 10 weeks

in another cohort of mice. Number of mice in each group: Chow, *n = 8/8* (Control/PAK1 KO);

HFD, *n = 8/8*. Mice had access to their respective diet and water ad libitum.

2.1.2. Double PAK1$^{-/-}$;PAK2$^{fl/fl}$;MyoD$^{iCre/+}$ mice

Double knockout mice with whole-body knockout of PAK1 and conditional, muscle-specific

knockout of PAK2, PAK1$^{-/-}$;PAK2$^{fl/fl}$;MyoD$^{iCre/+}$ were generated as previously described [13]. The

mice were on a mixed C57BL/6/FVB background. PAK1$^{-/-}$;PAK2$^{fl/fl}$;MyoD$^{iCre/+}$ were crossed with

PAK1$^{+/-}$;PAK2$^{fl/fl}$;MyoD$^{+/+}$ to generate littermate PAK1$^{-/-}$;PAK2$^{fl/fl}$;MyoD$^{iCre/+}$ (referred to as 1/m2

dKO), PAK1$^{-/-}$;PAK2$^{fl/fl}$;MyoD$^{+/+}$ (referred to as PAK1 KO), PAK1$^{+/-}$;PAK2$^{fl/fl}$;MyoD$^{iCre/+}$ (referred

to as PAK2 mKO), and PAK1$^{+/-}$;PAK2$^{fl/fl}$;MyoD$^{+/+}$ (referred to as controls) used for experiments.

Female and male mice were used for measurement of insulin-stimulated glucose uptake in isolated

muscle. The mice were 10-16 weeks of age at the time of tissue dissection and glucose uptake

measurements. Number of mice in each group: Control, *n = 6/4* (female/male); PAK1 KO, *n = 5/4*,

PAK2 mKO, *n = 6/4*, 1/m2 dKO, *n = 6/3*. Another cohort of mice was used for whole-body

metabolic measurements. The first measurement (insulin tolerance) was at 11-24 weeks of age and

the last measurement was at 23-33 weeks of age (home cage calorimetry). Number of mice in each

group: Control, *n = 9/11* (female/male); PAK1 KO, *n = 8/10*, PAK2 mKO, *n = 12/9*, 1/m2 dKO, *n*

= 9/14. For some of the metabolic measurements, only a subgroup of mice was used as indicated in

the relevant figure legends. Mice received standard rodent chow diet and water ad libitum.

2.2.Body composition

Body composition was analyzed using magnetic resonance imaging (EchoMRI-4in1TM, Echo

Medical System LLC, Texas, USA). Chow-fed PAK1 KO and control littermates were assessed at

7-19 weeks of age. HFD-fed PAK1 KO and control littermates were assessed 18-19 weeks into the

diet intervention (24-34 weeks of age). Chow-fed PAK1 KO, PAK2 mKO, 1/m2 dKO mice and

control littermates were assessed at 16-29 weeks of age.

2.3.Glucose tolerance test (GTT)

Glucose tolerance was assessed in 8-20 weeks of age chow-fed PAK1 KO mice and in week 14 of

the diet intervention of HFD-fed PAK1 KO mice (20-30 weeks of age). In chow-fed PAK1 KO,

PAK2 mKO, 1/m2 dKO mice and control littermates, glucose tolerance was assessed at 13-26

weeks of age. Prior to the test, chow- and HFD-fed PAK1 KO mice and control littermates fasted

for 12 hours from 10 p.m, while chow-fed PAK1 KO, PAK2 mKO, 1/m2 dKO mice and control

littermates fasted for 6 hours from 6 a.m. D-mono-glucose (2 g kg^{-1} body weight) was administered

intraperitoneal (i.p) and blood was collected from the tail vein and blood glucose concentration

determined at the indicated time points using a glucometer (Bayer Contour, Bayer, Switzerland).

Incremental Area Under the Curve (AUC) from the basal blood glucose concentration was

determined using the trapezoid rule. For measurement of plasma insulin, glucose was administered

i.p. on a separate experimental day (1-2 weeks after the GTT) and blood was sampled at time points

0 and 20 minutes, centrifuged and plasma was quickly frozen in liquid nitrogen and stored at -20°C

until processing. Plasma insulin was analyzed in duplicate (Mouse Ultrasensitive Insulin ELISA,

#80-INSTRU-E10, ALPCO Diagnostics, USA). Homeostatic model assessment of insulin

resistance (HOMA-IR) was calculated according to the formula: Fasting plasma insulin (mU L^{-1}) X

Fasting blood glucose (mM)/22.5.

2.4.Insulin tolerance test (ITT)

Insulin tolerance was assessed in 11-24 weeks of age chow-fed PAK1 KO, PAK2 mKO, 1/m2 dKO

mice and control littermates. Prior to the test, chow-fed PAK1 KO, PAK2 mKO, 1/m2 dKO mice

and control littermates fasted for 4 hours from 6 a.m. Insulin (0.5 U kg^{-1} body weight) was

administered i.p. and blood was collected from the tail vein and blood glucose concentration

determined using a glucometer (Bayer Contour, Bayer, Switzerland) at time point 0, 15, 30, 60, 90

and 120 minutes. For two female control mice and four female PAK2 mKO mice, the ITT had to be

stopped before the 120'-time point due to hypoglycemia (blood glucose <1.2 mM). Thus, blood

glucose was not measured in these mice for the last couple of time points.

2.5. Home cage indirect calorimetry

One week prior to the calorimetric measurements, chow-fed PAK1 KO, PAK2 mKO, 1/m2 dKO

mice and control littermates were single-housed in specialized cages for indirect gas calorimetry but

uncoupled from the PhenoMaster indirect calorimetry system (TSE PhenoMaster metabolic cage

systems, TSE Systems, Germany). After a 2-day acclimation period coupled to the PhenoMaster

indirect calorimetry system, oxygen consumption, CO_2 production, habitual activity (beam breaks)

and food intake were measured for 72 hours (TSE LabMaster V5.5.3, TSE Systems, Germany). On

day 2, mice fasted during the dark period followed by refeeding on day 3. Respiratory exchange

ratio (RER) was calculated as the ratio between CO_2 production and oxygen consumption. The mice

were 23-33 weeks of age.

2.6. Incubation of isolated muscles

Soleus and extensor digitorum longus (EDL) muscles were dissected from anaesthetized mice (6

mg pentobarbital sodium 100 g^{-1} body weight i.p.) and suspended at resting tension (4-5 mN) in

incubations chambers (Multi Myograph System, Danish Myo Technology, Denmark) in Krebs-

Ringer-Henseleit buffer with 2 mM pyruvate and 8 mM mannitol at 30°C, as described previously

[31]. Additionally, the Krebs-Ringer-Henseleit buffer was supplemented with 0.1% BSA (v/v).

Isolated muscles from female C57BL/6J mice were pre-incubated with 40 μM IPA-3 (Sigma-Aldrich) or as a control DMSO (0.25%) for 25 minutes followed by 30 minutes of insulin stimulation (60 nM; Actrapid, Novo Nordisk, Denmark). Isolated muscles from chow-PAK1 KO were pre-incubated for 30 minutes followed by 30 minutes of insulin stimulation (0.6 nm or 60 nM). Isolated muscles from chow-fed PAK1 KO, PAK2 mKO, 1/m2 dKO mice or control littermates were pre-incubated for 20 minutes followed by 20 minutes of insulin stimulation (60 nM). 2-Deoxyglucose (2DG) uptake was measured together with 1 mM 2DG during the last 10 min of the insulin stimulation period using 0.60-0.75 μCi mL^{-1} [3H]-2DG and 0.180-0.225 μCi mL^{-1} [14C]-mannitol radioactive tracers as described previously [31]. Tissue-specific [3H]-2DG accumulation with [14C]-mannitol as an extracellular marker was determined as previously described [32].

2.7.In vivo insulin-stimulated 2-Deoxyglucose uptake in PAK1 KO mice during a r.o. ITT

To determine 2DG uptake in skeletal muscle of PAK1 KO mice and littermate controls, [3H]-2DG (Perkin Elmer) was administered retro-orbitally (r.o.) in a bolus of saline containing 66.7 μCi mL^{-1} [3H]-2DG (∼32.4 Ci/mmol) corresponding to ∼10 μCi/mouse in chow-fed mice or ∼15 μCi/mouse in HFD-fed mice (6 μL g^{-1} body weight) as described [20]. The [3H]-2DG saline bolus was with or without insulin (Actrapid, Novo Nordisk, Denmark). Decreased insulin clearance has been observed by us [20] and others in obese rodent [33,34] and human [35] models. Therefore, to correct for changes in insulin clearance, 0.5 U kg^{-1} body weight of insulin was administered in chow-fed mice whereas only 60% of this dosage was administered to HFD-fed mice. Prior to stimulation, mice fasted for 4 hours from 07:00 and were anaesthetized (7.5/9 mg [Chow/HFD] pentobarbital sodium 100 g^{-1} body weight i.p.) 15 minutes before the r.o. injection. Blood samples were collected from the tail vein after the r.o. injection and analyzed for glucose concentration using a glucometer (Bayer Contour, Bayer, Switzerland) at the time points 0, 5 and 10 minutes. After 10 minutes,

skeletal muscle (gastrocnemius, quadriceps, triceps brachii and soleus) were excised, rinsed in

saline, and quickly frozen in liquid nitrogen and stored at -80°C until processing. Blood was

collected by punctuation of the heart, centrifuged and plasma was quickly frozen in liquid nitrogen

and stored at -80°C until processing. Plasma samples were analyzed for insulin concentration and

specific [3H]-2DG activity. Plasma insulin was analyzed in duplicate (Mouse Ultrasensitive Insulin

ELISA, #80-INSTRU-E10, ALPCO Diagnostics, USA). Tissue-specific 2DG-6-phosphate

accumulation was measured as described [36,37]. To determine 2DG clearance from the plasma

into the specific tissue, tissue-specific [3H]-2DG-6-P was divided by AUC of the plasma-specific

[3H]-2DG activity at the time points 0 and 10 minutes. To estimate tissue-specific glucose uptake

(glucose uptake index), clearance was multiplied by the average blood glucose levels for the time

points 0, 5, and 10 minutes. Tissue-specific 2DG clearance and glucose uptake were related to

analyzed muscle tissue weight and time.

2.8.Muscle molecular analysis

Prior to homogenization, gastrocnemius, quadriceps, and triceps brachii muscles were pulverized in

liquid nitrogen. All muscle were homogenized 2 x 30 sec at 30 Hz using a Tissuelyser II (Qiagen,

USA) in ice-cold homogenization buffer (10% (v/v) Glycerol, 1% (v/v) NP-40, 20 mM Na-

pyrophosphate, 150 mM NaCl, 50 mM HEPES (pH 7.5), 20 mM β-glycerophosphate, 10 mM NaF,

2mM PMSF, 1 mM EDTA (pH 8.0), 1 mM EGTA (pH 8.0), 2 mM Na3VO4, 10 μg mL^{-1}

Leupeptin, 10 μg mL^{-1} Aprotinin, 3 mM Benzamidine). After rotation end-over-end for 30 min at

4°C, lysate supernatants were collected by centrifugation (10,854-15,630 x g) for 15-20 min at 4°C.

2.8.1. Immunoblotting

Lysate protein concentration was determined using the bicinchoninic acid method using bovine

serum albumin (BSA) standards and bicinchoninic acid assay reagents (Pierce). Immunoblotting

samples were prepared in 6x sample buffer (340 mM Tris (pH 6.8), 225 mM DTT, 11% (w/v) SDS,

20% (v/v) Glycerol, 0.05% (w/v) Bromphenol blue). Protein phosphorylation (p) and total protein

expression were determined by standard immunoblotting technique loading equal amounts of

protein. The polyvinylidene difluoride membrane (Immobilon Transfer Membrane; Millipore) was

blocked in Tris-Buffered Saline with added Tween20 (TBST) and 2% (w/v) skim milk or 5% (w/v)

BSA protein for 15 minutes at room temperature, followed by incubation overnight at 4°C with a

primary antibody (Table 1). Next, the membrane was incubated with horseradish peroxidase-

conjugated secondary antibody (Jackson Immuno Research) for 1 hour at room temperature. Bands

were visualized using Bio-Rad ChemiDocTM MP Imaging System and enhanced

chemiluminescence (ECL+; Amersham Biosciences). Densitometric analysis was performed using

Image LabTM Software, version 4.0 (Bio-Rad, USA; RRID:SCR_014210). Coomassie brilliant

blue staining was used as a loading control [38].

2.9.Statistical analyses

Data are presented as mean ± S.E.M. or when applicable mean ± S.E.M. with individual data points

shown. Statistical tests varied according to the dataset being analyzed and the specific tests used are

indicated in the figure legends. Datasets were normalized by square root, log10, inverse or inverse

square root transformation if not normally distributed or unequal equal variance. If the null

hypothesis was rejected, Tukey's post hoc test was used to evaluate significant main effects of

genotype and significant interactions in ANOVAs. $P < 0.05$ was considered statistically significant.

$P<0.1$ was considered a tendency. Except for mixed-effects model analyses performed in GraphPad

Prism, version 8.2.1. (GraphPad Software, La Jolla, CA, USA; RRID:SCR_002798), all statistical

analyses were performed using Sigma Plot, version 13 (Systat Software Inc., Chicago, IL, USA;

RRID:SCR_003210). Due to missing values ascribed to hypoglycemia, differences between

genotypes and the effect of insulin administration were assessed with a mixed-effects model

analysis in Fig. 5F and S5F.

3. Results

3.1.Pharmacological inhibition of group I PAKs partially reduces insulin-stimulated glucose uptake in mouse soleus muscle

To investigate the involvement of group I PAKs in insulin-stimulated glucose uptake in mouse skeletal muscle, we first analyzed 2DG uptake in isolated soleus and EDL muscle in the presence of a pharmacological inhibitor, IPA-3. While glucose uptake *in vivo* is influenced by glucose delivery, GLUT4 translocation and muscle metabolism [39], glucose delivery is constant in isolated skeletal muscle and surface membrane GLUT4 is the limiting factor [40–42]. Therefore, 2DG uptake in isolated muscles likely reflects GLUT4 translocation. IPA-3 is a well-characterized inhibitor of group I PAKs (PAK1-3) [6,43] and shown to completely abolish insulin-stimulated GLUT4 translocation and glucose uptake in L6-GLUT4myc myoblasts and myotubes, respectively [6]. In soleus, 2DG uptake increased 4.5-fold upon maximal insulin-stimulation, an increase that was partly reduced (-20%) in IPA-3-treated muscles (Fig. 1A). IPA-3 did not significantly (p=0.080) impair insulin-stimulated (+2.4-fold) 2DG uptake in EDL muscle (Fig. 1B). We confirmed that IPA-3 treatment abolished insulin-stimulated phosphorylation of (p)PAK1 T423 in both soleus and EDL muscle (Fig. 1C+D). In contrast, insulin-stimulated pAkt T308 (Fig. 1E+F) and pAkt S473 (Fig. 1G+H) were unaffected by IPA-3 treatment, suggesting that IPA-3 did not interfere with insulin signalling to Akt. Altogether this suggests that group I PAKs are only partially involved in insulin-stimulated glucose uptake in isolated mouse muscle.

3.2.PAK1 knockout does not affect whole-body glucose tolerance or insulin-stimulated glucose uptake in isolated skeletal muscle

We next sought to confirm our findings in mice with whole-body lack of the PAK1 isoform (PAK1 KO) and therefore a complete knockout of PAK1 in skeletal muscle (Fig. 2A). In chow-fed mice,

fat mass, lean body mass, body weight and energy intake (Fig. 2B-C) were similar between whole-body PAK1 KO and littermate controls, as also reported previously [44]. During a GTT, the lack of PAK1 had no effect on the blood glucose response (Fig. 2D-E) or plasma insulin concentration (Fig. 2F). Additionally, HOMA-IR, a measure of basal glucose homeostasis (Fig. 2G), and both submaximal and maximal insulin-stimulated 2DG uptake in isolated soleus and EDL muscle were unaffected by PAK1 KO (Fig. 2H-I). Thus, unexpectedly, genetic ablation of PAK1 alone did not impair whole-body glucose tolerance, or skeletal muscle insulin sensitivity (submaximal insulin-stimulated glucose uptake) or insulin responsiveness (maximal insulin-stimulated glucose uptake) in divergence to previous reports [6,7,11].

3.3.PAK1 is dispensable for insulin-stimulated glucose uptake in lean or diet-induced insulin-resistant mice in vivo

Our findings on insulin-stimulated glucose uptake in isolated muscle from chow-fed PAK1 KO mice conflicted with a previous study reporting impaired glucose tolerance in PAK1 KO mice and defects in insulin-stimulated GLUT4 translocation in skeletal muscle *in vivo* [7]. Therefore, we further explored the effect of PAK1 KO on insulin-stimulated glucose uptake in skeletal muscle *in vivo*. Additionally, we fed a subgroup of PAK1 KO and control littermate mice a 60E% HFD for 21 weeks to investigate the role of PAK1 in insulin-resistant muscle. Insulin administration lowered blood glucose by 5.4 ± 0.5 mM (-47%) in chow-fed control mice (Fig. 3A). In HFD-fed control mice, blood glucose dropped 3.0 ± 0.9 mM (-26%) upon insulin administration (Fig. 3B). Lack of PAK1 had no impact on either basal blood glucose or whole-body insulin tolerance on either of the diets (Fig. 3A-B). Insulin increased glucose uptake in muscles from chow-fed (Gastrocnemius: 8.1-fold; Quadriceps: 8.5-fold, Triceps brachii: 12.3-fold; Soleus: 8.9-fold) and HFD-fed (Gastrocnemius: 3.5-fold; Quadriceps: 1.8-fold, Triceps brachii: 4.3-fold; Soleus: 11.6-fold) control mice (Fig. 3C-D). Consistent with our findings in isolated muscle, we observed no effect of PAK1

KO on basal or insulin-stimulated glucose uptake *in vivo* in muscle of either chow-fed mice or in

insulin-resistant muscles from HFD-fed mice. Like glucose uptake, 2DG clearance from the plasma

was unaffected by PAK1 KO in all of the investigated muscles (Fig. S1A-B). Importantly,

circulating [^{3}H]-2DG availability was unaffected by genotype on both of the diets (Fig. S1C). As in

chow-fed mice, lack of PAK1 in HFD-fed mice had no effect on fat mass, lean body mass, body

weight or energy intake (Fig. S1D-E). Similarly, whole-body glucose tolerance (Fig. S1F-G),

plasma insulin concentration during the GTT (Fig. S1H) and HOMA-IR (Fig. S1I) were unaffected

by PAK1 KO in HFD-fed mice. Thus, PAK1 is dispensable for *in vivo* insulin-stimulated muscle

glucose uptake in both the healthy lean and the diet-induced insulin-resistant state.

3.4. Whole-body substrate utilization is unaffected by genetic ablation of PAK1 and PAK2

Because of discrepancies in the data resulting from the use of a global pharmacological inhibitor of

group I PAKs and data resulting from PAK1 KO mice, we next sought to determine the relative

contribution and involvement of PAK1 and PAK2 in insulin signalling and glucose uptake in

skeletal muscle. Double knockout mice with whole-body knockout of PAK1 and muscle-specific

knockout of PAK2 (1/m2 dKO) were generated as previously described [13]. By crossing 1/m2

dKO mice with littermate controls, a cohort was generated consisting of whole-body PAK1 KO,

muscle-specific PAK2 (m)KO, 1/m2 dKO and littermate control mice. While no band for PAK1

could be detected in muscles lacking PAK1, muscles lacking PAK2 displayed only a partial

reduction in band intensity in the immunoblots for PAK2 (Fig. 4A). This is likely due to the fact

that PAK1 KO mice are whole-body knockouts, while PAK2 mKO mice are muscle-specific and

other cell types within skeletal muscle tissue could thus contribute to the signal obtained in the

PAK2 immunoblots. In a previous study, PAK3 was not detectable at the protein level in 1/m2 dKO

muscle [13].

As previously shown [13,45], 1/m2 dKO mice weighed less (-12%) than control littermates (Fig. 4B, Fig. S2A-B) due to reduced (-12%) lean body mass (Fig. 4C, Fig. S2C-D). Body weight and lean body mass decreased to the same extent in 1/m2 dKO mice, leaving lean body mass percentage largely unaffected (Fig. S2E-G). Using calorimetric chambers, we monitored whole-body metabolism for 72 hours during the light and dark period. On day 2, the mice fasted during the dark period followed by refeeding on day 3. Oxygen uptake (VO_2; Fig. 4D, Fig. S3A-B) and RER indicative of substrate utilization (Fig. 4E, Fig. S3C-D) were largely unaffected by genotype with only a slightly higher RER in mice lacking PAK2 (PAK2 mKO and 1/m2 dKO mice) upon fasting. Similar substrate utilization was obtained despite reduced habitual activity in the dark period in mice lacking PAK2, an effect largely driven by a decreased activity in 1/m2 dKO mice and increased activity in male PAK1 KO mice (Fig. 4F; Fig. S3E-F). Supporting the lower activity levels, energy intake was decreased (-11%) in 1/m2 dKO mice compared to PAK1 KO mice on day 1 (Fig. S3G) due to lower energy intake in the dark period (Fig. 4G). Upon refeeding, energy intake was reduced in mice lacking PAK2 (Fig. 4G; Fig. S3G) driven by a lower energy intake in the dark period in female mice lacking PAK2 (Fig. S3H-K). Altogether, these data suggest that lack of PAK1 and/or PAK2 are not compromising metabolic regulation during the light/dark period or in response to fasting/refeeding.

3.5. Glucose tolerance is reduced in mice lacking PAK2 in skeletal muscle

To test dependency on PAK1 and/or PAK2 in glucose handling and insulin sensitivity, we next investigated glucose and insulin tolerance in the 1/m2 dKO mouse strain. Blood glucose concentration in the fed state was similar between the genotypes (Fig. 5A, Fig. S4A-B). The blood glucose response to a glucose load was slightly increased in mice lacking PAK2 in skeletal muscle as evident by the increased area under the blood glucose curve (Fig. 5B-C). This was primarily driven by impaired glucose tolerance in female PAK2 mKO mice (Fig. S4C-F). Plasma insulin

concentration during the GTT was unaffected by lack of PAK1 (Fig. 5D) In contrast, plasma insulin

in male PAK2 mKO mice was slightly elevated compared to 1/m2 dKO mice and tended (p=0.055)

to be higher than control littermates (Fig. S4G-H). This indicates impaired insulin sensitivity in

male PAK2 mKO mice, but HOMA-IR was unaffected by lack of PAK1 and/or PAK2 (Fig. 5E,

Fig. S5A-B). In addition, even though fasted blood glucose immediately before an ITT was

modestly reduced in PAK2 mKO mice (Fig. S5C-E), the blood glucose response to an ITT was

largely unaffected by lack of either PAK1 and/or PAK2 (Fig. 5F, Fig. S5F-G). Thus, despite

slightly impaired glucose tolerance in mice lacking PAK2 in skeletal muscle, neither adverse effects

on plasma insulin nor defects in insulin sensitivity could be detected.

3.6.Insulin-stimulated glucose uptake relies partially on PAK2 in EDL, but not soleus muscle,

while PAK1 is not involved

To determine the relative contribution and involvement of PAK1 and PAK2 in glucose uptake in

skeletal muscle, we next investigated insulin-stimulated 2DG uptake in isolated soleus and EDL. To

our surprise, soleus muscle lacking PAK1 and PAK2 displayed normal insulin-stimulated 2DG

uptake compared to control littermates (Fig. 6A,C). In contrast, lack of PAK2 in EDL muscle

caused a modest reduction (PAK2 mKO: -18%; 1/m2 dKO: -12%) in insulin-stimulated 2DG

uptake (Fig. 6B,D). Thus, in oxidative soleus muscle, group I PAKs are surprisingly dispensable for

normal insulin-stimulated glucose uptake, whereas in glycolytic EDL muscle PAK2 plays a minor

role.

3.7.PAK2 regulates TBC1D4 protein expression and signalling

Lastly, we looked into the effects of PAK1 and PAK2 on insulin-stimulated signalling. All groups

displayed normal basal and insulin-stimulated pAkt S473 (Fig. 7A-B, Fig. S6A-B) and pAkt T308

(Fig. 7C-D, Fig. S6C-D) and Akt2 protein expression (Fig. S6E-F) compared to control littermates

in both soleus and EDL muscle. In contrast, lack of PAK2 increased protein expression of Akt's

downstream target, TBC1D4 in soleus muscle (PAK2 mKO: +47%; 1/m2 dKO: +20%) (Fig. 7E),

while reducing TBC1D4 expression in EDL (PAK2 mKO: -33%; 1/m2 dKO: -9%) (Fig. 7F). In

soleus, basal and insulin-stimulated pTBC1D4 T642 was similar in all groups (Fig. 7G, Fig. S6G),

suggesting that even with increased TBC1D4 expression, signalling through this pathway was

normal. Concomitant with the decreased TBC1D4 expression in EDL muscle, lack of PAK2

reduced insulin-stimulated pTBC1D4 T642 (Fig. 7H, Fig. S6H) driven by attenuated (-46%)

insulin-stimulated pTBC1D4 T642 in PAK2 mKO mice compared to control littermates (Fig. S6H).

Knockout of TBC1D4 has been associated with lower GLUT4 protein abundance in some muscles

[46,47]. Whereas GLUT4 protein expression was normal in soleus (Fig. 7I), GLUT4 protein

expression was mildly reduced in EDL in PAK2 mKO mice compared to littermate control (Fig.

7J). Thus, the slightly reduced insulin-stimulated glucose uptake in EDL muscle lacking PAK2 was

concomitant with downregulated TBC1D4 signalling and GLUT4 expression supporting a role for

PAK2, but not PAK1, in insulin-stimulated glucose uptake.

4. Discussion

We here undertook a systematic investigation into the requirement of the group I PAK isoforms in

muscle glucose uptake and whole-body metabolic regulation. In contrast to previous literature, our

results firmly show that PAK1 surprisingly is dispensable for insulin-stimulated glucose uptake in

skeletal muscle, while PAK2 may play a minor role. Using a cohort of whole-body PAK1 KO mice

and another cohort of transgenic mice lacking either PAK1 (whole-body), PAK2 (muscle-specific),

or jointly lacking both PAK1 and muscle PAK2, we show that PAK1 is not required in insulin-

stimulated muscle glucose uptake *in vivo* or in isolated muscles. In accordance, we found no effect

of whole-body PAK1 KO on glucose tolerance in either mice fed a standard chow diet (insulin

sensitive mice) or in mice fed a HFD (insulin-resistant mice). In contrast, PAK2 seemed partially

involved in insulin-stimulated glucose uptake, in glycolytic EDL muscle. This could potentially

explain the slightly impaired glucose tolerance with the muscle-specific knockout of PAK2 in mice.

Nevertheless, supporting only a minor role for skeletal muscle PAK2 in the whole-body substrate

utilization, RER was largely unaffected by lack of PAK1 and/or PAK2 and only slightly elevated in

mice lacking PAK2 when challenged by fasting.

In a previous study, the increase in GLUT4 abundance at the plasma membrane in response to

insulin measured was completely abrogated in PAK1 KO muscle as measured by subcellular

fractionation of homogenates of hindlimb skeletal muscles [7], suggesting that PAK1 is necessary

for insulin-stimulated GLUT4 translocation. Although glucose uptake was not assessed in that study

[7], this indicated a key role for PAK1 in regulating glucose uptake in mouse skeletal muscle.

Surprisingly, PAK1 was not required for insulin-stimulated glucose uptake in our hands.

Furthermore, in our study, both chow- and HFD-fed PAK1 KO mice displayed blood glucose

concentrations similar to control littermates during a GTT. This was in contrast to previous studies

reporting impaired glucose tolerance in chow-fed PAK1 KO mice [7,27] and elevated fasting blood

glucose in PAK1 KO mice fed a western diet (45E% fat) [44]. Instead, despite the previous finding

that insulin-stimulated GLUT4 translocation was unaffected by a 75% knockdown of PAK2 in L6-

GLUT4myc myoblasts [6], we found that muscle-specific PAK2 KO slightly impaired glucose

tolerance and insulin-stimulated glucose in mouse skeletal muscle. These discrepancies between our

and previous findings are difficult to delineate but might be due to methodological differences.

Wang et al. [7] used a crude fractionation method to measure GLUT4 translocation, whereas we

analyzed the direct outcome hereof: glucose uptake. Although the insulin-induced increase in 3-O-

methylglucose transport correlates with the increase in cell surface GLUT4 protein content in

human skeletal muscle strips [48], discrepancies between the presence of GLUT4 at the plasma

membrane and glucose uptake have occasionally been reported in cell culture studies [49–51].

Another potential explanatory factor could be that our studies were conducted in both female and

male mice, whereas past studies in PAK1 KO mice have been conducted in 4-6 months old male

mice [7,44]. However, our data suggest no major differences between female and male mice in the

response to lack of PAK1 and/or PAK2 on the whole-body metabolic parameters measured. Instead,

the discrepancies between our and previous findings could be due to an effect of age, as our studies

were conducted in mice at different ranges of age (10-37 of weeks age at the terminal experiment).

In fact, age-dependent myopathy and development of megaconial mitochondria have been reported

in the 1/m2 dKO mice [45]. Regardless, even though a role for group I PAKs in age-related insulin

resistance should be further investigated, our investigation suggests that group I PAKs are largely

dispensable in regulating whole-body glucose homeostasis or insulin-stimulated glucose uptake in

skeletal muscle.

In the current study, pharmacological inhibition of group I PAKs inhibited muscle glucose uptake in

response to insulin. Similarly, IPA-3 previously inhibited insulin-stimulated GLUT4 translocation

and glucose uptake in L6-GLUT4myc myoblasts and myotubes, respectively [6]. IPA-3 is a non-

ATP-competitive allosteric inhibitor of all group I PAKs (PAK1, 2, and 3). IPA-3 binds covalently

to the regulatory CRIB domain of group I PAKs, thereby preventing binding to PAK activators,

such as Rac1 [43]. Although IPA-3 is reported to be a highly selective and well-described inhibitor

of group I PAKs that does not affect other groups of PAKs or similar kinases tested [43],

pharmacological inhibitors often have off-target effects [52] which is a concern. It is also possible

that acute IPA-3-induced inhibition of group I PAKs elicits more potent effects compared with

jointly knockout of PAK1 and PAK2 because the transgenic manipulations have been present from

birth and may thus have resulted in compensatory changes. The development of inducible muscle-

specific group I PAK deficient models could help clarify this. Importantly, any possible

compensatory mechanisms cannot be via redundancy with group I PAKs, as PAK1 and PAK2 are

removed genetically, and even in 1/m2 dKO mice, PAK3 cannot be detected at the protein level

[13]. This emphasizes that group I PAKs are largely dispensable for insulin-stimulated glucose

uptake in skeletal muscle with only PAK2 playing a minor role.

Our hypothesis was that group I PAKs would be significantly involved in insulin-stimulated

glucose uptake because of the established necessity of their upstream activator, Rac1 [15–20]. Our

findings suggest that Rac1 does not exclusively mediate insulin-stimulated glucose uptake through

group I PAKs. Another downstream target of Rac1 is RalA. GLUT4 translocation induced by a

constitutively activated Rac1 mutant was abrogated in L6-GLUT4myc myoblasts upon RalA

knockdown [53] and, importantly, overexpression of a dominant-negative mutant of RalA reduced

GLUT4 translocation in response to insulin in mouse gastrocnemius muscle fibres [54].

Additionally, Rac1 is an essential component in the activation of the NADPH oxidase (NOX)

complex [55,56]. In L6-GLUT4myc myotubes, reactive oxygen species have been reported to

induce NOX2-dependent GLUT4 translocation in response to insulin [57]. Insulin-stimulated

NOX2 regulation in mature muscle remains to be investigated. However, a recent study suggested a

role for Rac1 in the regulation of muscle glucose uptake through activation of the NOX2 in

response to exercise [58]. Since Rac1 is required for both contraction- and insulin-stimulated

glucose uptake in isolated mouse muscle [18,59], Rac1 could also be involved in insulin-stimulated

glucose uptake via NOX2 activation. Consequently, future studies should aim to investigate other

players downstream of Rac1 since group I PAKs seem to be largely dispensable for glucose uptake

in mature skeletal muscle.

5. Conclusion

Based on our present findings, we conclude that PAK2 may be a requirement for insulin-stimulated

glucose uptake in EDL muscle. However, in contrast to previous reports, group I PAKs are largely

dispensable in the regulation of whole-body glucose homeostasis and insulin-stimulated glucose

uptake in mouse skeletal muscle.

Author contributions

LLVM: Conceptualization; Methodology; Formal analysis; Investigation; Writing - Original Draft;

Writing - Review & Editing; Visualization; Project administration; Funding acquisition. **MJ:**

Investigation; Writing - Review & Editing. **RK:** Methodology; Investigation; Writing - Review &

Editing. **GAJ:** Resources; Writing - Review & Editing. **ABM:** Investigation; Writing - Review &

Editing; Funding acquisition. **JRK:** Investigation; Writing - Review & Editing; Funding

acquisition. **A-ML:** Investigation; Writing - Review & Editing; Funding acquisition. **NRA:**

Investigation; Writing - Review & Editing. **PS:** Methodology; Investigation; Writing - Review &

Editing. **TEJ:** Investigation; Writing - Review & Editing; Funding acquisition. **RSK:** Resources;

Writing - Review & Editing; Funding acquisition. **EAR:** Conceptualization; Methodology; Writing

- Original Draft; Writing - Review & Editing; Supervision; Project administration; Funding

acquisition. **LS:** Conceptualization; Methodology; Investigation; Writing - Original Draft; Writing -

Review & Editing; Supervision; Project administration; Funding acquisition. LS is the guarantor of

this work and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Acknowledgements

504 We thank our colleagues, especially Jørgen Wojtaszewski and Bente Kiens, at the Section of

505 Molecular Physiology, Department of Nutrition, Exercise, and Sports (NEXS), Faculty of Science,

506 University of Copenhagen, for fruitful discussions on this topic. We acknowledge the skilled

technical assistance of Betina Bolmgren, Irene B. Nielsen, and Mona Ali (Section of Molecular Physiology, NEXS, Faculty of Science, University of Copenhagen, Denmark). The PAK1 KO founder mice were a kind gift from Debbie Thurmond (Department of Molecular & Cellular Endocrinology, Diabetes and Metabolism Research Institute, City of Hope/BRICA, USA). The graphical abstract was modified from Servier Medical Art, licensed under a Creative Common Attribution 3.0 Generic License (http://smart.servier.com/).

Grants

This study was supported by a PhD fellowship from The Lundbeck Foundation (grant 2015-3388 to LLVM); PhD scholarships from The Danish Diabetes Academy, funded by The Novo Nordisk Foundation (ABM and JRK); Postdoctoral research grant from the Danish Diabetes Academy, funded by the Novo Nordisk Foundation (grant NNF17SA0031406 to A-ML); National Institute of Arthritis and Musculoskeletal and Skin Diseases (grant AR046207 and AR070231 to RSK); The Danish Council for Independent Research, Medical Sciences (grant DFF-4004-00233 to LS, grant 6108-00203 to EAR); The Novo Nordisk Foundation (grant 10429 to EAR, grant 15182 to TEJ, grant NNF16OC0023418 and NNF18OC0032082 to LS).

Appendix A. Supplementary data

Supplementary table S1 and supplementary figures S1-6 are to be found in Appendix A.

Conflict of interest

None declared.

Data and Resource Availability

The datasets generated and analyzed during the current study are available from the corresponding

author upon reasonable request. No novel applicable resources were generated or analyzed during

the current study.

References

[1]   DeFronzo, R.A., Gunnarsson, R., Björkman, O., Olsson, M., Wahren, J., 1985. Effects of insulin on peripheral and splanchnic glucose metabolism in noninsulin-dependent (type II) diabetes mellitus. Journal of Clinical Investigation 76(1): 149–55, Doi: 10.1172/JCI111938.

[2]   Baron, A.D., Brechtel, G., Wallace, P., Edelman, S. V., 1988. Rates and tissue sites of non-insulin- and insulin-mediated glucose uptake in humans. American Journal of Physiology-Endocrinology and Metabolism 255(6): E769–74, Doi: 10.1152/ajpendo.1988.255.6.E769.

[3]   Kahn, S.E., 2003. The relative contributions of insulin resistance and beta-cell dysfunction to the pathophysiology of Type 2 diabetes. Diabetologia: 3–19, Doi: 10.1007/s00125-002-1009-0.

[4]   Lauritzen, H.P.M.M., 2013. Insulin- and Contraction-Induced Glucose Transporter 4 Traffic in Muscle. Exercise and Sport Sciences Reviews 41(2): 77–86, Doi: 10.1097/JES.0b013e318275574c.

[5]   Tsakiridis, T., Taha, C., Grinsteinl, S., Klip, A., 1996. Insulin activates a p21-activated kinase in muscle cells via phosphatidylinositol 3-kinase. Journal of Biological Chemistry 271(33): 19664–7, Doi: 10.1074/jbc.271.33.19664.

[6]   Tunduguru, R., Chiu, T.T., Ramalingam, L., Elmendorf, J.S., Klip, A., Thurmond, D.C., 2014. Signaling of the p21-activated kinase (PAK1) coordinates insulin-stimulated actin remodeling and glucose uptake in skeletal muscle cells. Biochemical Pharmacology 92(2): 380–8, Doi: 10.1016/j.bcp.2014.08.033.

[7]   Wang, Z., Oh, E., Clapp, D.W., Chernoff, J., Thurmond, D.C., 2011. Inhibition or ablation of p21-activated kinase (PAK1) disrupts glucose homeostatic mechanisms in vivo. Journal of Biological Chemistry 286(48): 41359–67, Doi: 10.1074/jbc.M111.291500.

[8]   Bagrodia, S., Cerione, R.A., 1999. PAK to the future. Trends in Cell Biology: 350–5, Doi: 10.1016/S0962-8924(99)01618-9.

[9]   Bokoch, G.M., 2003. Biology of the p21-Activated Kinases. Annual Review of Biochemistry 72(1): 743–81, Doi: 10.1146/annurev.biochem.72.121801.161742.

[10]   Hofmann, C., Shepelev, M., Chernoff, J., 2004. The genetics of Pak. Journal of Cell Science 117(19): 4343–54, Doi: 10.1242/jcs.01392.

[11]   Chiang, Y.A., Jin, T., 2014. p21-Activated protein kinases and their emerging roles in glucose homeostasis. American Journal of Physiology-Endocrinology and Metabolism 306(7): E707–22, Doi:

10.1152/ajpendo.00506.2013.

[12] Manser, E., Leung, T., Salihuddin, H., Zhao, Z., Lim, L., 1994. A brain serine/threonine protein
kinase activated by Cdc42 and Rac1. Nature 367(6458): 40–6, Doi: 10.1038/367040a0.

[13] Joseph, G.A., Lu, M., Radu, M., Lee, J.K., Burden, S.J., Chernoff, J., et al., 2017. Group I Paks
Promote Skeletal Myoblast Differentiation In Vivo and In Vitro. Molecular and Cellular Biology
37(4): e00222-16, Doi: 10.1128/MCB.00222-16.

[14] Arias-Romero, L.E., Chernoff, J., 2008. A tale of two Paks. Biology of the Cell 100(2): 97–108, Doi:
10.1042/BC20070109.

[15] Khayat, Z.A., Tong, P., Yaworsky, K., Bloch, R.J., Klip, A., 2000. Insulin-induced actin filament
remodeling colocalizes actin with phosphatidylinositol 3-kinase and GLUT4 in L6 myotubes. Journal
of Cell Science 113 Pt 2: 279–90.

[16] JeBailey, L., Wanono, O., Niu, W., Roessler, J., Rudich, A., Klip, A., 2007. Ceramide- and oxidant-
induced insulin resistance involve loss of insulin-dependent Rac-activation and actin remodeling in
muscle cells. Diabetes 56(2): 394–403, Doi: 10.2337/db06-0823.

[17] Ueda, S., Kitazawa, S., Ishida, K., Nishikawa, Y., Matsui, M., Matsumoto, H., et al., 2010. Crucial
role of the small GTPase Rac1 in insulin-stimulated translocation of glucose transporter 4 to the
mouse skeletal muscle sarcolemma. The FASEB Journal 24(7): 2254–61, Doi: 10.1096/fj.09-137380.

[18] Sylow, L., Jensen, T.E., Kleinert, M., Hojlund, K., Kiens, B., Wojtaszewski, J., et al., 2013. Rac1
Signaling Is Required for Insulin-Stimulated Glucose Uptake and Is Dysregulated in Insulin-Resistant
Murine and Human Skeletal Muscle. Diabetes 62(6): 1865–75, Doi: 10.2337/db12-1148.

[19] Sylow, L., Kleinert, M., Pehmøller, C., Prats, C., Chiu, T.T., Klip, A., et al., 2014. Akt and Rac1
signaling are jointly required for insulin-stimulated glucose uptake in skeletal muscle and
downregulated in insulin resistance. Cellular Signalling 26(2): 323–31, Doi:
10.1016/j.cellsig.2013.11.007.

[20] Raun, S.H., Ali, M., Kjøbsted, R., Møller, L.L. V., Federspiel, M.A., Richter, E.A., et al., 2018. Rac1
muscle knockout exacerbates the detrimental effect of high-fat diet on insulin-stimulated muscle
glucose uptake independently of Akt. The Journal of Physiology 596(12): 2283–99, Doi:
10.1113/JP275602.

[21] Sano, H., Kane, S., Sano, E., Mîinea, C.P., Asara, J.M., Lane, W.S., et al., 2003. Insulin-stimulated
Phosphorylation of a Rab GTPase-activating Protein Regulates GLUT4 Translocation. Journal of
Biological Chemistry 278(17): 14599–602, Doi: 10.1074/jbc.C300063200.

[22] Thong, F.S.L., Dugani, C.B., Klip, A., 2005. Turning signals on and off: GLUT4 traffic in the
insulin-signaling highway. Physiology: 271–84, Doi: 10.1152/physiol.00017.2005.

[23] Kramer, H.F., Witczak, C.A., Taylor, E.B., Fujii, N., Hirshman, M.F., Goodyear, L.J., 2006. AS160
Regulates Insulin- and Contraction-stimulated Glucose Uptake in Mouse Skeletal Muscle. Journal of
Biological Chemistry 281(42): 31478–85, Doi: 10.1074/jbc.M605461200.

[24] Benner, G.E., Dennis, P.B., Masaracchia, R.A., 1995. Activation of an S6/H4 Kinase (PAK 65) from
Human Placenta by Intramolecular and Intermolecular Autophosphorylation. Journal of Biological
Chemistry 270(36): 21121–8, Doi: 10.1074/jbc.270.36.21121.

[25] Gatti, A., Huang, Z., Tuazon, P.T., Traugh, J.A., 1999. Multisite autophosphorylation of p21-
activated protein kinase gamma-PAK as a function of activation. The Journal of Biological Chemistry
274(12): 8022–8, Doi: 10.1074/jbc.274.12.8022.

[26] Stierwalt, H.D., Ehrlicher, S.E., Bergman, B.C., Robinson, M.M., Newsom, S.A., 2018. Insulin-
stimulated Rac1-GTP binding is not impaired by palmitate treatment in L6 myotubes. Physiological
Reports 6(24): e13956, Doi: 10.14814/phy2.13956.

[27] Chiang, Y.A., Shao, W., Xu, X.X., Chernoff, J., Jin, T., 2013. P21-activated protein kinase 1 (Pak1)
mediates the cross talk between insulin and β-catenin on proglucagon gene expression and its ablation
affects glucose homeostasis in male C57BL/6 mice. Endocrinology 154(1): 77–88, Doi:
10.1210/en.2012-1781.

[28] Chiu, T.T., Patel, N., Shaw, A.E., Bamburg, J.R., Klip, A., 2010. Arp2/3- and Cofilin-coordinated
Actin Dynamics Is Required for Insulin-mediated GLUT4 Translocation to the Surface of Muscle
Cells. Molecular Biology of the Cell 21(20): 3529–39, Doi: 10.1091/mbc.e10-04-0316.

[29] Tunduguru, R., Zhang, J., Aslamy, A., Salunkhe, V.A., Brozinick, J.T., Elmendorf, J.S., et al., 2017.
The actin-related p41ARC subunit contributes to p21-activated kinase-1 (PAK1)-mediated glucose
uptake into skeletal muscle cells. Journal of Biological Chemistry 292(46): 19034–43, Doi:
10.1074/jbc.M117.801340.

[30] Allen, J.D., Jaffer, Z.M., Park, S.J., Burgin, S., Hofmann, C., Sells, M.A., et al., 2009. P21-activated
kinase regulates mast cell degranulation via effects on calcium mobilization and cytoskeletal
dynamics. Blood 113(12): 2695–705, Doi: 10.1182/blood-2008-06-160861.

[31] Viollet, B., Andreelli, F., Jørgensen, S.B., Perrin, C., Geloen, A., Flamez, D., et al., 2003. The AMP-
activated protein kinase α2 catalytic subunit controls whole-body insulin sensitivity. Journal of
Clinical Investigation 111(1): 91–8, Doi: 10.1172/JCI16567.

[32] Kjøbsted, R., Treebak, J.T., Fentz, J., Lantier, L., Viollet, B., Birk, J.B., et al., 2015. Prior AICAR
stimulation increases insulin sensitivity in mouse skeletal muscle in an AMPK-dependent manner.
Diabetes 64(6): 2042–55, Doi: 10.2337/db14-1402.

[33] Strömblad, G., Björntorp, P., 1986. Reduced hepatic insulin clearance in rats with dietary-induced
obesity. Metabolism 35(4): 323–7, Doi: 10.1016/0026-0495(86)90148-4.

[34] Brandimarti, P., Costa-Júnior, J.M., Ferreira, S.M., Protzek, A.O., Santos, G.J., Carneiro, E.M., et al.,
2013. Cafeteria diet inhibits insulin clearance by reduced insulindegrading enzyme expression and
mRNA splicing. Journal of Endocrinology 219(2): 173–82, Doi: 10.1530/JOE-13-0177.

[35] Jung, S.-H., Jung, C.-H., Reaven, G.M., Kim, S.H., 2018. Adapting to insulin resistance in obesity:
role of insulin secretion and clearance. Diabetologia 61(3): 681–7, Doi: 10.1007/s00125-017-4511-0.

[36] Ferre, P., Leturque, A., Burnol, A.-F., Penicaud, L., Girard, J., 1985. A method to quantify glucose
utilization in vivo in skeletal muscle and white adipose tissue of the anaesthetized rat. Biochemical
Journal 228(1): 103–10.

[37] Fueger, P.T., Bracy, D.P., Malabanan, C.M., Pencek, R.R., Wasserman, D.H., 2004. Distributed
control of glucose uptake by working muscles of conscious mice: roles of transport and
phosphorylation. American Journal of Physiology-Endocrinology and Metabolism 286(1): E77–84,
Doi: 10.1152/ajpendo.00309.2003.

[38] Welinder, C., Ekblad, L., 2011. Coomassie Staining as Loading Control in Western Blot Analysis.
Journal of Proteome Research 10(3): 1416–9, Doi: 10.1021/pr1011476.

[39] Wasserman, D.H., Kang, L., Ayala, J.E., Fueger, P.T., Lee-Young, R.S., 2011. The physiological
regulation of glucose flux into muscle in vivo. Journal of Experimental Biology 214(2): 254–62, Doi:
10.1242/jeb.048041.

[40] Hansen, P.A., Gulve, E.A., Marshall, B.A., Gao, J., Pessin, J.E., Holloszy, J.O., et al., 1995. Skeletal
muscle glucose transport and metabolism are enhanced in transgenic mice overexpressing the Glut4
glucose transporter. The Journal of Biological Chemistry 270(4): 1679–84, Doi:
10.1074/jbc.270.5.1679.

[41] Hansen, P.A., Marshall, B.A., Chen, M., Holloszy, J.O., Mueckler, M., 2000. Transgenic
Overexpression of Hexokinase II in Skeletal Muscle Does Not Increase Glucose Disposal in Wild-
type or Glut1-overexpressing Mice. Journal of Biological Chemistry 275(29): 22381–6, Doi:

10.1074/jbc.M001946200.

[42]   Zisman, A., Peroni, O.D., Abel, E.D., Michael, M.D., Mauvais-Jarvis, F., Lowell, B.B., et al., 2000.
Targeted disruption of the glucose transporter 4 selectively in muscle causes insulin resistance and
glucose intolerance. Nature Medicine 6(8): 924–8, Doi: 10.1038/78693.

[43]   Deacon, S.W., Beeser, A., Fukui, J.A., Rennefahrt, U.E.E., Myers, C., Chernoff, J., et al., 2008. An
Isoform-Selective, Small-Molecule Inhibitor Targets the Autoregulatory Mechanism of p21-
Activated Kinase. Chemistry and Biology 15(4): 322–31, Doi: 10.1016/j.chembiol.2008.03.005.

[44]   Ahn, M., Yoder, S.M., Wang, Z., Oh, E., Ramalingam, L., Tunduguru, R., et al., 2016. The p21-
activated kinase (PAK1) is involved in diet-induced beta cell mass expansion and survival in mice
and human islets. Diabetologia 59(10): 2145–55, Doi: 10.1007/s00125-016-4042-0.

[45]   Joseph, G.A., Hung, M., Goel, A.J., Hong, M., Rieder, M.-K., Beckmann, N.D., et al., 2019. Late-
onset megaconial myopathy in mice lacking group I Paks. Skeletal Muscle 9(1): 5, Doi:
10.1186/s13395-019-0191-4.

[46]   Lansey, M.N., Walker, N.N., Hargett, S.R., Stevens, J.R., Keller, S.R., 2012. Deletion of Rab GAP
AS160 modifies glucose uptake and GLUT4 translocation in primary skeletal muscles and adipocytes
and impairs glucose homeostasis. American Journal of Physiology-Endocrinology and Metabolism
303(10): E1273–86, Doi: 10.1152/ajpendo.00316.2012.

[47]   Wang, H.Y., Ducommun, S., Quan, C., Xie, B., Li, M., Wasserman, D.H., et al., 2013. AS160
deficiency causes whole-body insulin resistance via composite effects in multiple tissues.
Biochemical Journal 449(2): 479–89, Doi: 10.1042/BJ20120702.

[48]   Lund, S., Holman, G.D., Zierath, J.R., Rincon, J., Nolte, L.A., Clark, A.E., et al., 1965. Effect of
Insulin on GLUT4 CeU Surface Content and Turnover Rate in Human Skeletal Muscle as Measured
by the Exofacial Bis-Mannose Photolabeling Technique.

[49]   Somwar, R., Niu, W., Kim, D.Y., Sweeney, G., Randhawa, V.K., Huang, C., et al., 2001. Differential
Effects of Phosphatidylinositol 3-Kinase Inhibition on Intracellular Signals Regulating GLUT4
Translocation and Glucose Transport. Journal of Biological Chemistry 276(49): 46079–87, Doi:
10.1074/jbc.M109093200.

[50]   Somwar, R., Kim, D.Y., Sweeney, G., Huang, C., Niu, W., Lador, C., et al., 2001. GLUT4
translocation precedes the stimulation of glucose uptake by insulin in muscle cells: potential
activation of GLUT4 via p38 mitogen-activated protein kinase. The Biochemical Journal 359(Pt 3):
639–49, Doi: 10.1042/0264-6021:3590639.

[51]   Funaki, M., Randhawa, P., Janmey, P.A., 2004. Separation of Insulin Signaling into Distinct GLUT4
Translocation and Activation Steps. Molecular and Cellular Biology 24(17): 7567–77, Doi:
10.1128/mcb.24.17.7567-7577.2004.

[52]   Davies, S.P., Reddy, H., Caivano, M., Cohen, P., 2000. Specificity and mechanism of action of some
commonly used protein kinase inhibitors. Biochem. J 351: 95–105, Doi: 10.1517/13543776.11.3.405.

[53]   Nozaki, S., Ueda, S., Takenaka, N., Kataoka, T., Satoh, T., 2012. Role of RalA downstream of Rac1
in insulin-dependent glucose uptake in muscle cells. Cellular Signalling 24(11): 2111–7, Doi:
10.1016/j.cellsig.2012.07.013.

[54]   Takenaka, N., Sumi, Y., Matsuda, K., Fujita, J., Hosooka, T., Noguchi, T., et al., 2015. Role for RalA
downstream of Rac1 in skeletal muscle insulin signalling. The Biochemical Journal 469(3): 445–54,
Doi: 10.1042/BJ20150218.

[55]   Bedard, K., Krause, K.-H., 2007. The NOX Family of ROS-Generating NADPH Oxidases:
Physiology and Pathophysiology. Physiological Reviews 87(1): 245–313, Doi:
10.1152/physrev.00044.2005.

[56] Abo, A., Pick, E., Hall, A., Totty, N., Teahan, C.G., Segal, A.W., 1991. Activation of the NADPH
oxidase involves the small GTP-binding protein p21rac1. Nature 353(6345): 668–70, Doi:
10.1038/353668a0.

[57] Contreras-Ferrat, A., Llanos, P., Vásquez, C., Espinosa, A., Osorio-Fuentealba, C., Arias-Calderon,
M., et al., 2014. Insulin elicits a ROS-activated and an IP3-dependent Ca2+ release, which both
impinge on GLUT4 translocation. Journal of Cell Science 127(9): 1911–23, Doi: 10.1242/jcs.138982.

[58] Henríquez-Olguin, C., Knudsen, J.R., Raun, S.H., Li, Z., Dalbram, E., Treebak, J.T., et al., 2019.
Cytosolic ROS production by NADPH oxidase 2 regulates muscle glucose uptake during exercise.
Nature Communications 10(1): 4623, Doi: 10.1038/s41467-019-12523-9.

[59] Sylow, L., Jensen, T.E., Kleinert, M., Mouatt, J.R., Maarbjerg, S.J., Jeppesen, J., et al., 2013. Rac1 Is
a Novel Regulator of Contraction-Stimulated Glucose Uptake in Skeletal Muscle. Diabetes 62(4):
1139–51, Doi: 10.2337/db12-0491.

Figure Legends

Figure 1: Pharmacological inhibition of group I PAKs partially reduces insulin-stimulated glucose uptake in mouse soleus muscle. (A-B) Insulin-stimulated (60 nM) 2-Deoxyglucose (2DG) uptake in isolated soleus (A) and extensor digitorum longus (EDL, B) muscle $\pm$ 40 μM IPA-3 or as a control DMSO (0.25%). Isolated muscles were pre-incubated for 25 minutes followed by 30 minutes of insulin stimulation with 2DG uptake analyzed for the final 10 minutes of stimulation. **(C-H)** Quantification of phosphorylated (p)PAK1 T423, pAkt T308, and pAkt S473 in insulin-stimulated soleus (C, E, and G) and EDL (D, F, and H) muscle $\pm$ 40 μM IPA-3 or as a control DMSO (0.25%). Some of the data points were excluded due to the quality of the immunoblot, and the number of determinations was $n = 8/7$ for pAkt S473 and T308 in soleus muscle. **(I-J)** Representative blots showing pPAK1 T423, pAkt T308, and pAkt S473 in soleus (I) and EDL (J) muscle. Statistics were evaluated with a two-way repeated measures (RM) ANOVA. Main effects are indicated in the panels. Interactions in two-way RM ANOVA were evaluated by Tukey's post hoc test: Insulin stimulation vs. basal **/*** ($p<0.01/0.001$); IPA-3 vs. DMSO (#)/#/##/### ($p<0.1/0.05/0.01/0.001$). Unless otherwise stated previously in the figure legend, the number of determinations in each group: Soleus, $n = 9/8$ (DMSO/IPA-3); EDL, $n = 9/8$. Data are presented as mean $\pm$ S.E.M. with individual data points shown. Paired data points are connected with a straight line. A.U., arbitrary units.

Figure 2: PAK1 knockout does not affect whole-body glucose tolerance or insulin-stimulated glucose uptake in isolated skeletal muscle. (A) Representative blots showing PAK1 protein expression in gastrocnemius, quadriceps, and triceps brachii muscle from PAK1 knockout (KO) mice or control littermates. **(B)** Body composition (FM: Fat mass; LBM: Lean body mass; BW: Body weight) in gram in chow-fed PAK1 KO mice ($n = 12$) or control littermates ($n = 13$). The mice were 7-19 weeks of age. Statistics were evaluated with a Student's t-test. **(C)** Energy intake in

chow-fed PAK1 KO mice ($n = 8$) or control littermates ($n = 8$). Energy intake was monitored in a separate cohort of mice. Statistics were evaluated with a Student's t-test. **(D)** Blood glucose levels during a glucose tolerance test (GTT) in chow-fed PAK1 KO mice ($n = 9$) or control littermates ($n = 10$). The mice were 8-20 weeks of age. Statistics were evaluated with a two-way repeated measures (RM) ANOVA. **(E)** Incremental Area Under the Curve (AUC) for blood glucose levels during the GTT in panel D. Statistics were evaluated with a Student's t-test. **(F)** Plasma insulin values during a GTT in chow-fed PAK1 KO mice ($n = 10$) or control littermates ($n = 13$). The mice were 9-21 weeks of age. Statistics were evaluated with a two-way RM measures ANOVA. **(G)** Homeostatic Model Assessment of Insulin Resistance (HOMA-IR) in chow-fed PAK1 KO mice ($n=10$) or control littermates ($n=13$). Statistics were evaluated with a Student's t-test. **(H-I)** Submaximal (0.6 nM) and maximal (60 nM) insulin-stimulated 2-Deoxyglucose (2DG) uptake in isolated soleus (H) and extensor digitorum longus (EDL; I) muscle from PAK1 KO mice or littermate controls. Isolated muscles were pre-incubated for 30 minutes followed by 30 minutes of insulin stimulation with 2DG uptake analyzed for the final 10 minutes of stimulation. The mice were 12-24 weeks of age. The number of determinations in each group: Soleus-Control, $n = 6/7$ (Submax/max); Soleus-PAK1 KO, $n = 7/7$; EDL-Control, $n = 6/6$; EDL-PAK1 KO, $n = 7/7$. Statistics were evaluated with two two-way RM measures ANOVA. Main effects are indicated in the panels. Data are presented as mean $\pm$ S.E.M. or when applicable mean $\pm$ S.E.M. with individual data points shown. Paired data points are connected with a straight line.

Figure 3: PAK1 is dispensable for insulin-stimulated glucose uptake in lean or diet-induced insulin-resistant mice in vivo. (A-B) Blood glucose levels during a retro-orbital insulin tolerance test (r.o. ITT) in chow- (A) and HFD-fed (B) PAK1 knockout (KO) mice or control littermates. Chow-fed mice were investigated at 10-24 weeks of age. Mice fed a 60E% high-fat diet (HFD) for 21 weeks were investigated at 27-37 weeks of age. The number of mice in each group: Chow, $n =$

12/6 (Control/PAK1 KO); HFD, *n = 10/11*. Statistics were evaluated with a two-way repeated measures ANOVA. **(C-D)** Insulin-stimulated (Chow: 0.5 U kg^{-1} body weight; HFD: 60% of insulin administered to chow-fed mice) glucose uptake index in gastrocnemius (Gast), quadriceps (Quad), triceps brachii (Triceps) and soleus muscle from chow- (C) and HFD-fed (D) PAK1 KO mice or control littermates. The number of mice in each group: Chow-Saline, *n = 12/4* (Control/PAK1 KO); Chow-Insulin, *n = 12/6*; HFD-Saline, *n = 6/6*; HFD-Insulin, *n = 10/11*. Statistics were evaluated with a two-way ANOVA for each of the muscles. Main effects are indicated in the panels. Data are presented as mean $\pm$ S.E.M. or when applicable mean $\pm$ S.E.M. with individual data points shown.

Figure 4: Whole-body substrate utilization is unaffected by genetic ablation of PAK1 and PAK2. (A) Representative blots showing PAK1 and PAK2 protein expression in soleus and extensor digitorum longus (EDL) muscle from chow-fed whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (1/m2 dKO) mice or control littermates. **(B)** Body weight of chow-fed PAK1 KO (*n = 18*), PAK2 mKO (*n = 21*), 1/m2 dKO mice (*n = 23*) or control littermates (*n = 20*). The mice were 16-29 weeks of age. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with one-way ANOVA. (**C**) Body composition (FM: Fat mass; LBM: Lean body mass) in gram in chow-fed PAK1 KO (*n = 18*), PAK2 mKO (*n = 21*), 1/m2 dKO mice (*n = 23*) or control littermates (*n = 20*). Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. **(D-G)** Oxygen uptake (VO$_2$; D), respiratory exchange ratio (RER; E), activity (beam breaks; F), and energy intake (G), in chow-fed PAK1 KO (*n = 11*), PAK2 mKO (*n = 11*), 1/m2 dKO mice (*n =*

13) or control littermates (*n = 8; for energy intake, n = 7*) during the light and dark period recorded over a period of 72 hours. On day 2, the mice were fasted during the dark period and then refed on day 3. The mice were 23-33 weeks of age. Statistics were evaluated with two two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) in the light and dark period of day 1, respectively. Statistics for day 2 and 3 were evaluated similarly thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of the light and dark period were assessed with three two-way repeated measures (RM) ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. 1/m2 dKO) and 'Period' (Light vs. Dark) at day 1, 2 and 3, respectively. Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way (RM when applicable) ANOVA were evaluated by Tukey's post hoc test: Light vs. dark period */**/*** ($p<0.05/0.01/0.001$); Control vs. 1/m2 dKO †/††† ($p<0.05/0.001$); PAK1 KO vs. 1/m2 dKO ‡/‡‡/‡‡‡ ($p<0.05/0.01/0.001$); PAK2 mKO vs. 1/m2 dKO (\$)/\$/\$\$ ($p<0.1/0.05/0.01$). Data are presented as mean $\pm$ S.E.M. or when applicable mean $\pm$ S.E.M. with individual data points shown.

Figure 5: Glucose tolerance is reduced in mice lacking PAK2 in skeletal muscle. (A) Blood glucose concentration in the fed state (8 a.m.) in chow-fed whole-body PAK1 knockout (KO) (*n = 18*), muscle-specific PAK2 (m)KO (*n = 21*), PAK1/2 double KO (1/m2 dKO) mice (*n = 23*) or control littermates (*n = 20*). The mice were 17-30 weeks of age. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. **(B)** Blood glucose levels during a glucose tolerance test (GTT) in chow-fed PAK1 KO (*n = 18*), PAK2 mKO (*n = 19*), 1/m2 dKO mice (*n = 21*) or control littermates (*n = 19*). The mice were 13-26 weeks of age. Statistics were evaluated with six two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs.

PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) at time point 0, 15, 30, 60, 90 and 120, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of glucose administration were assessed with a two-way repeated measures (RM) ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. 1/m2 dKO) and 'Time' (0 vs. 15 vs. 30 vs. 60 vs. 90 vs. 120). **(C)** Incremental Area Under the Curve (AUC) for blood glucose levels during the GTT in panel B. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with one-way ANOVA. **(D)** Plasma insulin values during a GTT in chow-fed PAK1 KO (*n = 17*), PAK2 mKO (*n = 19*), 1/m2 dKO mice (*n = 22*) or control littermates (*n = 19*). The mice were 15-28 weeks of age. Statistics were evaluated with two two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) at time point 0 and 20, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of glucose administration were assessed with a two-way RM ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. 1/m2 dKO) and 'Time' (0 vs. 20). **(E)** Homeostatic Model Assessment of Insulin Resistance (HOMA-IR) in chow-fed PAK1 KO (*n = 18*), PAK2 mKO (*n = 19*), 1/m2 dKO mice (*n = 22*) or control littermates (*n = 20*). Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. **(F)** Blood glucose levels related to the basal concentration during an insulin tolerance test (ITT) in chow-fed PAK1 KO (*n = 15*), PAK2 mKO (*n = 18*), 1/m2 dKO mice (*n = 21*) or control littermates (*n = 19*). The mice were 11-24 weeks of age. For two female control mice and four female PAK2

mKO mice, the ITT had to be stopped before the 120'-time point due to hypoglycemia (blood glucose <1.2 mM), and thus blood glucose was not determined for these mice for the last couple of time points. Statistics were evaluated with five two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) at time point 15, 30, 60, 90 and 120, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of insulin administration were assessed with a mixed-effects model analysis to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. 1/m2 dKO) and 'Time' (0 vs. 15 vs. 30 vs. 60 vs. 90 vs. 120). Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way (RM when applicable) ANOVA were evaluated by Tukey's post hoc test: Effect of glucose/insulin administration vs. time point 0' */*** (p<0.05/0.001). Data are presented as mean ± S.E.M. or when applicable mean ± S.E.M. with individual data points shown. Paired data points are connected with a straight line.

Figure 6: Insulin-stimulated glucose uptake relies partially on PAK2 in EDL, but not soleus muscle, while PAK1 is not involved. (A-B) Insulin-stimulated (60 nM) 2-Deoxyglucose (2DG) uptake in isolated soleus (A) and extensor digitorum longus (EDL; B) muscle from whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (1/m2 dKO) mice or control littermates. Isolated muscles were pre-incubated for 20 minutes followed by 20 minutes of insulin stimulation with 2DG uptake analyzed for the final 10 minutes of stimulation. The mice were 10-16 weeks of age. Statistics were evaluated with two two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) in basal and insulin-stimulated samples, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of insulin stimulation were assessed with a two-way repeated measures (RM) ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. 1/m2 dKO) and 'Stimuli' (Basal vs. Insulin). **(C-D)** Δ2DG uptake in soleus (C)

and EDL (D) muscle from panel C-D. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) to assess of the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. The number of determinations in each group: Control, $n = 9/10$ (soleus/EDL); PAK1 KO, $n = 8/9$; PAK2 KO, $n = 10/10$; 1/m2 dKO, $n = 9/9$. Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way (RM when applicable) ANOVA were evaluated by Tukey's post hoc test: Insulin-stimulation vs. basal control *** ($p<0.001$); Control vs. PAK2 mKO £ ($p<0.05$); PAK1 KO vs. PAK2 mKO § ($p<0.05$). Data are presented as mean $\pm$ S.E.M. with individual data points shown. Paired data points are connected with a straight line.

Figure 7: PAK2 regulates TBC1D4 protein expression and signalling. (A-J) Quantification of phosphorylated (p)Akt S473, pAkt T308, pTBC1D4 T642 and total TBC1D4 and GLUT4 protein expression in insulin-stimulated (60 nM) soleus (A, C, E, G, and I) and extensor digitorum longus (EDL; B, D, F, H, and J) muscle from whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (1/m2 dKO) mice or control littermates. The mice were 10-16 weeks of age. Total protein expression is an average of the paired basal and insulin-stimulated sample. Some of the data points were excluded due to the quality of the immunoblot, so the number of determinations for GLUT4 in soleus muscle: Control, $n = 6$; PAK1 KO, $n = 5$; PAK2 KO, $n = 6$; 1/m2 dKO, $n = 6$. **(K-L)** Representative blots showing pAkt S473, pAkt T308, pTBC1D4 T642 and total PAK1, PAK2, Akt2, TBC1D4 and GLUT4 protein expression and coomassie staining as a loading control in soleus (K) and EDL (L) muscle. Statistics for phosphorylated proteins were evaluated with a two two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) in basal and insulin-stimulated samples, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between

genotypes and the effect of insulin stimulation were assessed with a two-way repeated measures (RM) ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. 1/m2 dKO) and 'Stimuli' (Basal vs. Insulin). Statistics for total protein expression were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way (RM when applicable) ANOVA were evaluated by Tukey's post hoc test: Control vs. PAK2 mKO £ (p<0.05). Unless otherwise stated previously in the figure legend, the number of determinations in each group: Control, *n = 9/10* (soleus/EDL); PAK1 KO, *n = 8/9*; PAK2 KO, *n = 10/9*; 1/m2 dKO, *n = 9/9*. Data are presented as mean ± S.E.M. with individual data points shown. Paired data points are connected with a straight line. A.U., arbitrary units.

Table 1. Antibody Table

Antibody name	Antibody ID (RRID)	Manufacturer; Catalog Number;	Species Raised in; Monoclonal or Polyclonal	Antibody dilution	Blocking buffer
Akt2	AB_2225186	Cell Signaling Technology; 3063	Rabbit, Monoclonal antibody	1:1000	2% milk
pAkt S473	AB_329825	Cell Signaling Technology; 9271	Rabbit; Polyclonal antibody	1:1000	2% milk
pAkt T308	AB_329828	Cell Signaling Technology; 9275	Rabbit; Polyclonal antibody	1:1000	2% milk
GLUT4	AB_2191454	Thermo Fisher Scientific; PA1-1065	Rabbit; Polyclonal antibody	1:1000	2% milk
PAK1	AB_330222	Cell Signaling Technology; 2602	Rabbit; Polyclonal antibody	1:1000	2% milk
PAK2	AB_2283388	Cell Signaling Technology; 2608	Rabbit; Polyclonal antibody	1:1000	2% milk
pPAK1/2 T423/402	AB_330220	Cell Signaling Technology; 2601	Rabbit; Polyclonal antibody	1:1000	5% BSA
TBC1D4	AB_492639	Millipore; 07-741	Rabbit; Polyclonal antibody	1:1000	2% milk
pTBC1D4 T642	AB_2651042	Cell Signaling Technology; 8881	Rabbit; Monoclonal antibody	1:1000	2% milk

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

Soleus

Main effect: Insulin-stimulation (p<0.001)

2DG uptake (µmol g^{-1} h^{-1})

Insulin (60 nM)   Control   PAK1 KO   PAK2 mKO   1/m2 dKO

B

EDL

Main effect: Muscle-specific PAK2 knockout at insulin-stimulation (p<0.05)

2DG uptake (µmol g^{-1} h^{-1})

£

*** *** *** ***

Insulin (60 nM)   Control   PAK1 KO   PAK2 mKO   1/m2 dKO

C

Soleus

Δ2DG uptake (µmol g^{-1} h^{-1})

Control   PAK1 KO   PAK2 mKO   1/m2 dKO

D

EDL

Main effect: Muscle-specific PAK2 knockout (p<0.01)

Δ2DG uptake (µmol g^{-1} h^{-1})

§

Control   PAK1 KO   PAK2 mKO   1/m2 dKO

Figure 7

Insulin-stimulated glucose uptake partly relies on p21-activated kinase (PAK)-2, but not PAK1, in mouse skeletal muscle

Lisbeth L. V. Møller[1], Merna Jaurji[1], Rasmus Kjøbsted[1], Giselle A. Joseph[2], Agnete B. Madsen[1], Jonas R. Knudsen[1], Annemarie Lundsgaard[1], Nicoline R. Andersen[1], Peter Schjerling[3,4], Thomas E. Jensen[1], Robert S. Krauss[2], Erik A. Richter[1*], and Lykke Sylow[1,*,#]

[1] Section of Molecular Physiology, Department of Nutrition, Exercise and Sports, Faculty of Science, University of Copenhagen, DK-2100 Copenhagen, Denmark

[2] Department of Cell, Developmental, and Regenerative Biology, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

[3] Institute of Sports Medicine, Department of Orthopaedic Surgery M, Bispebjerg Hospital, DK-2400 Copenhagen, Denmark

[4] Center for Healthy Aging, Faculty of Health Sciences, University of Copenhagen, DK-2100 Copenhagen, Denmark

[*] *Authors contributed equally*

[#]*Corresponding author and person to who reprint requests should be addressed:*

Lykke Sylow. E-mail: Lshansen@nexs.ku.dk. Tel.: +4520955250

Section of Molecular Physiology, Department of Nutrition, Exercise and Sports, August Krogh Building, 13 Universitetsparken, DK-2100 Copenhagen, Denmark

Short title: Group I PAKs and insulin-stimulated skeletal muscle glucose uptake

Table S1. Overview of the different cohorts of genetically modified mice

Cohort	Diet	Number of replicates	Age (weeks)						Terminal experiment
			Start diet intervention	Insulin tolerance	Glucose tolerance	Body composition	Home cage calorimetry	Tissue dissection	
Whole-body PAK1-/- mice	Chow	Control, n = 6/7 (female/male) PAK1 KO, n = 4/8			8-20	7-19		12-24	Insulin-stimulated glucose uptake in isolated muscle
Whole-body PAK1-/- mice	Chow	Control, n = 14/8 (female/male) PAK1 KO, n = 6/4						10-24	In vivo glucose uptake
Whole-body PAK1-/- mice	60E% HFD	Control, n = 7/7 (female/male) PAK1 KO, n = 11/5	6-16		20-30	24-34		27-37	In vivo glucose uptake
Double PAK1$^{-/-}$;PAK2$^{fl/fl}$;MyoD$^{iCre/+}$ mice	Chow	Control, n = 6/4 (female/male) PAK1 KO, n = 5/4 PAK2 mKO, n = 6/4 1/m2 dKO, n = 6/3						10-16	Insulin-stimulated glucose uptake in isolated muscle
Double PAK1$^{-/-}$;PAK2$^{fl/fl}$;MyoD$^{iCre/+}$ mice	Chow	Control, n = 9/11 (female/male) PAK1 KO, n = 8/10 PAK2 mKO, n = 12/9 1/m2 dKO, n = 9/14.		11-24	13-26	16-29	23-33		

Figure S1

Figure S1: (A-B) Insulin-stimulated (Chow: 0.5 U kg^{-1} body weight; HFD: 60% of insulin administered to chow-fed mice) 2-Deoxyglucose (2DG) clearance in gastrocnemius (Gast), quadriceps (Quad), triceps brachii (Triceps) and soleus muscle from chow- (A) and 60E% high-fat diet (HFD)-fed (B) PAK1 knockout (KO) mice or control littermates. The number of mice in each group: Chow-Saline, *n = 12/4* (Control/PAK1 KO); Chow-Insulin, *n = 12/6*; HFD-Saline, *n = 6/6*; HFD-Insulin, *n = 10/11*. Statistics were evaluated with a two-way ANOVA for each of the muscles. **(C)** Plasma [^{3}H] counts 10 minutes after retro-orbital (r.o.) administration of a bolus of saline containing [^{3}H]-labelled 2DG ([^{3}H]-2DG) with or without insulin. Statistics were evaluated with two two-way ANOVAs to test the factors 'stimuli' (basal vs. insulin) and 'genotype' (control vs. PAK1 KO) in chow-fed and HFD-fed mice, respectively. **(D)** Body composition (FM: Fat mass; LBM: Lean body mass; BW: Body weight) in gram in HFD-fed PAK1 KO mice (*n = 17*) or control littermates (*n = 14*). Body composition was assessed in week 18-19 of the diet intervention. The mice were 24-34 weeks of age. Statistics were evaluated with a Student's t-test. **(E)** Energy intake in HFD-fed PAK1 KO mice (*n = 8*) or control littermates (*n = 8*). Energy intake was monitored in a separate cohort of mice. Statistics were evaluated with a Student's t-test. **(F)** Blood glucose levels during a glucose tolerance test (GTT) in HFD-fed PAK1 KO mice (*n = 17*) or control littermates (*n = 13*). Glucose tolerance was assessed in week 14 of the diet intervention. The mice were 20-30 weeks of age. Statistics were evaluated with a two-way repeated measures (RM) ANOVA. **(G)** Incremental Area Under the Curve (AUC) for blood glucose levels during the GTT in panel F. Statistics were evaluated with a Student's t-test. **(H)** Plasma insulin values during a GTT in HFD-fed PAK1 KO mice (*n = 16*) or control littermates (*n = 11*). The plasma insulin response to glucose administration was measured in week 16 of the diet intervention. The mice were 22-32 weeks of age. Statistics were evaluated with a two-way RM ANOVA. **(I)** Homeostatic Model Assessment of Insulin Resistance (HOMA-IR) in HFD-fed PAK1 KO mice (*n = 16*) or control littermates (*n =*

11). Statistics were evaluated with a Student's t-test. Main effects are indicated in the panels. Data are presented as mean $\pm$ S.E.M. or when applicable mean $\pm$ S.E.M. with individual data points shown. Paired data points are connected with a straight line.

Figure S2

Figure S2: (A-B) Body weight of female (A) and male (B) whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (1/m2 dKO) mice or control littermates. **(C-D)** Body composition (FM: Fat mass; LBM: Lean body mass) in gram in female (C) and male (D) PAK1 KO, PAK2 mKO, 1/m2 dKO or control littermates. **(E-G)** Body composition (FM: Fat mass; LBM: Lean body mass) in percentage in both sexes combined (E) and in female (F) and male (G) PAK1 KO, PAK2 mKO, 1/m2 dKO or control littermates. The number of mice in each group: Control, $n = 9/11$ (female/male); PAK1 KO, $n = 8/10$; PAK2 mKO, $n = 12/9$; 1/m2 dKO, $n = 9/14$. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way ANOVA were evaluated by Tukey's post hoc test: Control vs. 1/m2 dKO †/†† ($p<0.05/0.01$); PAK1 KO vs. PAK2 mKO § ($p<0.05$); PAK1 KO vs. 1/m2 dKO ‡/‡‡/‡‡‡ ($p<0.05/0.01/0.001$); PAK2 mKO vs. 1/m2 dKO ($)/$$ ($p<0.1/0.01$). Data are presented as mean ± S.E.M. with individual data points shown.

Figure S3

Figure S3: (A-F) Oxygen uptake (VO$_2$), respiratory exchange ratio (RER), and activity (beam breaks) in female (A, C, and E) and male (B, D, and F) whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (1/m2 dKO) mice or control littermates during the light and dark period recorded over a period of 72 hours in calorimetric chambers. On day 2, mice fasted during the dark period and then refed on day 3. **(G-I)** Total energy intake on day 1 and day 3 in both sexes combined (G) and in female (H) and male (I) whole-body PAK1 KO, PAK2 mKO, 1/m2 dKO mice or control littermates recorded over a period of 72 hours in calorimetric chambers. **(J-K)** Energy intake in female (J) and male (K) whole-body PAK1 KO, Pak2 mKO, 1/m2 dKO mice or control littermates during the light and dark period recorded over a period of 72 hours in calorimetric chambers. On day 2, mice fasted during the dark period and were then refed on day 3. The number of mice in each group: Control, $n = 5/3$ (female/male; for energy intake, $n = 4/3$); PAK1 KO, $n = 6/5$; PAK2 mKO, $n = 6/5$; 1/m2 dKO, $n = 6/7$. **(A-F+J-K)** Statistics were evaluated with two two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) in the light and dark period of day 1, respectively. Statistics for day 2 and 3 were evaluated similarly thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of the light and the dark period were assessed with three two-way repeated measures (RM) ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. 1/m2 dKO) and 'Period' (Light vs. Dark) at day 1, 2 and 3, respectively. **(G-I)** Statistics were evaluated with two two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) at day 1 and day 2, respectively. Differences between genotypes and the day were assessed with one two-way repeated measures (RM) ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. 1/m2 dKO) and 'Day' (Day 1 vs. Day 3). Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way (RM when applicable) ANOVA were

evaluated by Tukey's post hoc test: Light vs. dark period */**/*** (p<0.05/0.01/0.001); Day 1 vs. Day 3 €€/€€€ (p<0.01/0.001); Control vs. PAK1 KO ¤¤ (p<0.01); Control vs. 1/m2 dKO † (p<0.05); PAK1 KO vs. PAK2 mKO §/§§ (p<0.05/0.01); PAK1 KO vs. 1/m2 dKO ‡/‡‡/‡‡‡ (p<0.05/0.01/0.001). Data are presented as mean ± S.E.M. with individual data points shown.

Figure S4

Figure S4: (A-B) Blood glucose concentration in the fed state (8 a.m.) in female (A) and male (B) whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (1/m2 dKO) mice or control littermates. The number of mice in each group: Control, $n = 9/11$ (female/male); PAK1 KO, $n = 8/10$; PAK2 mKO, $n = 12/9$; 1/m2 dKO, $n = 9/14$. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. **(C-D)** Blood glucose levels during a glucose tolerance test (GTT) in female (C) and male (D) PAK1 KO, PAK2 mKO, 1/m2 dKO mice or control littermates. The number of mice in each group: Control, $n = 8/11$ (female/male); PAK1 KO, $n = 8/10$; PAK2 mKO, $n = 10/9$; 1/m2 dKO, $n = 8/13$. Statistics were evaluated with six two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) at time point 0, 15, 30, 60, 90 and 120, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of glucose administration were assessed with a two-way repeated measures (RM) ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. 1/m2 dKO) and 'Time' (0 vs. 15 vs. 30 vs. 60 vs. 90 vs. 120). **(E-F)** Incremental Area Under the Curve (AUC) for blood glucose levels for females (E) and male (F) mice during the GTT in panel C-D. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. **(G-H)** Plasma insulin values during a GTT in female (G) and male (F) PAK1 KO, PAK2 mKO, 1/m2 dKO mice or control littermates. The number of mice in each group: Control, $n = 9/10$ (female/male); PAK1 KO, $n = 8/9$; PAK2 mKO, $n = 10/9$; 1/m2 dKO, $n = 9/13$. Statistics were evaluated with two two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and

'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) at time point 0 and 20, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of glucose administration were assessed with a two-way RM ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. 1/m2 dKO) and 'Time' (0 vs. 20). Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way (RM when applicable) ANOVA were evaluated by Tukey's post hoc test: Effect of glucose administration vs. time point 0' */*** ($p < 0.05/0.001$); Control vs. PAK2 mKO £ ($p < 0.05$); PAK1 KO vs. PAK2 mKO §§ ($p < 0.01$); PAK2 mKO vs. 1/m2 dKO \$ ($p < 0.05$). Data are presented as mean $\pm$ S.E.M. or when applicable mean $\pm$ S.E.M. with individual data points shown. Paired data points are connected with a straight line.

Figure S5

Figure S5: (A-B) Homeostatic Model Assessment of Insulin Resistance (HOMA-IR) in female (A) and male (B) whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (1/m2 dKO) mice or control littermates. The number of mice in each group: Control, *n = 9/11* (female/male); PAK1 KO, *n = 8/10*; PAK2 mKO, *n = 10/9*; 1/m2 dKO, *n = 9/13*. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Genotype differences were evaluated with a one-way ANOVA. **(C-E)** Basal blood glucose concentration (fasted state) immediately before an insulin tolerance test (ITT) in both sexes combined (C) and in female (D) and male (E) PAK1 KO, PAK2 mKO, 1/m2 dKO mice and control littermates. The number of mice in each group: Control, *n = 9/10* (female/male); PAK1 KO, *n = 6/9*; PAK2 mKO, *n = 10/8*; 1/m2 dKO, *n = 8/13*. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. **(F-G)** Blood glucose levels related to basal concentration during an ITT in female (F) and male (G) PAK1 KO, PAK2 mKO, 1/m2 dKO mice or control littermates. The number of mice in each group: Control, *n = 9/10* (female/male); PAK1 KO, *n = 6/9*; PAK2 mKO, *n = 10/8*; 1/m2 dKO, *n = 8/13*. For two female control mice and four female PAK2 mKO mice, the ITT had to be stopped before the 120'-time point due to hypoglycemia (blood glucose <1.2 mM), and thus blood glucose was not determined for these mice for the last couple of time points. Statistics were evaluated with five two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) at time point 15, 30, 60, 90 and 120, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of insulin administration were assessed with a two-way repeated measures (RM) ANOVA (mixed-effects

model analysis for female mice) to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. 1/m2 dKO) and 'Time' (0 vs. 15 vs. 30 vs. 60 vs. 90 vs. 120). Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way (RM when applicable) ANOVA were evaluated by Tukey's post hoc test: Effect of insulin administration vs. time point 0' **/*** (p<0.01/0.001); Control vs. PAK1 KO ¤/¤¤ (p<0.05/0.01); Control vs. PAK2 mKO (£)££ (p<0.1/0.01); Control vs. 1/m2 dKO ††† (p<0.001); PAK1 KO vs. 1/m2 dKO ‡ (p<0.05); PAK2 mKO vs. 1/m2 dKO $ (p<0.05). Data are presented as mean ± S.E.M. or when applicable mean ± S.E.M. with individual data points shown.

Figure S6: (A-D) Δ-phosphorylated (p)-Akt S473 and ΔpAkt T308 in insulin-stimulated (60 nM) soleus (A and C) and extensor digitorum longus (EDL; B and D) muscle from whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (1/m2 dKO) mice or control littermates from Fig. 7A-D. **(E-F)** Quantification of total Akt2 protein expression in soleus (E) and EDL (F) muscle from whole-body PAK1 KO, PAK2 mKO, 1/m2 dKO mice or control littermates. Total protein expression is an average of the paired basal and insulin-stimulated sample. **(G-H)** ΔpTBC1D4 T642 in insulin-stimulated (60 nM) soleus (G) and EDL (H) muscle from whole-body PAK1 KO, PAK2 mKO, 1/m2 dKO mice or control littermates from Fig. 7G-H. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. The number of determinations in each group: Control, *n = 9/10* (soleus/EDL); PAK1 KO, *n = 8/9*; PAK2 KO, *n = 10/9*; 1/m2 dKO, *n = 9/9*. Significant one-way ANOVA was evaluated by Tukey's post hoc test: Control vs. PAK2 mKO £ (p<0.05). Data are presented as mean $\pm$ S.E.M. with individual data points shown. A.U., arbitrary units.