Title:

Group I p21-activated kinases (PAKs) are largely dispensable for insulin-stimulated glucose

uptake in mouse skeletal muscle

Lisbeth L. V. Møller[1], Merna Jaurji[1], Rasmus Kjøbsted[1], Giselle A. Joseph[2], Agnete B. Madsen[1], Jonas R. Knudsen[1],

Annemarie Lundsgaard[1], Nicoline R. Andersen[1], Peter Schjerling[3,4], Thomas E. Jensen[1], Robert S. Krauss[2], Erik A.

Richter[1*], and Lykke Sylow[1*]

[1] Section of Molecular Physiology, Department of Nutrition, Exercise and Sports, Faculty of Science, University of

Copenhagen, Denmark

[2] Department of Cell, Developmental, and Regenerative Biology, Icahn School of Medicine at Mount Sinai, New York,

NY 10029 USA

[3] Institute of Sports Medicine, Department of Orthopedic Surgery M, Bispebjerg Hospital, Denmark

[4] Center for Healthy Aging, Faculty of Health Sciences, University of Copenhagen, Denmark

Corresponding authors and persons to who reprint requests should be addressed. Authors contributed

equally.

Lykke Sylow, Lshansen@nexs.ku.dk, phone +4520955250

Erik Arne Richter, Erichter@nexs.ku.dk, phone +4528751626

Universitetsparken 13, 2100 Copenhagen Oe, Denmark

Short title: Group I PAKs and insulin-stimulated skeletal muscle glucose uptake

Number of figures: 7 (+ 6 supplemental figures)

Non-standard abbreviations

2DG: 2-Deoxyglucose

AUC: Area under the curve

BCA: Bicinchoninic acid

BW: Body weight

dKO: Double knockout

EDL: Extensor digitorum longus

FM: Fat mass

GLUT4: Glucose transporter 4

GTT: Glucose tolerance test

HFD: High-fat diet

HOMA-IR: Homeostatic Model Assessment of Insulin Resistance

ITT: Insulin tolerance test

i.p.: Intraperitoneal

KO: Knockout

L6-GLUT4myc: Rat L6 skeletal muscle cells overexpressing myc-tagged GLUT4

LBM: Lean body mass

mKO: Muscle-specific knockout

NOX: NADPH oxidase

PAK: p21-activated kinase

PI3K: Phosphoinositide 3-kinase

RER: Respiratory exchange ratio

r.o.: Retro-orbital

VO_2: Oxygen uptake

Abstract

Glucose transport into skeletal muscle is essential for maintaining whole-body glucose homeostasis and accounts for the majority of glucose disposal in response to insulin. The group I p21-activated kinase (PAK) isoforms PAK1 and PAK2 are in skeletal muscle activated in response to insulin and evidence suggests that PAK1 is necessary for insulin-stimulated GLUT4 translocation. In accordance, insulin-induced PAK1 and PAK2 signalling are impaired in insulin-resistant skeletal muscle. However, the role of PAK1 and PAK2 in insulin-stimulated glucose uptake in mature skeletal muscle has not been determined. The aim of the present investigation was to determine the requirement for PAK1 and PAK2 in whole-body glucose homeostasis and insulin-stimulated glucose uptake in skeletal muscle. Therefore, glucose uptake was measured in isolated skeletal muscle incubated with a pharmacological inhibitor (IPA-3) of group I PAKs and in muscle from whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO and double whole-body PAK1 and muscle-specific PAK2 knockout mice. Whole-body respiratory exchange ratio, indicative of substrate utilization, was largely unaffected by lack of PAK1 and/or PAK2. Whole-body glucose tolerance was mildly impaired in PAK2 mKO, but not PAK1 KO mice. In contrast to a previous study of GLUT4 translocation in PAK1 KO mice, PAK1 KO muscles displayed normal insulin-stimulated glucose uptake *in vivo* and in isolated muscle. On the contrary, glucose uptake was slightly reduced in response to insulin in glycolytic extensor digitorum longus muscle lacking PAK2. In conclusion, the current study demonstrates that group I PAKs are largely dispensable for the regulation of whole-body glucose homeostasis and skeletal muscle glucose uptake. Thus, the present study challenges that group I PAKs, and especially PAK1, are necessary regulators of insulin-stimulated glucose uptake in skeletal muscle.

Introduction

Skeletal muscle acts as a major glucose sink for disposal of glucose from the blood and is therefore essential in maintaining whole-body glucose homeostasis. Skeletal muscle fulfils this function by increasing glucose uptake in response to insulin during conditions of elevated blood glucose. It is important to understand the mechanisms regulating glucose uptake by skeletal muscle, in part because skeletal muscles account for the majority of insulin-mediated whole-body glucose disposal [1,2] and muscle insulin resistance is an early defect in the pathophysiology of peripheral insulin resistance and type 2 diabetes mellitus [1,3].

Insulin stimulates glucose uptake in skeletal muscle by activation of a signalling cascade that leads to the translocation of glucose transporter (GLUT)4-containing vesicles to the sarcolemma and transverse tubuli [4]. This signalling cascade has been proposed to include activation of p21-activated kinase 1 (PAK1) downstream of PI3K [5–7]. PAKs are serine/threonine kinases and PAK1, PAK2 and PAK3 constitute the group I PAKs. Group I PAKs have been extensively studied as part of numerous signalling networks that regulate essential cellular activities, including cell proliferation, differentiation, apoptosis, and cytoskeleton dynamics [8,9]. The kinases are a downstream target of the Rho GTPases Cdc42 and Rac1 [10]. Previous studies suggest that only PAK1 and PAK2 are expressed in skeletal muscle, whereas PAK3 mRNA and protein expression is below the detection limit [6,8,11]. In muscle cells and mouse skeletal muscle, PAK1 are proposed to be required for GLUT4 translocation in response to insulin stimulation [6,7], likely downstream of Rac1 [12–17]. Thus, together with Akt, proposed to regulate GLUT4 translocation via phosphorylation of the Rab GTPase-activating protein TBC1D4 [18–20], Rac1-PAK1 activation is suggested to be necessary for insulin-stimulated GLUT4 translocation.

Upon activation of PAK1 and PAK2, conformational changes allow autophosphorylation of T423

and T402, respectively, thereby relieving the autoinhibition of PAK1 and PAK2 [10,21,22]. In

vastus lateralis muscle from subjects with obesity and type 2 diabetes, phosphorylation of PAK1/2

at T423/402 in response to insulin was 50% reduced compared to healthy controls [15]. Likewise,

insulin-stimulated pPAK1/2 T423/402 was diminished in palmitate-treated insulin-resistant L6

myotubes, even though upstream of PAK1/2, insulin-stimulated Rac1-GTP binding (i.e. activation)

was not impaired [23]. Together, these studies [15,23] associate dysregulated activity of PAK1 and

PAK2 with insulin resistance. In addition, a pharmacological inhibitor of group I PAKs, IPA-3

abolished insulin-stimulated GLUT4 translocation and glucose uptake into L6 myoblasts and

myotubes, respectively, overexpressing myc-tagged GLUT4 (L6-GLUT4myc) [6]. This indicates

that group I PAKs are required for insulin-stimulated glucose uptake. This effect has largely been

ascribed to PAK1, as whole-body genetic ablation of PAK1 in mice impaired glucose tolerance

[7,24] associated with a complete blockade of insulin-stimulated GLUT4 translocation in skeletal

muscle [7]. The suggested downstream mechanisms whereby PAK1 regulates GLUT4 translocation

include simultaneous cofilin-mediated actin depolymerization and N-WASP-cortactin-mediated

actin polymerization [6,25,26]. Further supporting PAK1 being the major isoform regulating

GLUT4 translocation, insulin-stimulated GLUT4 translocation was unaffected by a 75%

knockdown of PAK2 in L6-GLUT4myc myoblasts [6].

Although such studies implicate group I PAKs, and in particular PAK1, in the regulation of glucose

homeostasis and GLUT4 translocation in skeletal muscle, the relative role of PAK1 and PAK2 in

insulin-stimulated glucose uptake remains to be identified in mature skeletal muscle. Therefore, we

performed a systematic series of pharmacologic and genetic experiments to analyze the involvement

of group I PAKs in the regulation of insulin-stimulated glucose uptake in mouse skeletal muscle.

We hypothesized that group I PAKs, and in particular PAK1, would be necessary for glucose

uptake in response to insulin. Contradicting our hypothesis, our results revealed that group I PAKs

are largely dispensable for insulin-stimulated glucose uptake in skeletal muscle.

Results

Pharmacological inhibition of group I PAKs partially reduces insulin-stimulated glucose uptake in

mouse soleus muscle. PAK1, but not PAK2, has been suggested to regulate insulin-stimulated

GLUT4 translocation [6,7]. Whereas glucose uptake *in vivo* is influenced by glucose delivery,

GLUT4 translocation and muscle metabolism [27], glucose delivery is constant in isolated skeletal

muscle and surface membrane GLUT4 is the limiting factor [28–30]. Therefore, 2-Deoxyglucose

(2DG) uptake in isolated fibres likely reflects GLUT4 translocation. Consequently, to investigate

the involvement of group I PAKs in insulin-stimulated glucose uptake in mouse skeletal muscle, we

first analyzed 2DG uptake in isolated soleus and extensor digitorum longus (EDL) muscle in the

presence of a pharmacological inhibitor, IPA-3. IPA-3 is a well-characterized inhibitor of group I

PAKs (PAK1-3) [6,31] and shown to completely abolish insulin-stimulated GLUT4 translocation

and glucose uptake in L6-GLUT4myc myoblasts and myotubes, respectively [6]. In soleus, 2DG

uptake increased 4.5-fold upon maximal insulin-stimulation, an increase that was partly reduced (-

20%) in IPA-3-treated muscles (Fig. 1A). IPA-3 did not significantly (p=0.080) impair insulin-

stimulated (+2.4-fold) 2DG uptake in EDL muscle (Fig. 1B). We confirmed that IPA-3 treatment

completely abolished insulin-stimulated phosphorylation of (p)PAK1 T423 in both soleus and EDL

muscle (Fig. 1C+D). In contrast, insulin-stimulated pAkt T308 (Fig. 1E+F) and the mTORC2 site,

pAkt S473 (Fig. 1G+H) were unaffected by IPA-3 treatment, suggesting that IPA-3 did not interfere

with insulin signalling to Akt. Altogether this suggests that group I PAKs are only partially

involved in insulin-stimulated glucose uptake in isolated mouse muscle.

PAK1 knockout does not affect whole-body glucose tolerance or insulin-stimulated glucose uptake

in isolated skeletal muscle. Pharmacological inhibitors often have off-target effects [32]. Therefore,

we next sought to confirm our findings in mice with whole-body lack of the PAK1 isoform and

therefore a complete knockout of PAK1 in skeletal muscle (Fig. 2A). These whole-body PAK1

knockout (KO) mice have previously been reported to display abolished GLUT4 translocation in

skeletal muscle in response to insulin *in vivo* [7]. In chow-fed mice, fat mass, lean body mass, body

weight and energy intake (Fig. 2B-C) were similar between whole-body PAK1 KO and littermate

controls, as also reported previously [33]. During a glucose tolerance test (GTT), the lack of PAK1

had no effect on the blood glucose response (Fig. 2D-E) or plasma insulin concentration (Fig. 2F).

Additionally, Homeostatic Model Assessment of Insulin Resistance (HOMA-IR), a measure of

basal glucose homeostasis (Fig. 2G), and both submaximal and maximal insulin-stimulated 2DG

uptake in isolated soleus and EDL muscle were unaffected by PAK1 KO (Fig. 2H-I). Thus,

unexpectedly, genetic ablation of PAK1 alone did not impair whole-body glucose tolerance in

contrast to previous reports [6,9]. Neither did lack of PAK1 affect skeletal muscle insulin sensitivity

(submaximal insulin-stimulated glucose uptake) or insulin responsiveness (maximal insulin-

stimulated glucose uptake).

PAK1 is not required for insulin-stimulated glucose uptake in lean or diet-induced insulin-resistant

mice in vivo. Our findings on insulin-stimulated glucose uptake in isolated muscle from chow-fed

PAK1 KO mice conflicted with a previous study reporting impaired glucose tolerance in PAK1 KO

mice and defects in insulin-stimulated GLUT4 translocation in skeletal muscle *in vivo* [7].

Therefore, we further explored the effect of PAK1 KO on insulin-stimulated glucose uptake in

skeletal muscle *in vivo*. Additionally, we fed a subgroup of PAK1 KO and control littermate mice a

60E% high-fat diet (HFD) for 21 weeks to investigate the role of PAK1 in insulin-resistant muscle.

Insulin administration lowered blood glucose by 5.4 ± 0.5 mM (-47%) in chow-fed control mice

(Fig. 3A). In HFD-fed control mice, blood glucose dropped 3.0 ± 0.9 mM (-26%) upon insulin administration (Fig. 3B). Lack of PAK1 had no impact on either basal blood glucose or whole-body insulin tolerance on either of the diets (Fig. 3A-B). Insulin increased glucose uptake in muscles from chow-fed (Gastrocnemius: 8.1-fold; Quadriceps: 8.5-fold, Triceps brachii: 12.3-fold; Soleus: 8.9-fold) and HFD-fed (Gastrocnemius: 3.5-fold; Quadriceps: 1.8-fold, Triceps brachii: 4.3-fold; Soleus: 11.6-fold) control mice (Fig. 3C-D). Consistent with our findings in isolated muscle, we observed no effect of PAK1 KO on basal or insulin-stimulated glucose uptake *in vivo* in muscle of either chow-fed mice or in insulin-resistant muscles from HFD-fed mice. Like glucose uptake, 2DG clearance from the plasma was unaffected by PAK1 KO in all of the investigated muscles (Fig. S1A-B). Importantly, circulating [^{3}H]-2DG availability was unaffected by genotype on both of the diets (Fig. S1C). As in chow-fed mice, lack of PAK1 in HFD-fed mice had no effect on fat mass, lean body mass, body weight or energy intake (Fig. S1D-E). Similarly, whole-body glucose tolerance (Fig. S1F-G), plasma insulin concentration during the GTT (Fig. S1H) and HOMA-IR (Fig. S1I) were unaffected by PAK1 KO in HFD-fed mice. Thus, PAK1 is dispensable for *in vivo* insulin-stimulated muscle glucose uptake in both the healthy lean and the diet-induced insulin-resistant state.

Whole-body substrate utilization is unaffected by genetic ablation of PAK1 and PAK2. Because of discrepancies in the data resulting from the use of a global pharmacological inhibitor of group I PAKs and data resulting from PAK1 KO mice, we next sought to determine the relative contribution and involvement of PAK1 and PAK2 in insulin signalling and glucose uptake in skeletal muscle. Double knockout mice with whole-body knockout of PAK1 and muscle-specific knockout of PAK2 (d1/2 KO) were generated as previously described [8]. By crossing d1/2 KO mice with littermate controls, a cohort was generated consisting of whole-body PAK1 KO, muscle-specific PAK2 (m)KO, d1/2 KO and littermate control mice. While no band for PAK1 could be

detected in muscles lacking PAK1, muscles lacking PAK2 displayed only a partial reduction in

band intensity in the immunoblots for PAK2 (Fig. 4A). This is likely due to the fact that PAK1 KO

mice are whole-body knockouts, while PAK2 mKO mice are muscle-specific and other cell types

within skeletal muscle tissue could thus contribute to the signal obtained in the PAK2 immunoblots.

In a previous study, PAK3 was not detectable at the protein level in d1/2 KO muscle [8].

As previously shown [8,34], d1/2 KO mice weighed less (-12%) than control littermates (Fig. 4B,

Fig. S2A-B) due to reduced (-12%) lean body mass (Fig. 4C, Fig. S2C-D). Body weight and lean

body mass decreased to the same extent in d1/2 KO mice, leaving lean body mass percentage

largely unaffected (Fig. S2E-G). Using calorimetric chambers, we monitored whole-body

metabolism for 72 hours during the light and dark period. On day 2, the mice fasted during the dark

period followed by refeeding on day 3. Oxygen uptake (VO$_2$; Fig. 4D, Fig. S3A-B) and respiratory

exchange ratio indicative of substrate utilization (RER; Fig. 4E, Fig. S3C-D) were largely

unaffected by genotype with only a slightly higher RER in mice lacking PAK2 (PAK2 mKO and

d1/2 KO mice) upon fasting. Similar substrate utilization was obtained despite reduced habitual

activity in the dark period in mice lacking PAK2, an effect largely driven by a decreased activity in

d1/2KO mice and an increased activity in male PAK1 KO mice (Fig. 4F; Fig. S3E-F). Supporting

the lower activity levels, energy intake was decreased (-11%) in d1/2 KO mice compared to PAK1

KO mice on day 1 (Fig. S3G) due to lower energy intake in the dark period (Fig. 4G). Upon

refeeding, energy intake was reduced in mice lacking PAK2 (Fig 4G; Fig. S3G) driven by a lower

energy intake in the dark period in female mice lacking PAK2 (Fig S3H-K). Altogether, these data

suggest that lack of PAK1 and/or PAK2 are not compromising metabolic regulation during the

light/dark period or in response to fasting/refeeding.

Glucose tolerance is slightly reduced in mice lacking PAK2 in skeletal muscle. To test dependency

on PAK1 and/or PAK2 in glucose handling and insulin sensitivity, we next investigated glucose and

insulin tolerance in the d1/2 KO mouse strain. Blood glucose concentration in the fed state was

similar between the genotypes (Fig. 5A, Fig. S4A-B). The blood glucose response to a glucose load

was slightly increased in mice lacking PAK2 in skeletal muscle as evident by the increased area

under the blood glucose curve (Fig. 5B-C). This was primarily driven by impaired glucose tolerance

in female PAK2 mKO mice (Fig. S4C-F). Plasma insulin concentration during the GTT was

unaffected by lack of PAK1 (Fig. 5D) In contrast, plasma insulin in male PAK2 mKO mice was

slightly elevated compared to d1/2 KO mice and tended (p=0.055) to be higher than control

littermates (Fig. S4G-H). This indicates impaired insulin sensitivity in male PAK2 mKO mice, but

HOMA-IR was unaffected by lack of PAK1 and/or PAK2 (Fig. 5E, Fig. S5A-B). In addition, even

though fasted blood glucose immediately before an insulin tolerance test (ITT) was modestly

reduced in PAK2 mKO mice (Fig. S5C-E), the blood glucose response to an ITT was largely

unaffected by lack of either PAK1 and/or PAK2 (Fig. 5F, Fig. S5F-G). Thus, despite slightly

impaired glucose tolerance in mice lacking PAK2 in skeletal muscle, neither adverse effects on

plasma insulin nor defects in insulin sensitivity could be detected.

Insulin-stimulated glucose uptake relies partially on PAK2 in EDL, but not soleus muscle, while

PAK1 is not involved. To test the relative contribution and involvement of PAK1 and PAK2 in

glucose uptake in skeletal muscle, we next investigated insulin-stimulated 2DG uptake in isolated

soleus and EDL. To our surprise, soleus muscle lacking PAK1 and PAK2 displayed normal insulin-

stimulated 2DG uptake compared to control littermates (Fig. 6A,C). In contrast, lack of PAK2 in

EDL muscle caused a modest reduction (PAK2 mKO: -18%; d1/2KO: -12%) in insulin-stimulated

2DG uptake (Fig. 6B,D). Thus, in oxidative soleus muscle, group I PAKs are dispensable for

normal insulin-stimulated glucose uptake, whereas in glycolytic EDL muscle PAK2 plays a minor

role.

PAK2 regulates TBC1D4 protein expression and signalling. Lastly, we looked into the effects of

PAK1 and PAK2 on insulin-stimulated signalling. All groups displayed normal basal and insulin-

stimulated pAkt S473 (Fig. 7A-B, Fig. S6A-B) and pAkt T308 (Fig. 7C-D, Fig. S6C-D) and Akt2

protein expression (Fig. S6E-F) compared to control littermates in both soleus and EDL muscle. In

contrast, lack of PAK2 increased protein expression of Akt's downstream target, TBC1D4 in soleus

muscle (PAK2 mKO: +47%; d1/2KO: +20%) (Fig. 7E), while reducing TBC1D4 expression in

EDL (PAK2 mKO: -33%; d1/2KO: -9%) (Fig. 7F). In soleus, basal and insulin-stimulated

pTBC1D4 T642 was similar in all groups (Fig. 7G, Fig. S6G), suggesting that even with increased

TBC1D4 expression activation through this pathway was normal. Concomitant with the decreased

TBC1D4 expression in EDL muscle, lack of PAK2 reduced insulin-stimulated pTBC1D4 T642

(Fig. 7H, Fig. S6H) driven by attenuated (-46%) insulin-stimulated pTBC1D4 T642 in PAK2 mKO

mice compared to control littermates (Fig. S6H). Knockout of TBC1D4 has been associated with

lower GLUT4 protein abundance in some muscles [35,36]. Whereas GLUT4 protein expression was

normal in soleus (Fig. 7I), GLUT4 protein expression was mildly reduced in EDL in PAK2 mKO

mice compared to littermate control (Fig. 7J). Thus, the slightly reduced insulin-stimulated glucose

uptake in EDL muscle lacking PAK2 was concomitant with downregulated TBC1D4 signalling and

GLUT4 expression supporting a role for PAK2, but not PAK1, in insulin-stimulated glucose

uptake.

Discussion

We here undertook a systematic investigation into the requirement of the group I PAK isoforms in

muscle glucose uptake and whole-body metabolic regulation. In contrast to previous literature, our

results show that PAK1 is dispensable for insulin-stimulated glucose uptake in skeletal muscle,

while PAK2 may play a minor role. Using a cohort of whole-body PAK1 KO mice and another

cohort of transgenic mice lacking either PAK1 (whole-body), PAK2 (muscle-specific), or jointly

lacking both PAK1 and muscle PAK2, we show that PAK1 is not required in insulin-stimulated

muscle glucose uptake *in vivo* or in isolated muscles. In accordance, we found no effect of whole-

body PAK1 KO on glucose tolerance in either mice fed a standard chow diet (insulin sensitive

mice) or in mice fed a HFD (insulin-resistant mice). Moreover, In contrast, PAK2 seemed partially

involved in insulin-stimulated glucose uptake in EDL muscle. This could potentially explain the

slightly impaired glucose tolerance with the muscle-specific knockout of PAK2 in mice.

Nevertheless, supporting only a minor role for skeletal muscle PAK2 in the whole-body substrate

utilization, RER was largely unaffected by lack of PAK1 and/or PAK2 and only slightly elevated in

mice lacking PAK2 when challenged by fasting.

In a previous study, the increase in GLUT4 abundance at the plasma membrane in response to

insulin measured was completely abrogated in PAK1 KO muscle as measured by subcellular

fractionation of homogenates of hindlimb skeletal muscles [7], suggesting that PAK1 is necessary

for insulin-stimulated GLUT4 translocation. Although glucose uptake was not assessed in that study

[7], this indicated a key role for PAK1 in regulating glucose uptake in mouse skeletal muscle.

Surprisingly, PAK1 was not required for insulin-stimulated glucose uptake in our hands.

Furthermore, in our study, both chow- and HFD-fed PAK1 KO mice displayed blood glucose

concentrations similar to control littermates during a GTT. This was in contrast to previous studies

reporting impaired glucose tolerance in chow-fed PAK1 KO mice [7,24] and elevated fasting blood

glucose in PAK1 KO mice fed a western diet (45E% fat) [33]. Instead, despite the previous finding

that insulin-stimulated GLUT4 translocation was unaffected by a 75% knockdown of PAK2 in L6-

GLUT4myc myoblasts [6], we found that muscle-specific PAK2 KO slightly impairs glucose

tolerance and insulin-stimulated glucose in mouse skeletal muscle. These discrepancies between our

and previous findings are difficult to delineate but might be due to methodological differences. For

example, Wang et al. [7] used a crude fractionation method to measure GLUT4 translocation,

whereas we analyzed the direct outcome hereof: glucose uptake. Disagreements between the

presence of GLUT4 at the plasma membrane and glucose uptake have been reported in cell cultures

[37–39]. However, in human skeletal muscle strips, the insulin-induced increase in 3-O-

methylglucose transport correlates with the increase in cell surface GLUT4 protein content [40].

Another potential explanatory factor could be that our studies were conducted in both female and

male mice, whereas past studies in PAK1 KO mice have been conducted in 4-6 months old male

mice [7,33]. However, our data suggest no major differences between female and male mice in the

response to lack of PAK1 and/or PAK2 on the whole-body metabolic parameters measured. Instead,

the discrepancies between our and previous findings could be due to an effect of age, as our studies

were conducted in mice at different ranges of age (10-37 of weeks age at the terminal experiment).

In fact, age-dependent myopathy and development of megaconial mitochondria have been reported

in the d1/2 KO mice [34]. Regardless, even though a role for group I PAKs in age-related insulin

resistance should be further investigated, our investigation suggests that group I PAKs are

indispensable in regulating whole-body glucose homeostasis or insulin-stimulated glucose uptake in

skeletal muscle.

In the current study, pharmacological inhibition of group I PAKs inhibited muscle glucose uptake in

response to insulin. Similarly, IPA-3 previously inhibited insulin-stimulated GLUT4 translocation

and glucose uptake in L6-GLUT4myc myoblasts and myotubes, respectively [6]. However, these

inhibitory effects on glucose uptake might be unspecific as they were not recapitulated in mice

lacking PAK1, and only modest effects were found in mice lacking PAK2 in the present study. IPA-

3 is a non-ATP-competitive allosteric inhibitor of all group I PAKs (PAK1, 2, and 3). IPA-3 binds

covalently to the regulatory CRIB domain of group I PAKs, thereby preventing binding to PAK

activators, such as Rac1 [31]. Although IPA-3 is reported to be a highly selective and well-

described inhibitor of group I PAKs that does not affect other groups of PAKs or similar kinases

tested [31], off-target effects of pharmacological inhibitors are a concern. It is also possible that

acute IPA-3-induced inhibition of group I PAKs elicits more potent effects compared with jointly

knockout of PAK1 and PAK2 because the transgenic manipulations have been present from birth

and may thus have resulted in compensatory changes. The development of inducible muscle-

specific group I PAK deficient models could help clarify this. Importantly, any possible

compensatory mechanisms cannot be via redundancy with group I PAKs, as PAK1 and PAK2 are

removed genetically, and even in d1/2 KO mice, PAK3 cannot be detected at the protein level [8].

This emphasizes that group I PAKs are largely dispensable for insulin-stimulated glucose uptake in

skeletal muscle.

Our hypothesis was that group I PAKs would be significantly involved in insulin-stimulated

glucose uptake because of the established necessity of their upstream activator, Rac1 [12–17]. Our

findings suggest that Rac1 does not exclusively mediate insulin-stimulated glucose uptake through

group I PAKs. Another downstream target of Rac1 is RalA. GLUT4 translocation induced by a

constitutively activated Rac1 mutant was abrogated in L6-GLUT4myc myoblasts upon RalA

knockdown [41] and, importantly, overexpression of a dominant-negative mutant of RalA reduced

GLUT4 translocation in response to insulin mouse gastrocnemius muscle fibres [42]. Additionally,

Rac1 is also an essential component in the activation of the NADPH oxidase (NOX) complex

[43,44]. In L6-GLUT4myc myotubes, reactive oxygen species have been reported to induce NOX2-

dependent GLUT4 translocation in response to insulin [45]. Insulin-stimulated NOX2 regulation in

mature muscle remains to be investigated. However, a recent study suggested a role for Rac1 in the

regulation of muscle glucose uptake through activation of the NOX2 in response to exercise [46].

Since Rac1 is required for both contraction- and insulin-stimulated glucose uptake in isolated

mouse muscle [15,47], Rac1 could also be involved in insulin-stimulated glucose uptake via NOX2

activation. Consequently, future studies should aim to investigate other players downstream of Rac1

since group I PAKs seem to be largely dispensable for glucose uptake in mature skeletal muscle.

Based on our present findings, we conclude that even though PAK2 may be a minor requirement for

insulin-stimulated glucose uptake in EDL muscle, group I PAKs are largely dispensable in the

regulation of whole-body glucose homeostasis and insulin-stimulated glucose uptake in mouse

skeletal muscle.

Materials and Methods

Animal experiments. Female C57BL/6J mice (Taconic, Denmark) were used for all inhibitor

incubation studies. Mice received standard rodent chow diet (Altromin no. 1324; Brogaarden,

Denmark) and water ad libitum.

Whole-body PAK1$^{-/-}$ mice. Whole-body PAK1$^{-/-}$ mice on a C57BL/6J background were generated as

previously described [48]. The mice were obtained by heterozygous crossing. PAK1$^{-/-}$ mice

(referred to as PAK1 KO) and paired littermate PAK1$^{+/+}$ mice (referred to as controls) were used for

experiments. Female and male mice were used for measurements of body composition, glucose

tolerance and insulin-stimulated glucose uptake in isolated muscle. The mice were 12-24 weeks of

age at the time of tissue dissection and measurement of glucose uptake. Number of mice in each

group: Control, *n = 6/7* (female/male); PAK1 KO, *n = 4/8*. Mice received standard rodent chow

diet and water ad libitum.

For measurement of *in vivo* insulin-stimulated glucose uptake in chow- and 60E% HFD (no.

D12492; Brogaarden, Denmark)-fed PAK1 KO mice, mice were assigned to a chow or HFD group.

Chow-fed mice were 10-24 weeks of age at the time of glucose uptake measurements and tissue

dissection. Number of mice in each group: Control-Chow, $n = 14/8$ (female/male); PAK1 KO-

Chow, $n = 6/4$. For mice receiving HFD, the diet intervention started at 6-16 weeks of age and

lasted for 21 weeks. HFD-fed mice were used for body composition, glucose tolerance and *in vivo*

glucose uptake and were 27-37 weeks of age at the time of glucose uptake measurements and tissue

dissection. Number of mice in each group: Control-HFD, $n = 7/7$ (female/male); PAK1 KO-HFD, n

$= 11/5$. Energy intake was measured over a period of 10 weeks in another cohort of mice. Number

of mice in each group: Chow, $n = 8/8$ (Control/PAK1 KO); HFD, $n = 8/8$. Mice had access to their

respective diet and water ad libitum.

Double PAK1$^{-/-}$;PAK2$^{fl/fl}$;MyoD$^{iCre/+}$ mice. Double knockout mice with whole-body knockout of

PAK1 and conditional, muscle-specific knockout of PAK2, PAK1$^{-/-}$;PAK2$^{fl/fl}$;MyoD$^{iCre/+}$ were

generated as previously described [8]. The mice were on a mixed C57BL/6/FVB background.

PAK1$^{-/-}$;PAK2$^{fl/fl}$;MyoD$^{iCre/+}$ were crossed with PAK1$^{+/-}$;PAK2$^{fl/fl}$;MyoD$^{+/+}$ to generate littermate

PAK1$^{-/-}$;PAK2$^{fl/fl}$;MyoD$^{iCre/+}$ (referred to as d1/2 KO), PAK1$^{-/-}$;PAK2$^{fl/fl}$;MyoD$^{+/+}$ (referred to as

PAK1 KO), PAK1$^{+/-}$;PAK2$^{fl/fl}$;MyoD$^{iCre/+}$ (referred to as PAK2 mKO), and PAK1$^{+/-}$

;PAK2$^{fl/fl}$;MyoD$^{+/+}$ (referred to as controls) used for experiments. Female and male mice were used

for measurement of insulin-stimulated glucose uptake in isolated muscle. The mice were 10-16

weeks of age at the time of tissue dissection and glucose uptake measurements. Number of mice in

each group: Control, $n = 6/4$ (female/male); PAK1 KO, $n = 5/4$, PAK2 mKO, $n = 6/4$, d1/2KO, $n =$

$6/3$. Another cohort of mice was used for whole-body metabolic measurements. The first

measurement (insulin tolerance) was at 11-24 weeks of age and the last measurement was at 23-33

weeks of age (home cage calorimetry). Number of mice in each group: Control, $n = 9/11$

(female/male); PAK1 KO, $n = 8/10$, PAK2 mKO, $n = 12/9$, d1/2KO, $n = 9/14$. For some of the

metabolic measurements, only a subgroup of mice was used as indicated in the relevant figure legends. Mice received standard rodent chow diet and water ad libitum.

All animals were maintained on a 12:12-hour light-dark cycle and housed at 22°C (with allowed fluctuation of ±2°C) with nesting material. The mice were group-housed. All animal experiments complied with the European Convention for the protection of vertebrate animals used for experimental and other scientific purposes (No. 123, Strasbourg, France, 1985) and were approved by the Danish Animal Experimental Inspectorate.

Body composition. Body composition was analyzed using magnetic resonance imaging (EchoMRI-4in1TM, Echo Medical System LLC, Texas, USA). Chow-fed PAK1 KO and control littermates were assessed at 7-19 weeks of age. HFD-fed PAK1 KO and control littermates were assessed 18-19 weeks into the diet intervention (24-34 weeks of age). Chow-fed PAK1 KO, PAK2 mKO, d1/2 KO mice and control littermates were assessed at 16-29 weeks of age.

Glucose tolerance test (GTT). Prior to the test, chow- and HFD-fed PAK1 KO mice and control littermates fasted for 12 hours from 10 p.m, while chow-fed PAK1 KO, PAK2 mKO, d1/2 KO mice and control littermates fasted for 6 hours from 6 a.m. D-mono-glucose (2 g kg^{-1} body weight) was administered intraperitoneal (i.p) and blood was collected from the tail vein and blood glucose concentration determined at the indicated time points using a glucometer (Bayer Contour, Bayer, Switzerland). Incremental Area Under the Curve (AUC) from the basal blood glucose concentration was determined using the trapezoid rule. For measurement of plasma insulin, glucose was administered i.p. on a separate experimental day (1-2 weeks after the GTT) and blood was sampled at time points 0 and 20 minutes, centrifuged and plasma was quickly frozen in liquid nitrogen and stored at -20°C until processing. Plasma insulin was analyzed in duplicate (Mouse Ultrasensitive Insulin ELISA, #80-INSTRU-E10, ALPCO Diagnostics, USA). Homeostatic model assessment of

insulin resistance (HOMA-IR) was calculated according to the formula: Fasting plasma insulin (mU L^{-1}) X Fasting blood glucose (mM)/22.5. Glucose tolerance was assessed in 8-20 weeks of age chow-fed PAK1 KO mice and in week 14 of the diet intervention of HFD-fed PAK1 KO mice (20-30 weeks of age). In chow-fed PAK1 KO, PAK2 mKO, d1/2 KO mice and control littermates, glucose tolerance was assessed at 13-26 weeks of age.

Insulin tolerance test (ITT). Prior to the test, chow-fed PAK1 KO, PAK2 mKO, d1/2 KO mice and control littermates fasted for 4 hours from 6 a.m. Insulin (0.5 U kg^{-1} body weight) was administered i.p. and blood was collected from the tail vein and blood glucose concentration determined using a glucometer (Bayer Contour, Bayer, Switzerland) at time point 0, 15, 30, 60, 90 and 120 minutes. For two female control mice and four female PAK2 mKO mice, the ITT had to be stopped before the 120'-time point due to hypoglycemia (blood glucose <1.2 mM). Thus, blood glucose was not measured in these mice for the last couple of time points. Insulin tolerance was assessed in 11-24 weeks of age chow-fed PAK1 KO, PAK2 mKO, d1/2 KO mice and control littermates.

Home cage indirect calorimetry. One week prior to the calorimetric measurements, chow-fed PAK1 KO, PAK2 mKO, d1/2 KO mice and control littermates were single-housed in specialized cages for indirect gas calorimetry but uncoupled from the PhenoMaster indirect calorimetry system (TSE PhenoMaster metabolic cage systems, TSE Systems, Germany). After a 2-day acclimation period coupled to the PhenoMaster indirect calorimetry system, oxygen consumption, CO_2 production, habitual activity (beam breaks) and food intake were measured for 72 hours (TSE LabMaster V5.5.3, TSE Systems, Germany). On day 2, mice fasted during the dark period followed by refeeding on day 3. Respiratory exchange ratio (RER) was calculated as the ratio between CO_2 production and oxygen consumption. The mice were 23-33 weeks of age.

Incubation of isolated muscles. Soleus and EDL muscles were dissected from anaesthetized mice

(6 mg pentobarbital sodium 100 g^{-1} body weight i.p.) and suspended at resting tension (4-5 mN) in

incubations chambers (Multi Myograph System, Danish Myo Technology, Denmark) in Krebs-

Ringer-Henseleit buffer with 2 mM pyruvate and 8 mM mannitol at 30°C, as described previously

[49]. Additionally, the Krebs-Ringer-Henseleit buffer was supplemented with 0.1% BSA (v/v).

Isolated muscles from female C57BL/6J mice were pre-incubated with 40 μM IPA-3 (Sigma-

Aldrich) or as a control DMSO (0.25%) for 25 minutes followed by 30 minutes of insulin

stimulation (60 nM; Actrapid, Novo Nordisk, Denmark). Isolated muscles from chow-PAK1 KO

were pre-incubated for 30 minutes followed by 30 minutes of insulin stimulation (0.6 nm or 60

nM). Isolated muscles from chow-fed PAK1 KO, PAK2 mKO, d1/2 KO mice or control littermates

were pre-incubated for 20 minutes followed by 20 minutes of insulin stimulation (60 nM). 2DG

uptake was measured together with 1 mM 2DG during the last 10 min of the insulin stimulation

period using 0.6 μCi mL^{-1} [^{3}H]-2DG and 0.180 μCi mL^{-1} [^{14}C]-mannitol radioactive tracers as

described previously [49]. Tissue-specific [^{3}H]-2DG accumulation with [^{14}C]-mannitol as an

extracelllular marker was determined as previously described [50].

In vivo insulin-stimulated 2-Deoxyglucose uptake in PAK1 KO mice during a r.o. ITT. To

determine 2DG uptake in skeletal muscle of PAK1 KO mice and littermate controls, [^{3}H]-2DG

(Perkin Elmer) was administered retro-orbitally (r.o.) in a bolus of saline containing 66.7 μCi mL^{-1}

[^{3}H]-2DG (~32.4 Ci/mmol) corresponding to ~10 μCi/mouse in chow-fed mice or ~15 μCi/mouse

in HFD-fed mice (6 μL g^{-1} body weight) as described [17]. The [^{3}H]-2DG saline bolus was with or

without insulin (Actrapid, Novo Nordisk, Denmark). Decreased insulin clearance has been observed

by us [17] and others in obese rodent [51,52] and human [53] models. Therefore, to correct for

changes in insulin clearance, 0.5 U kg^{-1} body weight of insulin was administered in chow-fed mice

whereas only 60% of this dosage was administered to HFD-fed mice. Prior to stimulation, mice

fasted for 4 hours from 07:00 and were anaesthetized (7.5/9 mg [Chow/HFD] pentobarbital sodium

100 g^{-1} body weight i.p.) 15 minutes before the r.o. injection. Blood samples were collected from

the tail vein after the r.o. injection and analyzed for glucose concentration using a glucometer

(Bayer Contour, Bayer, Switzerland) at the time points 0, 5 and 10 minutes. After 10 minutes,

skeletal muscle (gastrocnemius, quadriceps, triceps brachii and soleus) were excised, rinsed in

saline, and quickly frozen in liquid nitrogen and stored at -80°C until processing. Blood was

collected by punctuation of the heart, centrifuged and plasma was quickly frozen in liquid nitrogen

and stored at -80°C until processing. Plasma samples were analyzed for insulin concentration and

specific [^{3}H]-2DG activity. Plasma insulin was analyzed in duplicate (Mouse Ultrasensitive Insulin

ELISA, #80-INSTRU-E10, ALPCO Diagnostics, USA). Tissue-specific 2DG-6-phosphate

accumulation was measured as described [54,55]. To determine 2DG clearance from the plasma

into the specific tissue, tissue-specific [^{3}H]-2DG-6-P was divided by AUC of the plasma-specific

[^{3}H]-2DG activity at the time points 0 and 10 minutes. To estimate tissue-specific glucose uptake

(glucose uptake index), clearance was multiplied by the average blood glucose levels for the time

points 0, 5, and 10 minutes. Tissue-specific 2DG clearance and glucose uptake were related to

analyzed muscle tissue weight and time.

Muscle analyses. Prior to homogenization, gastrocnemius, quadriceps, and triceps brachii muscles

were pulverized in liquid nitrogen. All muscle were homogenized 2 x 30 sec at 30 Hz using a

Tissuelyser II (Qiagen, USA) in ice-cold homogenization buffer (10% (v/v) Glycerol, 1% (v/v) NP-

40, 20 mM Na-pyrophosphate, 150 mM NaCl, 50 mM HEPES (pH 7.5), 20 mM β-

glycerophosphate, 10 mM NaF, 2mM PMSF, 1 mM EDTA (pH 8.0), 1 mM EGTA (pH 8.0), 2 mM

Na3VO4, 10 µg mL^{-1} Leupeptin, 10 µg mL^{-1} Aprotinin, 3 mM Benzamidine). After rotation end-

over-end for 30 min at 4°C, lysate supernatants were collected by centrifugation (10,854-15,630 x

g) for 15-20 min at 4°C.

Immunoblotting. Lysate protein concentration was determined using the bicinchoninic acid (BCA)

method using BSA standards and BCA assay reagents (Pierce). Immunoblotting samples were

prepared in 6x sample buffer (340 mM Tris (pH 6.8), 225 mM DTT, 11% (w/v) SDS, 20% (v/v)

Glycerol, 0.05% (w/v) Bromphenol blue). Protein phosphorylation (p) and total protein expression

were determined by standard immunoblotting technique loading equal amounts of protein. The

polyvinylidene difluoride membrane (Immobilon Transfer Membrane; Millipore) was blocked in

Tris-Buffered Saline with added Tween20 (TBST) and 2% (w/v) skim milk or 5% (w/v) BSA

protein for 15 minutes at room temperature, followed by incubation overnight at 4°C with a primary

antibody (Table 1). Next, the membrane was incubated with horseradish peroxidase-conjugated

secondary antibody (Jackson Immuno Research) for 1 hour at room temperature. Bands were

visualized using Bio-Rad ChemiDocTM MP Imaging System and enhanced chemiluminescence

(ECL+; Amersham Biosciences). Densitometric analysis was performed using Image LabTM

Software, version 4.0 (Bio-Rad, USA). Coomassie brilliant blue staining was used as a loading

control [56].

Statistical analyses. Data are presented as mean ± S.E.M. or when applicable mean ± S.E.M. with

individual data points shown. Statistical tests varied according to the dataset being analyzed and the

specific tests used are indicated in the figure legends. If the null hypothesis was rejected, Tukey's

post hoc test was used to evaluate significant main effects of genotype and significant interactions

in ANOVAs. P < 0.05 was considered statistically significant. P<0.1 was considered a tendency.

Except for mixed-effects model analyses performed in GraphPad Prism, version 8.2.1. (GraphPad

Software, La Jolla, CA, USA), all statistical analyses were performed using Sigma Plot, version 13

(Systat Software Inc., Chicago, IL, USA). Due to missing values ascribed to hypoglycemia,

differences between genotypes and the effect of insulin administration were assessed with a mixed-

effects model analysis in Fig. 5F and S5F.

Disclosure summary

No potential conflicts of interest relevant to this article were reported.

Acknowledgements

We thank our colleagues, especially Jørgen Wojtaszewski and Bente Kiens, at the Section of Molecular Physiology, Department of Nutrition, Exercise, and Sports (NEXS), Faculty of Science, University of Copenhagen, for fruitful discussions on this topic. We acknowledge the skilled technical assistance of Betina Bolmgren, Irene B. Nielsen, and Mona Ali (Section of Molecular Physiology, NEXS, Faculty of Science, University of Copenhagen, Denmark). The PAK1 KO mice were a kind gift from Debbie Thurmond (Department of Molecular & Cellular Endocrinology, Diabetes and Metabolism Research Institute, City of Hope/BRICA, USA).

Author contributions

LLVM, EAR, and LS conceptualized and designed the study. LLVM and LS conducted the experiments and analyzed the data, with experimental contribution and/or interpretation of data from MJ, RK, GAJ, ABM, JRK, A-ML, NRA, PS, TEJ, RSK. LLVM, EAR, and LS drafted the manuscript. All authors contributed to the final version of the manuscript. EAR and LS are the guaranteers of this work as, as such, have full access to all the data and take responsibility for the integrity of the data and the accuracy of the data analyses.

Grants

This study was supported by a PhD fellowship from The Lundbeck Foundation (grant 2015-3388 to LLVM); PhD scholarships from The Danish Diabetes Academy, funded by The Novo Nordisk Foundation (ABM and JRK); Postdoctoral research grant from the Danish Diabetes Academy, funded by the Novo Nordisk Foundation (grant NNF17SA0031406 to A-ML); National Institute of Arthritis and Musculoskeletal and Skin Diseases (grant AR046207 and AR070231 to RSK); The Danish Council for Independent Research, Medical Sciences (grant DFF-4004-00233 to LS, grant 6108-00203 to EAR); The Novo Nordisk Foundation (grant 10429 to EAR, grant 15182 to TEJ, grant NNF16OC0023418 and NNF18OC0032082 to LS).

References

1. DeFronzo RA, Gunnarsson R, Björkman O, Olsson M, Wahren J. Effects of insulin on peripheral and splanchnic glucose metabolism in noninsulin-dependent (type II) diabetes mellitus. J Clin Invest. 1985;76: 149–155. doi:10.1172/JCI111938

2. Baron AD, Brechtel G, Wallace P, Edelman S V. Rates and tissue sites of non-insulin- and insulin-mediated glucose uptake in humans. Am J Physiol Metab. 1988;255: E769–E774. doi:10.1152/ajpendo.1988.255.6.E769

3. Samuel VT, Petersen KF, Shulman GI. Lipid-induced insulin resistance: unravelling the mechanism. Lancet. England; 2010;375: 2267–2277. doi:10.1016/S0140-6736(10)60408-4

4. Lauritzen HPMM. Insulin- and Contraction-Induced Glucose Transporter 4 Traffic in Muscle. Exerc Sport Sci Rev. United States; 2013;41: 77–86. doi:10.1097/JES.0b013e318275574c

5. Tsakiridis T, Taha C, Grinsteinl S, Klip A. Insulin activates a p21-activated kinase in muscle cells via phosphatidylinositol 3-kinase. J Biol Chem. American Society for Biochemistry and Molecular Biology; 1996;271: 19664–19667. doi:10.1074/jbc.271.33.19664

6. Tunduguru R, Chiu TT, Ramalingam L, Elmendorf JS, Klip A, Thurmond DC. Signaling of the p21-activated kinase (PAK1) coordinates insulin-stimulated actin remodeling and glucose uptake in skeletal muscle cells. Biochem Pharmacol. NIH Public Access; 2014;92: 380–388. doi:10.1016/j.bcp.2014.08.033

7. Wang Z, Oh E, Clapp DW, Chernoff J, Thurmond DC. Inhibition or ablation of p21-activated kinase (PAK1) disrupts glucose homeostatic mechanisms in vivo. J Biol Chem. American Society for Biochemistry and Molecular Biology; 2011;286: 41359–41367. doi:10.1074/jbc.M111.291500

8. Joseph GA, Lu M, Radu M, Lee JK, Burden SJ, Chernoff J, et al. Group I Paks Promote Skeletal

Myoblast Differentiation In Vivo and In Vitro. Mol Cell Biol. American Society for Microbiology
(ASM); 2017;37: e00222-16. doi:10.1128/MCB.00222-16

9.   Chiang YA, Jin T. p21-Activated protein kinases and their emerging roles in glucose homeostasis.
Am J Physiol Metab. United States; 2014;306: E707–E722. doi:10.1152/ajpendo.00506.2013

10. Manser E, Leung T, Salihuddin H, Zhao Z, Lim L. A brain serine/threonine protein kinase activated
by Cdc42 and Rac1. Nature. Nature Publishing Group; 1994;367: 40–46. doi:10.1038/367040a0

11. Arvidson AR. A tale of two parks. Landsc Archit. 2008;98: 108–117. doi:10.1042/bc20070109

12. Khayat ZA, Tong P, Yaworsky K, Bloch RJ, Klip A. Insulin-induced actin filament remodeling
colocalizes actin with phosphatidylinositol 3-kinase and GLUT4 in L6 myotubes. J Cell Sci.
2000;113 Pt 2: 279–90. Available: http://www.ncbi.nlm.nih.gov/pubmed/10633079

13. JeBailey L, Wanono O, Niu W, Roessler J, Rudich A, Klip A. Ceramide- and oxidant-induced insulin
resistance involve loss of insulin-dependent Rac-activation and actin remodeling in muscle cells.
Diabetes. American Diabetes Association; 2007;56: 394–403. doi:10.2337/db06-0823

14. Ueda S, Kitazawa S, Ishida K, Nishikawa Y, Matsui M, Matsumoto H, et al. Crucial role of the small
GTPase Rac1 in insulin-stimulated translocation of glucose transporter 4 to the mouse skeletal muscle
sarcolemma. FASEB J. 2010;24: 2254–2261. doi:10.1096/fj.09-137380

15. Sylow L, Jensen TE, Kleinert M, Hojlund K, Kiens B, Wojtaszewski J, et al. Rac1 Signaling Is
Required for Insulin-Stimulated Glucose Uptake and Is Dysregulated in Insulin-Resistant Murine and
Human Skeletal Muscle. Diabetes. 2013;62: 1865–1875. doi:10.2337/db12-1148

16. Sylow L, Kleinert M, Pehmøller C, Prats C, Chiu TT, Klip A, et al. Akt and Rac1 signaling are jointly
required for insulin-stimulated glucose uptake in skeletal muscle and downregulated in insulin
resistance. Cell Signal. Elsevier Inc.; 2014;26: 323–331. doi:10.1016/j.cellsig.2013.11.007

17. Raun SH, Ali M, Kjøbsted R, Møller LL V, Federspiel MA, Richter EA, et al. Rac1 muscle knockout
exacerbates the detrimental effect of high-fat diet on insulin-stimulated muscle glucose uptake
independently of Akt. J Physiol. 2018;596: 2283–2299. doi:10.1113/JP275602

18. Sano H, Kane S, Sano E, Mîinea CP, Asara JM, Lane WS, et al. Insulin-stimulated Phosphorylation
of a Rab GTPase-activating Protein Regulates GLUT4 Translocation. J Biol Chem. United States;
2003;278: 14599–14602. doi:10.1074/jbc.C300063200

19. Thong FSL, Dugani CB, Klip A. Turning signals on and off: GLUT4 traffic in the insulin-signaling
highway. Physiology. American Physiological Society; 2005. pp. 271–284.
doi:10.1152/physiol.00017.2005

20. Kramer HF, Witczak CA, Taylor EB, Fujii N, Hirshman MF, Goodyear LJ. AS160 Regulates Insulin-
and Contraction-stimulated Glucose Uptake in Mouse Skeletal Muscle. J Biol Chem. 2006;281:
31478–31485. doi:10.1074/jbc.M605461200

21. Benner GE, Dennis PB, Masaracchia RA. Activation of an S6/H4 Kinase (PAK 65) from Human
Placenta by Intramolecular and Intermolecular Autophosphorylation. J Biol Chem. American Society
for Biochemistry and Molecular Biology; 1995;270: 21121–21128. doi:10.1074/jbc.270.36.21121

22. Gatti A, Huang Z, Tuazon PT, Traugh JA. Multisite autophosphorylation of p21-activated protein
kinase gamma-PAK as a function of activation. J Biol Chem. American Society for Biochemistry and
Molecular Biology; 1999;274: 8022–8. doi:10.1074/jbc.274.12.8022

23. Stierwalt HD, Ehrlicher SE, Bergman BC, Robinson MM, Newsom SA. Insulin-stimulated Rac1-
GTP binding is not impaired by palmitate treatment in L6 myotubes. Physiol Rep. United States;
2018;6: e13956. doi:10.14814/phy2.13956

24. Chiang YA, Shao W, Xu XX, Chernoff J, Jin T. P21-activated protein kinase 1 (Pak1) mediates the
cross talk between insulin and β-catenin on proglucagon gene expression and its ablation affects

glucose homeostasis in male C57BL/6 mice. Endocrinology. 2013;154: 77–88. doi:10.1210/en.2012-
1781

25.   Chiu TT, Patel N, Shaw AE, Bamburg JR, Klip A. Arp2/3- and Cofilin-coordinated Actin Dynamics
Is Required for Insulin-mediated GLUT4 Translocation to the Surface of Muscle Cells. Blanchoin L,
editor. Mol Biol Cell. 2010;21: 3529–3539. doi:10.1091/mbc.e10-04-0316

26.   Tunduguru R, Zhang J, Aslamy A, Salunkhe VA, Brozinick JT, Elmendorf JS, et al. The actin-related
p41ARC subunit contributes to p21-activated kinase-1 (PAK1)-mediated glucose uptake into skeletal
muscle cells. J Biol Chem. American Society for Biochemistry and Molecular Biology Inc.;
2017;292: 19034–19043. doi:10.1074/jbc.M117.801340

27.   Wasserman DH, Kang L, Ayala JE, Fueger PT, Lee-Young RS. The physiological regulation of
glucose flux into muscle in vivo. J Exp Biol. 2011;214: 254–262. doi:10.1242/jeb.048041

28.   Hansen PA, Gulve EA, Marshall BA, Gao J, Pessin JE, Holloszy JO, et al. Skeletal muscle glucose
transport and metabolism are enhanced in transgenic mice overexpressing the Glut4 glucose
transporter. J Biol Chem. 1995;270: 1679–84. doi:10.1074/jbc.270.5.1679

29.   Hansen PA, Marshall BA, Chen M, Holloszy JO, Mueckler M. Transgenic Overexpression of
Hexokinase II in Skeletal Muscle Does Not Increase Glucose Disposal in Wild-type or Glut1-
overexpressing Mice. J Biol Chem. JBC Papers in Press; 2000;275: 22381–22386.
doi:10.1074/jbc.M001946200

30.   Zisman A, Peroni OD, Abel ED, Michael MD, Mauvais-Jarvis F, Lowell BB, et al. Targeted
disruption of the glucose transporter 4 selectively in muscle causes insulin resistance and glucose
intolerance. Nat Med. 2000;6: 924–928. doi:10.1038/78693

31.   Deacon SW, Beeser A, Fukui JA, Rennefahrt UEE, Myers C, Chernoff J, et al. An Isoform-Selective,
Small-Molecule Inhibitor Targets the Autoregulatory Mechanism of p21-Activated Kinase. Chem
Biol. 2008;15: 322–331. doi:10.1016/j.chembiol.2008.03.005

32.   Davies SP, Reddy H, Caivano M, Cohen P. Specificity and mechanism of action of some commonly
used protein kinase inhibitors. Biochem J. 2000;351: 95–105. doi:10.1517/13543776.11.3.405

33.   Ahn M, Yoder SM, Wang Z, Oh E, Ramalingam L, Tunduguru R, et al. The p21-activated kinase
(PAK1) is involved in diet-induced beta cell mass expansion and survival in mice and human islets.
Diabetologia. NIH Public Access; 2016;59: 2145–2155. doi:10.1007/s00125-016-4042-0

34.   Joseph GA, Hung M, Goel AJ, Hong M, Rieder M-K, Beckmann ND, et al. Late-onset megaconial
myopathy in mice lacking group I Paks. Skelet Muscle. BioMed Central; 2019;9: 5.
doi:10.1186/s13395-019-0191-4

35.   Lansey MN, Walker NN, Hargett SR, Stevens JR, Keller SR. Deletion of Rab GAP AS160 modifies
glucose uptake and GLUT4 translocation in primary skeletal muscles and adipocytes and impairs
glucose homeostasis. Am J Physiol Metab. 2012;303: E1273–E1286.
doi:10.1152/ajpendo.00316.2012

36.   Wang HY, Ducommun S, Quan C, Xie B, Li M, Wasserman DH, et al. AS160 deficiency causes
whole-body insulin resistance via composite effects in multiple tissues. Biochem J. 2013;449: 479–
489. doi:10.1042/BJ20120702

37.   Somwar R, Niu W, Kim DY, Sweeney G, Randhawa VK, Huang C, et al. Differential Effects of
Phosphatidylinositol 3-Kinase Inhibition on Intracellular Signals Regulating GLUT4 Translocation
and Glucose Transport. J Biol Chem. United States; 2001;276: 46079–46087.
doi:10.1074/jbc.M109093200

38.   Somwar R, Kim DY, Sweeney G, Huang C, Niu W, Lador C, et al. GLUT4 translocation precedes the
stimulation of glucose uptake by insulin in muscle cells: potential activation of GLUT4 via p38
mitogen-activated protein kinase. Biochem J. England; 2001;359: 639–49. doi:10.1042/0264-

6021:3590639

39.   Funaki M, Randhawa P, Janmey PA. Separation of Insulin Signaling into Distinct GLUT4 Translocation and Activation Steps. Mol Cell Biol. American Society for Microbiology; 2004;24: 7567–7577. doi:10.1128/mcb.24.17.7567-7577.2004

40.   Lund S, Holman GD, Zierath JR, Rincon J, Nolte LA, Clark AE, et al. Effect of Insulin on GLUT4 CeU Surface Content and Turnover Rate in Human Skeletal Muscle as Measured by the Exofacial Bis-Mannose Photolabeling Technique. 1965.

41.   Nozaki S, Ueda S, Takenaka N, Kataoka T, Satoh T. Role of RalA downstream of Rac1 in insulin-dependent glucose uptake in muscle cells. Cell Signal. Pergamon; 2012;24: 2111–2117. doi:10.1016/j.cellsig.2012.07.013

42.   Takenaka N, Sumi Y, Matsuda K, Fujita J, Hosooka T, Noguchi T, et al. Role for RalA downstream of Rac1 in skeletal muscle insulin signalling. Biochem J. Portland Press Limited; 2015;469: 445–54. doi:10.1042/BJ20150218

43.   Bedard K, Krause K-H. The NOX Family of ROS-Generating NADPH Oxidases: Physiology and Pathophysiology. Physiol Rev. United States; 2007;87: 245–313. doi:10.1152/physrev.00044.2005

44.   Abo A, Pick E, Hall A, Totty N, Teahan CG, Segal AW. Activation of the NADPH oxidase involves the small GTP-binding protein p21rac1. Nature. 1991;353: 668–670. doi:10.1038/353668a0

45.   Contreras-Ferrat A, Llanos P, Vásquez C, Espinosa A, Osorio-Fuentealba C, Arias-Calderon M, et al. Insulin elicits a ROS-activated and an IP3-dependent Ca2+ release, which both impinge on GLUT4 translocation. J Cell Sci. Company of Biologists Ltd; 2014;127: 1911–1923. doi:10.1242/jcs.138982

46.   Henríquez-Olguin C, Knudsen JR, Raun SH, Li Z, Dalbram E, Treebak JT, et al. Cytosolic ROS production by NADPH oxidase 2 regulates muscle glucose uptake during exercise. Nat Commun. 2019;10: 4623. doi:10.1038/s41467-019-12523-9

47.   Sylow L, Jensen TE, Kleinert M, Mouatt JR, Maarbjerg SJ, Jeppesen J, et al. Rac1 Is a Novel Regulator of Contraction-Stimulated Glucose Uptake in Skeletal Muscle. Diabetes. 2013;62: 1139–1151. doi:10.2337/db12-0491

48.   Allen JD, Jaffer ZM, Park SJ, Burgin S, Hofmann C, Sells MA, et al. P21-activated kinase regulates mast cell degranulation via effects on calcium mobilization and cytoskeletal dynamics. Blood. 2009;113: 2695–2705. doi:10.1182/blood-2008-06-160861

49.   Viollet B, Andreelli F, Jørgensen SB, Perrin C, Geloen A, Flamez D, et al. The AMP-activated protein kinase α2 catalytic subunit controls whole-body insulin sensitivity. J Clin Invest. American Society for Clinical Investigation; 2003;111: 91–98. doi:10.1172/JCI16567

50.   Kjøbsted R, Treebak JT, Fentz J, Lantier L, Viollet B, Birk JB, et al. Prior AICAR stimulation increases insulin sensitivity in mouse skeletal muscle in an AMPK-dependent manner. Diabetes. 2015;64: 2042–2055. doi:10.2337/db14-1402

51.   Strömblad G, Björntorp P. Reduced hepatic insulin clearance in rats with dietary-induced obesity. Metabolism. 1986;35: 323–327. doi:10.1016/0026-0495(86)90148-4

52.   Brandimarti P, Costa-Júnior JM, Ferreira SM, Protzek AO, Santos GJ, Carneiro EM, et al. Cafeteria diet inhibits insulin clearance by reduced insulindegrading enzyme expression and mRNA splicing. J Endocrinol. 2013;219: 173–182. doi:10.1530/JOE-13-0177

53.   Jung S-H, Jung C-H, Reaven GM, Kim SH. Adapting to insulin resistance in obesity: role of insulin secretion and clearance. Diabetologia. 2018;61: 681–687. doi:10.1007/s00125-017-4511-0

54.   Ferre P, Leturque A, Burnol A-F, Penicaud L, Girard J. A method to quantify glucose utilization in vivo in skeletal muscle and white adipose tissue of the anaesthetized rat. Biochem J. 1985;228: 103–110.

681   55.   Fueger PT, Bracy DP, Malabanan CM, Pencek RR, Wasserman DH. Distributed control of glucose
682   uptake by working muscles of conscious mice: roles of transport and phosphorylation. Am J Physiol
683   Metab. 2004;286: E77–E84. doi:10.1152/ajpendo.00309.2003

684   56.   Welinder C, Ekblad L. Coomassie Staining as Loading Control in Western Blot Analysis. J Proteome
685   Res. 2011;10: 1416–1419. doi:10.1021/pr1011476

Figure Legends

Figure 1: Insulin-stimulated glucose uptake is reduced in IPA-3-treated mouse soleus muscle. (A-B) Insulin-stimulated (60 nM) 2-Deoxyglucose (2DG) uptake in isolated soleus (A) and extensor digitorum longus (EDL, B) muscle $\pm$ 40 µM IPA-3 or as a control DMSO (0.25%). Isolated muscles were pre-incubated for 25 minutes followed by 30 minutes of insulin stimulation with 2DG uptake analyzed for the final 10 minutes of stimulation. **(C-H)** Quantification of phosphorylated (p)PAK1 T423, pAkt T308, and pAkt S473 in insulin-stimulated soleus (C, E, and G) and EDL (D, F, and H) muscle $\pm$ 40 µM IPA-3 or as a control DMSO (0.25%). Some of the data points were excluded due to the quality of the immunoblot, and the number of determinations was $n = 8/7$ for pAkt S473 and T308 in soleus muscle. **(I-J)** Representative blots showing pPAK1 T423, pAkt T308, and pAkt S473 in soleus (I) and EDL (J) muscle. Statistics were evaluated with a two-way repeated measures (RM) ANOVA. Main effects are indicated in the panels. Interactions in two-way RM ANOVA were evaluated by Tukey's post hoc test: Insulin stimulation vs. basal **/*** ($p<0.01/0.001$); IPA-3 vs. DMSO (#)/#/##/### ($p<0.1/0.05/0.01/0.001$). Unless otherwise stated previously in the figure legend, the number of determinations in each group: Soleus, $n = 9/8$ (DMSO/IPA-3); EDL, $n = 9/8$. Data are presented as mean $\pm$ S.E.M. with individual data points shown. Paired data points are connected with a straight line. A.U., arbitrary units.

Figure 2: PAK1 knockout does not affect whole-body glucose homeostasis or insulin-stimulated glucose uptake in isolated skeletal muscle. (A) Representative blots showing PAK1 protein expression in gastrocnemius, quadriceps, and triceps brachii muscle from PAK1 knockout (KO) mice or control littermates. **(B)** Body composition (FM: Fat mass; LBM: Lean body mass; BW: Body weight) in gram in chow-fed PAK1 KO mice ($n = 12$) or control littermates ($n = 13$). The mice were 7-19 weeks of age. Statistics were evaluated with a Student's t-test. **(C)** Energy intake in chow-fed PAK1 KO mice ($n = 8$) or control littermates ($n = 8$). Energy intake was monitored in a separate cohort of mice. Statistics were evaluated with a Student's t-test. **(D)** Blood glucose levels during a glucose tolerance test (GTT) in chow-fed PAK1 KO mice ($n = 9$) or control littermates ($n = 10$). The mice were 8-20 weeks of age. Statistics were evaluated with a two-way repeated measures (RM) ANOVA. **(E)** Incremental Area Under the Curve (AUC) for blood glucose levels during the GTT in panel D. Statistics were evaluated with a Student's t-test. **(F)** Plasma insulin values during a GTT in chow-fed PAK1 KO mice ($n = 10$) or control littermates ($n = 13$). The mice were 9-21 weeks of age. Statistics were evaluated with a two-way RM measures ANOVA. **(G)** Homeostatic Model Assessment of Insulin Resistance (HOMA-IR) in chow-fed PAK1 KO mice ($n=10$) or control littermates ($n=13$). Statistics were evaluated

with a Student's t-test. **(H-I)** Submaximal (0.6 nM) and maximal (60 nM) insulin-stimulated 2-Deoxyglucose (2DG) uptake in isolated soleus (H) and extensor digitorum longus (EDL; I) muscle from PAK1 KO mice or littermate controls. Isolated muscles were pre-incubated for 30 minutes followed by 30 minutes of insulin stimulation with 2DG uptake analyzed for the final 10 minutes of stimulation. The mice were 12-24 weeks of age. The number of determinations in each group: Soleus-Control, *n = 6/7* (Submax/max); Soleus-PAK1 KO, *n = 7/7*; EDL-Control, *n = 6/6*; EDL-PAK1 KO, *n = 7/7*. Statistics were evaluated with two two-way RM measures ANOVA. Main effects are indicated in the panels. Data are presented as mean $\pm$ S.E.M. or when applicable mean $\pm$ S.E.M. with individual data points shown. Paired data points are connected with a straight line.

Figure 3: PAK1 is dispensable for *in vivo* insulin-stimulated glucose uptake in mouse skeletal muscle. (A-B) Blood glucose levels during a retro-orbital insulin tolerance test (r.o. ITT) in chow- (A) and HFD-fed (B) PAK1 knockout (KO) mice or control littermates. Chow-fed mice were investigated at 10-24 weeks of age. Mice fed a 60E% high-fat diet (HFD) for 21 weeks were investigated at 27-37 weeks of age. The number of mice in each group: Chow, *n = 12/6* (Control/PAK1 KO); HFD, *n = 10/11*. Statistics were evaluated with a two-way repeated measures ANOVA. **(C-D)** Insulin-stimulated (Chow: 0.5 U kg^{-1} body weight; HFD: 60% of insulin administered to chow-fed mice) glucose uptake index in gastrocnemius (Gast), quadriceps (Quad), triceps brachii (Triceps) and soleus muscle from chow- (C) and HFD-fed (D) PAK1 KO mice or control littermates. The number of mice in each group: Chow-Saline, *n = 12/4* (Control/PAK1 KO); Chow-Insulin, *n = 12/6*; HFD-Saline, *n = 6/6*; HFD-Insulin, *n = 10/11*. Statistics were evaluated with a two-way ANOVA for each of the muscles. Main effects are indicated in the panels. Data are presented as mean $\pm$ S.E.M. or when applicable mean $\pm$ S.E.M. with individual data points shown.

Figure 4: Whole-body energy utilization is unaffected by whole-body genetic ablation of PAK1 and muscle-specific genetic ablation of PAK2. (A) Representative blots showing PAK1and PAK2 protein expression in soleus and extensor digitorum longus (EDL) muscle from chow-fed whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (d1/2KO) mice or control littermates. **(B)** Body weight of chow-fed PAK1 KO (*n = 18*), PAK2 mKO (*n = 21*), d1/2 KO mice (*n = 23*) or control littermates (*n = 20*). The mice were 16-29 weeks of age. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with one-way ANOVA. (**C**) Body composition (FM: Fat mass; LBM: Lean body mass) in gram in chow-fed PAK1 KO (*n = 18*), PAK2 mKO (*n = 21*), d1/2 KO mice (*n = 23*)

or control littermates ($n = 20$). Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. **(D-G)** Oxygen uptake (VO$_2$; D), respiratory exchange ratio (RER; E), activity (beam breaks; F), and energy intake (G), in chow-fed PAK1 KO ($n = 11$), PAK2 mKO ($n = 11$), d1/2 KO mice ($n = 13$) or control littermates ($n = 8;$ for energy intake, $n = 7$) during the light and dark period recorded over a period of 72 hours. On day 2, the mice were fasted during the dark period and then refed on day 3. The mice were 23-33 weeks of age. Statistics were evaluated with two two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) in the light and dark period of day 1, respectively. Statistics for day 2 and 3 were evaluated similarly thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of the light and dark period were assessed with three two-way repeated measures (RM) ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. d1/2 KO) and 'Period' (Light vs. Dark) at day 1, 2 and 3, respectively. Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way (RM when applicable) ANOVA were evaluated by Tukey's post hoc test: Light vs. dark period */**/*** ($p<0.05/0.01/0.001$); Control vs. d1/2 KO †/††† ($p<0.05/0.001$); PAK1 KO vs. d1/2 KO ‡/‡‡/‡‡‡ ($p<0.05/0.01/0.001$); PAK2 mKO vs. d1/2 KO (\$)/\$/\$\$ ($p<0.1/0.05/0.01$). Data are presented as mean $\pm$ S.E.M. or when applicable mean $\pm$ S.E.M. with individual data points shown.

Figure 5: Mice lacking PAK2 in skeletal muscle are slightly glucose intolerant. (A) Blood glucose concentration in the fed state (8 a.m.) in chow-fed whole-body PAK1 knockout (KO) ($n = 18$), muscle-specific PAK2 (m)KO ($n = 21$), PAK1/2 double KO (d1/2KO) mice ($n = 23$) or control littermates ($n = 20$). The mice were 17-30 weeks of age. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. **(B)** Blood glucose levels during a glucose tolerance test (GTT) in chow-fed PAK1 KO ($n = 18$), PAK2 mKO ($n = 19$), d1/2 KO mice ($n = 21$) or control littermates ($n = 19$). The mice were 13-26 weeks of age. Statistics were evaluated with six two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) at time point 0, 15, 30, 60, 90 and 120, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of glucose administration were assessed with a two-way repeated measures (RM) ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. d1/2 KO) and 'Time' (0 vs. 15 vs. 30

vs. 60 vs. 90 vs. 120). **(C)** Incremental Area Under the Curve (AUC) for blood glucose levels during the GTT in panel B. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with one-way ANOVA. **(D)** Plasma insulin values during a GTT in chow-fed PAK1 KO ($n = 17$), PAK2 mKO ($n = 19$), d1/2 KO mice ($n = 22$) or control littermates ($n = 19$). The mice were 15-28 weeks of age. Statistics were evaluated with two two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) at time point 0 and 20, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of glucose administration were assessed with a two-way RM ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. d1/2 KO) and 'Time' (0 vs. 20). **(E)** Homeostatic Model Assessment of Insulin Resistance (HOMA-IR) in chow-fed PAK1 KO ($n = 18$), PAK2 mKO ($n = 19$), d1/2 KO mice ($n = 22$) or control littermates ($n = 20$). Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. **(F)** Blood glucose levels related to the basal concentration during an insulin tolerance test (ITT) in chow-fed PAK1 KO ($n = 15$), PAK2 mKO ($n = 18$), d1/2 KO mice ($n = 21$) or control littermates ($n = 19$). The mice were 11-24 weeks of age. For two female control mice and four female PAK2 mKO mice, the ITT had to be stopped before the 120'-time point due to hypoglycemia (blood glucose <1.2 mM), and thus blood glucose was not determined for these mice for the last couple of time points. Statistics were evaluated with five two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) at time point 15, 30, 60, 90 and 120, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of insulin administration were assessed with a mixed-effects model analysis to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. d1/2 KO) and 'Time' (0 vs. 15 vs. 30 vs. 60 vs. 90 vs. 120). Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way (RM when applicable) ANOVA were evaluated by Tukey's post hoc test: Effect of glucose/insulin administration vs. time point 0' */*** (p<0.05/0.001). Data are presented as mean ± S.E.M. or when applicable mean ± S.E.M. with individual data points shown. Paired data points are connected with a straight line.

Figure 6: Muscle-specific PAK2 knockout, but not knockout of PAK1, partly reduces glucose uptake in EDL muscle. (A-B) Insulin-stimulated (60 nM) 2-Deoxyglucose (2DG) uptake in isolated soleus (A) and extensor digitorum longus (EDL; B) muscle from whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO

(d1/2KO) mice or control littermates. Isolated muscles were pre-incubated for 20 minutes followed by 20 minutes of insulin stimulation with 2DG uptake analyzed for the final 10 minutes of stimulation. The mice were 10-16 weeks of age. Statistics were evaluated with two two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) in basal and insulin-stimulated samples, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of insulin stimulation were assessed with a two-way repeated measures (RM) ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. d1/2 KO) and 'Stimuli' (Basal vs. Insulin). **(C-D)** Δ2DG uptake in soleus (C) and EDL (D) muscle from panel C-D. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) to assess of the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. The number of determinations in each group: Control, $n = 9/10$ (soleus/EDL); PAK1 KO, $n = 8/9$; PAK2 KO, $n = 10/10$; PAK1/2 dKO, $n = 9/9$. Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way (RM when applicable) ANOVA were evaluated by Tukey's post hoc test: Insulin-stimulation vs. basal control *** ($p<0.001$); Control vs. PAK2 mKO £ ($p<0.05$); PAK1 KO vs. PAK2 mKO § ($p<0.05$). Data are presented as mean ± S.E.M. with individual data points shown. Paired data points are connected with a straight line.

Figure 7: Lack of PAK2 affects TBC1D4 protein expression and signalling. (A-J) Quantification of phosphorylated (p)Akt S473, pAkt T308, pTBC1D4 T642 and total TBC1D4 and GLUT4 protein expression in insulin-stimulated (60 nM) soleus (A, C, E, G, and I) and extensor digitorum longus (EDL; B, D, F, H, and J) muscle from whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (d1/2KO) mice or control littermates. The mice were 10-16 weeks of age. Total protein expression is an average of the paired basal and insulin-stimulated sample. Some of the data points were excluded due to the quality of the immunoblot, so the number of determinations for GLUT4 in soleus muscle: Control, $n = 6$; PAK1 KO, $n = 5$; PAK2 KO, $n = 6$; PAK1/2 dKO, $n = 6$. **(K-L)** Representative blots showing pAkt S473, pAkt T308, pTBC1D4 T642 and total PAK1, PAK2, Akt2, TBC1D4 and GLUT4 protein expression and coomassie staining as a loading control in soleus (K) and EDL (L) muscle. Statistics for phosphorylated proteins were evaluated with a two two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) in basal and insulin-stimulated samples, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of insulin stimulation were assessed with a two-way repeated measures (RM) ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. d1/2 KO) and 'Stimuli' (Basal vs. Insulin). Statistics for total protein expression were

evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way (RM when applicable) ANOVA were evaluated by Tukey's post hoc test: Control vs. PAK2 mKO £ ($p<0.05$). Unless otherwise stated previously in the figure legend, the number of determinations in each group: Control, $n = 9/10$ (soleus/EDL); PAK1 KO, $n = 8/9$; PAK2 KO, $n = 10/9$; PAK1/2 dKO, $n = 9/9$. Data are presented as mean $\pm$ S.E.M. with individual data points shown. Paired data points are connected with a straight line. A.U., arbitrary units.

Supplementary figures

Figure S1: (A-B) Insulin-stimulated (Chow: 0.5 U kg^{-1} body weight; HFD: 60% of insulin administered to chow-fed mice) 2-Deoxyglucose (2DG) clearance in gastrocnemius (Gast), quadriceps (Quad), triceps brachii (Triceps) and soleus muscle from chow- (A) and 60E% high-fat diet (HFD)-fed (B) PAK1 knockout (KO) mice or control littermates. The number of mice in each group: Chow-Saline, $n = 12/4$ (Control/PAK1 KO); Chow-Insulin, $n = 12/6$; HFD-Saline, $n = 6/6$; HFD-Insulin, $n = 10/11$. Statistics were evaluated with a two-way ANOVA for each of the muscles. **(C)** Plasma [^{3}H] counts 10 minutes after retro-orbital (r.o.) administration of a bolus of saline containing [^{3}H]-labelled 2DG ([^{3}H]-2DG) with or without insulin. Statistics were evaluated with two two-way ANOVAs to test the factors 'stimuli' (basal vs. insulin) and 'genotype' (control vs. PAK1 KO) in chow-fed and HFD-fed mice, respectively. **(D)** Body composition (FM: Fat mass; LBM: Lean body mass; BW: Body weight) in gram in HFD-fed PAK1 KO mice ($n = 17$) or control littermates ($n = 14$). Body composition was assessed in week 18-19 of the diet intervention. The mice were 24-34 weeks of age. Statistics were evaluated with a Student's t-test. **(E)** Energy intake in HFD-fed PAK1 KO mice ($n = 8$) or control littermates ($n = 8$). Energy intake was monitored in a separate cohort of mice. Statistics were evaluated with a Student's t-test. **(F)** Blood glucose levels during a glucose tolerance test (GTT) in HFD-fed PAK1 KO mice ($n = 17$) or control littermates ($n = 13$). Glucose tolerance was assessed in week 14 of the diet intervention. The mice were 20-30 weeks of age. Statistics were evaluated with a two-way repeated measures (RM) ANOVA. **(G)** Incremental Area Under the Curve (AUC) for blood glucose levels during the GTT in panel F. Statistics were evaluated with a Student's t-test. **(H)** Plasma insulin values during a GTT in HFD-fed PAK1 KO mice ($n = 16$) or control littermates ($n = 11$). The plasma insulin response to glucose administration was measured in week 16 of the diet intervention. The mice were 22-

32 weeks of age. Statistics were evaluated with a two-way RM ANOVA. **(I)** Homeostatic Model Assessment of Insulin Resistance (HOMA-IR) in HFD-fed PAK1 KO mice ($n = 16$) or control littermates ($n = 11$). Statistics were evaluated with a Student's t-test. Main effects are indicated in the panels. Data are presented as mean $\pm$ S.E.M. or when applicable mean $\pm$ S.E.M. with individual data points shown. Paired data points are connected with a straight line.

Figure S2: (A-B) Body weight of female (A) and male (B) whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (d1/2KO) mice or control littermates. **(C-D)** Body composition (FM: Fat mass; LBM: Lean body mass) in gram in female (C) and male (D) PAK1 KO, PAK2 mKO, d1/2 KO or control littermates. **(E-G)** Body composition (FM: Fat mass; LBM: Lean body mass) in percentage in both sexes combined (E) and in female (F) and male (G) PAK1 KO, PAK2 mKO, d1/2 KO or control littermates. The number of mice in each group: Control, $n = 9/11$ (female/male); PAK1 KO, $n = 8/10$; PAK2 mKO, $n = 12/9$; d1/2 KO, $n = 9/14$. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way ANOVA were evaluated by Tukey's post hoc test: Control vs. d1/2 KO †/†† (p<0.05/0.01); PAK1 KO vs. PAK2 mKO § (p<0.05); PAK1 KO vs. d1/2 KO ‡/‡‡/‡‡‡ (p<0.05/0.01/0.001); PAK2 mKO vs. d1/2 KO ($)/$$ (p<0.1/0.01). Data are presented as mean $\pm$ S.E.M. with individual data points shown.

Figure S3: (A-F) Oxygen uptake (VO$_2$), respiratory exchange ratio (RER), and activity (beam breaks) in female (A,C, and E) and male (B,D, and F) whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (d1/2KO) mice or control littermates during the light and dark period recorded over a period of 72 hours in calorimetric chambers. On day 2, mice fasted during the dark period and then refed on day 3. **(G-I)** Total energy intake on day 1 and day 3 in both sexes combined (G) and in female (H) and male (I) whole-body PAK1 KO, PAK2 mKO, d1/2 KO mice or control littermates recorded over a period of 72 hours in calorimetric chambers. **(J-K)** Energy intake in female (J) and male (K) whole-body PAK1 KO, Pak2 mKO, d1/2 KO mice or control littermates during the light and dark period recorded over a period of 72 hours in calorimetric chambers. On day 2, mice fasted during the dark period and were then refed on day 3. The number of mice in each group: Control, $n = 5/3$ (female/male; for energy intake, $n = 4/3$); PAK1 KO, $n = 6/5$; PAK2 mKO, $n = 6/5$; d1/2 KO, $n = 6/7$. **(A-F+J-K)** Statistics were evaluated with two two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) in the light and dark period of day 1, respectively. Statistics for day 2 and 3 were evaluated similarly thereby assessing the

relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of the light and the dark period were assessed with three two-way repeated measures (RM) ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. d1/2 KO) and 'Period' (Light vs. Dark) at day 1, 2 and 3, respectively. **(G-I)** Statistics were evaluated with two two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) at day 1 and day 2, respectively. Differences between genotypes and the day were assessed with one two-way repeated measures (RM) ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. d1/2 KO) and 'Day' (Day 1 vs. Day 3). Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way (RM when applicable) ANOVA were evaluated by Tukey's post hoc test: Light vs. dark period */**/*** ($p<0.05/0.01/0.001$); Day 1 vs. Day 3 €€/€€€ ($p<0.01/0.001$); Control vs. PAK1 KO ¤¤ ($p<0.01$); Control vs. d1/2 KO † ($p<0.05$); PAK1 KO vs. PAK2 mKO §/§§ ($p<0.05/0.01$); PAK1 KO vs. d1/2 KO ‡/‡‡/‡‡‡ ($p<0.05/0.01/0.001$). Data are presented as mean $\pm$ S.E.M. with individual data points shown.

Figure S4: (A-B) Blood glucose concentration in the fed state (8 a.m.) in female (A) and male (B) whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (d1/2KO) mice or control littermates. The number of mice in each group: Control, $n = 9/11$ (female/male); PAK1 KO, $n = 8/10$; PAK2 mKO, $n = 12/9$; d1/2 KO, $n = 9/14$. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. **(C-D)** Blood glucose levels during a glucose tolerance test (GTT) in female (C) and male (D) PAK1 KO, PAK2 mKO, d1/2 KO mice or control littermates. The number of mice in each group: Control, $n = 8/11$ (female/male); PAK1 KO, $n = 8/10$; PAK2 mKO, $n = 10/9$; d1/2 KO, $n = 8/13$. Statistics were evaluated with six two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) at time point 0, 15, 30, 60, 90 and 120, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of glucose administration were assessed with a two-way repeated measures (RM) ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. d1/2 KO) and 'Time' (0 vs. 15 vs. 30 vs. 60 vs. 90 vs. 120). **(E-F)** Incremental Area Under the Curve (AUC) for blood glucose levels for females (E) and male (F) mice during the GTT in panel C-D. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. **(G-H)** Plasma insulin values during a GTT in female (G) and male (F) PAK1 KO, PAK2 mKO, d1/2 KO mice or control littermates. The number of

mice in each group: Control, *n = 9/10* (female/male); PAK1 KO, *n = 8/9*; PAK2 mKO, *n = 10/9*; d1/2 KO, *n = 9/13*. Statistics were evaluated with two two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) at time point 0 and 20, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of glucose administration were assessed with a two-way RM ANOVA to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. d1/2 KO) and 'Time' (0 vs. 20). Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way (RM when applicable) ANOVA were evaluated by Tukey's post hoc test: Effect of glucose administration vs. time point 0' */*** ($p<0.05/0.001$); Control vs. PAK2 mKO £ ($p<0.05$); PAK1 KO vs. PAK2 mKO §§ ($p<0.01$); PAK2 mKO vs. d1/2 KO $ ($p<0.05$). Data are presented as mean ± S.E.M. or when applicable mean ± S.E.M. with individual data points shown. Paired data points are connected with a straight line.

Figure S5: (A-B) Homeostatic Model Assessment of Insulin Resistance (HOMA-IR) in female (A) and male (B) whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (d1/2KO) mice or control littermates. The number of mice in each group: Control, *n = 9/11* (female/male); PAK1 KO, *n = 8/10*; PAK2 mKO, *n = 10/9*; d1/2 KO, *n = 9/13*. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Genotype differences were evaluated with a one-way ANOVA. **(C-E)** Basal blood glucose concentration (fasted state) immediately before an insulin tolerance test (ITT) in both sexes combined (C) and in female (D) and male (E) PAK1 KO, PAK2 mKO, d1/2 KO mice and control littermates. The number of mice in each group: Control, *n = 9/10* (female/male); PAK1 KO, *n = 6/9*; PAK2 mKO, *n = 10/8*; d1/2 KO, *n = 8/13*. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. **(F-G)** Blood glucose levels related to basal concentration during an ITT in female (F) and male (G) PAK1 KO, PAK2 mKO, d1/2 KO mice or control littermates. The number of mice in each group: Control, *n = 9/10* (female/male); PAK1 KO, *n = 6/9*; PAK2 mKO, *n = 10/8*; d1/2 KO, *n = 8/13*. For two female control mice and four female PAK2 mKO mice, the ITT had to be stopped before the 120'-time point due to hypoglycemia (blood glucose <1.2 mM), and thus blood glucose was not determined for these mice for the last couple of time points. Statistics were evaluated with five two-way ANOVAs to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) at time point 15, 30, 60, 90 and 120, respectively, thereby assessing the relative contribution of PAK1 and PAK2. Differences between genotypes and the effect of insulin

administration were assessed with a two-way repeated measures (RM) ANOVA (mixed-effects model analysis for female mice) to test the factors 'Genotype' (Control vs. PAK1 KO vs. PAK2 mKO vs. d1/2 KO) and 'Time' (0 vs. 15 vs. 30 vs. 60 vs. 90 vs. 120). Main effects are indicated in the panels. Significant one-way ANOVA and interactions in two-way (RM when applicable) ANOVA were evaluated by Tukey's post hoc test: Effect of insulin administration vs. time point 0' **/*** ($p<0.01/0.001$); Control vs. PAK1 KO ¤/¤¤ ($p<0.05/0.01$); Control vs. PAK2 mKO (£)££ ($p<0.1/0.01$); Control vs. d1/2 KO ††† ($p<0.001$); PAK1 KO vs. d1/2 KO ‡ ($p<0.05$); PAK2 mKO vs. d1/2 KO $ ($p<0.05$). Data are presented as mean $\pm$ S.E.M. or when applicable mean $\pm$ S.E.M. with individual data points shown.

Figure S6: (A-D) Δ-phosphorylated (p)-Akt S473 and ΔpAkt T308 in insulin-stimulated (60 nM) soleus (A and C) and extensor digitorum longus (EDL; B and D) muscle from whole-body PAK1 knockout (KO), muscle-specific PAK2 (m)KO, PAK1/2 double KO (d1/2KO) mice or control littermates from Fig. 7A-D. **(E-F)** Quantification of total Akt2 protein expression in soleus (E) and EDL (F) muscle from whole-body PAK1 KO, PAK2 mKO, d1/2 KO mice or control littermates. Total protein expression is an average of the paired basal and insulin-stimulated sample. **(G-H)** ΔpTBC1D4 T642 in insulin-stimulated (60 nM) soleus (G) and EDL (H) muscle from whole-body PAK1 KO, PAK2 mKO, d1/2 KO mice or control littermates from Fig. 7G-H. Statistics were evaluated with a two-way ANOVA to test the factors 'PAK1' (PAK1+/- vs. PAK-/-) and 'PAK2' (PAK2$^{fl/fl}$;MyoD$^{+/+}$ vs. PAK2$^{fl/fl}$;MyoD$^{iCre/+}$) thereby assessing the relative contribution of PAK1 and PAK2, respectively. Differences between genotypes were evaluated with a one-way ANOVA. The number of determinations in each group: Control, *n = 9/10* (soleus/EDL); PAK1 KO, *n = 8/9*; PAK2 KO, *n = 10/9*; PAK1/2 dKO, *n = 9/9*. Significant one-way ANOVA was evaluated by Tukey's post hoc test: Control vs. PAK2 mKO £ ($p<0.05$). Data are presented as mean $\pm$ S.E.M. with individual data points shown. A.U., arbitrary units.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

Soleus

Main effect: Insulin-stimulation (p<0.001)

2DG uptake (µmol g^{-1} h^{-1})

Insulin (60 nM) — + Control — + PAK1 KO — + PAK2 mKO — + d1/2 KO

B

EDL

Main effect: Muscle-specific PAK2 knockout at insulin-stimulation (p<0.05)

2DG uptake (µmol g^{-1} h^{-1})

Insulin (60 nM) — + Control — + PAK1 KO — + PAK2 mKO — + d1/2 KO

C

Soleus

Δ2DG uptake (µmol g^{-1} h^{-1})

Control   PAK1 KO   PAK2 mKO   d1/2 KO

D

EDL

Main effect: Muscle-specific PAK2 knockout (p<0.01)

Δ2DG uptake (µmol g^{-1} h^{-1})

Control   PAK1 KO   PAK2 mKO   d1/2 KO

Figure 7

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

Table 1. Antibody Table

Antibody name	Antibody ID (RRID)	Manufacturer; Catalog number;	Species Raised in; Monoclonal or Polyclonal	Antibody dilution	Blocking buffer
Akt2	AB_2225186	Cell Signaling Technology; 3063	Rabbit; Monoclonal antibody	1:1000	2% milk
pAkt S473	AB_329825	Cell Signaling Technology; 9271	Rabbit; Polyclonal antibody	1:1000	2% milk
pAkt T308	AB_329828	Cell Signaling Technology; 9275	Rabbit; Polyclonal antibody	1:1000	2% milk
GLUT4	AB_2191454	Thermo Fisher Scientific; PA1-1065	Rabbit; Polyclonal antibody	1:1000	2% milk
PAK1	AB_330222	Cell Signaling Technology; 2602	Rabbit; Polyclonal antibody	1:1000	2% milk
PAK2	AB_2283388	Cell Signaling Technology; 2608	Rabbit; Polyclonal antibody	1:1000	2% milk
pPAK1/2 T423/402	AB_330220	Cell Signaling Technology; 2601	Rabbit; Polyclonal antibody	1:1000	5% BSA
TBC1D4	AB_492639	Millipore; 07-741	Rabbit; Polyclonal antibody	1:1000	2% milk
pTBC1D4 T642	AB_2651042	Cell Signaling Technology; 8881	Rabbit; Monoclonal antibody	1:1000	2% milk