The mycobacterial cell wall partitions the plasma membrane to organize its own synthesis

Alam García-Heredia[1], Takehiro Kado[2], Caralyn E. Sein[2], Julia Puffal[2†], Sarah H. Osman[2], Julius

Judd[3††], Todd A. Gray[3,4], Yasu S. Morita[1,2,*], M. Sloan Siegrist[1,2,*].

[1] Molecular and Cellular Biology Graduate Program, University of Massachusetts, Amherst, MA

01003, USA.

[2] Department of Microbiology, University of Massachusetts, Amherst, MA 01003, USA.

[3] Division of Genetics, Wadsworth Center, New York State Department of Health, Albany, NY

12208, USA.

[4] Department of Biomedical Sciences, University at Albany, Albany, NY 12208, USA.

[†] Present address: Department of Biochemistry and Molecular Biology, Rutgers University,

Robert Wood Johnson Medical School, Piscataway, NJ 08854, USA.

[††] Present address: Department of Molecular Biology and Genetics, Cornell University, Ithaca,

NY 14853, USA.

[*] Corresponding author. Email: siegrist@umass.edu (M.S.S.); ymorita@microbio.umass.edu

(Y.S.M.).

Abstract

Many antibiotics target the assembly of cell wall peptidoglycan, an essential, heteropolymeric

mesh that encases most bacteria. Different species have characteristic subcellular sites of

peptidoglycan synthesis that they must carefully maintain for surface integrity and, ultimately,

viability. In rod-shaped bacteria, cell wall elongation is spatially precise yet relies on a limited

pool of lipid-linked precursors that generate and are attracted to membrane disorder. By tracking

enzymes, substrates and products of peptidoglycan biosynthesis in *Mycobacterium smegmatis*,

we show that precursors are made in plasma membrane domains that are laterally and

biochemically distinct from sites of cell wall assembly. Membrane partitioning is required for

robust, orderly peptidoglycan synthesis, indicating that these domains help template

peptidoglycan synthesis. The cell wall-organizing protein DivIVA and the cell wall itself are

essential for domain homeostasis. Thus, the peptidoglycan polymer feeds back on its membrane

template to maintain an environment conducive to directional synthesis. We further show that

our findings are applicable to rod-shaped bacteria that are phylogenetically distant from *M.*

smegmatis, demonstrating that horizontal compartmentalization of precursors is a general feature

of bacillary cell wall biogenesis.

Main

The final lipid-linked precursor for peptidoglycan synthesis, lipid II, is made by the glycosyltransferase MurG in the inner leaflet of the plasma membrane. Lipid II is then flipped to the outer leaflet by MurJ where its disaccharide-pentapeptide cargo is inserted into the existing cell wall by membrane-bound transglycosylases and transpeptidases (Zhao et al., 2017). Early *in vitro* work in *Staphylococcus aureus* and *Escherichia coli* indicated that a fluid microenvironment might stimulate the activities of MurG and the upstream, lipid I synthase MraY (Norris & Manners, 1993). More recent *in vivo* data has localized *Bacillus subtilis* MurG to **r**egions of **i**ncreased **f**luidity (**RIF**s, [Muller et al., 2016; Strahl et al., 2014]), one of three classes of membrane domains that have been described in bacteria to date. In mycobacteria, **i**ntracellular **m**embrane **d**omains (IMD, formerly called PMf, [Morita et al., 2005]) can be separated from the conventional plasma membrane (PM-CW, for **p**lasma **m**embrane associated with **c**ell **w**all) by sucrose density gradient fractionation. The proteome and lipidome of IMD are distinct from those of the PM-CW (Hayashi et al., 2016; Morita et al., 2005). Reanalysis of our proteomics data (Hayashi et al., 2016) suggested that *Mycobacterium smegmatis* MurG is enriched in the IMD while sequentially-acting transglycosylases and transpeptidases associate with the PM-CW. While PM-CW-resident proteins distribute thorough the perimeter of live mycobacteria, IMD-resident proteins localize along the sidewall but are enriched toward sites of polar cell elongation (Hayashi et al., 2016; Hayashi et al., 2018). We also noted that polar enrichment of MurG resembles that of the validated IMD marker GlfT2 (Hayashi et al., 2016; Meniche et al., 2014), but that nascent peptidoglycan at the mycobacterial poles primarily abuts rather than colocalizes with GlfT2 (Hayashi et al., 2018). These observations suggested a model where lipid II synthesis is segregated from subsequent steps of cell wall assembly (Fig. 1A).

To test this model, we first expressed a functional MurG-Dendra2 fusion in *M. smegmatis*

(Figure 1–figure supplement 1A) and assayed its distribution in membrane fractions that had

been separated by density gradient (Fig. 1B). MurG-Dendra2, a peripheral membrane protein,

was enriched in both the cytoplasmic and IMD membrane fractions (Fig. 1C, top; Figure 1 –

figure supplement 2A). In intact cells, polar enrichment of MurG-Dendra2 was coincident with

that of the IMD marker mCherry-GlfT2 (Figure 2–figure supplement 1). This spatial relationship

was similar to that previously observed for MurG and GlfT2 (Hayashi et al., 2016; Hayashi et al.,

2018; Meniche et al., 2014).

Enzymes from the **p**enicillin **b**inding **p**roteins (PBPs) and **s**hape, **e**longation, **d**ivision and

sporulation (SEDS) families integrate the disaccharide-pentapeptide from lipid II into

peptidoglycan (Zhao et al., 2017). While our proteomics did not detect many polytopic

membrane proteins, including SEDS proteins, our PM-CW dataset was enriched for all known

PBPs (Hayashi et al., 2016). Fluorescent derivatives of β-lactam antibiotics, such as Bocillin-FL,

bind to PBPs and report transpeptidase-active enzymes. We incubated subcellular fractions from

wild-type *M. smegmatis* with Bocillin-FL and identified fluorescent proteins in the PM-CW (Fig.

1C, bottom; Figure 1 –figure supplement 2B). As expected for PBPs, the signal from these bands

was diminished by pre-treatment with ampicillin (Figure 1–figure supplement 3). We focused on

characterizing PonA1, an essential bifunctional transglycosylase/transpeptidase in *M. smegmatis*

(Hett et al., 2010; Kieser et al., 2015). Depletion of PonA1 (Hett et al., 2010) resulted in the loss

of the higher molecular band (Figure 1–figure supplement 3), confirming this protein is present

and active in the PM-CW (Fig. 1C, bottom). We next expressed a functional PonA1-mRFP

fusion in *M. smegmatis* (Kieser et al., 2015) and found that it was distributed around the cell

perimeter in a manner similar to the functional PM-CW marker PimE-GFP (Figure 1–figure

supplement 1B; Figure 2–figure supplement 1, [Hayashi et al., 2016]). Coexpression of MurG-

Dendra2 and PonA1-mRFP confirmed that the proteins are laterally segregated from one another

(Fig 2A). Together, our data show that MurG and PonA1 occupy membrane compartments that

are biochemically and spatially distinct.

The association of MurG with the IMD and of PonA1 with the PM-CW implied that the IMD is

the site of lipid II synthesis while the PM-CW is where peptidoglycan assembly takes place. We

refined an *in vitro* D-amino acid exchange assay to detect lipid-linked peptidoglycan precursors

from membrane fractions (Fig. 2B, [García-Heredia et al., 2018; Qiao et al., 2014]). In wild-type

cells, we detected biotinylated molecules in both the IMD and PM-CW (Fig. 2C; Figure 1–figure

supplement 2B). We hypothesized that the labeled species comprise precursors in both the inner

and outer leaflet of the plasma membrane. We and others have shown that depletion of MurJ

results in accumulation of biotinylated precursors (García-Heredia et al., 2018; Qiao et al., 2017).

By performing the D-amino acid exchange reaction on membrane fractions obtained from MurJ-

depleted *M. smegmatis* (Figure 1–figure supplement 1C), we found that precursors accumulate in

the IMD (Fig. 2C; Figure 1–figure supplement 2C). These results suggest that lipid II is made in

the IMD and transferred to the PM-CW in a MurJ-dependent manner.

Based on our biochemical data, we hypothesized that lipid II incorporation into the cell wall is

laterally segregated from its synthesis. We previously showed that alkynyl and azido D-amino

acid dipeptides (Liechti et al., 2014) incorporate into lipid-linked peptidoglycan precursors in *M.*

smegmatis (García-Heredia et al., 2018) and that metabolic labeling with alkynyl dipeptide

(alkDADA or EDA-DA, [Liechti et al., 2014]) is most intense in regions adjacent to the IMD

marker mCherry-GlfT2 (Hayashi et al., 2018). We labeled MurG-Dendra2-expressing *M.*

smegmatis with alkDADA and detected the presence of the alkyne by copper-catalyzed azide-

alkyne cycloaddition (CuAAC, [García-Heredia et al., 2018]). To distinguish extracellular

alkynes present in periplasmic lipid II and newly-polymerized cell wall from alkynes originating

from cytoplasmic lipid II, we selected picolyl azide-Cy3 as our label because of its poor

membrane permeability (Fig. 2B, [Yang & Hinner, 2015]). Using this optimized protocol, we

observed nascent peptidoglycan deposition at the polar tip, whereas MurG-Dendra2 was

proximal to this site (Fig. 2D). Our data suggest that lipid II synthesis is laterally partitioned

from the subsequent steps of peptidoglycan assembly. MurJ depletion reduced and delocalized

alkDADA-derived fluorescence (Figure 2–figure supplement 2), consistent with a gatekeeper

role for the flippase in both lateral membrane compartmentalization and flipping across the inner

membrane.

We next wanted to understand the effect of partitioning on cell wall synthesis. We opted to

perturb the plasma membrane with benzyl alcohol, which has been used to disperse membrane

domains in animal and bacterial cells, including *B. subtilis* RIFs (Muller et al., 2016; Nagy et al.,

2007). In *M. smegmatis,* we found that benzyl alcohol reduced the cellular material associated

with the IMD (Fig 3A, Figure 1–figure supplement 2D) and altered the distribution of FM4-64, a

non-specific lipophilic dye (Figure 3–figure supplement 1A) and of plasma membrane

glycolipids (Figure 3–figure supplement 2). However, benzyl alcohol did not alter labeling by N-

AlkTMM or O-AlkTMM (Figure 3–figure supplement 1B), probes that respectively mark the

noncovalent and covalent lipids of the outer 'myco' membrane (Foley et al., 2016). These

observations suggest that benzyl alcohol primarily affects the plasma membrane. MurG-Dendra2

was also notably less enriched in the IMD fraction following benzyl alcohol treatment (Fig. 1C,

top; Figure 1–figure supplement 2E) and, in live cells, at the poles (Fig. 2A). By contrast, benzyl

alcohol produced subtle changes in the subcellular distribution of active PBPs (Fig. 1C, bottom;

Figure 1–figure supplement 2D), although PonA1 shifted toward the poles in live cells (Fig. 2A).

Disruption of plasma membrane architecture was accompanied by dampening and delocalization

of peptidoglycan assembly (Fig. 3B) as well as a reduction in lipid precursor synthesis (Fig. 3C)

and halt in polar elongation (Figure 3–figure supplement 1B). The effects of benzyl alcohol were

reversible, as indicated by colony forming units and prompt recovery of peptidoglycan synthesis

(Figure 3–figure supplement 3). Dibucaine, a membrane-fluidizing anesthetic (Papahadjopoulos

et al., 1975), also delocalized IMD-resident proteins (Figure 3–figure supplement 4) and reduced

peptidoglycan synthesis (Fig. 3B). These data suggest that membrane architecture is essential

feature of both peptidoglycan synthesis and cell growth in *M. smegmatis*.

Benzyl alcohol delocalized or dampened cell wall assembly in other rod-shaped bacteria with

divergent envelope composition and modes of growth (Fig. 3D-E). Peptidoglycan synthesis was

less affected by benzyl alcohol in coccoid species, which lack MreB, and in rounded, A22-

treated *E. coli,* which have MreB inhibited (Fig. 3E). Thus, membrane partitioning is necessary

for effective, directional cell wall synthesis in rod-shaped bacteria.

How is the IMD partitioned away from the rest of the plasma membrane? The architecture of

eukaryotic membranes is influenced by transient links, or pinning, to the cytoskeleton (Fujimoto

& Parmryd, 2016; Liu et al., 2015). In *B. subtilis* and *E. coli,* actin homologs like MreB organize

RIFs (Oswald et al., 2016; Strahl et al., 2014) and direct peptidoglycan synthesis along the lateral

cell surface (Daniel & Errington, 2003; Iwai et al., 2002; Shi et al., 2018; Zhao et al., 2017).

While MreB is absent in mycobacteria, the essential tropomyosin-like protein DivIVA

concentrates cell wall assembly at the poles (Daniel & Errington, 2003; Iwai et al., 2002; Melzer

et al., 2018; Shi et al., 2018; Zhao et al., 2017). DivIVA depletion results in deformation and

rounding of mycobacterial cells (Kang et al., 2008). Given the similarities in DivIVA and MreB

function, we hypothesized that DivIVA creates and/or maintains the IMD. We used *M.*

smegmatis expressing DivIVA-eGFP-ID (in which DivIVA is fused to both eGFP and an

inducible degradation tag [Meniche et al., 2014]) to deplete DivIVA. Depletion of the protein

reduced the amount of IMD-associated cellular material (Fig. 4A, Figure 1–figure supplement

2F-G), altered the distribution of plasma membrane glycolipids (Figure 3–figure supplement 2),

and delocalized the IMD marker mCherry-GlfT2 from the poles (Fig. 4B). DivIVA

phosphorylation regulates MraY and/or MurG activity via an indirect, unknown mechanism (Jani

et al., 2010). Consistent with these data, we found that depletion of the protein reduced lipid-

linked peptidoglycan precursor abundance (Fig. 3C). The suppressive effects of benzyl alcohol

and DivIVA depletion on precursor abundance were not additive (Fig. 3C), suggesting that the

perturbations act on the same pathway.

As lipid II both generates and homes to disordered regions of model membranes (Ganchev et al.,

2006; Jia et al., 2011; Valtersson et al., 1985), the effect of DivIVA on precursors suggests that

concentrated peptidoglycan synthesis is a cause or a consequence (or both) of IMD/PM-CW

partitioning. In other organisms, lipid II production is required for MreB rotation (Dominguez-

Escobar et al., 2011; Garner et al., 2011; van Teeffelen et al., 2011) and to recruit MreB to the

plasma membrane (Schirner et al., 2015), so the precursor might also play an indirect role in

compartmentalizing the mycobacterial membrane via its influence on DivIVA. However, we

previously found that the IMD is biochemically intact after 8 hours of treatment with D-

cycloserine (Hayashi et al., 2018), an antibiotic that we have shown to block *M. smegmatis*

peptidoglycan precursor synthesis within 1 hour (García-Heredia et al., 2018). IMD-resident

proteins delocalize, but not until 6 hours of treatment. The persistence of IMD-resident proteins

and the time frame of delocalization indicate that concentrated lipid II synthesis is more likely a

consequence, rather than a cause, of mycobacterial membrane compartmentalization.

Our data supported a model in which DivIVA is necessary to maintain membrane partitioning,

which in turn supports efficient synthesis of peptidoglycan precursors and their precise

incorporation into the cell wall. However, we noted that eukaryotic membrane architecture can

also be influenced by transient interactions with external structures like extracellular matrix and

cellulose (Fujimoto & Parmryd, 2016; Liu et al., 2015). Furthermore, biophysical modeling of

bacteria suggests that osmotic pinning of the plasma membrane against the cell wall can induce

microphase separation (Mukhopadhyay et al., 2008). In mycobacteria, co-fractionation of the

plasma **m**embrane and **c**ell **w**all (*i.e.* PM-CW) upon mechanical cell lysis implies that they are

physically connected (Morita et al., 2005). We wondered whether the peptidoglycan polymer

itself might contribute to the membrane partitioning that organizes its synthesis. To test this

hypothesis, we spheroplasted (enzymatically removed the cell wall; Melzer et al., 2018) *M.*

smegmatis that expressed MurG-Dendra2 or DivIVA-eGFP-ID and mCherry-GlfT2. All fusions

were functional (Figure 1–figure supplement 1, [Hayashi et al., 2016; Meniche et al., 2014]).

Fractionated lysates from spheroplasted mycobacteria had more diffuse distribution of cellular

material and indistinct separation of the IMD and PM-CW fractions (Fig. 4A). Consistent with

the macroscopic appearance of the fractionated lysate, MurG-Dendra2 was distributed

throughout the gradient (Figure 4–figure supplement 1). MurG-Dendra2 and the IMD marker

mCherry-GlfT2 were also diffusely distributed around the periphery of spheroplasted cells while

DivIVA, a non-IMD protein, remained in foci (Fig. 4C-D). These experiments indicate that an

intact cell wall and DivIVA are both required for membrane compartmentalization in *M.*

smegmatis (Fig. 4E).

While peptidoglycan biogenesis is well known to vertically span the inner and outer leaflets of

the plasma membrane, here we demonstrate in *M. smegmatis* that it is also horizontally

partitioned. Partitioning of the mycobacterial membrane by DivIVA and the cell wall follows

similar logic to that of eukaryotic membranes, which can be compartmentalized by pinning to

cytoplasmic structures such as the actin cytoskeleton and to external structures like extracellular

matrix and cellulose (Fujimoto & Parmryd, 2016; Liu et al., 2015), and of model lipid bilayers,

which can be phase separated by adhesive forces (Gordon et al., 2008; Mukhopadhyay et al.,

2008). In mycobacteria, however, the membrane regions that promote cell wall synthesis are

segregated by the end product of the pathway (Fig. 4E). For these and other rod-shaped bacteria,

our model is that the membrane-cell wall axis is a self-organizing system in which directed cell

wall synthesis organizes the plasma membrane, and an organized plasma membrane in turn

makes cell wall elongation more efficient and precise.

Acknowledgments

We are grateful to Drs. James Chambers and Amy Burnside for microscopy and flow cytometry

guidance, Drs. Eric Strieter and Jiaan Liu and Ms. Sylvia Rivera and Katherine Chacón-Vargas

for technical assistance. We also thank Drs. Hesper Rego and Eric Rubin for *ponA1-mRFP M.*

smegmatis, Dr. Christopher Sassetti for the MurJ and MurG depletion strains, and Dr. Benjamin

Swarts for O-AlkTMM and N-AlkTMM. Research was supported by funds from the National

Institutes of Health (NIH) under awards R21 AI144748 (Y.S.M and M.S.S.), U01 CA221230

and NIH DP2 AI138238 (M.S.S.) and R03 AI140259-01 (Y.S.M.), and R01 AI097191 (J.J. and

T.A.G.). A.G.-H. was supported by an Honors Fellowship from Universidad Autónoma de

Nuevo León. J.P. is a recipient of the Science Without Boarders Fellowship from CAPES-Brazil

(0328-13-8). T.K. was supported by a postdoctoral fellowship from Uehara Memorial

Foundation.

Contributions

M.S.S., A.G.-H. and Y.S.M. designed the experiments and analyzed the data. A.G.-H.

performed most of the experiments. T.K. constructed the DivIVA-eGFP-ID/mCherry-GlfT2 *M.*

smegmatis mutant and helped with experiments from Figs. 3B, 4B and Figure 3–figure

supplement 4. J.P. performed immunoblots for Figure 1–figure supplement 2 and thin-layer

chromatography for Figure 3–figure supplement 2. C.E.S performed experiments for Fig. 4C.

S.H.O. did experiments for Fig. 4A and Figure 4–figure supplement 1. Y.S.M. did the glycolipid

analysis for Figure 1–figure supplement 1. J.J. and T.A.G. constructed the MurG-Dendra2

plasmid. M.S.S. and A.G.-H. wrote the manuscript. All authors reviewed and edited the

manuscript.

Competing interests

Authors declare no competing interests.

Data and materials availability

All data are available in the main text or the supplementary materials. Raw data is available at

open science framework through the next link:

https://osf.io/fm794/?view_only=7d5a24b82dd94744a46420aee09bad7f

Figure supplements

Supplementary information: Tables S1-4

References

Daniel, R. A., & Errington, J. (2003). Control of cell morphogenesis in bacteria: two distinct
ways to make a rod-shaped cell. *Cell, 113*(6), 767-776. doi:10.1016/s0092-
8674(03)00421-5
Dominguez-Escobar, J., Chastanet, A., Crevenna, A. H., Fromion, V., Wedlich-Soldner, R., &
Carballido-Lopez, R. (2011). Processive movement of MreB-associated cell wall

biosynthetic complexes in bacteria. *Science, 333*(6039), 225-228. doi:10.1126/science.1203466

Foley, H. N., Stewart, J. A., Kavunja, H. W., Rundell, S. R., & Swarts, B. M. (2016). Bioorthogonal Chemical Reporters for Selective In Situ Probing of Mycomembrane Components in Mycobacteria. *Angew Chem Int Ed Engl, 55*(6), 2053-2057. doi:10.1002/anie.201509216

Fujimoto, T., & Parmryd, I. (2016). Interleaflet Coupling, Pinning, and Leaflet Asymmetry-Major Players in Plasma Membrane Nanodomain Formation. *Front Cell Dev Biol, 4*, 155. doi:10.3389/fcell.2016.00155

Ganchev, D. N., Hasper, H. E., Breukink, E., & de Kruijff, B. (2006). Size and orientation of the lipid II headgroup as revealed by AFM imaging. *Biochemistry, 45*(19), 6195-6202. doi:10.1021/bi051913e

García-Heredia, A., Pohane, A. A., Melzer, E. S., Carr, C. R., Fiolek, T. J., Rundell, S. R., . . . Siegrist, M. S. (2018). Peptidoglycan precursor synthesis along the sidewall of pole-growing mycobacteria. *Elife, 7*. doi:10.7554/eLife.37243

Garner, E. C., Bernard, R., Wang, W., Zhuang, X., Rudner, D. Z., & Mitchison, T. (2011). Coupled, circumferential motions of the cell wall synthesis machinery and MreB filaments in B. subtilis. *Science, 333*(6039), 222-225. doi:10.1126/science.1203285

Gee, C. L., Papavinasasundaram, K. G., Blair, S. R., Baer, C. E., Falick, A. M., King, D. S., . . . Alber, T. (2012). A phosphorylated pseudokinase complex controls cell wall synthesis in mycobacteria. *Sci Signal, 5*(208), ra7. doi:10.1126/scisignal.2002525

Gordon, V. D., Deserno, M., Andrew, C. M. J., Egelhaaf, S. U., & Poon, W. C. K. (2008). Adhesion promotes phase separation in mixed-lipid membranes. *EPL (Europhysics Letters), 84*(4), 48003. doi:10.1209/0295-5075/84/48003

Hayashi, J. M., Luo, C. Y., Mayfield, J. A., Hsu, T., Fukuda, T., Walfield, A. L., . . . Morita, Y. S. (2016). Spatially distinct and metabolically active membrane domain in mycobacteria. *Proc Natl Acad Sci U S A, 113*(19), 5400-5405. doi:10.1073/pnas.1525165113

Hayashi, J. M., Richardson, K., Melzer, E. S., Sandler, S. J., Aldridge, B. B., Siegrist, M. S., & Morita, Y. S. (2018). Stress-Induced Reorganization of the Mycobacterial Membrane Domain. *MBio, 9*(1). doi:10.1128/mBio.01823-17

Hett, E. C., Chao, M. C., & Rubin, E. J. (2010). Interaction and modulation of two antagonistic cell wall enzymes of mycobacteria. *PLoS Pathog, 6*(7), e1001020. doi:10.1371/journal.ppat.1001020

Iwai, N., Nagai, K., & Wachi, M. (2002). Novel S-benzylisothiourea compound that induces spherical cells in Escherichia coli probably by acting on a rod-shape-determining protein(s) other than penicillin-binding protein 2. *Biosci Biotechnol Biochem, 66*(12), 2658-2662. doi:10.1271/bbb.66.2658

Jani, C., Eoh, H., Lee, J. J., Hamasha, K., Sahana, M. B., Han, J. S., . . . Kang, C. M. (2010). Regulation of polar peptidoglycan biosynthesis by Wag31 phosphorylation in mycobacteria. *BMC Microbiol, 10*, 327. doi:10.1186/1471-2180-10-327

Jia, Z., O'Mara, M. L., Zuegg, J., Cooper, M. A., & Mark, A. E. (2011). The effect of environment on the recognition and binding of vancomycin to native and resistant forms of lipid II. *Biophys J, 101*(11), 2684-2692. doi:10.1016/j.bpj.2011.10.047

Kang, C. M., Nyayapathy, S., Lee, J. Y., Suh, J. W., & Husson, R. N. (2008). Wag31, a homologue of the cell division protein DivIVA, regulates growth, morphology and polar

cell wall synthesis in mycobacteria. *Microbiology, 154*(Pt 3), 725-735. doi:10.1099/mic.0.2007/014076-0

Kieser, K. J., Boutte, C. C., Kester, J. C., Baer, C. E., Barczak, A. K., Meniche, X., . . . Rubin, E. J. (2015). Phosphorylation of the Peptidoglycan Synthase PonA1 Governs the Rate of Polar Elongation in Mycobacteria. *PLoS Pathog, 11*(6), e1005010. doi:10.1371/journal.ppat.1005010

Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol, 35*(6), 1547-1549. doi:10.1093/molbev/msy096

Liechti, G. W., Kuru, E., Hall, E., Kalinda, A., Brun, Y. V., VanNieuwenhze, M., & Maurelli, A. T. (2014). A new metabolic cell-wall labelling method reveals peptidoglycan in Chlamydia trachomatis. *Nature, 506*(7489), 507-510. doi:10.1038/nature12892

Liu, Z., Persson, S., & Sanchez-Rodriguez, C. (2015). At the border: the plasma membrane-cell wall continuum. *J Exp Bot, 66*(6), 1553-1563. doi:10.1093/jxb/erv019

Melzer, E. S., Sein, C. E., Chambers, J. J., & Sloan Siegrist, M. (2018). DivIVA concentrates mycobacterial cell envelope assembly for initiation and stabilization of polar growth. *Cytoskeleton (Hoboken), 75*(12), 498-507. doi:10.1002/cm.21490

Meniche, X., Otten, R., Siegrist, M. S., Baer, C. E., Murphy, K. C., Bertozzi, C. R., & Sassetti, C. M. (2014). Subpolar addition of new cell wall is directed by DivIVA in mycobacteria. *Proc Natl Acad Sci U S A, 111*(31), E3243-3251. doi:10.1073/pnas.1402158111

Morita, Y. S., Sena, C. B., Waller, R. F., Kurokawa, K., Sernee, M. F., Nakatani, F., . . . Maeda, Y. (2006). PimE is a polyprenol-phosphate-mannose-dependent mannosyltransferase that transfers the fifth mannose of phosphatidylinositol mannoside in mycobacteria. *Journal of Biological Chemistry*.

Morita, Y. S., Velasquez, R., Taig, E., Waller, R. F., Patterson, J. H., Tull, D., . . . McConville, M. J. (2005). Compartmentalization of lipid biosynthesis in mycobacteria. *J Biol Chem, 280*(22), 21645-21652. doi:10.1074/jbc.M414181200

Mukhopadhyay, R., Huang, K. C., & Wingreen, N. S. (2008). Lipid localization in bacterial cells through curvature-mediated microphase separation. *Biophys J, 95*(3), 1034-1049. doi:10.1529/biophysj.107.126920

Muller, A., Wenzel, M., Strahl, H., Grein, F., Saaki, T. N. V., Kohl, B., . . . Hamoen, L. W. (2016). Daptomycin inhibits cell envelope synthesis by interfering with fluid membrane microdomains. *Proc Natl Acad Sci U S A, 113*(45), E7077-e7086. doi:10.1073/pnas.1611173113

Nagy, E., Balogi, Z., Gombos, I., Akerfelt, M., Bjorkbom, A., Balogh, G., . . . Vigh, L. (2007). Hyperfluidization-coupled membrane microdomain reorganization is linked to activation of the heat shock response in a murine melanoma cell line. *Proc Natl Acad Sci U S A, 104*(19), 7945-7950. doi:10.1073/pnas.0702557104

Norris, V., & Manners, B. (1993). Deformations in the cytoplasmic membrane of Escherichia coli direct the synthesis of peptidoglycan. The hernia model. *Biophys J, 64*(6), 1691-1700. doi:10.1016/S0006-3495(93)81541-8

Oswald, F., Varadarajan, A., Lill, H., Peterman, E. J., & Bollen, Y. J. (2016). MreB-Dependent Organization of the E. coli Cytoplasmic Membrane Controls Membrane Protein Diffusion. *Biophys J, 110*(5), 1139-1149. doi:10.1016/j.bpj.2016.01.010

Paintdakhi, A., Parry, B., Campos, M., Irnov, I., Elf, J., Surovtsev, I., & Jacobs-Wagner, C. (2016). Oufti: an integrated software package for high-accuracy, high-throughput

quantitative microscopy analysis. *Mol Microbiol, 99*(4), 767-777.
doi:10.1111/mmi.13264
Papahadjopoulos, D., Jacobson, K., Poste, G., & Shepherd, G. (1975). Effects of local
anesthetics on membrane properties. I. Changes in the fluidity of phospholipid bilayers.
Biochim Biophys Acta, 394(4), 504-519. doi:10.1016/0005-2736(75)90137-6
Qiao, Y., Lebar, M. D., Schirner, K., Schaefer, K., Tsukamoto, H., Kahne, D., & Walker, S.
(2014). Detection of lipid-linked peptidoglycan precursors by exploiting an unexpected
transpeptidase reaction. *J Am Chem Soc, 136*(42), 14678-14681. doi:10.1021/ja508147s
Qiao, Y., Srisuknimit, V., Rubino, F., Schaefer, K., Ruiz, N., Walker, S., & Kahne, D. (2017).
Lipid II overproduction allows direct assay of transpeptidase inhibition by beta-lactams.
Nat Chem Biol, 13(7), 793-798. doi:10.1038/nchembio.2388
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., . . .
Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nat*
Methods, 9(7), 676-682. doi:10.1038/nmeth.2019
Schirner, K., Eun, Y. J., Dion, M., Luo, Y., Helmann, J. D., Garner, E. C., & Walker, S. (2015).
Lipid-linked cell wall precursors regulate membrane association of bacterial actin MreB.
Nat Chem Biol, 11(1), 38-45. doi:10.1038/nchembio.1689
Sena, C. B., Fukuda, T., Miyanagi, K., Matsumoto, S., Kobayashi, K., Murakami, Y., . . . Morita,
Y. S. (2010). Controlled expression of branch-forming mannosyltransferase is critical for
mycobacterial lipoarabinomannan biosynthesis. *Journal of Biological Chemistry*, jbc.
M109. 077297.
Shi, H., Bratton, B. P., Gitai, Z., & Huang, K. C. (2018). How to Build a Bacterial Cell: MreB as
the Foreman of E. coli Construction. *Cell, 172*(6), 1294-1305.
doi:10.1016/j.cell.2018.02.050
Strahl, H., Burmann, F., & Hamoen, L. W. (2014). The actin homologue MreB organizes the
bacterial cell membrane. *Nat Commun, 5*, 3442. doi:10.1038/ncomms4442
Valtersson, C., van Duyn, G., Verkleij, A. J., Chojnacki, T., de Kruijff, B., & Dallner, G. (1985).
The influence of dolichol, dolichol esters, and dolichyl phosphate on phospholipid
polymorphism and fluidity in model membranes. *J Biol Chem, 260*(5), 2742-2751.
Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/3919007
van Teeffelen, S., Wang, S., Furchtgott, L., Huang, K. C., Wingreen, N. S., Shaevitz, J. W., &
Gitai, Z. (2011). The bacterial actin MreB rotates, and rotation depends on cell-wall
assembly. *Proc Natl Acad Sci U S A, 108*(38), 15822-15827.
doi:10.1073/pnas.1108999108
Yang, N. J., & Hinner, M. J. (2015). Getting across the cell membrane: an overview for small
molecules, peptides, and proteins. *Methods Mol Biol, 1266*, 29-53. doi:10.1007/978-1-
4939-2272-7_3
Zhao, H., Patel, V., Helmann, J. D., & Dorr, T. (2017). Don't let sleeping dogmas lie: new views
of peptidoglycan synthesis and its regulation. *Mol Microbiol, 106*(6), 847-860.
doi:10.1111/mmi.13853

Main figures

Fig. 1. MurG is enriched in the IMD and PBPs associate with PM-CW. (A) Membrane-bound
steps of peptidoglycan synthesis with hypothesized partitioning into IMD and PM-CW. (B)
Bacteria are lysed by nitrogen cavitation and cell lysate is sedimented on a sucrose density
gradient. (C) Lysates from wild-type or MurG-Dendra2-expressing *M. smegmatis* were
fractionated as in (B) and separated by SDS-PAGE. Top, in-gel fluorescence shows MurG-
Dendra2 association with the IMD. Treatment with benzyl alcohol (BA) redistributed the protein
across the fractions. Bottom, wild-type *M. smegmatis* membrane fractions were incubated with
Bocillin-FL prior to SDS-PAGE. Labeled PBPs are enriched in PM-CW.

Fig. 2. Lipid II is synthesized in the IMD and trafficked to the PM-CW. (A) Left, conventional microscopy of *M. smegmatis* coexpressing PonA1-mRFP and MurG-Dendra2 treated +/- benzyl alcohol (BA). Right, fluorescence distribution of the fusion proteins. a.u., arbitrary units. (B) Top, detection scheme for lipid-linked peptidoglycan (PG) precursors from membrane fractions. Bottom, metabolic labeling scheme of mycobacterial cell wall synthesis (García-Heredia et al., 2018). (C) PG precursors are labeled as in (B), top. The labeled precursors are in the IMD and PM-CW of wild-type *M. smegmatis* but accumulate in the IMD upon MurJ depletion (García-Heredia et al., 2018). (D) *M. smegmatis* expressing MurG-Dendra2 were incubated with alkDADA. Surface-exposed alkynes on fixed cells were detected by CuAAC (García-Heredia et al., 2018). Cells imaged by structured illumination microscopy (SIM-E).

Fig. 3. Membrane perturbations disrupt peptidoglycan biogenesis in *M. smegmatis* and phylogenetically-distant bacilli. (A) Lysates from wild-type *M. smegmatis* treated +/- benzyl alcohol (BA) were sedimented in a sucrose density gradient. (B) Top, wild-type *M. smegmatis* was incubated or not with benzyl alcohol or dibucaine, then labeled with alkDADA. Bottom, the distribution of peptidoglycan labeling from wild-type *M. smegmatis* that was incubated with BA or dibucaine (DB) for the indicated time was quantitated as in Fig. 2A, except that signal intensity was not normalized. (C) DivIVA-eGFP-ID *M. smegmatis* was either treated with benzyl alcohol, depleted of DivIVA, or both, and the peptidoglycan precursors from whole cells were biotinylated as in Fig. 2B, top. (D) Phylogenetic tree constructed with 16S rDNA sequences (rate of mutation not considered). Taxonomic groups matched with colors to their levels with only diverging points shown. Shapes and growth modes illustrated for select species. (E) Left, different bacteria treated +/- benzyl alcohol followed by alkDADA incubation. Arrowheads highlight irregular patches of peptidoglycan. Insets are magnified. Where applicable, *E. coli* was preincubated with A22. Right, bacteria were treated with benzyl alcohol, translation-inhibiting kanamycin, or peptidoglycan-acting phosphomycin or ampicillin then labeled as in (B) and analyzed by flow cytometry (see Methods). Median fluorescence intensity (MFI) values were normalized to untreated controls and experiments were performed 3-9 times in triplicate. Error bars, +/- standard deviation of biological replicates. *, $p < 0.05$; **, $p < 0.005$; ***, $p < 0.0005$; ****, $p < 0.00005$, ratio paired t-tests and one-way ANOVA with Dunnet´s test for non-normalized MFI of biological replicates. Scale bars, 5 μm.

Fig. 4. Plasma membrane domains are maintained by DivIVA and the cell wall. (A) Lysates from MurG-Dendra2- or DivIVA-eGFP-ID-expressing *M. smegmatis* that had been depleted of DivIVA (Meniche et al., 2014) or spheroplasted (Melzer et al., 2018), were sedimented in a density gradient. (B). DivIVA was depleted or not in *M. smegmatis* expressing both DivIVA-eGFP-ID and mCherry-GlfT2. Normally, DivIVA localizes at the polar tip and mCherry-GlfT2 concentrates proximal to DivIVA. Depletion of DivIVA delocalizes mCherry-GlfT2. *M. smegmatis* expressing MurG-Dendra2 (C) or coexpressing mCherry-GlfT2 and DivIVA-eGFP-ID (D) were spheroplasted or not and imaged. In spheroplasted cells, the IMD-associated proteins distribute along the cell periphery. Arrows mark spheroplasts outside and within insets, which have increased size and brightness. (E) Model for self-organization of plasma membrane and cell wall in *M. smegmatis*. Brown line indicates the cell wall. Short brown lines perpendicular to the membrane and cell wall indicate that the cell wall is likely to be physically connected to the membrane in the PM-CW regions (Morita et al., 2005). All scale bars, 5 μm.

Methods

Bacterial strains and growth conditions

Mycobacterium smegmatis mc^{2}155 was grown in Middlebrook 7H9 growth medium (BD Difco, Franklin Lakes, NJ USA) supplemented with 0.4% (v/v) glycerol, 0.05% (v/v) Tween-80 (Sigma-Aldrich, St. Louis, MO USA) and 10% ADC (albumin-dextrose-catalase) as well as apramycin (50 µg/mL), kanamycin (25 µg/mL; Sigma-Aldrich, St. Louis, MO USA) and hygromycin (50 µg/mL) where appropriate. *Staphylococcus aureus* ATCC BA-1718 was grown in BHI (BD Difco, Franklyn Lakes NJ, USA); *Escherichia coli* K12 and *Bacillus subtilis* ZB307, in LB (VWR, Radnor PA, USA); *Caulobacter crescentus* NA 1000, in peptone yeast extract (BD Difco, Franklyn Lakes NJ, USA); and *Lactococcus lactis* NRRL B633, in MRS (Oxoid, Basingstoke, Hampshire UK). All bacteria were grown shaking at 37ºC with the exception of *C. crescentus*, which was incubated at 30ºC. See Table S3 for more information.

Mutant strain construction

To test the function of PimE-GFP-FLAG fusion, an expression vector for PimE-GFP-FLAG (pYAB186) (Hayashi et al., 2016) was electroporated into a *pimE* deletion mutant (Morita et al., 2006). Three independent colonies were picked for the phenotypic complementation of AcPIM6 biosynthetic defects. Lipid purification and analysis were performed as described previously (Morita et al., 2006).

The *murG* gene was amplified by PCR from *M. smegmatis* mc^{2}155 genomic DNA by PCR,

excluding the stop codon, and was inserted into pMSR in-frame with mycobacterial codon-

optimized Dendra2 using In-Fusion cloning (Takara Bio, Mountain View, CA USA). This

construct was transformed by electroporation into *M. smegmatis* mc^{2}155, where it integrates at

the L5 *attB* site, and was selected by apramycin treatment. Constitutive expression of the MurG-

Dendra2 fusion was achieved through the Psmyc promoter (GenBank: AF395207.1). The

plasmid construct was validated by Sanger sequencing.

To replace the endogenous *glfT2* gene with a gene encoding HA-mCherry-GlfT2 in the DivIVA-

eGFP-ID strain, we electroporated pMUM052 (Hayashi et al., 2016) into DivIVA-eGFP-ID *M.*

smegmatis, and positive clones were isolated using hygromycin resistance marker and SacB-

dependent sucrose sensitivity. Correct replacement of the *glfT2* gene was confirmed by PCR.

Generation of spheroplasts

To generate spheroplasts, we followed a previous protocol (Melzer et al., 2018). Briefly, wild-

type, MurG-Dendra2-expressing or mCherry-GlfT2/DivIVA-eGFP-ID-coexpressing *M.*

smegmatis was grown until log phase. Glycine (1.2% w/v final concentration) was added and the

culture was incubated for 24 h at 37ºC with shaking. Afterwards, the cells were washed with

sucrose-MgCl$_2$-maleic acid (SMM) buffer (pH 6.8) and harvested by centrifugation (4000 x *g* for

5 min). The pellet was resuspended in 7H9 medium where the water was replaced with SMM

buffer; the medium also was supplemented with glycine (1.2% final concentration) and lysozyme

(50 µg/mL final concentration). Bacteria were incubated another 24 h at 37ºC with shaking, and

then spheroplasts where either imaged by conventional fluorescence microscopy or lysed by

nitrogen cavitation immediately for subsequent biochemical analysis.

Membrane fractionation

Log phase *M. smegmatis* (that, where applicable, were untreated, treated with benzyl alcohol,

depleted for DivIVA or spheroplasted; see Table S1) were harvested by centrifugation and

washed in PBS + 0.05% Tween-80 (PBST). One gram of wet pellet was resuspended in 5 mL of

lysis buffer containing 25 mM HEPES (pH 7.4), 20% (w/v) sucrose, 2 mM EGTA, and a

protease inhibitor cocktail (Thermo Fisher Scientific, Waltham, MA USA) as described (Morita

et al., 2005). Bacteria were lysed using nitrogen cavitation at 2000 psi for 30 min three times.

Cell lysates were centrifuged at 3220 x g for 10 min at 4ºC twice to remove unlysed cells prior to

loading on a 20-50% sucrose gradient. Membrane-containing fractions were collected in 1 mL

after ultracentrifugation at 35,000 rpm on an SW-40 rotor (Beckman, Brea, CA USA) for 6 hours

at 4ºC and stored at -80ºC prior to analysis.

Detection of proteins in membrane fractions

MurG-Dendra2 and penicillin-binding proteins (PBPs) were detected by in-gel fluorescence. For

MurG-Dendra2, membrane fractions were incubated with an equal volume of 2x loading buffer

and then separated by SDS-PAGE on a 12% polyacrylamide gel. To detect PBPs, 50 µg total

protein from wild-type *M. smegmatis* membrane fractions were incubated with 40 µM of

Bocillin-FL (Thermo Fisher Scientific, Waltham, MA USA) for 30 min in the dark at 37ºC. An

equal volume of 2x loading buffer was then added and the mixture was boiled for 3 min at 95ºC

and then incubated on ice for 30 min. Membrane mixtures were separated on a 12%

polyacrylamide gel. Gels were washed in distilled water and imaged using an ImageQuant LAS

4000mini (GE Healthcare, Chicago, IL USA).

MurJ, PimB´ and MptA were detected by immunoblot. Briefly, cell lysate or membrane fraction

proteins were separated by SDS-PAGE on a 12% polyacrylamide gel and transferred to a PVDF

membrane. The membrane was blocked with 3% milk in PBS + 0.05% Tween-80 (PBST) and

then incubated overnight with primary antibodies (monoclonal mouse anti-FLAG [to detect

MurJ; Sigma], and polyclonal rabbit anti-PimB´ or anti-MptA antibodies (Sena et al., 2010).

Antibodies were detected with appropriate secondary antibodies conjugated to horseradish

peroxidase (GE Healthcare, Chicago, Il USA). Membranes were rinsed in PBS + 0.05% Tween-

20 and visualized by ECL in an ImageQuant LAS 4000mini (GE Healthcare, Chicago, IL USA)

as above.

Cell envelope labeling

Chemical probes used in this work include alkDADA (Liechti et al., 2014), N-AlkTMM and O-

AlkTMM (Foley et al., 2016) and FM4-64 FX (Invitrogen, Carlsbad, CA USA). AlkDADA was

synthesized by the Einstein Chemical Biology Core and the TMM probes were kind gifts of Dr.

Ben Swarts. Unless otherwise indicated, mid-log *M. smegmatis* or, where applicable, *B. subtilis,*

S. aureus, E. coli, L. lactis or *C. crescentus* were labeled with 2 mM alkDADA, 250 µM N-

AlkTMM, 50 µM O-AlkTMM or 5 µg/mL FM4-64FX for 15 min or 5 min in the case of *B.*

subtilis. Unless otherwise indicated and where applicable, the bacteria were pre-incubated in the

presence or absence of freshly-prepared chemicals (antibiotics or benzyl alcohol) before being

subjected to the probes (see Table S1 and S2). Cells were pelleted by centrifugation, washed in

PBST containing 0.01% BSA (PBSTB) and fixed for 10 min in 2% formaldehyde at room

temperature. For alkDADA and O-AlkTMM, cells were further washed twice with PBSTB and

subjected to CuAAC as described (García-Heredia et al., 2018). Unless otherwise specified,

picolyl azide-Cy3 was used in Fig. 2D; picolyl azide carboxyrhodamine 110 was used in Figs.

3B, 3D and Figure 2–figure supplement 2, Figure 3–figure supplement 1, and Figure 3–figure

supplement 3. Bacteria were then washed twice in PBSTB, once in PBST, and imaged (described

below) or subjected to flow cytometry (BD DUAL LSRFortessa, UMass Amherst Flow

Cytometry Core).

Microscopy and image analysis

Bacteria were imaged on agar pads by either conventional fluorescence microscopy (Nikon

Eclipse E600, Nikon Eclipse Ti or Zeiss Axioscope A1 with 100x objectives) or structured

illumination microscopy (Nikon SIM-E/A1R with SR Apo TIRF 100x objective, UMass

Amherst Light Microscopy Core).

To obtain the fluorescence intensity plots, the subcellular distribution of fluorescence was

quantitated from images obtained by conventional fluorescence microscopy. The images were

processed using Fiji and Oufti (Paintdakhi et al., 2016; Schindelin et al., 2012) as described

(García-Heredia et al., 2018). The signal was normalized to length and total fluorescence

intensity of the cell. Cells were oriented such that the brighter pole is on the right-hand of the

graph. The intensity plots from Fig. 2A were made from 42<n<56 cells; for Fig. 3B, from

14<n<70 cells.

Evaluation of membrane-bound peptidoglycan precursors

Wild-type or MurJ-depleted *M. smegmatis* (Gee et al., 2012) were grown to mid-log phase and

membrane fractions were isolated as above. Precursors were extracted from each membrane

fraction similar to previous publications (García-Heredia et al., 2018; Qiao et al., 2014; Qiao et

al., 2017). Briefly, glacial acetic acid was added to 500 µL of fractionated lysate to a final

volume of 1%. The sample was then transferred into a vial containing 500 µL of chloroform and

1 mL of methanol and left at room temperature for 1-2 hours with occasional vortexing. The

mixture was centrifuged at 21,000x *g* for 10 min and the supernatant was transferred into a vial

containing 500 µL of 1% glacial acetic acid (in water) and 500 µL chloroform, and vortexed for

1 min. The mixture was separated by centrifugation (900x *g* for 1 min at room temperature) and

the organic phase was collected. Where applicable, the white interface was reextracted to recover

additional organic material. The organic phase was dried under a nitrogen stream and the dried

lipids were resuspended in 12 µL of DMSO. D-amino acid-containing, lipid-linked

peptidoglycan precursors were biotinylated by subjecting organic extracts to an *in vitro* PBP4-

mediated exchange reaction with biotin-D-lysine (Chem-Impex International, Wood Dale, IL

USA; reagent was deprotected first) as described (Qiao et al., 2014). The products were

separated by SDS-PAGE on an 18%, polyacrylamide gels then transferred to a PVDF membrane,

blotted with streptavidin-HRP (diluted 1:10,000; Thermo Fisher Scientific, Waltham, MA USA)

and visualized by ECL as above.

Membrane glycolipid analysis

Sucrose density gradient fractions from wild-type *M. smegmatis* +/- 1 hour of 100 mM benzyl

alcohol treatment or DivIVA-eGFP-ID with DivIVA depleted or not (Meniche et al., 2014) were

subjected to lipid purification and analysis as previously described (Morita et al., 2005).

Phylogenetic analysis

The phylogenetic tree was made in Adobe Illustrator (version 23.0.3) based on a phylogenetic

tree generated with Mega (version 7.0.26; Kumar et al., 2018). Briefly, 16S rDNA sequences

were obtained from NCBI (see Table S2) and aligned using ClustalW. The phylogenetic tree was

generated using the Timura-Ney model with Gamma distribution and Bootstrap method (1000

replications). The taxonomic information was verified with the Interagency Taxonomic

Information System (available online https://www.itis.gov/).

Figure supplements

Figure 1–figure supplement 1. MurG-Dendra2 and PimE-GFP are functional, and MurJ can be
depleted. (A) MurG-Dendra2 expression rescues depletion of endogenous protein. MurG
depletion strain (Meniche et al., 2014) was transformed with a plasmid containing *murG-*
dendra2 and grown +/- 50 ng/mL anhydrotetracycline (ATC) to induce MurG-ID degradation.
(B) PimE-GFP is functional. PimE is a mannosyltransferase involved in phosphatidylinositol
mannoside (PIM) biosynthesis, converting AcPIM4 to more polar PIMs. Δ*pimE* was
complemented with the indicated expression vectors. Glycolipids were extracted, purified, and
separated by thin-layer chromatography. PIMs were visualized by orcinol staining. Similar to
PimE-FLAG (Morita et al., 2006), PimE-GFP can restore the production of AcPIM6. (C) MurJ is
depleted upon treatment with anhydrotetracycline (ATC). Depletion strain (MurJ-ID; carrying
FLAG tag) was grown +/- ATC. Insert, immunoblot of lysates showing that MurJ is degraded
after 8 hours of ATC treatment as reported (Gee et al., 2012).

Figure 1 –figure supplement 2. Western blot analysis of the IMD and the PM-CW membrane fractions separated by sucrose density sedimentation. Antibodies against PimB' and MptA or MptC were used to detect the IMD and PM-CW, respectively (Hayashi et al., 2016; Sena et al., 2010). Analyzed fractions shown here are the same fractions of: A) MurG-Dendra2-expressing cells as shown in Fig. 1C; B) wild-type cells as shown in Fig. 1C and Fig. 2C; C) MurJ-depleted cells as shown in Fig. 2C; D) wild-type cells treated with benzyl alcohol (BA) as shown in Fig. 1C; E) MurG-Dendra2-expressing cells treated with benzyl alcohol as shown in Fig. 1C; F) and G) are DivIVA-eGFP-ID where DivIVA was depleted or not depleted as in Fig 4A. Samples from F) and G) were concentrated 10-fold by precipitating proteins in chloroform and water. PimB' appears as ~45 kDa band in the IMD fractions but disappears upon BA treatment or DivIVA depletion. We do not yet know the reason for this disappearance. The band at ~37 kDa, which is visible especially in the cytoplasmic fractions of BA-treated cells is a non-specific protein (Hayashi et al., 2016). MptA and MptC appear as ~42 kDa band in the PM-CW fractions, which became fainter after BA treatment or depletion of DivIVA.

Figure 1–figure supplement 3. Visualization of PonA1 using Bocillin-FL. To assess specificity of the fluorescent signal, membrane fractions were pre-incubated +/- 16 µg/mL ampicillin and 5 µg/mL of the β-lactamase inhibitor clavulanate (García-Heredia et al., 2018) for 30 min at room temperature. 100 µg of total membrane proteins were incubated with 40 µM of Bocillin-FL for 30 min, then separated by SDS-PAGE (12% gel) and visualized by in-gel fluorescence to detect PBPs. WT, wild-type *M. smegmatis*; p$_{tet}$*ponA1* expressed (On) or depleted (Off) with addition of anhydrotetracycline (ATC) as described (Hett et al., 2010).

663 **Figure 2–figure supplement 1**. MurG-Dendra2 and PonA1-mRFP colocalize with IMD and
664 PM-CW reporters, respectively. *M. smegmatis* coexpressing MurG-Dendra2 and mCherry-GlfT2
665 or PonA1-mRFP and PimE-GFP were imaged by SIM-E. Scale bars, 5 μm. Graph, the
666 fluorescence intensity profiles of the proteins were quantitated as in Fig 2A from images
667 captured with conventional microscopy.

Figure 2–figure supplement 2. Depletion of MurJ alters amount and location of nascent peptidoglycan. Wild-type (left) or MurJ-ID (depletion strain; right) *M. smegmatis* in early log phase were treated +/ anhydrotetracycline (ATC) then incubated with alkDADA. Bacteria were washed, fixed, subjected to CuAAC and imaged by conventional fluorescence microscopy. Scale bar, 5 μm.

A

B

Figure 3–figure supplement 1. (A) Benzyl alcohol alters FM4-64FX distribution. Wild-type *M.*
smegmatis was labeled with FM4-64FX, then washed and incubated with benzyl alcohol. Cells
were then imaged with SIM-E. (B) Benzyl alcohol halts cell elongation but does not otherwise
impact the localization of mycomembrane probes. Wild-type *M. smegmatis* was incubated with
O-AlkTMM (left) or N-AlkTMM (right) to label covalent or noncovalent mycolates,
respectively, then washed and subjected or not to benzyl alcohol for 1 hour. Bacteria were
washed, fixed, subjected to CuAAC and imaged by conventional fluorescence microscopy. As a
control, bacteria were not treated with benzyl alcohol for 1 hour, *i.e.* fixed immediately after
probe incubation (bottom panel). All scale bars, 5 µm.

Figure 3–figure supplement 2. Benzyl alcohol and depletion of DivIVA affect the distribution of membrane glycolipids. Wild-type or DivIVA-eGFP-ID *M. smegmatis* were treated +/- benzyl alcohol or ATC, respectively. PIMs were visualized by orcinol staining as in Figure 1–figure supplement 1. AcPIM2 and AcPIM6 present during normal growth in the IMD and are diminished upon benzyl alcohol treatment. Depletion of DivIVA is accompanied by an enrichment of PIM2 species and depletion of PIM6 species. This change in lipids is currently under investigation.

Figure 3–figure supplement 3. *M. smegmatis* survives benzyl alcohol (BA) treatment. (A) *M. smegmatis* was treated with benzyl alcohol then washed in PBS. Ten-fold serial dilutions were spotted on LB agar. Shown are cells immediately after addition of benzyl alcohol (T_0) and after 1 hour of exposure. (B) *M. smegmatis* were treated with benzyl alcohol for 30 or 60 min, washed and incubated for 2 hours in fresh 7H9. Cells were then labeled with alkDADA for 15 min and then fixed and alkynes were detected by CuAAC to reveal active peptidoglycan metabolism. Scale bar 5 µm.

Figure 3–figure supplement 4. Dibucaine impacts the localization of MurG and GlfT2. *M. smegmatis* coexpressing the IMD-associated MurG-Dendra2 and mCherry-GlfT2 was treated or not with dibucaine. Images from conventional microscopy show these proteins lose their polar enrichment when treated with dibucaine. Scale bar, 5 μm. Right, the fluorescence intensity profiles of the fusion proteins were determined as in Figure 2A.

Figure 4–figure supplement 1. Cell wall is required for partitioning of MurG. *M. smegmatis* expressing MurG-Dendra2 were spheroplasted as in Fig. 4A and then lysed by nitrogen cavitation. The crude lysates were separated in a density gradient and blotted in a polyacrylamide gel. MurG-Dendra2 distributes widely along the membrane fractions consistent with the broad distribution of cellular material across the density gradient (see Fig. 4A), which contrasts with its normal partitioning (see Fig. 1C).

772 **Supplementary information**

Table S1. Incubation conditions		
Bacterium	Antibiotic or compound	Incubation conditions
M. smegmatis	100 mM benzyl alcohol	60 min at 37ºC with shaking
	50 mM dibucaine	180 min at 37ºC with shaking
DivIVA-eGFP-ID	50 ng/mL ATC	8 h, 14 to 16 h
MurJ-ID		8h
B. subtilis	24 µg/mL kanamycin	10 min at 37°C with shaking
	50 µg/mL phosphomycin	
	50 mM benzyl alcohol	
C. crescentus	5 µg/mL kanamycin	30 min at 30°C with shaking
	50 µg/mL phosphomycin	
	50 mM benzyl alcohol	
E. coli	20 µg/mL A22	90 min at 37°C with shaking. Cultures were back-diluted after 40 min of incubation to an OD_{600} ~0.3-0.5 in prewarmed medium + A22 and incubated until completion of 90 min
	16 µg/mL kanamycin	30 min at 37ºC with shaking
	50 µg/mL phosphomycin	
	50 mM benzyl alcohol	
S. aureus and *L. lactis*	50µg/mL phosphomycin	30 min at 37ºC with shaking
	20 µg/mL kanamycin	
	50 mM benzyl alcohol	
	50 µg/mL ampicillin (with 5 µg/mL clavulanate)	

Table S2. Incubation conditions of probes to metabolically label the cell envelope

Bacterium	Probe	Labeling conditions
M. smegmatis	alkDADA N-Alk-TMM O-Alk-TMM	Unless otherwise indicated, 15 min with shacking at 37ºC
E. coli *B. subtilis* *S. aureus* *L. lactis*	alkDADA	Unless otherwise indicated, 5 min with shacking at 37ºC
C. crescentus		Unless otherwise indicated, 5 min with shacking at 30ºC

Table S3. NCBI Accession numbers from 16S rDNA

Bacterium	NCBI accession number
M. smegmatis	AB305022.1
B. subtilis	NR_112116.2
S. aureus	NR_118997.2
L. lactis	FJ348447.1
E. coli	NR_024570.1
C. crescentus	NR_037099.1

 Table S4. Key Resources

Reagent type	Designation	Source or reference	Additional information
Strain (*M. smegmatis* mc^{2}155)	*M. smegmatis*	NC_008596 in GenBank	Wild-type *M. smegmatis*
Strain (*M. smegmatis*)	MurG-Dendra2	This study	The mutant was generated as described in supplementary material and methods.
Strain (*M. smegmatis*)	mCherry-GlfT2	(Hayashi et al., 2016)	See reference for details.
Strain (*M. smegmatis*)	PonA1-mRFP	(Kieser et al., 2015)	Obtained from Dr. Eric Rubin (Harvard SPH) and Dr. Hesper Rego (Yale Med)
Strain (*M. smegmatis*)	PimE-GFP	This study	The mutant was generated as described in supplementary material and methods.
Strain (*M. smegmatis*)	MurG-ID depletion strain	(Meniche et al., 2014)	Obtained from Dr. Chris Sassetti (U Mass Med)
Strain (*M. smegmatis*)	MurJ-ID (MviN) depletion strain	(Gee et al., 2012)	Obtained from Dr. Chris Sassetti (U Mass Med)
Strain (*M. smegmatis*)	p$_{tet}$*ponA1*	(Hett et al., 2010)	Obtained from Dr. Eric Rubin (Harvard SPH)
Strain (*M. smegmatis*)	DivIVA-eGFP-ID	(Meniche et al., 2014)	Obtained from Dr. Chris Sassetti (U Mass Med)
Strain (*M. smegmatis*)	mCherry-GlfT2/DivIVA-eGFP-ID	This study	
Strain (*B. subtilis* JH642)	*B. subtilis*	NZ_CP007800 in GeneBank	
Strain (*C. crescentus*)	*C. crescentus*	NA 1000	Obtained from Dr. Peter Chien (U Mass Amherst)
Strain (*E. coli* K12)	*E. coli K12*	MG1655	
Strain (*S. aureus*)	*S. aureus*	ATCC BA-1718	Obtained from Dr. Thai Thayumanavan (U Mass Amherst)
Strain (*L. lactis*)	*L. lactis lactis*	NRRL B633	
Chemical compound	alkyne-D-alanine-D-alanine (alkDADA or EDA-DA)	(Liechti et al., 2014)	Synthesized by the Chemical Synthesis Core Facility at Albert Einstein College of Medicine (NY, USA) following the referenced protocols
Chemical compound	O-alkyne-trehalose monomycolate (O-AlkTMM)	(Foley et al., 2016)	Obtained from Dr. Benjamin Swarts (Central Michigan University)

Chemical compound	N-alkyne-trehalose monomycolate (N-AlkTMM)	(Foley et al., 2016)	Obtained from Dr. Benjamin Swarts (Central Michigan University)
Software, algorithm	MATLAB codes	(García-Heredia et al., 2018)	Scripts designed for MATLAB to analyze the fluorescence profiles along a cell body from data collected in Oufti[28].
Chemical compound	Fmoc-D-Lys(biotinyl)-OH BDL precursor	Chem-Impex International (Wood Dale, Il USA)	Deprotected as described in[13] to yield BDL
Chemical compound	A22 (S-3,4-Dichlorobenzylisothiourea hydrochloride)	Sigma-Aldrich, St. Louis, MO USA	Dissolved in water and kept at -20ºC
Recombinant DNA reagent	PBP4 plasmid	(Qiao et al., 2014)	Obtained from Dr. Suzanne Walker (Harvard Med)

815