Membrane compartmentalization of mycobacterial peptidoglycan synthesis

Alam García-Heredia[1], Julia Puffal[2], Julius Judd[3] *, Todd A. Gray[3,4], Yasu S. Morita[1,2, †],

M. Sloan Siegrist[1,2, †].

Affiliations:

[1] Molecular and Cellular Biology Graduate Program, University of Massachusetts, Amherst, MA 01003, USA.

[2] Department of Microbiology, University of Massachusetts, Amherst, MA 01003, USA.

[3] Division of Genetics, Wadsworth Center, New York State Department of Health, Albany, NY 12208, USA.

[4] Department of Biomedical Sciences, University at Albany, Albany, NY 12208, USA.

* Current address: Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14853, USA.

† Corresponding author. Email: siegrist@umass.edu (M.S.S.); ymorita@microbio.umass.edu (Y.S.M.).

Abstract: Cell wall peptidoglycan, a mesh of polysaccharides crosslinked by short peptides, encases the bacterial cell and protects it from turgor pressure lysis. Peptidoglycan synthesis is an effective antibiotic target. Assembly of the biopolymer occurs in close association with the plasma membrane, but higher order organization of the process has not been described. In mycobacteria, intracellular membrane domains comprise biochemically and spatially distinct regions within the conventional plasma membrane. We find that lipid-linked peptidoglycan precursors are made in these domains and then trafficked to the conventional plasma membrane for insertion into the

cell wall. Disorganization of the membrane rapidly delocalizes and then halts

peptidoglycan assembly. Our data show that membrane compartmentalization is an

essential feature of mycobacterial cell wall biogenesis.

Main Text: Many antibiotics target peptidoglycan synthesis, a well-conserved pathway

that spans the cytoplasm, plasma membrane and periplasm. The polyprenol-linked,

disaccharide-pentapeptide monomer lipid II is completed by the glycosyltransferase

MurG in the inner leaflet of the plasma membrane (Fig. 1A). Lipid II is then flipped to the

outer leaflet by MurJ and integrated into the cell wall by membrane-bound

transglycosylases and transpeptidases from the penicillin-binding protein (PBP) and

shape, elongation, division, and sporulation (SEDS) families (*1-4*).

The plasma membrane is a heterogeneous mixture of lipids and proteins.

Mycobacteria, for example, have **i**ntracellular **m**embrane **d**omains (IMD, formerly called

the PMf (*5*) for **p**lasma **m**embrane **f**ree of cell wall) that are separable from the

conventional plasma membrane (designated the PM-CW, for **p**lasma **m**embrane tightly

associated with **c**ell **w**all) by sucrose density gradient fractionation. The proteome and

lipidome of IMD are distinct from PM-CW (*5, 6*). While PM-CW-resident proteins localize

along the perimeter of live mycobacteria, IMD-resident proteins localize along sidewall

but are enriched adjacent to sites of polar cell elongation (*6, 7*).

Our proteomics analysis indicated that MurG is present in the IMD while

sequentially-acting PBPs preferentially associate with the PM-CW (*6*). We also

observed *in situ* that the subpolar enrichment of MurG-RFP resembles that of the

validated IMD marker mCherry-GlfT2 or GlfT2-GFP (*6, 8*) but that nascent cell wall at

the mycobacterial poles primarily abuts rather than colocalizes with mCherry-GlfT2 (7).

These observations suggest that lipid II synthesis is biochemically and spatially

segregated from the subsequent steps of cell wall assembly (Fig. 1A).

We expressed a functional MurG-Dendra2 fusion in *Mycobacterium smegmatis*

(Fig. S1) and assayed its distribution in membrane fractions that had been separated by

density gradient (Fig. 1B). The fusion to MurG, a peripheral membrane protein, was

enriched in both the cytoplasmic and IMD membrane fractions (Fig. 1C; Fig. S2),

recapitulating the association predicted for the native protein (*6*). In intact cells, polar

enrichment of MurG-Dendra2 was coincident with that of mCherry-GlfT2 (Fig. 1D). The

spatial relationship between the proteins was similar to that of MurG-RFP and GlfT2-

GFP (*8*), suggesting that it is independent of the fluorescent protein tag.

The association of MurG with the IMD, but not with the PM-CW, implied that the

membrane domain is the site of lipid II synthesis. We refined an *in vitro* D-amino acid

exchange assay to detect lipid-linked peptidoglycan precursors from *M. smegmatis*

membrane fractions (Fig. 2A; *9, 10*). In wildtype cells, we detected biotinylated

molecules in both the IMD and PM-CW (Fig. 2B; Fig. S2). We hypothesized that the

labeled species comprised precursors in the inner leaflet of the plasma membrane as

well as lipid II that had been flipped to the outer leaflet. We, and others, have shown

that depletion of MurJ results in an accumulation of biotinylated precursors (*10, 11*; Fig.

S3). By performing the D-amino acid exchange reaction on membrane fractions, we

found that precursors accumulate in the IMD (Fig. 2B; Fig. S2). These results suggest

that lipid II is made in the IMD and transferred to the PM-CW in a MurJ-dependent

manner.

PBPs and SEDS proteins incorporate lipid II into peptidoglycan. Given the trafficking of precursors from the IMD to the PM-CW (Fig. 2B) and the association of cell wall fragments specifically with the PM-CW (*5*), we hypothesized that extracellular, peptidoglycan-acting enzymes function in the PM-CW. While our proteomics did not detect SEDS proteins, our PM-CW dataset was enriched for all of the known mycobacterial PBPs (*6*). Fluorescent derivatives of β-lactam antibiotics such as Bocillin-FL bind covalently to PBPs and can be used to image transpeptidase-active enzymes in both polyacrylamide gels and intact cells. We incubated membrane fractions from wildtype *M. smegmatis* with Bocillin-FL and identified fluorescent proteins in the PM-CW but not the IMD (Fig. 2B; Fig. S2). As expected for PBPs, the signal from these bands was diminished by pre-treatment with the β-lactam ampicillin (Fig. S4). We focused on characterizing PonA1, a bifunctional transglycosylase/transpeptidase that is essential for *M. smegmatis* growth (*12, 13*). Depletion of PonA1 (*12*) resulted in the loss of the highest molecular fluorescent band (Fig. S4), confirming that this protein is present and active specifically in the PM-CW (Fig. 2B). We next expressed a functional PonA1-mRFP fusion in *Mycobacterium smegmatis* (*13*) and found that it was more evenly distributed around the cell perimeter than MurG-Dendra2, and in a manner similar to the functional PM-CW marker PimE-GFP (Fig. 2C, Fig. S5; *6*). Together, our data show that MurG and PonA1 occupy spatially distinct compartments along the pathway of peptidoglycan synthesis.

Based on our biochemical data, we hypothesized that lipid II incorporation into the cell wall is laterally segregated from its synthesis. We previously showed that alkynyl and azido D-amino acid dipeptides (*14*) incorporate into lipid-linked

peptidoglycan precursors in *M. smegmatis* (*10*) and that metabolic labeling with alkynyl

dipeptide (alkDADA or EDA-DA) is most intense in regions adjacent to the IMD marker

mCherry-GlfT2 (*7*). We labeled MurG-Dendra2-expressing *M. smegmatis* with alkDADA

for ~1% of generation time and detected the presence of the alkyne by copper-

catalyzed azide-alkyne cycloaddition (CuAAC; Fig 2D; *10*). To tune our detection for

extracellular alkynes present in lipid II and newly-polymerized cell wall, we selected

picolyl azide-Cy3 as our label because the localized charge on the sulfonated cyanine

dye confers poor membrane permeability (*15*). Using this optimized protocol, we

observed nascent peptidoglycan deposition at the polar tip, whereas MurG-Dendra2

was more posterior (Fig. 2D). Our data support a model in which lipid II synthesis is

laterally and biochemically partitioned from the ensuing steps of peptidoglycan

assembly.

In the Gram-positive bacterium *Bacillus subtilis*, MurG associates with **r**egions of

increased **f**luidity (RIFs) in the plasma membrane that are marked by the accumulation

of certain lipophilic fluorescent dyes (*16*). Benzyl alcohol has been used to disperse

plasma membrane domains in plant, animal and bacterial cells, including *B. subtilis*

RIFs (*16-18*). Mycobacteria are also Gram-positive but have a second 'myco'

membrane that is covalently attached to the cell wall. We found that benzyl alcohol did

not alter labeling by NalkTMM or OalkTMM (Fig. S6), which respectively incorporate into

the noncovalent and covalent mycolates of the mycomembrane (*19*). It did, however,

alter the distribution of FM4-64 (Fig. S7), a dye used previously to label the plasma

membrane (*20*), and altered glycolipid abundance in the IMD (Fig. S8). MurG-Dendra2

was also notably less enriched in the IMD following benzyl alcohol treatment (Fig. 1C)

and, in live cells, at the cell poles (Fig. 3A). By contrast benzyl alcohol produced subtle

changes in the distribution of active PBPs (Fig. 2B), although PonA1 shifted toward the

poles in live cells (Fig. 3A). Disruption of plasma membrane organization by benzyl

alcohol was accompanied by delocalization of cell wall assembly within 5 min (Fig. S9)

as well as an overall reduction in synthesis (Fig. 3B, Figs. S9) and halt in cell elongation

(Fig. S6). The phenotypes were reversible: we recovered viable, peptidoglycan-

synthesizing cells following benzyl alcohol washout (Figs. S10). These data suggest

that membrane partitioning is an essential feature of both peptidoglycan synthesis and

cell growth in *M. smegmatis*.

We find that benzyl alcohol also inhibits cell wall assembly in *B. subtilis* (Fig. 3C),

demonstrating that plasma membrane architecture may optimize peptidoglycan

biogenesis in bacterial phyla with divergent cell envelope structures and modes of

growth. Moreover, our previous work suggests that the biosynthetic pathways for

phosphatidylethanolamine, a major phospholipid of mycobacterial plasma membrane,

phosphatidylinositol mannoside, a cell envelope glycolipid, and menaquinone, the

primary lipid electron carrier of the mycobacterial respiratory chain, are partitioned

across the IMD and PM-CW (*5, 6, 21*). In the absence of a standard set of membrane-

bound organelles, plasma membrane compartmentalization may be a general bacterial

strategy for organizing pathways with lipid-linked intermediates and enzymes.

References and Notes:

1.   C. L. Gee *et al.*, A phosphorylated pseudokinase complex controls cell wall

        synthesis in mycobacteria. *Sci Signal* **5**, ra7 (2012).

2.   A. J. Meeske *et al.*, SEDS proteins are a widespread family of bacterial cell wall

polymerases. *Nature* **537**, 634-638 (2016).

3.   K. Emami *et al.*, Corrigendum: RodA as the missing glycosyltransferase in

Bacillus subtilis and antibiotic discovery for the peptidoglycan polymerase

pathway. *Nature Microbiology* **2**, 17019 (2017).

4.   D. Arora *et al.*, The transpeptidase PbpA and non-canonical transglycosylase

RodA of Mycobacterium tuberculosis play important roles in regulating bacterial

cell lengths. *J Biol Chem*, jbc. M117. 811190 (2018).

5.   Y. S. Morita *et al.*, Compartmentalization of lipid biosynthesis in mycobacteria. *J*

Biol Chem **280**, 21645-21652 (2005).

6.   J. M. Hayashi *et al.*, Spatially distinct and metabolically active membrane domain

in mycobacteria. *Proc Natl Acad Sci U S A* **113**, 5400-5405 (2016).

7.   J. M. Hayashi *et al.*, Stress-Induced Reorganization of the Mycobacterial

Membrane Domain. *mBio* **9**, (2018).

8.   X. Meniche *et al.*, Subpolar addition of new cell wall is directed by DivIVA in

mycobacteria. *Proc Natl Acad Sci U S A* **111**, E3243-3251 (2014).

9.   Y. Qiao *et al.*, Detection of lipid-linked peptidoglycan precursors by exploiting an

unexpected transpeptidase reaction. *J Am Chem Soc* **136**, 14678-14681 (2014).

10.   A. García-Heredia *et al.*, Peptidoglycan precursor synthesis along the sidewall of

pole-growing mycobacteria. *eLife* **7**, (2018).

11.   Y. Qiao *et al.*, Lipid II overproduction allows direct assay of transpeptidase

inhibition by beta-lactams. *Nat Chem Biol* **13**, 793-798 (2017).

12. E. C. Hett *et al*., Interaction and modulation of two antagonistic cell wall enzymes of mycobacteria. *PLoS Pathog* **6**, e1001020 (2010).

13. K. J. Kieser *et al.*, Phosphorylation of the peptidoglycan synthase PonA1 governs the rate of polar elongation in mycobacteria. *PLoS Pathog* **11**, e1005010 (2015).

14. G. W. Liechti *et al.*, A new metabolic cell-wall labelling method reveals peptidoglycan in *Chlamydia trachomatis*. *Nature* **506**, 507-510 (2014).

15. N. J. Yang, M. J. Hinner, Getting across the cell membrane: an overview for small molecules, peptides, and proteins. *Methods in Molecular Biology (Clifton, N.J.)* **1266**, 29-53 (2015).

16. A. Muller *et al.*, Daptomycin inhibits cell envelope synthesis by interfering with fluid membrane microdomains. *Proc Natl Acad Sci U S A* **113**, E7077-e7086 (2016).

17. B. L. Orvar *et al.*, Early steps in cold sensing by plant cells: the role of actin cytoskeleton and membrane fluidity. *The Plant Journal* **23**, 785-794 (2000).

18. E. Nagy *et al.*, Hyperfluidization-coupled membrane microdomain reorganization is linked to activation of the heat shock response in a murine melanoma cell line. *Proc Natl Acad Sci U S A* **104**, 7945-7950 (2007).

19. H. N. Foley *et al*., Bioorthogonal chemical reporters for selective in situ probing of mycomembrane components in mycobacteria. *Angewandte Chemie* **128**, 2093-2097 (2016).

20. X. Zhou *et al.*, Sequential assembly of the septal cell envelope prior to V snapping in *Corynebacterium glutamicum*. *Nature Chemical Biology*, 1 (2019).

21.   J. Puffal *et al.*, Demethylmenaquinone Methyl Transferase Is a Membrane Domain-Associated Protein Essential for Menaquinone Homeostasis in *Mycobacterium smegmatis. Frontiers in Microbiology* **9**, 3145 (2018).

22.   Y. S. Morita *et al.*, PimE is a polyprenol-phosphate-mannose-dependent mannosyltransferase that transfers the fifth mannose of phosphatidylinositol mannoside in mycobacteria. *Journal of Biological Chemistry*, (2006).

23.   C. B. Sena *et al.*, Controlled expression of branch-forming mannosyltransferase is critical for mycobacterial lipoarabinomannan biosynthesis. *Journal of Biological Chemistry*, M109. 077297 (2010).

24.   J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nature Methods* **9**, 676-682 (2012).

25.   A. Paintdakhi *et al.*, Oufti: an integrated software package for high-accuracy, high-throughput quantitative microscopy analysis. *Mol Microbiol* **99**, 767-777 (2016).

Acknowledgments: We are grateful to Drs. James Chambers and Amy Burnside for microscopy and flow cytometry guidance, Drs. Eric Strieter and Jiaan Liu and Ms. Sylvia Rivera for technical assistance. We also thank Drs. Hesper Rego and Eric Rubin for ponA1-mRFP, and Dr. Christopher Sassetti for the MurJ and MurG depletion strains. We thank Dr. Benjamin Swarts for providing the OalkTMM and NalkTMM probes.

Funding: Research was supported by funds from the National Institutes of Health (NIH) under awards U01CA221230 and NIH DP2 AI138238 (M.S.S.) and R03 AI140259-01 (Y.S.M.), and R01 AI097191 (J.J. and T.A.G.). A.G.-H. was supported by an Honors

Fellowship from Universidad Autónoma de Nuevo León. J.P. is a recipient of the Science Without Boarders Fellowship from CAPES-Brazil (0328-13-8). **Author contributions:** Conceptualization: M.S.S., Y.S.M., A.G.-H. Formal analysis: A.G.-H. Resources: M.S.S., Y.S.M., T.A.G. and J.J. Writing, review and editing: M.S.S., A.G.-H.,Y.S.M., J.P., T.A.G. and J.J. **Competing interests:** Authors declare no competing interests. **Data and materials availability:** All data are available in the main text or the supplementary materials.

Supplementary Materials:

Materials and Methods

Figures S1-S10

Tables S1

Fig. 1. MurG is enriched in the IMD. (A) Membrane-bound steps of peptidoglycan synthesis with hypothesized partitioning into the IMD and PM-CW. NAM, N-acetylmuramic acid; NAG, N-acetylglucosamine; circles, amino acids: light green, L-ala; red, D-glu; deep blue, diaminopimelic acid; yellow, D-ala. (B) Plasma membrane fractionation. Bacteria are lysed by nitrogen cavitation and cell lysate is sedimented on a sucrose density gradient. (C) Lysates from MurG-Dendra2-expressing *M. smegmatis* were fractionated as in (B) and separated by SDS-PAGE. Protein detected by in-gel fluorescence. Incubation of bacteria with 100 mM benzyl alcohol (BA) for 1 hour decreased the enrichment of MurG-Dendra2 in the IMD. (D) *M. smegmatis* coexpressing MurG-Dendra2 and mCherry-GlfT2 was imaged by structured illumination

microscopy (SIM-E, left). Fluorescence distribution of the fusion proteins from 59 cells

was quantitated from parallel conventional fluorescence microscopy (right). Signal was

normalized to the length and total fluorescence intensity of the cell. Cells were oriented

such that the brighter pole is on the right hand side of the graph. a.u., arbitrary units,

scalebar 5 μm.

Fig. 2. Lipid II is synthesized in the IMD and trafficked to the PM-CW. (A) Detection of

lipid-linked peptidoglycan precursors from organic extracts of *M. smegmatis* membrane

fractions. Terminal D-alanines (yellow) of endogenous precursors are exchanged for

biotin-D-lysine (BDL; pink) via purified *S. aureus* PBP4. Biotinylated species are

detected by blotting with streptavidin-HRP. (B) Detection of peptidoglycan precursors

and PonA1 activity from density gradient fractions. Precursors are in both the IMD and

PM-CW in wildtype *M. smegmatis* but accumulate in the IMD upon MurJ depletion (*10*).

PonA1 binds Bocillin-FL in the PM-CW before or after 1 hour of 100 mM benzyl alcohol

treatment. Wildtype *M. smegmatis* membrane fractions (50 μg/mL of total protein each)

were incubated with 40 μM Bocillin-FL and separated by SDS-PAGE. Active PBPs

detected by in-gel fluorescence in PM-CW. PonA1 was identified in Fig. S3. (C) SIM-E

of *M. smegmatis* coexpressing PonA1-mRFP and PimE-GFP, scalebar 5 μm. (D) Top,

metabolic labeling of mycobacterial cell wall synthesis (*10*). Bottom, *M. smegmatis*

expressing MurG-Dendra2 were incubated with alkDADA for 2 min (~1% generation).

Surface-exposed alkynes on fixed cells were detected by CuAAC with picolyl azide-Cy3

(*10*). Cells imaged by SIM-E, scalebar 5 μm.

Fig. 3. Perturbation of plasma membrane organization disrupts peptidoglycan

biogenesis. (A) *M. smegmatis* coexpressing PonA1-mRFP and MurG-Dendra2 were

imaged +/- benzyl alcohol by SIM-E (left). Fluorescence distribution of the fusion

proteins from 42<n<56 cells was quantitated from parallel conventional fluorescence

microscopy (right). Signal was normalized as Fig. 1D, scalebar 5 µm. (B) Wildtype *M.*

smegmatis +/- benzyl alcohol were incubated with both azido D-amino acid dipeptide

and mycomembrane probe OalkTMM (*10, 19*) for 15 min and fixed in 2% formaldehyde.

Alkynes and azides were detected by sequential CuAAC reactions with picolyl azide

TAMRA and alkyne carboxyrhodamine 110 (Click Chemistry Tools) labels with a wash

step between, scalebar 5 µm. (C) *B. subtilis* were exposed to indicated antibiotics or

benzyl alcohol for 10 min then incubated with alkDADA for an additional 5 min. Cells

were fixed and alkynes were detected by CuAAC with picolyl azide CR 110. MFI,

median fluorescence intensity obtained by flow cytometry. Experiments performed three

times in triplicate. Error bars, +/- standard deviation of biological replicates. ***, $p <$

0.0005; ****, $p < 0.00005$, Tukey multiple comparison test.