1

Cold-induced switch in direction of chloroplast relocation occurs independently of

changes in endogenous phototropin expression

Yuta Fujii [1,2¶], Yuka Ogasawara [1,3¶], Yamato Takahashi [1,3¶], Momoko Sakata [1,4¶], Saori Tamura [1,4], and Yutaka Kodama [1,2,4*]

[1] *Center for Bioscience Research and Education, Utsunomiya University, Tochigi 321-8505, Japan*

[2] *United Graduate School of Agricultural Science, Tokyo University of Agriculture and Technology, Tokyo 183-8509, Japan*

[3] *Faculty of Agriculture, Utsunomiya University, Tochigi 321-8505, Japan*

[4] *Graduate School of Agricultural Science, Utsunomiya University, Tochigi 321-8505, Japan*

[*] Corresponding author

E-mail address: kodama@cc.utsunomiya-u.ac.jp (YK)

[¶]These authors contributed equally to this work.

Abstract

When exposed to fluctuating light intensity, chloroplasts move towards weak light (accumulation response), and away from strong light (avoidance response). In addition, cold treatment (5°C) induces the avoidance response even under weak light conditions (cold-avoidance response). These three responses are mediated by the phototropin (phot), which is a blue-light photoreceptor and has also been found to act as a thermosensory protein that perceives temperature variation. Our previous report indicated that cold-induced changes in phot biochemical activity initiate the cold-avoidance response. In this study, we further explored the induction mechanism of the cold-avoidance response in the liverwort *Marchantia polymorpha* and examined the relationship between changes in the amount of phot and the induction of the cold-avoidance response. The switch between the accumulation and avoidance responses occurs at a so-called 'transitional' light intensity. Our physiological experiments revealed that a cold-mediated decrease in the transitional light intensity leads to the induction of the cold-avoidance response. While artificial overexpression of phot decreased the transitional light intensity as much as cold treatment did, the amount of endogenous phot remained unchanged by cold treatment in wild-type *M. polymorpha*. Taken together, these findings show that the cold-avoidance response is initiated by a cold-mediated reduction of the transitional light intensity, independent of the amount of endogenous phot. This study provides a clue to understand the mechanism underlying the switch in direction of chloroplast relocation in response to light and temperature.

Introduction

Plants sense ambient light and temperature to adapt to an ever-changing environment. In response to the light and temperature changes, the intracellular chloroplast position is precisely controlled. Under weak light conditions (e.g., under canopy), chloroplasts move towards light to increase light capture for photosynthetic efficiency (accumulation response) [1]. Under strong light conditions (e.g., under direct sunlight), chloroplasts move away from light to avoid photodamage (avoidance response) [2]. Chloroplast position is also affected by temperature [3], notably cold. Under warm conditions, weak light induces the accumulation response; however, under cold conditions (5°C) the same light level induces chloroplast avoidance (Fig 1). This movement referred to as the cold-avoidance (or cold-positioning) response, has been studied in several plant species, including the moss *Funaria hygrometrica* [3], evergreen ferns (e.g., *Adiantum capillus-veneris*) [4], the liverwort *Marchantia polymorpha* [5] and the flowering plant *Arabidopsis thaliana* [6]. In *M. polymorpha*, the cold-avoidance response contributes to the reduction of photoinhibition under cold conditions [6].

Earlier studies showed that the accumulation, avoidance and cold-avoidance responses are all mediated by the blue-light (BL) photoreceptor phototropin (phot) [4,6–9]. Phot has two photosensory light, oxygen, or voltage (LOV) domains (LOV1 and LOV2) at its N-terminal region and a serine/threonine kinase domain at its C-terminal region. Phot perceives BL via its LOV domains and mediates the change of chloroplast position upon photoactivation of the LOV domains and kinase-mediated autophosphorylation [10]. A decrease in ambient temperature prolongs the lifetime of the photoactivated LOV2 domain, resulting in an increase in autophosphorylation [6]. The cold-induced enhancement of the photochemical and biochemical activities of phot (i.e., a prolonged lifetime of photoactivated LOV2 and increased autophosphorylation status) is essential for induction of the cold-avoidance response [6], and further study is required to fully understand this induction.

The switch in chloroplast behavior between the accumulation and avoidance

responses occurs at their 'transitional' light intensity (Fig 1A). In the cold-avoidance

response, because the chloroplast moves away from even weak BL (wBL) under cold

conditions [6], it is hypothesized that the cold-avoidance response is induced by a

cold-mediated reduction of the transitional light intensity (Fig 1B). However, this

hypothesis remains to be tested. Artificial overexpression of phot has been suggested to

decrease the transitional light intensity in *A. capillus-veneris* and *A. thaliana* [11,12]. In

addition, expression levels of endogenous phot are affected by both light [7,8,13–16]

and temperature [16] conditions. Based on these findings, we postulate that a variation

in the endogenous amount of phot may be involved in the induction of the

cold-avoidance response.

The goals of this study were two-fold: first, to determine if a cold-mediated

reduction of the transitional light intensity initiates the cold-avoidance response, and

second, to determine if changes in phot amounts play a role in the induction of the

cold-avoidance response. *M. polymorpha* was chosen for this study because it has a

single copy of the gene for phot (Mpphot) that can mediate the accumulation, avoidance,

and cold-avoidance responses [6,17].

Materials and Methods

Plasmid construction

To construct a plasmid to generate the complementation line of Mpphot-Citrine

(35S::Mpphot-Citrine/Mp*phot*^{KO}), we used the Gateway cloning system (Invitrogen)

and the binary vector pMpGWB306 [18], which includes *Citrine* (a gene for yellow

fluorescent protein) for fusion at the 3□-end of the target gene under the control of the

cauliflower mosaic virus (CaMV) 35S promoter. The LR reaction (Invitrogen) was

performed   with   a   mixture   of   pMpGWB306   and   the   donor   plasmid

pDONR207-MpPHOT [19], and the resulting plasmid pMpGWB306-MpPHOT was

used for transformation of *M. polymorpha*.

Plant materials and growth conditions

Thalli of *M. polymorpha* were asexually cultured on half-strength B5 medium under 75

μmol photons m^{-2} s^{-1} continuous white light (FL40SW, NEC Corporation). Light

intensity was measured by using a LI-250A light meter (LI-COR Biosciences). The

Mp*PHOT* knockout line (Mp*phot^{KO}*) was generated and kindly provided by Komatsu et

al. [17]. In this study, transformation of *M. polymorpha* was performed by the

G-AgarTrap method [20,21], which is an *Agrobacterium*-mediated transformation

method used for gemmalings. To generate the transformants, we used the binary vector

pMpGWB106-MpPHOT   [19]   for   Mpphot-Citrine   overexpression   lines

(35S::Mpphot-Citrine/WT), and pMpGWB306-MpPHOT for the complementation line

of   Mpphot   (35S::Mpphot-Citrine/Mp*phot^{KO}*).   The   Citrine-expressing   line

(35S::Citrine/WT) was generated in our previous study [20]. G2 gemmalings were used

throughout.

Observation of chloroplast positioning

One-day-old gemmalings were incubated in temperature-controlled incubators (IJ100,

Yamato Scientific Co., LTD) and light conditions were adjusted using blue

light-emitting diodes (LEDs; OptoSupply Limited) and a LI-250A light meter. To

produce polarized weak BL, 25 μmol photons m^{-2} s^{-1} of the blue LED was used as weak

BL [5,6], and it was filtered with a polarizing filter (HN38, Polaroid Corp.).

Chlorophyll fluorescence images were captured using a fluorescence microscope

MZ16F (Leica Microsystems) equipped with a DP73 digital camera (Olympus) and

quantification of chloroplast positioning was performed using the P/A ratio method with

ImageJ software, as previously described [5]. Briefly, fluorescence intensity was

measured at 30 points (0.625 µm each, equivalent to 1 pixel) along the anticlinal cell

walls (A) and 30 points (39.1 µm^2 each, equivalent to 10×10 pixels) along the

periclinal cell walls (P). This procedure was repeated five times, and then the P/A ratio

was calculated as a mean of the five P/A values. The standard deviation was calculated

using Excel software (Microsoft).

Immunoblot analysis

Immunoblot analysis was performed as previously described [6]. Briefly, four-day-old

gemmalings were collected in liquid nitrogen and kept at -80°C until use. The frozen

plants were homogenized in SDS buffer (2% SDS, 10% glycerol, 5% mercaptoethanol,

and 0.25 M Tris-HCl pH6.8) or Nonidet buffer (0.5% Nonidet P-40, 0.5M EDTA, 0.15

M NaCl, and 0.01 M Tris-HCl pH7.5) using a mortar and pestle and then boiled at 95°C

for 5 min. The samples were centrifuged at 14,000 g for 10 min, and the supernatants

used. Total protein amount was quantified by Bradford-XL assay (APRO Science).

Fifteen µg of protein was subjected to SDS-PAGE with 8% polyacrylamide gels and

transferred to a polyvinylidene difluoride (PVDF) membrane [6]. MpPHOT proteins

were detected with rabbit antibody raised against the custom-made synthetic peptide

(C+VDERAPPSKGSAKE) (Eurofins Genomics) and horseradish

peroxidase-conjugated anti-rabbit IgG (Thermo Scientific). Both antibodies were used

at dilution of 1:2000. For histone detection, anti-Histone H3 antibody (Abcam) at

dilution of 1:5000 and horseradish peroxide-conjugated anti-rabbit IgG (Thermo

Scientific) at dilution of 1:2000 were used. Chemiluminescent signal was detected using

ECL-Select (GE Healthcare) and Light Capture (ATTO). For the image of Rubisco

large subunit (RBCL), fifteen µg protein was subjected to SDS-PAGE with 8%

polyacrylamide gels and stained with conventional Coomassie brilliant blue (CBB)

method.

Results

Chloroplasts avoid weak blue light (wBL) under cold conditions

Using temperature-regulated microscopy with a microbeam irradiation system, we previously reported that under cold conditions (5°C), chloroplasts moved away from wBL that induced the accumulation response at 22°C [6]. However, in the study, physiological evidence of wBL avoidance under cold conditions was limited to microbeam analysis [6]. Here, we further analyzed whether chloroplasts move away from wBL under cold conditions.

In *M. polymorpha*, both cold and BL were required to elicit the cold-avoidance response [6]. Building on the result, to explore whether the cold-avoidance response is due to avoidance from wBL under cold conditions, we tested the effects of changing light direction. First, to induce the cold-avoidance response, wild-type (WT) cells were incubated under 25 µmol photons m^{-2} s^{-1} of wBL (BL25) with vertical irradiation at 5°C (Fig 2A). In WT cells, chloroplasts were positioned along the anticlinal cell walls as the cold-avoidance response (Fig 2B). When the light direction was changed from vertical to horizontal (Fig 2A), the chloroplasts along the anticlinal cell walls moved to the periclinal cell walls (Fig 2B), indicating chloroplast avoidance from BL25. To further confirm the results, we employed polarized wBL, because the vertically vibrating polarized wBL could induce chloroplast localization along the anticlinal wall by the accumulation response in *A. capillus-veneris* [22,23]. When 10-day-old sporeling cells of *M. polymorpha* were irradiated with vertically vibrating polarized wBL for 3 h at 22°C, chloroplasts localized along the anticlinal wall as the accumulation response (Fig 2C). When the cells were transferred to 5°C under the same polarized wBL condition, chloroplasts relocated from the anticlinal wall to the periclinal wall, indicating that chloroplasts avoided the polarized wBL at 5°C. These results confirm that the cold-avoidance response is due to avoidance from wBL under

cold conditions, which is consistent with the microbeam analysis in our previous study

[6].

Cold treatment decreases the transitional light intensity between the accumulation

and avoidance responses

We hypothesized that the cold-avoidance response is induced by a cold-mediated

decrease of the transitional light intensity (Fig 1). If this is the case, the accumulation

response would be induced under much weaker BL in cold conditions. To investigate

whether the accumulation response is induced at 5°C, WT cells were incubated under

different intensities of wBL (5, 10, 15 and 25 µmol photons m^{-2} s^{-1}; BL5, BL10, BL15

and BL25, respectively) (Fig 3A and 3B). Chloroplast position was quantified by a P/A

ratio method, developed in earlier studies [4,5]. The P/A ratio is an arbitrary unit

calculated by dividing the chlorophyll fluorescence intensity along the periclinal cell

wall by that along the anticlinal cell wall, with a higher value indicating periclinal

positioning of chloroplasts and vice versa. In WT cells, the cold-avoidance response

was observed at 5°C under BL10, BL15 and BL25 (Fig 3A), and at these higher BL

intensity levels, we observed a decrease in P/A ratios (Fig 3B). However, the

cold-avoidance response was not observed at 5°C under BL5 (Fig 3A) and the P/A ratio

was unchanged during the incubation of these samples (Fig 3B). These results suggest

that BL5 induces the accumulation response at 5°C. However, because similar

chloroplast positioning was observed under dark conditions at 5°C [6], it is also possible

that the cells were unable to respond to the extremely weak intensity of BL5. To clarify

whether the accumulation response is induced under BL5 or not, an additional test was

carried out. Following the induction of the cold-avoidance response at 5°C for 24 h

under BL25, the light intensity was changed to BL5 or dark conditions (Fig 3C). After

incubation for 24 h, BL5 conditions induced the accumulation response at 5°C, whereas

under dark conditions there was no change in chloroplast position (Fig 3D). These

results indicate that the accumulation response is induced under BL5 at 5°C. Taken

together, these results indicate that the transitional light intensity is modulated by the

temperature change (Fig 1).

Artificial overexpression of Mpphot in *M. polymorpha* induces the avoidance

response under wBL without cold treatment

We next considered the possibility that changes in endogenous phot expression may

play a role in the change in transitional light intensity, as previous studies using *A.*

capillus-veneris and *A. thaliana* suggest that a reduction in the transitional light

intensity can be the result of artificial overexpression of phot [11,12]. In *A.*

capillus-veneris at 25°C, transient overexpression of the Ac*PHOT2-GFP* gene induced

the avoidance response under wBL [11]. Similarly, the avoidance response was

observed under wBL in transgenic *A. thaliana* overexpressing the At*PHOT2-GFP* gene

at 23°C [12]. Consistent with these previous studies, we found that artificial

overexpression of Mpphot fused to the yellow fluorescent protein Citrine

(Mpphot-Citrine) in WT cells (35S::Mpphot-Citrine/WT) induced the avoidance

response under BL25 at 22°C (without cold treatment), and degree of the avoidance

response was dependent on Mpphot-Citrine expression levels (Fig 4A-4C). In the

overexpression line (OX2) under BL5, the accumulation response at 22°C and

cold-avoidance response at 5°C could be induced (Fig 4D and 4E), indicating that the

transitional intensity is reduced by overexpression of Mpphot-Citrine. Therefore, it is

conceivable that increase of the endogenous amount of Mpphot might play a role in the

induction of the cold-avoidance response.

Alteration of endogenous Mpphot is not required for the induction of the

cold-avoidance response

To examine the role of Mpphot expression in the induction of the cold-avoidance

response, the endogenous amount of Mpphot was determined using immunoblot

analysis. After gemmalings were transferred from the white light condition to BL25

condition at 22°C, the endogenous amount of Mpphot was increased at 6 h and the

increased amount was maintained until 24 h (S Fig 1). To avoid the effect of the light

condition change (i.e., from white light to BL25), the gemmalings were pre-incubated

under BL25 at 22°C for 24 h. When temperature was changed from 22°C to 5°C, the

endogenous amount of Mpphot was not changed at 3, 6, 9 h (Fig 5A). Note that this

experimental condition was not able to separate phosphorylated and dephosphorylated

forms of Mpphot. When we incubated the gemmalings under the same light and

temperature condition as shown in Fig 5A, the cold-avoidance response was induced

(Fig 5B), and the P/A ratio was decreased (Fig 5C). Unexpectedly the pre-incubation

under BL25 at 22°C for 24 h increased the P/A ratio than the white light condition at

22°C (see 0 h in Fig 5C and 0 h in Fig 3B). Anyway, an increase of the endogenous

amount of Mpphot is not required for induction of the cold-avoidance response in *M.*

polymorpha. To confirm this result, we examined the complementation lines of

Mpphot-Citrine under the control of the CaMV35S promoter with the genetic

background of Mp*phot*[KO] (35S::Mpphot-Citrine/Mp*phot*[KO]), as CaMV35S promoter

activity is reported to be suppressed by cold treatment in tobacco plants [24]. In

transgenic *M. polymorpha* expressing Citrine under the control of the CaMV35S

promoter (35S::Citrine/WT), we also confirmed that the promoter activity is slightly

suppressed by 5°C treatment in *M. polymorpha* (Fig 5D). In the complementation lines

(35S::Mpphot-Citrine/Mp*phot*[KO]), the cold-avoidance response was clearly observed

(Fig 5E), confirming that the induction of the cold-avoidance response does not require

any increase of Mpphot amount. Given that cold treatment increased the

autophosphorylation status of phot [6,25], we conclude that an increase of phot

autophosphorylation status decreases the transitional light intensity without alteration of

the endogenous amount of Mpphot.

Discussion

The transitional light intensity between the accumulation and avoidance responses

can be changed

Light-induced chloroplast movement is widely found in photosynthetic organisms from

algae to land plants [3]. The transitional light intensity between the accumulation and

avoidance responses differs among various plant species [26]. Furthermore, the

transitional light intensity can be reduced in response to low temperature, inducing the

cold-avoidance response [4–6]. In this study, we analyzed the light fluence rate

dependency of the cold-avoidance response. Under cold conditions, chloroplasts moved

in a light fluence rate-dependent manner (Fig 3A and 3B), as occurs during the

avoidance response at a standard growth temperature [27]. This result supports the

conclusion that there is a cold-induced reduction of the transitional light intensity (Fig

1). Because the cold-avoidance response has been found to moderate photodamage in *M.*

polymorpha [6], the plasticity of the transitional light intensity is vital for plants

exposed to low temperatures. In *A. thaliana*, the transitional light intensity is controlled

by two types of phot: Atphot1 and Atphot2. Atphot1 specifically regulates the

accumulation response, while Atphot2 regulates both the accumulation and avoidance

responses [7–9]. In this context, the coordination mechanism of the transitional light

intensity in *A. thaliana* appears to be more complex than that in *M. polymorpha*. Further

analysis of the transitional light intensity in various plant species and genetic mutants

may lead to a better understanding of the transition mechanism of chloroplast relocation

movements.

Photoactivated phototropin amount changes the transitional light intensity

Intracellular phot proteins are more frequently activated under strong light than weak

light conditions. Weak light irradiation at lower temperatures, prolongs the lifetime of

photoactivated phot [6], and like the effect of strong light irradiation at a standard

growth temperature, results in the increase of photoactivated phot population. Therefore,

we concluded that the intracellular amount of photoactivated phot determines the

transition between the accumulation and avoidance responses [6]. Furthermore,

considering the effect of phot overexpression on the transitional light intensity (Fig 4),

the transition mechanism may be based on the absolute amount of photoactivated phot

in the cell rather than the ratio of active/inactive phot. How an increase in

photoactivated phot proteins triggers the switch from the accumulation to avoidance

responses remains unknown. Because phot autophosphorylation is also increased under

strong-light/warm or weak-light/cold conditions [6,10,25], we speculate that to induce

the avoidance response, an increase of photoactivated phot amount may change the phot

phosphorylation status, possibly via intermolecular phosphorylation activity [28,29].

Notably, the recruitment of phot to microdomains at the plasma membrane is involved

in phot autophosphorylation [30]. Further autophosphorylation analysis regarding the

spatial density of photoactivated phot may clarify the switching mechanism between the

accumulation and avoidance responses.

Alteration of phot expression is not required for induction of the cold-avoidance

response in *M. polymorpha*

The cold-avoidance response is initiated by the temperature-dependent biochemistry of

the phot LOV2 domain [6]. However, in *A. thaliana*, alteration of the endogenous

amount of Atphot1 and Atphot2 in response to temperature was reported; lower

temperature decreases Atphot1 expression and increases Atphot2 expression [16]. In

this study, we showed that the cold-avoidance response can be induced regardless of the

change in the endogenous Mpphot amount (Fig 5A-5C) and even without its promoter

(Fig 5D and 5E), indicating that transcriptional regulation of Mpphot is not required for

the cold-avoidance response in *M. polymorpha*. Mpphot is similar to Atphot2, which

regulates both the accumulation and avoidance responses [17]. Consistent with Mpphot,

Atphot2 expression in *A. thaliana* was reported to be unchanged at a temperature range

from 4°C to 23°C [16]. Interestingly, cold treatment had a more significant impact on

the decrease of Atphot1 expression than the increase of Atphot2 expression [16],

implying that the expression of Atphot1 might be more sensitive to cold than that of

Atphot2. The gene duplication of phot [31] and functional differentiation between phot1

and phot2 may have enabled land plants to have different transitional light intensities

and chloroplast movements, allowing species to expand their range of habitat.

Acknowledgements

The authors thank Dr. Takayuki Kohchi (Kyoto University) and Dr. Kimitsune Ishizaki

(Kobe University) for providing *M. polymorpha* (WT and Mp*phot*^{KO}) and the binary

vectors (pMpGWB106 and pMpGWB306). The authors also thank Ms. Lee Kien Yong

(Utsunomiya University) for helping the preparation of the manuscript. This work was

supported by the Japan Society for the Promotion of Science KAKENHI [grant No.

18H02455 to Y.K.]; the Japan Science and Technology Agency, Exploratory Research

for Advanced Technology, Numata Organelle Reaction Cluster [grant No. JPMJER1602

to Y.K.]; and the Plant Transgenic Design Initiative of the University of Tsukuba to

Y.K.

References

1.　Gotoh E, Suetsugu N, Yamori W, Ishishita K, Kiyabu R, Fukuda M, et al. Chloroplast accumulation response enhances leaf photosynthesis and plant biomass production. Plant Physiol. 2018;178: 1358–1369. doi:10.1104/pp.18.00484

2.　Kasahara M, Kagawa T, Oikawa K, Suetsugu N, Miyao M, Wada M. Chloroplast avoidance movement reduces photodamage in plants. Nature. 2002;420: 829–832. doi:10.1038/nature01202.1.

3. Senn G. Die gestalts-und lageveränderung der pflanzen-chromatophoren.
Leipzig: Wilhelm-Engelmann; 1908.
4. Kodama Y, Tsuboi H, Kagawa T, Wada M. Low temperature-induced
chloroplast relocation mediated by a blue light receptor, phototropin 2, in fern
gametophytes. J Plant Res. 2008;121: 441–448. doi:10.1007/s10265-008-0165-9
5. Ogasawara Y, Ishizaki K, Kohchi T, Kodama Y. Cold-induced organelle
relocation in the liverwort Marchantia polymorpha L. Plant, Cell Environ.
2013;36: 1520–1528. doi:10.1111/pce.12085
6. Fujii Y, Tanaka H, Konno N, Ogasawara Y, Hamashima N, Tamura S, et al.
Phototropin perceives temperature based on the lifetime of its photoactivated
state. Proc Natl Acad Sci U S A. 2017;114: 9206–9211.
doi:10.1073/pnas.1704462114
7. Jarillo JA, Gabrys H, Capel J, Alonso JM, Ecker JR, Cashmore   a R.
Phototropin-related NPL1 controls chloroplast relocation induced by blue light.
Nature. 2001;410: 952–954. doi:10.1038/35073622
8. Kagawa T, Sakai T, Suetsugu N, Oikawa K, Ishiguro S, Kato T, et al.
Arabidopsis NPL1: a phototropin homolog controlling the chloroplast high-light
avoidance response. Science. 2001;291: 2138–2141.
doi:10.1126/science.291.5511.2138
9. Sakai T, Kagawa T, Kasahara M, Swartz TE, Christie JM, Briggs WR, et al.
Arabidopsis nph1 and npl1: blue light receptors that mediate both phototropism
and chloroplast relocation. Proc Natl Acad Sci U S A. 2001;98: 6969–6974.
doi:10.1073/pnas.101137598
10. Inoue S-I, Kinoshita T, Matsumoto M, Nakayama KI, Doi M, Shimazaki K-I.
Blue light-induced autophosphorylation of phototropin is a primary step for
signaling. Proc Natl Acad Sci U S A. 2008;105: 5626–5631.
doi:10.1073/pnas.0709189105
11. Kagawa T, Kasahara M, Abe T, Yoshida S, Wada M. Function analysis of
phototropin2 using fern mutants deficient in blue light-induced chloroplast
avoidance movement. Plant Cell Physiol. 2004;45: 416–426.
doi:10.1093/pcp/pch045
12. Kong SG, Kinoshita T, Shimazaki KI, Mochizuki N, Suzuki T, Nagatani A. The
C-terminal kinase fragment of Arabidopsis phototropin 2 triggers constitutive
phototropin responses. Plant J. 2007;51: 862–873.
doi:10.1111/j.1365-313X.2007.03187.x
13. Kanegae H, Tahir M, Savazzini F, Yamamoto K, Yano M, Sasaki T, et al. Rice
NPH1 homologues, OsNPH1a and OsNPH1b, are differently photoregulated.
Plant Cell Physiol. 2000;41: 415–423. doi:10.1093/pcp/41.4.415
14. Jain M, Sharma P, Tyagi SB, Tyagi AK, Khurana JP. Light regulation and
differential tissue-specific expression of phototropin homologues from rice
(Oryza sativa ssp. indica). Plant Sci. 2007;172: 164–171.
doi:10.1016/j.plantsci.2006.08.003
15. Suzuki H, Okamoto A, Kojima A, Nishimura T, Takano M, Kagawa T, et al.
Blue-light regulation of ZmPHOT1 and ZmPHOT2 gene expression and the
possible involvement of Zmphot1 in phototropism in maize coleoptiles. Planta.
2014;240: 251–261. doi:10.1007/s00425-014-2082-6
16. Łabuz J, Hermanowicz P, Gabryś H. The impact of temperature on blue light
induced chloroplast movements in Arabidopsis thaliana. Plant Sci. 2015;239:
238–249. doi:10.1016/j.plantsci.2015.07.013
17. Komatsu A, Terai M, Ishizaki K, Suetsugu N, Tsuboi H, Nishihama R, et al.
Phototropin encoded by a single-copy gene mediates chloroplast photorelocation
movements in the liverwort Marchantia polymorpha. Plant Physiol. 2014;166:
411–427. doi:10.1104/pp.114.245100

18. Ishizaki K, Nishihama R, Ueda M, Inoue K, Ishida S, Nishimura Y, et al.
Development of gateway binary vector series with four different selection
markers for the liverwort Marchantia polymorpha. PLoS One. 2015;10:
e0138876. doi:10.1371/journal.pone.0138876
19. Kodama Y. Time gating of chloroplast autofluorescence allows clearer
fluorescence imaging in planta. Abraham T, editor. PLoS One. 2016;11:
e0152484. doi:10.1371/journal.pone.0152484
20. Tsuboyama-Tanaka S, Kodama Y. AgarTrap-mediated genetic transformation
using intact gemmae/gemmalings of the liverwort Marchantia polymorpha L. J
Plant Res. 2015;128: 337–344. doi:10.1007/s10265-014-0695-2
21. Tsuboyama S, Kodama Y. AgarTrap protocols on your benchtop: Simple
methods for Agrobacterium-mediated genetic transformation of the liverwort
Marchantia polymorpha. Plant Biotechnol. 2018; in press.
22. Yatsuhashi H, Kadota A, Wada M. Blue- and red-light action in photoorientation
of chloroplasts in Adiantum protonemata. Planta. 1985;165: 43–50.
doi:10.1007/BF00392210
23. Kagawa T, Wada M. Polarized light induces nuclear migration in prothallial cells
of Adiantum capillus-veneris L. Planta An Int J Plant Biol. 1995;196: 775–780.
doi:10.1007/BF01106773
24. Schnurr JA, Guerra DJ. The CaMV-35S promoter is sensitive to shortened
photoperiod in transgenic tobacco. Plant Cell Rep. 2000;19: 279–282.
doi:10.1007/s002990050012
25. Okajima K, Kashojiya S, Tokutomi S. Photosensitivity of kinase activation by
blue light involves the lifetime of a cysteinyl-flavin adduct intermediate, S390, in
the photoreaction cycle of the LOV2 domain in phototropin, a plant blue light
receptor. J Biol Chem. 2012;287: 40972–40981. doi:10.1074/jbc.M112.406512
26. Königer M, Bollinger N. Chloroplast movement behavior varies widely among
species and does not correlate with high light stress tolerance. Planta. 2012;236:
411–426. doi:10.1007/s00425-012-1619-9
27. Kagawa T, Wada M. Chloroplast-avoidance response induced by high-fluence
blue light in prothallial cells of the fern Adiantum capillus-veneris as analyzed by
microbeam irradiation. Plant Physiol. 1999;119: 917–923.
doi:10.1104/pp.119.3.917
28. Cho HY, Tseng TS, Kaiserli E, Sullivan S, Christie JM, Briggs WR.
Physiological roles of the light, oxygen, or voltage domains of phototropin 1 and
phototropin 2 in arabidopsis. Plant Physiol. 2007;143: 517–529.
doi:10.1104/pp.106.089839
29. Kaiserli E, Sullivan S, Jones MA, Feeney KA, Christie JM. Domain swapping to
assess the mechanistic basis of Arabidopsis phototropin 1 receptor kinase
activation and endocytosis by blue light. Plant Cell. 2009;21: 3226–3244.
doi:10.1105/tpc.109.067876
30. Xue Y, Xing J, Wan Y, Lv X, Fan L, Zhang Y, et al. Arabidopsis Blue Light
Receptor Phototropin 1 Undergoes Blue Light-Induced Activation in Membrane
Microdomains. Mol Plant. 2018;11: 846–859. doi:10.1016/j.molp.2018.04.003
31. Li FW, Rothfels CJ, Melkonian M, Villarreal JC, Stevenson DW, Graham SW, et
al. The origin and evolution of phototropins. Front Plant Sci. 2015;6: 1–11.
doi:10.3389/fpls.2015.00637
439
440

Figure Legends

Fig 1. Schematic diagram showing the transitional light intensity of the
accumulation and avoidance responses. Solid vertical lines indicate points of the
transitional light intensity under warm (A; 22°C) and cold (B; 5°C) conditions.

Fig 2. Chloroplasts avoid weak blue light under cold conditions. (A) Diagram of the
experimental conditions used to induce the cold-avoidance response with vertical or
horizontal irradiation of blue-light. After incubation under 25 µmol photons m^{-2} s^{-1} of
blue-light (BL25) with vertical irradiation at 5°C for 24 h, the WT gemmalings were
irradiated by horizontal BL25. (B) Chlorophyll fluorescence images after irradiation of
vertical (left) or horizontal (right) BL as shown in (A). The white arrow indicates the
light direction. (C) Chlorophyll fluorescence images after irradiation of the polarized
weak BL. Ten-day-old WT sporelings were irradiated by vertically vibrating polarized
weak BL at 22°C. The cells were transferred to 5°C, followed by incubation for 24 h
and 60 h.

Fig 3. Cold-induced reduction of the transitional light intensity. (A) Chloroplast
positioning in WT gemmalings incubated at 5°C for 12 h with 5, 10, 15, and 25 µmol
photons m^{-2} s^{-1} of blue-light (BL5, BL10, BL15, and BL25, respectively). After
pre-incubation of 1-day-old WT gemmalings under BL25 at 22°C for 1 h to induce the
accumulation response, the cells were transferred to the indicated BL conditions at 5°C.
(B) The P/A ratios, calculated from chlorophyll fluorescence images after incubation for
0, 2, 6, 8, 12, and 24 h. Error bars indicate standard deviations. (C) Diagram of
experimental conditions used to confirm the cold-induced reduction of the transitional
intensity. After pre-incubation under BL25 at 5°C for 24 h, WT gemmalings were
transferred to BL5 or dark and incubated at 5°C for 24 h. (D) Representative

chlorophyll fluorescence images of WT gemmalings after pre-incubation (BL25) and

incubation under BL5 (from BL25 to BL5) or dark (from BL25 to Dark) as shown in

(C).

Fig 4. Overexpression of Mpphot-Citrine decreases the transitional light intensity.

(A) Immunoblot analysis of WT and Mpphot-Citrine overexpression lines

(35S::Mpphot-Citrine/WT: OX1, OX2, and OX3) with anti-Mpphot antibody. White

and black arrowheads indicate Mpphot-Citrine and Mpphot, respectively. Rubisco

large-subunit (RBCL) was shown as a loading control. (B) Representative images of

chlorophyll fluorescence in WT and Mpphot-Citrine overexpression lines. (C) The P/A

ratios calculated from the chlorophyll fluorescence images shown in (B) after the

incubation under BL25 at 22°C for 12 h. (D) Confirmation of the reduction of

transitional light intensity in OX2. In the 1-day-old gemmalings from OX2, the

accumulation response was induced under BL5 at 22°C (left panel). The cold-avoidance

response was induced under BL5 at 5°C for 24 h (right panel). (E) Time-course of the

cold-avoidance response induced under BL5 at 5°C in OX2. The P/A ratios were

calculated from the chlorophyll fluorescence images after incubation for 0, 6, 12, and 24

h.

Fig 5. Alteration of Mpphot expression is not required for the induction of the

cold-avoidance response. (A) Immunoblot analysis of endogenous MpPHOT amount

in 4-day-old WT gemmaling (white light for 3 days and BL25 for 24 h at 22°C),

incubated under BL25 at 5°C. The black arrowhead and asterisk indicate MpPHOT and

non-specific signal, respectively. Histone H3 protein is shown as a loading control. (B)

Representative chlorophyll fluorescence images of WT gemmalings incubated under the

same light and temperature conditions as shown in (A). (C) The P/A ratios, calculated

from chlorophyll fluorescence images after incubation for 0, 3, 6 and 9 h. Error bars

indicate standard deviations. (D) Immunoblot analysis of cold-induced repression of activity of CaMV35S promoter. A transformant expressing Citrine under the control of CaMV35S promoter (35S::Citrine/WT) was used. Coomassie Brilliant Blue (CBB)-stained Rubisco large-subunit (RBCL) was used as a loading control. (E) Chlorophyll fluorescence images showing the cold-avoidance response in complementation lines (35S::Mpphot-Citrine/Mp*phot*^{KO}). After incubation under BL25 at 22°C, the gemmalings were transferred to 5°C under the same light conditions.

S Fig 1. Effect of the light condition change on the endogenous Mpphot expression. Immunoblot analysis of endogenous Mpphot amount in WT gemmaling, incubated under BL25 at 22°C after culture under the white light condition (75 µmol photons m^{-2} s^{-1}) at 22°C for 3 days. The black arrowhead and asterisk indicate Mppot and non-specific signal, respectively. Histone H3 protein is shown as a loading control.

A Warm

Accumulation   Avoidance

B Cold

A

Vertical

Horizontal

B

Vertical | Horizontal

BL25 | BL25

Irradiation →

C

22°C

5°C

24 h | 60 h

A

22°C   5°C

BL25   BL5   BL10   BL15   BL25

B

BL5   BL10
BL15   BL25

P/A ratio

1.5

1.0

0.5

0 4 8 12 16 20 24 (h)

C

BL25 → BL5 / Dark

5°C for 24 h   5°C for 24 h

D

BL25   BL25→BL5   BL25→Dark

A

kDa   WT   OX1   OX2   OX3

250
150   ◁
100   ◀

RBCL

B

WT	OX1
22°C	22°C
OX2	OX3
22°C	22°C

C

P/A ratio

WT   OX1   OX2   OX3

D

BL5	BL5
22°C	5°C

E

P/A ratio

2.0
1.5
1.0
0.5

0   4   8   12   16   20   24 (h)

A

5°C

| | 0 | 3 | 6 | 9 | (h) |

Mpphot

Histone

B

| 0 h | 3 h | 6 h | 9 h |

C

P/A ratio

4.0
3.0
2.0
1.0

0   3   6   9 (h)

D

| | 0 | 5°C | | 22°C | | |
| | | 3 | 6 | 3 | 6 | (h) |

Citrine

RBCL

E

35S::Mpphot-Citrine/Mp*phot*^{KO}

| 22°C | 5°C |