The Group B Streptococcal surface antigen I/II protein, BspC, interacts with

host vimentin to promote adherence to brain endothelium and inflammation

during the pathogenesis of meningitis

Liwen Deng[1,2], Brady L. Spencer[2], Joshua A. Holmes[2], Rong Mu[1], Sara Rego[3], Thomas A.

Weston[1], Yoonsung Hu[4], Glenda F. Sanches[2,5], Sunghyun Yoon[6], Nogi Park[6], Prescilla E.

Nagao[5], Howard F. Jenkinson[3], Justin A. Thornton[4], Keun Seok Seo[6], Angela H. Nobbs[3] and

Kelly S. Doran[1,2]

[1]Department of Biology, San Diego State University, 5500 Campanile Dr. San Diego, CA,

92182, USA.

[2]Department of Immunology and Microbiology, University of Colorado School of Medicine,

Aurora, CO, 80045, USA.

[3]Bristol Dental School, University of Bristol, Lower Maudlin Street, Bristol BS1 2LY, UK.

[4]Department of Biological Sciences, Mississippi State University, Mississippi State, MS, 39762,

USA.

[5]Laboratory of Molecular Biology and Physiology of Streptococci, Instituto de Biologia Roberto

Alcantara Gomes, Rio de Janeiro State University, Rio de Janeiro, Brazil.

[6]Department of Basic Sciences, College of Veterinary Medicine, Mississippi State University,

Mississippi State, MS 39762, USA.

<u>SHORT TITLE</u>

The GBS adhesin BspC interacts with vimentin to promote meningitis

ABSTRACT

Streptococcus agalactiae (Group B *Streptococcus*, GBS) normally colonizes healthy adults but can cause invasive disease, such as meningitis, in the newborn. To gain access to the central nervous system, GBS must interact with and penetrate brain or meningeal blood vessels; however, the exact mechanisms are still being elucidated. Here, we investigate the contribution of BspC, an antigen I/II family adhesin, to the pathogenesis of GBS meningitis. Disruption of the *bspC* gene reduced GBS adherence to human cerebral microvascular endothelial cells (hCMEC), while heterologous expression of BspC in non-adherent *Lactococcus lactis* conferred bacterial attachment. In a murine model of hematogenous meningitis, mice infected with Δ*bspC* mutants exhibited lower mortality as well as decreased brain bacterial counts and inflammatory infiltrate compared with mice infected with WT GBS strains. Further, BspC was both necessary and sufficient to induce neutrophil chemokine expression. We determined that BspC interacts with the host cytoskeleton component vimentin, and confirmed this interaction using a bacterial two-hybrid assay, immunofluorescent staining, and imaging flow cytometry. Vimentin null mice were protected from WT GBS infection and also exhibited less inflammatory cytokine production in brain tissue. These results suggest that BspC and the vimentin interaction is critical for the pathogenesis of GBS meningitis.

AUTHOR SUMMARY

Group B *Streptococcus* (GBS) typically colonizes healthy adults but can cause severe disease in immune compromised individuals, including newborns. Despite wide-spread intrapartum antibiotic prophylaxis given to pregnant women, GBS remains a leading cause of neonatal meningitis. To cause meningitis, GBS must interact with and penetrate the blood-brain barrier (BBB), which separates bacteria and immune cells in the blood from the brain. In order to develop targeted therapies to treat GBS meningitis, it is important to understand the mechanisms of BBB crossing. Here, we describe the role of the GBS surface factor, BspC, in promoting meningitis and discover the host ligand for BspC, vimentin, which is an intermediate filament protein that is constitutively expressed by endothelial cells. We determined that BspC interacts with the C-terminal domain of cell-surface vimentin to promote bacterial attachment to brain endothelial cells and that purified BspC protein can induce immune signaling pathways. In a mouse model of hematogenous meningitis, we observed that a GBS mutant lacking BspC was less virulent compared to WT GBS and resulted in less inflammatory disease. We also observed that mice lacking vimentin were protected from GBS infection. These results reveal the importance of the BspC-vimentin interaction in the progression of GBS meningitis disease.

INTRODUCTION

Streptococcus agalactiae (Group B *Streptococcus*, GBS) is an opportunistic pathogen that asymptomatically colonizes the vaginal tract of up to 30% of healthy women. However, GBS possesses a variety of virulence factors and can cause severe disease when transmitted to susceptible hosts such as the newborn. Despite widespread intrapartum antibiotic administration to colonized mothers, GBS remains a leading cause of pneumonia, sepsis, and meningitis in neonates (1, 2). Bacterial meningitis is a life-threatening infection of the central nervous system (CNS) and is marked by transit of the bacterium across endothelial barriers, such as the blood-brain barrier (BBB) or the meningeal blood-cerebral spinal fluid barrier (mBCSFB). Both consist of a single layer of specialized endothelial cells that serve to maintain brain homeostasis and generally prevent pathogen entry into the CNS (3-5). Symptoms of bacterial meningitis may be due to the combined effect of bacterial adherence and brain penetration, direct cellular injury caused by bacterial cytotoxins, and/or activation of host inflammatory pathways that can disrupt brain barrier integrity and damage underlying nervous tissue. (6-8)

Bacterial meningitis typically develops as a result of the pathogen spreading from the blood to the meninges. In order to disseminate from the blood into the brain, GBS must first interact with barrier endothelial cells (9). A number of surface-associated factors that contribute to GBS-brain endothelium interactions have been described such as lipoteichoic acid (LTA) (10), pili (11), serine-rich repeat proteins (Srr) (12), and streptococcal fibronectin-binding protein (SfbA) (13). Pili, the Srr proteins, and SfbA have been shown to interact with extracellular matrix (ECM) components, which may help to bridge to host receptors such as integrins or other ECM

receptors. However, a direct interaction between a GBS adhesin and an endothelial cell receptor

has not been described.

Antigen I/II family (AgI/II) proteins are multifunctional adhesins that have been well

characterized as colonization determinants of oral streptococci (14). These proteins mediate

attachment of *Streptococcus mutans* and *Streptococcus gordonii* to tooth surfaces and can

stimulate an immune response from the colonized host (14). Genes encoding AgI/II polypeptides

are found in streptococcal species indigenous to the human mouth as well as other pathogenic

streptococci such as GBS, *S. pyogenes* (Group A *Streptococcus*, GAS), and *S. suis* (14, 15).

Intriguingly, the GAS AgI/II protein AspA (Group <u>A</u> *Streptococcus* <u>s</u>urface <u>p</u>rotein) is absent in

many GAS M serotypes and is found predominantly among M serotypes implicated in puerperal

sepsis and neonatal infections, including M2, M4, and M28. The gene encoding AspA is located

within an integrative and conjugative element designated region of difference 2 (RD2), which

likely originated in GBS and was acquired by invasive GAS serotypes through horizontal gene

transfer (Fig. 1A). It has been proposed that genes carried within RD2 may contribute to

pathogenicity of both GAS and GBS in pregnant women and newborns (16, 17). Supporting this,

AspA has been shown to facilitate GAS biofilm formation and virulence in a murine model of

GAS respiratory infection (18). Recently, *in silico* analysis has revealed four AgI/II gene

homologs in GBS, designated Group <u>B</u> *Streptococcus* <u>s</u>urface <u>p</u>roteins (BspA-D), that are

distributed among GBS of different capsular serotypes.

Previous work has shown that BspA and BspB are found in GBS strain NEM316. BspA has been

demonstrated to be important in biofilm formation as well as adherence to epithelial cells and

may play a role in facilitating colonization through its ability to bind to vaginal epithelium as

well as interact with the hyphal filaments of *Candida albicans* (15, 19), a frequent fungal

colonizer of the lower female reproductive tract. Other GBS strains contain the homolog BspC,

or in some cases BspD, which is over 99% identical to BspC, with the major difference being

that BspD is missing the leader peptide for targeting to the cell surface by the Sec translocation

machinery (15). To date, there have not been any studies examining the impact of the AgI/II

proteins on GBS invasive disease. Because the majority of GBS sequenced strains contain BspC,

we investigated the role of BspC in the pathogenesis of meningitis. Using targeted mutagenesis,

we show that BspC promotes adherence of bacteria to human cerebral microvascular endothelial

cells (hCMEC) and interacts with the host cytoskeleton component, vimentin. Additionally, we

found that BspC and vimentin contribute to the development of GBS meningitis in a mouse

infection model. Lastly, we observe that BspC stimulates inflammatory signaling from brain

endothelial cells *in vitro* and *in vivo* and that this immune signaling involves the NF-κB pathway.

RESULTS

Analysis of BspC domain architecture and construction of a *bspC* deletion strain.

BspC contains all six domains characteristic of the AgI/II protein family, and shares high

homology with other streptococcal AgI/II proteins, especially GAS AspA (Fig 1B). The

proposed domain organization of streptococcal AgI/II polypeptides comprises a stalk consisting

of the α-helical A (alanine-rich repeats) domain and the polyproline II (PPII) helical P domain,

separating the V (variable) domain and the C-terminal domain, which contains the LP*X*TG motif

required for cell wall anchorage (14). While the GBS BspC structure is not known, the structure

of several regions of the GBS homolog, BspA, has been solved (15). We generated a

hypothetical model of full length BspC using PyMOL (The PyMOL Molecular Graphics System,

Version 2.1 Schrödinger, LLC) for the purpose of showing the overall domain structure (Fig 1C).

Structures of individual BspC domains were generated using Phyre2 server (20). The V- and C-

domains were modeled on the V- and C-domains of BspA (PDB entries 5DZ8 and 5DZA,

respectively) (15), and approximately two-thirds of the A-domain sequence was modeled on

human fibrinogen (PDB entry 3GHG) (21). It was not possible to generate models for the N- and

P-domains, so the N-domain is shown as a sphere and the P-domain shown is a mirror image of

the A-domain.

We performed precise in-frame allelic replacement to generate a Δ*bspC* mutant in GBS strain

COH1, a hypervirulent GBS clinical isolate that is highly associated with meningitis (sequence

type [ST]-17, serotype III) (22, 23), using a method as described previously (10). Growth curve

analysis demonstrated that the Δ*bspC* mutant grew similarly to the WT parental strain under the

conditions used here (S1A Fig). Similarly, we observed no differences in hemolytic activity or

capsule abundance between the WT and mutant strains (S1B-D Fig). High-magnification

scanning electron microscopy (SEM) images of COH1 strains showed that the Δ*bspC* deletion

strain exhibits similar surface morphology to the isogenic wild type (Fig 2A and 2B). However,

lower-magnification SEM revealed that the Δ*bspC* mutant appeared to exhibit decreased

interaction between neighboring cells (Fig 2C and 2D).

BspC promotes bacterial adherence to endothelial cells *in vitro*.

Since AgI/II proteins are known to exhibit adhesive properties in other streptococci (14), we

hypothesized that BspC would contribute to GBS interaction with brain endothelium. Thus, we

characterized the ability of the Δ*bspC* mutant to attach to and invade hCMEC using our

established adherence and invasion assays (10, 24). The Δ*bspC* mutant exhibited a significant

decrease in adherence to hCMEC compared to WT GBS, and this defect was complemented

when BspC was expressed in the Δ*bspC* mutant strain (Fig 2E). This resulted in less recovery of

intracellular Δ*bspC* mutant (Fig. 2F), but together these results indicate that BspC contributes

primarily to bacterial attachment to hCMEC. To determine if BspC was sufficient to confer

adhesion, we heterologously expressed the GBS *bspC* gene in the non-adherent, non-pathogenic

bacterium *Lactococcus lactis*. BspC expression resulted in a significant increase in *L. lactis*

adherence to hCMEC compared to *L. lactis* containing the control vector, while invasion was not

affected (Fig 2G and 2H). These results demonstrate that BspC is both necessary and sufficient

to confer bacterial adherence to hCMEC.

BspC contributes to pathogenesis of GBS meningitis *in vivo*.

Our results thus far suggest a primary role for BspC in GBS adherence to brain endothelium. We

hypothesized that these *in vitro* phenotypes would translate into a diminished ability to penetrate

the BBB and produce meningitis *in vivo*. Using our standard model of GBS hematogenous

meningitis (10, 25, 26), mice were challenged with either WT GBS or the Δ*bspC* mutant as

described in the Methods. The WT GBS strain caused significantly higher mortality than the

isogenic Δ*bspC* mutant (Fig 3A). By 48 hours, 80% of mice infected with WT COH1 had

succumbed to death, while all of the mice infected with the Δ*bspC* mutant survived to the

experimental endpoint. In a subsequent experiment mice were infected with a lower dose of

either WT GBS or the Δ*bspC* mutant and were sacrificed at a defined endpoint (48 hrs) to

determine bacterial loads in blood, lung, and brain tissue. We recovered similar numbers of the

Δ*bspC* mutant strain from mouse blood and lung compared to WT, however we observed a

significant decrease in the amounts of the Δ*bspC* mutant recovered from the brain tissue (Fig

3B). To confirm these results using other GBS strains, we constructed *bspC* gene deletions as

described in the Methods in two other GBS strains: GBS 515 (serotype Ia) and meningeal isolate

90356 (serotype III). Mice infected with these mutant strains were also less susceptible to

infection and exhibited decreased bacterial loads in the brain compared to the isogenic parental

WT strain (S2 Fig).

As excessive inflammation is associated with CNS injury during meningitis, we performed

histological analysis of brains from infected animals. In WT infected mice we observed

leukocyte infiltration and meningeal thickening characteristic of meningitis that was absent in the

mice infected with the Δ*bspC* mutant strain (Fig 3C and 3D). Representative images from 8

mice, 4 infected with WT (Fig 3C) and 4 infected with the Δ*bspC* mutant (Fig 3D) are shown. In

subsequent experiments to quantify the inflammatory infiltrate, brain tissue from uninfected mice

and mice infected with either WT GBS or Δ*bspC* mutant GBS was processed and analyzed by

flow cytometry as described in the Methods. There was no significant difference in the numbers

of CD45 positive cells between the groups of mice, however within the CD11b positive

population we observed higher numbers of Ly6C positive and Ly6G positive cells in the brains

of animals infected with WT GBS compared to uninfected mice and mice infected with the

Δ*bspC* mutant strain (Fig 3E and 3F), indicating an increased population of monocytes and

neutrophils. Consistent with these results, we further observed that mice challenged with WT

GBS had significantly more neutrophil chemokine, KC, as well as the proinflammatory cytokine,

IL-1β, in brain homogenates than from Δ*bspC* mutant infected animals. (Fig 3G and 3H)

BspC is necessary and sufficient to induce neutrophil chemokine signaling.

To further characterize the role of BspC in stimulating immune signaling pathways we infected

hCMEC with WT GBS, the Δ*bspC* mutant, or the complemented strain. After four hours of

infection, we collected cells and isolated RNA for RT-qPCR analysis to quantify IL-8 and

CXCL-1 transcripts. We focused on the neutrophil chemokines IL-8 and CXCL-1 as these

cytokines are highly induced during bacterial meningitis (27) and we observed an increase in

neutrophilic infiltrate in brain tissue during the development of GBS meningitis in our mouse

model. Cells infected with WT GBS had significantly higher amounts of both transcripts

compared to cells infected with the Δ*bspC* mutant. Complementation of the *bspC* mutation

restored the ability of the bacteria to stimulate the expression of IL-8 and CXCL-1 (Fig 4A and

4B). Additionally, treatment of hCMEC with purified BspC protein resulted in increased

transcript abundance (Fig 4C and 4D) and protein secretion (Fig 4E and 4F) for both IL-8 and

CXCL-1 compared to untreated cells or treatment with control protein, CshA from *Streptococcus*

gordonii, that was similarly purified from *E. coli*.

Nuclear factor-κB (NF-κB) represents a family of inducible transcription factors, which

regulates a large array of genes involved in different processes of the immune and

inflammatory responses, including IL-8, CXCL-1 and IL-1 (28). To assess whether the NF- κB

pathway is activated by BspC, we utilized the Hela-57A NF- κB luciferase reporter cell line as

described previously (29). Cells infected with WT GBS had significantly higher luciferase

activity than uninfected and cells infected with Δ*bspC* mutant GBS, indicating that BspC

contributes NF- κB activation (Fig 4G). Additionally, immunofluorescent staining of hCMEC

revealed an increase in P65 expression, an indicator of NF- κB activation, during infection with

WT GBS but not in response to infection with Δ*bspC* mutant GBS (Fig 4I-4K).

BspC interacts with the host endothelial cytoskeletal component vimentin.

We next sought to identify the host protein receptor on brain endothelial cells that interact with

BspC. Membrane proteins of hCMEC were separated by 2-dimensional electrophoresis (2-DE),

then blotted to a PVDF membrane. Following incubation with biotinylated BspC protein, the

PVDF membrane was incubated with a streptavidin antibody conjugated to HRP. While many

proteins were detected on the Coomassie stained gel, biotinylated BspC protein specifically

interacted predominately with one spot in the PVDF membrane with molecular mass around 50-

55 kDa and isoelectric point (pI) around 5 (Fig 5A and 5B). The corresponding spots from the

Coomassie-stained 2-DE gel were excised and digested with trypsin (Fig 5C). Resulting peptides

were analyzed by liquid chromatography-tandem mass spectrometry. The spectra from the spot

yielded 158 peptide sequences, which matched to human vimentin. The molecular weight, 53.6

kDa, and calculated pI, 5.12, of vimentin match the value for the spot on the 2-DE gels. This

procedure was repeated for membrane proteins from another human brain endothelial cell line

(hBMEC) that has been used previously to study GBS interactions (30). Mass spectrometry

analysis also determined that BspC interacted with human vimentin (S3A-S3C Fig). The control

far Western blot with the streptavidin antibody conjugated to HRP did not show any

hybridization (S3D Fig).

To confirm these protein-protein interactions *in vivo*, we employed a bacterial two-hybrid system

(BACTH, "Bacterial Adenylate Cyclase-Based Two-Hybrid") (31). This system is based on the

interaction-mediated reconstitution of a cyclic adenosine monophosphate (cAMP) signaling

cascade in *Escherichia coli*, and has been used successfully to detect and analyze the interactions

between a number of different proteins from both prokaryotes and eukaryotes. (32) Using a

commercially available kit (Euromedex) according to manufactures directions and as described

previously (33), vimentin was cloned and fused to the T25 fragment as a N-terminal fusion (T25-

Vimentin), using the pKT25 plasmid, and *bspC* was cloned as a c-terminal fusion (BspC-T18)

using the pUT18 plasmid. To test for interaction these plasmids were transformed into an *E. coli*

strain lacking adenylate cyclase (*cyaA*). We observed blue colonies when grown on LB agar

plates containing X-gal, indicating β-galactosidase activity and a positive interaction between

Vimentin and BspC compared to the empty vector control (Fig 5D and 5E). A leucine zipper that

is fused to the T25 and T18 fragments served as the positive control for the system (Fig 5F). To

quantify β-galactosidase activity, cells grown to log phase were permeabilized with 0.1 % SDS

and toluene and enzymatic activity measured by adding ONPG as described previously (33). The

E. coli strain containing both vimentin and BspC expressing plasmids exhibited increased β-

galactosidase activity (Miller units) compared to the empty vector control strain (Fig 5G). These

results demonstrate a direct interaction between BspC and vimentin.

GBS co-localization with cell-surface vimentin is dependent on BspC.

To visualize the localization of WT GBS and vimentin in brain endothelial cells we performed

imaging flow cytometry of uninfected and WT GBS infected hCMEC. Cells were fixed,

permeabilized, and incubated with antibodies to vimentin and GBS. We observed that vimentin

protein was present throughout the cytoplasm of uninfected hCMEC (Fig 6A and 6B), but during

infection GBS co-localized with vimentin near the surface of infected cells (Fig 6C and 6D). To

further characterize the localization of GBS and vimentin, we performed immunofluorescent

staining of hCMEC infected with either WT GBS or the Δ*bspC* mutant. Following infection,

cells were fixed but not permeabilized to permit labeling of only extracellular bacteria and

surface expressed vimentin. We observed that surface vimentin of hCMEC co-localized with WT

GBS bacteria, while this was not seen for hCMEC infected with the Δ*bspC* mutant (Fig 6E-6J).

Vimentin contributes to the pathogenesis of GBS infection.

To first determine if BspC-mediated attachment to hCMEC is dependent on vimentin, we

assessed the effect of blocking the vimentin-GBS interaction by treating hCMEC with anti-

vimentin antibodies prior to infection with GBS. Pre-incubation with the mouse V9 antibody

(34), which reacts with the C-terminal of vimentin (AA392-466), reduced WT GBS adherence to

levels comparable to the adherence of the Δ*bspC* mutant (Fig 7A). Treatment with a vimentin

antibody that recognizes the N-terminal epitope (AA31-80), as well as with IgG isotype controls

did not alter adherence of the WT or Δ*bspC* mutant strains (Fig 7A and 7B). These results

indicate that the interaction between BspC and cell-surface vimentin is dependent on the C-

terminus of vimentin.

We also infected vimentin KO mice with WT GBS and observed that they were less susceptible

to GBS infection and exhibited increased survival compared to WT animals (Fig 7C). Further,

significantly less bacteria were recovered from the tissues of KO mice compared to the tissues of

WT mice (Fig 7D). Additionally, we detected significantly less KC and IL-1β in brain tissues of

vimentin KO mice infected with GBS compared to WT mice, suggesting that vimentin

contributes to the initiation of immune signaling pathways during GBS infection. Taken together

these results indicate the importance of vimentin in GBS dissemination into the brain and

meningitis disease progression.

DISCUSSION

Our studies reveal a unique requirement for the Group B streptococcal Antigen I/II protein,

BspC, to brain penetration by GBS, the leading agent of neonatal bacterial meningitis. A

decreased ability by the GBS Δ*bspC* mutant to attach to brain endothelium and induce neutrophil

chemoattractants *in vitro* was correlated with a reduced risk for development of meningitis and

markedly diminished lethality *in vivo*. We identified that BspC interacts directly with host

vimentin and that blocking this interaction abrogated BspC-mediated attachment to hCMEC.

Further, vimentin deficient mice infected with GBS exhibited decreased mortality, bacterial brain

loads, and cytokine production in brain tissue. These results corroborate the growing evidence

that this intermediate filament protein plays important roles in the pathogenesis of bacterial

infections (35), and provide new evidence for the pivotal role of the BspC adhesin in GBS CNS

disease (Fig 8).

The oral streptococcal AgI/II adhesins range in composition from 1310 - 1653 amino acid (AA)

residues, while GBS AgI/II proteins are smaller (826-932 AA residues) (36). The primary

sequences of AgI/II proteins are comprised of six distinct regions (Fig. 1B), several of which

have been shown to mediate the interaction to various host substrates. The *S. mutans* protein

SpaP as well as the *S. gordonii* proteins SspA and SspB have been demonstrated to interact

specifically with the innate immunity scavenger protein gp-340. (37) Recently, the GAS AgI/II

protein AspA, as well as BspA, the AgI/II homolog expressed mainly by the GBS strain

NEM316, have also been shown to bind to immobilized gp-340 (15, 18). Gp-340 proteins are

involved in various host innate defenses and are present in mucosal secretions, including saliva

in the oral cavity and bronchial alveolar fluid in the lung. They can form complexes with other

mucosal components such as mucins and function to trap microbes for clearance. However,

when gp-340 is immobilized, it can be used by bacteria as a receptor for adherence to the host

surface. (15, 38-41). There is evidence that the Variable (V) regions of SspB, SpaP, AspA and

BspA facilitate gp340-binding activity (15, 42-44). AgI/II family adhesins have also been shown

to interact with other host factors including fibronectin, collagen and β1 integrins to promote

host colonization (14, 45-47), demonstrating the multifactorial nature of these adhesins.

Here we show that GBS BpsC interacts with host vimentin, an important cytoskeletal protein

belonging to class III intermediate filaments. Vimentin is located in the cytoplasm and functions

as an intracellular scaffolding protein that maintains structural and mechanical cell integrity (48).

However, vimentin is also found on the surface of numerous cells such as T cells, platelets,

neutrophils, activated macrophages, vascular endothelial cells, skeletal muscle cells and brain

microvascular endothelial cells (49-56). Vimentin also mediates a variety of cell processes

including cell adhesion, immune signaling, and autophagy (51, 57, 58). Further, the role of cell

surface vimentin as an attachment receptor facilitating bacterial or viral entry, has been

previously documented for other pathogens (35, 59-61). The BspC domain that mediates the

vimentin interaction is currently under investigation. As the V domain is likely projected from

the cell surface and has been implicated in host interactions for other streptococci, we

hypothesize that this may be a critical domain for this interaction. Additionally, it is unknown if

GBS BspC interactions with host factors are similar to those of the AgI/II proteins from oral

streptococci, particularly since the respective V domains of these homologs are distant enough to

suggest different binding partners. However, as the V-domain of BspA protein from GBS, which

shares 96% identity to BspC, interacts with gp340, it is likely that BspC will similarly interact

with gp340. This requires further investigation as it may be critical to GBS colonization of

mucosal surfaces such as the gut and the vaginal tract.

There is a growing body of evidence that various bacteria can interact with vimentin to promote

their pathogenesis, including *Escherichia coli* K1, *Salmonella enterica*, GAS, and *Listeria*

monocytogenes (35, 53, 56, 62, 63). Interestingly, previous studies on the meningeal pathogen *E.*

coli K1 have demonstrated that the bacterial surface factor, IbeA, interacts with vimentin to

promote bacterial uptake into brain endothelial cells (56, 64). Similarly, while our study was

underway, it was reported that another bacterium capable of causing meningitis, *L.*

monocytogenes, uses InlF to interact with vimentin to promote brain invasion (63). Along with

our results presented here, this may suggest a common mechanism for meningeal bacterial

pathogens to penetrate the BBB and cause CNS disease. However, our analysis of these three

bacteria proteins showed no homology or predicted regions that might commonly interact with

vimentin. Furthermore, the interaction between the *E. coli* receptor IbeA and the *L.*

monocytogenes receptor InlF with cell-surface vimentin can be blocked by an antibody to the N-

terminal region of vimentin (56, 63), while we demonstrate that the interaction between BspC

and cell-surface vimentin can be blocked with an antibody to the C-terminal of vimentin. The

implications of this unique interaction between a bacterial receptor and the C-terminal of

vimentin remain to be explored.

Neuronal injury during bacterial meningitis involves both microbial and host factors, and

subsequent to attachment to the brain endothelium and penetration of the BBB, GBS stimulation

of host immune pathways is the next important step in the progression of meningitis. The release

of inflammatory factors by brain endothelial cells, microglia, astrocytes, and infiltrating immune

cells can exacerbate neuronal injury (9). Our data suggest that, like other streptococcal AgI/II

family polypeptides, BspC plays a role in immune stimulation. AgI/II family proteins contain

two antigenic regions (the antigens I/II and II) (65) and this ability to elicit an inflammatory

response makes SpaP, the *S. mutans* AgI/II protein, an attractive candidate for vaccine

development to prevent dental caries (14, 66). In this study we found that BspC can stimulate

NF-κB activation and the release of the proinflammatory cytokines IL-8 and CXCL-1 from

hCMEC. Both of these chemokines are major neutrophil recruiting chemoattractants and are the

most highly induced during GBS infection (26, 67). We observed that mice infected with GBS

mutants that lack BspC exhibited lower brain bacterial loads and less meningeal inflammation

compared to animals challenged with WT GBS. Interestingly, WT and Δ*bspC* mutant bacterial

loads were similar in the blood, indicating that BspC may not influence GBS survival and

proliferation in the blood; however further investigation is warranted.

This study demonstrates for the first time the importance of a streptococcal AgI/II protein, BspC,

in the progression of bacterial meningitis, as well as the unique requirement for vimentin to the

pathogenesis of CNS disease. Vimentin KO mice were markedly less susceptible to GBS

infection and exhibited reduced bacterial tissue load and inflammatory signaling. Vimentin is

also known to act as a scaffold for important signaling molecules and mediates the activation of a

variety of signaling pathways including NOD2 (nucleotide-binding oligomerization domain-

containing protein 2) and NLRP3 (nucleotide-binding domain, leucine-rich-containing family

pyrin domain-containing-3) that recognize bacterial peptidoglycan and activate inflammatory

response via NF-κB signaling (64, 68, 69). Thus continued investigation into the mechanisms of

how BspC-vimentin interactions dually promote bacterial attachment and immune responses will

provide important information that may inform future therapeutic strategies to limit GBS disease

progression.

MATERIALS AND METHODS

Bacterial strains, growth conditions and proteins.

GBS clinical isolate COH1 (serotype III) (70), 515 (serotype Ia) (19), the recent meningitis

isolate 90356 (serotype III) (71) and their isogenic Δ*bspC* mutants were used for the

experiments. GBS strains were grown in THB (Hardy Diagnostics) at 37°C, and growth was

monitored by measuring the optical density at 600 nm (OD_{600}). *Lactococcus lactis* strains were

grown in M17 medium (BD Biosciences) supplemented with 0.5% glucose at 30°C. For

antibiotic selection, 2 μg/mL chloramphenicol (Sigma) and 5 μg/mL erythromycin (Sigma) were

incorporated into the growth medium. BspC and CshA recombinant proteins were purified as

described previously (15, 72).

Targeted mutagenesis and complementation vector construction.

The Δ*bspC* mutant was generated in COH1and 90356 by in-frame allelic replacement with a

chloramphenicol resistance cassette by homologous recombination using a method previously

described (10). A knockout construct was generated by amplifying up- and down-flanking

regions of the *bspC* gene from COH1 genomic DNA using primer pairs of 5'flank-F

(GCAGACACCGATTGCACAAGC)/R   (GAAGGCGATCTTGCCCTCAA)   and   3'flank-F

(GTCAGCTATCGGTTTAGCAGG)/R(CTATACACGCCTACAGGTGTC).   The

chloramphenicol   resistance   (*cat*)   cassette   was   amplified   with   primers   Cat-F

(GAGGGCAAGATCGCCTTCATGGAGAAAAAAATCACTGGAT)   and   Cat-R

(CTGCTAAACCGATAGCTGACTTACGCCCCGCCCTGCCACT). Then the construct of two

flanks along with the *cat* cassette was amplified with a pair of nest primers, Nest-xhoI

(CCFCTCGAGGATGCTCAAGATGCACTCAC) and Nest-xbaI

(GCTCTAGACGAGCCAAATTACCCCTCCT), which was then cloned into the pHY304 vector

(73) and propagated in *E. coli* strain DH5α (74) prior to isolation and transformation to COH1

and 90356 GBS. A Δ*bspC* mutant had been generated previously in 515 (19). The complemented

strain of Δ*bspC* mutant in COH1 was generated by cloning *bspC* into pDCerm, an *E. coli*-GBS

shuttle expression vector. Gene *bspC* was amplified from GBS 515 genomic DNA using primers

pDC.*bspC*.F (TGGGTACCAGGAGAAAATATGTATAAAAATCAAAC) and pDC.*bspC*.R

(CCGGGAGCTCGCAGGTCCAGCTTCAAATC), designed to encode a *Kpn*I and *Sac*I

restriction site at its termini respectively. This *bspC* amplicon was then cloned into pDCerm and

propagated in *E. coli* strain Stellar^{TM} (ClonTech), prior to isolation and transformation into

COH1 GBS. A *L. lactis* strain expressing BspC had been generated previously (19).

Hemolysis assay.

GBS strains were grown to an OD$_{600}$ of 0.4, harvested by centrifugation, and resuspended in

PBS. A total of 1 x 10^8 CFU was added to fresh sheep blood (VWR) in V-bottom 96-well plates

(Corning). The plates were sealed and incubated at 37°C with agitation for 1 h. The plates were

centrifuged at 200 x *g* for 10 min, and 100 µl of the supernatant was transferred to a flat-bottom

96-well plate. The absorbance at 541 (*A*$_{541}$) was read, and percent hemolysis was calculated by

comparing the *A*$_{541}$ values for GBS-treated wells to the *A*$_{541}$ values for the wells with blood

incubated with water.

GBS capsule flow cytometry.

Flow cytometry to determine capsule expression was performed as described in (75). Briefly,

bacterial stocks were washed in sterile PBS containing 0.5% bovine serum albumin (BSA)

(VWR) then incubated with a purified monoclonal anti-serotype III antibody or a purified

monoclonal anti-serotype Ia isotype control at a 1:10,000 dilution, washed via centrifugation,

and labeled with a donkey anti-mouse IgM conjugated to AlexaFluor647 (Invitrogen) at a

1:2,000 dilution. All incubations were performed at 4°C with shaking. Samples were washed

again then resuspended and read on a FACScalibur flow cytometer (BD Biosciences), and

analyzed using FlowJo (v10) software. The monoclonal antibodies were provided by John

Kearney at the University of Alabama at Birmingham.

Scanning electron microscopy.

Bacteria were grown to log phase and were then fixed for 10 min using a one-step method with

2.5% glutaraldehyde, 1% osmium tetraoxide, 0.1M sodium cacodylate. Bacteria were collected

on 0.4 μM polycarbonate filters by passing the solution through a swinnex device outfitted on a

10 mL syringe. The filters were dehydrated through a series of increasing ethanol concentrations

and then dried in a Tousimis SAMRI-790 critical point drying machine. The dried filters were

mounted on SEM sample stubs with double-sided carbon tape, coated with 6nm platinum using a

Quorom Q150ts high-resolution coater and imaged with a FEI FEG450 scanning electron

microscope.

Cell lines and infection assays.

Cells of the well-characterized human cerebral microvascular endothelial cell line (hCMEC/D3),

referred to here as hCMEC were obtained from Millipore and were maintained in an EndoGRO-

MV complete medium kit supplemented with 1 ng/ml fibroblast growth factor-2 (FGF-2;

Millipore) (76-79). Hela57A were provided by Marijke Keestra-Gounder at the University of

Colorado, Anschutz Medical Campus and cultured in DMEM (Corning Cellgro) containing 10%

fetal bovine serum (Atlanta Biologicals). Assays to determine the total number of hCMEC

surface-adherent or intracellular bacteria were performed as describe previously (10). Briefly,

bacteria were grown to mid-log phase to infect hCMEC monolayers (1×10^5 CFU, at a

multiplicity of infection [MOI] of 1). Total cell-associated GBS and *L. lactis* were recovered

following a 30 min incubation, while intracellular GBS were recovered after 2 h infection and 2

h incubation with 100 μg gentamicin (Sigma) and 5 μg penicillin (Sigma) to kill all extracellular

bacteria. Cells were detached with 0.1 ml of 0.25% trypsin-EDTA solution and lysed with

addition of 0.4 ml of 0.025% Triton X-100 by vigorous pipetting. The lysates were then serially

diluted and plated on THB agar to enumerate bacterial CFU. For antibody pre-treatment assays,

hCMEC were incubated with 0.3μg/ml antibodies for 30 min prior to infection with GBS. The

mouse monoclonal antibody to vimentin, clone V9 (Abcam), the rabbit polyclonal antibody to N-

terminal vimentin (Sigma), and the isotype controls (VWR) were used. Total cell-associated

GBS were recovered following a 1h incubation. For luciferase assays, Hela57A cells were

infected with 1×10^6 CFU (MOI, 10) GBS for 90 min. Cells were then lysed and luciferase

activity quantified using a luciferase assay system (Promega) according to manufacturer's

instructions.

Ethics statement.

Animal experiments were approved by the committee on the use and care of animals at San

Diego State University (SDSU) protocol #16-10-021D and at University of Colorado School of

Medicine protocol #00316 and performed using accepted veterinary standards. The San Diego

State University and the University of Colorado School of Medicine are AAALAC accredited;

and the facilities meet and adhere to the standards in the "Guide for the Care and Use of

Laboratory Animals".

Mouse model of hematogenous GBS meningitis.

We utilized a mouse GBS infection model as described previously (10, 25, 26). Briefly, 8-week

old male CD-1 mice (Charles River), 129 WT, or 129S-Vim^{tm1Cba}/MesDmarkJ (Vimentin KO)

(Jackson Laboratory) were injected intravenously with 1×10^9 CFU of wild-type GBS or the

isogenic Δ$bspC$ mutant for a high dose challenge, or 1×10^8 CFU for a low dose challenge. At

the experimental endpoint mice were euthanized and blood, lung, and brain tissue were collected.

The tissue was homogenized, and the brain homogenates and lung homogenates as well as blood

were plated on THB agar for enumeration of bacterial CFU.

Histology.

Mouse brain tissue was frozen in OCT compound (Sakura) and sectioned using CM1950 cryostat

(Leica). Sections were stained using hematoxylin and eosin (Sigma) and images were taken

using a BZ-X710 microscope (Keyence).

Flow cytometry.

At 48 h post-infection with 1×10^8 CFU of GBS, mice were euthanized then perfused to replace

blood with PBS. The entire mouse brain was harvested from each animal and brains tissue was

processed with the Multi-Tissue Dissociation kit #1 following the Adult brain dissociation

protocol (Miltenyi Biotec). Cells were resuspended in MACS buffer (Miltenyi Biotec) and

incubated with antibodies to Ly6C conjugated to BV421, CD45 conjugated to PE, CD11b

conjugated to FITC, and Ly6G conjugated to APC (Invitrogen) at 1:200 dilution, UltraLeaf anti-

mouse CD16/CD32 Fc block (Biolegend) at 1:400 dilution, and fixable viability dye conjugated

to eFLuor506 (eBioscience) at 1:1000 dilution for 1 h, then fixed (eBioscience). Cells were

counted using a Countess automated cell counter (Invitrogen), read on a Fortessa X-20 flow

cytometer (BD Biosciences), and analyzed using FlowJo (v10) software. Gates were drawn

according to fluorescence minus one (FMO) controls.

RT-qPCR and ELISA.

GBS were grown to mid-log phase and 1×10^6 CFU (MOI, 10) were added to hCMEC

monolayers and incubated at 37°C with 5% CO_2 for 4 h. Cell supernatants were collected, the

cells were then lysed, and total RNA was extracted (Machery-Nagel) and cDNA was synthesized

(Quanta Biosciences) according to the manufacturers' instructions. Primers and primer

efficiencies for IL-8, CXCL-1, and GAPDH (glyceraldehyde-3-phosphate dehydrogenase) were

utilized as previously described (80). IL-8 and CXCL-1 from hCMEC supernatants, and KC and

IL-1β from mouse brain homogenates were detected by enzyme-linked immunosorbent assay

according to the manufacturer's instructions (R&D systems).

Immunofluorescent staining.

hCMEC were grown to confluency on collagenized coverslips (Fisher). Following a 1 h

infection, cells were washed with PBS to remove non-adherent bacteria and fixed with 4%

paraformaldehyde (Sigma) for 30 min. For Figure 4, cells were incubated with 1%BSA in PBS

with 0.01% Tween-20 (Research Products International) to block non-specific binding for 15

min, then incubated with a rabbit antibody to p65 (Sigma) at a 1:200 dilution overnight at 4°C.

Coverslips were then washed with PBS and incubated with donkey anti-rabbit conjugated to Cy3

(Jackson Immunoresearch) at a 1:500 dilution for 1 h at room temperature. For Figure 6, cells

were incubated with 1% BSA in PBS for 15 min, then with antibodies to vimentin (Abcam) and

GBS (Genetex) at a 1:200 dilution overnight at 4°C. Following washes with PBS and an

incubation with donkey anti-mouse conjugated to Cy3 and donkey anti-rabbit conjugated to 488

secondary antibodies (ThermoFisher) at a 1:500 dilution for 1 h at room temperature, coverslips

were washed with PBS and mounted onto glass microscopy slides (Fisher) with VECTASHIELD

mounting medium containing DAPI (Vector Labs). Cells were imaged using a BZ-X710

fluorescent microscope (Keyence).

2-dimensional electrophoresis (2-DE), Far Western blot, and mass spectrometry analysis.

Membrane proteins of hCMEC cells were enriched using a FOCUS membrane protein kit (G

Biosciences, St. Louis, MO), dissolved in rehydration buffer (7M urea, 2M thiolurea, 1% TBP,

and 0.2% ampholytes 3-10 NL), and quantified using 2D Quant kit (GE Healthcare, Piscataway,

NJ). Proteins (100μg) were loaded on 7-cm long immobilized pH gradient (IPG) strips with non-

linear (NL) 3-10 pH gradient (GE Healthcare). Isoelectric focusing was carried out in Multiphor

II electrophoresis system (GE Healthcare) in three running phase (phase 1: 250V/0.01h, phase 2:

3500V/1.5h, and phase 3: 3500V/ 4.5h). The second dimension SDS-PAGE was carried out

using 12.5% acrylamide gels in duplicate. One gel was stained with Coomassie Blue G250 (Bio-

Rad, Hercules, CA) for mass spectrometry analysis. The other gel was transferred to a PVDF

membrane for far Western blot analysis.

The PVDF membrane was denatured and renatured as described in (81), followed by

incubation in a blocking solution (5% skim milk in PBS) for 1 h. Recombinant BspC was

biotinylated using a EZ-Link Sulfo-NHS-Biotin kit (ThermoFisher Scientific, Waltham, MA).

The PVDF membrane was probed with the biotinylated BspC (100µg) in a blocking solution

overnight at 4°C. After washing three times with a washing buffer (PBS, 0.05% Tween-20), the

PVDF membrane was incubated with an antibody conjugated to streptavidin-horse radish

peroxidase (HRP). Interacting proteins were detected by adding enhanced chemiluminescence

(ECL) reagents (ThermoFisher Scientific) and visualized by x-ray film exposure. The protein

spots from far Western blot were aligned to the corresponding protein spots in the Coomassie

stained gel. The identified spots were excised and digested in gel with trypsin (Worthington,

Lakewood, NJ). Peptide mass spectra were collected on MALDI-TOF/TOF, (ABI 4700, AB

Systems, Foster City, CA) and protein identification was performed using the automated result

dependent analysis (RDA) of ABI GPS Explorer softwareV3.5. Spectra were analyzed by the

Mascot search engine using the Swiss protein database.

Bacterial two-hybrid assay.

A bacterial adenylate cyclase two-hybrid assay was performed as in (31) and following

manufacturer's instructions (Euromedex). Briefly, plasmids containing T25-Vimentin and BspC-

T18 were transformed into *E. coli* lacking *cyaA* and *E. coli* were plated on LB plates containing

X-gal (Sigma). To measure β -galactosidase activity, Miller assays were performed according to

standard protocols (82). Briefly, *E. coli* were grown in 0.5mM IPTG (Sigma), then permeabilized

with 0.1% SDS (Sigma) and toluene (Sigma). ONPG (Research Products International) was

added and absorbance was measured at 600, 550, and 420nm.

Imaging flow cytometry.

Following a 1 h infection with GBS, hCMEC were washed with PBS to remove non-adherent

GBS. Cells were collected using a cell scraper (VWR) and resuspended in PBS containing 10%

FBS and 1% sodium azide (Sigma). Cells were incubated with primary antibodies to vimentin

(Abcam) and GBS (Genetex) at 1:500 dilution for 1 h at 4°C followed by donkey anti-mouse

conjugated to Cy3 and donkey anti-rabbit conjugated to 488 (ThermoFisher) secondary

antibodies. Cells were then fixed (eBioscience) and analyzed using an ImageStream X imaging

flow cytometer (Amnis).

Data analysis.

GraphPad Prism version 7.0 was used for statistical analysis and statistical significance was

accepted at P values of <0.05. (∗, $P < 0.05$; ∗∗, $P < 0.005$; ∗∗∗, $P < 0.0005$; ∗∗∗∗, $P < 0.00005$).

Specific tests are indicated in figure legends.

ACKNOWLEDGEMENTS

We thank Dr. Marijke Keestra-Gounder (University of Colorado, Anschutz Medical Campus) for

the Hela57A reporter cell line, Tibor Pechan of the Mississippi State University Institute for

Genomics, Biocomputing & Biotechnology (IGBB) for assistance with the mass spectrometry,

Jeremy Rakhola at the Rocky Mountain Regional VA Medical Center (RMRVAMC) Flow Core

for imaging flow cytometry, and Catherine Back (University of Bristol, UK) for provision of

purified *S gordonii* CshA protein. Also to Katilynn Croom and Jeffrey Kavanugh for technical

support. This study was supported by the Rees Stealy Research Foundation/ SDSU Heart

Institute and San Diego Chapter ARCS Scholarships to L.D., the Coordenação de

Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001 to G.F.S.,

the NIH DE016690 to H.F.J., and the NIH/NINDS R01NS051247 to K.S.D.

    FIGURE LEGENDS

    Figure 1. Analysis of BspC domain architecture (A) Diagram of MGAS6180 region of difference

    2 (RD2). M2, M4, and M28 strains of Group A *Streptococcus* contain RD2, which was likely

    acquired from Group B *Streptococcus* through horizontal gene transfer and is absent in other

    common disease causing GAS *emm*-types. Open reading frames encoding LPXTG cell wall

    anchor domain-containing proteins are indicated with green arrows. Proteins AspA and R28 in

    GAS and their respective homologs in GBS, BspC and Rib, have been previously described. The

    other two cell wall anchor domain-containing proteins have not been characterized. (B)

    Schematic of sequence homologies across antigen I/II family proteins from select *Streptococcus*

    species. Six structural regions are shown: N, N-terminal region; A, alanine-rich repeats; V,

    variable region; P, proline-rich repeats; C, C-terminal region; and the cell-wall anchor-containing

    region. Numbers indicate percentage amino acid residue identities to BspC from *S. agalactiae*

    strain COH1 (NCBI Ref. Seq.: WP_000277676.1). The protein depicted are the antigen I/II

    proteins AspA from *S. pyogenes* strain M28 (WP_011285012.1), Pas from *S. intermedius*

    (WP_049476098.1), SspB from *S. gordonii* (WP_011999747), and SpaP from *S. mutans*

    (WP_024781655.1). (C) Hypothetical model of BspC generated using PyMOL.

    Figure 2. BspC bacterial adherence to endothelial cells *in vitro*. (A-D) Scanning electron

    microscopy images of WT (A and C) and Δ*bspC* mutant (B and D) GBS. Scale bar in high

    magnification images (A and B) is 1 μM. Scale bar in low magnification images (C and D) is 50

    μM. (E) Adherence of WT GBS, the Δ*bspC* mutant, and the Δ*bspC*p.*bspC* complemented strain

    to hCMEC was assessed after a 30 min incubation. Total cell-associated bacteria are shown. (F)

Invasion of WT GBS, the Δ*bspC* mutant, and the Δ*bspC*p.*bspC* strain was quantified after a 2 h

infection. (G) Adherence and (H) invasion of *Lactococcus lactis* containing the pMSP empty

vector or pMSP.*bspC* to hCMEC were quantified. Data indicates the percentage of the initial

inoculum that was recovered. Experiments were performed three times with each condition in

triplicate. Data from one representative experiment are shown and error bars represent the

standard deviation. Statistical analysis: (E and F) One-way ANOVA with Tukey's multiple

comparisons test. (G and H) Unpaired t test. $*$, $P < 0.05$.

Figure 3. BspC contributes to pathogenesis of GBS meningitis *in vivo*. (A) A Kaplan-Meier plot

showing survival of mice infected with 10^9 CFU of either WT GBS or the Δ*bspC* mutant. (B) 72

h after infection with 10^8 CFU of either WT GBS or the Δ*bspC* mutant, mice were euthanized

and bacterial loads in brain, lung, and blood were quantified. Lines indicate statistical median. (C

and D) H&E images showing the leptomeninges on the surface of brains from mice infected with

either WT GBS (C) or the Δ*bspC* mutant (D). Arrows indicate areas of meningeal thickening and

leukocyte infiltration. Scale bar is 100 μM. (E and F) Quantification of infiltrating immune cells

in the brains of mice infected with WT GBS or the Δ*bspC* mutant. The presence CD45+,

CD11b+, Ly6C+ (E), and CD45+, CD11b+, Ly6G+ (F) cells was determined using flow

cytometry. (G and H) ELISA was performed on mouse brain tissue homogenates to assess

cytokine protein levels. KC (G) and IL-1β (H) were quantified for brains from mice challenged

with either WT GBS or the Δ*bspC* mutant. Statistical analysis: (A) Log-rank test. (B) Two-way

ANOVA with Sidak's multiple comparisons test. (E and F) One-way ANOVA with Sidak's

multiple comparisons test. (G and H) Mann-Whitney test. $*$, $P < 0.0005$; $**$, $P < 0.005$; $***$, $P <$

0.0005; $****$, $P < 0.00005$.

Figure 4. BspC is necessary and sufficient to induce neutrophil chemokine signaling. (A and B)

RT-qPCR to assess IL-8 (A) and CXCL-1 (B) transcript levels in uninfected hCMEC or hCMEC

infected with WT GBS, the Δ*bspC* mutant, or the complemented strain. (C and D) RT-qPCR

analysis quantifying IL-8 (C) and CXCL-1 (D) transcripts in hCMEC treated with purified BspC

protein or the *S. gordonii* CshA protein. (E and F) ELISA to measure IL-8 (E) and CXCL-1 (F)

protein secretion by hCMEC treated with BspC or CshA. (G) Luciferase activity of uninfected

Hela-57A cells and cells infected with WT GBS or the Δ*bspC* mutant was assessed. Experiments

were performed at least three times with each condition in triplicate. Data from representative

experiments are shown and error bars represent the standard deviation of the mean for one

experiment. Data were analyzed with One-way ANOVA with Sidak's multiple comparisons test.

(H-J) Immunofluorescent staining showing p65 localization in uninfected hCMEC (H), hCMEC

infected with WT GBS (I), and hCMEC infected with the Δ*bspC* mutant (J). Following infection,

cells were fixed with formaldehyde and incubated with a rabbit antibody to p65. Nuclei were

labelled with DAPI. Scale bar is 50 µM. ∗, $P < 0.0005$; ∗∗, $P < 0.005$; ∗∗∗, $P < 0.0005$; ∗∗∗∗, *P*

< 0.00005.

Figure 5. BspC interacts with the host endothelial cytoskeletal component vimentin. (A) Far

Western blot analysis of hCMEC membrane proteins probed with biotinylated BspC. Membrane

proteins were extracted from hCMEC and separated by 2-DE in duplicate. One gel was

transferred to a PVDF membrane and probed with biotinylated BspC. The specific interaction of

BspC was detected by a streptavidin antibody conjugated to HRP and visualized by x-ray film

exposure. The other gel was stained with Coomassie blue. The spot identified from the x-ray film

was aligned to the Coomassie stained gel and the corresponding spot was excised and digested

with trypsin. (B) Electrospray ionization-tandem mass spectrometry analysis identifies the

protein vimentin. (C) Amino acid sequence of human vimentin. Underscored and bolded are the

peptide sequences identified from MS analysis. (D-F) Bacterial two-hybrid assay using *E. coli*

containing empty vector controls (D), the BspC and vimentin vectors (E), and the positive

control vectors (F). (G) β-galactosidase activity of *E. coli* was quantified using a Miller assay.

Experiments were performed at least three times with each condition in five replicates. Data

from one representative experiment is shown, error bars represent the standard deviation of the

mean. Data were analyzed using one-way ANOVA with Dunnett's multiple comparisons test.

****, $P < 0.00005$.

Figure 6. GBS co-localization with cell-surface vimentin is dependent on BspC. (A-D) Imaging

flow cytometry showing vimentin localization in uninfected hCMEC (A and B) and hCMEC

infected with WT GBS (C-D). Cells were fixed and permeabilized prior to staining with

antibodies to vimentin and the group B carbohydrate antigen to label cell-associated GBS. (E-J)

Immunofluorescent staining of hCMEC monolayers infected with WT GBS (E-G) or the Δ*bspC*

mutant (H-J). Following a one-hour infection, hCMEC were washed to remove nonadherent

bacteria then fixed and labelled with antibodies to vimentin and GBS. Nuclei were labelled with

DAPI.

Figure 7. Vimentin contributes to the pathogenesis of GBS infection. (A and B) hCMEC were

pre-incubated with monoclonal antibodies recognizing the C-terminal (A) or the N-terminal (B)

of vimentin or with an IgG control prior to a 30 min infection with either WT GBS or the Δ*bspC*

mutant. Experiments were performed three times with each condition at least in triplicate and a

data from representative experiments are shown. Error bars represent the standard deviation of

the mean. (C) Kaplan-Meier plot showing survival of WT and $Vim^{-/-}$ mice infected with 10^8 CFU

of WT GBS. (D) Tissue bacterial counts for WT and $Vim^{-/-}$ mice. (E and F) ELISA to quantify

KC (C) and IL-1β (D) protein in brain tissue homogenates. Statistical analysis: (A, B, and D)

Two-way ANOVA with Sidak's multiple comparisons test. (C) Log-rank test. (E and F) Mann-

Whitney test. ∗∗, $P < 0.005$; ∗∗∗, $P < 0.0005$; ∗∗∗∗, $P < 0.00005$.

Figure 8. Summary of the role of the BspC-vimentin interaction in promoting meningitis. BspC

interacts with vimentin on the surface of CMEC to promote GBS attachment and the production

of the neutrophil recruiting cytokines IL-8 and CXCL-1 through the NF-κB pathway.

Figure S1. (A) Growth curves for WT GBS and the Δ*bspC* mutant in THB. (B) Hemolysis assay

comparing hemolysis of sheep blood by WT GBS and the Δ*bspC* mutant. Representative data of

one of at least three independent experiments are shown. Error bars represent the standard

deviation of mean in one experiment. Data were analyzed using an unpaired t test. (C and D)

Flow cytometry using a monoclonal antibody to the serotype III capsule to determine the

presence of capsule in WT GBS (C) and the Δ*bspC* mutant (D).

Figure S2. (A) Kaplan-Meier plot showing survival of mice challenged with either WT 515 GBS

or the isogenic Δ*bspC* mutant. (B-D) Tissue bacterial counts for mice infected with WT 515 and

90356 GBS and the isogenic Δ*bspC* mutants. 48h post-infection, mice were sacrificed and

bacterial loads in brain (B), lung (C), and blood (D) were quantified. Statistical analysis: (A)

Log-rank test. (B-D) Two-way ANOVA with Sidak's multiple comparisons test. $*$, $P < 0.0005$;

$**$, $P < 0.005$.

Figure S3. (A) Far Western blot analysis of hBMEC membrane proteins using biotinylated BspC

protein. Two spots (I and II) were identified on the x-ray film and aligned to the Coomassie

stained gel. (B) Electrospray ionization-tandem mass spectrometry identifies spots I and II as

vimentin. (C) The amino acid sequence of human vimentin, with the peptide sequences identified

in the MS analysis underscored and bolded. (D) Control Far Western blot with the streptavidin

antibody conjugated to HRP only.

<u>REFERENCES</u>

1. Thigpen MC, Whitney CG, Messonnier NE, Zell ER, Lynfield R, Hadler JL, et al.

Bacterial meningitis in the United States, 1998-2007. N Engl J Med. 2011;364(21):2016-25.

2. Doran KS, Nizet V. Molecular pathogenesis of neonatal group B streptococcal infection:

no longer in its infancy. Mol Microbiol. 2004;54(1):23-31.

3. Abbott NJ, Patabendige AA, Dolman DE, Yusof SR, Begley DJ. Structure and function

of the blood-brain barrier. Neurobiol Dis. 2010;37(1):13-25.

4. Engelhardt B, Sorokin L. The blood-brain and the blood-cerebrospinal fluid barriers:

function and dysfunction. Semin Immunopathol. 2009;31(4):497-511.

5. Redzic Z. Molecular biology of the blood-brain and the blood-cerebrospinal fluid

barriers: similarities and differences. Fluids Barriers CNS. 2011;8(1):3.

6. Kim YS, Sheldon RA, Elliott BR, Liu Q, Ferriero DM, Tauber MG. Brain injury in

experimental neonatal meningitis due to group B streptococci. J Neuropathol Exp Neurol.

1995;54(4):531-9.

7. van Sorge NM, Doran KS. Defense at the border: the blood-brain barrier versus bacterial

foreigners. Future Microbiol. 2012;7(3):383-94.

8. Kim KS. Microbial translocation of the blood-brain barrier. Int J Parasitol.

2006;36(5):607-14.

9. Kim KS. Pathogenesis of bacterial meningitis: from bacteraemia to neuronal injury. Nat

Rev Neurosci. 2003;4(5):376-85.

10.   Doran KS, Engelson EJ, Khosravi A, Maisey HC, Fedtke I, Equils O, et al. Blood-brain

barrier invasion by group B Streptococcus depends upon proper cell-surface anchoring of

lipoteichoic acid. J Clin Invest. 2005;115(9):2499-507.

11.   Maisey HC, Quach D, Hensler ME, Liu GY, Gallo RL, Nizet V, et al. A group B

streptococcal pilus protein promotes phagocyte resistance and systemic virulence. FASEB J.

2008;22(6):1715-24.

12.   Seo HS, Mu R, Kim BJ, Doran KS, Sullam PM. Binding of glycoprotein Srr1 of

Streptococcus agalactiae to fibrinogen promotes attachment to brain endothelium and the

development of meningitis. PLoS Pathog. 2012;8(10):e1002947.

13.   Mu R, Kim BJ, Paco C, Del Rosario Y, Courtney HS, Doran KS. Identification of a

group B streptococcal fibronectin binding protein, SfbA, that contributes to invasion of brain

endothelium and development of meningitis. Infect Immun. 2014;82(6):2276-86.

14.   Brady LJ, Maddocks SE, Larson MR, Forsgren N, Persson K, Deivanayagam CC, et al.

The changing faces of Streptococcus antigen I/II polypeptide family adhesins. Mol Microbiol.

2010;77(2):276-86.

15.   Rego S, Heal TJ, Pidwill GR, Till M, Robson A, Lamont RJ, et al. Structural and

Functional Analysis of Cell Wall-anchored Polypeptide Adhesin BspA in Streptococcus

agalactiae. J Biol Chem. 2016;291(31):15985-6000.

16.   Sitkiewicz I, Green NM, Guo N, Mereghetti L, Musser JM. Lateral gene transfer of

streptococcal ICE element RD2 (region of difference 2) encoding secreted proteins. BMC

Microbiol. 2011;11:65.

17.   Zhang S, Green NM, Sitkiewicz I, Lefebvre RB, Musser JM. Identification and

characterization of an antigen I/II family protein produced by group A Streptococcus. Infect

Immun. 2006;74(7):4200-13.

18.   Franklin L, Nobbs AH, Bricio-Moreno L, Wright CJ, Maddocks SE, Sahota JS, et al. The

AgI/II family adhesin AspA is required for respiratory infection by Streptococcus pyogenes.

PLoS One. 2013;8(4):e62433.

19.   Pidwill GR, Rego S, Jenkinson HF, Lamont RJ, Nobbs AH. Coassociation between

Group B Streptococcus and Candida albicans Promotes Interactions with Vaginal Epithelium.

Infect Immun. 2018;86(4).

20.   Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. The Phyre2 web portal for

protein modeling, prediction and analysis. Nat Protoc. 2015;10(6):845-58.

21.   Kollman JM, Pandi L, Sawaya MR, Riley M, Doolittle RF. Crystal structure of human

fibrinogen. Biochemistry. 2009;48(18):3877-86.

22.   Davies HD, Jones N, Whittam TS, Elsayed S, Bisharat N, Baker CJ. Multilocus sequence

typing of serotype III group B streptococcus and correlation with pathogenic potential. J Infect

Dis. 2004;189(6):1097-102.

23.   Takahashi S, Adderson EE, Nagano Y, Nagano N, Briesacher MR, Bohnsack JF.

Identification of a highly encapsulated, genetically related group of invasive type III group B

streptococci. J Infect Dis. 1998;177(4):1116-9.

24.   Chi F, Wang L, Zheng X, Wu CH, Jong A, Sheard MA, et al. Meningitic Escherichia coli

K1 penetration and neutrophil transmigration across the blood-brain barrier are modulated by

alpha7 nicotinic receptor. PLoS One. 2011;6(9):e25016.

25.   Doran KS, Chang JC, Benoit VM, Eckmann L, Nizet V. Group B streptococcal beta-

hemolysin/cytolysin promotes invasion of human lung epithelial cells and the release of

interleukin-8. J Infect Dis. 2002;185(2):196-203.

26.   Banerjee A, Kim BJ, Carmona EM, Cutting AS, Gurney MA, Carlos C, et al. Bacterial

Pili exploit integrin machinery to promote immune activation and efficient blood-brain barrier

penetration. Nat Commun. 2011;2:462.

27.   Doran KS, Fulde M, Gratz N, Kim BJ, Nau R, Prasadarao N, et al. Host-pathogen

interactions in bacterial meningitis. Acta Neuropathol. 2016;131(2):185-209.

28.   Oeckinghaus A, Ghosh S. The NF-kappaB family of transcription factors and its

regulation. Cold Spring Harb Perspect Biol. 2009;1(4):a000034.

29.   Keestra AM, van Putten JP. Unique properties of the chicken TLR4/MD-2 complex:

selective lipopolysaccharide activation of the MyD88-dependent pathway. J Immunol.

2008;181(6):4354-62.

30.   Nizet V, Kim KS, Stins M, Jonas M, Chi EY, Nguyen D, et al. Invasion of brain

microvascular endothelial cells by group B streptococci. Infect Immun. 1997;65(12):5074-81.

31.   Karimova G, Gauliard E, Davi M, Ouellette SP, Ladant D. Protein-Protein Interaction:

Bacterial Two-Hybrid. Methods Mol Biol. 2017;1615:159-76.

32.   Dove SL. Studying protein-protein interactions using a bacterial two-hybrid system.

Methods Mol Biol. 2003;205:251-65.

33.   Holmes JA, Follett SE, Wang H, Meadows CP, Varga K, Bowman GR. Caulobacter

PopZ forms an intrinsically disordered hub in organizing bacterial cell poles. Proc Natl Acad Sci

U S A. 2016;113(44):12490-5.

34.   Tomiyama L, Kamino H, Fukamachi H, Urano T. Precise epitope determination of the

anti-vimentin monoclonal antibody V9. Mol Med Rep. 2017;16(4):3917-21.

35.   Mak TN, Bruggemann H. Vimentin in Bacterial Infections. Cells. 2016;5(2).

36.   Tettelin H, Masignani V, Cieslewicz MJ, Donati C, Medini D, Ward NL, et al. Genome

analysis of multiple pathogenic isolates of Streptococcus agalactiae: implications for the

microbial "pan-genome". Proc Natl Acad Sci U S A. 2005;102(39):13950-5.

37.   Leito JT, Ligtenberg AJ, Nazmi K, de Blieck-Hogervorst JM, Veerman EC, Nieuw

Amerongen AV. A common binding motif for various bacteria of the bacteria-binding peptide

SRCRP2 of DMBT1/gp-340/salivary agglutinin. Biol Chem. 2008;389(9):1193-200.

38.   Mollenhauer J, Wiemann S, Scheurlen W, Korn B, Hayashi Y, Wilgenbus KK, et al.

DMBT1, a new member of the SRCR superfamily, on chromosome 10q25.3-26.1 is deleted in

malignant brain tumours. Nat Genet. 1997;17(1):32-9.

39.   Holmskov U, Mollenhauer J, Madsen J, Vitved L, Gronlund J, Tornoe I, et al. Cloning of

gp-340, a putative opsonin receptor for lung surfactant protein D. Proc Natl Acad Sci U S A.

1999;96(19):10794-9.

40.   Madsen J, Mollenhauer J, Holmskov U. Review: Gp-340/DMBT1 in mucosal innate

immunity. Innate Immun. 2010;16(3):160-7.

41.   Stoddard E, Cannon G, Ni H, Kariko K, Capodici J, Malamud D, et al. gp340 expressed

on human genital epithelia binds HIV-1 envelope protein and facilitates viral transmission. J

Immunol. 2007;179(5):3126-32.

42.   Larson MR, Rajashankar KR, Patel MH, Robinette RA, Crowley PJ, Michalek S, et al.

Elongated fibrillar structure of a streptococcal adhesin assembled by the high-affinity association

of alpha- and PPII-helices. Proc Natl Acad Sci U S A. 2010;107(13):5983-8.

43.   Maddocks SE, Wright CJ, Nobbs AH, Brittan JL, Franklin L, Stromberg N, et al.

Streptococcus pyogenes antigen I/II-family polypeptide AspA shows differential ligand-binding

properties and mediates biofilm formation. Mol Microbiol. 2011;81(4):1034-49.

44.   Purushotham S, Deivanayagam C. The calcium-induced conformation and glycosylation

of scavenger-rich cysteine repeat (SRCR) domains of glycoprotein 340 influence the high

affinity interaction with antigen I/II homologs. J Biol Chem. 2014;289(32):21877-87.

45.   Petersen FC, Assev S, van der Mei HC, Busscher HJ, Scheie AA. Functional variation of

the antigen I/II surface protein in Streptococcus mutans and Streptococcus intermedius. Infect

Immun. 2002;70(1):249-56.

46.   Heddle C, Nobbs AH, Jakubovics NS, Gal M, Mansell JP, Dymock D, et al. Host

collagen signal induces antigen I/II adhesin and invasin gene expression in oral Streptococcus

gordonii. Mol Microbiol. 2003;50(2):597-607.

47.   Nobbs AH, Shearer BH, Drobni M, Jepson MA, Jenkinson HF. Adherence and

internalization of Streptococcus gordonii by epithelial cells involves beta1 integrin recognition

by SspA and SspB (antigen I/II family) polypeptides. Cell Microbiol. 2007;9(1):65-83.

48.   Ivaska J, Pallari HM, Nevo J, Eriksson JE. Novel functions of vimentin in cell adhesion,

migration, and signaling. Exp Cell Res. 2007;313(10):2050-62.

49.   Moisan E, Girard D. Cell surface expression of intermediate filament proteins vimentin

and lamin B1 in human neutrophil spontaneous apoptosis. J Leukoc Biol. 2006;79(3):489-98.

50.   Boilard E, Bourgoin SG, Bernatchez C, Surette ME. Identification of an autoantigen on

the surface of apoptotic human T cells as a new protein interacting with inflammatory group IIA

phospholipase A2. Blood. 2003;102(8):2901-9.

51.   Mor-Vaknin N, Punturieri A, Sitwala K, Markovitz DM. Vimentin is secreted by

activated macrophages. Nat Cell Biol. 2003;5(1):59-63.

52.   Podor TJ, Singh D, Chindemi P, Foulon DM, McKelvie R, Weitz JI, et al. Vimentin

exposed on activated platelets and platelet microparticles localizes vitronectin and plasminogen

activator inhibitor complexes on their surface. J Biol Chem. 2002;277(9):7529-39.

53.   Bryant AE, Bayer CR, Huntington JD, Stevens DL. Group A streptococcal myonecrosis:

increased vimentin expression after skeletal-muscle injury mediates the binding of Streptococcus

pyogenes. J Infect Dis. 2006;193(12):1685-92.

54.   Koudelka KJ, Destito G, Plummer EM, Trauger SA, Siuzdak G, Manchester M.

Endothelial targeting of cowpea mosaic virus (CPMV) via surface vimentin. PLoS Pathog.

2009;5(5):e1000417.

55.   Xu B, deWaal RM, Mor-Vaknin N, Hibbard C, Markovitz DM, Kahn ML. The

endothelial cell-specific antibody PAL-E identifies a secreted form of vimentin in the blood

vasculature. Mol Cell Biol. 2004;24(20):9198-206.

56.   Zou Y, He L, Huang SH. Identification of a surface protein on human brain

microvascular endothelial cells as vimentin interacting with Escherichia coli invasion protein

IbeA. Biochem Biophys Res Commun. 2006;351(3):625-30.

57.   Mendez MG, Kojima S, Goldman RD. Vimentin induces changes in cell shape, motility,

and adhesion during the epithelial to mesenchymal transition. FASEB journal : official

publication of the Federation of American Societies for Experimental Biology. 2010;24(6):1838-

51.

58.   Wang RC, Wei Y, An Z, Zou Z, Xiao G, Bhagat G, et al. Akt-mediated regulation of

autophagy and tumorigenesis through Beclin 1 phosphorylation. Science (New York, NY).

2012;338(6109):956-9.

59.   Yang J, Zou L, Yang Y, Yuan J, Hu Z, Liu H, et al. Superficial vimentin mediates

DENV-2 infection of vascular endothelial cells. Sci Rep. 2016;6:38372.

60.   Yu YT, Chien SC, Chen IY, Lai CT, Tsay YG, Chang SC, et al. Surface vimentin is

critical for the cell entry of SARS-CoV. J Biomed Sci. 2016;23:14.

61.   Du N, Cong H, Tian H, Zhang H, Zhang W, Song L, et al. Cell surface vimentin is an

attachment receptor for enterovirus 71. J Virol. 2014;88(10):5816-33.

62.   Murli S, Watson RO, Galan JE. Role of tyrosine kinases and the tyrosine phosphatase

SptP in the interaction of *Salmonella* with host cells. Cellular microbiology. 2001;3(12):795-810.

63.   Ghosh P, Halvorsen EM, Ammendolia DA, Mor-Vaknin N, O'Riordan MXD, Brumell

JH, et al. Invasion of the Brain by Listeria monocytogenes Is Mediated by InlF and Host Cell

Vimentin. MBio. 2018;9(1).

64.   Chi F, Jong TD, Wang L, Ouyang Y, Wu C, Li W, et al. Vimentin-mediated signalling is

required for IbeA+ *E. coli* K1 invasion of human brain microvascular endothelial cells. The

Biochemical journal. 2010;427(1):79-90.

65.   Russell MW, Bergmeier LA, Zanders ED, Lehner T. Protein antigens of Streptococcus

mutans: purification and properties of a double antigen and its protease-resistant component.

Infect Immun. 1980;28(2):486-93.

66.   Lehner T, Russell MW, Caldwell J, Smith R. Immunization with purified protein antigens

from Streptococcus mutans against dental caries in rhesus monkeys. Infect Immun.

1981;34(2):407-15.

67.   Doran KS, Liu GY, Nizet V. Group B streptococcal beta-hemolysin/cytolysin activates

neutrophil signaling pathways in brain endothelium and contributes to development of

meningitis. J Clin Invest. 2003;112(5):736-44.

68.   Stevens C, Henderson P, Nimmo ER, Soares DC, Dogan B, Simpson KW, et al. The

intermediate filament protein, vimentin, is a regulator of NOD2 activity. Gut. 2013;62(5):695-

707.

69.   dos Santos G, Rogel MR, Baker MA, Troken JR, Urich D, Morales-Nebreda L, et al.

Vimentin regulates activation of the NLRP3 inflammasome. Nature communications.

2015;6:6574.

70.   Wilson CB, Weaver WM. Comparative susceptibility of group B streptococci and

Staphylococcus aureus to killing by oxygen metabolites. J Infect Dis. 1985;152(2):323-9.

71.   Da Costa AF, Pereira CS, Santos Gda S, Carvalho TM, Hirata R, Jr., De Mattos-Guaraldi

AL, et al. Group B Streptococcus serotypes III and V induce apoptosis and necrosis of human

epithelial A549 cells. Int J Mol Med. 2011;27(5):739-44.

72.   Back CR, Sztukowska MN, Till M, Lamont RJ, Jenkinson HF, Nobbs AH, et al. The

Streptococcus gordonii Adhesin CshA Protein Binds Host Fibronectin via a Catch-Clamp

Mechanism. J Biol Chem. 2017;292(5):1538-49.

73.   Pritzlaff CA, Chang JC, Kuo SP, Tamura GS, Rubens CE, Nizet V. Genetic basis for the

beta-haemolytic/cytolytic activity of group B Streptococcus. Mol Microbiol. 2001;39(2):236-47.

74.   Woodcock DM, Crowther PJ, Doherty J, Jefferson S, DeCruz E, Noyer-Weidner M, et al.

Quantitative evaluation of Escherichia coli host strains for tolerance to cytosine methylation in

plasmid and phage recombinants. Nucleic Acids Res. 1989;17(9):3469-78.

75.   Park IH, Geno KA, Yu J, Oliver MB, Kim KH, Nahm MH. Genetic, biochemical, and

serological characterization of a new pneumococcal serotype, 6H, and generation of a

pneumococcal strain producing three different capsular repeat units. Clin Vaccine Immunol.

2015;22(3):313-8.

76.   Weksler BB, Subileau EA, Perriere N, Charneau P, Holloway K, Leveque M, et al.

Blood-brain barrier-specific properties of a human adult brain endothelial cell line. FASEB J.

2005;19(13):1872-4.

77.   Weksler B, Romero IA, Couraud PO. The hCMEC/D3 cell line as a model of the human

blood brain barrier. Fluids Barriers CNS. 2013;10(1):16.

78.   Llombart V, Garcia-Berrocoso T, Bech-Serra JJ, Simats A, Bustamante A, Giralt D, et al.

Characterization of secretomes from a human blood brain barrier endothelial cells in-vitro model

after ischemia by stable isotope labeling with aminoacids in cell culture (SILAC). J Proteomics.

2016;133:100-12.

79.   Ohtsuki S, Ikeda C, Uchida Y, Sakamoto Y, Miller F, Glacial F, et al. Quantitative

targeted absolute proteomic analysis of transporters, receptors and junction proteins for

validation of human cerebral microvascular endothelial cell line hCMEC/D3 as a human blood-

brain barrier model. Mol Pharm. 2013;10(1):289-96.

80.   van Sorge NM, Ebrahimi CM, McGillivray SM, Quach D, Sabet M, Guiney DG, et al.

Anthrax toxins inhibit neutrophil signaling pathways in brain endothelium and contribute to the

pathogenesis of meningitis. PLoS One. 2008;3(8):e2964.

81.   Wu Y, Li Q, Chen XZ. Detecting protein-protein interactions by Far western blotting.

Nature protocols. 2007;2(12):3278-84.

82.   Griffith KL, Wolf RE, Jr. Measuring beta-galactosidase activity in bacteria: cell growth,

permeabilization, and enzyme assays in 96-well arrays. Biochem Biophys Res Commun.

2002;290(1):397-402.

A

PI 3   Coomassie stain   PI 10

KDa

Far western blot with biotinylated BspC

Merge

B

Description	Accession	MW (kDa)	Calc PI	# AAs	# peptides	Coverage (%)
Vimentin	gi62414289	53.6	5.12	466	158	61.37

C

```
MSTRSVSSSS  YRRMFGGPGT  ASRPSSSRSY  VTTSTRTYSL  GSALRPSTSR  SLYASSPGGV
YATRSSAVRL  RSSVPGVRLL  QDSVDFSLAD  AINTEFKNTR  TNEKVELQEL  NDRFANYIDK
VRFLEQQNKI  LLAELEQLKG  QGKSRLGDLY  EEEMRELRRQ  VDQLTNDKAR  VEVERDNLAE
DIMRLREKLQ  EEMLQREEAE  NTLQSFRQDV  DNASLARLDL  ERKVESLQEE  IAFLKKLHEE
EIQELQAQIQ  EQHVQIDVDV  SKPDLTAALR  DVRQQYESVA  AKNLQEAEEW  YKSKFADLSE
AANRNNDALR  QAKQESTEYR  RQVQSLTCEV  DALKGTNESL  ERQMREMEEN  FAVEAANYQD
TIGRLQDEIQ  NMKEEMARHL  REYQDLLNVK  MALDIEIATY  RKLLEGEESR  ISLPLPNFSS
LNLRETNLDS  LPLVDTHSKR  TLLIKTVETR  DGQVINETSQ  HHDDLE
```

D

Empty vector

E

BspC-Vimentin

F

Positive control

G

β-galactosidase activty

	Phase contrast	Vimentin		Phase contrast	GBS	Vimentin	Overlay

No infection

GBS infection

WT GBS

ΔbspC

anti-Group B Strep

GBS

anti-vimentin

vimentin

merge

DAPI
GBS
vimentin