Microtubules form by progressively faster tubulin accretion, not by nucleation-elongation

Luke M. Rice[1], Michelle Moritz[2], David A. Agard[2]

[1] Departments of Biophysics and Biochemistry, UT Southwestern Medical Center, 5323 Harry Hines Blvd., Dallas TX 75390

[2] Department of Biochemistry and Biophysics, UC San Francisco, 600 16th St., San Francisco CA 94143

Correspondence: Luke.Rice@UTSouthwestern.edu, Agard@msg.ucsf.edu

Abstract

Microtubules are dynamic polymers that play fundamental roles in all eukaryotes. Despite their importance, how new microtubules form is poorly understood. Textbooks have focused on variations of a nucleation-elongation mechanism in which monomers rapidly equilibrate with an unstable oligomer (nucleus) that limits the rate of polymer formation; once formed, the polymer then elongates efficiently from this nucleus by monomer addition. Such models faithfully describe actin assembly, but they fail to account for how more complex polymers like hollow microtubules assemble. Here we articulate a new model for microtubule formation that has three key features: i) microtubules initiate via rectangular, sheet-like structures which grow faster the larger they become; ii) the dominant pathway proceeds via accretion, stepwise addition of longitudinal or lateral layers; iii) a 'straightening penalty' to account for the energetic cost of tubulin's curved-to-straight conformational transition. This model can quantitatively fit experimental assembly data, providing new insights into biochemical determinants and assembly pathways for microtubule nucleation.

Introduction

The microtubule cytoskeleton plays essential roles organizing the interior of eukaryotic cells: microtubules can act as "tracks" for motor-based transport, they form the basis of the mitotic spindle that mediates proper segregation of chromosomes during cell division, they help define cell polarization, and more (Barlan and Gelfand, 2017; Desai and Mitchison, 1997; Prosser and Pelletier, 2017). To fulfill these many roles, microtubules must assemble into a multitude of highly dynamic and spatially diverse networks. The dynamic instability of microtubules (Mitchison and Kirschner, 1984), which describes the switching of individual polymers between phases of growing and shrinking, allows wholesale reorganization of the microtubule network to occur rapidly and also provides a mechanism to efficiently search space (Holy and Leibler, 1994). Indeed, numerous regulatory factors allow cells to control how fast microtubules grow and how frequently they switch from growing to shrinking (catastrophe) or from shrinking back to growing (rescue) (Akhmanova and Steinmetz, 2015), and also to control the creation of new microtubules (Kollman et al., 2011; Roostalu and Surrey, 2017; Wieczorek et al., 2015), a process called nucleation.

Microtubules are hollow, cylindrical polymers formed from $\alpha\beta$-tubulin heterodimers that interact with each other in two ways: stronger head-to-tail (longitudinal) interactions between $\alpha\beta$-tubulins make up the straight protofilaments, and weaker side-to-side (lateral) interactions hold protofilaments together. The mechanisms underlying the dynamic instability of existing microtubules are increasingly well understood (Alushin et al., 2014; Driver et al., 2017; Duellberg et al., 2016; Gardner et al., 2011; Geyer et al., 2015; Grishchuk et al., 2005; Manka and Moores, 2018; Margolin et al., 2012; Piedra et al., 2016; VanBuren et al., 2005; VanBuren et al., 2002; Zhang et al., 2015), but models for how *existing* microtubules grow and shrink cannot specify how *new* microtubules form de novo. Thus, our understanding of the mechanisms that govern spontaneous microtubule formation remains relatively primitive (reviewed in

(Roostalu and Surrey, 2017)). Multiple factors contribute to this difficulty: first,

nucleation occurs very rarely and cannot truly be observed microscopically, so

measuring it directly is much harder than measuring the growing and shrinking of

existing microtubules. Second, the open, tube-like structure of the microtubule

poses unique complications. In most organisms, microtubules contain 13

protofilaments, although there are clear examples of specialized microtubules

containing 11 or 15 protofilaments (Burton et al., 1975; Chaaban et al., 2018;

Chalfie and Thomson, 1982; Davis and Gull, 1983; Kwiatkowska et al., 2006;

Saito and Hama, 1982; Tucker et al., 1992). Regardless of protofilament number,

the hollow nature of the microtubule means that it takes many more $\alpha\beta$-tubulin

subunits to close a tube than it does to make a minimal helical repeat for a

simpler, helical polymer like actin. Perhaps not surprisingly given this complexity,

there is no consensus about the sequence of intermediates that must be formed

(pathway) during the transition from individual subunits to a closed tube. To

further complicate matters, the structure of the free heterodimer also changes

during assembly, from a bent confirmation in solution to a straight conformation

within the final microtubule ((Ayaz et al., 2012; Nawrotek et al., 2011; Rice et al.,

2008); reviewed in (Brouhard and Rice, 2018)). Where in the assembly process

the straightening takes place, whether it is continuous or step-wise, how much

energy is required, and how this obligatory conformational change affects

nucleation all remain unknown. In addition to these factors, the GTPase activity

required for microtubule catastrophe can also antagonize nucleation by

destabilizing intermediate species.

An early study of the kinetics of spontaneous microtubule nucleation, which used

solution conditions that suppress catastrophe to simplify the behavior, posited

that nucleation proceeded sequentially through two discrete intermediates (Voter

and Erickson, 1984). The data were re-analyzed in later studies (Flyvbjerg et al.,

1996; Kuchnir Fygenson et al., 1995) that proposed a more complex pathway

containing four distinct nucleation intermediates (6-mer, 9-mer, 12-mer, and 15-

mer of $\alpha\beta$-tubulin). However, in this later analysis it was not possible to ascribe a

biochemical interpretation to the species or rate constants (Flyvbjerg et al.,

1996). Thus, the physical-chemical details of the underlying assembly process

continue to remain obscure.

In this work, we sought to obtain more explicit insight into the molecular

mechanisms that govern spontaneous microtubule assembly. Our primary goal

was to explore the simplest biochemical models that could faithfully explain the

experimental data. We began from the fundamental modes of $\alpha\beta$-tubulin

interactions within a microtubule: a single subunit (heterodimer) can make a

longitudinal interaction, a lateral interaction or make both simultaneously (corner

interaction). Starting with explicit simulations of the biochemical reactions that

can occur for all possible $\alpha\beta$-tubulin oligomers up to 12-mers, we observed that a

small subset of all possible oligomers became dominant. These dominant

oligomers were "rectangular", meaning that they maximized the number of

longitudinal and lateral contacts such that there were no unfilled corners. The

dominance of a small subset of species allowed us to develop a simpler,

approximate model based almost exclusively on rectangular oligomers. This

simplification allowed the model to access arbitrarily large oligomers in a way that

retained a connection to the fundamental biochemical interaction modes of the

individual subunits.

This approximate but biochemically faithful model provides a new way to view

microtubule nucleation. Rather than proceeding via a distinct rare, rate limiting

"nucleus", the model indicates that microtubule assembly involves the continuous

accretion of $\alpha\beta$-tubulins into a growing 2D lattice. Like crystallization processes,

the larger the assembly, the faster it grows. Our model reveals assembly

pathways for forming a new microtubule and provides new insights into how

exogeneous nucleators and other regulatory factors can accelerate microtubule

formation.

Results

Rectangular species dominate the assembly pathway

To obtain biochemical insight into possible pathways for spontaneous

microtubule assembly, we constructed an explicit model that includes all non-

redundant tubulin oligomers from dimers through dodecamers (Fig. 1A shows the

intermediates through hexamers). Here non-redundant means that we only

consider a single exemplar N-mer for each unique combination of longitudinal

and lateral contacts; statistical weighting of rate constants ensures that this

exemplar oligomer faithfully represents the ensemble of species it represents

(see example in Sup. Fig. 1). We defined the biochemical pathways that connect

different oligomers (arrows in Fig. 1A) by assuming that oligomers grow and

shrink via addition and loss of individual $\alpha\beta$-tubulin subunits. It was not practical

to explore larger oligomers in this formulation because of an explosion in the

number of species and reactions (reaching dodecamers requires ~200 distinct

species and ~600 reactions) (Fig. 1F).

We implemented the differential equations describing this discrete assembly

scheme in Berkeley Madonna (Marcoline et al., 2020) to simulate the assembly

pathway and kinetics. The only free parameters in this model are the on rate

constant k_f and the affinities for the three different $\alpha\beta-$tubulin interaction modes

(longitudinal, lateral, and corner (simultaneous longitudinal+lateral interaction

(Fig 1D). As is common, k_f was assumed (here we chose 10^6 $M^{-1}s^{-1}$ based on

experimental observations and consistent with work from others (Coombes et al.,

2013; Margolin et al., 2012; McIntosh et al., 2018; VanBuren et al., 2002;

Zakharov et al., 2015)). We explored different values for the three parameters,

seeking to identify conditions under which some appreciable concentration of

dodecamers would form. We were especially interested in the corner affinity,

because this is the interaction that would provide the driving force for microtubule

elongation. A relatively tight corner affinity (K_d^{corner} < 0.2 μM; Sup. Fig. 2) was

required for appreciable flux into the largest oligomers. With high-affinity corner

interactions, the simulations predominantly populated a small number of

'rectangular' oligomers, defined as oligomers wherein (i) every subunit makes at

least one longitudinal and one lateral contact to another, and (ii) there are no

'open' corner sites (Fig. 1B). More explicitly, what happens in the simulations is

that formation of an initial 2x2 oligomer forms relatively quickly. The next larger

species containing an additional 'singleton' tubulin are poorly populated because

of the weak longitudinal and lateral affinities, but those species also present an

open, high-affinity corner site. Filling this open site stabilizes the first subunit and

in the case of a 2x2 start, fills the next longitudinal or lateral layer resulting in a

3x2 or 2x3 assembly, respectively. On larger assemblies, additional corners are

vacant after the first, and these fill rapidly, essentially "zipping up" the layer. The

net consequence is that once initiated, layer completion occurs efficiently,

explaining why rectangular species dominate.

Only considering rectangular oligomers greatly reduces the number of species

and reactions required to reach much larger oligomers in simulations (e.g. Fig.

1C,F). However, in such an approximation the rates of adding or losing a layer

become context dependent, and depend on the height and width of the oligomer.

Using the steady-state approximation, we were able to derive a general recursive

expression for the effective 'forward' rate of adding of a longitudinal or lateral

layer (see Methods). An example of adding a longitudinal layer to a 2x2 oligomer

is illustrated in Fig. 1D. Considering the layer addition as a sequence of individual

steps also allowed us to derive an effective forward rate as well as a 'layer

equilibrium ratio', which describes the expected relative populations of oligomers

that differ by a single layer (Methods). Importantly, in these expressions both the

forward rate and equilibrium ratio for layer addition are defined in terms of the

biochemical properties of individual subunits (longitudinal, lateral and corner

affinities/rates). Consequently, and in contrast to what occurs with

phenomenological models, the reduction in the number of species and reactions

does not come at the expense of losing a connection to the fundamental

biochemical behavior. For the 12-mer simulated explicitly above, the layer model

provides a >10-fold reduction in species and >35-fold reduction in number of

reactions (Fig. 1F) (these reduction factors increase dramatically for larger

oligomer sizes).

We implemented the reduced, layer-based model into Berkeley Madonna to test

if it could recapitulate the results from our explicit simulations when using the

same input parameters for subunit affinities. Despite the dramatic reduction in

complexity, and with the exception of very low-abundance intermediate species,

the simplified layer-based simulations quantitatively reproduced the oligomer

concentrations and the kinetics of oligomer formation observed with the explicit

model (compare Fig. 1E to 1B). These initial benchmarking layer simulations only

considered oligomers up to dodecamers so that we could make a direct

comparison with the explicit model. In the next section, we expand the model to

include the much larger oligomers required to depict microtubule assembly.

A simplified accretion pathway for spontaneous microtubule assembly

We wrote a custom program to expand the layer model to arbitrary sized

rectangular oligomers. Whenever an oligomer gained a 13th protofilament, we

converted it into a microtubule. No additional penalty for seam formation was

included. We assumed that once formed, microtubules elongate by capturing αβ-

tubulin subunits one at a time in the open, pseudo-helical 'corner' sites on either

end. In order to minimize the number of adjustable parameters, the same rate

constants were used for events at either the plus- or minus-ends. These

additional simplifying assumptions allow the model to simulate spontaneous

microtubule assembly and the ensuing depletion of free αβ-tubulin subunitsin a

relatively natural way, without having to explicitly specify a particular nucleus or

assume a preferred assembly pathway.

We began testing our model by fitting it to experimental assembly curves that we

measured by light scattering (Fig. 2A). The model was able to fit individual curves

quite well (Fig. 2B), with minor discrepancies evident at low concentration. The

optimized affinities differed somewhat from fit to fit, but shared common features:

tight corner affinities (K_d^{corn} < 100 nM), weaker longitudinal affinities (K_d^{long} ~5-15

mM), and lateral affinities around 10-fold weaker than longitudinal affinities (Fig.

2B). The longitudinal affinity we obtained is in the range of that obtained from

biochemical models for microtubule growing and shrinking (e.g.(Gardner et al.,

2011; Kim and Rice, 2019; Piedra et al., 2016; VanBuren et al., 2002)). The

corner and lateral affinities are both about 10-fold stronger than in those models,

a difference that might reflect the different buffer conditions (high glycerol

concentration) in 'nucleation' assays relative to 'dynamics' assays. The corner

affinity is much stronger than the product of longitudinal and lateral affinities,

which is expected because the individual longitudinal and lateral affinities

represent a balance between (among other things) favorable protein:protein

contacts and the unfavorable entropic cost of losing rotational and translational

degrees of freedom -- when making a corner interaction the longitudinal and

lateral interfaces both contribute favorably while the entropic cost remains

approximately the same (see also (Erickson and Pantaloni, 1981)). Fits to an

independent set of assembly curves (Sup. Fig. 3A) yielded similar affinities (Sup.

Fig. 3B). We found it encouraging that the model could fit individual assembly

curves, but to be convincing a single parameter set should be able to fit all the

assembly curves simultaneously. Thus, we turned to global fitting where we

attempted to fit the set of assembly curves with one common parameter set.

In a global fit to the set of assembly curves (Fig. 2C), a limitation of the model

became apparent: even though each individual concentration could be well fit

(Fig. 2B), the model could not capture the correct behavior over the broad

concentration range measured (Fig. 2C). The model appeared to fit the

alternative dataset somewhat better (Sup. Fig. 3C), but it was still apparent that

the model was underestimating the concentration-dependence of the assembly

curves; an outlier curve also limited the concentration range available for fitting.

Because all layer transition rates scale with the chosen on-rate constant (see

derivations in Methods), changing the assumed on-rate does not improve the

global fits. The observed discrepancy between model and experiment suggests

that the model fails to capture some important aspect of assembly. In re-

examining our assumptions, we realized that it was too restrictive to consider

only monomer addition and loss as the 'elemental' reactions within the layer

assembly pathway. Indeed, a simple calculation demonstrates that any model

having these relative lateral and longitudinal affinities must also include a 'dimer

pathway' (Fig. 2D). Compared to the 'two step' monomer-based pathway in

which a first monomer associates weakly and only becomes 'locked in' by a

subsequent corner association (Fig. 2D, middle path), when the longitudinal

affinity is stronger than the lateral one as it is for microtubules it is more efficient

to initiate a lateral layer by adding a pre-formed longitudinal dimer (Fig. 2D, lower

path). This is true even though longitudinal dimers are present at very low

concentrations (roughly 1 part in a 1000 or ~10 nM) for the parameters we used.

Our 'monomer only' model global fit required a longitudinal affinity ~10-fold

stronger than the lateral one, but under those conditions additions of longitudinal

dimers should actually account for ~90% of the lateral layer initiations (Fig. 2E).

Thus, despite appearing reasonable, the fits do not make biochemical sense

because the model fails to account for a dimer pathway that has the potential to

dominate the flux through oligomers. It has been suggested that oligomer

associations may be important during spontaneous microtubule assembly

(Erickson and Pantaloni, 1981; Mozziconacci et al., 2008; Voter and Erickson,

1984), but to our knowledge these interactions have not previously been

implemented into kinetic models.

Incorporating a dimer pathway fails to recapitulate experimental data

without invoking a switch in lateral affinity

The cartoons and calculations presented in Fig. 2D,E illustrate how the rate of

initiating a new lateral layer will depend on the concentration of longitudinal

dimers of $\alpha\beta$-tubulin. By contrast, once the slow step of *initiating* a new layer has

occurred, *zipping up* via addition into favorable 'corner' (longitudinal + lateral)

sites should occur almost exclusively via $\alpha\beta$-tubulin 'monomers' because these

monomers far exceed longitudinal dimers in number/concentration. By a similar

logic, it is also not necessary to include a dimer pathway for initiating a

longitudinal layer: lateral dimers are very rare, and the 'upper' $\alpha\beta$-tubulin of a

longitudinal dimer provides no increased stabilization as a longitudinal layer

builds up.

Incorporating a parallel, dimer-based pathway for initiating lateral layers (Sup.

Fig. 4) requires that at each step we specify (i) the concentration of longitudinal

dimers, (ii) an association rate constant for the longitudinal dimers adding to a

complete layer, and (iii) a dissociation rate constant for the lateral 'unbinding' of a

longitudinal dimer. The concentration of longitudinal dimers is easily calculated

given the concentration of monomers and the equilibrium constant for

longitudinal interactions. Because off-rate constants tend to vary over a much

larger range than on-rate constants, for simplicity we used the same, roughly

diffusion limited association rate constant as we do for monomers. We derived a

formula that expresses the equilibrium constant for lateral association of a

longitudinal dimer solely in terms of biochemical properties of the monomers (see

Methods). Thus, we were able to parameterize the 'dimer pathway' purely in

terms of the fundamental monomer properties.

We first fit the dimer model to individual assembly curves, as we did for the

monomer model (Fig. 3A). The longitudinal and corner affinities for these dimer

model fits were comparable to those obtained from fits of the monomer model.

The lateral affinity (~25-30 M in the dimer model) was ~300-fold weaker than the

in the monomer model (~100 mM), reflecting a greatly diminished contribution of

the monomer pathway for lateral initiation. Although it was predicted to be

required biochemically, including this dimer pathway actually degraded the ability

of the model to recapitulate the shape of individual assembly curves compared to

the monomer-only fits: the simulated assembly curves displayed a less

pronounced lag phase and also appeared less cooperative (shallower sigmoid

transition; compare Fig. 3A to Fig. 2B). Global fits of the dimer model to the set of

assembly curves were also markedly poorer than for the monomer model (Fig.

3B). Thus, a more complete biochemical model for the assembly process

paradoxically degraded its predictive power.

Why was the monomer-only model able to fit the observed assembly curves so

much better than a model incorporating a dimer pathway? The presence of a

dimer pathway makes the effective rates of lateral and longitudinal growth much

more comparable to each other (Fig. 2D, compare top and bottom rate

expressions), whereas in the monomer-only model there is significantly more

contrast between the rates for longitudinal and lateral growth. We speculated that

the otherwise more accurate monomer+dimer model must still be missing a

critical feature that acts to increase the difference between longitudinal and

lateral growth.

What might be missing? Not having accounted for the effects of conformational

changes in $\alpha\beta$-tubulin that occur during microtubule formation (Ayaz et al., 2012;

Nawrotek et al., 2011; Pecqueur et al., 2012; Wang and Nogales, 2005) is a

strong candidate. Indeed, it is now well established that unpolymerized $\alpha\beta$-tubulin

subunits adopt a 'curved' conformation in solution and that the microtubule lattice

itself acts as an allosteric activator to 'straighten' $\alpha\beta$-tubulin during the assembly

process (Ayaz et al., 2012; Buey et al., 2006; Nawrotek et al., 2011; Rice et al.,

2008). That is, $\alpha\beta$-tubulin becomes progressively straighter as it becomes

increasingly surrounded by lateral neighbors (reviewed in (Brouhard and Rice,

2008)) (Fig. 4A). This straightening-induced conformational strain acts to weaken

lateral interactions in assemblies that require more straightening. To explore

these ideas in the context of our model, we introduced a very simple lateral

penalty to mimic the greater strain intrinsic to straighter conformations. We

assumed that two-wide assemblies could remain curved, whereas wider

assemblies would be forced to straighten, making the lateral interactions after the

first pairing in a given 'row' X-fold weaker.

We fit the monomer+dimer+penalty model to the data as before, testing a range

of penalty magnitudes (from 1 to 16 in Fig. 4B). The global fit to the experimental

curves improved as the magnitude of the lateral penalty increased, with the best

fits reached for penalty values greater than 4. Thus, penalizing lateral growth was

sufficient to 'correct' the deficiencies of the simple dimer pathway, resulting in

even better fits (Fig. 4B,C; penalty value = 8,) than for the monomer-only model

(see Fig. 3B,C). Beyond a value of 4, the improvement in global fit only weakly

depends on the precise value of the penalty (Fig. 4B). Fits to the alternative

dataset yielded similar but not identical affinities (Sup. Fig. 5). Other penalty

factors gave fits of comparable quality as long as the penalty factor was greater

than 4 (Fig. 4B), but yielded different longitudinal and lateral affinities, as

expected. Varying individual parameters shows that the fits are sensitive to small

(1.5-fold) changes in the parameter values (Sup. Fig. 6). Differences in the fitted

parameters for the primary and alternative dataset suggests that within a narrow

range there may be some compensation between corner and longitudinal

affinities (compare Fig. 4C to Sup. Fig. 5), Collectively, our computational

experiments led us to what we believe to be the first simple, biochemically-based

model that can accurately recapitulate the observed concentration-dependent

kinetics of spontaneous microtubule assembly without having to assume a

specific nucleus.

A kinetically frustrated pathway for spontaneous microtubule assembly

The model developed here provides a biochemical pathway that explains in a new way why the kinetics of microtubule initiation are slow relative to polymer elongation. As a first way to illustrate the assembly pathway that emerges from fitting our model to the experimental data, we plotted the total concentration and average size of intermediate oligomers as a function of time (Fig. 5A). The total concentration of oligomers peaks at around 200 s in the example shown, and the subsequent decrease is a consequence of the formation of microtubules and of the depletion of free $\alpha\beta$-tubulin. The average oligomer size increases gradually over several hundred seconds, reflecting the slow process of accretion and the low likelihood that large layers will be lost in 'backward' reactions.

The endpoint of microtubule formation – closure of the cylindrical lattice –requires a minimal width oligomer (13 in our model). Thus, we next examined the summed concentration of tubulin oligomers with a given width (lateral extent) as a function of time (Fig. 5B). This analysis shows that the concentration of oligomers decreases as their width increases – there are fewer 3-wide oligomers than 2-wide, fewer 4-wide then 3-wide, etc. (Fig. 5B). This decreasing concentration of wider oligomers happens because the transition from one width to the next accelerates throughout the assembly process. This acceleration can be understood as a statistical/kinetic consequence of the two-dimensional assembly pathway. No matter their height, all oligomers of a given width grow *taller* with the same characteristic rate because they present the same number of 'landing sites' for starting a new longitudinal layer. However, taller oligomers grow *wider* faster than shorter oligomers, because their increased height provides more 'landing sites' for starting a new lateral layer. Similarly, wider oligomers add longitudinal layers faster than the narrower ones for the same reason.

To provide more granular insight into the nature of the predominant microtubule

assembly intermediates, we examined the distribution of oligomer sizes at

different widths (Fig. 5C). For each oligomer width, and at the time the

concentration of oligomers of that width peaked, we plotted the most populated

height (filled circle) and the relative amount of taller or shorter oligomers (grey

teardrop/oval shapes, which are actually a set of circles with area scaled

proportionally to the concentration). The many parallel pathways in the assembly

process leads to an increasing spread in the range of oligomer heights as they

get wider. For example, the most populated 50% of all 3-wide oligomers have

heights between 7 and 11 (a height range of 5) whereas the most populated 50%

of all 12-wide oligomers span heights between 50 and 64 (a height range of 15,

three times greater than for 3-wides). The most populated immediate microtubule

precursor (12 subunits wide) contains almost 700 tubulins: 58 subunits tall x 12

wide. This is much larger than the number of subunits in the nucleus predicted by

phenomenological models (Flyvbjerg et al., 1996; Kuchnir Fygenson et al., 1995).

Also note that the rate of layer assembly cannot exceed the rate of elongation.

The layer-based assembly model thus provides a new, ensemble-based view of

the assembly pathway wherein many oligomers of varying heights contribute

appreciably to the formation of new microtubules.

Discussion

How existing microtubules grow and shrink is increasingly well-understood, but

the understanding of how new microtubules form spontaneously – a related but

mechanistically distinct process – lags behind (Roostalu and Surrey, 2017).

Many discussions of spontaneous microtubule formation have invoked the

concept of a 'critical nucleus', which is analogous to the transition-state of a

chemical reaction (Kollman et al., 2011; Kuchnir Fygenson et al., 1995; Roostalu

and Surrey, 2017; Voter and Erickson, 1984). This traditional view of the

assembly pathway assumes that (i) the formation and growth of early

intermediates is thermodynamically unfavorable, and (ii) above a critical size,

oligomer growth by monomer addition becomes favorable and this elongation

proceeds at a fixed rate. The polymerization kinetics of relatively simple helical

polymers like actin can be well-described by two-step, nucleation-elongation

models that invoke a small critical nucleus representing a 'mini-filament' (the

smallest oligomeric species that presents helical binding sites) (Sept and

McCammon, 2001; Tobacman and Korn, 1983). However, the spontaneous

assembly kinetics of hollow, cylindrical microtubules do not conform to the

simplest two-step, nucleation-elongation kinetics (Flyvbjerg et al., 1996; Kuchnir

Fygenson et al., 1995; Voter and Erickson, 1984). Phenomenological multi-step

nucleation-elongation mechanisms (Flyvbjerg et al., 1996) can fit assembly

curves but fail to provide biochemical insights into the assembly pathway.

Our model provides an alternative and conceptually distinct framework for

thinking about the pathway for microtubule formation (Fig. 6), and the

biochemical factors that govern the overall rate of assembly. From this comes the

understanding that there is not a singular, transition-state-like species (variously

estimated as 8-27 tubulins) that dictates the assembly kinetics, but instead what

we would traditionally call "nucleation" arises from an ensemble of accretion

events involving many species and hundreds of tubulin heterodimers. In what

follows, we summarize the main features of the predicted assembly pathway and

discuss implications for the multiple ways that a template or other regulatory

factor could accelerate the process.

The layer growth model predicts that a multitude of pathways and intermediates

contribute meaningfully to microtubule assembly. Stronger longitudinal

interactions cause the dominant rectangular intermediates to be taller than they

are wide. Early in the assembly pathway, transitions to larger oligomers become

energetically favorable (equilibrium ratio less than one) but still occur slowly

because the oligomers are small and because initiating the new layer depends

on a weak individual longitudinal or lateral interaction. The rate of transition

between oligomers accelerates as oligomers increase in size, not because of any

intrinsic changes in the underlying affinities, but because larger oligomers

present more sites on which the next layer can initiate. These size-dependent

changes in transition rates are a direct consequence of the two-dimensional

nature of the assembly pathway. Thus, even though nearly all of the layer

additions are energetically 'downhill', the repeated difficulty of initiating each new

layer contributes a series of progressively smaller delays into the assembly

sequence that in combination represent the effective "barrier". Conformational

'straightening' of $\alpha\beta$-tubulins adds an additional kind of frustration that makes

lateral growth of early oligomers even harder. Collectively, the ensemble of

initiation barriers creates a kind of 'frustration' that dictates unique kinetics for

spontaneous microtubule assembly.

This layer-based view of the assembly pathway provides new ways for thinking

about how regulatory factors might accelerate microtubule initiation. Our

modeling indicates that the kinetics of microtubule formation are dictated by the

need for *repeated* 'layer' transitions that are limited by the weaker lateral affinity

and by the straightening penalty (e.g. Fig. 7A). Classical "nucleators" like the γ-

tubulin ring complex (γ-TuRC) (Zheng et al., 1995) contain a lateral array 14 γ-

tubulins (Fig. 7B) that are thought to form a 'template' on which a microtubule

can elongate (Consolati et al., 2020; Kollman et al., 2015; Kollman et al., 2011;

Liu et al., 2020; Moritz et al., 1995; Wieczorek et al., 2020). We used our model

to explore how the potency of templating might depend on the width of the

template. We modified our simulations to include various width templating

oligomers (Fig. 7C), using a constant total amount of templating 'monomers'.

Note that in these simulations there are progressively lower concentrations of the

wider oligomers (e.g. half as many 4-wide oligomers as 2-wide, etc.). The affinity

of $\alpha\beta$-tubulins for γ-TuRC is unknown, so we explored a regime where γ:$\alpha\beta$

longitudinal interactions are identical to $\alpha\beta$:$\alpha\beta$ (Supplemental Fig. 7) and another

where $\gamma{:}\alpha\beta$ longitudinal interactions are weaker than for $\alpha\beta{:}\alpha\beta$ (Fig. 7). With the

stronger $\gamma{:}\alpha\beta$ interaction strength, even 2-wide "templates" potently accelerated

assembly (Supplemental Fig. 7). This is at odds with experimental data showing

that γ-tubulin small complexes (γ-TuSCs), which only contain two γ-tubulins, are

much poorer nucleators than γ-TuRCs (Kollman et al., 2015; Oegema et al.,

1999). By contrast, when $\gamma{:}\alpha\beta$ interactions were assumed to be weaker,

templating oligomers that were 2 subunits wide only modestly accelerated the

predicted assembly kinetics (1.1-fold faster characteristic time for assembly, $T_{0.1}$)

(Fig. 6C,D). A 3-wide oligomer was much more potent (1.7-fold faster) even

though in the simulations the concentration of 3-wide oligomers was less than

that of the 2-wide oligomer. This strong change in potency occurs because the

'hardest' transition in the assembly sequence is from two-wide oligomers to 3-

wide oligomers, providing an explanation for why γ-tubulin small complexes (γ-

TuSCs) only weakly stimulate assembly (Oegema et al., 1999). A recent study

provides experimental support for these prediction by showing that

$\gamma{:}\alpha\beta$ interactions are weaker than $\alpha\beta{:}\alpha\beta$ interactions (Thawani et al., 2020) In our

simulations, progressively wider templating arrays show even greater potency,

emphasizing that the barriers to lateral growth contribute throughout the entire

assembly pathway.

The idea that multiple reactions/transitions contribute to limit the overall rate of

microtubule formation also suggests that strategies besides templating could

enhance the rate of spontaneous assembly. The rates for layer additions depend

on how fast tubulins 'land' on the growing sheet, so enhancing this rate should

promote layer initiation and 'zippering'. For example, XMAP215 family

polymerases use an 'accelerated delivery' mechanism (Ayaz et al., 2014; Geyer

et al., 2018) to make microtubules grow faster, and this same activity likely

underlies the role for XMAP215 family proteins in microtubule nucleation

(Gunzelmann et al., 2018; Popov et al., 2002; Roostalu et al., 2015; Thawani et

al., 2018; Wieczorek et al., 2015). The efficiency of initiating new layers is

primarily limited by the relatively weak longitudinal or lateral affinities, especially

for the singleton tubulin that initiates each longitudinal layer. Thus, a factor that

stabilizes these associations should also increase the rate at which layer

transitions occur. Proteins like TPX2 and doublecortin bridge between

neighboring subunits in the lattice (Fourniol et al., 2010; Zhang et al., 2017); this

'molecular stapling' may help explain their documented roles in microtubule

"nucleation" (Moores et al., 2004; Roostalu et al., 2015). Finally, altering the

allosteric properties of $\alpha\beta$-tubulin itself should also profoundly affect microtubule

initiation. For example, and in keeping with predictions of our model, studies of

an $\alpha\beta$-tubulin 'conformation cycle' mutation (Geyer et al., 2015) that enhances

the strength of lateral interactions (Driver et al., 2017) (perhaps via reducing the

cost of $\alpha\beta$-tubulin straightening), greatly accelerates microtubule formation.

Similarly, small molecule compounds like taxol that stimulate microtubule

initiation and growth may do so by promoting straightening.

Conclusions and perspective

We described a relatively simple accretion model that explains spontaneous

microtubule assembly kinetics in terms of subunit biochemistry and an ensemble

of 'rectangular' intermediates. Whereas in typical 'nucleation-elongation'

paradigms the scarcity of the transition-state-like nucleus is what limits the rate of

polymer formation, our model is more akin to a process like crystal growth.

Indeed, in our accretion model most of the steps toward polymer formation are

energetically favorable but kinetically frustrated by the difficulty of initiating new

layers; this difficulty decreases as layers become larger because more landing

opportunities are presented. The accretion mechanism we describe is a direct

consequence of the two-dimensional nature of the hollow, cylindrical microtubule

lattice.

We did not consider the kinetic consequences of GTP hydrolysis within the

nascent polymer in our model, because it was not possible to account for the

stochastic hydrolysis within the context of our deterministic formalism. In keeping

with that, our assay conditions were chosen to suppress microtubule

catastrophe. We attempted to measure assembly curves in the presence of

GMPCPP, a hydrolysis-resistant GTP analog, but obtained variable results

because it was difficult to reliably eliminate contaminating amounts of

microtubule seeds from the initial reaction mixtures. Ignoring GTPase activity

represents a limitation of our model that we plan to address in future work. In

general, GTPase activity should antagonize assembly by preferentially

destabilizing intermediate oligomers, so the affinities we report may

underestimate the 'true' affinities. We still expect that growth through accretion

will represent the dominant pathway for assembly.

Finally, our model provides a unifying quantitative framework for understanding

the complex effects of regulatory molecules. Indeed, the 'zippering' transitions

between rectangular oligomers that dominate the initiation process also share

many features in common with microtubule elongation. Thus, our model also

explains why factors that regulate microtubule elongation and shrinking also

regulate spontaneous microtubule assembly.

561
562

Acknowledgements
This study was supported by grants from the National Science Foundation to
LMR (MCB-1615938 and MCB-1054947) and from the National Institutes of
Health to DAA (R01 GM031627, R35 GM118099 and P01 GM105537). DAA was
supported in part by the Howard Hughes Medical Institute. We thank T. Davis
and her colleagues for very helpful feedback on the manuscript.

569
570 **Author contributions**
571 MM collected the microtubule assembly data and helped write the manuscript.
572 LMR constructed models, fit the data, and drafted the manuscript. DAA analyzed
573 data, supervised the research, and helped write the manuscript.
574
575

Methods

Measurements of spontaneous assembly

The spontaneous assembly of phosphocellulose-purified porcine-brain tubulin was followed by turbidity at 350 nm basically as described previously (Gaskin et al., 1974; Voter and Erickson, 1984). Care was taken to remove microtubule seeds and inactive protein by cycling it through an additional polymerization/depolymerization step just before use: immediately prior to performing assays, tubulin at –80°C was rapidly thawed at 37°, placed on an ice-water slurry (0°C), and the buffer was adjusted to 80 mM K-PIPES, pH 6.8, 1mM EGTA, 4 mM $MgCl_2$ and 1 mM GTP. After 5 min incubation, 0.5 volume of 37°C glycerol was mixed in and the tubulin was allowed to polymerize at 37°C for 20 min. Microtubules were pelleted through a 37°C, 60% glycerol cushion in 50 mM K-MES, pH 6.6, 5 mM $MgSO_4$, 1 mM EGTA, 1 mM GTP in a TLA110 rotor (Beckman, Palo Alto, CA) at 80,000 rpm, 20 min. The pellets were resuspended at 37°C in assembly buffer (AB: 50 mM K-MES, pH6.6, 3.4 M glycerol, 5 mM $MgSO_4$, 1 mM EGTA, 1 mM GTP) using a warm Potter homogenizer. The resuspended microtubules were then depolymerized on an ice-water slurry for 20 min and centrifuged in a TLA100.3 rotor at 100,000 rpm for 35 min at 2°C to remove any remaining polymerized tubulin. The supernatant from this cold spin was removed and used for the assembly reactions. The tubulin was kept at 0°C as reaction mixtures were prepared. A series of samples was prepared at different tubulin concentrations in AB and kept at 0°C in thin-walled, 0.5 ml plastic tubes (PCR tubes, Stratagene, Cedar Creek, TX). Samples were rapidly heated to 37°C directly in cuvettes pre-warmed to 37°C in a Pelltier cell changer (took 30 sec), in a custom built spectrophotometer. The $A_{350 nm}$ was recorded approximately every 4 sec on 4 samples at a time until plateaus were reached. Recording on the spectrometer was started prior to transfer of tubulin to the cuvettes so that early lag-phase data were not missed. The time was adjusted to ignore the 30 sec heating step. Scaling analysis was performed using a customized program.

Explicit simulations in Berkeley Madonna

We constructed a model in Berkeley Madonna (Marcoline et al., 2020) to simulate αβ-tubulin self-assembly. The first step in this process was to define the set of αβ-tubulin oligomers and how they are related to each other by gain/loss of an αβ-tubulin though a longitudinal, lateral, or longitudinal+lateral ('corner') interaction. Starting from the two-dimensional nature of the microtubule lattice, we built a list of all possible configurations for an αβ-tubulin oligomer of a given size. Fig. 1 illustrates how there are two kinds of possible dimers of αβ-tubulin (longitudinal and lateral), three kinds of trimers of αβ-tubulin, etc. Larger oligomers can have multiple configurations that are energetically equivalent. To minimize the number of species and reactions considered in the model, we used a single exemplar species to represent each set of n-mers with the same number

of longitudinal and lateral contacts. For each exemplar species, population-
weighted average rate constants were used to account for the different properties
of each oligomer. Supplemental Figure 1 provides a worked example of this
approximation for a tetramer of $\alpha\beta$-tubulin. We then used the 'Chemical
Reactions' module in Berkeley Madonna (Marcoline et al., 2020) to build a set of
rate equations corresponding to each possible transition. We did not pursue
oligomers larger than 12-mers because the number of species and reactions
became intractable. This model provides a way to analyze the rate and extent to
which different species become populated for a given choice of the adjustable
parameters.
630
631 ***Derivation of layer transition rates and validation in Berkeley Madonna***
632 The explicit, monomer-based simulations in Berkeley Madonna (Marcoline et al.,
633 2020) indicated that a simpler reaction scheme that only considered transitions
634 between rectangular oligomers. We used the steady state approximation to
635 derive effective rate constants for these transitions, as described below.
636
We will illustrate the derivation of layer-based transition rate constants using a
specific example. Consider the layer-based reaction associated with adding a
longitudinal layer onto an oligomer that is, say, n tubulins tall and 4 tubulins wide
(in other words, the net reaction is (n,4) <--> (n+1,4). To simplify the notation, we
will call the starting species **4** and the final species **4'**; the intermediates with one,
two, and three tubulins on the layer will be called 4a, 4b, and 4c respectively. The
individual reactions are:
644

$$4 + \alpha\beta \rightleftharpoons 4a \qquad (1)$$
$$4a + \alpha\beta \rightleftharpoons 4b \qquad (2)$$
$$4b + \alpha\beta \rightleftharpoons 4c \qquad (3)$$
$$4c + \alpha\beta \rightleftharpoons 4' \qquad (4)$$

649
650
651 The associated rate equations for these reactions, which account for the
652 production and loss of intermediate species and also the fact that different
653 configurations present different numbers of binding sites (Sup. Fig. 8), are:
654

$$\frac{d}{dt}[4a] = 8k_f[4][\alpha\beta] - k_r^{Long}[4a] - \frac{6}{4}k_f[4a][\alpha\beta] + 2k_r^{Corner}[4b] \qquad (5)$$

$$\frac{d}{dt}[4b] = \frac{6}{4}k_f[4a][\alpha\beta] - 2k_r^{Corner}[4b] - \frac{4}{3}k_f[4b][\alpha\beta] + 2k_r^{Corner}[4c] \quad (6)$$

$$\frac{d}{dt}[4c] = \frac{4}{3}k_f[4b][\alpha\beta] - 2k_r^{Corner}[4c] - k_f[4c][\alpha\beta] \qquad (7)$$

$$\frac{d}{dt}[4'] = k_f[4c][\alpha\beta] \qquad (8)$$

659
660
661 Starting from the last intermediate 4c, we apply the steady-state approximation to
662 the intermediate species to obtain an expression for the net rate of producing 4'

663 from 4. Assuming that the concentration of 4c does not change with time, its time
664 derivative is zero, so:
665

$\frac{d}{dt}[4c] = 0$ (9)

$\Rightarrow \frac{4}{3}k_f[4b][\alpha\beta] - 2k_r^{Corner}[4c] - k_f[4c][\alpha\beta] = 0$ (10)

$\Rightarrow \frac{4}{3}k_f[4b][\alpha\beta] = [4c]\left(k_f[\alpha\beta] + 2k_r^{Corner}\right)$ (11)

$\Rightarrow [4c] = \frac{\frac{4}{3}k_f[\alpha\beta]}{\left(k_f[\alpha\beta] + 2k_r^{Corner}\right)}[4b]$ (12)

$\Rightarrow [4c] = W_3[4b], where\ W_3 = \frac{\frac{4}{3}k_f[\alpha\beta]}{\left(k_f[\alpha\beta] + 2k_r^{Corner}\right)}$ (13)

This yields an expression for the concentration of intermediate 4c in terms of its
predecessor 4b. The multiplicative factor W_3 is given by the pseudo-first-order
rate constant for forming 4c divided by the sum of the pseudo-first-order rate
constants for losing 4c (in the forward or reverse direction).
676

677 This process can be applied again to obtain an expression for the concentration
678 of intermediate 4b in terms of its predecessor 4a:
679

$\frac{d}{dt}[4b] = 0$ (14)

$\Rightarrow \frac{6}{4}k_f[4a][\alpha\beta] - 2k_r^{Corner}[4b] - \frac{4}{3}k_f[4b][\alpha\beta] + 2k_r^{Corner}[4c] = 0$ (15)

$\Rightarrow \frac{6}{4}k_f[4a][\alpha\beta] - 2k_r^{Corner}[4b] - \frac{4}{3}k_f[4b][\alpha\beta] + 2k_r^{Corner}W_3[4b] = 0$ (16)

$\Rightarrow [4b] = \frac{\frac{6}{4}k_f[\alpha\beta]}{\left(\frac{4}{3}k_f[\alpha\beta] + 2k_r^{Corner} - 2k_r^{Corner}W_3\right)}[4a]$ (17)

$\Rightarrow [4b] = W_2[4a], where\ W_2 = \frac{\frac{6}{4}k_f[\alpha\beta]}{\left(\frac{4}{3}k_f[\alpha\beta] + 2k_r^{Corner} - 2k_r^{Corner}W_3\right)}$ (18)

Where in (16) we eliminated [4c] by substituting $W_3[4b]$, and in (17) we simply
grouped terms in [4a] and [4b]. The weighting term W_2 is again given by a similar
ratio as for W3, but with an additional term that includes W3.
689

690 Applying the same procedure, we can obtain an expression for the concentration
691 of the first intermediate 4a in terms of its predecessor, the rectangular species 4:
692

$\frac{d}{dt}[4a] = 0$ (19)

$\Rightarrow [4a] = \frac{8k_f[\alpha\beta]}{\left(\frac{6}{4}k_f[\alpha\beta] + k_r^{Long} - 2k_r^{Corner}W_2\right)}[4]$ (20)

$\Rightarrow [4a] = W_1[4], where\ W_1 = \frac{8k_f[\alpha\beta]}{\left(\frac{6}{4}k_f[\alpha\beta] + k_r^{Long} - 2k_r^{Corner}W_2\right)}$ (21)

Finally, since the rate of finishing the layer is given by:

$\frac{d}{dt}[4'] = k_f[4c][\alpha\beta]$ (22)

$\frac{d}{dt}[4'] = k_f W_3 W_2 W_1 [4][\alpha\beta]$ (23)

Where in (23) we used (13) to eliminate [4c] for a term involving [4b], (18) to
eliminate [4b] for a term involving [4a], and (21) to eliminate [4a] for a term
involving [4]. This yields an apparent rate constant for adding a layer, the terms
of which only involve the fundamental rate constants and multiplicative factors
that account for the number of ways a particular reaction can happen. In the case
presented there are three such terms because the layer contains 4 tubulins.
Because k_r^{Corner} is small compared to $k_f[\alpha\beta]$, is can be neglected in the
denominator and so all weights excepting the first one will be numbers between 1
and 2. W_1 contains k_r^{Long} in the denomimator, which in general is large compared
to $k_f[\alpha\beta]$, so W_1 tends to be a small number that reflects the difficulty of initiating a
new layer. The general procedure is formulaic and can be applied to any size
layer.

The effective equilibrium ratio for the layer addition is given by

$$Layer\ equilibrium\ ratio\ = \frac{[4]}{[4']}$$

$$= \frac{K_d^{Long}(K_d^{Corner})^3}{[\alpha\beta]^4}$$

Finally, we compute an effective rate constant for layer loss by multiplying the
layer equilibrium ratio and the forward layer rate constant:

$k_r^{Layer} = \left(k_f^{Layer}\right)(Layer\ equilibrium\ ratio)$ (25)

$k_r^{Layer} = \left(k_f W_3 W_2 W_1 [\alpha\beta]\right)\left(\frac{K_d^{Long}(K_d^{Corner})^3}{[\alpha\beta]^4}\right)$ (26)

$k_r^{Layer} = \left(k_f W_3 W_2 W_1\right)\left(\frac{K_d^{Long}(K_d^{Corner})^3}{[\alpha\beta]^3}\right)$ (27)

The preceding derivation was performed for the specific case of adding a
longitudinal layer that is four subunits wide. For layers of size N, a similar logic
applies with the following differences: (i) the kinetic weighting factors for the first
through the (N-1)th subunit additions change to reflect the different number of
configurations (Sup. Fig. 8), and (ii) there is a different number of multiplicative
factors (W_is).

To determine if this reduced representation could accurately recapitulate the
timecourses for rectangular species in the explicit model, we constructed a new
model in Berkeley Madonna (Marcoline et al., 2020) (also truncated at 12-mers)
that considered monomer-based reactions to form the first rectangular
intermediate, a 2x2 tetramer of $\alpha\beta$-tubulin, but that only considered layer
transitions for larger species, using the layer rate laws described above. We input
expressions for the layer transition rates as derived above, and ran the
simulations using the same biochemical parameters used in the explicit
simulations. The layer model faithfully reproduced the results from the explicit
simulations (Fig. 1).
745
Layer-based simulations of microtubule self-assembly
To enable simulations of arbitrarily large oligomers, we wrote a custom computer
program that effectively performs the same calculations of Berkeley Madonna but
in a more automated way that makes it easier to incorporate a much larger
number of species and reactions. The program automates the generation of the
list of possible rectangular oligomers [(2,2), (3,2), (2,3), etc.] and the recursive
calculation of rate constants for transitions between them (as explained above
and in Sup. Fig. 8). The program then integrates the set of transition rate laws
using finite-difference methods. At each time point, the program uses longitudinal
and lateral affinities along with the concentration of 'free' tubulin (here free
means the population not incorporated into rectangular oligomers (2,2) or larger)
to calculate the concentration of free tubulin 'monomers' and of longitudinal and
lateral dimers, and trimers assuming these species are all in rapid equilibrium
(see box in Fig. 1E); using these quantities, the program then takes a small step
forward in time, updating quantities of all rectangular oligomers according to the
rate laws and the layer rate constants. This results in an updated set of
concentrations for the rectangular oligomers (and hence of free tubulin), and the
process is repeated. To account for microtubule formation, we simply converted
species that became 13-protofilaments wide into microtubules, and we assumed
that microtubules elongated by monomer addition into a 'corner' site. We did not
explicitly consider the seam or the variable configuration of the microtubule end,
and for simplicity we assumed plus- and minus-ends elongated with identical
kinetics. We used a downhill simplex algorithm to fit the simulated timecourses to
the experimental ones, using fraction of tubulin in rectangular oligomers as a
proxy for the normalized assembly data.
771
Derivation of the equilibrium constant for lateral association of a pre-
formed longitudinal dimer
To implement the dimer pathway (Fig. 2 and Sup. Fig. 4), we assume that the on-
rate constant for adding a dimer is the same as for adding a monomer. Given the
on-rate constant, we obtain the off-rate constant (for the initiating dimer to
dissociate) using the equilibrium constant for the dimer binding. We show here
that this new equilibrium constant can be calculated purely in terms of 'monomer'

properties. First, we write the dissociation constants in terms of free energies of
association, as follows:

$K_d^{Long} = e^{\Delta G^{Long} + \Delta S}$

$K_d^{Lat} = e^{\Delta G^{Lat} + \Delta S}$

$K_d^{Corner} = e^{\Delta G^{Long} + \Delta G^{Lat} + \Delta S}$

where we have separated out the entropic cost of subunit immobilization (loss of
rotational and translational degrees of freedom) as ΔS as has been done
previously (e.g. (Erickson and Pantaloni, 1981; VanBuren et al., 2002)). The
reason for doing this separation is that lateral association of a pre-formed
longitudinal dimer should have the same entropic 'cost' as for association of a
monomer (because the cost of losing rotational and translational degrees of
freedom is already built in to the longitudinal affinity). The K_d values are the fitted
parameters in the model.

What we wish to calculate is the dissociation constant for two simultaneous
lateral contacts:

$K_d^{2-Lat} = e^{2\Delta G^{lat} - \Delta S}$

We prove here that:

$K_d^{2-lat} = K_d^{Lat} * \left(K_d^{Corner} \middle/ K_d^{Long} \right)$

$K_d^{Lat} * \left(K_d^{Corner} \middle/ K_d^{Long} \right) = e^{\Delta G^{Lat} + \Delta S} * \left(e^{\Delta G^{Long} + \Delta G^{Lat} + \Delta S} \middle/ e^{\Delta G^{Long} + \Delta S} \right)$

$= e^{\Delta G^{Lat} + \Delta S} * \left(e^{\Delta G^{Long} + \Delta G^{Lat} + \Delta S - \Delta G^{Long} - \Delta S} \right)$

$= e^{\Delta G^{Lat} + \Delta S} * \left(e^{\Delta G^{Lat}} \right)$

$= e^{2*\Delta G^{Lat} + \Delta S}$

Simulating the effects of template-like nucleators
To investigate the effects of template-like nucleators in the model, we modified
the simulations to include an additional template species. We assumed a
constant total concentration of 'monomers' that were distributed into templating
oligomers of different widths. This assumption means that progressively wider
oligomers are present in the simulations at progressively lower concentrations;
for example, for a given concentration of templating monomers, there will be half
as many 4-wide oligomers as 2-wide, and so on. For calculating the rate
constants for the first layer addition onto the template, we assumed that only one
longitudinal interface of this templating oligomer (the plus end) was active for
nucleation.
In initial runs wherein interactions between γ- and αβ-tubulin were assumed to be
of identical affinity, we observed potent nucleation even from templates that were
only two subunits wide (Supplemental Fig. 7). Because two-wide nucleators like

the γ-TuSC are much less potent than 12-14-wide oligomers like the γ-TuRC

(Kollman et al., 2015; Oegema et al., 1999), we also performed simulations with

10-fold weaker interactions between γ- and $\alpha\beta$-tubulin than for $\alpha\beta$:$\alpha\beta$ interactions

(Fig. 6)

References

826 **References**
827
Akhmanova, A., and M.O. Steinmetz. 2015. Control of microtubule organization and
dynamics: two ends in the limelight. *Nat Rev Mol Cell Biol*. 16:711-726.
Alushin, G.M., G.C. Lander, E.H. Kellogg, R. Zhang, D. Baker, and E. Nogales. 2014.
High-resolution microtubule structures reveal the structural transitions in
alphabeta-tubulin upon GTP hydrolysis. *Cell*. 157:1117-1129.
Ayaz, P., S. Munyoki, E.A. Geyer, F.A. Piedra, E.S. Vu, R. Bromberg, Z. Otwinowski, N.V.
Grishin, C.A. Brautigam, and L.M. Rice. 2014. A tethered delivery mechanism
explains the catalytic action of a microtubule polymerase. *Elife*. 3:e03069.
Ayaz, P., X. Ye, P. Huddleston, C.A. Brautigam, and L.M. Rice. 2012. A TOG:alphabeta-
tubulin complex structure reveals conformation-based mechanisms for a
microtubule polymerase. *Science*. 337:857-860.
Barlan, K., and V.I. Gelfand. 2017. Microtubule-Based Transport and the Distribution,
Tethering, and Organization of Organelles. *Cold Spring Harb Perspect Biol*. 9.
Brouhard, G.J., and L.M. Rice. 2018. Microtubule dynamics: an interplay of
biochemistry and mechanics. *Nat Rev Mol Cell Biol*. 19:451-463.
Buey, R.M., J.F. Diaz, and J.M. Andreu. 2006. The nucleotide switch of tubulin and
microtubule assembly: a polymerization-driven structural change.
Biochemistry. 45:5933-5938.
Burton, P.R., R.E. Hinkley, and G.B. Pierson. 1975. Tannic acid-stained microtubules
with 12, 13, and 15 protofilaments. *J Cell Biol*. 65:227-233.
Chaaban, S., S. Jariwala, C.T. Hsu, S. Redemann, J.M. Kollman, T. Muller-Reichert, D.
Sept, K.H. Bui, and G.J. Brouhard. 2018. The Structure and Dynamics of C.
elegans Tubulin Reveals the Mechanistic Basis of Microtubule Growth. *Dev
Cell*. 47:191-204 e198.
Chalfie, M., and J.N. Thomson. 1982. Structural and functional diversity in the
neuronal microtubules of Caenorhabditis elegans. *J Cell Biol*. 93:15-23.
Consolati, T., J. Locke, J. Roostalu, Z.A. Chen, J. Gannon, J. Asthana, W.M. Lim, F.
Martino, M.A. Cvetkovic, J. Rappsilber, A. Costa, and T. Surrey. 2020.
Microtubule Nucleation Properties of Single Human gammaTuRCs Explained
by Their Cryo-EM Structure. *Dev Cell*. 53:603-617 e608.
Coombes, C.E., A. Yamamoto, M.R. Kenzie, D.J. Odde, and M.K. Gardner. 2013.
Evolving tip structures can explain age-dependent microtubule catastrophe.
Curr Biol. 23:1342-1348.
Davis, C., and K. Gull. 1983. Protofilament number in microtubules in cells of two
parasitic nematodes. *J Parasitol*. 69:1094-1099.
Desai, A., and T.J. Mitchison. 1997. Microtubule polymerization dynamics. *Annu Rev
Cell Dev Biol*. 13:83-117.
Driver, J.W., E.A. Geyer, M.E. Bailey, L.M. Rice, and C.L. Asbury. 2017. Direct
measurement of conformational strain energy in protofilaments curling
outward from disassembling microtubule tips. *Elife*. 6.
Duellberg, C., N.I. Cade, D. Holmes, and T. Surrey. 2016. The size of the EB cap
determines instantaneous microtubule stability. *Elife*. 5.

Erickson, H.P., and D. Pantaloni. 1981. The role of subunit entropy in cooperative
assembly. Nucleation of microtubules and other two-dimensional polymers.
Biophys J. 34:293-309.
Flyvbjerg, H., E. Jobs, and S. Leibler. 1996. Kinetics of self-assembling microtubules:
an "inverse problem" in biochemistry. *Proc Natl Acad Sci U S A*. 93:5975-
5979.
Fourniol, F.J., C.V. Sindelar, B. Amigues, D.K. Clare, G. Thomas, M. Perderiset, F.
Francis, A. Houdusse, and C.A. Moores. 2010. Template-free 13-protofilament
microtubule-MAP assembly visualized at 8 A resolution. *J Cell Biol*. 191:463-
470.
Gardner, M.K., B.D. Charlebois, I.M. Janosi, J. Howard, A.J. Hunt, and D.J. Odde. 2011.
Rapid microtubule self-assembly kinetics. *Cell*. 146:582-592.
Gaskin, F., C.R. Cantor, and M.L. Shelanski. 1974. Turbidimetric studies of the in vitro
assembly and disassembly of porcine neurotubules. *J Mol Biol*. 89:737-755.
Geyer, E.A., A. Burns, B.A. Lalonde, X. Ye, F.A. Piedra, T.C. Huffaker, and L.M. Rice.
2015. A mutation uncouples the tubulin conformational and GTPase cycles,
revealing allosteric control of microtubule dynamics. *Elife*. 4:e10113.
Geyer, E.A., M.P. Miller, C.A. Brautigam, S. Biggins, and L.M. Rice. 2018. Design
principles of a microtubule polymerase. *Elife*. 7.
Grishchuk, E.L., M.I. Molodtsov, F.I. Ataullakhanov, and J.R. McIntosh. 2005. Force
production by disassembling microtubules. *Nature*. 438:384-388.
Gunzelmann, J., D. Ruthnick, T.C. Lin, W. Zhang, A. Neuner, U. Jakle, and E. Schiebel.
2018. The microtubule polymerase Stu2 promotes oligomerization of the
gamma-TuSC for cytoplasmic microtubule nucleation. *Elife*. 7.
Holy, T.E., and S. Leibler. 1994. Dynamic instability of microtubules as an efficient
way to search in space. *Proc Natl Acad Sci U S A*. 91:5682-5685.
Kim, T., and L.M. Rice. 2019. Long-range, through-lattice coupling improves
predictions of microtubule catastrophe. *Mol Biol Cell*. 30:1451-1462.
Kollman, J.M., C.H. Greenberg, S. Li, M. Moritz, A. Zelter, K.K. Fong, J.J. Fernandez, A.
Sali, J. Kilmartin, T.N. Davis, and D.A. Agard. 2015. Ring closure activates
yeast gammaTuRC for species-specific microtubule nucleation. *Nat Struct Mol
Biol*. 22:132-137.
Kollman, J.M., A. Merdes, L. Mourey, and D.A. Agard. 2011. Microtubule nucleation by
gamma-tubulin complexes. *Nat Rev Mol Cell Biol*. 12:709-721.
Kuchnir Fygenson, D., H. Flyvbjerg, K. Sneppen, A. Libchaber, and S. Leibler. 1995.
Spontaneous nucleation of microtubules. *Phys Rev E Stat Phys Plasmas Fluids
Relat Interdiscip Topics*. 51:5058-5063.
Kwiatkowska, M., K. Poplonska, D. Stepinski, and Z. Hejnowicz. 2006. Microtubules
with different diameter, protofilament number and protofilament spacing in
Ornithogalum umbellatum ovary epidermis cells. *Folia Histochem Cytobiol*.
44:133-138.
Liu, P., E. Zupa, A. Neuner, A. Bohler, J. Loerke, D. Flemming, T. Ruppert, T. Rudack, C.
Peter, C. Spahn, O.J. Gruss, S. Pfeffer, and E. Schiebel. 2020. Insights into the
assembly and activation of the microtubule nucleator gamma-TuRC. *Nature*.
578:467-471.

Manka, S.W., and C.A. Moores. 2018. The role of tubulin-tubulin lattice contacts in
the mechanism of microtubule dynamic instability. *Nat Struct Mol Biol*.
25:607-615.
Marcoline, F., M. Grabe, S. Nayak, T. Zahnley, G. Oster, and R. Macey. 2020. Berkeley
Madonna User's Guide.
Margolin, G., I.V. Gregoretti, T.M. Cickovski, C. Li, W. Shi, M.S. Alber, and H.V.
Goodson. 2012. The mechanisms of microtubule catastrophe and rescue:
implications from analysis of a dimer-scale computational model. *Mol Biol
Cell*. 23:642-656.
McIntosh, J.R., E. O'Toole, G. Morgan, J. Austin, E. Ulyanov, F. Ataullakhanov, and N.
Gudimchuk. 2018. Microtubules grow by the addition of bent guanosine
triphosphate tubulin to the tips of curved protofilaments. *J Cell Biol*.
217:2691-2708.
Mitchison, T., and M. Kirschner. 1984. Dynamic instability of microtubule growth.
Nature. 312:237-242.
Moores, C.A., M. Perderiset, F. Francis, J. Chelly, A. Houdusse, and R.A. Milligan. 2004.
Mechanism of microtubule stabilization by doublecortin. *Mol Cell*. 14:833-
839.
Moritz, M., M.B. Braunfeld, J.W. Sedat, B. Alberts, and D.A. Agard. 1995. Microtubule
nucleation by gamma-tubulin-containing rings in the centrosome. *Nature*.
378:638-640.
Mozziconacci, J., L. Sandblad, M. Wachsmuth, D. Brunner, and E. Karsenti. 2008.
Tubulin dimers oligomerize before their incorporation into microtubules.
PLoS One. 3:e3821.
Nawrotek, A., M. Knossow, and B. Gigant. 2011. The determinants that govern
microtubule assembly from the atomic structure of GTP-tubulin. *J Mol Biol*.
412:35-42.
Oegema, K., C. Wiese, O.C. Martin, R.A. Milligan, A. Iwamatsu, T.J. Mitchison, and Y.
Zheng. 1999. Characterization of two related Drosophila gamma-tubulin
complexes that differ in their ability to nucleate microtubules. *J Cell Biol*.
144:721-733.
Pecqueur, L., C. Duellberg, B. Dreier, Q. Jiang, C. Wang, A. Pluckthun, T. Surrey, B.
Gigant, and M. Knossow. 2012. A designed ankyrin repeat protein selected to
bind to tubulin caps the microtubule plus end. *Proc Natl Acad Sci U S A*.
109:12011-12016.
Piedra, F.A., T. Kim, E.S. Garza, E.A. Geyer, A. Burns, X. Ye, and L.M. Rice. 2016. GDP-
to-GTP exchange on the microtubule end can contribute to the frequency of
catastrophe. *Mol Biol Cell*. 27:3515-3525.
Popov, A.V., F. Severin, and E. Karsenti. 2002. XMAP215 is required for the
microtubule-nucleating activity of centrosomes. *Curr Biol*. 12:1326-1330.
Prosser, S.L., and L. Pelletier. 2017. Mitotic spindle assembly in animal cells: a fine
balancing act. *Nat Rev Mol Cell Biol*. 18:187-201.
Rice, L.M., E.A. Montabana, and D.A. Agard. 2008. The lattice as allosteric effector:
structural studies of alphabeta- and gamma-tubulin clarify the role of GTP in
microtubule assembly. *Proc Natl Acad Sci U S A*. 105:5378-5383.

Roostalu, J., N.I. Cade, and T. Surrey. 2015. Complementary activities of TPX2 and
chTOG constitute an efficient importin-regulated microtubule nucleation
module. *Nat Cell Biol*. 17:1422-1434.
Roostalu, J., and T. Surrey. 2017. Microtubule nucleation: beyond the template. *Nat*
Rev Mol Cell Biol. 18:702-710.
Saito, K., and K. Hama. 1982. Structural diversity of microtubules in the supporting
cells of the sensory epithelium of guinea pig organ of Corti. *J Electron Microsc*
(Tokyo). 31:278-281.
Sept, D., and J.A. McCammon. 2001. Thermodynamics and kinetics of actin filament
nucleation. *Biophys J*. 81:667-674.
Thawani, A., R.S. Kadzik, and S. Petry. 2018. XMAP215 is a microtubule nucleation
factor that functions synergistically with the gamma-tubulin ring complex.
Nat Cell Biol. 20:575-585.
Thawani, A., M.J. Rale, N. Coudray, G. Bhabha, H.A. Stone, J.W. Shaevitz, and S. Petry.
2020. The transition state and regulation of gamma-TuRC-mediated
microtubule nucleation revealed by single molecule microscopy. *Elife*. 9.
Tobacman, L.S., and E.D. Korn. 1983. The kinetics of actin nucleation and
polymerization. *J Biol Chem*. 258:3207-3214.
Tucker, J.B., C.C. Paton, G.P. Richardson, M.M. Mogensen, and I.J. Russell. 1992. A cell
surface-associated centrosomal layer of microtubule-organizing material in
the inner pillar cell of the mouse cochlea. *J Cell Sci*. 102 (Pt 2):215-226.
VanBuren, V., L. Cassimeris, and D.J. Odde. 2005. Mechanochemical model of
microtubule structure and self-assembly kinetics. *Biophys J*. 89:2911-2926.
VanBuren, V., D.J. Odde, and L. Cassimeris. 2002. Estimates of lateral and
longitudinal bond energies within the microtubule lattice. *Proc Natl Acad Sci*
U S A. 99:6035-6040.
Voter, W.A., and H.P. Erickson. 1984. The kinetics of microtubule assembly. Evidence
for a two-stage nucleation mechanism. *J Biol Chem*. 259:10430-10438.
Wang, H.W., and E. Nogales. 2005. Nucleotide-dependent bending flexibility of
tubulin regulates microtubule assembly. *Nature*. 435:911-915.
Wieczorek, M., S. Bechstedt, S. Chaaban, and G.J. Brouhard. 2015. Microtubule-
associated proteins control the kinetics of microtubule nucleation. *Nat Cell*
Biol. 17:907-916.
Wieczorek, M., L. Urnavicius, S.C. Ti, K.R. Molloy, B.T. Chait, and T.M. Kapoor. 2020.
Asymmetric Molecular Architecture of the Human gamma-Tubulin Ring
Complex. *Cell*. 180:165-175 e116.
Zakharov, P., N. Gudimchuk, V. Voevodin, A. Tikhonravov, F.I. Ataullakhanov, and E.L.
Grishchuk. 2015. Molecular and Mechanical Causes of Microtubule
Catastrophe and Aging. *Biophys J*. 109:2574-2591.
Zhang, R., G.M. Alushin, A. Brown, and E. Nogales. 2015. Mechanistic Origin of
Microtubule Dynamic Instability and Its Modulation by EB Proteins. *Cell*.
162:849-859.
Zhang, R., J. Roostalu, T. Surrey, and E. Nogales. 2017. Structural insight into TPX2-
stimulated microtubule assembly. *Elife*. 6.
Zheng, Y., M.L. Wong, B. Alberts, and T. Mitchison. 1995. Nucleation of microtubule
assembly by a gamma-tubulin-containing ring complex. *Nature*. 378:578-583.

Figure 1

1005
Figure 1 An explicit biochemical model for early steps in spontaneous
microtubule nucleation.
A Cartoon of the model. Individual αβ-tubulin heterodimers (ovals) can associate
longitudinally (blue arrows) or laterally (orange arrows). As oligomers grow in
size, they can present sites where an associating tubulin would simultaneously
make longitudinal and lateral interactions ("corner"; bold black arrows). The
resulting collection of species and reactions was implemented in
BerkeleyMadonna (Macey et al., 2009) for simulations. Species up to 6-mers are
shown, but the model includes species up to 12-mers. Rapid growth of the
number of oligomers and reactions with size made it impractical to consider even
larger oligomers.
B (left) Plot showing that 'rectangular' species dominate the simulation. The four
dominant species are shown, with values in parentheses indicating the height
and width for the oligomers. (right) The same data plotted on a logarithmic
concentration scale. Other rectangular oligomers are much less common. Non-
rectangular oligomers are not shown, but are also much rarer than the dominant
oligomers shown.
C A simplified 'layer growth' model that only considers rectangular oligomers.
Arrows indicate longitudinal (blue) or lateral (orange) layer growth. The resulting
collection of species and reactions was implemented in BerkeleyMadonna for
simulations (see Methods)
D Cartoon of a 'layer' addition reaction with associated rate constants. The
equilibrium ratio and rate for layer addition can be defined in terms of underlying
longitudinal, lateral, and corner affinities and a single on rate constant. An
effective equilibrium constant and an approximation of the effective forward rate
constant for this layer addition are shown. See Methods for derivations.
E Similar to **B** but showing results from simulations of the simplified layer model.
This much simpler model with many fewer species and reactions quantitatively
reproduces results from the explicit model.
F (left) Plot of the species count for all possible oligomers (black) and for the
subset of rectangular oligomers (red). The extrapolation to a 40-mer emphasizes
the magnitude of the simplification. (right) Similar plot but for the number of
biochemical reactions.
1039

Figure 2 Expansion of the layer model to arbitrarily large oligomers for simulating
spontaneous microtubule nucleation. Oligomers are allowed to become arbitrarily
tall. We assumed that when oligomers reached a width of 13, they 'closed' into a
tube and elongated by monomer addition.
A Spontaneous microtubule nucleation kinetics measured by turbidity ($A_{350\ nm}$) at
multiple concentrations of αβ-tubulin (10, 11, 12, 13, 14, 15 µM).
B Fits of the model (black dotted line) to individual assembly curves (solid grey
line) at the indicated concentration. Fits to individual curves are good. Inset text
shows the resulting longitudinal, lateral, and 'corner' (simultaneous longitudinal
and lateral interactions) affinities, which differ somewhat from run to run.
C Global fit of the model (black dashed lines) to the full set of assembly data
(grey lines). As evidenced by the poor fit at low concentration, the model
underestimates the concentration-dependence of the progress curves.
D Cartoon illustrating that longitudinal dimers of $\alpha\beta$-tubulin should contribute to
lateral growth. The associated forward rates for the indicated reactions are
shown.
E The relative contribution of 'monomer' and 'dimer' initiation pathways can be
estimated as indicated based on the affinities of longitudinal and lateral
interactions. Plot predicting the fractional contribution of a 'dimer' pathway for
lateral growth in terms of the relative strength of longitudinal and lateral
interactions. The red star indicates the ratio obtained from the fit shown in **B**.
Those parameters predict that the 'dimer' pathway should dominate the
assembly, but it is not accounted for in our model.
1063

Figure 3 Incorporating a dimer pathway for lateral growth into the model
A Fits of the new dimer model (black dotted line) to individual assembly curves
(solid grey line) at the indicated concentrations. Fits to individual curves are
worse than for the monomer only model (compare to Fig. 2B). Inset text shows
the resulting longitudinal, lateral, and 'corner' (simultaneous longitudinal and
lateral interactions) affinities; the corner affinity varies almost two-fold over for 11
and 15 µM fits. The lateral affinity is ~200-fold weaker than for 'monomer model'
fits, reflecting the dominance of the dimer pathway in this model.
B Global fit of the model (black dashed lines) to the full set of assembly data
(grey lines). The fit is poor, with the model predicting too fast assembly at all
concentrations.
1075

A

Width = 2
Stronger Lateral
interaction

Width > 2
Weaker Lateral
Interaction

B

C

P = 8

K_d^{corn}: 11.5 nM
K_d^{long}: 5.9 mM
K_d^{late}: 1.95 M

Figure 4 A 'penalty' that mimics the cost of $\alpha\beta$-tubulin straightening markedly
improves the performance of the dimer model
A Cartoons illustrating $\alpha\beta$-tubulin straightening, which is thought to occur
progressively with width (the number of lateral neighbors). To model this in a very
simple way, we introduced a penalty to make lateral interactions weaker for the
3rd and successive lateral interactions.
B Performance of the 'dimer with penalty' model as a function of penalty
strength. The improved goodness of fit from the global optimization is not very
sensitive to the magnitude of the penalty, as long as the penalty is ~8 or greater.
C Illustration of the model fit to the experimental data for a penalty value of 8.
Inset shows fitted affinities (the corner affinity given here is for width=2
interactions, i.e. before application of the penalty).
1088
1089

A

B

C

Figure 5 Different ways of representing the predicted pathway for spontaneous
microtubule assembly
A The average size of intermediate oligomers (black curve) increases slowly over
time. The total concentration of intermediate oligomers increases and then
decreases, an effect that can be explained by depletion of free αβ-tubulin, and
the formation of microtubules.
B Summed concentration for all oligomers of a given width, plotted as a function
of time. Oligomers with width=2 dominate. Wider oligomers become
progressively rarer and peak at increasingly late times, reflecting the obstacles to
lateral growth. The vertically striped light grey curve behind the others shows the
extent of total assembly (right y-axis).
C Distribution of oligomer heights as a function of their width. The most
populated height is plotted as a filled circle for each width (different colors), at the
time that width peaks. Other height oligomers at a given width are plotted as
open circles, with the radius reflecting their concentration relative to the peak
species. The time to transition from one width to the next decreases as oligomers
get wider and taller, and the spread of oligomer heights increases as oligomers
get wider.
1108
1109

A

Nucleation-elongation:
Structural nucleus formed in one step

B

Phenomenological
Structural nucleus formed in multiple steps

elongation

1 αβ at a time

‘nucleus’

‘nucleus’

elongation

1 αβ at a time

*Steps from nucleus to a cylindrical
microtubule are not accounted for*

C

Accretion
No structural nucleus
‘Swath’ of pathways
Rate of assembly increases with size

elongation

1 αβ at a time

Figure 6 Summary of accretion model and comparison with nucleation-
elongation models
A Cartoon illustrating a classic nucleation-elongation model in which the rate of
polymer formation is limited by the concentration of a rare species (the nucleus)
that is in a rapid equilibrium with unpolymerized subunits (ovals). Such models
cannot account for the additional steps/species in between the nucleus and the
formation of the cylindrical microtubule.
B Cartoon illustrating a phenomenological model (Flyvbjerg et al., 1996) that
differs from nucleation-elongation model in that the nucleus is formed in multiple
steps. In this model the rate of polymer formation is also limited by the
concentration of a rare species (the nucleus), and there is no accounting for the
additional steps/species in between the nucleus and the formation of the
cylindrical microtubule.
C Cartoon illustrating the accretion model presented in this paper. In this model
there are many pathways to form a cylindrical microtubule, and all 'layer
accretion' steps are described. In this accretion model, and in contrast with
nucleation-elongation and phenomenological models, most of the steps toward
polymer formation are energetically favorable but kinetically frustrated by the
difficulty of initiating new layers. These barriers to initiation decrease as
oligomers become larger, as in crystal growth, and it is the ensemble of
pathways and barriers that define the kinetics of microtubule formation.

Figure 7 Implications for templating microtubule assembly

A Cartoon illustrating an intermediate stage of spontaneous nucleation. 2x2

oligomers of $\alpha\beta$-tubulin (left) assemble as described into rectangular

intermediates that tend to be taller than they are wide (middle) because of the

barriers to lateral growth. Arrows indicate multiple layer additions, each one

associated with a delay.

B When $\alpha\beta$-tubulin adds to a γ-TuRC-like template, cartooned here as a row of

blue ovals, the barrier to lateral growth is eliminated because 'corner' sites form

at the border of the $\alpha\beta$-tubulin oligomer. This accelerates the accretion process

(bolder arrows indicating faster transitions) and the resulting oligomers now tend

to be wider than they are tall

C Simulated nucleation curves in the presence of γ-tubulin templates of different

width assuming γ:$\alpha\beta$ interactions are weaker than $\alpha\beta$:$\alpha\beta$ interactions. The total

concentration of templating monomers is held constant. Templates that are only

two subunits wide (bright green) show a modest effect, but wider templates are

increasingly potent. The vertical line indicates the 175 s timepoint that we used

for analysis in the next panel.

D Quantification of the data from C. Wider templates are more potent than

narrower ones. Templating the earliest step in nucleation (from 2-wide to 3-wide)

yields the most dramatic decrease in $T_{0.1}$.

E Fraction of αβ-tubulin incorporated into microtubules at an early time (here,

175 s, see vertical line in panel C) plotted as a function of template width and

using data from C. Relative assembly at a given time represents the fraction

assembled for a given template divided by the fraction assembled in the

untemplated control. The changes in nucleation kinetics can result in large

changes in the overall degree of microtubule assembly at early times.

1157

Illustration of the meaning of 'non-redundant' oligomers

There are 6 distinct αβ-tubulin tetramers with 2 longitudinal bonds and 1 lateral bond

These 4 species present :
   1 corner (c),
   3 longitudinal (L), and
   5 lateral (l) binding sites

These 2 species present :
   2 corner (c),
   2 longitudinal (L), and
   4 lateral (l) binding sites

All 6 oligomers can be modeled using a single species
that presents the appropriate, population-weighted
number of different binding site types:

Population-weighted average:
   1.33 corner (c),
   2.67 longitudinal (L), and
   4.67 lateral (l) binding sites

Supplemental Figure 1 Illustration of the concept and implementation of 'non-
redundant' oligomers. Representative configurations of an $\alpha\beta$-tubulin tetramer
containing two longitudinal and one lateral bond are indicated. These are
indistinguishable from the standpoint of binding energy, but differ in terms of the
number and type of binding sites they present. Our algorithm simplifies these six
species into one and uses population weighted rate constants to capture the
'average' number and type of binding sites presented.
1165
1166

Supplemental Figure 2 In the explicit model, relatively tight corner affinity is
required for appreciable assembly into larger oligomers.
A Plots of the concentration of 'rectangular' oligomers vs time, for different values
of the corner affinity (color coding is given in panel C and is identical to that used
in Fig. 2B). Note different y-axis ranges for different plots. At corner affinity of 400
nM and weaker, assembly into species larger than a tetramer of $\alpha\beta$-tubulin (dark
blue curve) is negligeable.
B As in A, but using a log-scale y-axis to show the dramatic fall-off in the
concentration of larger species.
C Legend showing the correspondence between color and oligomer dimensions.
1177
1178

Supplemental Figure 3 Fit to an independent dataset

A Two independent data sets. The set plotted in grey is the one analyzed in the

main text. The set in black was collected independently. Tubulin concentrations

were 10, 11, 12, 13, 14, 15 μM for each. Most of the curves agree very closely,

with the exception of two outlier curves (10 and 12 μM, indicated with # and *,

respectively). The 10 μM curve was omitted from panel C.

B Fits of the layer model (black dotted line) to individual assembly curves (solid

grey line) at different concentrations. Fits to individual curves are good. Inset text

shows the resulting longitudinal, lateral, and 'corner' (simultaneous longitudinal

and lateral interactions) affinities, which are within 10% of those obtained in the

main text (see Fig. 2B).

C Global fit of the model (black dashed lines) to the alternate set of assembly

data (grey lines). The model underestimates the concentration-dependence of

the progress curves.

Illustration of the dimer pathway

Initiation by lateral addition
of a **longitudinal dimer**

'Zipping' of the layer by subsequent
addition of **monomers**

Supplemental Figure 4 Illustration of the dimer pathway.

Cartoon sequence of steps for adding a new lateral layer. In the dimer pathway,

a new lateral layer can be initiated by association of a pre-formed longitudinal

dimer (left). Because the concentration of these longitudinal dimers of $\alpha\beta$-tubulin

is much less than that of $\alpha\beta$-tubulin 'monomers', subsequent 'zipping' of the layer

occurs by monomer addition. See Methods for a derivation of the equilibrium

binding constant for the initiating assocition.

1200

1201

K_d^{corn}: 8.5 nM
K_d^{long}: 7.6 mM
K_d^{lat}: 2.15 M

Supplemental Figure 5 Fit of the dimer+penalty model to an independent
dataset
Global fit of the dimer+penalty model (black dotted lines) to the alternate set of
assembly data (grey lines). The global fit is good (with the exception of the 12 µM
outlier curve noted in Sup. Fig. 3). The fitted parameters are close to those
obtained from fitting to the original dataset, but for this dataset the corner affinity
is ~1.4-fold stronger and the longitudinal affinity is ~1.3-fold weaker.
1209

Supplemental Figure 6 Sensitivity of model predictions to variations in the
underlying parameters.
The dimer+penalty model was run with individual affinities made stronger (solid
lines) or weaker (dashed lines) by a factor of 1.5. Two different concentrations
are illustrated. The model predictions are comparably sensitive to variation in
corner (black) and longitudinal (blue) affinities, and less sensitive to variation in
the lateral affinity (orange).

Supplemental Figure 7 Simulations of templated microtubule assembly.
Simulated nucleation curves in the presence of γ-tubulin templates of different
width assuming γ:αβ interactions are the same strength as αβ:αβ interactions.
Compare to Fig. 6. The total concentration of templating monomers is held
constant. Templates that are only two subunits wide (bright green) show much
stronger nucleation than when we assumed weaker γ:αβ interactions (compare
green curve to the one in Fig. 6).
1224

Origin of the weighting factors used in the steady-state derivation (equations 5-8)

 To initiate a new longitudinal layer, there are 8 (=2*width) different places for an αβ-tubulin to associate. This explains the factor of 8 in equation 5.

Two 'singleton' configurations present one corner site and two present two corner sites.
On average there is 1.5 corner sites presented, hence the factor of 6/4 in equations 5 and 6.

 If the initiating αβ-tubulin is at the extreme left or extreme right, only 1 open corner site is presented.

 If the initiating αβ-tubulin is at either middle site, 2 open corner site are presented.

With two tubules added, two configurations present one corner site and one presents two
corner sites. On average there is 1.33 corner sites, hence the factor of 4/3 in equations 6 and 7.

 After the second addition, there are two configurations that present 1 open corner site

 But there is only one configuration that presents two open corner sites

After the third addition, there are only two configurations that each present 1 open corner site.
No weighting is applied.

The explanations above explain the weighting factors on k_f in Equations 5, 6, and 7. Weights for a layer of size N can be similarly obtained from a general expression. For a layer of size n, the weights for the first, second, third, etc associations are: 2n, 2(n-1)/n, 2(n-2)/(n-1), etc.

1225 **Supplemental Figure 8** Derivation of kinetic weighting factors
1226 Cartoons illustrating the sequence of intermediates for a 'layer addition',
1227 annotated to indicate the origin of the weighting factors applied in equations 5-8.
1228 Illustrations are made for the specific case of a layer of size 4, but the general
1229 expression for the coefficients is provided at the bottom of the figure.
1230
1231