1 **Microtubules form by progressively faster tubulin accretion, not by**
2 **nucleation-elongation**
3
4 Luke M. Rice[1], Michelle Moritz[2], David A. Agard[2,3]
5
6 [1] Departments of Biophysics and Biochemistry, UT Southwestern Medical
7 Center, 5323 Harry Hines Blvd., Dallas TX 75390
8
9 [2] Department of Biochemistry and Biophysics, UC San Francisco, 600 16th St.,
10 San Francisco CA 94143
11
12 [3] Howard Hughes Medical Institute, San Francisco, CA
13
14   Correspondence: Luke.Rice@UTSouthwestern.edu, Agard@msg.ucsf.edu
15
16

Abstract

Microtubules are highly dynamic polymers that play fundamental roles in all

eukaryotes. As the control of microtubule nucleation and regulation is central to

their function, there has long been interest in obtaining quantitative descriptions

of microtubule polymerization. Textbooks have focused on variations of a

nucleation-elongation mechanism: monomers are in rapid equilibrium with an

unstable oligomer (nucleus), but once the nucleus forms, the polymer grows by

monomer addition. While such mechanisms readily capture the actin assembly

process, they are inadequate to describe the physical mechanism by which the

much larger and more complex microtubules assemble. Here we develop a new

model for microtubule self-assembly that has three key features: i) microtubules

initiate via the formation of closed, 2D sheet-like structures which grow faster the

larger they become; ii) the dominant pathway proceeds via addition of complete

longitudinal or lateral layers; iii) as predicted by structural studies and the lattice

model, the formation of lateral interactions requires payment of an energetic

penalty for straightening early structures. This model quantitatively fits

experimental assembly data, and provides important insights into biochemical

determinants and assembly pathways for microtubule nucleation.

Introduction

The microtubule cytoskeleton plays essential roles organizing the interior of eukaryotic cells: microtubules can act as "tracks" for motor-based transport, they make up the mitotic spindle that mediates proper segregation of chromosomes during cell division, they help define cell polarization, and more (Desai and Mitchison, 1997; Prosser and Pelletier, 2017; Barlan and Gelfand, 2017). To fulfill these many roles, microtubules must assemble into a multitude of highly dynamic and spatially diverse networks. The dynamic instability of microtubules (Mitchison and Kirschner, 1984), which describes the switching of individual polymers between phases of growing and shrinking, allows wholesale reorganization of the microtubule network to occur rapidly and also provides a mechanism to efficiently search space (Holy and Leibler, 1994). Indeed, a host of regulatory factors allow cells to control how fast microtubules grow and how frequently they switch from growing to shrinking (catastrophe) or from shrinking back to growing (rescue) (Akhmanova and Steinmetz, 2015), and also to control the creation of new microtubules (Wieczorek et al., 2015; Kollman et al., 2011; Roostalu and Surrey, 2017), a process called nucleation.

Microtubules are hollow, cylindrical polymers formed from $\alpha\beta$-tubulin heterodimers that interact with each other in two ways: stronger head-to-tail (longitudinal) interactions between $\alpha\beta$-tubulins make up the straight protofilaments, and weaker side-to-side (lateral) interactions hold protofilaments together. The mechanisms underlying the dynamic instability of existing microtubules are increasingly well understood (Alushin et al., 2014; Zhang et al., 2015; Manka and Moores, 2018; Duellberg et al., 2016; Gardner et al., 2011; Geyer et al., 2015; Piedra et al., 2016; Driver et al., 2017; VanBuren et al., 2002; 2005; Grishchuk et al., 2005), but our understanding of spontaneous microtubule nucleation remains relatively primitive (reviewed in (Roostalu and Surrey, 2017)). Multiple factors contribute to this difficulty: first, nucleation occurs very rarely, so measuring it directly is much harder than measuring the growing and shrinking of

existing microtubules. Second, the open, tube-like structure of the microtubule

also poses unique challenges to assembly. In most organisms, microtubules

contain 13 protofilaments, although there are clear examples of specialized

microtubules containing 11 or 15 protofilaments (Chalfie and Thomson, 1982;

Davis and Gull, 1983; Kwiatkowska et al., 2006; Burton et al., 1975; Saito and

Hama, 1982; Tucker et al., 1992; Chaaban et al., 2018). Regardless of

protofilament number, the hollow nature of the microtubule means that it takes

many more $\alpha\beta$-tubulin subunits to close a tube than it does to make a minimal

helical repeat for a simpler, helical polymer like actin. Perhaps not surprisingly

given this structural complexity, there is no consensus for the sequence of

intermediates that must be formed (pathway) during the transition from individual

subunits to a closed tube. To complicate matters, the structure of the free

heterodimer also changes from a bent confirmation in solution to a straight

conformation within the final microtubule (reviewed in (Brouhard and Rice,

2018)). Once thought to be mediated by an allosteric transition within the

heterodimer, it is now appreciated that this change occurs on the growing

microtubule (lattice model) (Rice et al., 2008; Nawrotek et al., 2011; Ayaz et al.,

2012). However, where in the assembly process the straightening takes place,

how much energy is required, and how this obligatory conformational change

affects nucleation all remain unknown. In addition to these factors, the

biochemical processes underlying catastrophe also act to antagonize nucleation

by destabilizing nucleation intermediates.

An early study of the kinetics of spontaneous microtubule nucleation, which used

solution conditions that suppress catastrophe to simplify the behavior, posited

that nucleation proceeded sequentially through two discrete intermediates (Voter

and Erickson, 1984). The data from this pioneering study were later

demonstrated to display a property called phenomenological scaling (illustrated

in Fig. 1 using our own data), which allowed a more robust analysis that

suggested the presence of additional intermediates (Kuchnir Fygenson et al.,

1995; Flyvbjerg et al., 1996). The pathway proposed in this later work contains

four distinct nucleation intermediates (6-mer, 9-mer, 12-mer, and 15-mer of $\alpha\beta$-

tubulin); the largest intermediate was postulated to represent a 'turning point'

after which elongation occurred by repeated additions of individual $\alpha\beta$-tubulin

subunits. Demonstrating phenomenological scaling for the spontaneous

nucleation data was important because it pointed to a common molecular

mechanism over the entire concentration range measured, and because it

provided a way to assess the self-consistency of experimental measurements. At

the same time, the authors of that study also acknowledged that it was not

possible to ascribe a biochemical interpretation to the species and rate constants

of the abstracted kinetic model they put forth (Flyvbjerg et al., 1996). Indeed,

when this modeling is applied to our own data, it actually predicts a non-integral

number of steps, each of which add a non-integral number of $\alpha\beta$-tubulin

heterodimers per step. Thus, while it is able to fit the kinetic data, this type of

model cannot accurately capture the physical-chemical details of the underlying

assembly process. Moreover, kinetic models such as these do not provide

information on the specific geometry of the proposed intermediates.

In this work, we sought to obtain more explicit insight into the molecular

mechanisms of spontaneous microtubule assembly. While additional complexity

and parameters could always be added, our primary goal was to explore the

simplest biochemical models that could faithfully explain the experimental data.

We build from the basic modes of $\alpha\beta$-tubulin interactions within a microtubule: a

single subunit can make a longitudinal interaction, a lateral interaction or make

both simultaneously (corner interaction). Starting with explicit simulations of the

biochemical reactions that can occur for all possible $\alpha\beta$-tubulin oligomers up to

12-mers, we observed that a very small number of oligomers represent the

dominant species. These oligomers are the "rectangular" ones in which the

number of longitudinal and lateral contacts is maximized, such that they have no

unfilled corners. Using this key insight, we developed an approximate model that

only considered these dominant species, allowing us to include arbitrarily large

oligomers while retaining a connection to the biochemical properties of individual

subunits. Fitting this approximate but biochemically faithful model to experimental

time courses for spontaneous microtubule nucleation yielded a new way to view

microtubule nucleation. Rather than proceeding via a rare, rate limiting

"nucleus", our model indicates that microtubule assembly involves the continuous

accretion of $\alpha\beta$-tubulins into a growing 2D lattice. Like crystallization processes,

the larger the assembly, the faster it grows. Beyond this, our model reveals the

dominant assembly pathways for forming a new microtubule, and provides new

insights into how exogeneous nucleators and other regulatory factors can

accelerate microtubule formation.

Results

Closed species dominate the assembly pathway

While phenomenological models for microtubule assembly can recapitulate

observed assembly kinetics, they do not provide a direct connection to the

sequence of assembly intermediates or to the underlying biochemical affinities.

To provide insight into the biochemical determinants of $\alpha\beta$-tubulin self-assembly,

we constructed an explicit model that includes all non-redundant tubulin

oligomers from dimers through dodecamers (Fig. 2A shows all intermediates

through hexamers). Here non-redundant means that we only consider a single

exemplar n-mer for each unique combination of longitudinal and lateral contacts

(Sup. Fig. 1 explains this using an example). We defined the biochemical

pathways that connect different oligomers (arrows in Fig. 2A) by assuming that

oligomers grow and shrink via addition and loss of individual $\alpha\beta$-tubulin subunits.

An explosion in the number of species and reactions with increasing size made it

impractical to explore oligomers larger than dodecamers (reaching dodecamers

requires ~200 species and ~600 reactions) (Fig. 2F).

We implemented the differential equations describing this discrete assembly

scheme in Berkeley Madonna (Macey et al., 2009) to simulate the assembly

pathway and kinetics. The only free parameters in this model are the affinities for

the three different $\alpha\beta$–tubulin interaction modes (longitudinal, lateral, and corner

(simultaneous longitudinal+lateral interaction) (Fig 2D); as is common, the on-

rate constant k_f was assumed (here we chose 10^6/sec based on experimental

observations and consistent with work from others (VanBuren et al., 2002;

Margolin et al., 2012; Zakharov et al., 2015; Coombes et al., 2013; McIntosh et

al., 2018)). We sought to identify conditions under which some appreciable

concentration of dodecamers would form, revealing that a relatively tight corner

affinity ($K_d^{corner} < 0.2$ μM; Sup. Fig. 2) was required for flux into the largest

oligomers. Importantly, the basic behavior is invariant to the choice of on-rate

constant, although the value chosen does affect the best-fit affinities. Under

conditions of high-affinity corner interactions, the simulations predominantly

populated a small number of 'closed' oligomers (Fig. 2B). More explicitly what

happens is that formation of an initial 2x2 oligomer occurs relatively quickly.

Because of the weak longitudinal and lateral affinities, the next larger species

containing a 'singleton' tubulin is poorly populated but presents an open, high-

affinity corner site. Filling this stabilizes the first subunit and in the case of a 2x2

start, fills the next longitudinal or lateral layer. On larger assemblies, additional

corners are vacant, which rapidly fill, essentially "zipping up" the layer. The net

consequence is that closed species predominate, and that once initiated, layer

completion occurs efficiently. This suggests that only considering transitions

between closed oligomers could provide a much simpler assembly pathway still

capable of quantitatively capturing the intricacies of microtubule assembly

behavior. Importantly, such a scheme allows much larger oligomers to be

simulated while maintaining a manageable number of reactions (e.g. Fig. 2C,F).

Whereas the reaction rates for each elemental reaction in the complete model

(above) are invariant, in the reduced model the rates for adding or losing a layer

are context dependent because they depend on the size and shape of the

oligomer. Using the steady-state approximation, we derived an expression for the

effective 'forward' rate constant governing addition of a longitudinal or lateral

layer (Fig. 2D shows a simple example, the general expression and its derivation

are provided in Methods). Considering the layer addition as a sequence of

individual steps also allowed us to derive a 'layer equilibrium constant', which

describes the expected populations of oligomers differing by a layer (Methods).

Notably, both the rate constant and equilibrium constant for layer addition are

defined in terms of the biochemical properties of individual subunits.

Consequently, and in contrast to what occurs with phenomenological models, the

reduction in the number of species and reactions (>10-fold for species and >35-

fold for reactions up to 12-mers) afforded by the layer pathway does not come at

the expense of a lost connection to the fundamental biochemical behavior.

We implemented the reduced, layer-based model into Berkeley Madonna to test

if it could recapitulate the results from our explicit simulations when using the

same input parameters for subunit affinities. Despite the dramatic reduction in

complexity, and with the exception of very low-abundance intermediate species,

the simplified layer-based simulations quantitatively reproduced the oligomer

concentrations and the kinetics of oligomer formation observed with the explicit

model (compare Fig. 2E to 2B). These initial benchmarking layer simulations only

considered oligomers up to dodecamers so that we could make a direct

comparison with the explicit model, but the formulaic nature of the oligomer sizes

and transition rates in the reduced model means that this approach can easily be

extended to much larger oligomers (Fig. 2F). Thus, the layer model provides a

tractable way to simulate the pathway and kinetics of microtubule assembly in

terms of the elemental biochemical interactions.

A simplified accretion pathway for spontaneous microtubule assembly

We wrote a custom program to expand the layer model to arbitrary sized oligomers without having to specify each one manually. Whenever an oligomer gained a 13th protofilament, we converted it into a microtubule. No additional penalty for seam formation was included. We assumed that once formed, microtubules elongate by capturing $\alpha\beta$-tubulin subunits one at a time in the open, pseudo-helical 'corner' sites on either end. In order to minimize the number of adjustable parameters, the same rate constants were used for events at either the plus- or minus-ends. These additional simplifying assumptions allow us to simulate spontaneous microtubule assembly (within the constraints of the layer model) without having to specify a particular nucleus or preferred assembly pathway.

We began testing our model by fitting it to a subset of our experimental assembly curves (taken from Fig. 1). The model fit these individual curves quite well (Fig. 3A), with minor discrepancies evident at low concentration. The optimized affinities differed somewhat from fit to fit, but shared common features: tight corner affinities ($K_d^{corn} < 100$ nM), weaker longitudinal affinities ($K_d^{long} \sim 5\text{-}15$ mM), and lateral affinities around 10-fold weaker than longitudinal affinities (Fig. 3A). We found it encouraging that the model could fit individual assembly curves, but to be convincing a single parameter set should be able to fit all the assembly curves simultaneously.

In a global fit to the set of assembly curves (Fig. 3B), a limitation of the model became apparent: it could not simultaneously capture the correct behavior over the broad concentration range measured (Fig 3B), even though each individual concentration could be well fit (Fig 3A). Because all transition rates scale with the chosen on-rate constant (see derivations in Methods), changing the on-rate would not improve the global fits. The concentration dependence can be quantified in part from the slope of a plot of the $\log(T_{0.1})$ vs log(concentration)

(inset Fig 1C, Fig 3C). The steeper this slope, the greater the spread in lag times

with concentration. This discrepancy between model (slope = 3.1) and

experiment (slope = 3.8) suggests that the model is missing some key behavior.

In re-examining our assumptions, we realized that it was too restrictive to only

consider monomer addition and loss as possible 'elemental' reactions within the

layer assembly pathway. In fact, a simple calculation demonstrates that a 'dimer

pathway' must be considered. Compared to the 'two step' monomer-based

pathway in which a first monomer associates weakly and only becomes 'locked

in' by a subsequent corner association (Fig. 3D, middle path), when the

longitudinal affinity is stronger than the lateral one it is more efficient to initiate a

lateral layer by adding a pre-formed longitudinal dimer (Fig. 3D, lower path). This

is true even though longitudinal dimers are present at very low concentrations

(roughly 1 part in a 1000 or ~10 nM). Our 'monomer only' model fit required a

longitudinal affinity ~10-fold stronger than the lateral one, but under those

conditions the dimer pathway, which was not accounted for in the monomer

model, should account for ~90% of the lateral layer initiations (Fig. 3E). Thus, the

reasonable looking fits do not make biochemical sense, because the monomer

model ignored a pathway that has the potential to dominate the flux through

oligomers.

Incorporating a dimer pathway fails to recapitulate experimental data

without invoking a switch in lateral affinity

The cartoons and calculations presented in Fig. 3D,E illustrate how the rate of

initiating a new lateral layer will be very sensitive to the concentration of $\alpha\beta$-

tubulin longitudinal dimers. By contrast, once the slow step of initiating a new

layer has occurred, zipping up occurs almost exclusively via addition of $\alpha\beta$-

tubulin 'monomers', which far exceed longitudinal dimers in

number/concentration. It is not necessary to include a dimer pathway for initiating

a longitudinal layer: lateral dimers are very rare, and the 'upper' $\alpha\beta$-tubulin of a

longitudinal dimer provides no increased stabilization as a longitudinal layer

builds up.

Incorporating a parallel, dimer-based pathway for initiating lateral layers requires

that at each step we specify (i) the concentration of longitudinal dimers, (ii) an

association rate constant for the longitudinal dimers adding to a complete layer,

and (iii) a dissociation rate constant for the lateral 'unbinding' of a longitudinal

dimer (Sup. Fig. 3). The concentration of longitudinal dimers is easily calculated

given the concentration of monomers and the equilibrium constant for

longitudinal interactions. Because off-rate constants tend to vary over a much

larger range than on-rate constants, for simplicity we used the same, roughly

diffusion limited association rate constant as we do for monomers. We derived a

formula that expresses the equilibrium constant for lateral association of a

longitudinal dimer solely in terms of biochemical properties of the monomers

(Sup Fig. 3). Parameterizing the 'dimer pathway' in terms of monomer

biochemistry thus allows us to incorporate a higher-order reaction without losing

connection to the fundamental monomer properties.

We first fit the dimer model to individual assembly curves, as we did for the

monomer model (Fig. 4A). The longitudinal and corner affinities for these dimer

model fits were comparable to those obtained from fits of the monomer model.

The lateral affinity (~25-30 M in the dimer model) was ~30-fold weaker than the

in the monomer model (~100 mM), reflecting a greatly diminished contribution of

the monomer pathway for lateral initiation. While required biochemically,

including a dimer pathway worsened the ability to capture the shape of individual

assembly curves compared to the monomer-only fits. For the dimer pathway, the

simulated assembly curves displayed a less pronounced lag phase and also

appeared less cooperative (shallower sigmoid transition; compare Fig. 4A to Fig.

3A). Global fits of the dimer model to the set of assembly curves were markedly

poorer than for the monomer model – in addition to showing a too shallow

transition after the lag phase, the concentration-dependence of the characteristic

time ($T_{0.1}$) for assembly was too low (Fig 4C). Thus, a more complete

biochemical model for the assembly process paradoxically degraded its

predictive power.

Why was the monomer-only model able to fit the observed assembly curves so

much better than a model incorporating a dimer pathway? The presence of a

dimer pathway makes the effective rates of lateral and longitudinal growth much

more comparable to each other (Fig. 3D, compare top and bottom rate

expressions), whereas in the monomer-only model there is significantly more

contrast between the rates for longitudinal and lateral growth. We speculated that

the otherwise more exact monomer+dimer model must still be missing a critical

feature that acts to penalize lateral growth.

Previous structural studies provide a physical justification for such a penalty

(Pecqueur et al., 2012; Nawrotek et al., 2011; Ayaz et al., 2012; Wang and

Nogales, 2005). Indeed, it is now well established that unpolymerized $\alpha\beta$-tubulin

subunits adopt a 'curved' conformation in solution and that the microtubule lattice

itself acts as an allosteric activator to 'straighten' $\alpha\beta$-tubulin during the assembly

process (Buey et al., 2006; Rice et al., 2008; Ayaz et al., 2012; Nawrotek et al.,

2011). That is, $\alpha\beta$-tubulin becomes progressively straighter as it becomes

increasingly surrounded by lateral neighbors (reviewed in (Brouhard and Rice,

2018)) (Fig 5A); this straightening-induced conformational strain should work

against lateral interactions, weakening those contacts. To explore these ideas in

the context of our model, we introduced a very simple lateral penalty to mimic the

greater strain intrinsic to straighter conformations: we assumed that the lateral

interactions after the first pairing in a given 'row' were X-fold weaker.

We fit the monomer+dimer+penalty model to the data as before, testing a range

of penalty magnitudes (from 1 to 16 in Fig. 5B). The global fit to the experimental

curves improved as the magnitude of the lateral penalty increased, with the best

fits reached for penalty values greater than 4. Thus, penalizing lateral growth was

sufficient to 'correct' the problems associated with the presence of a dimer

pathway, and this correction only weakly depends on the value of the penalty, as

long as the penalty is sufficiently large (Fig. 5B). A representative global fit

(penalty value = 8, Fig. 5C,D) shows that the fit to the data is even better than for

the monomer-only model (see Fig. 3B,C). Runs with other penalty factors gave

fits of comparable quality, but as expected yielded somewhat different

longitudinal and lateral affinities. Collectively, our computational experiments led

us to what we believe to be the first simple, biochemically-based model that can

recapitulate the observed concentration-dependent kinetics of spontaneous

microtubule assembly.

A kinetically frustrated pathway for spontaneous microtubule assembly

The model developed here provides a biochemically-based pathway for

microtubule assembly and explains why the kinetics of initial assembly are slow

relative to polymer elongation. To describe this pathway, in what follows we focus

on the width (lateral extent) of oligomers because the endpoint of the process --

closure of the cylindrical lattice – typically requires a minimal width (fixed at 13-

wide in our model).

The summed concentration of different width tubulin oligomers as a function of

time reveals that the concentration of oligomers decreases as width increases –

there are fewer 3-wide oligomers than 2-wide, fewer 4-wide then 3-wide, etc.

(Fig. 6A). This means that the transition from one width to the next accelerates

throughout the assembly process. This acceleration can be understood as a

statistical/kinetic consequence of the two-dimensional assembly pathway. No

matter their height, all oligomers of a given width grow taller with the same

characteristic rate because they present the same number of 'landing sites' for

initiating a new longitudinal layer. However, taller oligomers grow wider faster

than shorter oligomers, because increased height provides more 'landing sites'

for initiating a new lateral layer. Similarly the wider oligomers add longitudinal

layers faster than the narrower ones. Thus, the two-dimensional nature of the

assembly pathway provides a simple explanation for why the assembly process

accelerates as it progresses.

To provide more granular insight into the nature of the predominant microtubule

assembly intermediates, we examined the distribution of oligomer sizes (Fig. 6B).

For each oligomer width, at the time the concentration of oligomers of that width

peaked, we plotted the most populated height (filled circle) and the relative

amount of taller or shorter oligomers (grey teardrop/oval shapes, which are

actually a set of circles with area scaled proportionally to the concentration). The

many parallel pathways in the assembly process leads to an increasing spread in

the range of oligomer heights as they get wider. For example, the most populated

50% of all 3-wide oligomers have heights between 7 and 11 (a height range of 5)

whereas the most populated 50% of all 12-wide oligomers span heights between

50 and 64 (a height range of 15, three times larger than for 3-wides). The most

populated immediate microtubule precursor (12 subunits wide) contains almost

700 tubulins: 58 subunits tall x 12 wide. This is much larger than the 12-18

subunits predicted by the Flyvbjerg nucleation-elongation model discussed above

(Fig 1). Also note that at no time prior to forming a closed microtubule, does the

rate of assembly equal the rate of elongation. The layer-based assembly model

thus provides an ensemble-based view of the assembly pathway wherein many

oligomers of varying height contribute appreciably to the formation of new

microtubules.

While the concentration of 2x2 species determines the point of entry for the

accretion-based assembly pathway, it is the straightening-induced reduction in

the kinetics of lateral growth that most clearly causes the lag phase. And it is the

kinetic consequences of this straightening penalty that are actually being

approximated in the classical nucleation-elongation models. Thus, the apparent

"nucleus" size (Fig. 1), is much closer to the number of tubulins in polymers of

width = 3 (~27, see Fig. 6), than with the number of tubulins in the 12-wide

immediate microtubule precursor species (~700, see Fig. 6). However, it should

be clear from the pronounced height variability (Fig 6B) that this is a statistical

average and does not represent a discrete nucleus.

Discussion

Many discussions of spontaneous microtubule formation have invoked the

concept of a 'critical nucleus', which is analogous to the transition-state of a

chemical reaction (Kollman et al., 2011). This traditional view of the assembly

pathway assumes that (i) the formation and growth of early intermediates is

thermodynamically unfavorable, and (ii) above a critical size, oligomer growth by

monomer addition becomes favorable and this elongation proceeds at a fixed

rate. The polymerization kinetics of relatively simple helical polymers like actin

can be well-described by two-step, nucleation-elongation models that invoke a

small critical nucleus representing a 'mini-filament' (the smallest oligomeric

species that presents helical binding sites) (Tobacman and Korn, 1983; Sept and

McCammon, 2001). In contrast to simpler filaments like actin, the spontaneous

assembly kinetics of hollow, cylindrical microtubules do not conform to the

simplest two-step, nucleation-elongation view (Kuchnir Fygenson et al., 1995;

Voter and Erickson, 1984; Flyvbjerg et al., 1996). Indeed, the size of the smallest

physically realistic oligomer that presents pseudo-helical binding sites (~26

tubulins) is ~2-fold larger than the size of the critical nucleus predicted from

kinetic analyses (~12-15 tubulins), and moreover both the time-dependence of

assembly and the concentration-dependence of the characteristic time ($T_{0.1}$) are

greater than would be expected for the two kinetic steps of the most basic

nucleation-elongation mechanism. Even more complex, multi-step nucleation-

elongation mechanisms (Flyvbjerg et al., 1996) fail to provide useful insights into

the assembly pathway.

Our model provides an alternative conceptual framework for thinking about the

biochemical pathway for microtubule formation, and the factors that limit the

overall rate of assembly. Indeed, because there is no singular, transition-state-

like species that dictates the assembly kinetics, we suggest that the term

"nucleation" is potentially misleading when considering microtubule assembly

and should be avoided. In what follows, we summarize the main features of the

predicted assembly pathway, and discuss implications for the multiple ways that

a template or other regulatory factor could accelerate the process.

The layer growth model predicts that a multitude of pathways contribute

meaningfully to microtubule assembly. Stronger longitudinal interactions cause

these intermediates to be taller than they are wide. Early in the assembly

pathway, transitions to larger oligomers become energetically favorable but

nevertheless occur slowly because adding a new layer depends on a weak and

unfavorable initiating longitudinal or lateral interaction. The rate of transition

between oligomers accelerates with increasing size, not because of any intrinsic

changes in affinity, but because larger oligomers present more sites on which to

stochastically initiate the next layer. These size-dependent changes in transition

rates are a specific consequence of the two-dimensional nature of the assembly

pathway. Even though nearly all of the layer additions are energetically 'downhill',

the repeated difficulty of initiating each new layer contributes a series of

progressively smaller delays into the assembly sequence. The overall ensemble

of initiation barriers thus creates a kind of 'frustration' that dictates the unique

kinetics of spontaneous microtubule assembly.

This layer-based view of the assembly pathway provides new ways for thinking

about how regulatory factors might accelerate microtubule initiation. Our

modeling indicates that the kinetics of microtubule formation are dictated by the

need for *repeated* 'layer' transitions that are limited by the weaker lateral affinity

and by the straightening penalty (e.g. Fig. 7A). Classical "nucleators" like the γ-

tubulin ring complex (γ-TuRC) (Zheng et al., 1995) contain a lateral array of 12-

14 γ-tubulins that are thought to form a 'template' on which a microtubule can

elongate (Kollman et al., 2015; Moritz et al., 1995; 2000; Kollman et al., 2011).

We wondered how the potency of templating might depend on the width of the

template. To investigate this question in the context of our model, we modified

our simulations to include various width templating oligomers (Fig. 7B), using a

constant total amount of templating 'monomers'. Thus, in the simulations there

are progressively lower and lower concentrations of the wider oligomers (e.g. half

as many 4-wide oligomers as 2-wide). The intrinsic affinity of $\alpha\beta$-tubulins for

γ is unknown, so we explored a regime where γ:$\alpha\beta$ longitudinal

interactions are identical to $\alpha\beta$:$\alpha\beta$ (Supplemental Fig. 4) and another where γ:$\alpha\beta$

longitudinal interactions are weaker than for $\alpha\beta$:$\alpha\beta$ (Fig. 7). With the stronger

γ:$\alpha\beta$ interaction strength, even 2-wide "templates" potently accelerated

assembly. This is at odds with experimental data showing that γ-tubulin small

complexes (γ-TuSCs), which only contain two γ-tubulins, are much poorer

nucleators than γ-TuRCs (Oegema et al., 1999; Kollman et al., 2015). By

contrast, when γ:$\alpha\beta$ interactions were weaker, templating oligomers that were 2

subunits wide only modestly accelerated the predicted assembly kinetics (1.1-

fold faster characteristic time for assembly, $T_{0.1}$) (Fig. 7C,D,E). A 3-wide oligomer

was much more potent (1.7-fold faster) even though in the simulations the

concentration of 3-wide oligomers was less than that of the 2-wide oligomer. This

strong change in potency occurs because the 'hardest' transition in the assembly

sequence is from two-wide oligomers to 3-wide oligomers, providing an

explanation for why γ-tubulin small complexes (γ-TuSCs) only weakly stimulate

assembly (Oegema et al., 1999). Progressively wider templating arrays show

even greater potency, emphasizing that the barriers to lateral growth contribute

throughout the entire assembly pathway.

The idea that multiple reactions/transitions contribute to limit the overall rate of

microtubule formation also suggests that strategies besides templating could

enhance the rate of spontaneous assembly. The rates for layer additions depend

on how fast tubulins 'land' on the growing sheet, so enhancing this rate should

promote layer initiation and 'zippering'. For example, XMAP215 family

polymerases use an 'accelerated delivery' mechanism (Ayaz et al., 2014; Geyer

et al., 2018) to make microtubules grow faster, and this same activity likely

underlies the role for XMAP215 family proteins in microtubule nucleation

(Roostalu et al., 2015; Wieczorek et al., 2015; Flor-Parra et al., 2018; Thawani et

al., 2018; Gunzelmann et al., 2018; Popov et al., 2002). The efficiency of

initiating new layers is primarily limited by the relatively weak longitudinal or

lateral affinities, especially for the singleton tubulin that initiates each layer. Thus,

a factor that stabilizes these associations should also increase the rate at which

layer transitions occur. Proteins like TPX2 and doublecortin bridge between

neighboring subunits in the lattice (Fourniol et al., 2010; Zhang et al., 2017); this

'molecular stapling' may help explain their documented roles in microtubule

"nucleation" (Roostalu et al., 2015; Moores et al., 2004). Finally, altering the

allosteric properties of $\alpha\beta$-tubulin itself should also profoundly effect microtubule

initiation. For example, and in keeping with predictions of our model, studies of

an $\alpha\beta$-tubulin 'conformation cycle' mutation (Geyer et al., 2015) that enhances

the strength of lateral interactions (Driver et al., 2017)(perhaps via reducing the

cost of $\alpha\beta$-tubulin straightening), greatly accelerates microtubule formation.

Similarly, small molecule compounds like taxol that stimulate microtubule

initiation and growth may do so by promoting straightening.

Conclusions and perspective

We described a relatively simple accretion model that explains spontaneous

microtubule assembly kinetics in terms of subunit biochemistry and a sequence

of 'closed' intermediates. Whereas in typical 'nucleation-elongation' paradigms

the scarcity of the transition-state-like nucleus is what limits the rate of polymer

formation, our model is more akin to a process like crystal growth. Indeed, in our

accretion model most of the steps toward polymer formation are energetically

favorable but kinetically frustrated by the difficulty of initiating a new layer; this

difficulty decreases as layers become larger because more landing opportunities

are presented. The accretion mechanism we describe is a direct consequence of

the two-dimensional nature of the hollow, cylindrical microtubule lattice.

We did not consider the kinetic consequences of GTP hydrolysis within the

nascent polymer in our model, because it was not possible to account for the

stochastic hydrolysis within the context of our deterministic formalism. In keeping

with that, our assay conditions were chosen to suppress microtubule

catastrophe. While GTPase activity should frustrate assembly by preferentially

destabilizing the smallest and least stable oligomers, we still expect that growth

through accretion as outlined here will represent the dominant pathway for

assembly.

Finally, our model provides a unifying quantitative framework for understanding

the complex effects of regulatory molecules. Indeed, the 'zippering' transitions

between closed oligomers that dominate the initiation process also share many

features in common with microtubule elongation. Thus, our model also explains

why factors that regulate microtubule elongation and shrinking also regulate

spontaneous microtubule assembly.

536
537

538 **Acknowledgements**
539 This study was supported by grants from the National Science Foundation to
540 LMR (MCB-1615938 and MCB-1054947) and from the National Institutes of

544 Author contributions
545 MM collected the microtubule assembly data and helped write the manuscript.
546 LMR constructed models, fit the data, and drafted the manuscript. DAA analyzed
547 data, supervised the research, and helped write the manuscript.
548
549

Methods

Measurements of spontaneous assembly

The spontaneous assembly of phosphocellulose-purified porcine-brain tubulin was followed by turbidity at 350 nm basically as described previously (Voter and Erickson, 1984; Gaskin et al., 1974) Care was taken to remove microtubule seeds and inactive protein by cycling it through an additional polymerization/depolymerization step just before use: immediately prior to performing assays, tubulin at –80°C was rapidly thawed at 37°, placed on an ice-water slurry (0°C), and the buffer was adjusted to 80 mM K-PIPES, pH 6.8, 1mM EGTA, 4 mM MgCl$_2$ and 1 mM GTP. After 5 min incubation, 0.5 volume of 37°C glycerol was mixed in and the tubulin was allowed to polymerize at 37°C for 20 min. Microtubules were pelleted through a 37°C, 60% glycerol cushion in 50 mM K-MES, pH 6.6, 5 mM MgSO$_4$, 1 mM EGTA, 1 mM GTP in a TLA110 rotor (Beckman, Palo Alto, CA) at 80,000 rpm, 20 min. The pellets were resuspended at 37°C in assembly buffer (AB: 50 mM K-MES, pH6.6, 3.4 M glycerol, 5 mM MgSO$_4$, 1 mM EGTA, 1 mM GTP) using a warm Potter homogenizer. The resuspended microtubules were then depolymerized on an ice-water slurry for 20 min and centrifuged in a TLA100.3 rotor at 100,000 rpm for 35 min at 2°C to remove any remaining polymerized tubulin. The supernatant from this cold spin was removed and used for the assembly reactions. The tubulin was kept at 0°C as reaction mixtures were prepared. A series of samples was prepared at different tubulin concentrations in AB and kept at 0°C in thin-walled, 0.5 ml plastic tubes (PCR tubes, Stratagene, Cedar Creek, TX). Samples were rapidly heated to 37°C directly in cuvettes pre-warmed to 37°C in a Pelltier cell changer (took 30 sec), in a custom built spectrophotometer. The A$_{350\,nm}$ was recorded approximately every 4 sec on 4 samples at a time until plateaus were reached. Recording on the spectrometer was started prior to transfer of tubulin to the cuvettes so that early lag-phase data were not missed. The time was adjusted to ignore the 30 sec heating step. Scaling analysis was performed using a customized program.

Explicit simulations in Berkeley Madonna

We constructed a model in Berkeley Madonna (Macey et al., 2009) to simulate αβ-tubulin self-assembly. The first step in this process was to define the set of αβ-tubulin oligomers and how they are related to each other by gain/loss of an αβ-tubulin though a longitudinal, lateral, or longitudinal+lateral ('corner') interaction. Starting from the two-dimensional nature of the microtubule lattice, we built a list of all possible configurations for an αβ-tubulin oligomer of a given size. Fig. 2 illustrates how there are two kinds of possible dimers of αβ-tubulin (longitudinal and lateral), three kinds of trimers of αβ-tubulin, etc. Larger oligomers can have multiple configurations that are energetically equivalent. To minimize the number of species and reactions considered in the model, we used a single exemplar species to represent each set of n-mers with the same number

of longitudinal and lateral contacts. For each exemplar species, population-
weighted average rate constants were used to account for the different properties
of each oligomer. Supplemental Figure 1 provides a worked example of this
approximation for a tetramer of $\alpha\beta$-tubulin. We then used the 'Chemical
Reactions' module in Berkeley Madonna (Macey et al., 2009) to build a set of
rate equations corresponding to each possible transition. We did not pursue
oligomers larger than 12-mers because the number of species and reactions
became intractable. This model provides a way to analyze the rate and extent to
which different species become populated for a given choice of the adjustable
parameters.
604
Derivation of layer transition rates and validation in Berkeley Madonna
The explicit, monomer-based simulations in Berkeley Madonna (Macey et al.,
2009) indicated that a simpler reaction scheme that only considered transitions
between 'closed' (rectangular) oligomers. We used the steady state
approximation to derive effective rate constants for these transitions, as
described below.
611
We will illustrate the derivation of layer-based transition rate constants using a
specific example. Consider the layer-based reaction associated with adding a
longitudinal layer onto an oligomer that is, say, n tubulins tall and 4 tubulins wide
(in other words, the net reaction is (n,4) <--> (n+1,4). To simplify the notation, we
will call the starting species **4** and the final species **4'**; the intermediates with one,
two, and three tubulins on the layer will be called 4a, 4b, and 4c respectively. The
individual reactions are:
619
620   $4 + \alpha\beta \rightleftharpoons 4a$ (1)
621   $4a + \alpha\beta \rightleftharpoons 4b$ (2)
622   $4b + \alpha\beta \rightleftharpoons 4c$ (3)
623   $4c + \alpha\beta \rightleftharpoons 4'$ (4)
624
625
626   The associated rate equations for these reactions, which account for the
627   production and loss of intermediate species, are:
628
629   $\frac{d}{dt}[4a] = 8k_f[4][\alpha\beta] - k_r^{Long}[4a] - \frac{6}{4}k_f[4a][\alpha\beta] + 2k_r^{Corner}[4b]$ (5)
630   $\frac{d}{dt}[4b] = \frac{6}{4}k_f[4a][\alpha\beta] - 2k_r^{Corner}[4b] - \frac{4}{3}k_f[4b][\alpha\beta] + 2k_r^{Corner}[4c]$ (6)
631   $\frac{d}{dt}[4c] = \frac{4}{3}k_f[4b][\alpha\beta] - 2k_r^{Corner}[4c] - k_f[4c][\alpha\beta]$ (7)
632   $\frac{d}{dt}[4'] = k_f[4c][\alpha\beta]$ (8)
633
634
635   Starting from the last intermediate 4c, we apply the steady-state approximation to
636   the intermediate species to obtain an expression for the net rate of producing 4'

637 from 4. Assuming that the concentration of 4c does not change with time, its time
638 derivative is zero, so:
639

$\frac{d}{dt}[4c] = 0$ (9)

$\Rightarrow \frac{4}{3}k_f[4b][\alpha\beta] - 2k_r^{Corner}[4c] - k_f[4c][\alpha\beta] = 0$ (10)

$\Rightarrow \frac{4}{3}k_f[4b][\alpha\beta] = [4c]\left(k_f[\alpha\beta]+2k_r^{Corner}\right)$ (11)

$\Rightarrow [4c] = \dfrac{\frac{4}{3}k_f[\alpha\beta]}{\left(k_f[\alpha\beta]+2k_r^{Corner}\right)}[4b]$ (12)

$\Rightarrow [4c] = W_3[4b]\,,where\ W_3 = \dfrac{\frac{4}{3}k_f[\alpha\beta]}{\left(k_f[\alpha\beta]+2k_r^{Corner}\right)}$ (13)

This yields an expression for the concentration of intermediate 4c in terms of its
predecessor 4b. The multiplicative factor W_3 is given by the pseudo-first-order
rate constant for forming 4c divided by the sum of the pseudo-first-order rate
constants for losing 4c (in the forward or reverse direction).
650

651 This process can be applied again to obtain an expression for the concentration
652 of intermediate 4b in terms of its predecessor 4a:
653

$\frac{d}{dt}[4b] = 0$ (14)

$\Rightarrow \frac{6}{4}k_f[4a][\alpha\beta] - 2k_r^{Corner}[4b] - \frac{4}{3}k_f[4b][\alpha\beta] + 2k_r^{Corner}[4c] = 0$ (15)

$\Rightarrow \frac{6}{4}k_f[4a][\alpha\beta] - 2k_r^{Corner}[4b] - \frac{4}{3}k_f[4b][\alpha\beta] + 2k_r^{Corner}W_3[4b] = 0$ (16)

$\Rightarrow [4b] = \dfrac{\frac{6}{4}k_f[\alpha\beta]}{\left(\frac{4}{3}k_f[\alpha\beta]+2k_r^{Corner} - 2k_r^{Corner}W_3\right)}[4a]$ (17)

$\Rightarrow [4b] = W_2[4a]\,,where\ W_2 = \dfrac{\frac{6}{4}k_f[\alpha\beta]}{\left(\frac{4}{3}k_f[\alpha\beta]+2k_r^{Corner} - 2k_r^{Corner}W_3\right)}$ (18)

Where in (16) we eliminated [4c] by substituting W_3[4b], and in (17) we simply
grouped terms in [4a] and [4b]. The weighting term W_2 is again given by a similar
ratio as for W_3, but with an additional term that includes W_3.
663

664 Applying the same procedure, we can obtain an expression for the concentration
665 of the first intermediate 4a in terms of its predecessor, the closed rectangular
666 species 4:
667

668 $\frac{d}{dt}[4a] = 0$ (19)

669 $\Rightarrow [4a] = \dfrac{8k_f[\alpha\beta]}{\left(\frac{6}{4}k_f[\alpha\beta]+k_r^{Long} - 2k_r^{Corner}W_2\right)}[4]$ (20)

670 $\Rightarrow [4a] = W_1[4], where\ W_1 = \dfrac{8k_f[\alpha\beta]}{\left(\frac{6}{4}k_f[\alpha\beta]+k_r^{Long} - 2k_r^{Corner}W_2\right)}$ (21)

671
672
673   Finally, since the rate of finishing the layer is given by:
674

676    $$\frac{d}{dt}[4'] = k_f[4c][\alpha\beta] \tag{22}$$

677    $$\frac{d}{dt}[4'] = k_f W_3 W_2 W_1[4][\alpha\beta] \tag{23}$$
675
678
Where in (23) we used (13) to eliminate [4c] for a term involving [4b], (18) to
eliminate [4b] for a term involving [4a], and (21) to eliminate [4a] for a term
involving [4]. This yields an apparent rate constant for adding a layer, the terms
of which only involve the fundamental rate constants and multiplicative factors
that account for the number of ways a particular reaction can happen. In the case
presented there are three such terms because the layer contains 4 tubulins.
Because k_r^{Corner} is small compared to $k_f[\alpha\beta]$, is can be neglected in the
denominator and so all weights excepting the first one will be numbers between 1
and 2. W_1 contains k_r^{Long} in the denomimator, which in general is large compared
to $k_f[\alpha\beta]$, so W_1 tends to be a small number that reflects the difficulty of initiating a
new layer. The general procedure is formulaic and can be applied to any size
layer.
691
692   The effective equilibrium dissociation constant for the layer addition is given by
693

694    $$K_d^{Layer} = \frac{[4']}{[4]} = \frac{K_d^{Long}(K_d^{Corner})^3}{[\alpha\beta]^4} \tag{24}$$
695
696   Finally, we compute an effective rate constant for layer loss by multiplying the
697   equilibrium dissociation constant and the forward rate constant:
698

700    $$k_r^{Layer} = \left(k_f^{Layer}\right)\left(K_d^{Layer}\right) \tag{25}$$

701    $$k_r^{Layer} = \left(k_f W_3 W_2 W_1[\alpha\beta]\right)\left(\frac{K_d^{Long}(K_d^{Corner})^3}{[\alpha\beta]^4}\right) \tag{26}$$

699    $$k_r^{Layer} = \left(k_f W_3 W_2 W_1\right)\left(\frac{K_d^{Long}(K_d^{Corner})^3}{[\alpha\beta]^3}\right) \tag{27}$$

702
To determine if this reduced representation could accurately recapitulate the
timecourses for rectangular species in the explicit model, we constructed a new
model in Berkeley Madonna (Macey et al., 2009) (also truncated at 12-mers) that
considered monomer-based reactions to form the first rectangular intermediate, a
2x2 tetramer of $\alpha\beta$-tubulin, but that only considered layer transitions for larger
species. We input expressions for the layer transition rates as derived above,
and ran the simulations using the same biochemical parameters used in the

explicit simulations. The layer model faithfully reproduced the results from the
explicit simulations (Fig. 2).
712
Derivation of the equilibrium constant for lateral association of a pre-
formed longitudinal dimer
To implement the dimer pathway (Fig. 3 and Sup. Fig. 3), we assume that the on-
rate constant for adding a dimer is the same as for adding a monomer. Given the
on-rate constant, we obtain the off-rate constant (for the initiating dimer to
dissociate) using the equilibrium constant for the dimer binding. We show here
that this new equilibrium constant can be calculated purely in terms of 'monomer'
properties. First, we write the dissociation constants in terms of free energies of
association, as follows:

$K_d^{Long} = e^{\Delta G^{Long} + \Delta S}$

$K_d^{Lat} = e^{\Delta G^{Lat} + \Delta S}$

$K_d^{Corner} = e^{\Delta G^{Long} + \Delta G^{Lat} + \Delta S}$

where we have separated out the entropic cost of subunit immobilization (loss of
rotational and translational degrees of freedom) as ΔS as has been done
previously (e.g. (Erickson and Pantaloni, 1981; VanBuren et al., 2002)). The
reason for doing this separation is that lateral association of a pre-formed
longitudinal dimer should have the same entropic 'cost' as for association of a
monomer (because the cost of losing rotational and translational degrees of
freedom is already built in to the longitudinal affinity). The K_d values are the fitted
parameters in the model.
734
735 What we wish to calculate is the dissociation constant for two simultaneous
736 lateral contacts:

737 $K_d^{2-Lat} = e^{2\Delta G^{lat} - \Delta S}$

738
We prove here that:

$K_d^{2-lat} = K_d^{Lat} * \left(K_d^{Corner} \Big/ K_d^{Long} \right)$

$K_d^{Lat} * \left(K_d^{Corner} \Big/ K_d^{Long} \right) = e^{\Delta G^{Lat} + \Delta S} * \left(e^{\Delta G^{Long} + \Delta G^{Lat} + \Delta S} \Big/ e^{\Delta G^{Long} + \Delta S} \right)$

$= e^{\Delta G^{Lat} + \Delta S} * \left(e^{\Delta G^{Long} + \Delta G^{Lat} + \Delta S - \Delta G^{Long} - \Delta S} \right)$

$= e^{\Delta G^{Lat} + \Delta S} * \left(e^{\Delta G^{Lat}} \right)$

$= e^{2 * \Delta G^{Lat} + \Delta S}$

745
Layer-based simulations of microtubule self-assembly
To enable simulations of arbitrarily large oligomers, we wrote a custom computer
program to implement and automate the construction of the list of possible
oligomers and the recursive calculation of rate constants for transitions between
them, and to integrate the rate laws using finite-difference methods. To simulate

spontaneous microtubule assembly, we converted species that reached 13-
protofilaments wide into microtubules, and we assumed that microtubules
elongated by monomer addition into a 'corner' site. We did not explicitly consider
the seam, and for simplicity we assumed plus- and minus-ends elongated with
identical kinetics. We used a downhill simplex algorithm to fit the simulated
timecourses to the experimental ones.
757
Simulating the effects of template-like nucleators
To investigate the effects of template-like nucleators in the model, we modified
the simulations to include an additional template species. We assumed a
constant total concentration of 'monomers' that were distributed into templating
oligomers of different widths. This assumption means that progressively wider
oligomers are present in the simulations at progressively lower concentrations;
for example, for a given concentration of templating monomers, there will be half
as many 4-wide oligomers as 2-wide, and so on. For calculating the rate
constants for the first layer addition onto the template, we assumed that only one
longitudinal interface of this templating oligomer (the plus end) was active for
nucleation.
In initial runs wherein interactions between γ- and $\alpha\beta$-tubulin were assumed to be
of identical affinity, we observed potent nucleation even from templates that were
only two subunits wide (Supplemental Fig. 4). Because two-wide nucleators like
the γ-TuSC are much less potent than 12-14-wide oligomers like the γ-TuRC
(Oegema et al., 1999; Kollman et al., 2015), we also performed simulations with
10-fold weaker interactions between γ- and $\alpha\beta$-tubulin than for $\alpha\beta{:}\alpha\beta$ interactions
(Fig. 7)
776
777

References

Akhmanova, A., and M.O. Steinmetz. 2015. Control of microtubule organization and dynamics: two ends in the limelight. *Nat. Rev. Mol. Cell Biol.* 16:711–726. doi:10.1038/nrm4084.

Alushin, G.M., G.C. Lander, E.H. Kellogg, R. Zhang, D. Baker, and E. Nogales. 2014. High-Resolution Microtubule Structures Reveal the Structural Transitions in αβ-Tubulin upon GTP Hydrolysis. *Cell*. 157:1117–1129. doi:10.1016/j.cell.2014.03.053.

Ayaz, P., S. Munyoki, E.A. Geyer, F.-A. Piedra, E.S. Vu, R. Bromberg, Z. Otwinowski, N.V. Grishin, C.A. Brautigam, and L.M. Rice. 2014. A tethered delivery mechanism explains the catalytic action of a microtubule polymerase. *Elife*. 3:e03069. doi:10.7554/eLife.03069.

Ayaz, P., X. Ye, P. Huddleston, C.A. Brautigam, and L.M. Rice. 2012. A TOG:αβ-tubulin complex structure reveals conformation-based mechanisms for a microtubule polymerase. *Science*. 337:857–860. doi:10.1126/science.1221698.

Barlan, K., and V.I. Gelfand. 2017. Microtubule-Based Transport and the Distribution, Tethering, and Organization of Organelles. *Cold Spring Harb Perspect Biol*. 9:a025817. doi:10.1101/cshperspect.a025817.

Brouhard, G.J., and L.M. Rice. 2018. Microtubule dynamics: an interplay of biochemistry and mechanics. *Nat. Rev. Mol. Cell Biol.* 19:451–463. doi:10.1038/s41580-018-0009-y.

Buey, R.M., J.F. Díaz, and J.M. Andreu. 2006. The nucleotide switch of tubulin and microtubule assembly: a polymerization-driven structural change. *Biochemistry*. 45:5933–5938. doi:10.1021/bi060334m.

Burton, P.R., R.E. Hinkley, and G.B. Pierson. 1975. Tannic acid-stained microtubules with 12, 13, and 15 protofilaments. *J. Cell Biol.* 65:227–233.

Chaaban, S., S. Jariwala, C.-T. Hsu, S. Redemann, J.M. Kollman, T. Müller-Reichert, D. Sept, K.H. Bui, and G.J. Brouhard. 2018. The Structure and Dynamics of C. elegans Tubulin Reveals the Mechanistic Basis of Microtubule Growth. *Dev. Cell*. 47:191–204.e8. doi:10.1016/j.devcel.2018.08.023.

Chalfie, M., and J.N. Thomson. 1982. Structural and functional diversity in the neuronal microtubules of Caenorhabditis elegans. *J. Cell Biol.* 93:15–23.

Coombes, C.E., A. Yamamoto, M.R. Kenzie, D.J. Odde, and M.K. Gardner.

2013. Evolving tip structures can explain age-dependent microtubule
catastrophe. *Curr. Biol.* 23:1342–1348. doi:10.1016/j.cub.2013.05.059.

Davis, C., and K. Gull. 1983. Protofilament number in microtubules in cells of two
parasitic nematodes. *J. Parasitol.* 69:1094–1099.

Desai, A., and T.J. Mitchison. 1997. Microtubule polymerization dynamics. *Annu.*
Rev. Cell Dev. Biol. 13:83–117. doi:10.1146/annurev.cellbio.13.1.83.

Driver, J.W., E.A. Geyer, M.E. Bailey, L.M. Rice, and C.L. Asbury. 2017. Direct
measurement of conformational strain energy in protofilaments curling
outward from disassembling microtubule tips. *Elife.* 6:576.
doi:10.7554/eLife.28433.

Duellberg, C., N.I. Cade, D. Holmes, and T. Surrey. 2016. The size of the EB cap
determines instantaneous microtubule stability. *Elife.* 5:711.
doi:10.7554/eLife.13470.

Erickson, H.P., and D. Pantaloni. 1981. The role of subunit entropy in
cooperative assembly. Nucleation of microtubules and other two-dimensional
polymers. *Biophys. J.* 34:293–309. doi:10.1016/S0006-3495(81)84850-3.

Flor-Parra, I., A.B. Iglesias-Romero, and F. Chang. 2018. The XMAP215
Ortholog Alp14 Promotes Microtubule Nucleation in Fission Yeast. *Curr. Biol.*
28:1681–1691.e4. doi:10.1016/j.cub.2018.04.008.

Flyvbjerg, H., E. Jobs, and S. Leibler. 1996. Kinetics of self-assembling
microtubules: an "inverse problem" in biochemistry. *Proc. Natl. Acad. Sci.*
U.S.A. 93:5975–5979.

Fourniol, F.J., C.V. Sindelar, B. Amigues, D.K. Clare, G. Thomas, M. Perderiset,
F. Francis, A. Houdusse, and C.A. Moores. 2010. Template-free 13-
protofilament microtubule-MAP assembly visualized at 8 A resolution. *J. Cell*
Biol. 191:463–470. doi:10.1083/jcb.201007081.

Gardner, M.K., M. Zanic, C. Gell, V. Bormuth, and J. Howard. 2011.
Depolymerizing kinesins Kip3 and MCAK shape cellular microtubule
architecture by differential control of catastrophe. *Cell.* 147:1092–1103.
doi:10.1016/j.cell.2011.10.037.

Gaskin, F., C.R. Cantor, and M.L. Shelanski. 1974. Turbidimetric studies of the in
vitro assembly and disassembly of porcine neurotubules. *J. Mol. Biol.*
89:737–755.

Geyer, E.A., A. Burns, B.A. Lalonde, X. Ye, F.-A. Piedra, T.C. Huffaker, and L.M.
Rice. 2015. A mutation uncouples the tubulin conformational and GTPase

cycles, revealing allosteric control of microtubule dynamics. *Elife*. 4:3389.
doi:10.7554/eLife.10113.

Geyer, E.A., M.P. Miller, C.A. Brautigam, S. Biggins, and L.M. Rice. 2018.
Design principles of a microtubule polymerase. *Elife*. 7:604.
doi:10.7554/eLife.34574.

Grishchuk, E.L., M.I. Molodtsov, F.I. Ataullakhanov, and J.R. McIntosh. 2005.
Force production by disassembling microtubules. *Nature*. 438:384–388.
doi:10.1038/nature04132.

Gunzelmann, J., D. Rüthnick, T.-C. Lin, W. Zhang, A. Neuner, U. Jäkle, and E.
Schiebel. 2018. The microtubule polymerase Stu2 promotes oligomerization
of the γ-TuSC for cytoplasmic microtubule nucleation. *Elife*. 7:934.
doi:10.7554/eLife.39932.

Holy, T.E., and S. Leibler. 1994. Dynamic instability of microtubules as an
efficient way to search in space. *Proc. Natl. Acad. Sci. U.S.A.* 91:5682–5685.

Kollman, J.M., A. Merdes, L. Mourey, and D.A. Agard. 2011. Microtubule
nucleation by γ-tubulin complexes. *Nat. Rev. Mol. Cell Biol.* 12:709–721.
doi:10.1038/nrm3209.

Kollman, J.M., C.H. Greenberg, S. Li, M. Moritz, A. Zelter, K.K. Fong, J.-J.
Fernandez, A. Sali, J. Kilmartin, T.N. Davis, and D.A. Agard. 2015. Ring
closure activates yeast γTuRC for species-specific microtubule nucleation.
Nat. Struct. Mol. Biol. 22:132–137. doi:10.1038/nsmb.2953.

Kuchnir Fygenson, D., H. Flyvbjerg, K. Sneppen, A. Libchaber, and S. Leibler.
1995. Spontaneous nucleation of microtubules. *Phys. Rev. E*. 51:5058–5063.
doi:10.1103/PhysRevE.51.5058.

Kwiatkowska, M., K. Popłońska, D. Stepiński, and Z. Hejnowicz. 2006.
Microtubules with different diameter, protofilament number and protofilament
spacing in Ornithogalum umbellatum ovary epidermis cells. *Folia Histochem.*
Cytobiol. 44:133–138.

Macey, R., G. Oster, and T. Zahnley. 2009. Berkeley Madonna User's Guide.
University of California, Department of Molecular and Cellular Biology.

Manka, S.W., and C.A. Moores. 2018. The role of tubulin-tubulin lattice contacts
in the mechanism of microtubule dynamic instability. *Nat. Struct. Mol. Biol.*
25:607–615. doi:10.1038/s41594-018-0087-8.

Margolin, G., I.V. Gregoretti, T.M. Cickovski, C. Li, W. Shi, M.S. Alber, and H.V.
Goodson. 2012. The mechanisms of microtubule catastrophe and rescue:

implications from analysis of a dimer-scale computational model. *Mol. Biol.*
Cell. 23:642–656. doi:10.1091/mbc.E11-08-0688.

McIntosh, J.R., E. O'Toole, G. Morgan, J. Austin, E. Ulyanov, F. Ataullakhanov,
and N. Gudimchuk. 2018. Microtubules grow by the addition of bent
guanosine triphosphate tubulin to the tips of curved protofilaments. *J. Cell*
Biol. 265:jcb.201802138–2708. doi:10.1083/jcb.201802138.

Mitchison, T., and M. Kirschner. 1984. Dynamic instability of microtubule growth.
Nature. 312:237–242.

Moores, C.A., M. Perderiset, F. Francis, J. Chelly, A. Houdusse, and R.A.
Milligan. 2004. Mechanism of microtubule stabilization by doublecortin. *Mol.*
Cell. 14:833–839. doi:10.1016/j.molcel.2004.06.009.

Moritz, M., M.B. Braunfeld, J.W. Sedat, B. Alberts, and D.A. Agard. 1995.
Microtubule nucleation by gamma-tubulin-containing rings in the centrosome.
Nature. 378:638–640. doi:10.1038/378638a0.

Moritz, M., M.B. Braunfeld, V. Guénebaut, J. Heuser, and D.A. Agard. 2000.
Structure of the gamma-tubulin ring complex: a template for microtubule
nucleation. *Nature Cell Biology.* 2:365–370. doi:10.1038/35014058.

Nawrotek, A., M. Knossow, and B. Gigant. 2011. The determinants that govern
microtubule assembly from the atomic structure of GTP-tubulin. *J. Mol. Biol.*
412:35–42. doi:10.1016/j.jmb.2011.07.029.

Oegema, K., C. Wiese, O.C. Martin, R.A. Milligan, A. Iwamatsu, T.J. Mitchison,
and Y. Zheng. 1999. Characterization of two related Drosophila gamma-
tubulin complexes that differ in their ability to nucleate microtubules. *J. Cell*
Biol. 144:721–733.

Pecqueur, L., C. Duellberg, B. Dreier, Q. Jiang, C. Wang, A. Plückthun, T.
Surrey, B. Gigant, and M. Knossow. 2012. A designed ankyrin repeat protein
selected to bind to tubulin caps the microtubule plus end. *Proc. Natl. Acad.*
Sci. U.S.A. 109:12011–12016. doi:10.1073/pnas.1204129109.

Piedra, F.-A., T. Kim, E.S. Garza, E.A. Geyer, A. Burns, X. Ye, and L.M. Rice.
2016. GDP-to-GTP exchange on the microtubule end can contribute to the
frequency of catastrophe. *Mol. Biol. Cell.* 27:3515–3525.
doi:10.1091/mbc.E16-03-0199.

Popov, A.V., F. Severin, and E. Karsenti. 2002. XMAP215 is required for the
microtubule-nucleating activity of centrosomes. *Curr. Biol.* 12:1326–1330.

Prosser, S.L., and L. Pelletier. 2017. Mitotic spindle assembly in animal cells: a

fine balancing act. *Nat. Rev. Mol. Cell Biol.* 18:187–201.
doi:10.1038/nrm.2016.162.

Rice, L.M., E.A. Montabana, and D.A. Agard. 2008. The lattice as allosteric
effector: structural studies of alphabeta- and gamma-tubulin clarify the role of
GTP in microtubule assembly. *Proc. Natl. Acad. Sci. U.S.A.* 105:5378–5383.
doi:10.1073/pnas.0801155105.

Roostalu, J., and T. Surrey. 2017. Microtubule nucleation: beyond the template.
Nat. Rev. Mol. Cell Biol. 18:702–710. doi:10.1038/nrm.2017.75.

Roostalu, J., N.I. Cade, and T. Surrey. 2015. Complementary activities of TPX2
and chTOG constitute an efficient importin-regulated microtubule nucleation
module. *Nature Cell Biology*. 17:1422–1434. doi:10.1038/ncb3241.

Saito, K., and K. Hama. 1982. Structural diversity of microtubules in the
supporting cells of the sensory epithelium of guinea pig organ of Corti. *J
Electron Microsc (Tokyo)*. 31:278–281.

Sept, D., and J.A. McCammon. 2001. Thermodynamics and Kinetics of Actin
Filament Nucleation. *Biophys. J.* 81:667–674. doi:10.1016/S0006-
3495(01)75731-1.

Thawani, A., R.S. Kadzik, and S. Petry. 2018. XMAP215 is a microtubule
nucleation factor that functions synergistically with the γ-tubulin ring complex.
Nature Cell Biology. 20:575–585. doi:10.1038/s41556-018-0091-6.

Tobacman, L.S., and E.D. Korn. 1983. The kinetics of actin nucleation and
polymerization. *J. Biol. Chem.* 258:3207–3214.

Tucker, J.B., C.C. Paton, G.P. Richardson, M.M. Mogensen, and I.J. Russell.
1992. A cell surface-associated centrosomal layer of microtubule-organizing
material in the inner pillar cell of the mouse cochlea. *J. Cell. Sci.* 102 (Pt
2):215–226.

VanBuren, V., D.J. Odde, and L. Cassimeris. 2002. Estimates of lateral and
longitudinal bond energies within the microtubule lattice. *Proc. Natl. Acad.
Sci. U.S.A.* 99:6035–6040. doi:10.1073/pnas.092504999.

VanBuren, V., L. Cassimeris, and D.J. Odde. 2005. Mechanochemical model of
microtubule structure and self-assembly kinetics. *Biophys. J.* 89:2911–2926.
doi:10.1529/biophysj.105.060913.

Voter, W.A., and H.P. Erickson. 1984. The kinetics of microtubule assembly.
Evidence for a two-stage nucleation mechanism. *J. Biol. Chem.* 259:10430–
10438.

Wang, H.-W., and E. Nogales. 2005. Nucleotide-dependent bending flexibility of
tubulin regulates microtubule assembly. *Nature*. 435:911–915.
doi:10.1038/nature03606.

Wieczorek, M., S. Bechstedt, S. Chaaban, and G.J. Brouhard. 2015. Microtubule-
associated proteins control the kinetics of microtubule nucleation. *Nature Cell
Biology*. 17:907–916. doi:10.1038/ncb3188.

Zakharov, P., N. Gudimchuk, V. Voevodin, A. Tikhonravov, F.I. Ataullakhanov,
and E.L. Grishchuk. 2015. Molecular and Mechanical Causes of Microtubule
Catastrophe and Aging. *Biophys. J.* 109:2574–2591.
doi:10.1016/j.bpj.2015.10.048.

Zhang, R., G.M. Alushin, A. Brown, and E. Nogales. 2015. Mechanistic Origin of
Microtubule Dynamic Instability and Its Modulation by EB Proteins. *Cell*.
162:849–859. doi:10.1016/j.cell.2015.07.012.

Zhang, R., J. Roostalu, T. Surrey, and E. Nogales. 2017. Structural insight into
TPX2-stimulated microtubule assembly. *Elife*. 6:1518.
doi:10.7554/eLife.30959.

Zheng, Y., M.L. Wong, B. Alberts, and T. Mitchison. 1995. Nucleation of
microtubule assembly by a gamma-tubulin-containing ring complex. *Nature*.
378:578–583. doi:10.1038/378578a0.

973
974
975

Figure 1 Spontaneous microtubule nucleation: assay and scaling behavior.
A Time course of spontaneous microtubule assembly measured by turbidity
(absorbance at A350 nm) at ab-tubulin concentrations of (10, 11, 12, 13, 14, and
15 μM).
B The same data as in **A** but with the turbidity values of each curve normalized
by the corresponding plateau (steady-state) value achieved. Inset shows the
same data plotted on logarithmic axes. The horizontal red line indicates 10%
assembly.
C Data from **B** but with the time values for each curve normalized by the
corresponding $T_{0.1}$ (time taken to reach 10% assembly). Inset shows how $T_{0.1}$
depends on concentration. The collapse of all curves after this scaling procedure
indicates that a common mechanism operates over the entire concentration
range (Flyvbjerg et al., 1996).
D Cartoon of a mechanism to explain spontaneous microtubule nucleation
previously postulated based on this kind of scaling analysis. The basic idea is
that nucleation proceeds through a small number of discrete intermediate
species: for the scaling analysis of the data presented in A the sequence of
oligomers would be 8-mer, 12-mer, 16-mer (in other words, two intermediate
species precede the 16-mer nucleus).
960

Figure 2

Figure 2 An explicit biochemical model for early steps in spontaneous
microtubule nucleation.
A Cartoon of the model. Individual ab-tubulin heterodimers (ovals) can associate
longitudinally (blue arrows) or laterally (orange arrows). As oligomers grow in
size, they can present sites where an associating tubulin would simultaneously
make longitudinal and lateral interactions ("corner"; bold black arrows). The
resulting collection of species and reactions was implemented in
BerkeleyMadonna (Macey et al., 2009) for simulations. Species up to 6-mers are
shown, but the model includes species up to 12-mers. Rapid growth of the
number of oligomers and reactions with size made it impractical to consider even
larger oligomers.
B (left) Plot showing that 'rectangular' species dominate the simulation. The four
dominant species are shown, with values in parentheses indicating the height
and width for the oligomers. (right) The same data plotted on a logarithmic
concentration scale. Other closed oligomers are much less common. Non-
rectangular oligomers are not shown, but are also much rarer than the dominant
oligomers shown.
C A simplified 'layer growth' model that only considers rectangular oligomers.
Arrows indicate longitudinal (blue) or lateral (orange) layer growth. The resulting
collection of species and reactions was implemented in BerkeleyMadonna for
simulations (see Methods)
D Cartoon of a 'layer' addition reaction with associated rate constants. The
equilibrium constant and rate constant for layer addition can be defined in terms
of underlying longitudinal, lateral, and corner affinities and a single on rate
constant. An effective equilibrium constant and an approximation of the effective
forward rate constant for this layer addition are shown. See Methods for
derivations.
E Similar to **B** but showing results from simulations of the simplified layer model.
This much simpler model with many fewer species and reactions quantitatively
reproduces results from the explicit model.
F (left) Plot of the species count for all possible oligomers (black) and for the
subset of closed oligomers (red). The extrapolation to a 40-mer emphasizes the
magnitude of the simplification. (right) Similar plot but for the number of
biochemical reactions.
995

A

10 µM

K_d^{corn}: 17 nM
K_d^{long}: 13 mM
K_d^{lat}: 130 mM

13 µM

K_d^{corn}: 52 nM
K_d^{long}: 7 mM
K_d^{lat}: 83 mM

15 µM

K_d^{corn}: 68 nM
K_d^{long}: 5 mM
K_d^{lat}: 90 mM

B

Global fit

C

	Concentration-dependence of $T_{0.1}$
Exp.	3.8
Sim.	3.1

D

rate ~ $[\alpha\beta]^2/K_d^{long}$

rate ~ $[\alpha\beta]^2/K_d^{lat}$

rate ~ $[\alpha\beta]^2/K_d^{long}$

Missing 'dimer' pathway

E

If $K_d^{long} < K_d^{lat}$, the 'dimer pathway' should be majority contributor to lateral growth

K_d^{Long}/K_d^{Lat}

Figure 3 Expansion of the layer model to arbitrarily large oligomers for
simulating spontaneous microtubule nucleation. Oligomers are allowed to
become arbitrarily tall. We assumed that when oligomers reached a width of 13,
they 'closed' into a tube and elongated by monomer addition.
A Fits of the model (black dotted line) to individual assembly curves (solid grey
line) at different concentrations. Fits to individual curves are good. Inset text
shows the resulting longitudinal, lateral, and 'corner' (simultaneous longitudinal
and lateral interactions) affinities, which differ somewhat from run to run.
B Global fit of the model (black dashed lines) to the full set of assembly data
(grey lines). As evidenced by the poor fit at low concentration and by the
concentration-dependence of the $T_{0.1}$ (see **C**), the model underestimates the
concentration-dependence of the progress curves.
C Tabulation of the concentration-dependence of $T_{0.1}$ for the experimental data
and for the model fit.
D (left) Cartoon illustrating that longitudinal dimers of $\alpha\beta$-tubulin could contribute
to lateral growth. The associated forward rate constant for this case is also
shown.
E The relative contribution of 'monomer' and 'dimer' initiation pathways can be
estimated as indicated based on the affinities of longitudinal and lateral
interactions. Plot predicting the fractional contribution of a 'dimer' pathway for
lateral growth in terms of the relative strength of longitudinal and lateral
interactions. The red star indicates the ratio obtained from the fit shown in **B**.
Those parameters predict that the 'dimer' pathway should dominate the
assembly, but it is not accounted for in our model.
1020

A

10 μM

K_d^{corn}: 5 nM
K_d^{long}: 6.4 mM
K_d^{late}: 32.4 M

12 μM

K_d^{corn}: 6 nM
K_d^{long}: 5.6 mM
K_d^{late}: 27.6 M

15 μM

K_d^{corn}: 6 nM
K_d^{long}: 5.5 mM
K_d^{late}: 24 M

B

C

	Concentration-dependence of $T_{0.1}$
Exp.	**3.8**
Sim.	**3.1**

Figure 4 Incorporating a dimer pathway for lateral growth into the model
A Fits of the new dimer model (black dotted line) to individual assembly curves
(solid grey line) at different concentrations. Fits to individual curves are worse
than for the monomer only model. Inset text shows the resulting longitudinal,
lateral, and 'corner' (simultaneous longitudinal and lateral interactions) affinities;
these differ mainly in the magnitude of the lateral affinity, which is much weaker
than for the monomer only model.
B Global fit of the model (black dashed lines) to the full set of assembly data
(grey lines). The fit is very poor, with the model predicting too fast assembly at all
concentrations.
C Tabulation of the concentration-dependence of $T_{0.1}$ for the experimental data
and for the model fit. Consistent with the poor fit, the dimer model fails to capture
the correct concentration dependence. The dimer model thus also
underestimates the concentration-dependence of the progress curves.
1035
1036

A

Width = 2
Stronger Lateral
interaction

Width > 2
Weaker Lateral
Interaction

B

C

P = 8

D

	Concentration-dependence of $T_{0.1}$
Exp.	3.8
Sim.	3.5

Figure 5 A 'penalty' that mimics the cost of $\alpha\beta$-tubulin straightening markedly
improves the performance of the dimer model
A Cartoons illustrating $\alpha\beta$-tubulin straightening, which is thought to occur
progressively with width (the number of lateral neighbors). To model this in a very
simple way, we introduced a penalty to make lateral interactions weaker for the
3^{rd} and successive lateral interactions.
B Performance of the 'dimer with penalty' model as a function of penalty
strength. The improved goodness of fit from the global optimization is not very
sensitive to the magnitude of the penalty, as long as the penalty is ~8 or greater.
C Illustration of the model fit to the experimental data for a penalty value of 8.
1047
1048

A

B

Figure 6 Two complementary views of the predicted pathway for spontaneous
microtubule assembly
A Summed concentration for all oligomers of a given width, plotted as a function
of time. Oligomers with width=2 dominate. Wider oligomers become
progressively rarer, and peak at increasingly late times. The vertically striped light
grey curve behind the others shows the extent of total assembly (right y-axis).
B Distribution of oligomer heights as a function of their width. The most populated
height is plotted as a filled circle for each width (different colors), at the time that
width peaks. Other height oligomers at a given width are plotted as open circles,
with the radius reflecting their concentration relative to the peak species. The
time to transition from one width to the next decreases as oligomers get wider
and taller, and the spread of oligomer heights increases as oligomers get wider.
1061
1062

Figure 7 Implications for templating microtubule assembly
A Cartoon illustrating an intermediate stage of spontaneous nucleation. 2x2
oligomers of $\alpha\beta$-tubulin (left) assemble as described into rectangular
intermediates that tend to be taller than they are wide (middle) because of the
barriers to lateral growth. Arrows indicate multiple layer additions, each one
associated with a delay.
B When $\alpha\beta$-tubulin adds to a γ-TuRC-like template, cartooned here as a row of
blue ovals, the barrier to lateral growth is eliminated because 'corner' sites form
at the border of the $\alpha\beta$-tubulin oligomer. This accelerates the accretion process
(bolder arrows indicating faster transitions) and the resulting oligomers now tend
to be wider than they are tall
C Simulated nucleation curves in the presence of γ-tubulin templates of different
width assuming $\gamma{:}\alpha\beta$ interactions are weaker than $\alpha\beta{:}\alpha\beta$ interactions. The total
concentration of templating monomers is held constant. Templates that are only
two subunits wide (bright green) show a modest effect, but wider oligomers are
increasingly potent.
D Quantification of the data from C. Wider templates are more potent than
narrower ones. Templating the earliest step in nucleation (from 2-wide to 3-wide)
yields the most dramatic decrease in $T_{0.1}$.
E Relative extent of assembly at an early time (here, 175 s) plotted as a function
of template width and using data from C. Relative assembly at a given time
represents the fraction assembled for a given template divided by the fraction
assembled in the untemplated control. The changes in nucleation kinetics can
result in large changes in the overall degree of microtubule assembly at early
times.
1088

Illustration of the meaning of 'non-redundant' oligomers

There are 6 distinct αβ-tubulin tetramers with 2 longitudinal bonds and 1 lateral bond

These 4 species present :
   1 corner (c),
   3 longitudinal (L), and
   5 lateral (l) binding sites

These 2 species present :
   2 corner (c),
   2 longitudinal (L), and
   4 lateral (l) binding sites

Population-weighted average:
   1.33 corner (c),
   2.67 longitudinal (L), and
   4.67 lateral (l) binding sites

Supplemental Figure 1 Illustration of the concept and implementation of 'non-
redundant' oligomers. Representative configurations of an $\alpha\beta$-tubulin tetramer
containing two longitudinal and one lateral bond are indicated. These are
indistinguishable from the standpoint of binding energy, but differ in terms of the
number and type of binding sites they present. Our algorithm simplifies these six
species into one and uses population weighted rate constants to capture the
'average' number and type of binding sites presented.
1096
1097

Supplemental Figure 2 In the explicit model, relatively tight corner affinity is
required for appreciable assembly into larger oligomers.
A Plots of the concentration of 'rectangular' oligomers vs time, for different
values of the corner affinity (color coding is given in panel C and is identical to
that used in Fig. 2B). Note different y-axis ranges for different plots. At corner
affinity of 400 nM and weaker, assembly into species larger than a tetramer of
$\alpha\beta$-tubulin (dark blue curve) is negligeable.
B As in A, but using a log-scale y-axis to show the dramatic fall-off in the
concentration of larger species.
C Legend showing the correspondence between color and oligomer dimensions.
1108
1109

Illustration of the dimer pathway

Initiation by lateral addition
of a longitudinal dimer

'Zipping' of the layer by subsequent
addition of monomers

Supplemental Figure 3 Illustration of the dimer pathway.
Cartoon sequence of steps for adding a new lateral layer. In the dimer pathway, a
new lateral layer can be initiated by association of a pre-formed longitudinal
dimer (left). Because the concentration of these longitudinal dimers of $\alpha\beta$-tubulin
is much less than that of $\alpha\beta$-tubulin 'monomers', subsequent 'zipping' of the layer
occurs by monomer addition. See Methods for a derivation of the equilibrium
binding constant for the initiating assocition.
1117
1118

Supplemental Figure 4 Simulations of templated microtubule assembly.
Simulated nucleation curves in the presence of γ-tubulin templates of different
width assuming $\gamma{:}\alpha\beta$ interactions are the same strength as $\alpha\beta{:}\alpha\beta$ interactions.
Compare to Fig. 7. The total concentration of templating monomers is held
constant. Templates that are only two subunits wide (bright green) show very
stronger nucleation effect than when we assumed weaker $\gamma{:}\alpha\beta$ interactions
(compare to Fig. 7).
1126
1127