Soil type and interannual variation shape unique fungal pathogen communities on nearby

populations of a California native bunchgrass

Johannah E. Farner[1], Erin R. Spear[1,2], Erin A. Mordecai[1]

[1]Biology Department, Stanford University, Stanford, California 94305 USA; [2] Biology

Department, Regis University, Denver, Colorado 80221 USA

Author for correspondence:

Johannah Farner

E-mail: jfarner@stanford.edu

Phone: 1 (805) 317 – 5432

Abstract

The role of infectious disease in regulating host populations is increasingly recognized, but how environmental conditions affect pathogen communities and infection levels remains poorly understood. Over three years, we compared foliar disease burden, fungal pathogen community composition, and plant tissue chemistry in the perennial bunchgrass *Stipa pulchra* on nearby serpentine and nonserpentine soils that have distinct chemical properties and plant communities. We found that serpentine and nonserpentine *S. pulchra* experienced consistent, low disease pressure associated with distinct communities of pathogens, that interannual fungal species turnover differed with soil type, and that plant chemistry reflected soil chemistry. The results indicate that this species experiences minimal foliar disease associated with diverse fungal pathogen communities that are structured by environmental conditions. Highly variable conditions associated with fungal species turnover may create a refuge from disease outbreaks for *S. pulchra*, contributing to the low disease burden observed on this and other Mediterranean grassland species.

Key words: California grasslands, leaf spot pathogen, plant fungal pathogens, serpentine soil, Stipa pulchra

Introduction

How environmental heterogeneity affects species interactions, and ultimately population growth, remains a fundamental ecological question. Infectious disease, which arises from interactions between hosts, parasites, and the environment, is particularly likely to vary across populations in differing environments. Both biotic factors, such as the phylogenetic structure of the host community, and abiotic factors, such as elevation, have been associated with the presence of specific diseases and the severity of their impacts on their hosts (Abbate and Antonovics 2014; Parker *et al.* 2015). Additionally, mounting evidence suggests that specific characteristics of pathogen community structures, including the identities and abundances of the diseases present in a host population, mediate the outcomes of host – pathogen interactions (Seabloom *et al.* 2015; Borer *et al.* 2016). Together, these observations highlight the importance of understanding interactions between environmental variation, pathogen community composition, and pathogen pressure on host populations to assess the potential for disease to regulate host population dynamics. For plant hosts, such assessments are important both in conservation and agricultural contexts because identifying populations at high risk for severe disease outbreaks and applying preventative measures may prevent local species extinctions and improve crop yields, respectively. While studies have frequently explored the interactions of single host – pathogen pairs under differing abiotic conditions (e.g., Abbate and Antonovics 2014) or the effect of varying pathogen community composition on disease burden (e.g., Telfer *et al.* 2010), relationships between abiotic variables and pathogen community structures remain largely unexplored in natural systems (but see Spear 2017).

Fungal pathogens are particularly important to study in this context because they cause the majority of known plant diseases, have an increasing rate of emergence, and pose a substantial threat to plant health and population persistence in natural and agricultural settings (Agrios 2005; Fisher *et al.* 2012). Dramatic diebacks of North American chestnut trees and oaks caused by fungal pathogens (Anagnostakis 1987; Rizzo and Garbelotto 2003) and economically devastating crop losses caused by fungal rusts shared among wild and domesticated plants (Loehrer and Schaffrath 2011; Nazareno *et al*. 2018) demonstrate this threat. These examples indicate that understanding the communities of fungal pathogens that infect wild plants is important to developing effective protection strategies for both natural biodiversity and human food supplies. However, we lack basic information about the identities, diversity, and effects of

the fungal pathogens that infect most plant species (Burdon and Laine 2019). Therefore, there is

a pressing need for systematic surveys of these pathogen communities and their variation across

environmental conditions in natural systems.

California grasslands are an ideal setting in which to study interactions between

environmental variation, fungal pathogen community composition, and disease burden because

these plant communities are widespread, exist across a variety of environmental gradients, and

include distinct plant community subtypes (Eviner 2016). Here, we take advantage of the natural

occurrence of the native perennial bunchgrass *Stipa pulchra* on distinct soil types—serpentine,

hosting diverse, native-dominated grassland communities, and a nonserpentine greenstone

(hereafter, nonserpentine) dominated by exotic plants—to understand the extent to which soil

type and time influence the landscape of disease encountered by a single plant species, including

foliar fungal pathogen community composition and disease severity.

Differences in plant community composition and soil chemistry suggest that *S. pulchra*

populations growing in distinct soil types are likely to host distinct fungal pathogen communities

and/or experience differing disease burdens, particularly if the phenotype of *S. pulchra* is

influenced by the soil in which it grows. *S. pulchra* naturally occurs in both nonserpentine soils

and in relatively rare serpentine soils that host grassland plant communities. These two soil types

differ in their chemistry and host distinct plant communities, resulting in a suite of environmental

differences that may influence fungal pathogen infection, reproduction, and persistence (Harrison

and Viers 2007; Burdon and Laine 2019). In contrast to the majority of California grasslands,

which are heavily invaded by exotic species, serpentine grasslands have a diverse array of native

species adapted to chemically and physically harsh soils that are low in essential plant nutrients

(N, P, K, Ca), high in heavy metals that are potentially toxic to plants (Co, Ni, Cr, Fe), and rocky

and shallow, with poor moisture retention (Harrison and Viers 2007; Oze *et al.* 2008; Eviner,

2016).

Soil properties determine a variety of factors likely to influence the occurrence and

outcomes of plant fungal infections, including soil moisture levels and plant chemistry and

physiology (Cunningham *et al.* 1999; McElrone *et al.* 2005; Smith 2007). Spores of different

fungal species vary in both their ability to survive in dry and moist environments and their

tolerance of heavy metals (Coley-Smith and Cooke 1971; Iram *et al.* 2009). Moreover, plants of

the same species grown in drier and lower-nutrient soils can have thicker epidermises and higher

concentrations of phenolic compounds, many of which have antifungal properties (Osbourne

1996; Cunningham *et al.* 1999). Finally, soil chemistry may influence pathogen infection either

through hyperaccumulation of heavy metals, which can bolster chemical defenses against natural

enemies, or by limiting common mechanisms of plant defense (Springer *et al.* 2006; Rascio and

Navari-Izzo 2011).

In this study, we ask: (1) how does disease burden differ on plants growing in different

soil types? (2) how does the species composition of fungal pathogen communities differ on

plants growing in different soil types? and (3) by what mechanisms—including host community

composition and plant tissue chemistry—might soil type affect disease burden and pathogen

composition? We hypothesize that both disease burden and pathogen species composition differ

on *S. pulchra* individuals growing in serpentine versus nonserpentine soils, either because of

differences in the composition of the surrounding plant community, differences in plant

physiological responses, or both. We further hypothesize that the soil type *S. pulchra* grows in

alters foliar tissue chemistry in ways that may affect disease burden, with higher heavy metal

content and/or lower C, N, and P content in serpentine plants. If so, these foliar chemistry

differences may translate into either (a) higher disease burden in serpentine plants relative to

nonserpentine plants, facilitated by negative impacts of limited plant resource availability on

plant health (as in Springer *et al.,* 2006), or (b) lower disease burden in serpentine plants due to

heightened chemical and/or physiological plant defenses in response to serpentine soil properties.

In addition to contributing novel information about previously uncharacterized plant fungal

pathogen communities in California grasslands, the results of this study will aid in assessing

disease risk for plant populations in natural and agricultural contexts by providing information

about the spatial and temporal scales over which soil type and other environmental factors can

affect the spread of outbreaks.

Materials and Methods

All surveys and sample collection took place at Jasper Ridge Biological Preserve (JRBP),

a 485-hectare site in San Mateo County, CA managed by Stanford University (37.4° N, 122.2°

W). JRBP has a Mediterranean climate with cool winters (mean 9.2°C), warm summers (mean

20.1°C), and annually variable precipitation that predominantly occurs in the winter and spring

(averaging 622.5 mm per year) (Ackerly *et al.* 2002). The growing season occurs during the

rainy season and plants senesce in early summer. This study was conducted during the growing

seasons from mid-April to mid-May in 2015, 2016, and 2017. There was highly variable annual

precipitation across the three years (491 mm, 604 mm, and 985 mm in 2015, 2016, and 2017,

respectively; Weather Underground).

Substantial populations of *S. pulchra* occur in both serpentine and nonserpentine

grasslands at JRBP (McNaughton 1968). The serpentine plant communities at JRBP are more

diverse than the nearby nonserpentine plant communities and include many native grasses and

forbs, in contrast to the invasive-dominated nonserpentine grasslands (McNaughton 1968;

Huenneke *et al.* 1990; Field *et al.* 1996). Plant species commonly found in serpentine grasslands

at JRBP include *Stipa pulchra*, *Elymus glaucus*, *Elymus multisetus, Eschscholzia californica*, and

the invasive grass *Bromus hordeaceus* (McNaughton 1968; Hobbs and Mooney 1985). The

nonnative annual species that dominate the nonserpentine grasslands but are absent from

serpentine grasslands include *Avena barbata, Avena fatua*, *Bromus diandrus,* and *Erodium*

botrys (McNaughton 1968; Hobbs and Mooney 1985; Field *et al.* 1996).

Quantification of disease

To assess plant disease burden, we quantified the percentage of living leaf area exhibiting

symptoms of fungal disease in serpentine and nonserpentine populations of *S. pulchra*. We

selected three serpentine grassland sites and three nonserpentine grassland sites such that each

serpentine site was paired with a nearby (~170m away) nonserpentine site (Fig. 1a). Individual

sites within each soil type were >300m apart and the maximum distance between any two sites

was 1600m (Fig. 1a). Each year, we randomly placed four 5-meter transects at each 40m x 40m

site and assessed the infection status of the *S. pulchra* individual nearest each meter mark,

resulting in five individual plants per transect surveyed each year (N = 60 plants per soil type per

year). A single surveyor visually estimated the percentage of living foliar tissue damaged by

pathogens for six arbitrarily-selected leaves per plant. We used printed field reference guides

showing known amounts of damage on leaves to standardize estimates. Damage was indicated

by spots and lesions which ranged in color from dark brown to pale yellow (Fig. 1b). For damage

estimates of 5% or greater, values were determined to the nearest 5%. Lower damage levels were

described as either 1% or 0.1%. All survey sites were located along the ridgetop at JRBP, where

serpentine and nonserpentine (greenstone) soils are present in discrete adjacent bands (Fig. 1a)

(Oze *et al.*,2004). The relative flatness of the ridgetop ensured that all survey sites were at

roughly the same elevation and had similar slopes, aspects, and water availability (Oze *et al.*

2008).

We used analysis of variance (ANOVA) to test for effects of soil type, year, soil type x

year, and site on log-transformed mean percent disease damage by transect in serpentine and

nonserpentine *S. pulchra* (N = 36 serpentine and 36 nonserpentine transects). When ANOVA

results were statistically significant, we used t-tests for every possible pairwise combination,

with Bonferroni adjustments of p-values to account for multiple comparisons, to identify the

interactions contributing to the significant result.

Isolation and molecular identification of foliar fungal pathogens

To identify the fungal foliar pathogens infecting serpentine and nonserpentine *S. pulchra,*

we harvested one segment of symptomatic leaf tissue for culturing and identification of the

putatively causal fungal pathogen(s) from each surveyed individual with disease. We excised,

surface sterilized, and plated a 2mm^2 square of tissue from each sample at the leading edge of

infection, to improve the chances of isolating the causal fungal pathogen, on malt extract agar

with 2% chloramphenicol to prevent bacterial growth. We observed the plates for growth for

eight weeks and isolated morphologically distinct hyphae into axenic culture. Six segments of

surface sterilized, asymptomatic tissue were plated as a control and showed no growth. We

extracted and sequenced genomic DNA from the ITS region for each isolate. Fungal isolation

and sequencing methods with ITS1F and ITS4 primers followed Spear and Mordecai 2018.

It is possible that the growing medium used may have led to selective isolation and

sequencing of fungi other than those causing symptoms. Our use of small segments of leaf tissue

from the advancing margin of disease was intended to prevent this outcome under the

assumption that symptom-causing organisms should be dominant in visibly diseased tissue and

particularly at its expanding margin, and our fungal cultures were observed growing directly out

of lesions. Inoculation trials on a variety of host species described in Spear and Mordecai 2018

confirmed pathogenicity for 27% of tested fungi isolated from this grassland system under this

protocol.

We estimated taxonomic placement of fungal isolates by grouping sequences into

operational taxonomic units with 97% sequence similarity (OTUs), a proxy for species

delineation based on the range of intraspecific ITS divergence (O'Brien *et al.* 2005), and then

comparing assembled DNA sequences to the entries in the UNITE fungal database. To prepare

sequences for taxonomic placement, we first trimmed forward and reverse Sanger sequencing

reads to a maximum error rate of 5% and then automatically assembled bidirectional reads into

consensus sequences with a minimum of 20% overlap and 85% sequence similarity in Geneious

(Kearse *et al.* 2012), discarding consensus sequences less than 450 bp in length and/or less than

60% in read quality. We grouped the remaining internal transcribed spacer (ITS) sequences into

OTUs using USEARCH 10.0.240 (Edgar 2010) and estimated the taxonomic placement of each

OTU in mothur v.1.40.5 (http://mothur.org/; Schloss 2009) using the naïve Bayes classification

approach (classify.seqs with options: method='wang', cutoff=0, iters=200). We used a

bootstrapping confidence score of at least 80% to assign a species name and at least 60% to

assign taxonomy at higher ranks.

Analyses of fungal community composition

To characterize foliar fungal community composition in serpentine and nonserpentine *S.*

pulchra populations, we compared the diversity and assessed the similarity of communities of

fungi cultured from the diseased tissue of surveyed plants. To describe fungal community

diversity, we calculated observed taxa richness and Fisher's alpha, a widely used measure of

species richness that is robust to uneven sample sizes (Fisher *et al.* 1943; Magurran 2013);

generated taxa accumulation curves to understand sampling efficacy (Oksanen *et al.* 2016); and

counted the number of fungal isolates in each genus, family, order, and class from each grassland

type. We compared fungal community composition across grassland types and across years

visually, using non-metric multidimensional scaling (NMDS), and statistically, using

permutational multivariate analyses of variance (PERMANOVA) with pairwise Adonis

functions (Anderson 2001; Arbizu 2016; Oksanen *et al.* 2016). To assess spatial turnover in

fungal communities, we included a site distance factor in the statistical analyses, with

comparisons between sites <600m apart categorized as "near" and comparisons between sites

>600m apart categorized as "far" (Fig. 1a). To account for low culturing success size in some

transects, we considered the combined isolates from two nearby transects at each site in each

year (N = 18 serpentine and 18 nonserpentine communities, generated from 36 total transects per

soil type) to be distinct communities for NMDS visualization.

For ANOVA and PERMANOVA analyses we considered the combined isolates from

each site in each year to be distinct communities (N = 9 serpentine and 9 nonerspentine

communities). When PERMANOVA results were statistically significant, we used pairwise

PERMANOVAs with Bonferroni adjustments to p-values to clarify which fungal communities

were significantly different from one another (Arbizu 2016). For the NMDS and PERMANOVA

analyses, we used the function vegdist with the abundance-based Chao method, accounting for

unsampled species, to create a matrix of pairwise community dissimilarities (Chao *et al.* 2016;

Oksanen *et al.* 2016). We also used the Morisita-Horn index, which is independent of sample

size and diversity, to make pairwise comparisons of similarity within and between the serpentine

and nonserpentine fungal communities from each year (N = 265 isolates, 200 bootstrap

replicates) (Wolda 1981; Jost *et al.* 2011). Finally, we used ANOVA to test for effects of soil

type, year, site distance, year x soil type, and year x site distance interactions on fungal diversity

(Fisher's alpha) and richness at the site level.

To perform these fungal community analyses, we used the packages BiodiversityR (Kindt

and Coe 2005), bipartite (Dormann *et al.* 2008), fossil (Vavrek 2011), rich (Rossi 2011), SpadeR

(Chao *et al.* 2016), and vegan (Oksanen *et al.* 2016) with custom commands (Gardener 2014) in

R version 3.4.1.

Chemical analyses of leaf tissue

We analyzed C, N, Ca, Cr, Mg, Ni, and P content of the foliar tissue of *S. pulchra* grown

in serpentine and nonserpentine soils *in situ* and in pots (as part of a separate field experiment;

see Methods S1). In 2016, we collected the youngest fully expanded and entirely asymptomatic

leaf of each individual surveyed in the field (N = 60 serpentine, 58 nonserpentine). We dried the

samples at room temperature for three months, then at 60°C for 48 hours, ground them to a

powder, and then dried them again at 60°C for 24 hours. We measured C and N content of these

samples with a Carlo Erba NA-1500 elemental analyzer using atropine standards.

To compare Ca, Cr, Mg, Ni, and P content, which requires more tissue biomass than

could be collected from plants in the field, we used a set of plants grown in pots in the

greenhouse and then moved to JRBP for a separate experiment (Methods S1). In total, we grew

480 plants in serpentine soil sourced from our survey sites at JRBP and 480 plants in

nonserpentine greenhouse soil (PROMIX HP Mycorrhizae TM High Porosity Professional

Growing Medium, Premier Horticulture Inc.). Interactions with soil microbes can affect plant

defenses against consumers (Pineda *et al.* 2010). Therefore, to ensure that serpentine- and

nonserpentine-grown plants encountered similar soil biota, we triple autoclave-sterilized

serpentine soil (40-minute cycles at 120°C) and then immediately inoculated it with greenhouse

soil in a ratio of 93:7 by weight (serpentine soil treatment; Heinze *et al.* 2015) and mixed

sterilized serpentine soil into greenhouse soil in the same ratio (nonserpentine soil treatment, to

account for possibility of microorganisms surviving autoclaving).

Both our choice of growing medium and the effects of autoclave sterilization on physical

and chemical soil properties may have affected our results. Our methods should result in a soil

environment in which naturally occurring mutualists of *S. pulchra* found in serpentine soils are

replaced with the biota found in the commercial house soil we used. However, artificial

modification of plant-soil microbe interactions may alter plant nutrient uptake (Khan 2005). To

avoid such effects, we might have used non-sterilized serpentine and nonserpentine soils from

our field sites. However, the volume of soil required for the expanded experiment these plants

were a part of, along with the sensitive nature of our study area, precluded this possibility.

Results from studies that show effects of autoclaving on extractable levels of the

elements we measured suggest that this method is more likely to increase than decrease the

similarity of nutrient availability in serpentine and nonserpentine soils. For example, autoclaving

has been shown to increase extractable Ca:Mg and decrease extractable Ni (Tadros 1957; Wolf *et*

al. 1989; Abou-Shanab *et al.* 2003). Therefore, any significant differences observed between the

foliar chemistry of *S. pulchra* grown in autoclaved serpentine and unsterilized nonserpentine

soils that indicate plant tissue chemistry matching soil chemistry should not be due to our use of

autoclave sterilization.

Elemental content of these plants was measured via inductively coupled plasma optical

emission spectrometry (ICP-OES). Because the minimum amount of leaf tissue for which some

heavy metals can be detected via ICP-OES is 0.5g and most of our *S. pulchra* had a total

aboveground biomass of less than 0.5g, each measurement represents the combined elemental

content of equal weights of oven-dried foliar tissue from two plants. Plants were randomly

selected for analysis from among all individuals with an aboveground biomass of at least 0.29g.

All asymptomatic leaves were removed from each plant's base and processed as described

above. A 0.5g portion of each pooled sample was digested in 10ml of 70% trace metal quality

nitric acid with microwave assistance. Digested samples were diluted to 16.7% acid

concentration with DI water and the concentrations of Ca, Cr, Mg, Ni, and P were measured in

parts per billion by a Thermo Scientific iCAP 6300 spectrometer (Thermo Scientific, Waltham,

MA, USA) (Cantarelli *et al.* 2010).

To assess the potential for soil chemistry to mediate plant-pathogen interactions, we

related soil type to leaf tissue chemistry of *in situ* and potted *S. pulchra*, and chemistry of *in situ*

plants to disease burden. We compared the mean mol/kg dried weight of leaf tissue for Ca, Cr,

Mg, and Ni for potted serpentine and nonserpentine plants with Wilcoxon rank sum tests. We

also compared mean C : N and Ca : Mg ratios that are indicative of plant health (Ciompi *et al.*

1996; Brady *et al.* 2005). We used Pearson's product-moment correlation to test for significant

monotonic relationships between the log-transformed mean amount of diseased tissue observed

on individual plants in the field and their foliar C and N content.

All statistical analyses were done in R version 3.4.1. In addition to the R packages

specified in the fungal community analyses methods, we used the packages cowplot (Wilke

2019), data.table (Dowle and Srinivasan 2019), RColorBrewer (Neuwirth 2014), reshape2

(Wickham 2007), tidyverse (Wickham 2017), and xtable (Dahl *et al*. 2019) to manipulate, plot,

and export data and results.

Results

Pathogen damage

Pathogen damage was similarly ubiquitous yet low-burden in serpentine and

nonserpentine soils. Every plant surveyed exhibited evidence of foliar fungus-caused disease.

The mean percentage of diseased leaf area observed across all years was 1.63 for serpentine and

1.66 for nonserpentine *S. pulchra,* respectively (Fig. 1c), and the mean proportion of surveyed

leaves with pathogen damage was 0.83 for serpentine and 0.85 for nonserpentine plants.

ANOVA results indicated a significant effect of year on log-transformed mean percent damage

(df = 2, SS = 9.41, F-value = 5.375, p-value = 7.04×10^{-3}) (Table 1). The only significant

Bonferroni-adjusted p-value for t-tests for every possible pairwise combination of years was the

comparison between 2015 and 2017 (2015 mean percent damage = 1.05, 2017 = 1.94, adjusted

p-value = 5.013 x 10^{-3}).

Fungal pathogen community

We isolated 267 unique fungal isolates from 258 plated symptomatic tissue pieces with

fungal growth (144 nonserpentine and 114 serpentine tissue pieces from 36 nonserpentine and 34

serpentine transects, out of 360 total tissue pieces). Of the 267 unique fungal isolates, we

successfully sequenced 256 isolates. The sequenced isolates clustered into 30 operational

taxonomic units (OTUs) based on 97% sequence similarity, representing 23 genera, 15 families,

8 orders, and 5 fungal classes. Due to a lack of close sequence matches in the UNITE database

and/or a lack of taxonomic information for close sequence matches, we were unable to identify

18 OTUs to the fungal species level, and four of these to the genus level (Table 2). The

taxonomic results indicate that many of the OTUs represent close relatives of known plant

pathogens including *Alternaria, Cladosporium, Drechslera, Phaeosphaeria, Pyrenophora,*

Stemphylium, Ramularia, and *Wojnowiciella* spp. (Table 2) (Kirk *et al.* 2008; Stukenbrock and

McDonald 2008; Havis *et al.* 2015; Marin-Felix *et al.* 2017, 2019). The largest OTU contained

48 isolates and the smallest OTUs consisted of one isolate each. OTUs are hereafter referred to

as "species" (O'Brien et al. 2005).

The serpentine fungal community was more diverse than the nonserpentine community at

every taxonomic level (Tables 2, S1-S5). Twenty-four fungal species were isolated from

serpentine *S. pulchra* (Fisher's alpha = 9.324, 95% CI = 4.664, 13.983) and 22 from

nonserpentine (Fisher's alpha = 7.261, 95% CI = 3.597, 10.925); 16 of these species (53%) were

shared (Fig. 2; Tables 1, 2). ANOVA results showed no significant relationship between soil

type, year, or year x soil type interactions and fungal richness or Fisher's alpha at the site level

(Tables S6, S7). Novel genera were isolated for both community types in every year of surveying

(Table S5). Species accumulation curves did not approach horizontal asymptotes for either soil

type, and the serpentine curve had a steeper slope than the nonserpentine curve (Fig. S1). These

curves suggest that neither fungal community was fully described, and that there were more

unsampled fungal species associated with the plants growing in serpentine soil than those in

nonserpentine soil.

Both fungal communities were dominated by a few abundant species (isolated >10

times), but the numerically dominant species differed between nonserpentine and serpentine

soils: these were *Alternaria* cf. *infectoria* and *Ramularia* cf. *proteae*, respectively (Table 2; Fig.

S2). The nonserpentine community included five abundant species, representing 62% of the

isolates, while the serpentine community included two abundant species, representing 51% of

serpentine isolates (Table 2; Fig S2). The serpentine community had a higher proportion of rare

species (<3 isolates) than the nonserpentine community, 63% versus 36%, respectively. One

Drechslera species was abundant in the nonserpentine community and rare in the serpentine

(Table 2; Fig. S2).

Soil type and year significantly affected fungal community composition, based on two-

way PERMANOVA analysis of site-level fungal communities and NMDS visualization of

community similarity (soil type: F-value = 5.112, R^2 = 0.163 p = 0.003; year: F-value = 5.655,

R^2 = 0.360 p = 0.001; Fig. 3; Table 4). Serpentine fungal communities were generally more

similar to one another than they were to nonserpentine fungal communities, and vice versa (Fig.

3; Tables 2, 3). Community composition differed significantly between 2015 and 2017, and 2016

and 2017, but not between 2015 and 2016 (Table S8).

Serpentine community composition was more consistent across years than

nonserpentine fungal community composition (Fig. 3; Tables 2, 3). While nonserpentine fungal

community similarity was unpredictable between years, with some very dissimilar communities

(Morisita-Horn overlap = 0.295) and some very similar ones (Morisita-Horn overlap = 0.882),

serpentine fungal communities were always relatively similar between years, with Morisita-Horn

overlap ranging from 0.707 to 0.786. Morisita-Horn overlap for nonserpentine and serpentine

fungal communities in each year was also variable, ranging from 0.535 to 0.725 (Table 3).

Leaf tissue chemistry and disease burden

As expected from soil chemistry differences, leaf chemistry in plants grown on serpentine

versus nonserpentine soil differed in mean C, N, Ca, Mg, Ni, and P content, and in Ca:Mg ratio.

Serpentine plants had significantly lower mean Ca, C, N, and P content and Ca:Mg ratio than

nonserpentine plants and significantly higher Ni and Mg (Fig. 4; Table S9). Cr content and C:N

ratio were similar for plants from both soil types. The results of Pearson's product-moment

correlation tests relating log-transformed mean percent symptomatic tissue to percent C, percent

N, and C:N ratio in asymptomatic foliar tissue were not statistically significant for serpentine or

nonserpentine plants (Fig. 5).

Discussion

Here we show that a diverse suite of foliar fungi was associated with ubiquitous, low-
burden damage on host individuals across growing seasons and environmental conditions. Soil
type and time appeared to shape the composition and diversity of fungal pathogen communities
in this system. These results provide some of the first evidence that local growing conditions
structure the pathogen communities infecting a single host species even over relatively small
spatial scales of a few hundred meters (Figs. 1a, 3; Table 3).

Compared to other plant species' interactions with both pathogenic and non-pathogenic
fungi, *S. pulchra* appears to host an unusually species-rich fungal community within its diseased
foliar tissue. Where we isolated 30 fungal species, past studies that described fungal pathogen
diversity for more than 100 different plant hosts consistently found fewer than 20 fungal
pathogens per species, and almost all hosts had fewer than ten pathogens (Mitchell and Power
2003; Hantsch *et al.* 2013; Rottstock *et al.* 2014). The level of fungal diversity we documented is
also high compared to that of fungal endophyte communities associated with asymptomatic plant
tissues: of 35 plant species analyzed by Arnold and colleagues, only five were found to host 30
or more endopyhtes (Arnold and Lutzoni 2007; Del Olmo-Ruiz and Arnold 2014). The
exceptional diversity of the foliar fungi that inhabit *S. pulchra* is further indicated both by
species accumulation curves that suggest many more species in this community remain to be
discovered (Fig. S1) and by the lack of close sequence matches in genetic databases for some of
the species isolated (Table 2). These findings suggest that the characteristics that justified the
identification of the California Floristic Province as a biodiversity hotspot may extend to the
fungal diversity associated with this flora (Myers *et al.* 2000).

Although disease burden was similar in both soil types, fungal communities varied by
soil type in both their composition and their stability over time (Figs. 1c, 2, 3, S2; Tables 1-4).
Higher fungal diversity and lower species turnover between years in the serpentine align with
patterns in serpentine versus nonserpentine grassland plant community diversity and temporal
stability (Fernandez-Going *et al.* 2012). The results thus imply that pathogen sharing among
hosts in native-dominated serpentine grasslands versus nonnative-dominated nonserpentine
grasslands leads to different plant-fungal interactions based on soil type. That the chemical

composition of *S. pulchra* reflected the chemistry of the soil in which it grew (except for Cr

levels) suggests that soil influence on plant phenotype may also contribute to differences

between fungal communities (Fig. 4, Table S9) (Oze *et al.* 2008).

Both the close proximity of the sampled nonserpentine and serpentine plants and the

finding that fungal communities were more similar between more distant sites of the same soil

type than they were between nearby sites of different soil types suggest that environmental

filtering, rather than dispersal limitation, plays a key role in fungal community assembly (Table

4). In addition to its effects on tissue chemistry and plant community composition, soil type

might impact disease dynamics if differing water availability between serpentine and

nonserpentine soils alters leaf physiology in ways that affect infection processes. We hypothesize

that some or all of these factors influence the composition of fungal communities and their

variation across environments.

Differences in nonserpentine and serpentine fungal communities may result from

differences in the plant communities found on these soils. Serpentine *S. pulchra* pathogen

communities were more diverse than nonserpentine communities at the species, genus, family,

order, and class levels (Fig. S2; Tables 1, S1-S5). Results from previous studies that demonstrate

a positive relationship between host diversity and pathogen diversity indicate that higher plant

species diversity in serpentine grasslands might contribute to higher fungal diversity if native

forbs and bunchgrasses that specialize on serpentine soils share foliar fungal pathogens with *S.*

pulchra (Hechinger and Lafferty 2005; Spear 2017). Research on five common invasive and

native grasses (including *S. pulchra*) at JRBP shows that multi-host foliar fungal pathogens

dominate this system, supporting this hypothesis (Spear and Mordecai 2018).

Differing heavy metal content in the leaves of nonserpentine and serpentine *S. pulchra*

may influence fungal community assembly through a variety of mechanisms. Soil Ca content is

known to influence the outcomes of plant-pathogen interactions through its roles in cellular and

biochemical pathways in both the plant immune system (Levine *et al.* 1996; Romeis *et al.* 2001)

and in fungal pathogen virulence, growth, and reproduction (e.g., Shaw and Hoch 2000; Uhm *et*

al. 2003; Brandhorst *et al.* 2005). Fungal pathogen interspecific variation in reliance on Ca for

successful infection and/or fungal response to Ca-dependent plant immune defenses could

therefore explain differences in nonserpentine and serpentine fungal community composition.

Previous studies showed that *Hesperolinon* sp. infection frequency and severity by the fungal

rust *Melampsora lini* in California grasslands depended on plant tissue Ca levels linked to

serpentine soil chemistry (Springer *et al.* 2006; Springer 2009). Varying tolerances of heavy

metals other than Ca among fungi may act as additional filters on the species present in

nonserpentine and serpentine communities (Iram *et al.* 2009).

Alternatively, serpentine and nonserpentine fungal communities could differ due to

genetic divergence between the *S. pulchra* populations found in these soils. The gene-for-gene

model of the evolution of virulence hypothesizes that a given pathogen can infect a given host

only if the pathogen does not possess an avirulence gene recognized by a matching resistance

gene in the host (Flor 1956; Parker and Gilbert 2004). To date, there have been few studies of

genetic differences between serpentine and nonserpentine populations of the same plant species.

However, genetic and chemical analyses of serpentine and nonserpentine populations of the

native forb *Lasthenia californica* at JRBP show that these populations represent two

phylogenetically distinct species, each of which accumulates ions in significantly different

concentrations and has a unique flavonoid profile (Desrochers and Bohm 1993; Chan *et al.* 2002;

Rajakaruna *et al.* 2003). Since many flavonoids have antifungal properties, serpentine and

nonserpentine *S. pulchra* could also contain different genotypes with different potential for

resistance to any given fungus due to divergent resistance gene and/or flavonoid profiles (Parker

and Gilbert 2004; Treutter 2006).

Species turnover between years contributed substantially to fungal diversity in this

system (Fig. 3; Tables 1, S5). Interannual variation in these communities may be due to highly

variable yearly precipitation (2015 rainfall = 491 mm; 2016 = 604 mm, 2017 = 985 mm)

(Weather Underground). Timing and amount of precipitation might alter infection dynamics by

washing away fungal spores, transmitting spores to new hosts, and/or influencing plant

community composition (Hobbs and Mooney 1991; Madden 1997). Although few studies track

interannual variation in fungal communities on plants, the year-to-year dissimilarity of those

hosted by *S. pulchra* is similar to that observed across a diverse taxonomic range of plants

including ferns, legumes, and shrubs (Olmo-Ruiz and Arnold 2014; Cotton *et al.* 2015; Zhang *et*

al. 2016). In this context, our results add to a growing body of evidence for the general

importance of storage effects—in which temporally variable environmental conditions act as

ecological niches that facilitate species diversity—in plant-associated fungal communities

(Warner and Chesson 1985). This finding highlights the necessity of long-term data collection to
accurately characterize plant-fungal interaction networks.

476   Contrary to our hypothesis, differences in soil chemistry that extended to plant chemistry
did not affect disease burden (Figs. 1, 4; Table S9). Elevated heavy metal content in serpentine-
grown plants was far below "hyperaccumulation" levels that have been shown to inhibit
pathogenic infection in plants (e.g., 1.92×10^{-2} mol Cr/kg required for hyperaccumulation versus
3.30×10^{-4} mol Cr/kg observed in serpentine *S. pulchra*) (Jaffre *et al.* 1976; Brooks *et al.* 1977;
Boyd *et al.* 1994) (Table S9), but the consistently low disease burden observed in both soil types,
combined with evidence from recent work that symptoms in the amounts observed inflict only
minimal plant fitness costs, suggests that *S. pulchra* is able to effectively resist or suppress
damage by foliar pathogens such that heightened defenses via opportunistic hyperaccumulation
would provide little to no fitness benefit (Spear and Mordecai 2018; Uricchio *et al.* 2019). The
consistent presence of low amounts of foliar disease on *S. pulchra* implies a coevolutionary
relationship between this bunchgrass and its foliar fungi that has led to species interactions
nearer the commensal than the parasitic end of the symbiotic spectrum.

489   The variation in fungal community composition we observed on a single host species
with soil chemistry and over time suggests possible mechanisms preventing foliar disease from
regulating plant host populations in California grasslands by impacting host fitness. Transitions
between distinct soil types over spatial scales within the range of fungal spore dispersal may
prevent disease outbreaks by hampering the spread of pathogens that are more or less virulent
depending on leaf tissue chemistry. Variable climatic conditions favoring successful infection by
different fungal species each growing season (Table S5), combined with plant senescence during
the dry season each year, suggest temporal barriers to pathogens evolving high virulence on both
nonserpentine and serpentine plants.

498   The processes that limit disease burden and impact in *S. pulchra* suggest that climate and
growth form may explain differences results from studies in tropical rainforest systems that
report large fungal pathogen burdens on host populations (e.g., Augspurger and Kelly 1984;
Spear *et al.* 2015). For example, seasonal senescence and local environmental barriers to
transmission in semi-arid and Mediterranean systems may limit the severity of individual disease
outbreaks, in contrast to year-round suitable climates and potentially more homogenous biotic

and abiotic environments in tropical systems. Life histories that promote the accumulation of

large disease burdens, along with relatively high and constant humidity, may be important

determinants of the potential for fungal pathogens to regulate plant host populations. This

observation implies potential climate- and life history-based limits on mechanisms through

which disease is hypothesized to structure plant communities, including enemy escape effects

(Keane and Crawley 2002), density- or distance-dependent Janzen-Connell effects (Janzen 1970;

Connell 1971; Mordecai 2011), and epidemic-driven community shifts (Day and Monk 1974).

Clarifying the specific mechanisms that drive differences in serpentine and nonserpentine

fungal communities will require inoculation experiments to test how plant susceptibility to

infection by different fungal species changes with plant tissue chemistry and plant genetics.

Additionally, characterization of comprehensive plant-pathogen networks including all plant

species co-occurring with *S. pulchra* in serpentine and nonserpentine grasslands will improve our

understanding of the role of the surrounding plant community in structuring fungal pathogen

communities.

Conclusions

Fungi associated with foliar disease in the native bunchgrass *S. pulchra* are relatively

benign, likely cause minimal fitness impacts (Spear and Mordecai 2018), and contribute

substantially to the overall biodiversity of California grasslands. The high taxonomic diversity

and rarity of serpentine fungi demonstrates that this community is potentially at risk of species

loss via host-parasite coextinction as nonnative annual grasses continue to expand their ranges at

the cost of native plants, and human activities including development and increased C and N

deposition contribute to the loss of serpentine grasslands that already make up <2% of the

California's surface area (Huenneke *et al.* 1990; Vallano *et al.* 2012). This work demonstrates

that, in addition to >200 endemic plant species, California's serpentine grasslands support unique

communities of leaf-infecting fungi, providing additional motivation for active conservation of

native grassland communities based on the far-reaching loss of biodiversity across trophic levels

associated with their disappearance (Sprent 1992; Dunn *et al.* 2009).

Acknowledgements

We thank Nona Chiariello, Caroline Daws, Joe Wan, Peter Vitousek, Scott Fendorf, Lawrence Uricchio, Guangchao Li, Douglas Turner, Virginia Walbot, Cody Hamilton, and the Mordecai and Peay Lab Groups for their help. This work was supported by the Jasper Ridge Kennedy Endowment, the National Science Foundation (DEB-1518681), Stanford Vice Provost for Undergraduate Education, and Stanford Bio-X.

References

Abbate JL, Antonovics J. 2014. Elevational disease distribution in a natural plant–pathogen
system: insights from changes across host populations and climate. *Oikos* **123:** 1126–1136.

Abou-Shanab RA, Angle JS, Delorme TA, Chaney RL, Van Berkum P, Moawad H, hanem
K, Ghozlan HA. 2003. Rhizobacterial effects on nickel extraction from soil and uptake by
Alyssum murale. *New Phytologist* **158:** 219–224.

Ackerly DD, Knight CA, Weiss SB, Barton K, Starmer KP. 2002. Leaf size, specific leaf area
and microhabitat distribution of chaparral woody plants: contrasting patterns in species
level and community level analyses. *Oecologia* **130:** 449–457.

Agrios GN. 2005. *Plant Pathology.* Burlington, MA, USA. Elsevier Academic Press.

Anagnostakis, SL. 1987. Chestnut blight: the classical problem of an introduced pathogen.
Mycologia **79(1):** 23-37.

Anderson MJ. 2001. A new method for non-parametric multivariate analysis of variance.
*Austral **Ecology** **26:** 32–46.

Arbizu PM. 2016. *How can I do PerMANOVA pairwise contrasts in R?*[WWW document]
Researchgate 2016.
URLhttps://www.researchgate.net/post/How_can_I_do_PerMANOVA_pairwise_contrasts_
in_R. [accessed 21 June 2019].

Augspurger CK, Kelly, CK. 1984. Pathogen mortality of tropical tree seedlings: experimental
studies of the effects of dispersal distance, seedling density, and light conditions. *Oecologia*
61: 211-217.

Borer ET, Laine A, Seabloom EW. 2016. A multiscale approach to plant disease using the
metacommunity concept. *Annual Review of Phytopathology* **54:** 397–418.

Boyd RS, Shaw JJ, Martens SN. 1994. Nickel hyperaccumulation defends *Streptanthus*
polygaloides (Brassicaceae) against pathogens. American Journal of Botany 81:294–300.

Brady KU, Kruckeberg AR, Bradshaw HD. 2005. Evolutionary ecology of plant adaptation to
serpentine soils. *Annual Review of Ecology, Evolution, and Systematics* **36:** 243–266.

Brandhorst TT, Gauthier GM, Stein RA, Klein BS. 2005. Calcium binding by the essential

virulence factor BAD-1 of *Blastomyces dermatitidis*. *Journal of Biological Chemistry* **280:**

42156–42163.

Brooks RR, Lee J, Reeves RD, Jaffre T. 1977. Detection of nickeliferous rocks by analysis of

herbarium specimens of indicator plants. *Journal of Geochemical Exploration* **7:** 49–57.

Burdon JJ, Laine A-L. 2019. *Evolutionary Dynamics of Plant-Pathogen Interactions.*

Cambridge, UK. Cambridge University Press.

Cantarelli MA, Pellerano RG, Del Vitto LA, Marchevsky EJ, Camiña JM. 2010.

Characterisation of two South American food and medicinal plants by chemometric

methods based on their multielemental composition. *Phytochemical Analysis* **21:** 550–555.

Chan R, Baldwin BG, Ornduff R. 2002. Cryptic goldfields: A molecular phylogenetic

reinvestigation of *Lasthenia californica* sensu lato and close relatives (Compositae:

Heliantheae sensu lato). *American Journal of Botany* **89:** 1103–1112.

Chao A, MA KH, Hsieh TC, Chiu C-H. 2016. SpadeR: Species prediction and diversity

estimation with R.

Ciompi S, Gentili E, Guidi L, Soldatini GF. 1996. The effect of nitrogen deficiency on leaf gas

exchange and chlorophyll fluorescence parameters in sunflower. *Plant Science* **118:** 177–

184.

Coley-Smith JR, Cooke RC. 1971. Survival and germination of fungal sclerotia. *Annual Review*

of Phytopathology **9:** 65–92.

Connell JH. 1971. On the role of natural enemies in preventing competitive exclusion in some

marine animals and in rain forest trees. In Den Boer PJ, Gradwell GR, eds. *Dynamics of*

numbers in populations. Proceedings of the Advanced Study Institute. Wageningen, The

Netherlands: Centre for Agricultural Publication and Documentation, 298–312.

Cunningham SA, Summerhayes B, Westoby M. 1999. Evolutionary divergences in leaf

structure and chemistry, comparing rainfall and soil nutrient gradients. *Ecology* **69:** 569–

588.

Dahl DB, Scott D, Roosen C, Magnusson A, Swinton J. 2019. xtable: Export Tables to LaTeX or HTML. R package version 1.8-4. https://CRAN.R-project.org/package=xtable

Day FP, Monk CD. 1974. Vegetation patterns on a southern Appalachian watershed. *Ecology* **55:** 1064–1074.

Desrochers AM, Bohm BA. 1993. Flavonoid variation in *Lasthenia californica* (Asteraceae). *Biochemical Systematics and Ecology* **21:** 449–453.

Dormann CF, Gruber B, Fruend J. 2008. 2008. Introducing the bipartite package: analysing ecological networks. *R news* **8:** 8–11.

Dowle M, Srinivasan A. 2019. data.table: Extension of 'data.frame'. R package version 1.12.0. https://CRAN.R-project.org/package=data.table

Dunn RR, Harris NC, Colwell L, Koh P, Sodhi NS. 2009. The sixth mass coextinction: are most endangered species parasites and mutualists? *Proceedings of the Royal Society B: Biological Sciences* **276:** 3037–3045.

Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26(19):** 2460-2461.

Eviner VT. 2016. Grasslands. In Mooney H, Zavaleta E, eds. *Ecosystems of California*. Oakland, CA, USA. University of California Press, 449–477.

Fernandez-Going BM, Anacker BL, Harrison SP. 2012. Temporal variability in California grasslands: soil type and species functional traits mediate response to precipitation. *Ecology* **93 (9):** 2104 – 2114.

Field CB, Chapin FS, Chiariello NR, Holland EA, Mooney HA. 1996. The Jasper Ridge CO_2 Experiment: Design and Motivation. In: Koch GW, Mooney HA, eds. *Carbon dioxide and terrestrial ecosystems.* San Diego, CA, USA: Academic Press, 121-145.

Fisher MC, Henk DA, Briggs CJ, Brownstein JS, Madoff LC, McCraw SL, Gurr SJ. 2012. Emerging fungal threats to animal, plant and ecosystem health. *Nature* **484:** 186–194.

Fisher RA, Corbet AS, Williams CB. 1943. The relation between the number of species and
the number of individuals in a random sample of an animal population. *Journal of Animal*
Ecology **12:** 42–58.

Flor HH. 1956. The complementary genic systems in flax and flax rust. *Advances in Genetics* **8:**
29–54.

Gardener M. 2014. *Community ecology: analytical methods using R and Excel*. Exeter, UK:
Pelagic Publishing.**Hansen RA, Coleman DC. 1998.** Litter complexity and composition are
determinants of the diversity and species composition of oribatid mites (Acari: Oribatida) in
litterbags. *Applied Soil Ecology* **9:** 17–23.

Hantsch L, Braun U, Scherer-Lorenzen M, Bruelheid H. 2013. Species richness and species
identity effects on occurrence of foliar fungal pathogens in a tree diversity experiment.
Ecosphere **4(7):** 81.

Harrison SP, Viers JH. 2007. Serpentine Grasslands. In: Stromberg MR, Corbin JD, D'Antonio
CM, eds. *California Grasslands: Ecology and Management*. Berkeley, CA, USA:
University of California Press, 145-155.

Havis ND, Brown JKM, Clemente G, Frei P, Jedryczka M, Kaczmarek J, Kaczmarek M,
Matusinsky P, McGraan GRD, Pereyra S, *et al.* 2015. *Ramularia collo-cygni*—an
emerging pathogen of barley crops. *Phytopathology.* **105(7):** 895–904.

Hechinger RF, Lafferty KD. 2005. Host diversity begets parasite diversity: bird final hosts and
trematodes in snail intermediate hosts. *Proceedings of the Royal Society B: Biological*
Sciences **272:**1059–1066.

Heinze J, Bergmann J, Rillig MC, Joshi J. 2015. Negative biotic soil-effects enhance
biodiversity by restricting potentially dominant plant species in grasslands. *Perspectives in*
Plant Ecology, Evolution and Systematics **17:** 227–235.

Hobbs RJ, Mooney HA. 1985. Community and population dynamics of serpentine annual
grassland in relation to gopher disturbances. *Oecologia* **67:** 342–351.

Hobbs RJ, Mooney HA. 1991. Effects of rainfall variability and gopher disturbance on
serpentine annual grassland dynamics. *Ecology* **72:** 59–68.

Huenneke LF, Hamburg SP, Koide R, Mooney HA, Vitousek PM. 1990. Effects of soil

resources on plant invasion and community structure in Californian serpentine grassland.

Ecology **71:** 478–491.

Iram S, Ahmad I, Javed B, Yaqoob S, Akhtar K, Kazmi MR, Badar-uz-Zaman. 2009.

Fungal tolerance to heavy metals. *Pakistan Journal of Botany* **41:** 2583–2594.

Janzen DH. 1970. Herbivores and the number of tree species in tropical forests. *The American*

Naturalist **104:** 501–528.

Jost L, Chao A, Chazdon RL. 2011. Compositional similarity and beta diversity. In: McGill BJ,

Magurran AE, editors. *Biological diversity: frontiers in measurement and assessment.*

Oxford, UK: Oxford University Press, 66-84.

Keane RM, Crawley MJ. 2002. Exotic plant invasions and the enemy release hypothesis.

Trends in Ecology and Evolution **17:** 164–170.

Kearse MR, Moir A, Wilson S, Stones-Havas M, Cheung S, Sturrock S, Buxton A, Cooper

S, Markowitz C, Duran T et al. 2012. Geneious Basic: an integrated and extendable

desktop software platform for the organization and analysis of sequence data.

Bioinformatics **28(12):** 1647-1649.

Khan AG. 2005. Role of soil microbes in the rhizospheres of plants growing on trace

metal contaminated soils in phytoremediation. *Journal of Trace Elements in Medicine and*

Biology. **18**: 355–364.

Kindt R, Coe R. 2005. BiodiversityR. Tree diversity analysis. A manual and software for

common statistical methods for ecological and biodiversity studies. World Agroforestry

Centre (ICRAF), Nairobi.

Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008. *Ainsworth and Bisby's Dictionry of the*

Fungi. Oxon, UK: CAB International.

Lee SC, Lee YH. 1998. Calcium/calmodulin-dependent signaling for appressorium formation in

the plant pathogenic fungus *Magnaporthe grisea. Molecules & Cells* **8:** 698–704.

Levine A, Pennell RI, Alvarez ME, Palmer R, Lamb C. 1996. Calcium-mediated apoptosis in
a plant hypersensitive disease resistance response. *Current Biology* **6:** 427–437.

Loehrer M, Schaffrath U. 2011. Soybean rust – meet a prominent challenge in soybean
cultivation. In *Ng T-B*, editor. *Soybean – Biochemistry, Chemistry, and Physiology.* London,
UK: InTech, 83–100 .

Madden, LV. 1997. No effects of rain on splash dispersal of fungal pathogens. *Canadian*
Journal of Plant Pathology **19:** 225–230.

Magurran AE. 2013. *Measuring Biological Diversity*. Malden, MA, USA: Blackwell Science
Ltd.

Marin-Felix Y, Groenewald JZ, Cai L, Chen Q, Marincowitz S, Barnes I, Bensch K, Braun
U, Camporesi E, Damm U, *et al.* 2017. Genera of phytopathogenic fungi: GOPHY 1.
Studies in Mycology **86:** 99–216.

Marin-Felix Y, Hernández-Restrepo M, Wingield MJ, Akulov A, Carnegie AJ,
Cheewangkoon R, Gramaje D, Groenewald JZ, Guarnaccia V, Halleen F, *et al.* 2019.
Genera of phytopathogenic: GOPHY 2. *Studies in Mycology* **92:** 47–133.

McElrone AJ, Reid CD, Hoye KA, Hart E, Jackson RB. 2005. Elevated CO_2 reduces disease
incidence and severity of a red maple fungal pathogen via changes in host physiology and
leaf chemistry. *Global Change Biology* **11:** 1828–1836.

McNaughton SJ. 1968. Structure and function in California grasslands. *Ecology* **49:** 962–972.

Mitchell CE, Power AO. 2003. Release of invasive plants from fungal and viral pathogens.
Nature **421:** 625–627.

Mordecai EA. 2011. Pathogen impacts on plant communities: unifying theory, concepts, and
empirical work. *Ecological Monographs* **81:** 429–441.

Nazareno ES, Li F, Smith M, Park RF, Kianian SF, Figureoa M. 2018. *Puccinia coronata* f.
sp. *avenae*: a threat to global oat production. *Molecular Plant Pathology* **19(5):** 1047–1060.

Neuwirth, E. 2014. RColorBrewer: ColorBrewer Palettes. R package version 1.1-2.
https://CRAN.R-project.org/package=RColorBrewer

O'Brien HE, Parrent JL, Jackson JA, Moncalvo J, Vilgalys R. 2005. Fungal community
analysis by large-scale sequencing of environmental samples. *Applied and Environmental*
Microbiology **71:** 5544–5550.

Oksanen J, Kindt R, Legendre P, O'Hara B. 2016. Vegan: community ecology package. R
package.

Osbourne AE. 1996. Preformed antimicrobial compounds and plant defense against fungal
attack. *The Plant Cell* **8:** 1821–1831.

Oze C, Fendorf S, Bird DK, Coleman RG. 2004. Chromium geochemistry in serpentinized
ultramafic rocks and serpentine soils from the Franciscan complex of California. *American*
Journal of Science **304:** 67–101.

Oze C, Skinner C, Schroth AW, Coleman RG. 2008. Growing up green on serpentine soils:
biogeochemistry of serpentine vegetation in the Central Coast Range of California. *Applied*
Geochemistry **23:** 3391–3403.

Parker IM, Gilbert GS. 2004. The evolutionary ecology of novel plant-pathogen interactions.
Annual Review of Ecology, Evolution, and Systematics **35:** 675–700.

Parker IM, Saunders M, Bontrager M, Weitz AP, Hendricks R, Magarey R, Suiter K,
Gilbert GS. 2015. Phylogenetic structure and host abundance drive disease pressure in
communities. *Nature* **520:** 542–544.

Pineda A, Zheng S-J, van Loon JJA, Pieterse CMJ, Dicke M. 2010. Helping plants to deal
with insects: the role of beneficial soil-borne microbes. *Trends in Plant Science* **15:** 507-
514.

Rajakaruna N, Baldwin BG, Chan R, Desrochers AM, Bohm BA, Whitton J. 2003. Edaphic
races and phylogenetic taxa in the *Lasthenia californica* complex (Asteraceae: Heliantheae):
an hypothesis of parallel evolution. *Molecular Ecology* **12:** 1675–1679.

Rascio N, Navari-Izzo F. 2011. Heavy metal hyperaccumulating plants: how and why do they

do it? and what makes them so interesting? *Plant Science* **180:** 169–181.

Rizzo DM, Garbelotto M. 2003. Sudden oak death: endangering California nad Oregon forest
ecosystems. *Frontiers in Ecology and the Environment* **1(5):** 197-204.

Romeis T, Ludwig AA, Martin R, Jones JDG. 2001. Calcium-dependent protein kinases play
an essential role in a plant defence response. *EMBO Journal* **20:** 5556–5567.

Rossi J-P. 2011. rich: an R package to analyse species richness. *Diversity* **3:** 112–120.

Rottstock T, Joshi J, Kummer V, Fischer M. 2014. Higher plant diversity promotes higher
diversity of fungal pathogens, while it decreases pathogen infection per plant. *Ecology* **95(7):**
1907-1917.

Seabloom EW, Borer ET, Gross K, Kendig AE, Lacroix C, Mitchell CE, Mordecai EA,
Power AG. 2015. The community ecology of pathogens: coinfection, coexistence and
community composition. *Ecology Letters* **18:** 401–415.

Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Sahl JW. 2009.
Introducing mothur: open-source, platform-independent, community-supported software for
describing and comparing microbial communities. *Applied and Environmental*
Microbiology **75(23):** 7537-7541.

Shaw BD, Hoch HC. 2000. Ca^{2+} regulation of *Phyllosticta ampelicida* pycnidiospore
germination and appressorium formation. *Fungal Genetics and Biology* **31:** 43–53.

Smith V. 2007. Host resource supplies influence the dynamics and outcome of infectious
disease. *Integrative and Comparative Biology* **47:** 310–316.

Spear ER. 2017. Phylogenetic relationships and spatial distributions of putative fungal
pathogens of seedlings across a rainfall gradient in Panama. *Fungal Ecology* **26:** 65–73.

Spear ER, Coley PD, Kursar TA. 2015. Do pathogens limit the distributions of tropical trees
across a rainfall gradient? *Journal of Ecology* 103:165–174.

Spear ER, Mordecai EA. 2018. Foliar pathogens do not stabilize coexistence of competing
species in a California grassland. *Ecology* **99(10):** 2250–2259.

Sprent JFA. 1992. Parasites lost? *International Journal for Parasitology* **22:** 139–151.

Springer YP. 2009. Do extreme environments provide a refuge from pathogens? A phylogenetic
test using serpentine flax. *American Journal of Botany* **96:** 2010–2021.

Springer YP, Hardcastle BA, Gilbert GS. 2006. Soil calcium and plant disease in serpentine
ecosystems: a test of the pathogen refuge hypothesis. *Oecologia* **151:** 10–21.

Stukenbrock EH, McDonald BA. 2008. The origins of plant pathogens in agro-ecosystems.
Annual Review of Phytopathology. **46:** 75–100.

Tadros TM. 1957. Evidence of the presence of an edapho-biotic factor in the problem of
serpentine tolerance. *Ecology* **38:** 14–23.

Telfer S, Lambin X, Birtles R, Beldomenico P, Burthe S, Paterson S, Begon M. 2010.
Species interactions in a parasite community drive infection risk in a wildlife population.
Science **330:** 243–247.

Treutter D. 2006. Significance of flavonoids in plant resistance: a review. *Environmental*
Chemistry Letters **4:** 147–157.

Uhm K-H, Ahn I-P, Kim S, Lee Y-H. 2003. Calcium/calmodulin-dependent signaling for
prepenetration development in *Colletotrichum gloeosporioides*. *Phytopathology* **93:** 82–87.

UNITE Community. 2017a. Full mothur UNITE+INSD dataset 2. Version

01.12.2017. https://doi.org/10.15156/BIO/587480

UNITE Community. 2017b. UNITE general FASTA release. Version 01.12.2017.

https://doi.org/10.15156/BIO/587475

Uricchio LH, Daws SC, Spear ER, Mordecai EA. 2019. Exotic species dominate due to niche
overlap in a complex grassland community. *The American Naturalist* **193(2):** 213-226.

Vallano DM, Selmants PC, Zavaleta ES. 2012. Simulated nitrogen deposition enhances the
performance of an exotic grass relative to native serpentine grassland competitors. *Plant*
Ecology **213:** 1015–1026.

Vavrek MJ. 2011. Fossil: palaeoecological and palaeogeographical analysis tools.
Palaeontologia Electronica **14**.

Warner RR, Chesson PL. 1985. Coexistence mediated by recruitment fluctuations: a field
guide to the storage effect. *The American Naturalist* **125:** 769–787.

Weather Underground. 2018. Jasper Ridge Biological Preserve KCAWOODS19.
https://www.wunderground.com/personal-weather-station/dashboard?ID=KCAWOODS19.

Wickham H. 2007. Reshaping data with the reshape package. *Journal of Statistical Software* **21(**
12): 1-20.
800
Wickham H. 2017. tidyverse: Easily Install and Load the 'Tidyverse'. R package version 1.2.1.
https://CRAN.R-project.org/package=tidyverse**Wilke CO. 2019.** cowplot: Streamlined Plot
Theme and Plot Annotations for 'ggplot2'. R package version 0.9.4. https://CRAN.R-
project.org/package=cowplot**Wolda H. 1981.** Similarity indices, sample size and diversity.
Oecologia **50:** 296–302.

Wolf DC, Dao TH, Scott HD, Lavy TL. 1989. Influence of sterilization methods on selected
soil microbiological, physical, and chemical properties. *Journal of Environment Quality*
18(1): 39-44.

    Figures

Figure 1. Pathogen damage on *Stipa pulchra* was consistently low across three years and both grassland types. Panel **(a)** maps the locations of surveyed nonserpentine sites (blue) and serpentine sites (green) along the ridgetop at JRBP. Panel **(b)** shows an example of fungal-caused pathogen foliar damage on *S. pulchra.* Panel **(c)** shows mean percentage (bars) and two standard errors (error bars) of diseased leaf area for surveyed serpentine and nonserpentine *S. pulchra,* by year.

Figure 2. Pathogens isolated from *Stipa pulchra* growing in nonserpentine and serpentine grasslands. Bipartite network showing species interactions between *S. pulchra* plants (left) growing in serpentine grasslands (green) and nonserpentine grasslands (blue) and fungal species cultured from symptomatic leaf tissue (white bars on the right). The length of each white bar correlates with the number of isolates in the operational species unit (based on 97% sequence similarity). The thickness of lines connecting left and right bars represents the number of times a particular fungal species was isolated from *S. pulchra* in a particular grassland type. The nonserpentine fungi came from 15 genera, 8 families, 4 orders, and 3 classes; the serpentine fungi came from 20 genera, 14 families, 7 orders, and 5 classes (Tables 1, S1-S5).

Figure 3. Distinct fungal communities isolated from *Stipa pulchra* growing in nearby serpentine and nonserpentine sites across years. Non-metric multidimensional scaling visualization of serpentine and nonserpentine fungal community similarity. Each point corresponds to the combined fungal community of two transects at the same site in the same year. Serpentine communities are shown in green and nonserpentine communities are shown in blue. Circles, triangles, and squares represent fungal communities sampled in 2015, 2016, and 2017, respectively. Ellipses enclose 95% confidence intervals for ordination of serpentine and nonserpentine communities. Community dissimilarity increases with the distance between points. PERMANOVA analysis showed a significant effect of soil type on fungal community composition (F = 3.623, R^2 = 0.0618, p = 0.001).

Figure 4. *Stipa pulchra* **grown in nonserpentine soil had higher C, N, P, Ca, and Ca:Mg and lower Mg and Ni than those grown in serpentine soil.** Box plots comparing elemental content of dried foliar tissue from nonserpentine- and serpentine-grown *S. pulchra*. In each plot, the thick black bar represents the median, the height of the box represents the interquartile range (IQR), the whiskers represent the maximum and minimum points within 1.5*IQR of the 25th and 75th percentiles, and the circles represent points outside this range. An asterisk above a panel indicates a statistically significant difference between serpentine- and nonserpentine-grown plants ($p < 0.05$) indicated by a Wilcoxon ranked-sum test. Panel **(a)** shows percent C of dried foliar tissue (W = 2713, p-value = 1.65×10^{-7}); panel **(b)**, percent N (W = 2346, p-value = 0.0011); panel **(c)**, C:N ratio (W = 1398, p-value = 0.066); panel **(d)**, moles P per kilogram dried leaf tissue (W = 390, p-value = 2.83×10^{-9}); panel **(e)**, moles Ca per kilogram dried leaf tissue (W = 400, p-value = 1.45×10^{-11}); panel **(f)**, Ca:Mg ratio (W = 400, p-value = 1.451×10^{-11}); panel **(g)**, moles Cr per kilogram dried leaf tissue (W = 192.5, p-value = 0.8498); panel **(h)**, moles Mg per kilogram dried leaf tissue (W = 1, p-value = 2.90×10^{-11}); and panel **(i)**, moles Ni per kilogram dried leaf tissue (W = 0, p-value = 6.79×10^{-8}).

Figure 5. C, N, and C:N do not predict the mean foliar pathogen damage on *Stipa pulchra*
in serpentine or nonserpentine grasslands. Foliar percent C (Panels **(a)** and **(b)**), percent N
(Panels **(c)** and **(d)**) and C:N ratio (Panels **(e)** and **(f)**) versus mean percentage of diseased leaf
area for serpentine and nonserpentine *S. pulchra* plants.

881　**Tables**

882　**Table 1. Pathogen damage on *Stipa pulchra* varied significantly by year, but not soil type.**

Variables	df	SS	MSE	F value	p-value
soil type	1	6.57×10^{-1}	6.57×10^{-1}	7.50×10^{-1}	0.39
year	2	9.41	4.71	5.37	$7.04 \times 10^{-3}*$
site	4	6.00	1.50	1.71	1.59×10^{-1}
soil type x year	2	1.80	9.02×10^{-1}	1.03	3.63×10^{-1}

ANOVA results for log-transformed transect-level measurements of percentage of symptomatic
living tissue on serpentine and nonserpentine *S. pulchra* (N = 3 sites per soil treatment per year,
with 12 transects per site per year, for a total of 36 serpentine and 36 nonserpentine transects
over all three years). The first column lists the factor(s) tested; the second column degrees of
freedom (df); the third column the sum of squares (SS); the fourth column the mean squared
error (MSE); the fifth column the F-value; and the 6th column the p-value. P-values < 0.05 are
marked with an asterisk.

 Table 2. Fungal species isolated from symptomatic *Stipa pulchra* leaves were unevenly

 distributed across soil types. Each row represents one of the species isolated. The columns

 indicate the isolation frequency for each species in each soil type.

Species	Nonserpentine	Serpentine
Alternaria cf. *infectoria*	26	21
Keissleriella sp.	19	8
Drechslera cf. *fugax*	17	1
Stemphylium sp.	15	8
Ramularia cf. *proteae*	11	37
Drechslera sp.	9	1
Neoascochyta sp.	7	3
Cladosporium sp.	6	3
Alternaria cf. *eureka*	5	2
Parastagonospora cf. *avenae*	5	0
Pyrenophera cf. *chaeotmioides*	5	2
Pyrenophera cf. *lolii*	4	3
Phaeosphaerieaceae sp. 2	3	5
Pleosporaceae sp.	3	2
Drechslera cf. *nobleae*	1	5
Epicoccum sp.	1	0
Heyderia sp.	1	1
Parastagonospora cf. *cumpignensis*	1	0
Parastagonospora sp.	1	0
Phaeosphaeria sp.	1	2
Podospora sp.	1	0
Phaeosphaerieaceae sp. 1	1	0
Chalastospora cf. *gossypii*	0	1
Hyphodiscus cf. *brevicollaris*	0	1
Nemania cf. *serpens*	0	1
Phaeophlebiopsis sp.	0	1
Plectania sp.	0	2
Pseudoseptoria sp.	0	1
Helotiales sp.	0	1
Wojnowiciella sp.	0	1

Table 3. Fungal community diversity was higher and communities were distinct on *Stipa pulchra* plants growing in serpentine versus nonserpentine grasslands, across all years.

Fungal community pair	Fisher's alpha	Shared species	Morisita-Horn Similarity Index
All years serpentine (24)	9.324	16	0.675
All years nonserpentine (22)	7.261	(53.0%)	
2015 serpentine (11)	7.955	5	0.535
2015 nonserpentine (12)	5.205	(27.8%)	
2016 serpentine (16)	7.897	11	0.766
2016 nonserpentine (13)	5.861	(61.1%)	
2017 serpentine (12)	6.784	8	0.725
2017 nonserpentine (12)	5.136	(50.0%)	
2015 serpentine (11)	7.955	7	0.707
2016 serpentine (16)	7.897	(35.0%)	
2015 serpentine (11)	7.955	5	0.721
2017 serpentine (12)	6.784	(27.8%)	
2016 serpentine (16)	7.897	7	0.786
2017 serpentine (12)	6.784	(33.3%)	
2015 nonserpentine (12)	5.205	6	0.668
2016 nonserpentine (13)	5.861	(31.5%)	
2015 nonserpentine (12)	5.205	5	0.295
2017 nonserpentine (12)	5.136	(26.3%)	
2016 nonserpentine (13)	5.861	7	0.882
2017 nonserpentine (12)	5.136	(38.9%)	

Fungal community similarity by soil type and year. The leftmost column lists the fungal communities considered, with the number of species in each community in parentheses. The second column lists Fisher's alpha for each community. The third column shows the number of shared species and, in parentheses, the percentage of total observed species that were shared for each pair of communities. The rightmost column lists the estimated Morisita-Horn community overlap value, based on absolute species abundances, for each pair of communities. The Morisita-Horn index ranges from 0 to 1, with 1 indicating complete overlap.

Table 4. PERMANOVA results show significant effects of both soil type and year on fungal foliar community composition (N = 18 serpentine and 18 nonserpentine sites).

	df	SS	MSE	F-Value	p-value
soil type	1	0.5700	0.56997	5.1119	0.003*
year	2	1.2611	0.63055	5.6552	0.001*
distance	1	0.0767	0.07667	0.5850	0.675
soil type x year	2	0.3339	0.16697	1.4975	0.224
distance x year	2	0.1697	0.08485	0.6473	0.734

From left to right, columns list factors tested, degrees of freedom (df), sum of squares (SS), mean squared error (MSE), F-value, and p-value. Statistically significant p-values ($p < 0.05$) are marked with an asterisk. The distance factor compares communities at near (<600m) vs. far (>600m) distances from one another.

Supporting Information

Fig. S1 Species accumulation curves for serpentine (green) and nonserpentine (blue) fungal

communities isolated from *Stipa pulchra* leaves.

Fig. S2 Fungal species abundance by rank for serpentine (green) and nonserpentine (blue) fungal

communities on *Stipa pulchra* leaves.

Table S1 Genus counts for serpentine and nonserpentine fungal communities isolated from *Stipa*

pulchra leaves.

Table S2 Family counts for nonserpentine and serpentine fungal communities isolated from

Stipa pulchra leaves.

Table S3 Order counts for nonserpentine and serpentine fungal communities isolated from *Stipa*

pulchra leaves.

Table S4 Class counts for nonserpentine and serpentine fungal communities isolated from *Stipa*

pulchra leaves.

Table S5 Presence and absence of genera in nonserpentine and serpentine fungal communities

isolated from *Stipa pulchra* leaves, by year.

Table S6 ANOVA table for effects of soil type, year, and soil type x year interactions on site-

level fungal community diversity in terms of Fisher's alpha.

Table S7 ANOVA table for effects of soil type, year, and soil type x year interactions on site

fungal species richness.

Table S8 Pairwise PERMANOVA table for effects of year on fungal community composition.

Table S9 Results of chemical analysis of serpentine and nonserpentine *Stipa pulchra* foliar

tissue.

Methods S1 Details regarding *Stipa pulchra* plants used for ICP-OES analysis of Ca, Cr, Mg, Ni,

and P.

953   **Supporting Figures and Tables**

Figure S1. Species accumulation curves for serpentine (green) and nonserpentine (blue)
fungal communities isolated from *Stipa pulchra* leaves. The cumulative number of unique
fungal OTUs (based on 97% sequence similarity; y-axis) is plotted against the number of *S.*
pulchra leaf tissue pieces sampled (x-axis).

962
Figure S2. Fungal species abundance by rank for serpentine (green) and nonserpentine

(blue) fungal communities on *Stipa pulchra* leaves. Each point represents the number of times

a single species was isolated from *S. pulchra* leaves in each soil type.

967 **Table S1.** Genus counts for serpentine and nonserpentine fungal communities isolated from

968 *Stipa pulchra* leaves.

Genus	Nonserpentine	Serpentine	Hosts	References
Alternaria	31	23		
Drechslera	27	7		
Keissleriella	19	8		
Stemphylium	15	8		
Ramularia	11	37		
Pyrenophera	9	5		
Neoascochyta	7	3		
Parastagonospora	7	0		
Cladosporium	6	3		
Phaeosphaerieaceae	4	5		
unknown Pleosporaceae	3	2		
Epicoccum	1	0		
Heyderia	1	1		
Phaeosphaeria	1	2		
Podospora	1	0		
Chalastospora	0	1		
Hyphodiscus	0	1		
Nemania	0	1		
Phaeophlebiopsis	0	1		
Plectania	0	2		
Pseudoseptoria	0	1		
unkown Helotiales	0	1		
Wojnowiciella	0	1		

970 Numbers in each column represent the number of times a genus was isolated from *S. pulchra*

971 leaves in each soil type.

973   **Table S2.** Family counts for nonserpentine and serpentine fungal communities isolated from

974   *Stipa pulchra* leaves.

Family	Nonserpentine	Serpentine
Pleosporaceae	85	45
Lentitheciaceae	19	8
Phaeosphaeriaceae	12	8
Mycosphaerellaceae	11	37
Didymellaceae	8	3
Cladosporiaceae	6	3
Helotiaceae	1	1
Lasiosphaeriaceae	1	0
Dothioraceae	0	1
Hyaloscyphaceae	0	1
Meruliaceae	0	1
Pleosporales *incertae sedis*	0	1
Sacrosomataceae	0	2
unknown Helotiales	0	1
Xylariaceae	0	1

976   Numbers in each column represent the number of times a family was isolated from *S. pulchra*

977   leaves in each soil type.

989 **Table S3**. Order counts for nonserpentine and serpentine fungal communities isolated from *Stipa*

990 *pulchra* leaves.

Order	Nonserpentine	Serpentine
Pleosporales	124	65
Capnodiales	17	40
Helotiales	1	3
Sordariales	1	0
Dothideales	0	1
Pezizales	0	2
Polyporales	0	1
Xylariales	0	1

992 Numbers in each column represent the number of times an order was isolated from *S. pulchra*

993 leaves in each soil type.

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009 **Table S4.** Class counts for nonserpentine and serpentine fungal communities isolated from *Stipa*

1010 *pulchra* leaves.

Class	Nonserpentine	Serpentine
Dothideomycetes	141	106
Leotiomycetes	1	3
Sordariomycetes	1	1
Agaricomycetes	0	1
Pezizomycetes	0	2

1011

1012 Numbers in each column represent the number of times a class was isolated from *S. pulchra*

1013 leaves in each soil type.

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030 **Table S5.** Presence (P, shaded) or absence (A, unshaded) of genera in nonserpentine and

1031 serpentine fungal communities isolated from *Stipa pulchra* leaves, by year.

1032

Genus	Nonserpentine 2015	Nonserpentine 2016	Nonserpentine 2017	Serpentine 2015	Serpentine 2016	Serpentine 2017
Alternaria	P	P	P	P	P	P
Chalastospora	A	A	A	P	A	A
Cladosporium	P	A	A	P	A	P
Drechslera	P	P	P	P	P	P
Epicoccum	A	A	P	A	A	A
Heyderia	A	A	P	A	A	P
Hyphodiscus	A	A	A	A	P	A
Keissleriella	P	P	P	P	P	P
Nemania	A	A	A	A	A	P
Neoascochyta	A	P	P	A	P	P
Parastagonospora	P	A	P	A	A	A
Phaeophlebiopsis	A	A	A	A	P	A
Phaeosphaeria	A	P	A	P	P	A
Plectania	A	P	A	A	A	P
Podospora	A	P	A	A	A	A
Pseudoseptoria	A	A	A	A	P	A
Pyrenophera	P	P	P	P	P	P
Ramularia	P	P	A	P	P	P
Stemphylium	A	P	P	P	A	P
unknown Phaeosphaerieaceae	P	P	A	A	P	A
unknown Pleosporaceae	A	P	P	A	P	P
unkown Helotiales	A	A	A	A	P	A
Wojnowiciella	A	A	A	P	A	A

1033

1034

1035   **Table S6.** ANOVA table for effects of soil type, year, and soil type x year interactions on site-

1036   level fungal community diversity in terms of Fisher's alpha.

	df	SS	MSE	F value	p-value
Soil type	1	0.098369	0.098369	0.009695	0.923189
Year	2	3.629916	1.814958	0.178881	0.838395
Soil type x Year	2	17.34903	8.674514	0.854953	0.449656

1037

1038   Columns, from left to right, list the variable tested, the degrees of freedom, the sum of squares,

1039   the mean squared error, the F value, and the p-value.

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055 **Table S7.** ANOVA table for effects of soil type, year, and soil type x year interactions on site

1056 fungal species richness.

	df	SS	MS	F value	p-value
Soil type	1	6.722222	6.722222	1.070796	0.32117
Year	2	3.111111	1.555556	0.247788	0.784422
Soil type x year	2	8.444444	4.222222	0.672566	0.528628

1057

1058 Columns, from left to right, list the variable tested, the degrees of freedom, the sum of squares,

1059 the mean squared error, the F value, and the p-value.

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074 **Table S8.** Pairwise PERMANOVA table for effects of year on fungal community composition.

1075 From left to right, columns list the pair of years being compared, the F-value, R^2, and the

1076 Bonferroni-adjusted p-value. P-values < 0.05 are marked with an asterisk.

Pair	F-value	R^2	Bonferroni p-value
2015 vs 2016	2.616237	0.207371	0.114
2015 vs 2017	5.983453	0.374353	0.009*
2016 vs 2017	4.396858	0.304303	0.018*

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

Table S9. Results of chemical analysis of serpentine and nonserpentine *S. pulchra* foliar tissue. The Serpentine and Nonserpentine columns report elemental content in mean mol/kg dried weight leaf tissue derived from each soil type, except for C and N, which are reported as percentage of leaf tissue by weight. The standard error of each mean is listed in parentheses. W, in the third column, is the Wilcoxon-rank sum test statistic comparing serpentine and nonserpentine mean values. Asterisks denote statistically significant p-values ($p < 0.05$).

Element	Serpentine		Nonserpentine		W	p-value	
C	38.97	(3.44×10^{-1})	41.48	(2.35×10^{-1})	2713	1.65×10^{-7}	*
Ca	6.50×10^{-2}	(1.89×10^{-3})	1.8×10^{-1}	(4.78×10^{-3})	400	1.45×10^{-11}	*
Cr	3.30×10^{-4}	(6.16×10^{-5})	3.10×10^{-4}	(5.73×10^{-5})	192.5	8.50×10^{-1}	
Mg	1.40×10^{-1}	(5.10×10^{-3})	7.5×10^{-2}	(1.84×10^{-3})	1	2.90×10^{-11}	*
N	1.44	(4.96×10^{-2})	1.61	(3.84×10^{-2})	2346	1.10×10^{-3}	*
Ni	3.50×10^{-4}	(2.83×10^{-5})	6.57×10^{-5}	(4.89×10^{-6})	0	6.79×10^{-08}	*
P	2.70×10^{-2}	(8.43×10^{-4})	4.2×10^{-2}	(1.50×10^{-3})	389	2.83×10^{-09}	*
Ca:Mg	4.70×10^{-1}	(6.55×10^{-2})	2.37	(7.20×10^{-3})	0	1.45×10^{-11}	*
C:N	28.72	(8.74×10^{-1})	26.54	(6.18×10^{-1})	1398	6.62×10^{-2}	

Methods S1

Details regarding Stipa pulchra plants used for ICP-OES analysis of Ca, Cr, Mg, Ni, and P

We sourced foliar tissue for ICP-OES analysis of Ca, Cr, Mg, Ni, and P from *S. pulchra*

plants initially grown in individual pots in greenhouse conditions and then set out at Jasper Ridge

Biological Preserve (JRBP) for a separate experiment designed to assess fungal infection over

geographic space in a natural setting. Potted plants were left in the field for 12 weeks (3/9/2017 –

5/26/2017) and then returned to the greenhouse, surveyed for symptoms of foliar disease,

separated into above- and below-ground portions, and dried. Only 22 of 480 nonserpentine-

grown plants and 33 of 480 serpentine-grown plants had symptomatic leaf tissue; these plants

were excluded from the pool of samples for chemical analysis.

1123

1124

1125