Habitat type and interannual variation shape unique fungal pathogen communities on a

California native bunchgrass

Johannah E. Farner[1], Erin R. Spear[1,2], Erin A. Mordecai[1]

[1]Biology Department, Stanford University, Stanford, California 94305 USA; [2] Biology

Department, Regis University, Denver, Colorado 80221 USA

Author for correspondence:

Johannah Farner

E-mail: jfarner@stanford.edu

Phone: 1 (805) 317 – 5432

Abstract

The role of infectious disease in regulating host populations is increasingly recognized,

but how environmental conditions affect pathogen communities and infection levels remains

poorly understood. Over three years, we compared foliar disease burden, fungal pathogen

community composition, and plant tissue chemistry in the perennial bunchgrass *Stipa pulchra*

occurring in adjacent serpentine and nonserpentine grassland habitats with distinct soil types and

plant communities. We found that serpentine and nonserpentine *S. pulchra* experienced

consistent, low disease pressure associated with distinct fungal pathogen communities with high

interannual species turnover, and that plant chemistry differed with habitat type. The results

indicate that this species experiences minimal foliar disease associated with diverse fungal

pathogen communities that are structured across landscapes by spatially and temporally variable

conditions. Highly variable conditions associated with fungal species turnover may create a

refuge from disease outbreaks for *S. pulchra*, contributing to the low disease burden observed on

this and other Mediterranean grassland species.

Key words: California grasslands, leaf spot pathogen, plant fungal pathogens, serpentine soil,
Stipa pulchra

Introduction

How environmental heterogeneity affects species interactions, and ultimately population

growth rates and structures, remains a fundamental ecological question. Infectious disease, which

arises from interactions between hosts, parasites, and the environment, is particularly likely to

vary across populations in differing environments. Biotic factors, such as the phylogenetic

structure of the host community, and abiotic factors, such as elevation, have been associated with

the presence of specific diseases and the severity of their impacts on their hosts (Abbate and

Antonovics 2014; Parker *et al.* 2015). Additionally, mounting evidence suggests that specific

characteristics of pathogen community structures, including the identities and abundances of the

diseases present in a host population, mediate the outcomes of host – pathogen interactions

(Seabloom *et al.* 2015; Borer *et al.* 2016).

Together, these observations highlight the importance of understanding interactions

between environmental variation, pathogen community composition, and pathogen pressure on

host populations to assess the potential for disease to regulate host population dynamics.

For plant hosts, such assessments are important both in conservation and agricultural contexts

because identifying populations at high risk for severe disease outbreaks and applying

preventative measures may prevent local species extinctions and improve crop yields. While

studies have frequently explored the interactions of single host – pathogen pairs under differing

abiotic conditions (e.g., Abbate and Antonovics 2014), the effect of varying pathogen

community composition on disease burden and relationships between habitat characteristics and

pathogen community structures remain largely unexplored in natural systems (but see Rottstock

et al. 2014 and Spear 2017).

Fungal pathogens are particularly important to study in this context because they cause

the majority of known plant diseases, have an increasing rate of emergence, and pose a

substantial threat to plant health and population persistence in natural and agricultural settings

(Agrios 2005; Fisher *et al.* 2012). Dramatic diebacks of North American chestnut trees and oaks

caused by fungal pathogens (Anagnostakis 1987; Rizzo and Garbelotto 2003) and economically

devastating crop losses caused by fungal rusts shared among wild and domesticated plants

(Loehrer and Schaffrath 2011; Nazareno *et al*. 2018) demonstrate this threat. These examples

indicate that understanding the communities of fungal pathogens that infect wild plants is

important for developing effective protection strategies for both natural biodiversity and human

food supplies. However, we lack basic information about the identities, diversity, and effects of

the fungal pathogens that infect most plant species (Burdon and Laine 2019). Therefore, there is

a pressing need for systematic surveys of these pathogen communities and their variation across

natural habitats.

Evidence that abiotic and biotic factors drive both natural fungal endophyte community

assembly in healthy leaves and fungal foliar disease dynamics in agricultural systems suggests

that different habitats are likely to affect fungal foliar pathogen community composition and

disease burden. For example, the repeated finding that foliar endophyte community structures

change with temperature, humidity, and rainfall (Cordier *et al.* 2012, Zimmerman and Vitousek

2012, Bálint *et al.*, 2015, Millberg *et al.* 2015, Giauque and Hawkes 2016, Barge *et al.*, 2019)

suggests that climate is likely to also shape fungal pathogen communities. Evidence from the

plant pathology literature supports this expectation: for foliar fungal pathogens of wheat, barley,

and lettuce, among others, infection dynamics are closely linked to temperature and moisture

availability (Shaw 1986, Moss *et al.* 1987, Eversmyer *et al.* 1988, Huber and Gillespie 1992,

Magarey et al. 2004, Bernard *et al.* 2013, Clarkson *et al.* 2014, Velásquez *et al.* 2018).

Additionally, local habitat characteristics, such as soil and plant physical and chemical

properties, plant community composition, and plant genetics, can determine the availability and

quality of susceptible and competent hosts for different fungal pathogen species (Snoeijers *et al.*

2000, Mitchell *et al.* 2002, Solomon *et al.* 2003, Bolton *et al.* 2008, Rottstock et al. 2014,

Velásquez *et al.* 2018). Differences in soil nutrient content, plant nutrient content, and plant

genetic backgrounds are known to influence the identities and abundances of fungal endophytes

that inhabit the leaves of individual host species (Eschen *et al.* 2010, Busby *et al.* 2014). The

likely relevance of these same factors to fungal pathogens is supported by extensive work in

agricultural systems wherein nitrogen fertilization and plant genetic diversity are both associated

with disease prevalence and severity (Zhu *et al.* 2000, Solomon *et al.* 2003, Verosoglou *et al.*

2013).

Notably, general understanding of the factors that shape foliar endophyte communities is

hampered by the short lengths of most observation periods, which frequently result in annual or

seasonal snapshots of fungal community composition that cover only one or two years (but see

Giauque and Hawkes, 2016). Extended observation periods can provide the context necessary to

determine whether such snapshots are broadly representative versus exceptional, a key step for
assessing the overall strength of different predictors for infection patterns.

Here, we examine the effects of habitat type on the burden and species composition of
natural plant pathogens infecting a single host species over a span of three years. Specifically, we
take advantage of the natural occurrence of the California native perennial bunchgrass *Stipa*
pulchra in two unique, adjacent grassland types to understand the extent to which habitat type
and time influence the landscape of disease encountered by a single plant species, including
foliar fungal pathogen community composition and disease severity. Stark differences in soil
properties and plant community structures between serpentine and nonserpentine greenstone
(hereafter, nonserpentine) grassland habitats result in a suite of environmental differences that
may influence fungal pathogen infection, reproduction, and persistence (Harrison and Viers
2007; Burdon and Laine 2019). Additionally, high interannual variation in climate conditions
and plant community composition in California grasslands may lead to temporal variation in
infection dynamics through impacts on fungal growth and survival (Hobbs and Mooney 1991,
Fernandez-Going *et al.* 2012). Therefore, in this study we ask: (1) how does disease burden
differ on plants growing in different habitat types and over time? (2) how does the species
composition of fungal pathogen communities differ on plants growing in different habitat types
and over time? and (3) are there differences in plant tissue chemistry between habitat types that
correspond to differences in disease burden?

Because of differences between serpentine and nonserpentine plant community
composition, plant physiology, and plant genetics, we hypothesize that both disease burden and
pathogen species composition differ on *S. pulchra* individuals growing in these two habitats.
Moreover, because abiotic and biotic conditions vary substantially between growing seasons in
California grasslands, we hypothesize that disease burden and pathogen species composition
differ interannually on *S. pulchra*. Finally, we hypothesize that habitat type influences *S. pulchra*
foliar tissue chemistry in ways that may affect disease burden, with lower C and N content in
serpentine plants growing in nutrient-poor soils. If so, these foliar chemistry differences may
translate into either (a) higher disease burden in serpentine plants relative to nonserpentine
plants, facilitated by negative impacts of limited plant resource availability on plant health (as in
Springer *et al.,* 2006), or (b) lower disease burden in serpentine plants due to lower plant nutrient
content limiting resources available for fungal growth.

By contributing novel information about previously uncharacterized plant fungal

pathogen communities in California grasslands, the results of this study will provide new insight

into the diversity and natural history of the fungi that infect the leaves of wild grasses in

temperate climates. Further, the results of this study will aid in assessing disease risk for plant

populations in natural and agricultural contexts by providing information about the spatial and

temporal scales over which environmental heterogeneity can affect the species composition and

severity of foliar pathogen infection.

Materials and Methods

Study system

All surveys and sample collection took place at Jasper Ridge Biological Preserve (JRBP),

a 485-hectare site in San Mateo County, CA managed by Stanford University (37.4° N, 122.2°

W). JRBP has a Mediterranean climate with cool winters (mean 9.2°C), warm summers (mean

20.1°C), and annually variable precipitation that predominantly occurs in the winter and spring

(averaging 622.5 mm per year) (Ackerly *et al.* 2002). The growing season occurs during the

rainy season and plants senesce in early summer. This study was conducted during the growing

seasons from mid-April to mid-May in 2015, 2016, and 2017. There was highly variable annual

precipitation across the three years (491 mm, 604 mm, and 985 mm in 2015, 2016, and 2017,

respectively; Weather Underground).

S. pulchra occurs at substantial densities in both serpentine and nonserpentine grasslands

at JRBP (McNaughton 1968). The extent of genetic differentiation between serpentine and

nonserpentine *S. pulchra* is unknown. While there are no physical barriers at this site to prevent

gene flow between individuals in different habitat types, there is high genetic divergence

between serpentine and nonserpentine populations of the forb *Lasthenia californica* occurring in

the same grasslands at JRBP (Chan *et al.* 2002). The individuals studied here may thus represent

either a single continuous population or multiple populations that are locally adapted to different

habitat types.

The serpentine plant communities at JRBP are more diverse than the nearby

nonserpentine plant communities and include many native grasses and forbs, in contrast to the

invasive-dominated nonserpentine grasslands (McNaughton 1968; Huenneke *et al.* 1990; Field *et*

al. 1996). Plant species commonly found in serpentine grasslands at JRBP include native *Stipa*

pulchra, *Elymus glaucus*, *Elymus multisetus, Eschscholzia californica*, and the invasive grass

Bromus hordeaceus (McNaughton 1968; Hobbs and Mooney 1985). The nonnative annual

species that dominate the nonserpentine grasslands but are absent from serpentine grasslands

include *Avena barbata, Avena fatua, Bromus diandrus,* and *Erodium botrys* (McNaughton 1968;

Hobbs and Mooney 1985; Field *et al.* 1996). The serpentine and nonserpentine grasslands at

JRBP share a land use history of livestock grazing that ceased in 1960, when conservation of

these areas became a management priority (Bocek and Reese 1992).

All survey sites were located along the ridgetop at JRBP, where serpentine and

nonserpentine (greenstone) soils are present in discrete adjacent bands (Fig. 1a) (Oze *et al.*

2004). Chemical analyses of the soils on this ridgetop show consistent, significant differences in

serpentine and nonserpentine soil chemistry, with the serpentine soils enriched in heavy metals

and trace elements (Fe, Cr, Ni, Mn, and Co), and impoverished in essential plant nutrients (Streit

et al. 1993, Oze *et al.* 2004, 2008). The relative flatness of the ridgetop ensured that all survey

sites were at roughly the same elevation and had similar slopes, aspects, and water availability

(Oze *et al.* 2008).

Quantification of disease

To assess plant disease burden, we quantified the percentage of living leaf area exhibiting

symptoms of fungal disease in serpentine and nonserpentine populations of *S. pulchra*. We

selected three serpentine grassland sites and three nonserpentine grassland sites such that each

serpentine site was paired with a nearby (~170m away) nonserpentine site (Fig. 1a). Individual

sites within each habitat type were >300m apart and the maximum distance between any two

sites was 1600m (Fig. 1a). Each year, we randomly placed four 5-meter transects at each 40m x

40m site and assessed the infection status of the *S. pulchra* individual nearest each meter mark,

resulting in five individual plants per transect surveyed each year (N = 60 plants per habitat type

per year). The infection status of individual plants was not tracked across years, although some

plants could have been sampled repeatedly by chance. Following the protocol for estimation of

foliar infection severity on grasses described in Mitchell *et al.* 2002 and Spear and Mordecai

2018, a single surveyor visually estimated the percentage of living foliar tissue damaged by

pathogens for six arbitrarily-selected leaves per plant. We used printed field reference guides

showing known amounts of damage on leaves to standardize estimates. Damage was indicated

by spots and lesions which ranged in color from dark brown to pale yellow (Fig. 1b). Senesced

tissue and herbivore damage (commonly in the form of scrapes and holes in the leaf surface) was

excluded from the analyzed leaf area. For damage estimates of 5% or greater, values were

determined to the nearest 5%. Damage levels below 5% were assigned values of 1% or 0.1%.

We used analysis of variance (ANOVA), with individual plants nested within sites, to test

for effects of habitat type, year, habitat type x year, and site on log-transformed mean percent

disease damage on serpentine and nonserpentine *S. pulchra* (N = 36 serpentine and 36

nonserpentine transects). When ANOVA results were statistically significant, we used t-tests for

every possible pairwise combination, with Bonferroni adjustments of p-values to account for

multiple comparisons, to identify the interactions contributing to the significant result.

Isolation and molecular identification of foliar fungal pathogens

To identify the fungal foliar pathogens infecting serpentine and nonserpentine *S. pulchra*

over all years*,* we harvested one segment of symptomatic leaf tissue for culturing and

identification of the putatively causal fungal pathogen(s) from each individual included in the

damage survey with symptoms of disease. This tissue segment came from the first surveyed leaf

with disease symptoms on each plant. We excised, surface sterilized, and plated a 4mm^2 square

of tissue from each sample at the leading edge of infection, to improve the chances of isolating

the causal fungal pathogen, on malt extract agar with 2% chloramphenicol to prevent bacterial

growth. While it is possible that the growing medium used may have led to selective isolation

and sequencing of fungi other than those causing symptoms, our use of small segments of leaf

tissue from the advancing margin of disease was intended to prevent this outcome under the

assumption that symptom-causing organisms should be dominant in visibly diseased tissue and

particularly at its expanding margin. We observed the plates for growth for eight weeks and

isolated morphologically distinct hyphae into axenic culture. Six segments of surface sterilized,

asymptomatic tissue were plated as a control and showed no growth. We extracted and

sequenced genomic DNA from the ITS region for each isolate. Fungal isolation and sequencing

methods with ITS1F and ITS4 primers followed Spear and Mordecai 2018.

We estimated taxonomic placement of fungal isolates by grouping sequences into

operational taxonomic units with 97% sequence similarity (OTUs), a proxy for species

delineation based on the range of intraspecific ITS divergence (O'Brien *et al.* 2005), and then

comparing assembled DNA sequences to the entries in the UNITE fungal database. To prepare
sequences for taxonomic placement, we first trimmed forward and reverse Sanger sequencing
reads to a maximum error rate of 5% and then automatically assembled bidirectional reads into
consensus sequences with a minimum of 20% overlap and 85% sequence similarity in Geneious
(Kearse *et al.* 2012), discarding consensus sequences less than 450 bp in length and/or less than
60% in read quality. We grouped the remaining internal transcribed spacer (ITS) sequences into
OTUs using USEARCH 10.0.240 (Edgar 2010) and estimated the taxonomic placement of each
OTU in mothur v.1.40.5 (http://mothur.org/; Schloss 2009) using the naïve Bayes classification
approach (classify.seqs with options: method='wang', cutoff=0, iters=200). We used a
bootstrapping confidence score of at least 80% to assign a species name and at least 60% to
assign taxonomy at higher ranks.

Analyses of fungal community composition

To characterize foliar fungal community composition in serpentine and nonserpentine *S.*
pulchra populations, we compared the diversity and assessed the similarity of communities of
fungi cultured from the diseased tissue of surveyed plants. To describe fungal community
diversity, we calculated observed taxa richness and Fisher's alpha, a widely used measure of
species richness that is robust to uneven sample sizes (Fisher *et al.* 1943; Magurran 2013);
generated taxa accumulation curves to understand sampling efficacy (Oksanen *et al.* 2016); and
counted the number of fungal isolates in each genus, family, order, and class from each grassland
type. We compared fungal community composition across grassland types and across years
visually, using non-metric multidimensional scaling (NMDS), and statistically, using
permutational multivariate analyses of variance (PERMANOVA) with pairwise Adonis
functions for non-normalized OTU read counts in the R package vegan (Anderson 2001; Arbizu
2016; Oksanen *et al.* 2016). To assess spatial turnover in fungal communities, we included a site
distance factor in the statistical analyses, with comparisons between sites <600m apart
categorized as "near" and comparisons between sites >600m apart categorized as "far" (Fig. 1a).
To account for low culturing success in some transects, we considered the combined isolates
from two nearby transects at each site in each year (N = 18 serpentine and 18 nonserpentine
communities, generated from 36 total transects per habitat type) to be distinct communities for
NMDS visualization.

For ANOVA and PERMANOVA analyses we considered the combined isolates from
each site in each year to be distinct communities (N = 9 serpentine and 9 nonerspentine
communities). When PERMANOVA results were statistically significant, we used pairwise
PERMANOVAs with Bonferroni adjustments to p-values to clarify which fungal communities
were significantly different from one another (Arbizu 2016). For the NMDS and PERMANOVA
analyses, we used the function vegdist with the abundance-based Chao method, accounting for
unsampled species, to create a matrix of pairwise community dissimilarities (Chao *et al.* 2016;
Oksanen *et al.* 2016). We also used the Morisita-Horn index, which is independent of sample
size and diversity, to make pairwise comparisons of similarity within and between the serpentine
and nonserpentine fungal communities from each year (N = 265 isolates, 200 bootstrap
replicates), and within the entire fungal community between years (Wolda 1981; Jost *et al.*
2011). Finally, we used ANOVA to test for effects of habitat type, year, site distance, year x
habitat type, and year x site distance interactions on fungal diversity (Fisher's alpha) and
richness at the site level.

To perform these fungal community analyses, we used the packages BiodiversityR (Kindt
and Coe 2005), bipartite (Dormann *et al.* 2008), fossil (Vavrek 2011), rich (Rossi 2011), SpadeR
(Chao *et al.* 2016), and vegan (Oksanen *et al.* 2016) with custom commands (Gardener 2014) in
R version 3.4.1.

Chemical analyses of leaf tissue

In 2016, we analyzed C and N content of the foliar tissue of the *S. pulchra* plants
surveyed in serpentine and nonserpentine grasslands. Because both leaf age and infection status
might influence leaf chemistry, we collected the youngest fully expanded and entirely
asymptomatic leaf of each individual surveyed in the field (N = 60 serpentine, 58 nonserpentine).
We dried the samples at room temperature for three months, then at 60°C for 48 hours, ground
them to a powder, and then dried them again at 60°C for 24 hours. We measured C and N
content of these samples with a Carlo Erba NA-1500 elemental analyzer using atropine
standards.

To assess the potential for habitat type to mediate plant-pathogen interactions through its
influence on plant tissue chemistry, we related habitat type to leaf tissue chemistry of *in situ S.*
pulchra, and chemistry of these plants to disease burden. We compared the mean percentage of

C and N per dried weight of leaf tissue for surveyed serpentine and nonserpentine plants with

Wilcoxon rank sum tests. We also compared mean C : N ratio, which is indicative of plant health

(Ciompi *et al.* 1996; Brady *et al.* 2005). We used Pearson's product-moment correlation to test

for significant monotonic relationships between the log-transformed mean amount of diseased

tissue observed on individual plants in the field and their foliar C and N content.

All statistical analyses were done in R version 3.4.1. In addition to the R packages

specified in the fungal community analyses methods, we used the packages cowplot (Wilke

2019), data.table (Dowle and Srinivasan 2019), RColorBrewer (Neuwirth 2014), reshape2

(Wickham 2007), tidyverse (Wickham 2017), and xtable (Dahl *et al*. 2019) to manipulate, plot,

and export data and results.

Results

Pathogen damage

Pathogen damage was similarly ubiquitous yet low-burden in serpentine and

nonserpentine grassland habitats. Every plant surveyed exhibited evidence of foliar fungus-

caused disease. The mean percentage of diseased leaf area observed across all years was 1.63 for

serpentine and 1.66 for nonserpentine *S. pulchra,* respectively (Fig. 1c), and the mean proportion

of surveyed leaves with pathogen damage was 0.83 for serpentine and 0.85 for nonserpentine

plants. Year significantly affected log-transformed mean percent damage (ANOVA: df = 2, SS =

9.41, F-value = 5.375, p-value = 7.04×10^{-3}) (Table 1). There were significant differences in

percent damage between was between 2015 and 2016 and between 2015 and 2017 (2015 mean

percent damage = 1.05, 2016 = 1.96, 2017 = 1.94, t-values = -2.66, -3.70, Bonferonni-adjusted p-

values = 2.5×10^{-3}, 8.1×10^{-4}).

Fungal pathogen community

We isolated 267 unique fungal isolates from 258 plated symptomatic tissue pieces with

fungal growth (144 nonserpentine and 114 serpentine tissue pieces from 36 nonserpentine and 34

serpentine transects, out of 360 total tissue pieces). Of the 267 unique fungal isolates, we

successfully sequenced 256 isolates. The sequenced isolates clustered into 30 operational

taxonomic units (OTUs) based on 97% sequence similarity, representing 23 genera, 15 families,

8 orders, and 5 fungal classes. We were unable to identify 18 OTUs to the fungal species level,

and four of these to the genus level due to a lack of close sequence matches in the UNITE

database and/or a lack of taxonomic information for close sequence matches (Table 2). The

taxonomic results indicate that many of the OTUs represent close relatives of known plant

pathogens including *Alternaria, Cladosporium, Drechslera, Phaeosphaeria, Pyrenophora,*

Stemphylium, Ramularia, and *Wojnowiciella* spp. (Table 2) (Kirk *et al.* 2008; Stukenbrock and

McDonald 2008; Havis *et al.* 2015; Marin-Felix *et al.* 2017, 2019). The most common OTU was

represented by 48 isolates and the least common OTUs were represented by one isolate each.

OTUs are hereafter referred to as "species" (O'Brien et al. 2005).

332   Fungal community diversity was similar between serpentine and nonserpentine habitats

and between years (Table 2). Although fungal richness was higher at every taxonomic level in

the serpentine relative to the nonserpentine, these differences were not statistically significant

(Tables 2, S1-S5). Twenty-four fungal species were isolated from serpentine *S. pulchra* (Fisher's

alpha = 9.324, 95% CI = 4.664, 13.983) and 22 from nonserpentine (Fisher's alpha = 7.261, 95%

CI = 3.597, 10.925); 16 of these species (53%) were shared (Fig. 2; Tables 1, 2). Habitat type,

year, and year x habitat type were not significantly related to fungal richness or Fisher's alpha at

the site level (ANOVA: Tables S6, S7). Novel genera were isolated for both community types in

every year of surveying, with only seven of 30 (23%) observed species and six of 23 genera

(26%) isolated in all three years (Table S5). Species accumulation curves did not approach

horizontal asymptotes for either habitat type, suggesting that neither fungal community was fully

described (Fig. S1).

344   Both fungal communities were dominated by a few abundant species (isolated >10

times), but the most abundant species differed between nonserpentine and serpentine habitats:

these were *Alternaria* cf. *infectoria* and *Ramularia* cf. *proteae*, respectively (Table 2; Fig. S2).

The nonserpentine community included five abundant species, representing 62% of the isolates,

while the serpentine community included two abundant species, representing 51% of serpentine

isolates (Table 2; Fig S2). The serpentine community had a higher proportion of rare species (<3

isolates) than the nonserpentine community, 63% versus 36%, respectively. One *Drechslera*

species was abundant in the nonserpentine community and rare in the serpentine (Table 2; Fig.

S2).

353   Habitat type and year significantly affected fungal community composition, based on

two-way PERMANOVA analysis of site-level fungal communities and NMDS visualization of

community similarity (habitat type: F-value = 5.112, R^2 = 0.163, p = 0.003; year: F-value =

5.655, R^2 = 0.360, p = 0.001; Fig. 3; Table 4). Serpentine fungal communities were generally

more similar to one another than they were to nonserpentine fungal communities, and vice versa

(Fig. 3; Tables 2, 3). Community composition differed significantly between 2015 and 2017, and

2016 and 2017, but not between 2015 and 2016 (Table S8). Differences in fungal community

composition between years did not correspond to differences in disease severity between years

(Figure 1c, Table S8).

Interpretation of the significance of variation in values of Morisita-Horn community

overlap that fall between zero (no overlap) and one (complete overlap) is a longstanding

challenge in community ecology (Horn 1966, Rodrigues and Vieira 2010). Qualitatively,

estimates of the Morisita-Horn similarity index for communities within habitat types suggest that

the amount of interannual species turnover is more consistent within serpentine than

nonserpentine communities (Fig. 3, Tables 2, 3). While serpentine community overlap between

years ranged from 0.707 to 0.786, nonserpentine community overlap ranged between 0.295 and

0.882 (Table 3). Within-year variation in species turnover between habitat types was

comparatively moderate, with Morisita-Horn overlap ranging from 0.535 to 0.725 (Table 3).

Leaf tissue chemistry and disease burden

As expected from differences in soil type between habitats, leaf chemistry in plants

occurring on serpentine versus nonserpentine soils differed in mean C and N content. Serpentine

plants had significantly lower mean C and N content than nonserpentine plants (Fig. 4; Table

S9). C:N ratio was similar for plants in both habitat types. Log-transformed mean percent

symptomatic tissue was not significantly associated with percent C, percent N, or C:N ratio in

asymptomatic foliar tissue for serpentine or nonserpentine plants, based on Pearson's product-

moment correlation tests (Fig. 4).

Discussion

Here we show that a diverse suite of foliar fungi was associated with ubiquitous, low-

burden damage on host individuals across growing seasons and habitat types. Habitat type

appeared to shape fungal community composition in this system, but not foliar disease burden. In

contrast, time appeared to shape both fungal community composition and disease burden.

However, interannual variation in pathogen community composition did not correspond to yearly

differences in disease burden (Figs. 1c, 3, Table S8). While C and N content measured in young,

healthy leaves differed between habitat types, variation in leaf chemistry did not explain

variation in infection severity. The results provide some of the first evidence that local growing

conditions structure the pathogen communities infecting a single host species even over

relatively small areas of a few hundred meters, where hosts and pathogens experience similar

climate conditions (Figs. 1a, 3; Table 3). Overall, the findings imply that environmental

heterogeneity within landscapes may both generate barriers to fungal disease spread in wild plant

populations and bolster fungal diversity.

Fungal community composition

The close proximity of the sampled nonserpentine and serpentine plants and the finding

that habitat type, but not distance between sites, predicted fungal community composition

suggest that habitat filtering, rather than dispersal limitation, plays a key role in fungal

community assembly (Table 4). Fungal communities varied between habitats in their

composition, and the stability of these communities over time varied between habitats (Figs. 1c,

2, 3, S2; Tables 1-4). The results imply that differences in environmental conditions—both biotic

and abiotic—beyond climate averages can lead to differences in plant-fungal interactions in

distinct habitat types. In this system, influential habitat characteristics might include differences

in pathogen sharing among hosts in native-dominated serpentine grasslands versus nonnative-

dominated nonserpentine grasslands, and in plant genotypes occurring in different habitats.

Differences in plant nutrient content between habitat types suggest that soil influence on plant

phenotype may also contribute to differences between fungal communities (Fig. 4). In addition to

C and N differences, potted *S. pulchra* plants grown from seed and serpentine soils collected in

these sites had significantly lower Ca and P content and higher Ni and Mg content than similar

plants grown in greenhouse soils, potentially suggesting additional phenotypic differences

between serpentine and nonserpentine plants that may affect pathogen community composition

(Farner *et al.*, unpublished data). Finally, habitat type might impact disease dynamics if differing

water availability between serpentine and nonserpentine soils alters leaf physiology in ways that

affect infection processes. We hypothesize that some or all of these factors influence the

composition of fungal communities and their variation across environments.

Differences in nonserpentine and serpentine fungal communities may result from
differences in the plant communities found on these soils. The presence of additional fungal
taxonomic diversity and the relatively constrained amount of species turnover between years in
the serpentine align with patterns in serpentine versus nonserpentine grassland plant community
diversity and temporal stability (Fig. S2; Tables 1, 3, S1-S5) (Fernandez-Going *et al.* 2012).
Results from previous studies that demonstrate a positive relationship between host diversity and
pathogen diversity indicate that higher plant species diversity in serpentine grasslands might
contribute to higher fungal diversity if native forbs and bunchgrasses that specialize on
serpentine soils share foliar fungal pathogens with *S. pulchra* (Hechinger and Lafferty 2005;
Spear 2017). Research on five common invasive and native grasses (including *S. pulchra*) at
JRBP shows that multi-host foliar fungal pathogens dominate this system, supporting this
hypothesis (Spear and Mordecai 2018).

Differing chemistry between the leaves of nonserpentine and serpentine *S. pulchra* may
influence fungal community assembly through a variety of mechanisms. Fertilization with
nitrogen, which was elevated in nonserpentine compared to serpentine leaf tissue (Fig. 4b),
differentially impacts foliar disease severity depending on host and fungal pathogen identities
(Dordas 2008, Verosoglou *et al.* 2013). The mechanisms by which nitrogen mediates plant-
fungal pathogen interaction outcomes are unknown, but might involve impacts on plant immune
pathways and/or fungal pathogen virulence, growth, and reproduction. Eschen *et al.* propose
fungal interspecific variation in nutritional requirements as an explanation for correlations
between soil and plant nitrogen content and fungal endophyte community composition in
Cirsium arvense (Eschen *et al.* 2010). While our results are qualitatively similar to those of
Eschen *et al.*, greenhouse inoculation experiments with different fungal pathogens and plants of
known tissue chemistries are needed to test this hypothesis.

Importantly, our study design limits our mechanistic understanding of how variation in
plant chemistry and fungal communities between habitats may be related. Measurements of leaf
chemistry taken from young, healthy leaves may not be representative of the chemistry of the
older, larger leaves that both fungi and surveyors may be more likely to encounter. Further, the
low number of sampled fungal communities for which we measured plant chemistry prevents
statistical analysis of relationships between fungal community composition and plant nutrient
content. However, our hypothesis that differences in plant tissue chemistry between serpentine

and nonserpentine habitats shape fungal pathogen presence is supported by previous studies that

showed *Hesperolinon* sp. infection frequency and severity by the fungal rust *Melampsora lini* in

California grasslands depended on plant tissue Ca levels linked to serpentine soil chemistry

(Springer *et al.* 2006; Springer 2009).

Serpentine and nonserpentine fungal communities may also differ due to genetic

divergence between the *S. pulchra* populations found in these soils. The gene-for-gene model of

the evolution of virulence hypothesizes that a given pathogen can infect a given host only if the

pathogen does not possess an avirulence gene recognized by a matching resistance gene in the

host (Flor 1956; Parker and Gilbert 2004). To date, there have been few studies of genetic

differences between serpentine and nonserpentine populations of the same plant species.

However, genetic and chemical analyses of serpentine and nonserpentine populations of the

native forb *Lasthenia californica* at JRBP show that these populations represent two

phylogenetically distinct species, each of which accumulates ions in significantly different

concentrations and has a unique flavonoid profile (Desrochers and Bohm 1993; Chan *et al.* 2002;

Rajakaruna *et al.* 2003). Since many flavonoids have antifungal properties, serpentine and

nonserpentine *S. pulchra* could also contain different genotypes with different potential for

resistance to any given fungus due to divergent resistance gene and/or flavonoid profiles (Parker

and Gilbert 2004; Treutter 2006).

In addition to fungal species turnover between habitat types, interannual turnover

contributed substantially to fungal diversity in this system, with only seven of 30 total species

isolated in all three years (Fig. 3; Tables 1, S5). Interannual variation in these communities may

be driven by a variety of mechanisms, including highly variable yearly precipitation (2015

rainfall = 491 mm; 2016 = 604 mm, 2017 = 985 mm) (Weather Underground). Timing and

amount of precipitation might alter infection dynamics by washing away fungal spores and/or

transmitting spores to new hosts (Madden 1997). Additionally, yearly variation in plant

population sizes, occurrences of different plant genotypes, and plant community composition

may also drive temporal variation in fungal communities (Hobbs and Mooney 1991). Although

few studies track interannual variation in fungal communities on plants, the year-to-year

dissimilarity of those hosted by *S. pulchra* is similar to that observed across a diverse taxonomic

range of plants including ferns, legumes, and shrubs (Olmo-Ruiz and Arnold 2014; Cotton *et al.*

2015; Zhang *et al.* 2016). In this context, our results are consistent with a growing body of

evidence for the general importance of storage effects—in which temporally variable conditions
act as ecological niches that facilitate species diversity—in plant-associated fungal communities
(Warner and Chesson 1985). This finding highlights the necessity of long-term data collection to
accurately characterize plant-fungal interaction networks.

Foliar disease burden

Time, but not habitat type, shaped disease burden, but mean levels of foliar infection
observed were low even at their peak of 1.9% symptomatic leaf area in 2017. This level of
damage is similar to that observed on other grass species in nonserpentine grasslands at JRBP
(Spear and Mordecai 2018), and on taxonomically diverse tropical forest plants in Mexico and
Hawaii (García-Guzmán and Dirzo 2001, Funk and Throop 2010). Recent work in the JRBP
grassland system shows that symptoms in the amounts observed inflict only minimal plant
fitness costs, suggesting that *S. pulchra* is able to effectively resist or suppress damage by foliar
pathogens in both serpentine and nonserpentine grasslands (Spear and Mordecai 2018; Uricchio
et al. 2019). These expectations align with assessments in wheat-fungal pathogen systems that
indicate yield reductions of $\leq$ 1% for each percent of symptomatic leaf area (Milus 1993, Khan *et*
al. 1997). The finding that between-habitat differences in foliar C and N content do not
correspond to differences in infection severity aligns with the finding that infection severity did
not differ between habitats. Additionally, similar C:N ratios suggesting that serpentine and
nonserpentine are equally healthy correspond to the consistently low damage observed across
habitats. However, the limitations described above to interpreting observed relationships between
plant chemistry and fungal community composition apply equally to observed relationships
between plant chemistry and disease severity.

The variation in fungal community composition we observed on a single host species
with habitat type and over time suggests possible mechanisms preventing severe outbreaks of
foliar disease in California grasslands. Transitions between distinct soil types over distances
within the range of fungal spore dispersal may prevent the spread of pathogens that are sensitive
to leaf tissue chemistry. Interannually variable conditions favoring successful infection by
different fungal species each growing season (Table S5), combined with plant senescence during
the dry season each year, suggest temporal barriers to pathogens evolving high virulence on both
nonserpentine and serpentine plants.

Clarifying the specific mechanisms that drive differences in serpentine and nonserpentine

fungal communities will require inoculation experiments to test how plant susceptibility to

infection by different fungal species changes with plant tissue chemistry and plant genetics.

Additionally, characterization of comprehensive plant-pathogen networks including all plant

species co-occurring with *S. pulchra* in serpentine and nonserpentine grasslands will improve our

understanding of the role of the surrounding plant community in structuring fungal pathogen

communities.

Fungal diversity

Compared to other plant species' interactions with both pathogenic and non-pathogenic

fungi, *S. pulchra* appears to host an unusually species-rich community of culturable fungi within

its diseased foliar tissue. Where we isolated 30 fungal species, past studies that described fungal

pathogen diversity for more than 100 different plant hosts consistently found fewer than 20

fungal pathogens per species, and almost all hosts had fewer than ten pathogens (Mitchell and

Power 2003; Hantsch *et al.* 2013; Rottstock *et al.* 2014). The level of fungal diversity we

documented is also high compared to that of culturable fungal endophyte communities associated

with asymptomatic plant tissues: of 35 plant species analyzed by Arnold and colleagues, only

five were found to host 30 or more endopyhtes (Arnold and Lutzoni 2007; Del Olmo-Ruiz and

Arnold 2014). Notably, studies that utilize high-throughput sequencing techniques to observe

both culturable and unculturable endophytic fungi frequently report hundreds of fungal OTUS

colonizing single host species (e.g., Jumpponen and Jones 2009, Zimmerman and Vitousek

2012). However, such results are difficult to compare to our own given the mismatch in the

detection capabilities of the methods used. Nonetheless, the exceptional diversity of the

culturable foliar fungi that inhabit *S. pulchra* is further indicated both by species accumulation

curves that suggest many more species in this community remain to be discovered (Fig. S1) and

by the lack of close sequence matches in genetic databases for some of the species isolated

(Table 2). These findings suggest that the characteristics that make the California Floristic

Province a biodiversity hotspot may extend to the fungal diversity associated with this flora

(Myers *et al.* 2000).

Conclusions

Fungi associated with foliar disease in the native bunchgrass *S. pulchra* are relatively

benign, likely cause minimal fitness impacts (Spear and Mordecai 2018), and contribute
substantially to the overall biodiversity of California grasslands. Variation between habitats
appears to play an important role in supporting the fungal diversity observed here, and may also
reduce disease risk for *S. pulchra*. The high taxonomic diversity and rarity of serpentine fungi
demonstrates that this community is potentially at risk of species loss via host-parasite
coextinction as nonnative annual grasses continue to expand their ranges at the cost of native
plants, and human activities including development and increased C and N deposition contribute
to the loss of serpentine grasslands that already make up <2% of California's surface area
(Huenneke *et al.* 1990; Vallano *et al.* 2012). This work demonstrates that, in addition to >200
endemic plant species, California's serpentine grasslands support unique communities of leaf-
infecting fungi, providing additional motivation for active conservation of native grassland
communities based on the far-reaching loss of biodiversity across trophic levels associated with
their disappearance (Sprent 1992; Dunn *et al.* 2009).

Acknowledgements
We thank Nona Chiariello, Caroline Daws, Joe Wan, Peter Vitousek, Scott Fendorf, Lawrence
Uricchio, Guangchao Li, Douglas Turner, Virginia Walbot, Cody Hamilton, and the Mordecai
and Peay Lab Groups for their help. This work was supported by the Jasper Ridge Kennedy
Endowment, the National Science Foundation (DEB-1518681), Stanford Vice Provost for
Undergraduate Education, and Stanford Bio-X.

References

Abbate JL, Antonovics J. 2014. Elevational disease distribution in a natural plant–pathogen
system: insights from changes across host populations and climate. *Oikos* **123:** 1126–1136.

Ackerly DD, Knight CA, Weiss SB, Barton K, Starmer KP. 2002. Leaf size, specific leaf area
and microhabitat distribution of chaparral woody plants: contrasting patterns in species
level and community level analyses. *Oecologia* **130:** 449–457.

Agrios GN. 2005. *Plant Pathology.* Burlington, MA, USA. Elsevier Academic Press.

Anagnostakis, SL. 1987. Chestnut blight: the classical problem of an introduced pathogen.
Mycologia **79(1):** 23-37.

Anderson MJ. 2001. A new method for non-parametric multivariate analysis of variance.
*Austral **Ecology** **26:** 32–46.

Arbizu PM. 2016. *How can I do PerMANOVA pairwise contrasts in R?*[WWW document]
Researchgate 2016.
URLhttps://www.researchgate.net/post/How_can_I_do_PerMANOVA_pairwise_contrasts_
in_R. [accessed 21 June 2019].

Bálint M, Bartha L, O'Hara RB, Olson MS, Otte J, Pfenninger M, Roberston AL, Tiffin P,
Schmitt I. 2015. Relocation, high-latitude warming and host genetic identity shape the
foliar fungal microbiome of poplars. *Molecular Ecology* **24 (1):** 235-248.

Barge EG, Leopold DR, Peay KG, Newcombe G, Busby PE. 2019. Differentiating spatial
from environmental effects on foliar fungal communities of *Populus trichocarpa. Journal of*
Biogeography **46(9):** 2001-2011.

Bernard F, Sache I, Suffert F, Chelle M. 2013. The development of a foliar fungal pathogen
does react to leaf temperature! *New Phytologist* **198(1):** 232-240.

Borer ET, Laine A, Seabloom EW. 2016. A multiscale approach to plant disease using the
metacommunity concept. *Annual Review of Phytopathology* **54:** 397–418.

Brady KU, Kruckeberg AR, Bradshaw HD. 2005. Evolutionary ecology of plant adaptation to
serpentine soils. *Annual Review of Ecology, Evolution, and Systematics* **36:** 243–266.

Brandhorst TT, Gauthier GM, Stein RA, Klein BS. 2005. Calcium binding by the essential
virulence factor BAD-1 of *Blastomyces dermatitidis. Journal of Biological Chemistry* **280:**
42156–42163.

Burdon JJ, Laine A-L. 2019. *Evolutionary Dynamics of Plant-Pathogen Interactions.*

Cambridge, UK. Cambridge University Press.

Chan R, Baldwin BG, Ornduff R. 2002. Cryptic goldfields: A molecular phylogenetic

reinvestigation of *Lasthenia californica* sensu lato and close relatives (Compositae:

Heliantheae sensu lato). *American Journal of Botany* **89:** 1103–1112.

Chao A, MA KH, Hsieh TC, Chiu C-H. 2016. SpadeR: Species prediction and diversity

estimation with R.

Ciompi S, Gentili E, Guidi L, Soldatini GF. 1996. The effect of nitrogen deficiency on leaf gas

exchange and chlorophyll fluorescence parameters in sunflower. *Plant Science* **118:** 177–

184.

Clarkson JP, Fawcett L, Anthony SG, Young C, Pöggeler S. 2014. A model for *Sclerotinia*

sclerotiorum infection and disease development in lettuce, based on the effects of

temperature, relative humidity and ascospore density. *PLoS ONE* **9(4):** e94049.

Cordier T, Robin C, Capdeveille X, Fabreguettes O, Desprez-Loustau M, Vacher C. 2012.

The composition of phyllosphere fungal assemblages of European beech (*Fagus sylvatica*)

varies significantly along an elevation gradient. *New Phytologist* **196:** 510–519.

Dahl DB, Scott D, Roosen C, Magnusson A, Swinton J. 2019. xtable: Export Tables to LaTeX

or HTML. R package version 1.8-4. https://CRAN.R-project.org/package=xtable

Desrochers AM, Bohm BA. 1993. Flavonoid variation in *Lasthenia californica* (Asteraceae).

Biochemical Systematics and Ecology **21:** 449–453.

Dordas C. 2008. Role of nutrients in controlling plant diseases in sustainable agriculture. A

review. *Agronomy for Sustainable Development.* **28(1):** 33-46.

Eschen R, Hunt S, Mykura C, Gange AC, Sutton BC. 2010. The foliar endophytic community

composition in *Cirsium arvense* is affected by myccorhizal colonization and soil nutrient

content. *Fungal Biology* **114:** 991-998.

Dormann CF, Gruber B, Fruend J. 2008. 2008. Introducing the bipartite package: analysing

ecological networks. *R news* **8:** 8–11.

Dowle M, Srinivasan A. 2019. data.table: Extension of 'data.frame'. R package version 1.12.0.

https://CRAN.R-project.org/package=data.table.

Dunn RR, Harris NC, Colwell L, Koh P, Sodhi NS. 2009. The sixth mass coextinction: are

most endangered species parasites and mutualists? *Proceedings of the Royal Society B:*

Biological Sciences **276:** 3037-3045.

Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*
26(19): 2460-2461.

Eversmyer MG, Kramer CL, Hassan ZM. 1988. Environmental influences on the
establishment of *Puccinia recondita* infection structures. *Plant Disease* **72:** 409-412.

Fernandez-Going BM, Anacker BL, Harrison SP. 2012. Temporal variability in California
grasslands: soil type and species functional traits mediate response to precipitation. *Ecology*
93 (9): 2104 – 2114.

Field CB, Chapin FS, Chiariello NR, Holland EA, Mooney HA. 1996. The Jasper Ridge CO_2
Experiment: Design and Motivation. In: Koch GW, Mooney HA, eds. *Carbon dioxide and*
terrestrial ecosystems. San Diego, CA, USA: Academic Press, 121-145.

Fisher MC, Henk DA, Briggs CJ, Brownstein JS, Madoff LC, McCraw SL, Gurr SJ. 2012.
Emerging fungal threats to animal, plant and ecosystem health. *Nature* **484:** 186–194.

Fisher RA, Corbet AS, Williams CB. 1943. The relation between the number of species and
the number of individuals in a random sample of an animal population. *Journal of Animal*
Ecology **12:** 42–58.

Flor HH. 1956. The complementary genic systems in flax and flax rust. *Advances in Genetics* **8:**
29–54.

Funk JL, Throop HL. 2010. Enemy release and plant invasion: patterns of defensive traits and
leaf damage in Hawaii. *Oecologia* **162(4):** 815-823.

García-Guzmán G, Dirzo R. 2001. Patterns of leaf-pathogen infection in the understory of a
Mexican rain forest: incidence, spatiotemporal variation, and mechanisms of infection.
American Journal of Botany **88(4):** 634-645.

Gardener M. 2014. *Community ecology: analytical methods using R and Excel*. Exeter, UK:
Pelagic Publishing.**Hansen RA, Coleman DC. 1998.** Litter complexity and composition are
determinants of the diversity and species composition of oribatid mites (Acari: Oribatida) in
litterbags. *Applied Soil Ecology* **9:** 17–23.

Giauque H, Hawkes CV. 2016. Historical and current climate drive spatial and temporal
patterns in fungal endophyte diversity. *Fungal Ecology* **20:** 108-114.

Hantsch L, Braun U, Scherer-Lorenzen M, Bruelheid H. 2013. Species richness and species
identity effects on occurrence of foliar fungal pathogens in a tree diversity experiment.
Ecosphere **4(7):** 81.

Harrison SP, Viers JH. 2007. Serpentine Grasslands. In: Stromberg MR, Corbin JD, D'Antonio

CM, eds. *California Grasslands: Ecology and Management.* Berkeley, CA, USA:

University of California Press, 145-155.

Havis ND, Brown JKM, Clemente G, Frei P, Jedryczka M, Kaczmarek J, Kaczmarek M,

Matusinsky P, McGraan GRD, Pereyra S, *et al.* 2015. *Ramularia collo-cygni*—an

emerging pathogen of barley crops. *Phytopathology.* **105(7):** 895–904.

Hechinger RF, Lafferty KD. 2005. Host diversity begets parasite diversity: bird final hosts and

trematodes in snail intermediate hosts. *Proceedings of the Royal Society B: Biological*

Sciences **272:**1059–1066.

Hobbs RJ, Mooney HA. 1985. Community and population dynamics of serpentine annual

grassland in relation to gopher disturbances. *Oecologia* **67:** 342–351.

Hobbs RJ, Mooney HA. 1991. Effects of rainfall variability and gopher disturbance on

serpentine annual grassland dynamics. *Ecology* **72:** 59–68.

Huber L, Gillespie TJ. 1992. Modeling leaf wetness in relation to plant disease epidemiology.

Annual Review of Phytopathology. **30:** 553-577.

Huenneke LF, Hamburg SP, Koide R, Mooney HA, Vitousek PM. 1990. Effects of soil

resources on plant invasion and community structure in Californian serpentine grassland.

Ecology **71:** 478–491.

Jost L, Chao A, Chazdon RL. 2011. Compositional similarity and beta diversity. In: McGill BJ,

Magurran AE, editors. *Biological diversity: frontiers in measurement and assessment.*

Oxford, UK: Oxford University Press, 66-84.

Jumpponen A, Jones KL. 2009. Massively parallel 454 sequencing indicates hyperdiverse

fungal communities in temperate *Quercus macrocarpa* phyllosphere. *New Phytologist*

184(2): 438-448.

Kearse MR, Moir A, Wilson S, Stones-Havas M, Cheung S, Sturrock S, Buxton A, Cooper

S, Markowitz C, Duran T et al. 2012. Geneious Basic: an integrated and extendable

desktop software platform for the organization and analysis of sequence data.

Bioinformatics **28(12):** 1647-1649.

Khan MA, Trevathan LE, Robbins JT. 1997. Quantitative relationship between leaf rust and

wheat yield in Mississippi. *Plant Disease* **81(7):** 769-772.

Kindt R, Coe R. 2005. BiodiversityR. Tree diversity analysis. A manual and software for

common statistical methods for ecological and biodiversity studies. World Agroforestry

Centre (ICRAF), Nairobi.

Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008. *Ainsworth and Bisby's Dictionry of the*

Fungi. Oxon, UK: CAB International.

Lee SC, Lee YH. 1998. Calcium/calmodulin-dependent signaling for appressorium formation in

the plant pathogenic fungus *Magnaporthe grisea. Molecules & Cells* **8:** 698–704.

Levine A, Pennell RI, Alvarez ME, Palmer R, Lamb C. 1996. Calcium-mediated apoptosis in

a plant hypersensitive disease resistance response. *Current Biology* **6:** 427–437.

Loehrer M, Schaffrath U. 2011. Soybean rust − meet a prominent challenge in soybean

cultivation. In *Ng T-B*, editor. *Soybean – Biochemistry, Chemistry, and Physiology.* London,

UK: InTech, 83–100 .

Madden, LV. 1997. No effects of rain on splash dispersal of fungal pathogens. *Canadian*

Journal of Plant Pathology **19:** 225–230.

Magarey RD, Sutton TB, Thayer CL. 2005. A simple generic infection model for foliar fungal

plant pathogens. *Phytopathology* **95(1):** 92-100.

Magurran AE. 2013. *Measuring Biological Diversity*. Malden, MA, USA: Blackwell Science

Ltd.

Marin-Felix Y, Groenewald JZ, Cai L, Chen Q, Marincowitz S, Barnes I, Bensch K, Braun

U, Camporesi E, Damm U, *et al.* **2017.** Genera of phytopathogenic fungi: GOPHY 1.

Studies in Mycology **86:** 99–216.

Marin-Felix Y, Hernández-Restrepo M, Wingield MJ, Akulov A, Carnegie AJ,

Cheewangkoon R, Gramaje D, Groenewald JZ, Guarnaccia V, Halleen F, *et al.* **2019.**

Genera of phytopathogenic: GOPHY 2. *Studies in Mycology* **92:** 47–133.

McNaughton SJ. 1968. Structure and function in California grasslands. *Ecology* **49:** 962–972.

Millberg H, Boberg J. 2015. Changes in fungal community of Scots pine (*Pinus sylvestris*)

needles along a latitudinal gradient in Sweden. *Fungal Ecology* **17:** 126-139.

Milus EA. 1993. Effects of leaf rust and septoria leaf blotch on yield and test weight of wheat in

Arkansas. *Plant Disease* **78(1):** 55-59.

Mitchell CE, Power AO. 2003. Release of invasive plants from fungal and viral pathogens.

Nature **421:** 625–627.

Mitchell CE, Tilman D, Groth JV. 2002. Effects of grassland plant species diversity,

abundance, and composition on foliar fungal disease. *Ecology* **83(6):** 1713-1726.

Moss MA. 1987. Environmental conditions conducive to infection of Ryegrass by *Pyricularia grisea. Phytopathology* **77(6):** 863-866.

Nazareno ES, Li F, Smith M, Park RF, Kianian SF, Figureoa M. 2018. *Puccinia coronata* f. sp. *avenae*: a threat to global oat production. *Molecular Plant Pathology* **19(5):** 1047–1060.

Neuwirth, E. 2014. RColorBrewer: ColorBrewer Palettes. R package version 1.1-2. https://CRAN.R-project.org/package=RColorBrewer

O'Brien HE, Parrent JL, Jackson JA, Moncalvo J, Vilgalys R. 2005. Fungal community analysis by large-scale sequencing of environmental samples. *Applied and Environmental Microbiology* **71:** 5544–5550.

Oksanen J, Kindt R, Legendre P, O'Hara B. 2016. Vegan: community ecology package. R package.

Oze C, Fendorf S, Bird DK, Coleman RG. 2004. Chromium geochemistry in serpentinized ultramafic rocks and serpentine soils from the Franciscan complex of California. *American Journal of Science* **304:** 67–101.

Oze C, Skinner C, Schroth AW, Coleman RG. 2008. Growing up green on serpentine soils: biogeochemistry of serpentine vegetation in the Central Coast Range of California. *Applied Geochemistry* **23:** 3391–3403.

Parker IM, Gilbert GS. 2004. The evolutionary ecology of novel plant-pathogen interactions. *Annual Review of Ecology, Evolution, and Systematics* **35:** 675–700.

Parker IM, Saunders M, Bontrager M, Weitz AP, Hendricks R, Magarey R, Suiter K, Gilbert GS. 2015. Phylogenetic structure and host abundance drive disease pressure in communities. *Nature* **520:** 542–544.

Pineda A, Zheng S-J, van Loon JJA, Pieterse CMJ, Dicke M. 2010. Helping plants to deal with insects: the role of beneficial soil-borne microbes. *Trends in Plant Science* **15:** 507-514.

Rajakaruna N, Baldwin BG, Chan R, Desrochers AM, Bohm BA, Whitton J. 2003. Edaphic races and phylogenetic taxa in the *Lasthenia californica* complex (Asteraceae: Heliantheae): an hypothesis of parallel evolution. *Molecular Ecology* **12:** 1675–1679.

Rizzo DM, Garbelotto M. 2003. Sudden oak death: endangering California nad Oregon forest ecosystems. *Frontiers in Ecology and the Environment* **1(5):** 197-204.

Romeis T, Ludwig AA, Martin R, Jones JDG. 2001. Calcium-dependent protein kinases play
an essential role in a plant defence response. *EMBO Journal* **20:** 5556–5567.

Rossi J-P. 2011. rich: an R package to analyse species richness. *Diversity* **3:** 112–120.

Rottstock T, Joshi J, Kummer V, Fischer M. 2014. Higher plant diversity promotes higher
diversity of fungal pathogens, while it decreases pathogen infection per plant. *Ecology* **95(7):**
1907-1917.

Seabloom EW, Borer ET, Gross K, Kendig AE, Lacroix C, Mitchell CE, Mordecai EA,
Power AG. 2015. The community ecology of pathogens: coinfection, coexistence and
community composition. *Ecology Letters* **18:** 401–415.

Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Sahl JW. 2009.
Introducing mothur: open-source, platform-independent, community-supported software for
describing and comparing microbial communities. *Applied and Environmental*
Microbiology **75(23):** 7537-7541.

Shaw BD, Hoch HC. 2000. Ca^{2+} regulation of *Phyllosticta ampelicida* pycnidiospore
germination and appressorium formation. *Fungal Genetics and Biology* **31:** 43–53.

Shaw MW. 1986. Effects of temperature and leaf wetness on *Pyrenophera teres* growing on
barley cv. Sonja. *Plant Pathology* **35(3):** 294-309.

Snoeijers SS, Pérez-García A, Joosten MHAJ, De Wit PJGM. 2000. The effect of nitrogen on
disease development and gene expression in bacterial and fungal plant pathogens. *Journal*
of Plant Pathology **106:** 493-506.

Solomon PS, Tan KC, Oliver RP. 2004. The nutrient supply of pathogneic fungi; a fertile field
for study. *Molecular Plant Pathology* **4(3):** 203-210.

Spear ER. 2017. Phylogenetic relationships and spatial distributions of putative fungal
pathogens of seedlings across a rainfall gradient in Panama. *Fungal Ecology* **26:** 65–73.

Spear ER, Mordecai EA. 2018. Foliar pathogens do not stabilize coexistence of competing
species in a California grassland. *Ecology* **99(10):** 2250–2259.

Sprent JFA. 1992. Parasites lost? *International Journal for Parasitology* **22:** 139–151.

Springer YP. 2009. Do extreme environments provide a refuge from pathogens? A phylogenetic
test using serpentine flax. *American Journal of Botany* **96:** 2010–2021.

Springer YP, Hardcastle BA, Gilbert GS. 2006. Soil calcium and plant disease in serpentine
ecosystems: a test of the pathogen refuge hypothesis. *Oecologia* **151:** 10–21.

Stukenbrock EH, McDonald BA. 2008. The origins of plant pathogens in agro-ecosystems.
Annual Review of Phytopathology. **46:** 75–100.

Telfer S, Lambin X, Birtles R, Beldomenico P, Burthe S, Paterson S, Begon M. 2010.
Species interactions in a parasite community drive infection risk in a wildlife population.
Science **330:** 243–247.

Treutter D. 2006. Significance of flavonoids in plant resistance: a review. *Environmental*
Chemistry Letters **4:** 147–157.

Uhm K-H, Ahn I-P, Kim S, Lee Y-H. 2003. Calcium/calmodulin-dependent signaling for
prepenetration development in *Colletotrichum gloeosporioides*. *Phytopathology* **93:** 82–87.

UNITE Community. 2017a. Full mothur UNITE+INSD dataset 2. Version
01.12.2017. https://doi.org/10.15156/BIO/587480

UNITE Community. 2017b. UNITE general FASTA release. Version 01.12.2017.
https://doi.org/10.15156/BIO/587475

Uricchio LH, Daws SC, Spear ER, Mordecai EA. 2019. Exotic species dominate due to niche
overlap in a complex grassland community. *The American Naturalist* **193(2):** 213-226.

Vallano DM, Selmants PC, Zavaleta ES. 2012. Simulated nitrogen deposition enhances the
performance of an exotic grass relative to native serpentine grassland competitors. *Plant*
Ecology **213:** 1015–1026.

Vavrek MJ. 2011. Fossil: palaeoecological and palaeogeographical analysis tools.
Palaeontologia Electronica **14**.

Velásquez AC, Castroverde CDM, He SY. 2018. Plant-pathogen warfare under changing
climate conditions. *Current Biology* **28(10):** R619-R634.

Verosoglou SD, Barto EK, Menexes G, Rillig MC. 2013. Fertilization affects severity of
disease caused by fungal plant pathogens. *Plant Pathology* **62(5):** 961-969.

Warner RR, Chesson PL. 1985. Coexistence mediated by recruitment fluctuations: a field
guide to the storage effect. *The American Naturalist* **125:** 769–787.

Weather Underground. 2018. Jasper Ridge Biological Preserve KCAWOODS19.
https://www.wunderground.com/personal-weather-station/dashboard?ID=KCAWOODS19.

Wickham H. 2007. Reshaping data with the reshape package. *Journal of Statistical Software* **21(**
12): 1-20.

Wickham H. 2017. tidyverse: Easily Install and Load the 'Tidyverse'. R package version 1.2.1.

https://CRAN.R-project.org/package=tidyverse

Wilke CO. 2019. cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2'. R

package version 0.9.4. https://CRAN.R-project.org/package=cowplot

Wolda H. 1981. Similarity indices, sample size and diversity. *Oecologia* **50:** 296–302.

Zimmerman NB, Vitousek PM. 2012. Fungal endophyte communities reflect environmental

structuring across a Hawaiian landscape. *Proceedings of the National Academy of Sciences*

109: 13022-13027.

Figure legends

Figure 1. Pathogen damage on *Stipa pulchra* was consistently low across three years and

both grassland types. Panel **(a)** maps the locations of surveyed nonserpentine sites (blue) and

serpentine sites (green) along the ridgetop at JRBP. Panel **(b)** shows an example of fungal-

caused pathogen foliar damage on *S. pulchra.* Panel **(c)** shows mean percentage (bars) and two

standard errors (error bars) of diseased leaf area for surveyed serpentine and nonserpentine *S.*

pulchra, by year.

Figure 2. Pathogens isolated from *Stipa pulchra* growing in nonserpentine and serpentine

grasslands. Bipartite network showing species interactions between *S. pulchra* plants (left)

growing in serpentine grasslands (green) and nonserpentine grasslands (blue) and fungal species

cultured from symptomatic leaf tissue (white bars on the right). The length of each white bar

correlates with the number of isolates in the operational species unit (based on 97% sequence

similarity). The thickness of lines connecting left and right bars represents the number of times a

particular fungal species was isolated from *S. pulchra* in a particular grassland type. The

nonserpentine fungi came from 15 genera, 8 families, 4 orders, and 3 classes; the serpentine

fungi came from 20 genera, 14 families, 7 orders, and 5 classes (Tables 1, S1-S5).

Figure 3. Distinct fungal communities were isolated from *Stipa pulchra* growing in nearby

serpentine and nonserpentine sites across years. Non-metric multidimensional scaling

visualization of serpentine and nonserpentine fungal community similarity. Each point

corresponds to the combined fungal community of two transects at the same site in the same

year. Serpentine communities are shown in green and nonserpentine communities are shown in

blue. Circles, triangles, and squares represent fungal communities sampled in 2015, 2016, and

2017, respectively. Ellipses enclose 95% confidence intervals for ordination of serpentine and

nonserpentine communities. Community dissimilarity increases with the distance between

points. PERMANOVA analysis showed a significant effect of soil type on fungal community

composition ($F = 3.623$, $R^2 = 0.0618$, $p = 0.001$).

Figure 4. *Stipa pulchra* foliar C and N content differs between habitat types, but does not

predict mean pathogen damage. Foliar percent C (Panel **(a)**), percent N (Panel **(b)**, and C:N

ratio (Panels **(c)** versus mean percentage of diseased leaf area for serpentine and nonserpentine *S.*

pulchra plants.

Figures

Figure 1.

Alternaria cf. eureka

Alternaria cf. infectoria

Cladosporium sp.

Drechslera cf. fugax

Drechslera cf. nobleae
Drechslera sp.

Keissleriella sp.

Neoascochyta sp.
Parastagonospora sp. 1

Pyrenophera cf. avenae

Pyrenophera cf. lolii

Ramularia cf. proteae

Stemphylium sp.

unknown Phaeosphaeriaceae 2
unknown Pleosporaceae

884   **Figure 2.**

Figure 3.

901   **Figure 4.**

  Tables

905 **Table 1. Pathogen damage on *Stipa pulchra* varied significantly by year, but not habitat**

906 **type.**

Variables	df	SS	MSE	F value	p-value
habitat type	1	5.8	5.81	2.63	1.05×10^{-1}
year	2	31.4	15.68	7.10	9.53×10^{-4}*
site	4	20.2	5.05	2.29	5.98×10^{-2}
habitat type x year	2	9.9	4.96	2.25	1.07×10^{-1}

ANOVA results for log-transformed transect-level measurements of percentage of symptomatic

living tissue on serpentine and nonserpentine *S. pulchra* (N = 3 sites per habitat type per year,

with 12 transects per site per year, for a total of 36 serpentine and 36 nonserpentine transects

over all three years). The first column lists the factor(s) tested; the second column degrees of

freedom (df); the third column the sum of squares (SS); the fourth column the mean squared

error (MSE); the fifth column the F-value; and the 6th column the p-value. P-values < 0.05 are

marked with an asterisk.

Table 2. Fungal species isolated from symptomatic *Stipa pulchra* leaves were unevenly

distributed across habitat types. Each row represents one of the species isolated. The columns

indicate the isolation frequency for each species in each soil type.

Species	Nonserpentine	Serpentine
Alternaria cf. *infectoria*	26	21
Keissleriella sp.	19	8
Drechslera cf. *fugax*	17	1
Stemphylium sp.	15	8
Ramularia cf. *proteae*	11	37
Drechslera sp.	9	1
Neoascochyta sp.	7	3
Cladosporium sp.	6	3
Alternaria cf. *eureka*	5	2
Parastagonospora cf. *avenae*	5	0
Pyrenophera cf. *chaeotmioides*	5	2
Pyrenophera cf. *lolii*	4	3
Phaeosphaerieaceae sp. 2	3	5
Pleosporaceae sp.	3	2
Drechslera cf. *nobleae*	1	5
Epicoccum sp.	1	0
Heyderia sp.	1	1
Parastagonospora cf. *cumpignensis*	1	0
Parastagonospora sp.	1	0
Phaeosphaeria sp.	1	2
Podospora sp.	1	0
Phaeosphaerieaceae sp. 1	1	0
Chalastospora cf. *gossypii*	0	1
Hyphodiscus cf. *brevicollaris*	0	1
Nemania cf. *serpens*	0	1
Phaeophlebiopsis sp.	0	1
Plectania sp.	0	2
Pseudoseptoria sp.	0	1
Helotiales sp.	0	1
Wojnowiciella sp.	0	1

922 **Table 3. Fungal communities were distinct on *Stipa pulchra* plants growing in serpentine**

923 **versus nonserpentine grasslands, across all years.**

Fungal community pair	Fisher's alpha	Shared species	Morisita-Horn Similarity Index
All years serpentine (24)	9.324	16	0.675
All years nonserpentine (22)	7.261	(53.0%)	
2015 serpentine (11)	7.955	5	0.535
2015 nonserpentine (12)	5.205	(27.8%)	
2016 serpentine (16)	7.897	11	0.766
2016 nonserpentine (13)	5.861	(61.1%)	
2017 serpentine (12)	6.784	8	0.725
2017 nonserpentine (12)	5.136	(50.0%)	
2015 serpentine (11)	7.955	7	0.707
2016 serpentine (16)	7.897	(35.0%)	
2015 serpentine (11)	7.955	5	0.721
2017 serpentine (12)	6.784	(27.8%)	
2016 serpentine (16)	7.897	7	0.786
2017 serpentine (12)	6.784	(33.3%)	
2015 nonserpentine (12)	5.205	6	0.668
2016 nonserpentine (13)	5.861	(31.5%)	
2015 nonserpentine (12)	5.205	5	0.295
2017 nonserpentine (12)	5.136	(26.3%)	
2016 nonserpentine (13)	5.861	7	0.882
2017 nonserpentine (12)	5.136	(38.9%)	

Fungal community similarity by habitat type and year. The leftmost column lists the fungal

communities considered, with the number of species in each community in parentheses. The

second column lists Fisher's alpha for each community. The third column shows the number of

shared species and, in parentheses, the percentage of total observed species that were shared for

each pair of communities. The rightmost column lists the estimated Morisita-Horn community

overlap value, based on absolute species abundances, for each pair of communities. The

Morisita-Horn index ranges from 0 to 1, with 1 indicating complete overlap.

Table 4. PERMANOVA results show significant effects of both habitat type and year on

fungal foliar community composition (N = 18 serpentine and 18 nonserpentine sites).

	df	SS	MSE	F-Value	p-value
habitat type	1	0.5700	0.56997	5.1119	0.003*
year	2	1.2611	0.63055	5.6552	0.001*
distance	1	0.0767	0.07667	0.5850	0.675
habitat type x year	2	0.3339	0.16697	1.4975	0.224
habitat type x distance	1	0.0515	0.05147	0.3927	0.815
distance x year	2	0.1697	0.08485	0.6473	0.734

935 From left to right, columns list factors tested, degrees of freedom (df), sum of squares (SS), mean

936 squared error (MSE), F-value, and p-value. Statistically significant p-values (p < 0.05) are

937 marked with an asterisk. The distance factor compares communities at near (<600m) vs. far

938 (>600m) distances from one another.

Supporting Information

Fig. S1 Species accumulation curves for serpentine (green) and nonserpentine (blue) fungal

communities isolated from *Stipa pulchra* leaves.

Fig. S2 Fungal species abundance by rank for serpentine (green) and nonserpentine (blue) fungal

communities on *Stipa pulchra* leaves.

Table S1 Genus counts for serpentine and nonserpentine fungal communities isolated from *Stipa*

pulchra leaves.

Table S2 Family counts for nonserpentine and serpentine fungal communities isolated from

Stipa pulchra leaves.

Table S3 Order counts for nonserpentine and serpentine fungal communities isolated from *Stipa*

pulchra leaves.

Table S4 Class counts for nonserpentine and serpentine fungal communities isolated from *Stipa*

pulchra leaves.

Table S5 Presence and absence of genera in nonserpentine and serpentine fungal communities

isolated from *Stipa pulchra* leaves, by year.

Table S6 ANOVA table for effects of habitat type, year, and habitat type x year interactions on

site-level fungal community diversity in terms of Fisher's alpha.

Table S7 ANOVA table for effects of habitat type, year, and habitat type x year interactions on

site fungal species richness.

Table S8 Pairwise PERMANOVA table for effects of year on fungal community composition.

Table S9 Results of chemical analysis of serpentine and nonserpentine *Stipa pulchra* foliar

tissue.

985   **Supporting Figures and Tables**

987   **Figure S1. Species accumulation curves for serpentine (green) and nonserpentine (blue)**

988   **fungal communities isolated from *Stipa pulchra* leaves.** The cumulative number of unique

989   fungal OTUs (based on 97% sequence similarity; y-axis) is plotted against the number of *S.*

990   *pulchra* leaf tissue pieces sampled (x-axis).

994
995 **Figure S2. Fungal species abundance by rank for serpentine (green) and nonserpentine**

996 **(blue) fungal communities on *Stipa pulchra* leaves.** Each point represents the number of times

997 a single species was isolated from *S. pulchra* leaves in each habitat type.

999 **Table S1.** Genus counts for serpentine and nonserpentine fungal communities isolated from

1000 *Stipa pulchra* leaves.

Genus	Nonserpentine	Serpentine
Alternaria	31	23
Drechslera	27	7
Keissleriella	19	8
Stemphylium	15	8
Ramularia	11	37
Pyrenophera	9	5
Neoascochyta	7	3
Parastagonospora	7	0
Cladosporium	6	3
Phaeosphaerieaceae	4	5
unknown Pleosporaceae	3	2
Epicoccum	1	0
Heyderia	1	1
Phaeosphaeria	1	2
Podospora	1	0
Chalastospora	0	1
Hyphodiscus	0	1
Nemania	0	1
Phaeophlebiopsis	0	1
Plectania	0	2
Pseudoseptoria	0	1
unkown Helotiales	0	1
Wojnowiciella	0	1

1001

1002 Numbers in each column represent the number of times a genus was isolated from *S. pulchra*

1003 leaves in each habitat type.

1004

 Table S2. Family counts for nonserpentine and serpentine fungal communities isolated from

1006 *Stipa pulchra* leaves.

Family	Nonserpentine	Serpentine
Pleosporaceae	85	45
Lentitheciaceae	19	8
Phaeosphaeriaceae	12	8
Mycosphaerellaceae	11	37
Didymellaceae	8	3
Cladosporiaceae	6	3
Helotiaceae	1	1
Lasiosphaeriaceae	1	0
Dothioraceae	0	1
Hyaloscyphaceae	0	1
Meruliaceae	0	1
Pleosporales *incertae sedis*	0	1
Sacrosomataceae	0	2
unknown Helotiales	0	1
Xylariaceae	0	1

1007

1008 Numbers in each column represent the number of times a family was isolated from *S. pulchra*

1009 leaves in each habitat type.

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021   **Table S3**. Order counts for nonserpentine and serpentine fungal communities isolated from *Stipa*

1022   *pulchra* leaves.

Order	Nonserpentine	Serpentine
Pleosporales	124	65
Capnodiales	17	40
Helotiales	1	3
Sordariales	1	0
Dothideales	0	1
Pezizales	0	2
Polyporales	0	1
Xylariales	0	1

1023

1024   Numbers in each column represent the number of times an order was isolated from *S. pulchra*

1025   leaves in each habitat type.

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041   **Table S4.** Class counts for nonserpentine and serpentine fungal communities isolated from *Stipa*

1042   *pulchra* leaves.

Class	Nonserpentine	Serpentine
Dothideomycetes	141	106
Leotiomycetes	1	3
Sordariomycetes	1	1
Agaricomycetes	0	1
Pezizomycetes	0	2

1043

1044   Numbers in each column represent the number of times a class was isolated from *S. pulchra*

1045   leaves in each habitat type.

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062   **Table S5.** Presence (P, shaded) or absence (A, unshaded) of genera in nonserpentine and

1063   serpentine fungal communities isolated from *Stipa pulchra* leaves, by year.

1064

Genus	Nonserpentine 2015	Nonserpentine 2016	Nonserpentine 2017	Serpentine 2015	Serpentine 2016	Serpentine 2017
Alternaria	P	P	P	P	P	P
Chalastospora	A	A	A	P	A	A
Cladosporium	P	A	A	P	A	P
Drechslera	P	P	P	P	P	P
Epicoccum	A	A	P	A	A	A
Heyderia	A	A	P	A	A	P
Hyphodiscus	A	A	A	A	P	A
Keissleriella	P	P	P	P	P	P
Nemania	A	A	A	A	A	P
Neoascochyta	A	P	P	A	P	P
Parastagonospora	P	A	P	A	A	A
Phaeophlebiopsis	A	A	A	A	P	A
Phaeosphaeria	A	P	A	P	P	A
Plectania	A	P	A	A	A	P
Podospora	A	P	A	A	A	A
Pseudoseptoria	A	A	A	A	P	A
Pyrenophera	P	P	P	P	P	P
Ramularia	P	P	A	P	P	P
Stemphylium	A	P	P	P	A	P
unknown Phaeosphaerieaceae	P	P	A	A	P	A
unknown Pleosporaceae	A	P	P	A	P	P
unkown Helotiales	A	A	A	A	P	A
Wojnowiciella	A	A	A	P	A	A

1065
1066
1067
1068

1069   **Table S6.** ANOVA table for effects of habitat type, year, and habitat type x year interactions on

1070   site-level fungal community diversity in terms of Fisher's alpha.

	df	SS	MSE	F value	p-value
Habitat type	1	0.10	0.10	0.010	0.92
Year	2	3.63	1.82	0.18	0.84
Habitat type x Year	2	17.35	8.68	0.86	0.45

1071

1072   Columns, from left to right, list the variable tested, the degrees of freedom, the sum of squares,

1073   the mean squared error, the F value, and the p-value.

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096 **Table S7.** ANOVA table for effects of habitat type, year, and habitat type x year interactions on

1097 site fungal species richness.

	df	SS	MS	F value	p-value
Habitat type	1	6.72	6.72	1.07	0.32
Year	2	3.11	1.56	0.25	0.78
Habitat type x year	2	8.44	4.22	0.67	0.53

1098

1099 Columns, from left to right, list the variable tested, the degrees of freedom, the sum of squares,

1100 the mean squared error, the F value, and the p-value.

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122 **Table S8.** Pairwise PERMANOVA table for effects of year on fungal community composition.

1123 From left to right, columns list the pair of years being compared, the F-value, R^2, and the

1124 Bonferroni-adjusted p-value. P-values < 0.05 are marked with an asterisk.

Pair	F-value	R^2	Bonferroni p-value
2015 vs 2016	2.616	0.207	0.114
2015 vs 2017	5.983	0.374	0.009*
2016 vs 2017	4.397	0.304	0.018*

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148 **Table S9.** Results of chemical analysis of serpentine and nonserpentine *S. pulchra* foliar tissue.

1149 The Serpentine and Nonserpentine columns report elemental content in percentage of leaf tissue

1150 by dried weight. The standard error of each mean is listed in parentheses. W, in the third column,

1151 is the Wilcoxon-rank sum test statistic comparing serpentine and nonserpentine mean values.

1152 Asterisks denote statistically significant p-values ($p < 0.05$).

1153

Element	Serpentine		Nonserpentine		W	p-value	
C	38.97	(3.44×10^{-1})	41.48	(2.35×10^{-1})	2713	1.65×10^{-7}	*
N	1.44	(4.96×10^{-2})	1.61	(3.84×10^{-2})	2346	1.10×10^{-3}	*
C:N	28.72	(8.74×10^{-1})	26.54	(6.18×10^{-1})	1398	6.62×10^{-2}	

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175