Habitat type and interannual variation shape unique fungal pathogen communities on a California native bunchgrass

Johannah E. Farner[1], Erin R. Spear[1,2], Erin A. Mordecai[1]

[1]Biology Department, Stanford University, Stanford, California 94305 USA; [2]Smithsonian Tropical Research Institute, Panama City, Panama, Republic of Panama

Author for correspondence:

Johannah Farner

E-mail: jfarner@stanford.edu

Phone: 1 (805) 317 – 5432

Abstract

The role of infectious disease in regulating host populations is increasingly recognized, but how environmental conditions affect pathogen communities and infection levels remains poorly understood. Over three years, we compared foliar disease burden, fungal pathogen community composition, and foliar chemistry in the perennial bunchgrass *Stipa pulchra* occurring in adjacent serpentine and nonserpentine grassland habitats with distinct soil types and plant communities. We found that serpentine and nonserpentine *S. pulchra* experienced consistent, low disease pressure associated with distinct fungal pathogen communities with high interannual species turnover. Additionally, plant chemistry differed with habitat type. The results indicate that this species experiences minimal foliar disease associated with diverse fungal communities that are structured across landscapes by spatially and temporally variable conditions. Distinct fungal communities associated with different growing conditions may shield *S. pulchra* from large disease outbreaks, contributing to the low disease burden observed on this and other Mediterranean grassland species.

Introduction

How environmental heterogeneity affects species interactions, and ultimately population growth rates and structures, remains a fundamental ecological question. Infectious disease, which arises from interactions between hosts, parasites, and the environment, is particularly likely to vary across populations in differing environments. Biotic factors, such as the phylogenetic structure of the host community, and abiotic factors, such as elevation, have been associated with the presence of specific diseases and the severity of their impacts on their hosts (Abbate and Antonovics 2014; Parker *et al.* 2015). Additionally, mounting evidence suggests that specific characteristics of pathogen community structures mediate the outcomes of host – pathogen interactions. The identities and abundances of the pathogen species present can shape infection dynamics both through pathogen – pathogen interactions, which can range from facilitative to antagonistic depending on the community context, and because pathogen species vary in their transmissibility, virulence, and sensitivity to different environmental conditions (Agrios, 2005; Seabloom *et al.* 2015; Borer *et al.* 2016; Dallas et al. 2019).

Separate observations of abiotic and biotic impacts on infection outcomes combine to highlight the importance of understanding interactions between environmental variation, pathogen community composition, and pathogen pressure on host populations. For plant hosts, identifying the conditions that place populations at high risk for severe disease outbreaks and applying preventative measures may prevent local species extinctions and improve crop yields. While studies have frequently explored the interactions of single host – pathogen pairs under differing abiotic conditions (e.g., Abbate and Antonovics 2014), both the effect of varying pathogen community composition on disease burden and the relationships between habitat characteristics and pathogen community structures remain largely unexplored in natural systems

(but see Rottstock *et al.* 2014 and Spear 2017).

Fungal pathogens are particularly important to study in this context because they cause the majority of known plant diseases, have an increasing rate of emergence, and pose a substantial threat to plant health and population persistence in natural and agricultural settings (Agrios 2005; Fisher *et al.* 2012). Dramatic diebacks of North American chestnut trees and oaks caused by fungal pathogens (Anagnostakis 1987; Rizzo and Garbelotto 2003) and economically devastating crop losses caused by fungal rusts shared among wild and domesticated plants (Loehrer and Schaffrath 2011; Nazareno *et al.* 2018) demonstrate this threat. These examples indicate that understanding the communities of fungal pathogens that infect wild plants is important for developing effective protection strategies for both natural biodiversity and human food supplies. However, we lack basic information about the identities, diversity, and effects of the fungal pathogens that infect most plant species (Burdon and Laine 2019). Therefore, there is a pressing need for systematic surveys of these pathogen communities and their variation across natural habitats.

Evidence that abiotic and biotic factors drive both natural fungal endophyte community assembly in healthy leaves and fungal foliar disease dynamics in agricultural systems suggests that different habitats are likely to affect fungal foliar pathogen community composition and disease burden (Agrios 2005, Arnold 2007, Burdon and Laine 2019). For example, the repeated finding that foliar endophyte community structures change with temperature, humidity, and rainfall (Cordier *et al.* 2012, Zimmerman and Vitousek 2012, Bálint *et al.*, 2015, Millberg *et al.* 2015, Giauque and Hawkes 2016, Barge *et al.*, 2019) suggests that climate is likely to also shape fungal pathogen communities. Evidence from the plant pathology literature supports this expectation: for foliar fungal pathogens of wheat, barley, and lettuce, among others, infection

dynamics are closely linked to temperature and moisture availability (Shaw 1986, Moss and Trevathan 1987, Eversmyer *et al.* 1988, Huber and Gillespie 1992, Magarey et al. 2005, Bernard *et al.* 2013, Clarkson *et al.* 2014, Velásquez *et al.* 2018). Timing and amount of precipitation can alter infection dynamics by washing away fungal spores and/or transmitting spores to new hosts (Madden 1997).

Additionally, local habitat characteristics, such as soil and plant physical and chemical properties, plant community composition, and plant genetics, can determine the availability and quality of susceptible and competent hosts for different fungal pathogen species (Snoeijers *et al.* 2000, Mitchell *et al.* 2002, Solomon *et al.* 2003, Bolton *et al.* 2008, Rottstock et al. 2014, Velásquez *et al.* 2018). Differences in soil nutrient content, plant nutrient content, and plant genetic backgrounds can influence the identities and abundances of fungal endophytes that inhabit the leaves of individual host species (Eschen *et al.* 2010, Busby *et al.* 2014). However, the short one- to two-year lengths of most observation periods hamper a more general understanding of the factors that shape foliar endophyte communities by generating snapshots of fungal community composition absent the context needed to determine whether observed patterns are broadly representative (but see Giauque and Hawkes 2016). Extensive work in agricultural systems wherein nitrogen fertilization and plant genetic diversity are both associated with disease prevalence and severity support the likely relevance of local habitat factors to fungal pathogens (Zhu *et al.* 2000, Solomon *et al.* 2003, Verosoglou *et al.* 2013).

Here, we examine the effects of habitat type on the burden and species composition of natural fungal plant pathogens infecting a single host species over a span of three years. Specifically, we take advantage of the natural occurrence of the California native perennial bunchgrass *Stipa pulchra* in two unique, adjacent grassland types to understand the extent to

which habitat type and time influence the landscape of disease that a single plant species encounters, including foliar fungal pathogen community composition and disease severity. Stark differences in soil properties and plant community structures between serpentine and nonserpentine greenstone (hereafter, nonserpentine) grassland habitats result in a suite of environmental differences that may influence fungal pathogen infection, reproduction, and persistence (Harrison and Viers 2007; Burdon and Laine 2019). Additionally, high interannual variation in climate conditions and plant community composition in California grasslands may lead to temporal variation in infection dynamics through impacts on fungal growth and survival (Hobbs and Mooney 1991, Fernandez-Going *et al.* 2012).

We hypothesize that both disease burden and pathogen species composition differ on *S. pulchra* individuals across habitat types and years. In turn, we hypothesize that habitat type influences *S. pulchra* foliar tissue chemistry in ways that may affect plant – fungal pathogen interactions, with lower C and N content in serpentine plants growing in nutrient-poor soils. If so, these foliar chemistry differences may translate into differences in fungal community composition and either (a) higher disease burden in serpentine plants relative to nonserpentine plants, facilitated by negative impacts of limited plant resource availability on plant health (as in Springer *et al.,* 2006), or (b) lower disease burden in serpentine plants due to lower plant nutrient content limiting resources available for fungal growth. By contributing novel information about previously uncharacterized plant fungal pathogen communities in California grasslands, the results of this study will provide new insight into the diversity and natural history of the fungi that infect the leaves of wild grasses in temperate climates. Further, the results of this study will aid in assessing disease risk for plant populations in natural and agricultural contexts by providing information about the spatial and temporal scales over which environmental

heterogeneity can affect the species composition and severity of foliar pathogen infection.

Materials and Methods

Study system

All surveys and sample collection took place at Jasper Ridge Biological Preserve (JRBP), a 485-hectare site in San Mateo County, CA managed by Stanford University (37.4° N, 122.2° W). JRBP has a Mediterranean climate with cool winters (mean 9.2°C), warm summers (mean 20.1°C), and annually variable precipitation that predominantly occurs in the winter and spring (averaging 622.5 mm per year) (Ackerly *et al.* 2002). The growing season occurs during the rainy season and plants senesce in early summer. This study was conducted during the growing seasons from mid-April to mid-May in 2015, 2016, and 2017. There was highly variable annual precipitation across the three years (491 mm, 604 mm, and 985 mm in 2015, 2016, and 2017, respectively; Weather Underground).

S. pulchra occurs at substantial densities in both serpentine and nonserpentine grasslands at JRBP (McNaughton 1968). The serpentine plant communities at JRBP are more diverse than the nearby nonserpentine plant communities and include many native grasses and forbs, in contrast to the invasive-dominated nonserpentine grasslands (McNaughton 1968; Huenneke *et al.* 1990; Field *et al.* 1996). Plant species commonly found in serpentine grasslands at JRBP include native *Stipa pulchra*, *Elymus glaucus*, *Elymus multisetus, Eschscholzia californica*, and the invasive grass *Bromus hordeaceus* (McNaughton 1968; Hobbs and Mooney 1985). The nonnative annual species that dominate the nonserpentine grasslands but are absent from serpentine grasslands include *Avena barbata, Avena fatua*, *Bromus diandrus,* and *Erodium botrys* (McNaughton 1968; Hobbs and Mooney 1985; Field *et al.* 1996). The serpentine and nonserpentine grasslands at JRBP share a land use history of livestock grazing that ceased in

1960, when conservation of these areas became a management priority (Bocek and Reese 1992).

All survey sites were located along the ridgetop at JRBP, where serpentine and nonserpentine (greenstone) soils are present in discrete adjacent bands (Fig. 1a) (Oze *et al.* 2004). Chemical analyses of the soils on this ridgetop show consistent, significant differences in serpentine and nonserpentine soil chemistry, with the serpentine soils enriched in heavy metals and trace elements (Fe, Cr, Ni, Mn, and Co), and impoverished in essential plant nutrients (Streit *et al.* 1993, Oze *et al.* 2004, 2008). The relative flatness of the ridgetop ensured that all survey sites were at roughly the same elevation and had similar slopes, aspects, and water availability (Oze *et al.* 2008).

Quantification of disease

To assess plant disease burden, we quantified the percentage of living leaf area exhibiting symptoms of fungal disease in serpentine and nonserpentine populations of *S. pulchra*. We selected three serpentine grassland sites and three nonserpentine grassland sites such that each serpentine site was paired with a nearby (~170m away) nonserpentine site (Fig. 1a). Individual sites within each habitat type were >300m apart and the maximum distance between any two sites was 1600m (Fig. 1a). Each year, we randomly placed four 5-meter transects at each 40m x 40m site and assessed the infection status of the *S. pulchra* individual nearest each meter mark, resulting in five individual plants per transect surveyed each year (N = 60 plants per habitat type per year). The infection status of individual plants was not tracked across years, although some plants could have been sampled repeatedly by chance. Following the protocol for estimation of foliar infection severity on grasses described in Mitchell *et al.* 2002 and Spear and Mordecai 2018, a single surveyor visually estimated the percentage of living foliar tissue damaged by pathogens for six arbitrarily-selected leaves per plant. We used printed field reference guides

showing known amounts of damage on leaves to standardize estimates. Damage was indicated by spots and lesions which ranged in color from dark brown to pale yellow (Fig. 1b). Senesced tissue and herbivore damage (commonly in the form of scrapes and holes in the leaf surface) was excluded from the analyzed leaf area. For damage estimates of 5% or greater, values were determined to the nearest 5%. Damage levels below 5% were assigned values of 1% or 0.1%.

We used analysis of variance (ANOVA), with individual plants nested within sites and sites nested within habitat types, to test for separate effects of habitat type and year, as well as interactions between habitat type x year and habitat type x site, on log-transformed mean percent disease damage on serpentine and nonserpentine *S. pulchra* (N = 36 serpentine and 36 nonserpentine transects). When ANOVA results were statistically significant, we used t-tests for every possible pairwise combination, with Bonferroni adjustments of p-values to account for multiple comparisons, to identify the interactions contributing to the significant result.

Isolation and molecular identification of foliar fungal pathogens

To identify the fungal foliar pathogens infecting serpentine and nonserpentine *S. pulchra* over all years, we harvested one segment of symptomatic leaf tissue for culturing and identification of the putatively causal fungal pathogen(s) from each individual included in the three-year damage survey with symptoms of disease. This tissue segment came from the first surveyed leaf with disease symptoms on each plant. We excised, surface sterilized, and plated a 4mm^2 square of tissue from each sample at the leading edge of infection, to improve the chances of isolating the causal fungal pathogen, on malt extract agar with 2% chloramphenicol to prevent bacterial growth. While it is possible that the growing medium used may have led to selective isolation and sequencing of fungi other than those causing symptoms, our use of small segments of leaf tissue from the advancing margin of disease was intended to prevent this outcome under

the assumption that symptom-causing organisms should be dominant in visibly diseased tissue and particularly at its expanding margin. We observed the plates for growth for eight weeks and isolated morphologically distinct hyphae into axenic culture. Six segments of surface sterilized, asymptomatic tissue were plated as a control and showed no growth. We extracted and sequenced genomic DNA from the ITS region for each isolate. Fungal isolation and sequencing methods with ITS1F and ITS4 primers followed Spear and Mordecai 2018.

We estimated taxonomic placement of fungal isolates by grouping sequences into operational taxonomic units with 97% sequence similarity (OTUs), a proxy for species delineation based on the range of intraspecific ITS divergence (O'Brien *et al.* 2005), and then comparing assembled DNA sequences to the entries in the UNITE fungal database. To prepare sequences for taxonomic placement, we first trimmed forward and reverse Sanger sequencing reads to a maximum error rate of 5% and then automatically assembled bidirectional reads into consensus sequences with a minimum of 20% overlap and 85% sequence similarity in Geneious (Kearse *et al.* 2012), discarding consensus sequences less than 450 bp in length and/or less than 60% in read quality. We grouped the remaining internal transcribed spacer (ITS) sequences into OTUs using USEARCH 10.0.240 (Edgar 2010) and estimated the taxonomic placement of each OTU in mothur v.1.40.5 (http://mothur.org/; Schloss 2009) using the naïve Bayes classification approach (classify.seqs with options: method = 'wang', cutoff = 0, iters = 200). We used a bootstrapping confidence score of at least 80% to assign a species name and at least 60% to assign taxonomy at higher ranks.

Analyses of fungal community composition

To characterize foliar fungal community composition in serpentine and nonserpentine *S. pulchra* populations, we compared the diversity and assessed the similarity of communities of

fungi cultured from the diseased tissue of surveyed plants. Additionally, we compared the identified fungal taxa to previously reported plant - fungal genus associations compiled in the USDA-ARS plant - fungal database (Farr and Rossman 2020), the Genera of Phytopathogenic Fungi (GOPHY) database (Marin-Felix *et al.*, 2017, 2019), and the publications indexed in Web of Science and Google Scholar. To describe fungal community diversity, we calculated observed taxa richness and Fisher's alpha, a widely used measure of species richness that is robust to uneven sample sizes (Fisher *et al.* 1943; Magurran 2013); generated taxa accumulation curves to understand sampling efficacy (Oksanen *et al.* 2016); and counted the number of fungal isolates in each genus, family, order, and class from each grassland type. We compared fungal community composition across grassland types and across years visually, using non-metric multidimensional scaling (NMDS), and statistically, using permutational multivariate analyses of variance (PERMANOVA) with pairwise Adonis functions for non-normalized OTU read counts in the R package vegan (Anderson 2001; Arbizu 2016; Oksanen *et al.* 2016). To assess spatial turnover in fungal communities, we included a site distance factor in the statistical analyses, with comparisons between sites <600m apart categorized as "near" and comparisons between sites >600m apart categorized as "far" (Fig. 1a). The PERMANOVA tested for effects of habitat type, year, distance between sites, and all possible interactions between each of these three factors, on fungal community composition. We used the function 'betadisper' with the 'anova' method to assess the potential for group differences in dispersion to affect the PERMANOVA results. To account for low culturing success in some transects, we considered the combined isolates from two nearby transects at each site in each year (N = 18 serpentine and 18 nonserpentine communities, generated from 36 total transects per habitat type) to be distinct communities for NMDS visualization.

For ANOVA and PERMANOVA analyses we considered the combined isolates from each site in each year to be distinct communities (N = 9 serpentine and 9 nonerspentine communities). When PERMANOVA results were statistically significant, we used pairwise PERMANOVAs with Bonferroni adjustments to p-values to clarify which fungal communities were significantly different from one another (Arbizu 2016). Additionally, we used the function 'betadisper' in vegan to assess the potential for differences in dispersion to affect the PERMANOVA results (Oksanen *et al.* 2016). For the NMDS and PERMANOVA analyses, we used the function vegdist with the abundance-based Chao method, accounting for unsampled species, to create a matrix of pairwise community dissimilarities (Chao *et al.* 2016; Oksanen *et al.* 2016). We also used the Morisita-Horn index, which is independent of sample size and diversity, to make pairwise comparisons of similarity within and between the serpentine and nonserpentine fungal communities from each year (N = 265 isolates, 200 bootstrap replicates), and within the entire fungal community between years (Wolda 1981; Jost *et al.* 2011). Finally, we used ANOVA to test for effects of habitat type, year, site distance, year x habitat type, and year x site distance interactions on fungal diversity (Fisher's alpha) and richness at the site level.

To perform these fungal community analyses, we used the packages BiodiversityR (Kindt and Coe 2005), bipartite (Dormann *et al.* 2008), fossil (Vavrek 2011), rich (Rossi 2011), SpadeR (Chao *et al.* 2016), and vegan (Oksanen *et al.* 2016) with custom commands (Gardener 2014) in R version 3.4.1.

Chemical analyses of leaf tissue

In 2016, we analyzed C and N content of the foliar tissue of the *S. pulchra* plants surveyed in serpentine and nonserpentine grasslands. Because both leaf age and infection status might influence leaf chemistry, we collected the youngest fully expanded and entirely

asymptomatic leaf of each individual surveyed in the field (N = 60 serpentine, 58 nonserpentine).

We dried the samples at room temperature for three months, then at 60°C for 48 hours, ground

them to a powder, and then dried them again at 60°C for 24 hours. We measured C and N

content of these samples with a Carlo Erba NA-1500 elemental analyzer using atropine

standards.

To assess the potential for habitat type to mediate plant-pathogen interactions through its

influence on plant tissue chemistry, we related habitat type to leaf tissue chemistry of *in situ S.

pulchra*, and chemistry of these plants to disease burden and fungal species abundances. To

relate leaf tissue chemistry to observed damage levels, we compared the mean percentage of C

and N per dried weight of leaf tissue for surveyed serpentine and nonserpentine plants with

Wilcoxon rank sum tests. We also compared mean C:N ratio, which is indicative of plant health

(Ciompi *et al.* 1996; Brady *et al.* 2005). We used Pearson's product-moment correlation to test

for significant monotonic relationships between the log-transformed mean amount of diseased

tissue observed on individual plants in the field and their foliar C and N content. Additionally,

we used multivariate generalized linear models with resampling-based hypothesis testing to test

for fungal community responses to tissue chemistry. For fungal species represented by five or

more isolates in 2016, we tested for significant effects of foliar tissue C and N percentages, C:N

ratio, and habitat type on fungal species observations. For these analyses, we used the functions

'manyglm' and 'anova' (nBoot = 999), with p-values adjusted for multiple comparisons, in the R

package mvabund (Wang et al. 2012).

All statistical analyses were done in R version 3.4.1. In addition to the R packages

specified above, we used the packages cowplot (Wilke 2019), data.table (Dowle *et al.* 2019),

RColorBrewer (Neuwirth 2014), reshape2 (Wickham 2007), tidyverse (Wickham 2017), and

xtable (Dahl *et al*. 2019) to manipulate, plot, and export data and results.

Results

Pathogen damage

Pathogen damage was similarly ubiquitous yet low-burden in serpentine and nonserpentine grassland habitats. Every plant surveyed exhibited evidence of foliar fungus-caused disease. The mean percentage of diseased leaf area observed across all years was 1.63 for serpentine and 1.66 for nonserpentine *S. pulchra,* respectively (Fig. 1c), and the mean proportion of surveyed leaves with pathogen damage was 0.83 for serpentine and 0.85 for nonserpentine plants. Year significantly affected log-transformed mean percent damage (ANOVA: df = 2, SS = 9.41, F-value = 5.375, p-value = 7.04 x 10^{-3}) (Table 1). There were significant differences in percent damage between was between 2015 and 2016 and between 2015 and 2017 (2015 mean percent damage = 1.05; 2016 = 1.96; 2017 = 1.94; t-values = -2.66, -3.70; Bonferonni-adjusted p-values = 2.5 x 10^{-3}, 8.1 x 10^{-4}).

Fungal pathogen community

We isolated 267 unique fungal isolates from 258 plated symptomatic tissue pieces with fungal growth (144 nonserpentine and 114 serpentine tissue pieces from 36 nonserpentine and 34 serpentine transects, out of 360 total tissue pieces). Of the 267 unique fungal isolates, we successfully sequenced 256 isolates. The sequenced isolates clustered into 30 operational taxonomic units (OTUs) based on 97% sequence similarity, representing 23 genera, 15 families, 8 orders, and 5 fungal classes, primarily the Dothideomycetes. We were unable to identify 18 OTUs to the fungal species level, and four of these to the genus level, due to a lack of close sequence matches in the UNITE database and/or a lack of taxonomic information for close

sequence matches (Table 2). The most common OTU was represented by 48 isolates and the least common OTUs were represented by one isolate each. OTUs are hereafter referred to as "species" (O'Brien et al. 2005).

All genera identified include fungal species previously isolated from plant hosts, and the majority of these genera include known plant pathogens (Table 2). The taxonomic results indicate that many of the species represent close relatives of major agricultural plant pathogens including *Alternaria, Cladosporium, Drechslera, Phaeosphaeria, Pyrenophora, Stemphylium, Ramularia,* and *Wojnowiciella* spp. (Table 2) (Kirk *et al.* 2008; Stukenbrock and McDonald 2008; Havis *et al.* 2015; Marin-Felix *et al.* 2017, 2019; Farr and Rossman 2020). Inoculation experiments described in Spear and Mordecai 2018 previously confirmed the pathogenicity of *Parastagonospora* and *Pyrenophera* species on *S. pulchra* leaves. Six genera, *Heyderia, Hyphodiscus, Nemania, Phaeophlebiopsis, Plectania,* and *Wojnowiciella,* are here reported for the first time from hosts in the Poaceae; six genera, *Heyderia, Hyphodiscus, Keissleriella, Phaeophlebiopsis, Plectania,* and *Podospora* are reported for the first time in association with plant disease symptoms; and two additional genera, *Chalastospora* and *Nemania,* are reported for the first time in association with foliar disease symptoms (Table 2) (United States Department of Agriculture 1960; Goetz and Dugan 2006; Marin-Felix *et al.* 2017, 2019; Golzar *et al.* 2019; Adikaram and Yakandawala 2020; Farr and Rossman 2020).

Fungal community diversity was similar between serpentine and nonserpentine habitats and between years (Table 2). Although fungal richness was higher at every taxonomic level in the serpentine relative to the nonserpentine, these differences were not statistically significant (Tables 2, S1-S5). Twenty-four fungal species were isolated from serpentine *S. pulchra* (Fisher's alpha = 9.324, 95% CI = 4.664, 13.983) and 22 from nonserpentine (Fisher's alpha = 7.261, 95%

CI = 3.597, 10.925); 16 of these species (53%) were shared (Fig. 2; Tables 1, 2). Habitat type, year, and year x habitat type were not significantly related to fungal richness or Fisher's alpha at the site level (ANOVA: Tables S6, S7). Novel genera were isolated for both community types in every year of surveying, with only seven of 30 (23%) observed species and six of 23 genera (26%) isolated in all three years (Table S5). Species accumulation curves did not approach horizontal asymptotes for either habitat type, suggesting that neither fungal community was fully described (Fig. S1).

Both fungal communities were dominated by a few abundant species (isolated >10 times), but the most abundant species differed between nonserpentine and serpentine habitats: these were *Alternaria* cf. *infectoria* and *Ramularia* cf. *proteae*, respectively (Table 2; Fig. S2). The nonserpentine community included five abundant species, representing 62% of the isolates, while the serpentine community included two abundant species, representing 51% of serpentine isolates (Table 2; Fig S2). The serpentine community had a higher proportion of rare species (<3 isolates) than the nonserpentine community, 63% versus 36%, respectively. One *Drechslera* species was abundant in the nonserpentine community and rare in the serpentine (Table 2; Fig. S2).

Habitat type and year significantly affected fungal community composition, based on two-way PERMANOVA analysis of site-level fungal communities and NMDS visualization of community similarity (habitat type: F-value = 5.112, R^2 = 0.163, p = 0.003; year: F-value = 5.655, R^2 = 0.360, p = 0.001; Fig. 3; Table 4). Fungal communities had homogeneous dispersions across habitat types (average distance to median: serpentine = 0.53, nonserpentine = 0.56; ANOVA F-value = 1.10, p-value = 0.3) and years (average distance to median: 2015 = 0.38, 2016 = 0.34, 2017 = 0.27; ANOVA F-value = 0.93, p-value = 0.42), indicating that

significant PERMANOVA results were due to true differences in centroid positions. Serpentine

fungal communities were generally more similar to one another than they were to nonserpentine

fungal communities, and vice versa (Fig. 3; Tables 2, 3). Community composition differed

significantly between 2015 and 2017, and 2016 and 2017, but not between 2015 and 2016 (Table

S8). Differences in fungal community composition between years did not correspond to

differences in disease severity between years (Figure 1c, Table S8).

Interpretation of the statistical significance of differences in Morisita-Horn community

overlap that fall between zero (no overlap) and one (complete overlap) is a longstanding

challenge in community ecology (Horn 1966, Rodrigues and Vieira 2010). Qualitatively,

estimates of the Morisita-Horn similarity index for communities within habitat types suggest that

the amount of interannual species turnover is more consistent within serpentine than

nonserpentine communities (Fig. 3, Tables 2, 3). While serpentine community overlap between

years ranged from 0.707 to 0.786, nonserpentine community overlap ranged between 0.295 and

0.882 (Table 3). Within-year variation in species turnover between habitat types was

comparatively moderate, with Morisita-Horn overlap ranging from 0.535 to 0.725 (Table 3).

Leaf tissue chemistry and plant – pathogen interactions

As expected from differences in soil type between habitats, leaf chemistry in plants

occurring on serpentine versus nonserpentine soils differed in mean C and N content. Serpentine

plants had significantly lower mean C and N content than nonserpentine plants (Fig. 4; Table

S9). C:N ratio was similar for plants in both habitat types. Log-transformed mean percent

symptomatic tissue was not significantly associated with percent C, percent N, or C:N ratio in

asymptomatic foliar tissue for serpentine or nonserpentine plants, based on Pearson's product-

moment correlation tests (Fig. 4). For the fungal community consisting of the seven species

isolated five or more times in 2016 (N = 74 isolates), all of which occurred at least once in each habitat type, species composition differed between habitats, but was not associated with foliar percent C, percent N, or C:N ratio (Table S10).

Discussion

Here we show that a diverse suite of foliar fungi is associated with ubiquitous, low-burden damage on host individuals across growing seasons and habitat types. Habitat type appeared to shape fungal community composition in this system, but not foliar disease burden. In contrast, sampling year affected both fungal community composition and disease burden. However, this interannual variation in disease burden was not mediated by differences in pathogen community composition (Figs. 1c, 3; Table S8). While C and N content measured in young, healthy leaves differed between habitat types, variation in leaf chemistry did not explain variation in infection severity or in fungal community composition (Fig. 4; Tables S9-10). The results provide some of the first evidence that local growing conditions structure the pathogen communities infecting a single host species even over relatively small areas of a few hundred meters, where hosts and pathogens experience similar climate conditions (Figs. 1a, 3; Table 3). Overall, the findings imply that environmental heterogeneity within landscapes may both generate barriers to fungal disease spread in wild plant populations and bolster fungal diversity.

Foliar disease burden

Mean levels of foliar infection were low even at their peak of 1.9% symptomatic leaf area in 2017 (Fig. 1c; Table 1). This level of damage is similar to that observed on other grass species in nonserpentine grasslands at JRBP (Spear and Mordecai 2018), and on taxonomically diverse tropical forest plants in Mexico and Hawaii (García-Guzmán and Dirzo 2001, Funk and Throop 2010). Recent work in the JRBP grassland system shows that symptoms in the amounts observed

inflict only minimal plant fitness costs, suggesting that *S. pulchra* is able to effectively resist or suppress damage by foliar pathogens in both serpentine and nonserpentine grasslands (Spear and Mordecai 2018; Uricchio *et al.* 2019). These expectations align with assessments in wheat – fungal pathogen systems that indicate yield reductions of ≤ 1% for each percent of symptomatic leaf area (Milus 1993, Khan *et al.* 1997). The finding that between-habitat differences in foliar C and N content do not correspond to differences in infection severity aligns with the finding that infection severity did not differ between habitats (Figs. 1, 4). Additionally, similar C:N ratios suggesting that serpentine and nonserpentine plants are equally healthy correspond to the consistently low damage observed across habitats (Fig. 4; Table S9). However, the observational study design and use of young, healthy leaves for the tissue chemistry measurements limit interpretation of the observed relationships between leaf chemistry and disease severity. Because both tissue age and fungal infections can affect plant chemistry (Paul 1989; Donaldson *et al.* 2006), the leaf chemistry measurements may not accurately represent the chemical environment encountered by the pathogens that caused the observed damage.

Fungal community composition

Our results suggest that habitat filtering, rather than dispersal limitation, plays a key role in fungal community assembly. The sampled nonserpentine and serpentine plants are in close proximity, yet habitat type—not distance between sites—predicted fungal community composition (Fig. 3; Tables 4, S10) (Martiny *et al.* 2006). Fungal community composition, and its relative stability over time, varied between habitats (Figs. 1c, 2, 3, S2; Tables 1-4). The results imply that abiotic and biotic differences in local environmental conditions can lead to differences in plant-fungal interactions in distinct habitat types. We hypothesize that influential habitat characteristics in this system include differences in pathogen sharing among the predominantly

native hosts in serpentine grasslands versus the predominantly nonnative hosts in nonserpentine grasslands, in plant phenotypes, and/or in plant genotypes.

Plant hosts may differ in competence for different fungal species, and fungal species may in turn vary in transmissibility to *S. pulchra*. These two factors, and their interaction, likely contribute to differences in pathogen community composition on *S. pulchra* between serpentine and nonserpentine habitats. Serpentine grassland plant communities are generally more taxonomically diverse and stable across years than nonserpentine grasslands (Fernandez-Going *et al.* 2012). Qualitatively, our results suggest that fungal communities follow the same pattern of higher diversity and lower species turnover between years in serpentine areas, indicating potential links between plant and pathogen community structures (Fig. S2; Tables 1, 3, S1-S5). Research on five common invasive and native grasses (including *S. pulchra*) at JRBP shows that multi-host foliar fungal pathogens dominate this system, and that *S. pulchra* shares different pathogens with different co-occurring plant species, supporting this hypothesis (Spear and Mordecai 2018).

Differing nonserpentine and serpentine *S. pulchra* phenotypes might also influence fungal community assembly through a variety of mechanisms, including impacts of plant chemistry on plant immune pathways and/or fungal pathogen virulence, growth, and reproduction. Although differences in plant C and N content did not affect the distributions of commonly isolated fungal species found in both habitat types (Tables 2, S10), the limitations of the foliar chemistry measurements described above apply equally here. Importantly, the impacts of leaf C and N content on the distributions of rare species, many of which appear to be habitat specialists, remain unassessed (Tables 2, S10). Evidence that serpentine soil chemistry impacts on plant Ca content affect the frequency and severity of *Hesperolinon spp.* infection by the

fungal rust pathogen *Melampsora lini* indicates the possible importance of habitat influence on elements other than C and N in *S. pulchra* tissues (Springer *et al.* 2006; Springer 2009). Potted *S. pulchra* plants grown from seed and serpentine soils collected in these sites had significantly lower Ca and P content and higher Ni and Mg content than similar plants grown in greenhouse soils, potentially suggesting additional phenotypic differences between serpentine and nonserpentine plants that may affect fungal community composition (Farner *et al.*, unpublished data).

Overall, further investigation is needed to test the hypothesis that habitat-associated plant phenotypic differences shape *S. pulchra* foliar fungal communities. Future studies should include analyses of more fungal species, more elements, and more plant traits, including leaf physiological characteristics that might respond to elevated water stress in serpentine habitats. Based on the genetic foundations of the complex set of physical, chemical, and genetic defenses that define plant immune profiles, investigation into the extent of genetic divergence between serpentine and nonserpentine *S. pulchra* could also clarify the mechanisms that structure its fungal communities (Dodds and Rathjen 2010; Łaźniewska *et al.* 2012). While there are no physical barriers to prevent gene flow between individuals in different habitat types, the occurrence of phylogenetically and chemically distinct serpentine and nonserpentine populations of the native forb *Lasthenia californica* occurring in the same grasslands at JRBP suggests the potential for similar structure in the focal *S. pulchra* population (Desrochers and Bohm 1993; Chan *et al.* 2002; Rajakaruna *et al.* 2003).

In addition to turnover between habitat types, interannual fungal species turnover contributed substantially to fungal diversity in this system, with only seven of 30 total species isolated in all three years (Fig. 3; Tables 1, S5). Interannual variation in these communities may

be driven by a variety of mechanisms, including high yearly variation in precipitation (2015 rainfall = 491 mm; 2016 = 604 mm, 2017 = 985 mm) (Weather Underground), plant population sizes, occurrences of different plant genotypes, and plant community composition (Hobbs and Mooney 1991). Although few studies track interannual variation in fungal communities on plants, the year-to-year dissimilarity of those hosted by *S. pulchra* is similar to that observed across a diverse taxonomic range of plants including ferns, legumes, and shrubs (Del Olmo-Ruiz and Arnold 2014; Cotton *et al.* 2015; Zhang *et al.* 2016). In this context, our results are consistent with a growing body of evidence for the general importance of storage effects—in which temporally variable conditions act as ecological niches that facilitate species diversity—in plant-associated fungal communities (Warner and Chesson 1985). This finding highlights the necessity of long-term data collection to accurately characterize plant-fungal interaction networks.

The variation in fungal community composition we observed on a single host species with habitat type and over time suggests possible mechanisms preventing severe outbreaks of foliar disease in California grasslands, where uniformly low disease burden was observed in this, and other, studies (e.g., Spear and Mordecai 2018). Transitions between distinct soil types over distances within the range of fungal spore dispersal may prevent the spread of pathogens that are sensitive to leaf tissue chemistry or have limited host ranges. Interannually variable conditions favoring successful infection by different fungal species each growing season suggest temporal barriers to pathogens evolving high virulence on both nonserpentine and serpentine plants (Table S5). Clarifying the specific mechanisms that drive differences in serpentine and nonserpentine fungal communities will require inoculation experiments to test how plant susceptibility to infection by different fungal species changes with plant tissue chemistry and plant genetics.

Characterization of comprehensive plant-pathogen interaction networks including all plant species co-occurring with *S. pulchra* in serpentine and nonserpentine grasslands will improve our understanding of the role of the surrounding plant community in structuring fungal pathogen communities.

Fungal diversity

Our work underscores the sparsity of current scientific understanding of plant – fungal interactions in natural systems. The novel plant host and plant disease associations reported here for nine fungal genera highlight the wealth of plant – fungal interactions that remain to be described (Table 2). Because this study is observational, the pathogenicity on *S. pulchra* of many of the fungal species isolated—even those from genera including well-known plant pathogens— remains unknown. It is possible that some of the isolated fungal taxa are not primary pathogens on this species, but endophytes that opportunistically act as secondary pathogens. Inoculation experiments and high-throughput sequencing studies to characterize and compare entire fungal communities inhabiting symptomatic and healthy *S. pulchra* leaf tissue will shed light on the ecology of the isolated taxa and on the potential for interactions between co-occurring fungal species to impact plant health.

These open questions notwithstanding, *S. pulchra* appears to host an unusually species-rich community of culturable fungi within its diseased foliar tissue when compared to other plant species' interactions with both pathogenic and non-pathogenic fungi. Where we isolated 30 fungal species*,* past studies that described fungal pathogen diversity for more than 100 different plant hosts consistently found fewer than 20 fungal pathogens per species, and almost all hosts had fewer than ten pathogens (Mitchell and Power 2003; Hantsch *et al.* 2013; Rottstock *et al.* 2014). The level of fungal diversity we documented is also high compared to that of culturable

fungal endophyte communities associated with asymptomatic plant tissues: of 35 plant species

analyzed by Arnold and colleagues, only five were found to host 30 or more endopyhtes (Arnold

and Lutzoni 2007; Del Olmo-Ruiz and Arnold 2014). Notably, studies that utilize high-

throughput sequencing techniques to observe both culturable and unculturable endophytic fungi

frequently report hundreds of fungal OTUs colonizing single host species (e.g., Jumpponen and

Jones 2009, Zimmerman and Vitousek 2012). However, such results are difficult to compare to

our own given the mismatch in the detection capabilities of the methods used. Nonetheless, the

exceptional diversity of the culturable foliar fungi that inhabit *S. pulchra* is further indicated by

species accumulation curves that suggest many more species in this community remain to be

discovered (Fig. S1), by the lack of close sequence matches in genetic databases for some of the

species isolated (Table 2), and by the isolation of multiple genera previously unassociated with

plant hosts in the Poaceae (Table 2). These findings suggest that the characteristics that make the

California Floristic Province a biodiversity hotspot may extend to the fungal diversity associated

with this flora (Myers *et al.* 2000).

Conclusions

Fungi associated with foliar disease in the native bunchgrass *S. pulchra* are relatively

benign, likely cause minimal fitness impacts (Spear and Mordecai 2018), and contribute

substantially to the overall biodiversity of California grasslands. Variation between habitats

appears to play an important role in supporting the fungal diversity observed here, and may also

reduce disease risk for *S. pulchra*. The high taxonomic diversity and rarity of serpentine fungi

demonstrates that this community is potentially at risk of species loss via host-parasite

coextinction as nonnative annual grasses continue to expand their ranges at the cost of native

plants, and human activities including development and increased C and N deposition contribute

to the loss of serpentine grasslands that already make up <2% of California's surface area (Huenneke *et al.* 1990; Vallano *et al.* 2012). This work demonstrates that, in addition to >200 endemic plant species, California's serpentine grasslands support unique communities of leaf-infecting fungi, providing additional motivation for active conservation of native grassland communities based on the far-reaching loss of biodiversity across trophic levels associated with their disappearance (Sprent 1992; Dunn *et al.* 2009).

Acknowledgements

We thank Nona Chiariello, Caroline Daws, Joe Wan, Peter Vitousek, Scott Fendorf, Lawrence Uricchio, Guangchao Li, Douglas Turner, Virginia Walbot, Cody Hamilton, and the Mordecai and Peay Lab Groups for their help. This work was supported by the Jasper Ridge Kennedy Endowment, the National Science Foundation (DEB-1518681), Stanford Vice Provost for Undergraduate Education, and Stanford Bio-X.

References

Abbate JL, Antonovics J. 2014. Elevational disease distribution in a natural plant–pathogen system: insights from changes across host populations and climate. *Oikos* **123:** 1126–1136.

Ackerly DD, Knight CA, Weiss SB, Barton K, Starmer KP. 2002. Leaf size, specific leaf area and microhabitat distribution of chaparral woody plants: contrasting patterns in species level and community level analyses. *Oecologia* **130:** 449–457.

Adikaram NKB, Yakandawala DMD. 2020. A checklist of plant pathogenic fungi and Oomycota in Sri Lanka. *Ceylon Journal of Science* **49(1):** 93-123.

Agrios GN. 2005. *Plant Pathology.* Burlington, MA, USA. Elsevier Academic Press.

Anagnostakis, SL. 1987. Chestnut blight: the classical problem of an introduced pathogen. *Mycologia* **79(1):** 23-37.

Anderson MJ. 2001. A new method for non-parametric multivariate analysis of variance. *Austral **Ecology** 26:* 32–46.

Arbizu PM. 2016. *How can I do PerMANOVA pairwise contrasts in R?*[WWW document] Researchgate 2016. URLhttps://www.researchgate.net/post/How_can_I_do_PerMANOVA_pairwise_contrasts_in_R. [accessed 21 June 2019].

Arnold AE. 2007. Understanding the diversity of foliar endophytic fungi: progress, challenges, and frontiers. *Fungal Biology Reviews* **21:** 51-66.

Arnold AE, Lutzoni F. 2007. Diversity and host range of foliar endophytes: are tropical leaves biodiversity hotspots? *Ecology* **88 (3):** 541-549.

Bálint M, Bartha L, O'Hara RB, Olson MS, Otte J, Pfenninger M, Roberston AL, Tiffin P, Schmitt I. 2015. Relocation, high-latitude warming and host genetic identity shape the

foliar fungal microbiome of poplars. *Molecular Ecology* **24 (1):** 235-248.

Barge EG, Leopold DR, Peay KG, Newcombe G, Busby PE. 2019. Differentiating spatial

from environmental effects on foliar fungal communities of *Populus trichocarpa*. *Journal of*

Biogeography **46(9):** 2001-2011.

Bernard F, Sache I, Suffert F, Chelle M. 2013. The development of a foliar fungal pathogen

does react to leaf temperature! *New Phytologist* **198(1):** 232-240.

Bocek B, Reese E. 1992. Land use history of Jasper Ridge Biological Preserve. *Stanford*

University Research Report **8.**

Bolton MD, Kolmer JA, Garvin DF. 2008. Wheat leaf rust caused by *Puccinia triticina*.

Molecular Plant Pathology **9(5):** 563-575.

Borer ET, Laine A, Seabloom EW. 2016. A multiscale approach to plant disease using the

metacommunity concept. *Annual Review of Phytopathology* **54:** 397–418.

Brady KU, Kruckeberg AR, Bradshaw HD. 2005. Evolutionary ecology of plant adaptation to

serpentine soils. *Annual Review of Ecology, Evolution, and Systematics* **36:** 243–266.

Brandhorst TT, Gauthier GM, Stein RA, Klein BS. 2005. Calcium binding by the essential

virulence factor BAD-1 of *Blastomyces dermatitidis*. *Journal of Biological Chemistry* **280:**

42156–42163.

Burdon JJ, Laine A-L. 2019. *Evolutionary Dynamics of Plant-Pathogen Interactions.*

Cambridge, UK. Cambridge University Press.

Busby PE, Newcombe G, Dirzo R, Whitham TG. 2014. Differentiating genetic and

environmental drivers of plant-pathogen community interactions. *Journal of Ecology*

102(5): 1300-1309.

Chan R, Baldwin BG, Ornduff R. 2002. Cryptic goldfields: A molecular phylogenetic

reinvestigation of *Lasthenia californica* sensu lato and close relatives (Compositae: Helantheae sensu lato). *American Journal of Botany* **89:** 1103–1112.

Chao A, MA KH, Hsieh TC, Chiu C-H. 2016. SpadeR: Species prediction and diversity estimation with R.

Ciompi S, Gentili E, Guidi L, Soldatini GF. 1996. The effect of nitrogen deficiency on leaf gas exchange and chlorophyll fluorescence parameters in sunflower. *Plant Science* **118:** 177–184.

Clarkson JP, Fawcett L, Anthony SG, Young C, Pöggeler S. 2014. A model for *Sclerotinia sclerotiorum* infection and disease development in lettuce, based on the effects of temperature, relative humidity and ascospore density. *PLoS ONE* **9(4):** e94049.

Cordier T, Robin C, Capdeveille X, Fabreguettes O, Desprez-Loustau M, Vacher C. 2012. The composition of phyllosphere fungal assemblages of European beech (*Fagus sylvatica*) varies significantly along an elevation gradient. *New Phytologist* **196:** 510–519.

Cotton TEA, Fitter AH, Miller RM, Dumbrell AJ, Helgason T. 2015. Fungi in the future: interannual variation and effects of atmospheric change on afbuscular mycorrhizal fungal communities. *New Phytologist* **205(4):** 1598-1607.

Dahl DB, Scott D, Roosen C, Magnusson A, Swinton J. 2019. xtable: Export Tables to LaTeX or HTML. R package version 1.8-4. https://CRAN.R-project.org/package=xtable

Dallas TA, Laine A-L, Ovaskainen O. 2019. Detecting parasite associations within multi-species host and parasite communities. *Proceedings of the Royal Society B: Biological Sciences* **286(1912):** 20191109.

Del Olmo-Ruiz M, Arnold AE. 2014. Interannual variation and host affiliations of endophytic fungi associated with ferns at La Selva, Costa Rica. *Mycologia* **106(1):** 8-21.

Desrochers AM, Bohm BA. 1993. Flavonoid variation in *Lasthenia californica* (Asteraceae). *Biochemical Systematics and Ecology* **21:** 449–453.

Dodds PN, Rathjen JP. 2010. Plant immunity: towards and integrated view of plant-pathogen interactions. *Nature Reviews Genetics* **11(8):** 539-548.

Donaldson JR, Stevens MT, Barnhill HR, Lindroth RL. 2006. Age-related shifts in leaf chemistry of clonal aspen (*Populus tremuloides*). *Journal of Chemical Ecology* **32(7):** 1415-1429.

Dordas C. 2008. Role of nutrients in controlling plant diseases in sustainable agriculture. A review. *Agronomy for Sustainable Development.* **28(1):** 33-46.

Dormann CF, Gruber B, Fruend J. 2008. 2008. Introducing the bipartite package: analysing ecological networks. *R news* **8:** 8–11.

Dowle M, Srinivasan A, Gorecki J, Chirico M, Stetsenko P, Short T, Lianoglou S, Antonyan E, Bonsch M, Parsonage H, *et al*. 2019. data.table: Extension of 'data.frame'. R package version 1.12.8. https://CRAN.R-project.org/package=data.table.

Dunn RR, Harris NC, Colwell L, Koh P, Sodhi NS. 2009. The sixth mass coextinction: are most endangered species parasites and mutualists? *Proceedings of the Royal Society B: Biological Sciences* **276:** 3037-3045.

Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26(19):** 2460-2461.

Eschen R, Hunt S, Mykura C, Gange AC, Sutton BC. 2010. The foliar endophytic community composition in *Cirsium arvense* is affected by myccorhizal colonization and soil nutrient content. *Fungal Biology* **114:** 991-998.

Eversmyer MG, Kramer CL, Hassan ZM. 1988. Environmental influences on the

establishment of *Puccinia recondita* infection structures. *Plant Disease* **72:** 409-412.

Farr DF, Rossman AY. 2020. Fungal Databases, U.S. National Fungus Collections, ARS, USDA. https://nt.ars-grin.gov/fungaldatabases/

Fernandez-Going BM, Anacker BL, Harrison SP. 2012. Temporal variability in California grasslands: soil type and species functional traits mediate response to precipitation. *Ecology* **93 (9):** 2104 – 2114.

Field CB, Chapin FS, Chiariello NR, Holland EA, Mooney HA. 1996. The Jasper Ridge CO_2 Experiment: Design and Motivation. In: Koch GW, Mooney HA, eds. *Carbon dioxide and terrestrial ecosystems.* San Diego, CA, USA: Academic Press, 121-145.

Fisher MC, Henk DA, Briggs CJ, Brownstein JS, Madoff LC, McCraw SL, Gurr SJ. 2012. Emerging fungal threats to animal, plant and ecosystem health. *Nature* **484:** 186–194.

Fisher RA, Corbet AS, Williams CB. 1943. The relation between the number of species and the number of individuals in a random sample of an animal population. *Journal of Animal Ecology* **12:** 42–58.

Flor HH. 1956. The complementary genic systems in flax and flax rust. *Advances in Genetics* **8:** 29–54.

Funk JL, Throop HL. 2010. Enemy release and plant invasion: patterns of defensive traits and leaf damage in Hawaii. *Oecologia* **162(4):** 815-823.

García-Guzmán G, Dirzo R. 2001. Patterns of leaf-pathogen infection in the understory of a Mexican rain forest: incidence, spatiotemporal variation, and mechanisms of infection. *American Journal of Botany* **88(4):** 634-645.

Gardener M. 2014. *Community ecology: analytical methods using R and Excel*. Exeter, UK: Pelagic Publishing.**Hansen RA, Coleman DC. 1998.** Litter complexity and composition are

determinants of the diversity and species composition of oribatid mites (Acari: Oribatida) in litterbags. *Applied Soil Ecology* **9:** 17–23.

Giauque H, Hawkes CV. 2016. Historical and current climate drive spatial and temporal patterns in fungal endophyte diversity. *Fungal Ecology* **20:** 108-114.

Goetz J, Dugan FM. 2006. *Alternaria malorum*: a mini-review with new records for hosts and pathogenicity. *Pacific Northwest Fungi* **1(3):** 1-8.

Golzar H, Thomas G, Jayasena KW, Wright D, Wang C, Kehoe M. 2019. Neoascochyta species cause leaf scorch on wheat in Australia. *Australasian Plant Disease Notes* **14(1):** 1-5.

Hantsch L, Braun U, Scherer-Lorenzen M, Bruelheid H. 2013. Species richness and species identity effects on occurrence of foliar fungal pathogens in a tree diversity experiment. *Ecosphere* **4(7):** 81.

Harrison SP, Viers JH. 2007. Serpentine Grasslands. In: Stromberg MR, Corbin JD, D'Antonio CM, eds. *California Grasslands: Ecology and Management*. Berkeley, CA, USA: University of California Press, 145-155.

Havis ND, Brown JKM, Clemente G, Frei P, Jedryczka M, Kaczmarek J, Kaczmarek M, Matusinsky P, McGraan GRD, Pereyra S, *et al.* 2015. *Ramularia collo-cygni*—an emerging pathogen of barley crops. *Phytopathology*. **105(7):** 895–904.

Hechinger RF, Lafferty KD. 2005. Host diversity begets parasite diversity: bird final hosts and trematodes in snail intermediate hosts. *Proceedings of the Royal Society B: Biological Sciences* **272:**1059–1066.

Hobbs RJ, Mooney HA. 1985. Community and population dynamics of serpentine annual grassland in relation to gopher disturbances. *Oecologia* **67:** 342–351.

Hobbs RJ, Mooney HA. 1991. Effects of rainfall variability and gopher disturbance on serpentine annual grassland dynamics. *Ecology* **72:** 59–68.

Horn HS. 1966. Measurement of "overlap" in comparative ecological studies. *The American Naturalist* **100(914):** 419-424.

Huber L, Gillespie TJ. 1992. Modeling leaf wetness in relation to plant disease epidemiology. *Annual Review of Phytopathology.* **30:** 553-577.

Huenneke LF, Hamburg SP, Koide R, Mooney HA, Vitousek PM. 1990. Effects of soil resources on plant invasion and community structure in Californian serpentine grassland. *Ecology* **71:** 478–491.

Jost L, Chao A, Chazdon RL. 2011. Compositional similarity and beta diversity. In: McGill BJ, Magurran AE, editors. *Biological diversity: frontiers in measurement and assessment.* Oxford, UK: Oxford University Press, 66-84.

Jumpponen A, Jones KL. 2009. Massively parallel 454 sequencing indicates hyperdiverse fungal communities in temperate *Quercus macrocarpa* phyllosphere. *New Phytologist* **184(2):** 438-448.

Kearse MR, Moir A, Wilson S, Stones-Havas M, Cheung S, Sturrock S, Buxton A, Cooper S, Markowitz C, Duran T et al. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28(12):** 1647-1649.

Khan MA, Trevathan LE, Robbins JT. 1997. Quantitative relationship between leaf rust and wheat yield in Mississippi. *Plant Disease* **81(7):** 769-772.

Kindt R, Coe R. 2005. BiodiversityR. Tree diversity analysis. A manual and software for common statistical methods for ecological and biodiversity studies. World Agroforestry

Centre (ICRAF), Nairobi.

Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008. *Ainsworth and Bisby's Dictionry of the Fungi.* Oxon, UK: CAB International.

Łaźniewska J, Macioszek VK, Kononowicz AK. 2012. Plant-fungus interface: the role of surface structures in plant resistance and susceptibility to pathogenic fungi. *Physiological and Molecular Plant Pathology* **78:** 24-30.

Lee SC, Lee YH. 1998. Calcium/calmodulin-dependent signaling for appressorium formation in the plant pathogenic fungus *Magnaporthe grisea. Molecules & Cells* **8:** 698–704.

Levine A, Pennell RI, Alvarez ME, Palmer R, Lamb C. 1996. Calcium-mediated apoptosis in a plant hypersensitive disease resistance response. *Current Biology* **6:** 427–437.

Loehrer M, Schaffrath U. 2011. Soybean rust – meet a prominent challenge in soybean cultivation. In *Ng T-B*, editor. *Soybean – Biochemistry, Chemistry, and Physiology.* London, UK: InTech, 83–100 .

Madden, LV. 1997. No effects of rain on splash dispersal of fungal pathogens. *Canadian Journal of Plant Pathology* **19:** 225–230.

Magarey RD, Sutton TB, Thayer CL. 2005. A simple generic infection model for foliar fungal plant pathogens. *Phytopathology* **95(1):** 92-100.

Magurran AE. 2013. *Measuring Biological Diversity*. Malden, MA, USA: Blackwell Science Ltd.

Marin-Felix Y, Groenewald JZ, Cai L, Chen Q, Marincowitz S, Barnes I, Bensch K, Braun U, Camporesi E, Damm U, *et al.* **2017.** Genera of phytopathogenic fungi: GOPHY 1. *Studies in Mycology* **86:** 99–216.

Marin-Felix Y, Hernández-Restrepo M, Wingield MJ, Akulov A, Carnegie AJ,

Cheewangkoon R, Gramaje D, Groenewald JZ, Guarnaccia V, Halleen F, *et al.* 2019. Genera of phytopathogenic: GOPHY 2. *Studies in Mycology* **92:** 47–133.

Martiny JBH, Bohannan BJM, Brown JH, Colwell RK, Fuhrman JA, Green JL, Horner-Devine MC, Kane M, Krumins JA, Kuske CR, *et al.* 2006. Microbial biogeography: putting microorganisms on the map. *Nature Reviews Microbiology* **4(2):** 102-112.

McNaughton SJ. 1968. Structure and function in California grasslands. *Ecology* **49:** 962–972.

Millberg H, Boberg J, Stenlid J. 2015. Changes in fungal community of Scots pine (*Pinus sylvestris)* needles along a latitudinal gradient in Sweden. *Fungal Ecology* **17:** 126-139.

Milus EA. 1993. Effects of leaf rust and septoria leaf blotch on yield and test weight of wheat in Arkansas. *Plant Disease* **78(1):** 55-59.

Mitchell CE, Power AO. 2003. Release of invasive plants from fungal and viral pathogens. *Nature* **421:** 625–627.

Mitchell CE, Tilman D, Groth JV. 2002. Effects of grassland plant species diversity, abundance, and composition on foliar fungal disease. *Ecology* **83(6):** 1713-1726.

Moss MA. 1987. Environmental conditions conducive to infection of Ryegrass by *Pyricularia grisea. Phytopathology* **77(6):** 863-866.

Myers N, Mittermeir RA, Mittermeir CG, da Fonseca, GAB, Kent J. 2000. Biodiversity hotspots for conservation priorities. *Nature* **403(6772):** 853-858.

Nazareno ES, Li F, Smith M, Park RF, Kianian SF, Figureoa M. 2018. *Puccinia coronata* f. sp. *avenae*: a threat to global oat production. *Molecular Plant Pathology* **19(5):** 1047–1060.

Neuwirth, E. 2014. RColorBrewer: ColorBrewer Palettes. R package version 1.1-2. https://CRAN.R-project.org/package=RColorBrewer

O'Brien HE, Parrent JL, Jackson JA, Moncalvo J, Vilgalys R. 2005. Fungal community

analysis by large-scale sequencing of environmental samples. *Applied and Environmental Microbiology* **71:** 5544–5550.

Oksanen J, Kindt R, Legendre P, O'Hara B. 2016. Vegan: community ecology package. R package version 2.5-5. https://cran.r-project.org/web/packages/vegan/index.html

Oze C, Fendorf S, Bird DK, Coleman RG. 2004. Chromium geochemistry in serpentinized ultramafic rocks and serpentine soils from the Franciscan complex of California. *American Journal of Science* **304:** 67–101.

Oze C, Skinner C, Schroth AW, Coleman RG. 2008. Growing up green on serpentine soils: biogeochemistry of serpentine vegetation in the Central Coast Range of California. *Applied Geochemistry* **23:** 3391–3403.

Parker IM, Gilbert GS. 2004. The evolutionary ecology of novel plant-pathogen interactions. *Annual Review of Ecology, Evolution, and Systematics* **35:** 675–700.

Parker IM, Saunders M, Bontrager M, Weitz AP, Hendricks R, Magarey R, Suiter K, Gilbert GS. 2015. Phylogenetic structure and host abundance drive disease pressure in communities. *Nature* **520:** 542–544.

Paul ND. 1989. Effects of fungal pathogens on nitrogen, phosphorous, and sulphur relations of individual plants and populations. In: Boddy L, Marchant R, Read DJ, eds. *Nitrogen, Phosphorus, and Sulphur Utilization by Fungi.* Cambridge, UK: Cambridge University Press, 155-180.

Pineda A, Zheng S-J, van Loon JJA, Pieterse CMJ, Dicke M. 2010. Helping plants to deal with insects: the role of beneficial soil-borne microbes. *Trends in Plant Science* **15:** 507-514.

Rajakaruna N, Baldwin BG, Chan R, Desrochers AM, Bohm BA, Whitton J. 2003. Edaphic

races and phylogenetic taxa in the *Lasthenia californica* complex (Asteraceae: Heliantheae): an hypothesis of parallel evolution. *Molecular Ecology* **12:** 1675–1679.

Rodrigues FL, Vieira JP. 2010. Feeding strategy of *Menticirrhus americanus* and *Menticirrhus littoralis* (Perciformes: Sciaenidae) juveniles in a sandy beach surf zone of southern Brazil. *Zoologia* **27(6):** 873-880.

Rizzo DM, Garbelotto M. 2003. Sudden oak death: endangering California nad Oregon forest ecosystems. *Frontiers in Ecology and the Environment* **1(5):** 197-204.

Romeis T, Ludwig AA, Martin R, Jones JDG. 2001. Calcium-dependent protein kinases play an essential role in a plant defence response. *EMBO Journal* **20:** 5556–5567.

Rossi J-P. 2011. rich: an R package to analyse species richness. *Diversity* **3:** 112–120.

Rottstock T, Joshi J, Kummer V, Fischer M. 2014. Higher plant diversity promotes higher diversity of fungal pathogens, while it decreases pathogen infection per plant. *Ecology* **95(7):** 1907-1917.

Seabloom EW, Borer ET, Gross K, Kendig AE, Lacroix C, Mitchell CE, Mordecai EA, Power AG. 2015. The community ecology of pathogens: coinfection, coexistence and community composition. *Ecology Letters* **18:** 401–415.

Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Sahl JW. 2009. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and Environmental Microbiology* **75(23):** 7537-7541.

Shaw BD, Hoch HC. 2000. Ca^{2+} regulation of *Phyllosticta ampelicida* pycnidiospore germination and appressorium formation. *Fungal Genetics and Biology* **31:** 43–53.

Shaw MW. 1986. Effects of temperature and leaf wetness on *Pyrenophera teres* growing on

barley cv. Sonja. *Plant Pathology* **35(3):** 294-309.

Snoeijers SS, Pérez-García A, Joosten MHAJ, De Wit PJGM. 2000. The effect of nitrogen on disease development and gene expression in bacterial and fungal plant pathogens. *Journal of Plant Pathology* **106:** 493-506.

Solomon PS, Tan KC, Oliver RP. 2003. The nutrient supply of pathogneic fungi; a fertile field for study. *Molecular Plant Pathology* **4(3):** 203-210.

Spear ER. 2017. Phylogenetic relationships and spatial distributions of putative fungal pathogens of seedlings across a rainfall gradient in Panama. *Fungal Ecology* **26:** 65–73.

Spear ER, Mordecai EA. 2018. Foliar pathogens do not stabilize coexistence of competing species in a California grassland. *Ecology* **99(10):** 2250–2259.

Sprent JFA. 1992. Parasites lost? *International Journal for Parasitology* **22:** 139–151.

Springer YP. 2009. Do extreme environments provide a refuge from pathogens? A phylogenetic test using serpentine flax. *American Journal of Botany* **96:** 2010–2021.

Springer YP, Hardcastle BA, Gilbert GS. 2006. Soil calcium and plant disease in serpentine ecosystems: a test of the pathogen refuge hypothesis. *Oecologia* **151:** 10–21.

Stukenbrock EH, McDonald BA. 2008. The origins of plant pathogens in agro-ecosystems. *Annual Review of Phytopathology.* **46:** 75–100.

Streit B, Hobbs RJ, Streit S. 1993. Plant distributions and soil chemistry at a serpentine/non-serpentine boundary in California. In *Markert BA*, editor. *Plants as biomonitors:indicators for heavy metals in the terrestrial environment.* Weinheim, Germany: VCH, 167-168 .

Telfer S, Lambin X, Birtles R, Beldomenico P, Burthe S, Paterson S, Begon M. 2010. Species interactions in a parasite community drive infection risk in a wildlife population. *Science* **330:** 243–247.

Treutter D. 2006. Significance of flavonoids in plant resistance: a review. *Environmental Chemistry Letters* **4:** 147–157.

Uhm K-H, Ahn I-P, Kim S, Lee Y-H. 2003. Calcium/calmodulin-dependent signaling for prepenetration development in *Colletotrichum gloeosporioides*. *Phytopathology* **93:** 82–87.

UNITE Community. 2017a. Full mothur UNITE+INSD dataset 2. Version 01.12.2017. https://doi.org/10.15156/BIO/587480

UNITE Community. 2017b. UNITE general FASTA release. Version 01.12.2017. https://doi.org/10.15156/BIO/587475

United States Department of Agriculture. 1960. Index of plant diseases in the United States. Agricultural handbook #165. U.S. Government Printing Office, Washington, DC.

Uricchio LH, Daws SC, Spear ER, Mordecai EA. 2019. Exotic species dominate due to niche overlap in a complex grassland community. *The American Naturalist* **193(2):** 213-226.

Vallano DM, Selmants PC, Zavaleta ES. 2012. Simulated nitrogen deposition enhances the performance of an exotic grass relative to native serpentine grassland competitors. *Plant Ecology* **213:** 1015–1026.

Vavrek MJ. 2011. Fossil: palaeoecological and palaeogeographical analysis tools. *Palaeontologia Electronica* **14**.

Velásquez AC, Castroverde CDM, He SY. 2018. Plant-pathogen warfare under changing climate conditions. *Current Biology* **28(10):** R619-R634.

Verosoglou SD, Barto EK, Menexes G, Rillig MC. 2013. Fertilization affects severity of disease caused by fungal plant pathogens. *Plant Pathology* **62(5):** 961-969.

Wang Y, Naumann U, Wright ST, Warton DI. 2012. mvabund – an R package for model-based analysis of multivariate abundance data. *Methods in Ecology and Evolution* **3:** 471-

474.

Warner RR, Chesson PL. 1985. Coexistence mediated by recruitment fluctuations: a field
guide to the storage effect. *The American Naturalist* **125:** 769–787.

Weather Underground. 2018. Jasper Ridge Biological Preserve KCAWOODS19.
https://www.wunderground.com/personal-weather-station/dashboard?ID=KCAWOODS19

Wickham H. 2007. Reshaping data with the reshape package. *Journal of Statistical Software*
21(12): 1-20.

Wickham H. 2017. tidyverse: Easily Install and Load the 'Tidyverse'. R package version 1.2.1.
https://CRAN.R-project.org/package=tidyverse.

Wilke CO. 2019. cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2'. R
package version 0.9.4. https://CRAN.R-project.org/package=cowplot.

Wolda H. 1981. Similarity indices, sample size and diversity. *Oecologia* **50:** 296–302.

Zhang Y, Ni J, Tang F, Pei K, Luo Y, Jian L, Sun L, Liang Y. 2016. Root-associated fungi of
Vaccinium carlesii in subtoprical forests of China: intra- and inter-annual variability and
impacts of human disturbances. *Scientific Reports* **6(1):** 22399.

Zhu Y, Chen H, Fan J, Wang Y, Li Y, Chen J, Fan J, Yang S, Hu L, Leung H, *et al.* **2000.**
Genetic diversity and disease control in rice. *Nature* **406(6797):** 718-722.

Zimmerman NB, Vitousek PM. 2012. Fungal endophyte communities reflect environmental
structuring across a Hawaiian landsYcape. *Proceedings of the National Academy of
Sciences* **109:** 13022-13027.

Figure legends

Figure 1. Pathogen damage on *Stipa pulchra* was consistently low across three years and both grassland types. (A) Locations of surveyed nonserpentine sites (blue) and serpentine sites (green) along the ridgetop at JRBP. **(B)** Example of fungal-caused pathogen foliar damage on *S. pulchra.* **(C)** Mean percentage (bars) and two standard errors (error bars) of diseased leaf area for surveyed serpentine and nonserpentine *S. pulchra,* by year. Damage was similar in both habitat types, but differed interannually. The asterisk indicates significantly lower damage in 2015 than in subsequent years of the survey.

Figure 2. Pathogens isolated from *Stipa pulchra* growing in nonserpentine and serpentine grasslands. Bipartite network showing species interactions between *S. pulchra* plants (left) growing in serpentine grasslands (green) and nonserpentine grasslands (blue) and fungal species cultured from symptomatic leaf tissue (white bars on the right). The length of each white bar indicates the number of isolates in the operational species unit (based on 97% sequence similarity). The thickness of lines connecting left and right bars represents the number of times a particular fungal species was isolated from *S. pulchra* in a particular grassland type. The nonserpentine fungi came from 15 genera, 8 families, 4 orders, and 3 classes; the serpentine fungi came from 20 genera, 14 families, 7 orders, and 5 classes (Tables 1, S1-S5).

Figure 3. Distinct fungal communities were isolated from *Stipa pulchra* growing in nearby serpentine and nonserpentine sites across years. Non-metric multidimensional scaling visualization of serpentine and nonserpentine fungal community similarity. Each point corresponds to the combined fungal community of two transects at the same site in the same year. Serpentine communities are shown in green and nonserpentine communities are shown in blue. Circles, triangles, and squares represent fungal communities sampled in 2015, 2016, and

2017, respectively. Ellipses enclose 95% confidence intervals for ordination of serpentine and nonserpentine communities. Community similarity decreases with the distance between points. PERMANOVA analysis showed a significant effect of soil type on fungal community composition ($F = 3.623$, $R^2 = 0.0618$, $p = 0.001$).

Figure 4. *Stipa pulchra* **foliar C and N content differs between habitat types, but does not predict mean pathogen damage. (A)** Foliar percent C, **(B)** percent N, and **(C)** C:N ratio versus mean percentage of diseased leaf area for serpentine and nonserpentine *S. pulchra* plants. Wilcoxon rank-sum tests indicated significantly higher foliar percent C and foliar percent N in nonserpentine compared to serpentine plants. Pearson's correlation tests showed no relationships between foliar tissue chemistry and the amount of symptomatic leaf tissue observed on individual plants.

Supporting Information

Fig. S1 Species accumulation curves for serpentine (green) and nonserpentine (blue) fungal communities isolated from *Stipa pulchra* leaves, for all samples over all years.

Fig. S2 Fungal species abundance by rank for serpentine (green) and nonserpentine (blue) fungal communities on *Stipa pulchra* leaves.

Table S1 Genus counts for serpentine and nonserpentine fungal communities isolated from *Stipa pulchra* leaves.

Table S2 Family counts for nonserpentine and serpentine fungal communities isolated from *Stipa pulchra* leaves.

Table S3 Order counts for nonserpentine and serpentine fungal communities isolated from *Stipa pulchra* leaves.

Table S4 Class counts for nonserpentine and serpentine fungal communities isolated from *Stipa pulchra* leaves.

Table S5 Presence and absence of genera in nonserpentine and serpentine fungal communities isolated from *Stipa pulchra* leaves, by year.

Table S6 ANOVA table for effects of habitat type, year, and habitat type x year interactions on site-level fungal community diversity in terms of Fisher's alpha.

Table S7 ANOVA table for effects of habitat type, year, and habitat type x year interactions on site fungal species richness.

Table S8 Pairwise PERMANOVA table for effects of year on fungal community composition.

Table S9 Results of chemical analysis of serpentine and nonserpentine *Stipa pulchra* foliar tissue.

Table S10 Analysis of deviance table for multivariate test of fungal community responses to

foliar C and N content.

Tables

Table 1. Pathogen damage on *Stipa pulchra* varied significantly by year, but not by habitat type. ANOVA results for log-transformed transect-level measurements of percentage of symptomatic living tissue on serpentine and nonserpentine *S. pulchra* (N = 3 sites per habitat type per year, with 12 transects per site per year, for a total of 36 serpentine and 36 nonserpentine transects over all three years). The first column lists the factor(s) tested; the second column degrees of freedom (df); the third column the sum of squares (SS); the fourth column the mean squared error (MSE); the fifth column the F-value; and the 6th column the p-value. P-values < 0.05 are marked with an asterisk.

Variables	df	SS	MSE	F value	p-value
habitat type	1	5.8	5.81	2.63	0.105
year	2	31.4	15.68	7.10	9.53×10^{-4}*
habitat type x site	4	20.2	5.05	2.29	0.0598
habitat type x year	2	9.9	4.96	2.25	0.107

Table 2. Fungal species isolated from symptomatic *Stipa pulchra* leaves were unevenly distributed across habitat types. Each row represents one of the species isolated. The columns indicate the isolation frequency for each species in each soil type. Asterisks indicate fungal species in genera not previously reported from plant hosts in the Poaceae; daggers indicate species not previously reported in association with any plant disease symptoms; double daggers indicate species in genera not previously reported in association with foliar disease symptoms.

Species	Nonserpentine	Serpentine
Alternaria cf. *infectoria*	26	21
Keissleriella sp. †	19	8
Drechslera cf. *fugax*	17	1
Stemphylium sp.	15	8
Ramularia cf. *proteae*	11	37
Drechslera sp.	9	1
Neoascochyta sp.	7	3
Cladosporium sp.	6	3
Alternaria cf. *eureka*	5	2
Parastagonospora cf. *avenae*	5	0
Pyrenophera cf. *chaeotmioides*	5	2
Pyrenophera cf. *lolii*	4	3
Phaeosphaerieaceae sp. 2	3	5
Pleosporaceae sp.	3	2
Drechslera cf. *nobleae*	1	5
Epicoccum sp.	1	0
Heyderia sp. *†	1	1
Parastagonospora cf. *cumpignensis*	1	0
Parastagonospora sp.	1	0
Phaeosphaeria sp.	1	2
Podospora sp.†	1	0
Phaeosphaerieaceae sp. 1	1	0
Chalastospora cf. *gossypii* ‡	0	1
Hyphodiscus cf. *brevicollaris* *†	0	1
Nemania cf. *serpens* *‡	0	1
Phaeophlebiopsis sp. *†	0	1
Plectania sp. *†	0	2
Pseudoseptoria sp.	0	1
Helotiales sp.	0	1
Wojnowiciella sp.*	0	1

Table 3. Fungal communities were distinct on *Stipa pulchra* plants growing in serpentine versus nonserpentine grasslands, across all years. Fungal community similarity by habitat type and year. The leftmost column lists the fungal communities considered, with the number of species in each community in parentheses. The second column lists Fisher's alpha for each community. The third column shows the number of shared species and, in parentheses, the percentage of total observed species that were shared for each pair of communities. The rightmost column lists the estimated Morisita-Horn community overlap value, based on absolute species abundances, for each pair of communities. The Morisita-Horn index ranges from 0 to 1, with 1 indicating complete overlap.

Fungal community pair	Fisher's alpha	Shared species	Morisita-Horn Similarity Index
All years serpentine (24)	9.324	16	0.675
All years nonserpentine (22)	7.261	(53.0%)	
2015 serpentine (11)	7.955	5	0.535
2015 nonserpentine (12)	5.205	(27.8%)	
2016 serpentine (16)	7.897	11	0.766
2016 nonserpentine (13)	5.861	(61.1%)	
2017 serpentine (12)	6.784	8	0.725
2017 nonserpentine (12)	5.136	(50.0%)	
2015 serpentine (11)	7.955	7	0.707
2016 serpentine (16)	7.897	(35.0%)	
2015 serpentine (11)	7.955	5	0.721
2017 serpentine (12)	6.784	(27.8%)	
2016 serpentine (16)	7.897	7	0.786
2017 serpentine (12)	6.784	(33.3%)	
2015 nonserpentine (12)	5.205	6	0.668
2016 nonserpentine (13)	5.861	(31.5%)	
2015 nonserpentine (12)	5.205	5	0.295
2017 nonserpentine (12)	5.136	(26.3%)	
2016 nonserpentine (13)	5.861	7	0.882
2017 nonserpentine (12)	5.136	(38.9%)	

Table 4. PERMANOVA results show significant effects of both habitat type and year on fungal foliar community composition (N = 18 serpentine and 18 nonserpentine sites). From left to right, columns list factors tested, degrees of freedom (df), sum of squares (SS), mean squared error (MSE), F-value, and p-value. Statistically significant p-values (p < 0.05) are marked with an asterisk. The distance factor compares communities at near (<600m) vs. far (>600m) distances from one another.

	df	SS	MSE	F-Value	p-value
habitat type	1	0.5700	0.56997	5.1119	0.003*
year	2	1.2611	0.63055	5.6552	0.001*
distance	1	0.0767	0.07667	0.5850	0.675
habitat type x year	2	0.3339	0.16697	1.4975	0.224
habitat type x distance	1	0.0515	0.05147	0.3927	0.815
distance x year	2	0.1697	0.08485	0.6473	0.734

Figures

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Supporting Information

Figure S1. Species accumulation curves for serpentine (green) and nonserpentine (blue) fungal communities isolated from *Stipa pulchra* leaves over three years of surveys. The cumulative number of unique fungal OTUs (based on 97% sequence similarity; y-axis) is plotted against the number of *S. pulchra* leaf tissue pieces sampled (x-axis).

Figure S2. Fungal species abundance by rank for serpentine (green) and nonserpentine (blue) fungal communities on *Stipa pulchra* leaves. Each point represents the number of times a single species was isolated from *S. pulchra* leaves in each habitat type.

Table S1. Genus counts for serpentine and nonserpentine fungal communities isolated from *Stipa pulchra* leaves. Numbers in each column represent the number of times a genus was isolated from *S. pulchra* leaves in each habitat type.

Genus	Nonserpentine	Serpentine
Alternaria	31	23
Drechslera	27	7
Keissleriella	19	8
Stemphylium	15	8
Ramularia	11	37
Pyrenophera	9	5
Neoascochyta	7	3
Parastagonospora	7	0
Cladosporium	6	3
Phaeosphaerieaceae	4	5
unknown Pleosporaceae	3	2
Epicoccum	1	0
Heyderia	1	1
Phaeosphaeria	1	2
Podospora	1	0
Chalastospora	0	1
Hyphodiscus	0	1
Nemania	0	1
Phaeophlebiopsis	0	1
Plectania	0	2
Pseudoseptoria	0	1
unkown Helotiales	0	1
Wojnowiciella	0	1

Table S2. Family counts for nonserpentine and serpentine fungal communities isolated from *Stipa pulchra* leaves. Numbers in each column represent the number of times a family was isolated from *S. pulchra* leaves in each habitat type.

Family	Nonserpentine	Serpentine
Pleosporaceae	85	45
Lentitheciaceae	19	8
Phaeosphaeriaceae	12	8
Mycosphaerellaceae	11	37
Didymellaceae	8	3
Cladosporiaceae	6	3
Helotiaceae	1	1
Lasiosphaeriaceae	1	0
Dothioraceae	0	1
Hyaloscyphaceae	0	1
Meruliaceae	0	1
Pleosporales *incertae sedis*	0	1
Sacrosomataceae	0	2
unknown Helotiales	0	1
Xylariaceae	0	1

Table S3. Order counts for nonserpentine and serpentine fungal communities isolated from *Stipa pulchra* leaves. Numbers in each column represent the number of times an order was isolated from *S. pulchra* leaves in each habitat type.

Order	Nonserpentine	Serpentine
Pleosporales	124	65
Capnodiales	17	40
Helotiales	1	3
Sordariales	1	0
Dothideales	0	1
Pezizales	0	2
Polyporales	0	1
Xylariales	0	1

Table S4. Class counts for nonserpentine and serpentine fungal communities isolated from *Stipa pulchra* leaves. Numbers in each column represent the number of times a class was isolated from *S. pulchra* leaves in each habitat type.

Class	Nonserpentine	Serpentine
Dothideomycetes	141	106
Leotiomycetes	1	3
Sordariomycetes	1	1
Agaricomycetes	0	1
Pezizomycetes	0	2

Table S5. Presence (P, shaded) or absence (A, unshaded) of genera in nonserpentine and serpentine fungal communities isolated from *Stipa pulchra* leaves, by year.

Genus	Nonserpentine 2015	Nonserpentine 2016	Nonserpentine 2017	Serpentine 2015	Serpentine 2016	Serpentine 2017
Alternaria	P	P	P	P	P	P
Chalastospora	A	A	A	P	A	A
Cladosporium	P	A	A	P	A	P
Drechslera	P	P	P	P	P	P
Epicoccum	A	A	P	A	A	A
Heyderia	A	A	P	A	A	P
Hyphodiscus	A	A	A	A	P	A
Keissleriella	P	P	P	P	P	P
Nemania	A	A	A	A	A	P
Neoascochyta	A	P	P	A	P	P
Parastagonospora	P	A	P	A	A	A
Phaeophlebiopsis	A	A	A	A	P	A
Phaeosphaeria	A	P	A	P	P	A
Plectania	A	P	A	A	A	P
Podospora	A	P	A	A	A	A
Pseudoseptoria	A	A	A	A	P	A
Pyrenophera	P	P	P	P	P	P
Ramularia	P	P	A	P	P	P
Stemphylium	A	P	P	P	A	P
unknown Phaeosphaerieaceae	P	P	A	A	P	A
unknown Pleosporaceae	A	P	P	A	P	P
unkown Helotiales	A	A	A	A	P	A
Wojnowiciella	A	A	A	P	A	A

Table S6. ANOVA table for effects of habitat type, year, and habitat type x year interactions on site-level fungal community diversity in terms of Fisher's alpha. Columns, from left to right, list the variable tested, the degrees of freedom, the sum of squares, the mean squared error, the F value, and the p-value.

	df	SS	MSE	F value	p-value
Habitat type	1	0.10	0.10	0.010	0.92
Year	2	3.63	1.82	0.18	0.84
Habitat type x Year	2	17.35	8.68	0.86	0.45

Table S7. ANOVA table for effects of habitat type, year, and habitat type x year interactions on site fungal species richness. Columns, from left to right, list the variable tested, the degrees of freedom, the sum of squares, the mean squared error, the F value, and the p-value.

	df	SS	MS	F value	p-value
Habitat type	1	6.72	6.72	1.07	0.32
Year	2	3.11	1.56	0.25	0.78
Habitat type x year	2	8.44	4.22	0.67	0.53

Table S8. Pairwise PERMANOVA table for effects of year on fungal community composition. From left to right, columns list the pair of years being compared, the F-value, R^2, and the Bonferroni-adjusted p-value. P-values < 0.05 are marked with an asterisk.

Pair	F-value	R^2	Bonferroni p-value
2015 vs 2016	2.616	0.207	0.114
2015 vs 2017	5.983	0.374	0.009*
2016 vs 2017	4.397	0.304	0.018*

Table S9. Results of chemical analysis of serpentine and nonserpentine *S. pulchra* foliar tissue. The Serpentine and Nonserpentine columns report elemental content in percentage of leaf tissue by dried weight. The standard error of each mean is listed in parentheses. W, in the third column, is the Wilcoxon-rank sum test statistic comparing serpentine and nonserpentine mean values. Asterisks denote statistically significant p-values ($p < 0.05$).

Element	Serpentine		Nonserpentine		W	p-value	
C	38.97	(3.44×10^{-1})	41.48	(2.35×10^{-1})	2713	1.65×10^{-7}	*
N	1.44	(4.96×10^{-2})	1.61	(3.84×10^{-2})	2346	1.10×10^{-3}	*
C:N	28.72	(8.74×10^{-1})	26.54	(6.18×10^{-1})	1398	6.62×10^{-2}	

Table S10. Analysis of deviance table for multivariate test of fungal community responses, based on fungal species presence/absence data, to foliar percent C, foliar percent N, foliar C:N ratio, and habitat type. Only fungal species represented by five or more isolates in 2016 were considered.

	df	deviance	p-value
Habitat type	96	20.93	0.01*
Foliar %C	95	7.98	0.35
Foliar %N	94	9.02	0.33
Foliar C:N	93	6.896	0.63