The genomic landscape of metastatic castration-resistant prostate cancers reveals multiple

distinct genotypes with potential clinical impact

Lisanne F. van Dessel[1,a], Job van Riet[1,2,3,a], Minke Smits[4], Yanyun Zhu[5,6], Paul Hamberg[7], Michiel S. van der Heijden[8,9,10], Andries M. Bergman[5,10], Inge M. van Oort[11], Ronald de Wit[1], Emile E. Voest[6,8,10], Neeltje Steeghs[8,10], Takafumi N. Yamaguchi[12], Julie Livingstone[12], Paul C. Boutros[12,13,14,15,16,17], John W.M. Martens[1,8], Stefan Sleijfer[1,8], Edwin Cuppen[18,19], Wilbert Zwart[5,6,20], Harmen J.G. van de Werken[2,3], Niven Mehra[4,b], Martijn P. Lolkema[1,8,b,*]

[1] Department of Medical Oncology, Erasmus MC Cancer Institute, Erasmus University Medical Center Rotterdam, Dr. Molewaterplein 40, 3015 GD, Rotterdam, the Netherlands.

[2] Cancer Computational Biology Center, Erasmus MC Cancer Institute, Erasmus University Medical Center Rotterdam, Dr. Molewaterplein 40, 3015 GD, Rotterdam, the Netherlands.

[3] Department of Urology, Erasmus MC Cancer Institute, Erasmus University Medical Center Rotterdam, Dr. Molewaterplein 40, 3015 GD, Rotterdam, the Netherlands.

[4] Department of Medical Oncology, Radboud University Nijmegen Medical Center, Geert Grooteplein Zuid 10, 6525 GA, Nijmegen, the Netherlands.

[5] Division on Oncogenomics, the Netherlands Cancer Institute, Plesmanlaan 121, 1066 CX, Amsterdam, the Netherlands.

[6] Oncode Institute, the Netherlands.

[7] Department of Internal Medicine, Franciscus Gasthuis & Vlietland, Kleiweg 500, 3045 PM, Rotterdam, the Netherlands.

[8] Center for Personalized Cancer Treatment, Rotterdam, the Netherlands.

[9] Division of Molecular Carcinogenesis, the Netherlands Cancer Institute, Plesmanlaan 121, 1066 CX, Amsterdam, the Netherlands.

[10] Department of Medical Oncology, the Netherlands Cancer Institute, Plesmanlaan 121, 1066 CX, Amsterdam, the Netherlands.

[11] Department of Urology, Radboud University Nijmegen Medical Center, Geert Grooteplein Zuid 10, 6525 GA, Nijmegen, the Netherlands.

[12] Informatics and Biocomputing Program, Ontario Institute for Cancer Research, Toronto, Ontario, Canada.

[13] Department of Medical Biophysics, University of Toronto, Toronto, Ontario, Canada.

[14] Department of Pharmacology and Toxicology, University of Toronto, Toronto, Ontario, Canada.

[15] Department of Human Genetics, University of California Los Angeles, Los Angeles, CA, USA.

[16] Department of Urology, University of California, Los Angeles, Los Angeles, CA, USA.

[17] Jonsson Comprehensive Cancer Centre, University of California, Los Angeles, Los Angeles, CA, USA.

[18] Center for Molecular Medicine and Oncode Institute, University Medical Center Utrecht, Heidelberglaan 100, 3584 CX, Utrecht, the Netherlands.

[19] Hartwig Medical Foundation, Science Park 408, 1098 XH, Amsterdam, the Netherlands.

[20] Laboratory of Chemical Biology and Institute for Complex Molecular Systems, Department of Biomedical Engineering, Eindhoven University of Technology, PO Box 513, 5600 MB, Eindhoven, the Netherlands.

[a,b] These authors contributed equally

[*] Corresponding author

Address for correspondence: Martijn P. Lolkema. Department of Medical Oncology, Erasmus MC

Cancer Institute, Room NT-537, P.O. Box 2040, 3000 CA, Rotterdam, The Netherlands. Tel.: +31 10

704 19 06. E-mail: m.lolkema@erasmusmc.nl

Keywords: prostate cancer, metastasis, cancer genomics, homologous recombination deficiency

Abstract

Metastatic castration-resistant prostate cancer (mCRPC) has a highly complex genomic landscape.

With the recent development of novel treatments, accurate stratification strategies are needed. Here

we present the whole-genome sequencing (WGS) analysis of fresh-frozen metastatic biopsies from

197 mCRPC patients. Using unsupervised clustering based on genomic features, we define eight

distinct genomic clusters. We observe potentially clinically relevant genotypes, including microsatellite

instability (MSI), homologous recombination deficiency (HRD) enriched with genomic deletions and

BRCA2 aberrations, a tandem duplication genotype associated with *CDK12$^{-/-}$* and a chromothripsis-

enriched subgroup. Our data suggests that stratification on WGS characteristics may improve

identification of MSI, *CDK12$^{-/-}$* and HRD patients. From WGS and ChIP-seq data, we show the

potential relevance of recurrent alterations in non-coding regions identified with WGS and highlight

the central role of AR signaling in tumor progression. These data underline the potential value of

using WGS to accurately stratify mCRPC patients into clinically actionable subgroups.

Introduction

Prostate cancer is known to be a notoriously heterogeneous disease and the genetic basis for this interpatient heterogeneity is poorly understood[1,2]. The ongoing development of new therapies for metastatic prostate cancer that target molecularly defined subgroups further increases the need for accurate patient classification and stratification[3–5]. Analysis of whole-exome sequencing data of metastatic prostate cancer tumors revealed that 65% of patients had actionable targets in non-androgen receptor related pathways, including PI3K, Wnt and DNA repair[6]. Several targeted agents involved in these pathways, including mTOR/AKT pathway inhibitors[7] and PARP inhibitors[8], are currently in various phases of development and the first clinical trials show promising results. Therefore, patients with metastatic prostate cancer could benefit from better stratification to select the most appropriate therapeutic option. More extensive analysis using whole-genome sequencing (WGS) based classification of tumors may be useful to improve selection of patients for different targeted therapies. The comprehensive nature of WGS has many advantages including the detection of mutational patterns, as proven by the successful treatment of patients with high tumor mutational burden with immune checkpoint blockade therapy[9–12]. Moreover, WGS unlike exome sequencing, can detect structural variants and aberrations in non-coding regions, both important features of prostate cancer.

The stratification of prostate cancer patients, based on differences in the mutational landscape of their tumors, has mainly focused on mutually-exclusive mutations, copy number alterations or distinct patterns in RNA-sequencing caused by the abundant TMPRSS2-ERG fusion, which is recurrent in 50% of primary prostate tumors[6,13–18]. More recently, WGS of metastatic prostate cancer tumors demonstrated that structural variants arise from specific alterations such as $CDK12^{-/-}$ and $BRCA2^{-/-}$ genotypes and are strongly associated with genome-wide events such as large tandem duplications or small genomic deletions, respectively[19–23]. Advances in WGS and interpretation have revealed rearrangement signatures in breast cancer relating to disease stage, homologous recombination deficiency (HRD) and $BRCA1/BRCA2$ defects based on size and type of structural variant[22,24]. Thus, WGS enables the identification of patterns of DNA aberrations (i.e. genomic scars) that may profoundly improve classification of tumors that share a common etiology, if performed in a sufficiently powered dataset.

In this study, we analyzed the WGS data obtained from 197 metastatic castration-resistant

prostate cancer (mCRPC) patients. We describe the complete genomic landscape of mCRPC,

including tumor specific single and multi-nucleotide variants (SNVs and MNVs), small insertions and

deletions (InDels), copy number alterations (CNAs), mutational signatures, kataegis, chromothripsis

and structural variants (SVs). Next, we compared the mutational frequency of the detected driver

genes and genomic subgroups with an unmatched WGS cohort of primary prostate cancer (n = 210),

consisting of exclusively of Gleason score 6-7 tumors[15,25]. We investigated the presence of possible

driver genes by analyzing genes with enriched (non-synonymous) mutational burdens and recurrent

or high-level copy number alterations[26,27]. By utilizing various basic genomic features reflecting

genomic instability and employing unsupervised clustering, we were able to define eight distinct

genomic subgroups of mCRPC patients. We combined our genomic findings with AR, FOXA1 and

H3K27me ChIP-seq data and confirmed that important regulators of AR-mediated signaling are

located in non-coding regions with open chromatin and highlight the central role of AR signaling in

tumor progression.

Results

Characteristics of the mCRPC cohort and sequencing approach

We analyzed fresh-frozen metastatic tumor samples and matched blood samples from 197

castration-resistant prostate cancer (CRPC) patients using WGS generating to date the largest WGS

dataset for mCRPC (Figure 1a). Clinical details on biopsy site, age and previous treatments of the

included patients are described in Figure 1b-c and Supplementary table 2. WGS data was sequenced

to a mean coverage of 104X in tumor tissues and 38X in peripheral blood (Supplementary figure 1a).

The median estimated tumor cell purity using *in silico* analysis of our WGS data was 62% (range: 16-

96%; Supplementary figure 1b). Tumor cell purity correlated weakly with the frequency of called SNVs

(Spearman correlation; rho = 0.2; *p* = 0.005), InDels (Spearman correlation; rho = 0.35; *p* < 0.001),

MNVs (Spearman correlation; rho = 0.25; *p* < 0.001) and structural variants (Spearman correlation;

rho   =   0.22;   *p*   =   0.002;   Supplementary   figure   1c).

Landscape of mutational and structural variants in mCRPC

The median tumor mutational burden (TMB) at the genomic level (SNVs and InDels per Mbp)

was 2.7 in our mCRPC cohort, including 14 patients with high TMB (> 10). We found a median of

6,621 single-nucleotide variants (SNVs; IQR: 5,048-9,109), 1,008 small insertions and deletions

(InDels; IQR: 739-1,364), 55 multi-nucleotide variants (MNVs; IQR: 34-86) and 224 SVs (IQR: 149-

370) per patient (Supplementary figure 2a-c). We observed a highly complex genomic landscape

consisting of multiple driver mutations and structural variants in our cohort.

We confirmed that known driver genes of prostate cancer were enriched for non-synonymous

mutations (Figure 2 and Supplementary figure 2e)[13,15,28]. In total, we detected 11 genes enriched with

non-synonymous mutations: *TP53, AR, FOXA1, SPOP, PTEN, ZMYM3, CDK12, ZFP36L2, PIK3CA*

and *APC. ATM* was mutated in 11 samples, but after multiple-testing correction appeared not to be

enriched.

Our copy number analysis revealed distinct amplified genomic regions including 8q and Xq

and deleted regions including 8p, 10q, 13q and 17p (Supplementary figure 2d). Well-known prostate

cancer driver genes[8,16], such as *AR*, *PTEN*, *TP53* and *RB1*, are located in these regions. In addition

to large-scale chromosomal copy number alterations, we could identify narrow genomic regions with

recurrent copy number alterations across samples which could reveal important prostate cancer driver

genes (Supplementary table 3).

TMPRSS2-ERG gene fusions were the most common fusions in our cohort (n = 84 out of

197; 42.6%) and were the majority of *ETS* family fusions (n = 84 out of 95; 88.4%; Figure 2 and

Supplementary figure 3). This is comparable to primary prostate cancer, where *ETS* fusions are found

in approximately 50% of tumors[13,15]. The predominant break point was located upstream of the

second exon of *ERG*, which preserves its *ETS*-domain in the resulting fusion gene.

In 42 patients (21.3%), we observed regional hypermutation (kataegis; Figure 2 and

Supplementary figure 4). In addition, we did not observe novel mutational signatures specific for

metastatic disease or possible pre-treatment histories (Supplementary figure 5)[29].

To further investigate whether our description of the genome-wide mutational burden and

observed alterations in drivers and/or subtype-specific genes in mCRPC were metastatic specific, we

compared our data against an unmatched WGS cohort of primary prostate cancer (n = 210)[15,25],

consisting of Gleason score 6-7 disease. Comparison of the median genome-wide TMB (SNVs and

InDels per Mbp) revealed that the TMB was roughly 3.8 times higher in mCRPC (Figure 3a) and the

frequency of structural variants was also higher (Figure 3b) between disease stages, increasing as

disease progresses. Analysis on selected driver and subtype-specific genes showed that the

mutational frequency of several genes (*AR*, *TP53*, *MYC*, *ZMYM3*, *PTEN*, *PTPRD*, *ZFP36L2*,

ADAM15, *MARCOD2*, *BRIP1*, *APC*, *KMT2C*, *CCAR2*, *NKX3-1*, *C8orf58* and *RYBP*) was significantly

altered (q ≤ 0.05) between the primary and metastatic cohorts (Figure 3c-e). All genes for which we

observed significant differences in mutational frequency, based on coding mutations, were enriched in

mCRPC (Figure 3d). We did not identify genomic features that were specific for the metastatic setting,

beyond androgen deprivation therapy-specific aberrations revolving *AR* (no aberrations in hormone-

sensitive setting versus 137 aberrations in castration-resistant setting). We cannot exclude from these

data that matched sample analysis or larger scale analysis could reveal such aberrations.

We next determined whether previous treatments affect the mutational landscape. Using

treatment history information, we grouped prior secondary anti-hormonal therapy, taxane-based

chemotherapy and systemic radionucleotide therapy into different groups (Supplementary figure 6).

This analysis did not reveal systematic biases due to pre-treatment in aberrations, such as TMB,

kataegis, chromothripsis, *ETS* fusions, or somatically altered genes (Supplementary table 3).

The role of the AR-pathway in mCRPC

Focusing on the AR-pathway revealed that aberrant AR signaling occurred in 80% of our patients. In 57.3% of patients both *AR* and the *AR*-enhancer (~66.13 Mb on chromosome X; located about 629 kbp upstream of the *AR* gene[20]) were affected (Figure 4a). In an additional 6.6% and 14.7% of tumors only *AR* gene alterations or *AR*-enhancer amplification occurred, respectively. The percentage of mCRPC patients with the exclusive *AR*-enhancer amplification (29 out of 197; 14.7%) versus exclusively *AR*-locus amplification (13 out of 197; 6.6%) is similar to previous observations which showed 21 out of 94 CRPC patients (10.3%) with exclusively *AR*-enhancer amplification versus 4 out of 94 CRPC patients (4.3%) with exclusively AR-locus amplification[20]. Thus, concurrent amplification of the *AR* gene and the *AR*-enhancer was not necessarily of equal magnitude, which resulted in differences in copy number enrichment of these loci (Figure 4b).

To date no AR ChIP-seq data has been reported in human mCRPC samples and evidence of increased functional activity of the amplified enhancer thus far is based on cell line models[30]. To resolve this, we performed AR ChIP-seq on two selected mCRPC patient samples with *AR*-enhancer amplification based on WGS data. As controls we used two prostate cancer cell-lines (LNCaP and VCaP) and three independent primary prostate cancer samples that did not harbor copy number alterations at this locus (Supplementary figure 7)[31]. We observed active enhancer regions (H3K27ac) in the castration-resistant setting, co-occupied by AR and FOXA1, at the amplified *AR*-enhancer, which is substantially stronger when compared to the hormone-sensitive primary prostate cancer samples without somatic amplifications (Figure 4c and Supplementary figure 7). Furthermore, a recurrent focal amplification in a non-coding region was observed at 8q24.21 near *PCAT1*. This locus bears similar epigenetic characteristics to the *AR*-enhancer with regard to H3K27ac and, to a lesser extent, binding of AR and/or FOXA1 in the mCRPC setting (Figure 4d and Supplementary figure 7).

WGS-based stratification defines genomic subgroups in mCRPC

Our comprehensive WGS data and large sample size enabled us to perform unsupervised clustering on several WGS characteristics to identify genomic scars that can define subgroups of mCRPC patients. We clustered our genomic data using the total number of SVs, relative frequency of SV events (translocations, inversions, insertions, tandem duplications and deletions), genome-wide

TMB encompassing SNV, InDels and MNV, and tumor ploidy. Prior to clustering, we subdivided

tandem duplications and deletions into two major categories based on the respective genomic size of

the aberration (smaller and larger than 100 kbp) since previous studies revealed distinctions based on

similar thresholds for these structural variants in relation to specific mutated genes[19–21,32]. Similarly,

we observed a difference in genomic size and number in our subgroups of mCRPC patients

(Supplementary figure 8).

This analysis defined eight distinct subgroups (Figure 5-6 and Supplementary figure 8-11): A)

Microsatellite Instability (MSI) signature with high TMB and association with mismatch repair

deficiency; B) Tandem duplication (> 100 kbp) phenotype associated with biallelic CDK12 inactivation;

D) Homologous Recombination Deficiency (HRD) features with many (> 100 kbp) deletions and

association with (somatic) mutations in BRCAness-associated genes; this was supported by high HR-

deficiency scores (CHORD; Supplementary figure 8-9); F) chromothripsis; C, E, G, H); non-significant

genomic signature without any currently known biological association. Table 1 summarizes the key

features of each subgroup.

Clusters A and B represent previously identified genomic subgroups (MSI and CDK12[-/-])

[6,19,21,33]. In cluster B only two patients were allocated to this subgroup without a specific mutation in

the identifying gene. The well-known mismatch repair genes: MLH1, MSH2 and MSH6 are among the

cluster-specific mutated genes in cluster A (Figure 6a). Twelve out of these thirteen patients had at

least one inactivating alteration in one of these genes (Figure 6b). Interestingly, cluster B (CDK12[-/-])

harbors two patients without non-synonymous CDK12 mutation or copy number alteration; the cause

of their tandem duplication phenotype is currently unknown (Figure 6b). Cluster D shows significant

features of HRD, specifically biallelic BRCA2 inactivation (Supplementary figure 13), mutational

signature 3, enrichment of deletions (< 100 kbp) and is supported by high HR-deficiency scores

(CHORD) (Supplementary figure 8-9)[22,34]. Remarkably, seven out of twenty-two patients did not have

a biallelic BRCA2 inactivation. However, four of these patients showed at least one (deleterious)

aberration in other BRCAness-related genes (Figure 6b)[35]. Cluster F was enriched for chromothripsis

events, however we could not reproduce a previous finding suggesting chromothripsis was associated

with inversions and p53 inactivation in prostate cancer[21]. Apart from the chromothripsis events, no

clear gene aberration was associated with this cluster (Figure 6b). In the remaining patients, there

were no distinct genomic signatures or biologic rationale for patient clustering (cluster C, E, G, H). In

cluster C, conjoint aberrations of *BRCA1* and *TP53* were observed in one patient with a high HR-

deficiency prediction score (CHORD), which is known to lead to a small tandem duplication

phenotype (< 100 kbp)[32]. Two other patients within cluster C displayed a weak CHORD scoring

associated with HR-deficiency, however no additional definitive evidence was found for a *BRCA1*

loss-of-function mutation within these patients.

In addition to our unsupervised clustering approach, we clustered our samples using the

clustering scheme proposed by TCGA (Supplementary figure 12a), which defines seven clusters

based on coding mutations and copy number aberrations in *SPOP*, *FOXA1*, *IDH1* and *ETS* family

gene fusions (and overexpression) per promiscuous partner (*ERG*, *ETV1*, *ETV4* and *FLI1*)[13].

Unfortunately, we currently lack matched mRNA-sequencing data in our cohort and therefore cannot

observe overexpression of fused *ETS* family members which restricted us to only characterize the

genomic breaks of these promiscuous partners. Without incorporation of *ETS* family overexpression,

this proposed clustering scheme categorizes 61% of mCRPC into these seven groups versus 68% of

the original cohort containing primary prostate cancer described by TCGA (Supplementary figure

12b)[13]. There was no significant correlation between the TCGA clustering scheme and our defined

genomic subtypes such as MSI, BRCAness or $CDK12^{-/-}$. In addition, we did not detect statistical

enrichment or depletion ($q \leq 0.05$) between these supervised clusters and additional mutated genes,

kataegis and chromothripsis, only the known enrichment of homozygous *CHD1* deletions in the

SPOP-cluster was observed[13].

Performing unsupervised clustering and principal component analysis on the primary prostate

cancer and metastatic cohorts revealed no striking primary-only genomic subgroup nor did we detect

the presence of the mCRPC-derived genomic subgroups in the primary prostate cancer cohort

(Supplementary figure 14). This could reflect the absence of *CDK12* mutations and the presence of

only three sporadic *BRCA2* mutated samples (1%) in the primary prostate cancer cohort.

Furthermore, only one sample (1%) with MSI-like and high TMB (> 10), respectively, was observed in

the primary cancer cohort. Indeed, there is a striking difference in the mutational load between both

disease settings.

Discussion

We performed WGS of metastatic tumor biopsies and matched normal blood obtained from 197 patients with mCRPC to provide an overview of the genomic landscape of mCRPC. The size of our cohort enables classification of patients into distinct disease subgroups using unsupervised clustering. Our data suggest that classification of patients using genomic events, as detected by WGS, improves patient stratification, specifically for clinically actionable subgroups such as BRCA deficient and MSI patients. Furthermore, we confirm the central role of AR signaling in mCRPC that mediates its effect through regulators located in non-coding regions and the apparent difference in primary versus metastatic prostate cancers.

The classification of patients using WGS has the advantage of being, in theory, more precise in determining genomically defined subgroups in prostate cancer compared to analyses using targeted panels consisting of a limited number of genes, or exome sequencing. The identification of subgroups based on predominant phenotypic characteristics encompassing genomic signatures may be clinically relevant and our clustering analysis refines patient classification. In cluster A, we observed a high TMB which has been associated in other tumor types with a high sensitivity to immune check-point inhibitors[9,11,12]. Clinical trials using pembrolizumab in selected mCRPC patients are underway (KEYNOTE-028, KEYNOTE-199)[36,37]. Interestingly, in both cluster B and cluster D, we identified patients that do not have the defining biallelic *CDK12* or *BRCA2* (somatic) mutation. Such patients might be deemed false-negatives or false-positives when using FDA-approved assays (BRCAnalysis™ and FoundationFocus™), currently used in breast cancer diagnosis, to predict response to poly(ADP-ribose) polymerase (PARP) inhibitors and/or platinum compounds based on the presence of *BRCA1/2* mutations. The first clinical trials combining PARP inhibitors with AR-targeted therapies in mCRPC show promising results[8]. Thus, WGS based stratification may improve the patient classification of DNA repair deficient tumors as it uses the genome wide scars caused by defective DNA repair to identify tumors that have these deficiencies.

The use of WGS also allowed us to gain more insight into the role of non-coding regions of the genome in prostate cancer. We confirmed the amplification of a recently reported *AR*-enhancer[20,21,30]. In line with the cell line-based observations, we show AR binding at these mCRPC-

specific enhancer regions, providing the first clinical indication that *AR*-enhancer amplification also

increases AR signaling in mCRPC tumors. These findings are supported by previous studies

demonstrating that this amplification ultimately resulted in significantly elevated expression of *AR*

itself[20,21,30]. Furthermore, we confirm a recurrent focal amplification near *PCAT1*, which shows robust

chromatin binding for AR in mCRPC samples, providing clinical proof-of-concept of a functional

enhancer that is also active and AR-bound in cell line models. Recent research elucidated to the

functional importance of this region in regulating *MYC* expression in prostate cancer, which could

highlight a putative role of this somatically-acquired amplification[31]. However, the WGS and ChIP-seq

data presented here are not conclusive in elucidating the definitive role of this amplified region in

regulating *MYC* expression and further mechanistic studies are needed to establish a potential link to

MYC regulation.

In addition, *PCAT1* is a long non-coding RNA which is known to be upregulated in prostate

cancer and negatively regulates *BRCA2* expression while positively affecting *MYC* expression[38,39].

Combining our WGS approach with AR, FOXA1 and H3K27ac ChIP-seq data, we identify non-coding

regions affecting both *AR* itself, and possibly *MYC*, through *AR*-enhancer amplification as a potential

mechanism contributing to castration resistance.

A potential pitfall of our clustering analysis is the selection of features used; for this we made

a number of assumptions based on the literature and distribution of the structural variants within our

cohort[19–21,32]. As the input features and weights for clustering analysis are inherent to the clustering

outcome, we performed additional clustering analyses using various combinations of these features

and applied alternative approaches but did not detect striking differences compared to the current

approach. Another potential pitfall of the employed hierarchical clustering scheme is that patients are

only attributed to a single cluster. An example of this can be seen in cluster A where a patient is

grouped based on its predominant genotype (MSI) and associated mutations in MMR-related genes

(*MLH1*, *POLE*, *POLD3* and *BLM*), but this sample also displays an increased number of structural

variants and increased ploidy status and harbors a pathogenic *BRCA2* mutation. However, it is

missing the characteristic number of genomic deletions (< 100 kbp) and BRCA mutational signature

associated with *BRCA2*$^{-/-}$ samples that define cluster D. Despite these pitfalls we conclude that

unbiased clustering contributes towards improved classification of patients.

The CPCT-02 study was designed to examine the correlation of genomic data with treatment outcome after biopsy at varying stages of disease. Our cohort contains patients with highly variable pre-treatment history and since the treatments for mCRPC patients nowadays significantly impacts overall survival, the prognosis of patients differs greatly. Therefore, correlation between genomic data and clinical endpoints, such as survival is inherently flawed due to the very heterogeneous nature of the patient population. Moreover, our analysis comparing primary and metastatic samples shows a significant increase in the number of genomic aberrations with advancing disease, meaning that the difference in timing of the biopsies may bias the prognostic value of the data. In future studies we plan to gather all known clinically defined prognostic information and determine whether the genomic subtypes increase the ability to predict outcome. Unfortunately, some clinical parameters with prognostic importance such as ethnicity will not be available due to ethical regulations. Moreover, we will increase the sample size, in order to correlate genomic features to clinical parameters to better determine whether the subtypes we identified are stable over time. Therefore, we are currently unable to present meaningful correlations between clinical endpoints and the clusters we identified.

Overall, we show the added value of WGS-based unsupervised clustering in identifying patients with genomic scars who are eligible for specific therapies. Since our clustering method does not rely on one specific genetic mutation we are able to classify patients even when WGS (or our methodology) does not find conclusive evidence for (bi-allelic) mutations in the proposed gene-of-interest. Further research should validate clinical response and outcome on specific therapies in matched subgroups. This study also shows that a large population of mCRPC patients do not fall into an as-of-yet clinically-relevant or biologically-clear genotype and further research can help elucidate the oncogenic driver events and provide new therapeutic options.

 METHODS

Methods, including statements of data availability and any associated accession codes and references, are available at https://doi.org/XXXXXXXXXXXXX.

References

1. Boyd, L. K., Mao, X. & Lu, Y. J. The complexity of prostate cancer: Genomic alterations and heterogeneity. *Nature Reviews Urology* (2012). doi:10.1038/nrurol.2012.185

2. Wei, L. *et al.* Intratumoral and Intertumoral Genomic Heterogeneity of Multifocal Localized Prostate Cancer Impacts Molecular Classifications and Genomic Prognosticators. *Eur. Urol.* (2017). doi:10.1016/j.eururo.2016.07.008

3. Mullane, S. A. & Van Allen, E. M. Precision medicine for advanced prostate cancer. *Current Opinion in Urology* (2016). doi:10.1097/MOU.0000000000000278

4. Ciccarese, C. *et al.* Prostate cancer heterogeneity: Discovering novel molecular targets for therapy. *Cancer Treatment Reviews* (2017). doi:10.1016/j.ctrv.2017.02.001

5. Shtivelman, E., Beer, T. M. & Evans, C. P. Molecular pathways and targets in prostate cancer. *Oncotarget* (2014). doi:10.18632/oncotarget.2406

6. Robinson, D. *et al.* Integrative clinical genomics of advanced prostate cancer. *Cell* **161**, 1215–1228 (2015).

7. Chow, H. *et al.* A phase 2 clinical trial of everolimus plus bicalutamide for castration-resistant prostate cancer. *Cancer* (2016). doi:10.1002/cncr.29927

8. Clarke, N. *et al.* Olaparib combined with abiraterone in patients with metastatic castration-resistant prostate cancer: a randomised, double-blind, placebo-controlled, phase 2 trial. *Lancet Oncol.* (2018). doi:10.1016/S1470-2045(18)30365-6

9. Yarchoan, M., Hopkins, A. & Jaffee, E. M. Tumor Mutational Burden and Response Rate to PD-1 Inhibition. *N. Engl. J. Med.* (2017). doi:10.1056/NEJMc1713444

10. Chan, T. A. *et al.* Development of tumor mutation burden as an immunotherapy biomarker: Utility for the oncology clinic. *Annals of Oncology* (2019). doi:10.1093/annonc/mdy495

11. Rizvi, N. A. *et al.* Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science (80-.).* (2015). doi:10.1126/science.aaa1348

12. Samstein, R. M. *et al.* Tumor mutational load predicts survival after immunotherapy across

multiple cancer types. *Nature Genetics* (2019). doi:10.1038/s41588-018-0312-8

13.   Cancer Genome Atlas Research Network *et al.* The Molecular Taxonomy of Primary Prostate

Cancer. *Cell* **163**, 1011–25 (2015).

14.   Angeles, A. K., Bauer, S., Ratz, L., Klauck, S. M. & Sültmann, H. Genome-Based

Classification and Therapy of Prostate Cancer. *Diagnostics* **8**, (2018).

15.   Fraser, M. *et al.* Genomic hallmarks of localized, non-indolent prostate cancer. *Nature* **541**,

359–364 (2017).

16.   Schoenborn, J. R., Nelson, P. & Fang, M. Genomic profiling defines subtypes of prostate

cancer with the potential for therapeutic stratification. *Clin. Cancer Res.* (2013).

doi:10.1158/1078-0432.CCR-12-3606

17.   Nam, R. K. *et al.* Expression of TMPRSS2 ERG gene fusion in prostate cancer cells is an

important prognostic factor for cancer progression. *Cancer Biol. Ther.* (2007).

doi:10.4161/cbt.6.1.3489

18.   Tomlins, S. A. *et al.* Role of the TMPRSS2-ERG gene fusion in prostate cancer. *Neoplasia*

(2008).

19.   Wu, Y.-M. *et al.* Inactivation of CDK12 Delineates a Distinct Immunogenic Class of Advanced

Prostate Cancer Article Inactivation of CDK12 Delineates a Distinct Immunogenic Class of

Advanced Prostate Cancer. *Cell* **173**, 1770-1782.e14 (2018).

20.   Viswanathan, S. R. *et al.* Structural Alterations Driving Castration-Resistant Prostate Cancer

Revealed by Linked-Read Genome Sequencing. *Cell* **174**, 433-447.e19 (2018).

21.   Quigley, D. A. *et al.* Genomic Hallmarks and Structural Variation in Metastatic Prostate

Cancer. *Cell* **174**, 758-769.e9 (2018).

22.   Nik-Zainal, S. *et al.* Landscape of somatic mutations in 560 breast cancer whole-genome

sequences. *Nature* **534**, 47–54 (2016).

23.   Taylor, R. A. *et al.* Germline BRCA2 mutations drive prostate cancers with distinct evolutionary

trajectories. *Nat. Commun.* (2017). doi:10.1038/ncomms13671

24.   Davies, H. *et al.* Whole-genome sequencing reveals breast cancers with mismatch repair

deficiency. *Cancer Res.* (2017). doi:10.1158/0008-5472.CAN-17-1083

25.   Espiritu, S. M. G. *et al.* The Evolutionary Landscape of Localized Prostate Cancers Drives

Clinical Aggression. *Cell* (2018). doi:10.1016/j.cell.2018.03.029

26.   Martincorena, I. *et al.* Universal Patterns of Selection in Cancer and Somatic Tissues. *Cell*

393   **171**, 1029-1041.e21 (2017).

394   27.   Mermel, C. H. *et al.* GISTIC2.0 facilitates sensitive and confident localization of the targets of

395   focal somatic copy-number alteration in human cancers. *Genome Biol.* **12**, R41 (2011).

396   28.   Armenia, J. *et al.* The long tail of oncogenic drivers in prostate cancer. *Nat. Genet.* (2018).

397   doi:10.1038/s41588-018-0078-z

398   29.   Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**, 415–

399   421 (2013).

400   30.   Takeda, D. Y. *et al.* A Somatically Acquired Enhancer of the Androgen Receptor Is a

401   Noncoding Driver in Advanced Prostate Cancer. *Cell* (2018). doi:10.1016/j.cell.2018.05.037

402   31.   Mazrooei, P. *et al.* Somatic Mutations and Risk-Variants Converge on Cis-Regulatory

403   Elements to Reveal the Cancer Driver Transcription Regulators in Primary Prostate Tumors.

404   *SSRN Electron. J.* (2018). doi:10.2139/ssrn.3245213

405   32.   Menghi, F. *et al.* The Tandem Duplicator Phenotype Is a Prevalent Genome-Wide Cancer

406   Configuration Driven by Distinct Gene Mutations. *Cancer Cell* **34**, 197-210.e5 (2018).

407   33.   Pritchard, C. C. *et al.* Complex MSH2 and MSH6 mutations in hypermutated microsatellite

408   unstable advanced prostate cancer. *Nat. Commun.* **5**, (2014).

409   34.   Polak, P. *et al.* A mutational signature reveals alterations underlying deficient homologous

410   recombination repair in breast cancer. *Nat. Genet.* (2017). doi:10.1038/ng.3934

411   35.   Lord, C. J. & Ashworth, A. BRCAness revisited. *Nat. Rev. Cancer* **16**, 110–120 (2016).

412   36.   Hansen, A. R. *et al.* Pembrolizumab for advanced prostate adenocarcinoma: Findings of the

413   KEYNOTE-028 study. *Ann. Oncol.* (2018). doi:10.1093/annonc/mdy232

414   37.   De Bono, J. S. *et al.* KEYNOTE-199: Pembrolizumab (pembro) for docetaxel-refractory

415   metastatic castration-resistant prostate cancer (mCRPC). *J. Clin. Oncol.* (2018).

416   doi:10.1200/JCO.2018.36.15_suppl.5007

417   38.   Prensner, J. R. *et al.* The Long Non-Coding RNA PCAT-1 Promotes Prostate Cancer Cell

418   Proliferation through cMyc. *Neoplasia* **16**, 900–908 (2014).

419   39.   Prensner, J. R. *et al.* PCAT-1, a long noncoding RNA, regulates BRCA2 and controls

420   homologous recombination in cancer. *Cancer Res.* **74**, 1651–1660 (2014).

40.   Priestley, P. *et al.* Pan-cancer whole genome analyses of metastatic solid tumors. *bioRxiv*

415133 (2018). doi:10.1101/415133

41.   Broad Institute. Picard tools. *https://broadinstitute.github.io/picard/* (2016). Available at:

https://broadinstitute.github.io/picard/%5Cnhttp://broadinstitute.github.io/picard/.

42.   Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**, 285–

291 (2016).

43.   Casper, J. *et al.* The UCSC Genome Browser database: 2018 update. *Nucleic Acids Res.* **46**,

D762–D769 (2018).

44.   Forbes, S. A. *et al.* COSMIC: Somatic cancer genetics at high-resolution. *Nucleic Acids Res.*

45, D777–D783 (2017).

45.   Tamborero, D. *et al.* Cancer Genome Interpreter annotates the biological and clinical

relevance of tumor alterations. *Genome Med.* **10**, 25 (2018).

46.   Griffith, M. *et al.* CIViC is a community knowledgebase for expert crowdsourcing the clinical

interpretation of variants in cancer. *Nat. Genet.* **49**, 170–174 (2017).

47.   Harrow, J. *et al.* GENCODE: The reference human genome annotation for the ENCODE

project. *Genome Res.* **22**, 1760–1774 (2012).

48.   Davies, H. *et al.* HRDetect is a predictor of BRCA1 and BRCA2 deficiency based on

mutational signatures. *Nat. Med.* **23**, 517–525 (2017).

49.   Huber, W., Toedling, J. & Steinmetz, L. M. Transcript mapping with high-density

oligonucleotide tiling arrays. *Bioinformatics* **22**, 1963–1970 (2006).

50.   Gel, B. & Serra, E. karyoploteR: an R/Bioconductor package to plot customizable genomes

displaying arbitrary data. *Bioinformatics* **33**, 3088–3090 (2017).

51.   Blokzijl, F., Janssen, R., van Boxtel, R. & Cuppen, E. MutationalPatterns: Comprehensive

genome-wide analysis of mutational processes. *Genome Med.* (2018). doi:10.1186/s13073-

018-0539-0

52.   Gaujoux, R. & Seoighe, C. A flexible R package for nonnegative matrix factorization. *BMC*

Bioinformatics **11**, (2010).

53.   Cortés-ciriano, I., Lee, J., Xi, R., Jain, D. & Jung, Y. L. Comprehensive analysis of

chromothripsis in 2 , 658 human cancers using whole-genome sequencing. (2018).

54.   Hahsler, M., Hornik, K. & Buchta, C. Getting Things in Order⬚: An Introduction to the R

Package seriation. *J. Stat. Softw.* **25**, 1–27 (2008).

55. Venables, W. N. & Ripley, B. D. *Modern Applied Statistics with S.* (2002). doi:10.1007/978-0-

387-21706-2

56. Abhishek A. Singh, Karianne Schuurman, Ekaterina Nevedomskaya, Suzan Stelloo, Simon

Linder, Marjolein Droog, Yongsoo Kim, Joyce Sanders, Henk van der Poel, Andries M

Bergman, Lodewyk FA Wessels, W. Z. Optimized ChIP-seq method facilitates transcription

factor profiling in human tumors. *Life Sci. Alliance* **In Press**,

57. Zwart, W. *et al.* A carrier-assisted ChIP-seq method for estrogen receptor-chromatin

interactions from breast cancer core needle biopsy samples. *BMC Genomics* (2013).

doi:10.1186/1471-2164-14-232

58. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics **25**, 1754–1760 (2009).

59. Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, (2008).

60. Kumar, V. *et al.* Uniform, optimal signal processing of mapped deep-sequencing data. *Nat.*

Biotechnol. **31**, 615–622 (2013).

61. Stelloo, S. *et al.* Endogenous androgen receptor proteomic profiling reveals genomic

subcomplex involved in prostate tumorigenesis. *Oncogene* **37**, 313–322 (2018).

62. Kim, S. *et al.* Strelka2: Fast and accurate variant calling for clinical sequencing applications.

doi.org 192872 (2017). doi:10.1101/192872

63. Ha, G. *et al.* TITAN: inference of copy number architectures in clonal cell populations from

tumor whole-genome sequence data. *Genome Res.* (2014). doi:10.1101/gr.180281.114

64. Rausch, T. *et al.* DELLY: Structural variant discovery by integrated paired-end and split-read

analysis. *Bioinformatics* (2012). doi:10.1093/bioinformatics/bts378

Acknowledgements

This publication and the underlying study have been made possible partly on the basis of the data

that Hartwig Medical Foundation and the Center of Personalized Cancer Treatment (CPCT) have

made available to the study. We would like to thank the local principal investigators of all contributing

centers for their help with patient enrollment (listed in Supplementary table 1). We would also like to

thank Tesa M. Severson for her help with the computational analyses of the ChIP-seq data, Suzan

Stelloo for providing ChIP-seq results on cell lines and Arne van Hoeck for providing the CHORD

(HR-deficiency) prediction scores. We thank Dr. Joost van Rosmalen for his advises on the statistical

analyses. In addition, we would like to thank the Barcode for Life foundation for making this research

possible. Figure 1 was created with BioRender.com.

Author contributions

LFVD, JVR, MS, MPL and HJGVDW wrote the manuscript, which all authors critically reviewed. JVR

and HJGVDW performed the bioinformatics analyses and visualization. LFVD, MPL and NM managed

clinical data assessment. PW and WZ performed ChIP-seq experiments and data analyses. NM,

MPL, PH, AMB, MSVDH, IMVO, and RDW are clinical contributors. MPL, MSVDH, EV, NS, JWMM

and SS are part of the CPCT-02 study. EPJGC coordinated the sequencing of samples and

contributed to the bioinformatics analyses. TNY, JL and PCB assisted with acquisition of the primary

prostate cancer cohort and subsequent analysis.

Grants

This work was supported in parts by a KWF-Alpe d'HuZes project [NKI 2014-7080], a grant from

Astellas Pharma [Lolkema/NL-72-RG-11] and a Johnson & Johnson grant [212082PCR3014]. This

work was supported by the NIH/NCI under award number P30CA016042. HJGVDW, JVR and the

Erasmus MC Cancer Computational Biology Center (CCBC) were financed through a grant from the

Daniel den Hoed foundation.

Competing interests

All authors declare no competing interests.

Additional information

Supplementary information is available for this paper at https://doi.org/XXXXXXXXX.

 Figure legends

Figure 1 - Overview of study design and patient cohort (n = 197)

(a) Flowchart of patient inclusion. From the CPCT-02 cohort, patients with metastatic prostate

cancer were selected. Patients were excluded if data from metastatic samples were not

available and if clinical data indicated that patients had hormone-sensitive or neuro-endocrine

prostate cancer or unknown disease status at the time of analysis.

(b) Overview of the biopsy sites. Number of biopsies per metastatic site analyzed with WGS.

(c) Age of patients at biopsy. Bee-swarm boxplot with notch of the patient age distribution.

Boxplot depicts the upper and lower quartiles, with the median shown as a solid line; whiskers

indicate 1.5 times the interquartile range (IQR). Data points outside the IQR are shown.

Figure 2 - mCRPC shows several recurrent somatic alterations affecting several oncogenic

pathways

Based on dN/dS (q ≤ 0.1) and GISTIC2 focal peak (q ≤ 0.1) criteria, we show the genes and focal

genomic foci which are recurrently mutated, amplified or deleted in our mCRPC cohort of 197

patients. The upper track (top bar plot) displays tumor mutational burden (TMB) per SNV (blue), InDel

(yellow) and MNV (orange) category in coding regions (square root scale). Samples are sorted based

on mutual-exclusivity of the depicted genes and foci. The heatmap displays the type of mutation(s)

per sample; (light-)green or (light-)red backgrounds depict copy number aberrations whilst the inner

square depicts the type of (coding) mutation(s). Relative proportions of mutational categories (coding

mutations [SNV, InDels and MNV] (yellow), SV (blue), deep amplifications [high-level amplifications

resulting in many additional copies] (green) and deep deletions [high-level losses resulting in (near)

homozygous losses] (red)) per gene and foci are shown in the bar plot next to the heatmap. Narrow

GISTIC2 peaks covering ≤ 3 genes were reduced to gene-level rows if one of these genes is present

in the dN/dS (q ≤ 0.1) analysis or is a known oncogene or tumor-suppressor. For GISTIC2 peaks

covering multiple genes, only deep amplifications and deep deletions are shown. Recurrent aberrant

focal genomic foci in gene deserts are annotated with their nearest gene. Significance scores (-

1*log10(q)) of the dN/dS and GISTIC2 analysis are shown on the outer-right bar plots; bars in the

GISTIC2 significance plot are colored red if these foci were detected as a recurrent focal deletion and

green if detected as a recurrent focal gain. Per sample, the presence of (predicted) *ETS* fusions

(green), chromothripsis (light pink), kataegis (red), CHORD prediction score (HR-deficiency) (pink

gradient), MSI status (dark blue) and biopsy location are shown as bottom tracks.

Figure 3 - Comparison of the mutational landscape between primary prostate cancer and

mCRPC

(a) Tumor mutational burden (SNVs and InDels per Mbp) from a primary prostate cancer ($n =$

210) and the CPTC-02 mCRPC cohort ($n = 197$). Bee-swarm boxplot with notch of the tumor

mutational burden. Boxplot depicts the upper and lower quartiles, with the median shown as a

solid line; whiskers indicate 1.5 times the interquartile range (IQR). Data points outside the

IQR are shown. Statistical significance was tested with Wilcoxon rank-sum test and $p \leq 0.001$

is indicated as ***.

(b) Frequency of structural variant events from an unmatched cohort of primary prostate cancer

($n = 210$) and the CPTC-02 mCRPC cohort ($n = 197$). Boxplot depicts the upper and lower

quartiles, with the median shown as a solid line; whiskers indicate 1.5 times the interquartile

range (IQR). Data points outside the IQR are shown. Statistical significance was tested with

Wilcoxon rank-sum test and $p \leq 0.001$ is indicated as ***.

(c) Comparison of the mutational frequencies for driver genes detected by dN/dS and/or

GISTIC2, or subtype-specific genes, enriched in mCRPC relative to primary prostate cancer

or vice-versa. The difference in relative mutational frequency is shown on the x-axis and the

adjusted p-value (two-sided Fisher's Exact Test with BH correction) is shown on the y-axis.

Size of the dot is proportional to the absolute difference in mutational frequency between both

the cohorts. Symbols of genes with p-values below 0.05 are depicted in black and additional

genes-of-interests are highlighted in grey. The general genomic foci of the gene and absolute

number of samples with an aberration per cohort in primary prostate cancer and mCRPC,

respectively, is shown below the gene symbol. This analysis was performed on coding

mutations, gains and deletions per gene.

(d) Same as in (c) but using only coding mutations.

(e) Overview of the mutational categories (coding mutations [yellow], deletions [red] and

amplifications [green]) of the driver genes detected by dN/dS and/or GISTIC2, or subtype-

specific genes, enriched in mCRPC relative to primary prostate cancer ($q \leq 0.05$). For each

gene the frequency in primary prostate cancer is displayed followed by the frequency in

mCRPC.

Figure 4 - WGS reveals novel insight into the various (non-coding) aberrations affecting AR

regulation

(a) Mutational overview of top recurrently-mutated genes affecting *AR* regulation and their

putative enhancer foci (as detected by GISTIC2). The first track represents the number of

genomic mutations per Mbp (TMB) per SNV (blue), InDels (yellow) and MNV (orange)

category (square root scale). Samples are sorted based on mutual-exclusivity of the depicted

genes and foci. The heatmap displays the type of mutation(s) per sample, (light-)green or

(light-)red backgrounds depict copy number aberrations whilst the inner square depicts the

type of (coding) mutation(s). Relative proportions of mutational categories (coding mutations

[SNV, InDels and MNV] (yellow), SV (blue), deep amplifications (green) and deep deletions

(red)) per gene and foci are shown in the bar plot next to the heatmap. The presence of

chromothripsis (light pink), kataegis (red), CHORD prediction score (HR-deficiency) (pink

gradient), MSI status (dark blue) and biopsy location are shown as bottom tracks.

(b) Overview of the copy number deviations between putative enhancer and gene regions for *AR*

and *MYC*. Samples were categorized as enhancer- (blue) or gene- (red) enriched if enhancer-

to-gene ratio deviated > 1 studentized residual (residual in standard deviation units) from a

1:1 ratio.

(c) Copy number and ChIP-seq profiles surrounding the *AR* and *PCAT1/MYC* gene loci (with

1.25 additional Mbp up-/downstream). The upper panel displays the selected genomic

window and the overlapping genes. The first and second track display the aggregated mean

copy number (per 0.1Mbp window) of the enhancer- and gene-enriched samples,

respectively. These profiles identify distinct amplified regions (indicated by red asterisk) in

proximity to the respective gene bodies. The 3^{th} to 8^{th} tracks represent AR ChIP-seq profiles

(median read-coverage per 0.1Mbp windows) in two mCRPC patients (# 3 and 4), LNCaP (#

5) and LNCaP with R1881 treatment (# 6), VCaP (# 7) and bicalutamide-resistant VCaP (# 8).

The 9^{th} to 11^{th} tracks represent FOXA1 ChIP-seq profiles (median read-coverage per 0.1Mbp

windows) in two mCRPC patients (# 9 and 10) and LNCaP with R1881 treatment (# 11). The

12th to 14th tracks represent H3K27ac ChIP-seq profiles (median read-coverage per 0.1Mbp

windows) in two mCRPC patients (# 12 and 13) and LNCaP with R1881 treatment (# 14)

reflecting active enhancer regions.

ChIP-seq peaks (MACS/MACS2; $q < 0.01$) are shown as black lines per respective sample.

Figure 5 - Unsupervised clustering of mCRPC reveals distinct genomic phenotypes

(a) Dendrogram of unsupervised clustering with optimal leaf ordering. Top eight clusters are

highlighted and denoted based on order of appearance (left to right): A to H. Y-axis displays

clustering distance (Pearson correlation; ward.D).

(b) Number of genomic mutations per Mbp (TMB) of SNV (blue), InDels (yellow) and MNV

(orange) categories. All genome-wide somatic mutations were taken into consideration

(square root scale).

(c) Absolute number of unique structural variants per sample. Cumulative frequency of

inversions, tandem duplication, deletions, insertions and translocations.

(d) Relative frequency per structural variant category. Tandem Duplications and Deletions are

subdivided into > 100 kbp and < 100 kbp categories. This track shows if an enrichment for

particular category of (somatic) structural variant can be detected, which in turn, can be

indicative for a specific mutational aberration.

(e) Relative genome-wide ploidy status, ranging from 0 to ≥ 7 copies. This track shows the

relative percentage of the entire genome which is (partially) deleted (ploidy < 2 per diploid

genome) or amplified (ploidy > 2 per diploid genome).

(f) Relative contribution to mutational signatures (COSMIC) summarized per proposed etiology.

This track displays the proposed etiology of each SNV based on their mutational contexts.

(g) Relative frequency of different SNV mutational changes.

(h) HR-deficient prediction score as assessed by CHORD. The binary prediction score of

CHORD (ranging from 0 to 1) is shown, in which higher scores reflect more evidence for HR-

deficiency in a given sample.

(i) MSI status as determined using a stringent threshold of MSI characteristics[40].

(j) Presence of a fusion with a member of the *ETS* transcription factor family. Green color

indicates a possible fusion.

(k) Presence of chromothripsis. Pink color indicates presence of chromothripsis as estimated by

ShatterSeek.

(l) Presence of kataegis. Red color indicates presence of one or more regions showing kataegis.

(m) General biopsy location.

Figure 6 - Distinct genomic phenotypes in mCRPC are enriched by mutually-exclusive

aberrations in key pathways

(a) Cluster-specific enrichment of mutated genes (multiple colors), chromothripsis (light pink) and

structural variants (light blue) (Fisher's Exact Test with BH correction; $q \leq 0.05$). Percentages

to the left of the black line represent the relative mutational frequency in mCRPC samples

which are not present in the respective cluster, whilst the percentages to the right of the black

line represent the relative mutational frequency present in the samples from the tested

cluster.

(b) Genomic overview with biologically-relevant genes in the clusters A, B, D and F with

mutational enrichment of genes or large-scale events. The first track represents the number

of genomic mutations per Mbp (TMB) per SNV (blue), InDels (yellow) and MNV (orange)

category (square-root scale). The second track represents the absolute number of unique

structural variants (green) per sample. The third track represents the relative frequency per

structural variant category. Tandem Duplications and Deletions are subdivided into > 100 kbp

and < 100 kbp categories. The fourth track represents relative genome-wide ploidy status,

ranging from 0 to ≥ 7 copies. The fifth track represents the relative contribution to mutational

signatures (COSMIC) summarized per proposed etiology. The sixth track displays somatic

mutations in the relevant genes found in at least one cluster. The lower tracks represent

presence of *ETS* fusions (green), chromothripsis (pink), kataegis (red), CHORD prediction

scores (HR-deficiency) (pink gradient) and MSI status (blue) based on a threshold of MSI

characteristics.

Table 1 - Cluster characteristics

Overview of the distinctive characteristics for each cluster (A-H).

Table 1: Cluster characteristics

	Number of patients (n; % of cohort)	Tumor mutational burden (CDS)	SNV/InDel ratio	Number of structural variants	Main structural variant category or differentiating category	Ploidy status	Main mutational signature	Top 3 cluster-specific aberrations (% of cluster)	ETS-fusions (n)	Chromothripsis (n)	Kataegis (n)
Cluster A	13 (6.6)	36,88	0,99	149	None	1,92	MSI	MSH6 (69.2) JAK1 (69.2) CIC (58.3)	3	1	6
Cluster B	13 (6.6)	2,44	7,07	669	Tandem duplications (>100 kb)	2,39	N/A	CDK12 (84.6) FGF3 (69.2) FGF4 (69.2)	2	0	1
Cluster C	15 (7.6)	3,00	6,73	237	Tandem duplications (<100 kb)	3,19	N/A	None	7	1	2
Cluster D	22 (11.2)	4,39	7,28	323	Deletions (>100kb)	2,16	BRCA	BRCA2 (68.2)	7	5	5
Cluster E	55 (27.9)	2,12	7,13	178	None	3,24	N/A	None	25	8	13
Cluster F	20 (10.15)	2,51	6,15	399,5	None	3,35	N/A	Chromothripsis (80)	10	16	7
Cluster G	34 (17.3)	2,12	6,13	221,5	None	2,98	N/A	None	23	8	5
Cluster H	25 (12.7)	2,30	5,81	201	Insertions	1,97	N/A	None	16	7	3

All numbers are median of the cluster, unless otherwise indicated.

CDS, coding sequence

Methods

Patient cohort and study procedures

Patients with metastatic prostate cancer were recruited under the study protocol (NCT01855477) of

the Center for Personalized Cancer Treatment (CPCT). This consortium consists of 41 hospitals in

The Netherlands (Supplementary table 1). This CPCT-02 protocol was approved by the medical

ethical committee (METC) of the University Medical Center Utrecht and was conducted in accordance

with the Declaration of Helsinki. Patients were eligible for inclusion if the following criteria were met: 1)

age ≥ 18 years; 2) locally advanced or metastatic solid tumor; 3) indication for new line of systemic

treatment with registered anti-cancer agents; 4) safe biopsy according to the intervening physician.

For the current study, patients were included for biopsy between 03 May 2016 and 28 May 2018.

Data were excluded of patients with the following characteristics: 1) hormone-sensitive prostate

cancer; 2) neuro-endocrine prostate cancer (as assessed by routine diagnostics); 3) unknown disease

status; 4) prostate biopsy (Figure 1a). All patients provided written informed consent before any study

procedure. The study procedures consisted of the collection of matched peripheral blood samples for

reference DNA and image-guided percutaneous biopsy of a single metastatic lesion. Soft tissue

lesions were biopsied preferentially over bone lesions. The clinical data provided by CPCT have been

locked at 1st of July 2018.

Collection and sequencing of samples[40]

Blood samples were collected in CellSave preservative tubes (Menarini-Silicon Biosystems,

Huntington Valley, PA, USA) and shipped by room temperature to the central sequencing facility at

the Hartwig Medical Foundation. Tumor samples were fresh-frozen in liquid nitrogen directly after the

procedure and send to a central pathology tissue facility. Tumor cellularity was estimated by

assessing a hematoxylin-eosin (HE) stained 6 micron section. Subsequently, 25 sections of 20 micron

were collected for DNA isolation. DNA was isolated with an automated workflow (QiaSymphony)

using the DSP DNA Midi kit for blood and QiaSymphony DSP DNA Mini kit for tumor samples

according to the manufacturer's protocol (Qiagen). DNA concentration was measured by Qubit™

fluorometric quantitation (Invitrogen, Life Technologies, Carlsbad, CA, USA). DNA libraries for

Illumina sequencing were generated from 50-100 ng of genomic DNA using standard protocols

(Illumina, San Diego, CA, USA) and subsequently whole-genome sequenced in a HiSeq X Ten

system using the paired-end sequencing protocol (2x150bp). Whole-genome alignment (GRCh37),

somatic variants (SNV, InDel (max. 50 bp), MNV), structural variant and copy number calling and *in*

silico tumor cell percentage estimation were performed in a uniform manner as detailed by Priestley et

al. (2019)[40]. Mean read coverages of reference and tumor BAM were calculated using Picard Tools

(v1.141;   CollectWgsMetrics)   based   on   GRCh37[41].

Additional annotation of somatic variants and heuristic filtering

In addition, heuristic filtering removed somatic SNV, InDel and MNV variants based on the following

criteria: 1) minimal alternative reads observations ≤ 3; 2) gnomAD exome (ALL) allele frequency ≥

0.001 (corresponding to ~62 gnomAD individuals); and 3) gnomAD genome (ALL) ≥ 0.005 (~75

gnomAD individuals)[42]. gnomAD database v2.0.2 was used. Per gene overlapping a genomic variant,

the most deleterious mutation was used to annotate the overlapping gene. Structural variants, with

BAF ≥ 0.1, were further annotated by retrieving overlapping and nearest up- and downstream

annotations using custom R scripts based on GRCh37 canonical UCSC promoter and gene

annotations with respect to their respective up- or downstream orientation (if known)[43]. Only potential

fusions with only two different gene-partners were considered (e.g. *TMPRSS2-ERG*); structural

variants with both breakpoints falling within the same gene were simply annotated as structural

variant mutations. Fusion annotation from the COSMIC (v85), CGI and CIVIC databases were used to

assess known fusions[44–46]. The COSMIC (v85), OncoKB (July 12, 2018), CIVIC (July 26, 2018), CGI

(July 26, 2018) and the list from Martincorena et al. (dN/dS) were used to classify known oncogenic or

cancer-associated genes[26,44–46].

Ploidy and copy number analysis

Ploidy and copy number (CN) analysis was performed by a custom pipeline as detailed by Priestley et

al. (2019)[40]. Briefly, this pipeline combines B-allele frequency (BAF), read depth and structural

variants to estimate the purity and CN profile of a tumor sample. Recurrent focal and broad CN

alterations were identified by GISTIC2.0 (v2.0.23)[27]. GISTIC2.0 was run with the following

parameters: a) genegistic 1; b) gcm extreme; c) maxseg 4000; d) broad 1; e) brlen 0.98; f) conf 0.95;

g) rx 0; h) cap 3; i) saveseg 0; j) armpeel 1; k) smallmem 0; l) res 0.01; m) ta 0.1; n) td 0.1; o)

savedata 0; p) savegene 1; q) gvt 0.1. Categorization of shallow and deep CN aberration per gene

was based on thresholded GISTIC2 calls. Focal peaks detected by GISTIC2 were re-annotated,

based on overlapping genomic coordinates, using custom R scripts and UCSC gene annotations.

GISTIC2 peaks were annotated with all overlapping canonical UCSC genes within the wide peak

limits. If a GISTIC2 peak overlapped with ≤ 3 genes, the most-likely targeted gene was selected

based on oncogenic or tumor suppressor annotation in the COSMIC (v85), OncoKB (July 12, 2018),

CIVIC (July 26, 2018) and CGI (July 26, 2018) lists[26,44–46]. Peaks in gene deserts were annotated with

their nearest gene.

Estimation of tumor mutational burden

The mutation rate per megabase (Mbp) of genomic DNA was calculated as the total genome-wide

amount of SNV, MNV and InDels divided over the total amount of callable nucleotides (ACTG) in the

human reference genome (hg19) FASTA sequence file:

729
$$TMB_{genomic} = \frac{(SNV_g + MNV_g + InDels_g)}{\left(\frac{2858674662}{10^6}\right)} \tag{1}$$

730   The mutation rate per Mbp of coding mutations was calculated as the amount of coding SNV, MNV

731   and InDels divided over the summed lengths of distinct non-overlapping coding regions, as

732   determined on the subset of protein-coding and fully supported (TSL = 1) transcripts in GenCode v28

733   (hg19)[47]:

734
$$TMB_{coding} = \frac{(SNV_c + MNV_c + InDels_c)}{\left(\frac{28711682}{10^6}\right)} \tag{2}$$

MSI and HR-deficiency prediction

HR-deficiency/BRCAness was estimated using the CHORD classifier (Nguyen, van Hoeck and

Cuppen, manuscript in preparation). This classifier was based on the HRDetect[48] algorithm, however,

redesigned to improve its performance beyond primary BC. The binary prediction score (ranging from

zero to one) was used to indicate BRCAness level within sample. To elucidate the potential target

gene(s) in the HR-deficient samples (Figure 4), we used the list of BRCAness genes from Lord et al.

(2016)[35].

MSI status was determined based on the following criteria; if a sample contained more than 11,436

genomic InDels (max. 50 bp, with repeat-stretches of ≥ 4 bases, repeat length sequence between 2

and 4, or if these consists of a single repeat sequence which repeats ≥ 5 times), the sample was

designated as MSI.[40]

Detection of (onco-)genes under selective pressure

To detect (onco-)genes under tumor-evolutionary mutational selection, we employed a Poisson-based

dN/dS model (192 rate parameters; under the full trinucleotide model) by the R package dndscv

(v0.0.0.9)[26]. Briefly, this model tests the normalized ratio of non-synonymous (missense, nonsense

and splicing) over background (synonymous) mutations whilst correcting for sequence composition

and mutational signatures. A global q-value ≤ 0.1 (with and without taking InDels into consideration)

was used to identify statistically-significant (novel) driver genes.

Identification of hypermutated foci (kataegis)

Putative kataegis events were detected using a dynamic programming algorithm which determines a

globally optimal fit of a piecewise constant expression profile along genomic coordinates as described

by Huber et al. and implemented in the tilingarray R package (v1.56.0)[49]. Only SNVs were used in

detecting kataegis. Each chromosome was assessed separately and the maximum number of

segmental breakpoints was based on a maximum of five consecutive SNVs (max. 5000 segments per

chromosome). Fitting was performed on $\log_{10}$-transformed intermutational distances. Per segment, it

was assessed if the mean intermutational distance was ≤ 2000 bp and at least five SNVs were used

in the generation of the segment. A single sample with > 200 distinct observed events was set to zero

observed events as this sample was found to be hypermutated throughout the entire genome rather

than locally. Kataegis was visualized using the R package karyoploteR (v1.4.1).[50]

Mutational signatures analysis

Mutational signatures analysis was performed using the MutationalPatterns R package (v1.4.2)[51]. The

thirty consensus mutational signatures, as established by Alexandrov et. al, (matrix Sij; i = 96; number

of trinucleotide motifs; j = number of signatures) were downloaded from COSMIC (as visited on 23-05-

2018)[29]. Mutations (SNVs) were categorized according to their respective tri-nucleotide context (hg19)

into a mutational spectrum matrix Mij (i = 96; number of trinucleotide contexts; j = number of

samples) and subsequently, per sample a constrained linear combination of the thirty consensus

mutational signatures was constructed using non-negative least squares regression implemented

in the R package pracma (v1.9.3).

Between two and fifteen custom signatures were assessed using the NMF package (v0.21.0) with

1000 iterations[52]. By comparing the cophenetic correlation coefficient, residual sum of squares and

silhouette, we opted to generate five custom signatures. Custom signatures were correlated to

existing   (COSMIC)   signatures   using   cosine   similarity.

Detection of chromothripsis-like events

Rounded absolute copy number (excluded Y chromosome) and structural variants (BAF ≥ 0.1) were

used in the detection of chromothripsis-like events by the Shatterseek software (v0.4) using default

parameters[53]. As a precise standardized definition of chromothripsis has not yet been fully

established, and as per the author's instruction, we performed visual inspection of reported

chromothripsis-like events after dynamically adapting criteria thresholds (taking the recommended

thresholds into consideration). We opted to use the following criteria: a) Total number of

intrachromosomal structural variants involved in the event ≥ 25; b) max. number of oscillating CN

segments (2 states) ≥ 7 or max. number of oscillating CN segments (3 states) ≥ 14; c) Total size of

chromothripsis event ≥ 20 Mbp; d) Satisfying the test of equal distribution of SV types ($p > 0.05$); and

e) Satisfying the test of non-random SV distribution within the cluster region or chromosome ($p \leq$

0.05).

Unsupervised clustering of mCRPC WGS characteristics

Samples were clustered using the Euclidian distance of the Pearson correlation coefficient (1-*r*) and

Ward.D hierarchical clustering based on five basic whole genome characteristics; number of

mutations per genomic Mbp (SNV, InDel and MNV), mean genome-wide ploidy, number of structural

variants and the relative frequencies of structural variant categories (inversions, tandem duplications

(larger and smaller than 100 kbp), deletions (larger and smaller than 100 kbp), insertions and

interchromosomal translocations). Data was scaled but not centered (root mean square) prior to

calculating Pearson correlation coefficients. After clustering, optimal leaf ordering (OLO) was

performed using the seriation package (v1.2.3)[54]. The elbow method was employed to determine

optimal number of discriminating clusters (Supplementary figure 10) using the factoextra package

(v1.0.5). Bootstrapping was performed using the pvclust package (v2.0) with 5000 iterations.

Cluster-specific enrichment of aberrant genes (either through SV, deep copy number alteration or

coding SNV/InDel/MNV), kataegis, chromothripsis, GISTIC2 peaks and predicted fusions) between

clusters was tested using a two-sided Fisher's Exact Test and Benjamini-Hochberg correction.

A principal component analysis (with scaling and centering) using the prcomp R package[55] was

performed on the chosen genomic features and cos^2 values for each feature per principal component

were retrieved to determine the importance of each feature per respective principal component.

To test the robustness of our clustering, we performed unsupervised clustering, and also other

techniques, using various combinations of structural variants and clustering mechanisms as a

surrogate for different genome-instability metrics but this analysis did not reveal any striking new

clusters.

Supervised clustering based on mutually-exclusive aberrations

Samples were sorted on mutual-exclusivity of *SPOP*, *FOXA1* and *IDH1* coding mutations and copy

number aberrations and *ETS* family gene fusions (and overexpression) per promiscuous partner

(*ERG*, *ETV1*, *ETV4* and *FLI1*) as defined in primary prostate cancer.[13] Supplementary table S1A of

the article The Molecular Taxonomy of Primary Prostate Cancer[13] was used to determine the relative

frequency and mutational types of each of the respective primary prostate cancer within the TCGA

cohort. In addition, as the TCGA cohort did not denote high-level/deep amplifications, we did not

incorporate these either in this analysis.

Correlation of the detection rate of genomic aberrations versus tumor cell percentages

Absolute counts of SNV, InDels, MNV and SV were correlated to the *in silico* estimated tumor cell

percentage using Spearman's correlation coefficient.

Correlation of pre-treatment history with detected aberrations and WGS characteristics

Pre-treatment history of patients were summarized into 10 groups:

1. Only chemo-treatment (with radio-nucleotides).

2. Only chemo-treatment (without radio-nucleotides).

3. Only radio-nucleotides.

4. Only secondary anti-hormonal therapy (with radio-nucleotides).

5. Only secondary anti-hormonal therapy (without radio-nucleotides).

6. Secondary anti-hormonal therapy + one chemo-treatment (with radio-nucleotides)

7. Secondary anti-hormonal therapy + two chemo-treatment (with radio-nucleotides)

8. Secondary anti-hormonal therapy + one chemo-treatment (without radio-nucleotides)

9. Secondary anti-hormonal therapy + two chemo-treatment (without radio-nucleotides)

10. No additional treatment after androgen deprivation therapy.

Association with mutated genes, presence of chromothripsis, presence of kataegis, MSI-status and

genomic subtypes was tested with a two-sided Fisher's exact test with Benjamini-Hochberg

correction.

ChIP-seq experimental set-up and analysis

ChIP-seq cell culturing

VCaP cells were incubated in RPMI medium in additional with 10% fetal bovine serum (FBS).

Bicalutamide resistant VCaP cells (VCaP-Bic) were cultured in RPMI medium supplemented with 10%

dextran charcoal-stripped bovine serum (DCC) and 10-6M bicalutamide. VCaP cells were hormone

deprived in RPMI medium with 10% DCC for 3 days before the ChIP-seq experiment.

ChIP-seq and peak calling analysis

For both cell and tissue ChIPs, 5 µg of antibody and 50 µl of magnetic protein A or G beads (10008D

or 10009D, ThermoFisher Scientific) were used per IP. The following antibodies were used: Foxa1/2

(M-20, sc-6554 Santa Cruz Biotechnology), AR (N-20, sc-816 Santa Cruz Biotechnology), and

H3K27ac (39133, Active Motif). ChIP-seq was performed as described previously[56]. In brief, fresh

frozen tissue was cryosectioned into 30-µm-thick slices and stored at –80°C till processing. Samples

were fixed using 2 mM DSG (20593; Thermo Fisher Scientific) in solution A (50 mM Hepes-KOH, 100

mM NaCl, 1 mM EDTA, 0.5 mM EGTA) while rotating for 25 min at room temperature, followed by the

addition of 1% formaldehyde and another 20 min incubation at room temperature. The reaction was

quenched by adding a surplus of glycine. Subsequently, tissue sections were pelleted and washed

with cold PBS. Tissue was disrupted using a motorized pellet pestle (Sigma-Aldrich) to disrupt the

tissue in cold PBS and obtain a cell suspension, after which the nuclei were isolated and the

chromatin was sheared. During immunoprecipitation, human control RNA (4307281; Thermo Fisher

Scientific) and recombinant Histone 2B (M2505S; New England Biolabs) were added as carriers, as

described previously[57].

Immunoprecipitated DNA was processed for sequencing using standard protocols and sequenced on

an Illumina HiSeq 2500 with 65bp single end reads. Sequenced samples were aligned to the

reference human genome (Ensembl release 55: Homo sapiens GRCh 37.55) using Burrows-Wheeler

Aligner (BWA, v0.5.10)[58], reads with a mapping quality > 20 were used for further downstream

analysis.

For the tissues, peak calling was performed using MACS2[59] with option --nomodel. In addition, peaks

were called against matched input using DFilter[60] in the refine setting with a bandwidth of 50 and a

kernel size of 30. Only peaks that were shared between the two algorithms were considered.

For the cell lines, peaks were obtained with MACS (v1.4; $p \leq 10^{-7}$).

The AR and FOXA1 ChIP-seq data for LNCAP with/-out R1881 was obtained from GSE94682[61]. The

H3K27ac   ChIP-seq   data   for   LNCAP   was   obtained   from   GSE114737[56].

Determining enrichment of enhancer to gene ratios

Absolute copy numbers segments overlapping the gene loci and putative enhancer region (as

detected by GISTIC2; focal amplification peaks with a width < 5000 bp) were retrieved per sample. If

regions overlapped multiple distinct copy number segments, the maximum copy number value of the

overlapping segments was used to represent the region. Samples with gene-to-enhancer ratios

deviating > 1 studentized residual from equal 1:1 gene-to-enhancer ratios (linear model: $\log_2$(copy

number of enhancer) - $\log_2$(copy number of gene locus) ~ 0) were categorized as gene or enhancer

enriched. Based on the direction of the ratio, samples were either denoted as enhancer (if positive

ratio) or gene (if negative ratio) enriched.

Comparison of unmatched primary prostate cancer and mCRPC

Mutational frequencies of the drivers (dN/dS and or GISTIC2) and subtype-specific genes were

compared to a separate (unmatched) cohort of primary prostate cancer (n = 210) focusing on Gleason

score (GS) of 3 + 3, 3 + 4 or 4 + 3, as described by Fraser et al. (2017) and Espiritu et al. (2018)[15,25].

Briefly, whole-genome sequencing reads were mapped to the human reference genome (hg19) using

BWA[58] (v0.5.7) and downstream analysis was performed using Strelka[62] (v.1.0.12) for mutational

calling using a matched-normal design (SNVs and InDels), copy number alterations were estimated

with TITAN[63] (v1.11.0) and SNP array data as described in Espiritu et al. (2018)[25] with Delly[64] (v0.5.5

and v0.7.8) was used for detecting structural variants (translocations, inversions, tandem-duplications

and deletions). Large insertion calls and overall ploidy was not available for the primary prostate

cancer cohort.

Tumor mutational burden (TMB) was calculated by dividing the number of SNVs and InDels by the

total amount of callable bases in the human reference genome (hg19), identical to **equation 1**. MNV

calls were not available for the primary prostate cancer cohort.

Multiple aberrations per gene within a sample were summarized as a single mutational event, *e.g.* a

deletion and mutation in *PTEN* would only count for a single mutation sample in the sample. Only

non-synonymous mutations and gains / deletions overlapping with coding regions were used.

Statistically significant differences in mutational frequencies were calculated using a two-sided

Fisher's Exact test with Benjamini-Hochberg correction.

The primary prostate cancer dataset was clustered together with the mCRPC cohort using the

Euclidian distance of the Pearson correlation coefficient (1-r) and Ward.D hierarchical clustering

based on three basic whole genome characteristics which were available for all samples; number of

mutations per genomic Mbp (SNVs and InDels), number of structural variants and the relative

frequencies of structural variant categories (inversions, tandem duplications (larger and smaller than

100 kbp), deletions (larger and smaller than 100 kbp), and interchromosomal translocations).

Data availability

The data that support the findings of this study are available from Hartwig Medical Foundation, which were used under data request number DR-011 for the current study. Both WGS and clinical data is freely available for academic use from the Hartwig Medical Foundation through standardized procedures and request forms can be found at https://www.hartwigmedicalfoundation.nl[40]. The ChIP-seq profiles (aligned reads and MACS/MACS2 peaks) as analyzed and shown in this manuscript have been deposited on GEO under accession number: GSE138168.

Code availability

All tools and scripts used for processing of the WGS data are available at https://github.com/hartwigmedical/ and/or can be provided by authors upon request.

a

(CPCT-02 study)
Metastatic prostate cancer patients included for biopsy
n = 341

Non-metastatic biopsy site (i.e. prostate)
n = 3

No biopsy taken
n = 24

Fresh-frozen biopsy and blood control taken
n = 314

Successful biopsy (TC% ≥30%)
n = 218

Failed biopsy (TC% < 30%)
n = 96

No WGS due to insufficient tumor DNA quantity
n = 1

WGS biopsy (114x) and blood control (38x)
n = 217

non-mCRPC (Neuro-endocrine/HSPC/Unknown)
n = 20

Final inclusion of mCRPC (adenocarcinoma)
n = 197

b

Liver
(*n* = 29)

Lung
(*n* = 3)

Bone
(*n* = 70)

Soft tissue
(*n* = 14)

Lymph node
(*n* = 81)

c

Age at biopsy

mCRPC cohort(n = 197)