Draft Genome Assembly and Annotation of Red Raspberry *Rubus Idaeus*

Running title: The red raspberry genome sequencing

Haley Wight[1,2], Junhui Zhou[1], Muzi Li[1,2], Sridhar Hannenhalli[1,2], Stephen M. Mount[1,2] and Zhongchi Liu[1]*

1.Dept. of Cell Biology and Molecular Genetics, University of Maryland, College Park, MD 20742

2.Center for Bioinformatics and Computational Biology, University of Maryland, College Park, MD20742

Emails: haleywight18@gmail.com**, junhui55@umd.edu, limuzi92@terpmail.umd.edu, sridhar@umiacs.umd.edu, smount@umd.edu, zliu@umd.edu***

*Corresponding author
Zhongchi Liu
Dept. of Cell Biology and Molecular Genetics
University of Maryland, College Park, MD 20742
Tel: 301-405-1586
Fax: 301-314-9489
zliu@umd.edu

Abstract

The red raspberry, *Rubus idaeus*, is widely distributed in all temperate regions of Europe, Asia,

and North America and is a major commercial fruit valued for its taste, high antioxidant and

vitamin content. However, *Rubus* breeding is a long and slow process hampered by limited

genomic and molecular resources. Genomic resources such as a complete genome sequencing

and transcriptome will be of exceptional value to improve research and breeding of this high

value crop. Using a hybrid sequence assembly approach including data from both long and short

sequence reads, we present the first assembly of the *Rubus idaeus* genome (Joan J. variety). The

de novo assembled genome consists of 2,145 scaffolds with a genome completeness of 95.3%

and an N50 score of 638 KB. Leveraging a linkage map, we anchored 80.1% of the genome onto

seven chromosomes. Using over 1 billion paired-end RNAseq reads, we annotated 35,566

protein coding genes with a transcriptome completeness score of 97.2%. The *Rubus idaeus*

genome provides an important new resource for researchers and breeders.

Key words: red raspberry, genome assembly, annotation, genome comparison

Introduction

The red raspberry, *Rubus idaeus*, is widely distributed in all temperate regions of Europe, Asia, and North America and has been used as food and medicine since 4[th] century AD (Graham et al., 2004). Often dubbed "European red raspberry", *Rubus idaeus* is a globally commercialized specialty fruit crop with a large number of commercial varieties, high price, and increasing consumer demands. Owing to its health promoting value, unique flavor, and attractive appearance, *Rubus idaeus* sales have recently climbed by 8.4% with world production over 795, 000 tons (Darnell et al., 2006)(Barney et al., 2007; Food and Agriculture Organization of the United Nations Statistics Division (FAOSTAT)). In addition to its economic and health-promoting value, the red raspberry plants possess interesting and sometimes unique biological characteristics such as cold hardiness, aggregate fruits, perennial roots and biennial canes, either summer-bearing or ever-bearing flowering/fruiting, and large numbers of hybrids and cultivars. However, red raspberry breeding and research has fallen behind relative to other special fruit crops due to poor seed germination, an absence of reference genome, and limited transcriptome data (Graham and Woodhead, 2009; Hyun et al., 2014). With the recent publication of a high quality black raspberry (*Rubus occidentalis)* genome (VanBuren et al., 2018), this red raspberry genome allows comparative genomics, genetic breeding, and gene identification of this globally commercialized berry.

Rubus idaeus is a member of the economically important *Rosaceae* family that also includes rose, peach, apple, cherry, pear, almond, strawberry, and blackberry. Up to now, the genomes of several *Rosaceae* family members have been sequenced, including *Rubus occidentalis* (black raspberry) (VanBuren et al., 2016, 2018)*, Malus* x *domestica* (apple) (Daccord et al., 2017; Velasco et al., 2010), *Prunus persica* (peach) (Ahmad et al., 2011; Verde et al., 2013), *Pyrus bretschneideri* (Chinese pear) and *Pyrus communi* (Chagne et al., 2014) (Chagné et al., 2014), *Fragaria vesca* (woodland strawberry) (Edger et al., 2018; Shulaev et al., 2011), *Potentilla micranthia* (mock strawberry) (Buti et al., 2018), and *Rosa chinensis* (Chinese rose) (Hibrand Saint-Oyant et al., 2018; Raymond et al., 2018). Due to the small genome size, wide variety of fruit types (pomes, drupes, achenes, hips, follicles and capsules), and plant growth habits (ranging from herbaceous to cane, bush and tree forms), *Rosaceous* genomes offer one of the best systems for the comparative studies in genome evolution and development (Xiang et al.,

2017). The availability of whole-genome sequences of key diploid species such as *Rubus idaeus*

in this family will be crucial to these efforts.

Here, we report a draft genome assembly of red raspberry, *Rubus idaeus (*Joan J. variety)*,* using

long reads of single-molecular real-time (SMRT) Pacific Biosciences sequencing as well as high

coverage Illumina short reads. The resulting draft genome is 300 Mbp in size with a BUSCO-

calculated genome completeness score of 95.3% and contains 2,145 scaffolds with a N50 of 638

Kb. Using RNA-seq data from dissected fruit tissues at two developmental stages, we annotated

the genome yielding 35,566 protein coding genes with a BUSCO-calculated transcriptome

completeness score of 97.2%. We anchored the genome to two previously published high

density linkage maps of *Rubus idaeus* (Ward et al., 2013), facilitating future marker

development, breeding, and identification of genes controlling useful trait characteristics. Future

comparative analysis, bolstered by this reference sequence, will enable the study of the complex

evolution of morphological diversity in fleshy fruits of *Rosaceae*.

Materials and methods

Plant material and DNA sequencing

Joan J., a high-yielding, thornless, early primocane raspberry variety was chosen for genome

sequencing. The Joan J. variety of *Rubus idaeus* was obtained from Appalachian Fruit Research

Station of USDA ARS. Its genomic DNA was extracted from young leaves using the

NucleoSpin®⬚Plant⬚II⬚Midi kit (MACHEREY-NAGEL, Düren, Germany). DNA was

sequenced at the Genomics Resource Center of the University of Maryland School of Medicine's

Institute of Genome Sciences. Specifically, a long read (5-20kb) PacBio genomic library was

constructed using SMRTBell Template Prep Kit and sequenced on two SMRT cells on the

PacBio Sequel System, generating 1,305,619 sequence reads with an average length of 9,879 bp

(Supplementary Table 1). At the same time, a DNA-seq library was constructed using TruSeq

DNA Library Pre Kits (Illumina) and then sequenced on Illumina HiSeq4000 platform in a single

lane, yielding 249,081,860 reads of PE150 (Supplementary Table 1).

Analysis of the Illumina DNA-Sequencing Data

PCR adapter sequences were removed using cutadapt (Martin, 2013). Jellyfish (Marçais and

Kingsford, 2011) was then used to perform the k-mer distribution analysis with k=31

(Supplementary Figure 1).

Genome Assembly

The genome assembly pipeline is shown in Supplementary Figure 2. A mixture of Illumina short

reads and Pacbio long reads (Supplemental Table 1) were assembled into contigs using

MaSuRCA; an assembler which combines the efficiency of de Bruijn graph and Overlap-Layout-

Consensus approaches (Zimin et al., 2013). The specific settings used in the configuration file

other than default were PE= pe 180 20, JF_SIZE = 200000000 and SOAP_ASSEMBLY=0.

Subsequently, Redundans was used to remove heterozygous contigs using an all versus all BLAT

approach (Pryszcz and Gabaldón, 2016). The Redundans pipeline also performed scaffolding

using a mixture of Illumina short reads and Pacbio long reads (Pryszcz and Gabaldón, 2016).

The genomes of *Potentilla micranthia* (Buti et al., 2018)*, Rubus occidentalis* (VanBuren et al.,

2016), and *Fragaria vesca* (Edger et al., 2018) were leveraged to improve scaffolding using

MeDuSa (Bosi et al., 2015). Scaffolds with less than 10X coverage were removed and scaffolds

with more than 500 consecutive N's were split. Bowtie2 version 2.3.0 (Langmead and Salzberg,

2012) was used to map the Illumina reads back onto the genome prior to Pilon with maximum

fragment length to be 1000 and default settings otherwise. The mapping rate was 97.8% which

further validates assembly quality. Pilon (Walker et al., 2014) was then used for one iteration to

correct bases, fix misassembly and fill assembly gaps using the diploid parameter. Repeats were

then softmasked by first creating a custom repeat library with RepeatModeler -1.0.11

(http://www.repeatmasker.org/RepeatModeler/) using the NCBI engine option and then using

RepeatMasker (http://www.repeatmasker.org). Lastly, Haplomerger2 (Huang et al., 2017) split

the resulting assembly into two sub-assemblies to further remove hetereozygosity.

Sample collection and RNA-sequencing

Raspberry fruit from the Joan J. variety was dissected and separated into three tissues: ovary

wall, seed (or ovule), and receptacle. The fruit was collected at two developmental stages, 0 and

12 DPA. Three biological replicates for above 6 tissues were obtained (Supplementary Table 2).

Each tissue was homogenized in the presence of liquid nitrogen. Total RNA extraction was

performed following a previously published protocol (Jones et al., 1997) with few modifications.

The CTAB solution (3% CTAB, 100 mM Tris-HCl pH 8.0, 1.5 M NaCl, 20 mM EDTA,

5%PVP, and 1% β-mercaptoethanol made just before use) was added. 10 M Licl solution was

mixed with total RNA for two days to precipitate RNA. The total RNA samples were eluted in

DEPC-treated H_2O and stored in -80 ^{o}C.

Total RNA was shipped to the Weill Cornell's Genomics and Epigenemics Core Facility, where

polyA was isolated and RNA-seq libraries made using Tru-Seq RNA Library Prep Kit.

Subsequently, the RNA-seq libraries were sequenced on Illumina HiSeq4000, yielding a total of

1,057,377,357 reads; an average of 96.24% of these reads mapped to the genome

(Supplementary Table 2).

120
121
Genome annotation

Repeat Masker was used with a custom repeat library built with Repeat Modeler to soft-mask the

genome, and then a combination of *ab initio* and alignment guided assembly was employed to

annotate the soft-masked genome. The Illumina Reads of RNA-seq data described above were

trimmed with Trimmomatic (Bolger et al., 2014). RNA-Seq reads were mapped onto the draft

genome sequence using Bowtie2 (Langmead et al., 2009). The bam file obtained was used to

generate the training set for the gene prediction of BRAKER1 pipeline (Hoff et al., 2016).

Candidate transcripts containing no known protein domains by Interproscan5 (Jones et al., 2014)

were removed from the final set (13.96% percent decrease).

Trinity was then used to assemble the transcriptome on both genome guided and *de novo* settings

(Grabherr et al., 2011). Prior to trinity assembly, reads were normalized using the perl script

provided by Trinity and aligned using Bowtie2 (Grabherr et al., 2011; Langmead et al., 2009).

Trinity assemblies were amassed into a comprehensive transcriptome database using PASA

(Haas et al., 2003). Lastly, cd-hit-v4.6.8 (Li and Godzik, 2006) was used to cluster transcriptome

assemblies from the resulting PASA and BRAKER1 assemblies with over 95% identity into

unigenes. Unigenes that did not map to the genome, had no RNA-seq evidence, and had no

known protein domains or orthologues were removed.

Blast2GOPro version 5.1.1 was used to associate Gene Ontology (GO) terms to the resulting

transcripts (Supplementary Data 1). Protein sequences were searched against the non-redundant

(nr) database protein database from NCBI using BLASTP with an e-value cutoff of 1.0E-3

(Conesa et al., 2005). InterProScan was run using default databases in order to assign putative

domains to each transcript.

GO enrichment

GO enrichment tests were performed to understand potential function of *Rubus* specific genes.

GO term enrichment p-values were calculated using the Fisher's exact test in the TopGO R

package (http://bioconductor.org/packages/release/bioc/html/topGO.html). P-values were then

adjusted using R's FDR method.

Anchorage to linkage maps

BLAT was run with default settings to identify unique and complete matches to each marker

(Kent, 2002). After preparing the input files from BLAT (Supplementary Data 2),

pseudochromosomes were then constructed using ALLMAPS with default parameters (Tang et

al., 2015). Each genetic map was given a weight of 1. Chimeric scaffolds were manually broken

at positions with low coverage, correcting many misassemblies. The seven pseudochromosomes

were then constructed by integrating 98% of the markers from the genetic map.

Comparative genomics

Orthology was established using OrthoFinder-1.1.2 (Emms and Kelly, 2015) using default

parameters to infer a rooted species tree and identify orthologous gene groups. Subsequent to the

gene trees Orthofinder also produced the species tree. The resulting orthogroups and species tree

were then visualized with UpSetR (Conway et al., 2017) and an adjacent phylogenetic tree

visualized with iTOL (Letunic and Bork, 2016) (Figure 2A). A Circos plot (Krzywinski et al.,

2009) was created by creating links between every gene pair determined to be orthologs (Figure

2B-D). Syntenic orthologues were established by using MCScanX (Wang et al., 2012) with

settings -s 5. An all by all BLASTp (Boratyn et al., 2013) query with an e-value cutoff of 1e-10

was performed and used as a basis for MCScanX with default parameters to identify syntenic

gene regions.

Results and Discussion

Genome assembly and annotation

Rubus idaeus is a diploid species (2n=2x=14) with an estimated genome size of 293 Mbp based

on flow cytometry analysis (Graham and Woodhead, 2009). We first sequenced the *Rubus*

idaeus genome using 120X Illumina coverage (Supplementary Table 1). The distribution of k-

mers indicates that the *Rubus idaeus* genome is approximately 303 Mbp (Methods), and the

bimodal distribution of 31-mers (Supplemental Figure 1) suggests significant polymorphism and

heterozygosity in the genome.

To overcome the issue of heterozygosity for genome assembly, a hybrid genome assembly

approach was used taking advantage of both the sequencing depth and accuracy offered by the

Illumina platform (at 120X coverage) and the sequence length offered by the PacBio platform (at

26X overage) (Supplemental Table 1). The pipeline of the assembly is outlined in Supplemental

Figure 2. We used Redundans (Pryszcz and Gabaldón, 2016) and Haplomerger2 (Huang et al.,

2017) tools to correct for heterozygosity. A comparative genomic approach (Bosi et al., 2015;

Pop et al., 2004) was used as part of the genome assembly. Specifically, the most recently

assembled genomes of closely related species *Potentilla micranthia* (Buti et al., 2018)*, Rubus*

occidentalis (VanBuren et al., 2016), and *Fragaria vesca* (Edger et al., 2018) were leveraged to

improve scaffolding using MeDuSa (Bosi et al., 2015). The resulting *R. idaeus* genome assembly

is 300 Mbp in size, containing 2,145 scaffolds with a N50 of 638 Kb (Table 1). To assess the

completeness of the genome, BUSCO v.3.0.2 (Simão et al., 2015) was used to locate the

presence or absence of the embryophyta_odb9 (plant) dataset. The BUSCO Completeness Score

reached 95.3% (Table 1), which validates the good assembly quality.

To annotate the *Rubus idaeus* genome, a transcriptome was generated from 1,057,377,357

Illumina RNA-seq reads pooled from 18 RNA-seq libraries derived from three different fruit

tissues (ovary wall, ovule/seed, receptacle) at two developmental stages (0 and 12 Days Post-

Anthesis or DPA) in three biological replicates (Supplemental Table 2). A combination of *ab*

initio and alignment guided assembly was employed to annotate the genome (soft-masked for

repeats). This resulted in 35,566 protein coding genes with a BUSCO-calculated transcriptome

completeness score of 97.2% (Table 1). The high completeness score indicates that transcripts

from almost all genes expressed in these tissues have been sequenced. Finally, Blast2GO was

used to associate Gene Ontology (GO) terms to the annotated genes (Supplementary Data 1).

Anchoring scaffolds to genetic maps

The scaffolds were anchored onto pseudochromosomes (Figure 1) taking advantage of two

previous genetic linkage maps. They are respectively the 'Heritage' and 'Tulameen' variety-

based linkage maps that collectively contained 4225 markers. As a result, the

pseudochromosomes contain 80.1% of the assembly (ie. at 240 Mb). The average magnitude of

the Pearson correlation coefficient between the physical and map locations is 0.92 showing a

high consistency between the genome and previously published linkage maps (Figure 1;

Supplementary Data 2).

Comparative Genomics

Orthologous gene groups were established from 10 angiosperms using OrthoFinder-1.1.2 (Emms

and Kelly, 2015); these include 9 members of the *Rosaceae* family (*Prunus persica*, *Pyrus*

communis, *Malus* x *domestica*, *Rosa chinesis*, *Rosa multiflora*, *Rubus occidentalis*, *Rubus*

idaeus, *Fragaria vesca*, *Potentilla micrantha*) and the model organism *Arabidopsis thaliana*,

used here as an outlier species to root the tree. The resulting phylogenetic tree (Figure 2A) is

consistent with previously published phylogenetic analyses of the *Rosaceae* family (Xiang et al.,

2017). In total 25,193 orthogroups were established (Supplementary Data 3). As shown in Figure

2A, 10,205 orthogroups contained proteins from all 9 *Rosaceae* species as well as *Arabidopsis*.

Interestingly, many specific orthogroups (1,878) are unique to *Malus* x *domestica* and *Pyrus*

communis. Both species belong to the subfamily *Maleae*, which has undergone a whole genome

duplication, at its origin (Daccord et al., 2017; Wu et al., 2013; Xiang et al., 2017). The large

number of orthogroups shared between *Malus* x *domestica* and *Pyrus communis* suggests that

substantial diversification occurred after whole genome duplication (WGD) within the *Maleae*

subfamily, which may have contributed to the subfamily's pome fruit type (Velasco et al., 2010;

Xiang et al., 2017). Expectedly, all members of the *Rosaceae* family share many orthogroups

(1,420) that are distinct from *Arabidopsis thaliana*. Members of the same genus also show a high

number of common gene families. Specifically, there are 1,071 and 775 orthogroups limited to

the *Rosa* and *Rubus* genera, respectively (Figure 2A, Supplementary Data 3). As *Rubus* is one of

the largest and most morphologically diverse genus in the *Rosaceae* family (Alice and Campbell,

1999), we examined GO term enrichment among the 775 *Rubus*-specific orthogroups

(Supplementary Data 4). Significantly enriched GO terms include chromatin assembly, RNA-

splicing, and fungal-type cell wall organization, suggesting that *Rubus*-specific genes are

involved in gene regulation and defense.

242
Strawberry and raspberry share the same base chromosome number (n=7), with estimated

divergence time of 75 million years (Xiang et al., 2017). *Rubus occidentalis* and *Rubus idaeus*,

on the other hand, are closely related species. Syntenic blocks revealed a high collinearity

between *Rubus idaeus* and *Rubus occidentalis* and between *Rubus idaeus* and *F. vesca* (Figure

2B and C). *R. occidentalis* had 25,289 gene pairs represented within 1,596 collinear blocks with

R. idaeus. F. vesca and *R. idaeus* shared 17,769 syntenic gene pairs within 887 collinear blocks.

This high degree of synteny helps validate the *Rubus idaeus* assembly. When compared with the

more distant peach genome, *Prunus persica*, which has a different base chromosome number

(n=8), collinearity decreases slightly: *P. persica* and *R. idaeus* share 17,064 gene pairs on 877

collinear regions. Although there is lower collinearity, there are strikingly large conserved

syntenic blocks. For example, a large portion of *R. idaeus* chromosome 7 is syntenic to *P.*

persica chromosome 2 while a smaller portion of *R. idaeus* chromosome 7 syntenic to *P. persica*

7 (Figure 2D).

256
To facilitate future functional studies of raspberry development, the *Rubus idaeus* genome

assembly version 1 file, total transcript version 1 file, and annotation version 1 gff3 file are

provided as Supplementary Data 5, 6, and 7 respectively. The Transcription Factors (TFs) and

major hormonal pathway genes of *R. idaeus* are also identified and provided as Supplementary

Data 8. Together with the GO assignment (Supplementary Data 1), linkage between physical and

genetic markers (Supplementary Data 2), and ortholog assignment of nine *Rosaceae* species

(Supplementary Data 3), these new genomic resources will assist raspberry research and

breeding.

Supplemental Information

Supplementary Table 1. Summary statistics of DNA sequence data for *Rubus idaeus* genome

assembly

Supplementary Table 2. Summary statistics of RNA-seq data for *Rubus idaeus* fruit tissues.

Supplementary Figure 1. Bimodal K-mer distribution of *Rubus idaeus* (variety Joan J.) genome.

Supplemental Figure 2. Genome assembly pipeline.

Supplementary Data 1: GO annotations associated with *Rubus idaeus* transcripts

Supplementary Data 2: Correlation between scaffold positions and genetic markers

Supplementary Data 3: Orthology clustering of *Rosaceae* species and *Arabidopsis*

Supplementary Data 4: GO enrichment of *Rubus*-specific genes

Supplementary Data 5: *Rubus idaeus*_genome_v1.fa.gz

Supplementary Data 6: *Rubus idaeus*_transcript_v1.fa.gz

Supplementary Data 7: *Rubus idaeus*_annotation_v1.gff3

Supplementary Data 8: Orthologs of known *Arabidopsis* transcription factors and hormone

related genes

Funding

H.W. is supported by the NSF Computation and Mathematics for Biological Networks Research

Traineeship (NSF_NRT 1632976). The work is supported by NSF grants (IOS1444987) to S.H.

and Z.L.

Conflict of interest: None declared.

Availability of supporting data

The genomic DNA-sequencing and RNA-sequencing data supporting the results of this article

are available at Sequence Read Archive of NCBI with accession numbers SRP4284044 and

SRP153061 respectively.

Acknowledgements

We would like to thank Drs. Ann Callahan and Chris Dardick at USDA ARS for the Joan J.

Rubus idaeus plants and Miss Anuhyea Pulapaka for help with the genome assembly.

References

Ahmad, R., Parfitt, D.E., Fass, J., Ogundiwin, E., Dhingra, A., Gradziel, T.M., Lin, D., Joshi, N.A., Martinez-Garcia, P.J., and Crisosto, C.H. (2011). Whole genome sequencing of peach (Prunus persica L.) for SNP identification and selection. BMC Genomics *12*, 569.

Alice, L.A., and Campbell, C.S. (1999). Phylogeny of Rubus (rosaceae) based on nuclear ribosomal DNA internal transcribed spacer region sequences. Am. J. Bot. *86*, 81–97.

Barney, D.L., Bristow, P., Cogger, C., Fitzpatrick, S.M., Hart, J., Kaufman, D., Miles, C., Miller, T., Moore, P.P., Murray, T. and Rempel, H. (2007). Commercial red raspberry production in the Pacific Northwest. *Pacific Northwest Ext. Publ. PNW*, *598*.

Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics (Oxford, England) *30*, 2114–2120.

Boratyn, G.M., Camacho, C., Cooper, P.S., Coulouris, G., Fong, A., Ma, N., Madden, T.L., Matten, W.T., McGinnis, S.D., Merezhuk, Y., et al. (2013). BLAST: a more efficient report with usability improvements. Nucleic Acids Res *41*, W29–W33.

Bosi, E., Donati, B., Galardini, M., Brunetti, S., Sagot, M.-F., Lió, P., Crescenzi, P., Fani, R., and Fondi, M. (2015). MeDuSa: a multi-draft based scaffolder. Bioinformatics *31*, 2443–2451.

Buti, M., Moretto, M., Barghini, E., Mascagni, F., Natali, L., Brilli, M., Lomsadze, A., Sonego, P., Giongo, L., Alonge, M. and Velasco, R., (2018). The genome sequence and transcriptome of Potentilla micrantha and their comparison to Fragaria vesca (the woodland strawberry). *GigaScience*, *7*(4), pp.1-14.

Chagné, D., Crowhurst, R.N., Pindo, M., Thrimawithana, A., Deng, C., Ireland, H., Fiers, M., Dzierzon, H., Cestaro, A., Fontana, P., et al. (2014). The Draft Genome Sequence of European Pear (Pyrus communis L. "Bartlett"). PLoS ONE *9*, e92644–e92644.

Conesa, A., Gotz, S., Garcia-Gomez, J.M., Terol, J., Talon, M., and Robles, M. (2005). Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. Bioinformatics *21*, 3674–3676.

Conway, J.R., Lex, A., and Gehlenborg, N. (2017). UpSetR: an R package for the visualization of intersecting sets and their properties. Bioinformatics *33*, 2938–2940.

Daccord, N., Celton, J.-M., Linsmith, G., Becker, C., Choisne, N., Schijlen, E., van de Geest, H., Bianco, L., Micheletti, D., Velasco, R., et al. (2017). High-quality de novo assembly of the apple genome and methylome dynamics of early fruit development. Nature Genetics *49*, 1099–1106.

Darnell, R.L., Alvarado, H.E., Williamson, J.G., Brunner, B., Plaza, M., and Negrón, E. (2006). Annual, Off-season Raspberry Production in Warm Season Climates. HortTechnology *16*, 92–97.

Edger, P.P., VanBuren, R., Colle, M., Poorten, T.J., Wai, C.M., Niederhuth, C.E., Alger, E.I., Ou, S., Acharya, C.B., Wang, J., et al. (2018). Single-molecule sequencing and optical mapping

yields an improved genome of woodland strawberry (Fragaria vesca) with chromosome-scale contiguity. GigaScience *7*, 1–7.

Emms, D.M., and Kelly, S. (2015). OrthoFinder: solving fundamental biases in whole genome comparisons dramatically improves orthogroup inference accuracy. Genome Biology *16*, 157.

Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., et al. (2011). Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nature Biotechnology *29*, 644–652.

Graham J., Woodhead M. (2009) Raspberries and Blackberries: The Genomics of *Rubus*. In: Folta K.M., Gardiner S.E. (eds) Genetics and Genomics of Rosaceae. Plant Genetics and Genomics: Crops and Models, vol 6. Springer, New York, NY

Graham, J., Smith, K., MacKenzie, K., Jorgenson, L., Hackett, C., and Powell, W. (2004). The construction of a genetic linkage map of red raspberry (Rubus idaeus subsp. idaeus) based on AFLPs, genomic-SSR and EST-SSR markers. Theor Appl Genet *109*, 740–749.

Haas, B.J., Delcher, A.L., Mount, S.M., Wortman, J.R., Smith, R.K., Hannick, L.I., Maiti, R., Ronning, C.M., Rusch, D.B., Town, C.D., et al. (2003). Improving the Arabidopsis genome annotation using maximal transcript alignment assemblies. Nucleic Acids Research *31*, 5654–5666.

Hibrand Saint-Oyant, L., Ruttink, T., Hamama, L., Kirov, I., Lakhwani, D., Zhou, N.N., Bourke, P.M., Daccord, N., Leus, L., Schulz, D., et al. (2018). A high-quality genome sequence of Rosa chinensis to elucidate ornamental traits. Nature Plants *4*, 473–484.

Hoff, K.J., Lange, S., Lomsadze, A., Borodovsky, M., and Stanke, M. (2016). BRAKER1: Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS: Table 1. Bioinformatics *32*, 767–769.

Huang, S., Kang, M., and Xu, A. (2017). HaploMerger2: rebuilding both haploid sub-assemblies from high-heterozygosity diploid genome assembly. Bioinformatics *33*, 2577–2579.

Hyun, T.K., Lee, S., Rim, Y., Kumar, R., Han, X., Lee, S.Y., Lee, C.H., and Kim, J.-Y. (2014). De-novo RNA Sequencing and Metabolite Profiling to Identify Genes Involved in Anthocyanin Biosynthesis in Korean Black Raspberry (Rubus coreanus Miquel). PLoS One *9(2), e88292.*

Jones, C.S., Iannetta, P.P.M., Woodhead, M., Davies, H.V., McNicol, R.J., and Taylor, M.A. (1997). The isolation of RNA from raspberry (Rubus idaeus) fruit. Molecular Biotechnology *8*, 219–221.

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A., Nuka, G., et al. (2014). InterProScan 5: genome-scale protein function classification. Bioinformatics (Oxford, England) *30*, 1236–1240.

Kent, W.J. (2002). BLAT--the BLAST-like alignment tool. Genome Research *12*, 656–664.

Krzywinski, M., Schein, J., Birol, I., Connors, J., Gascoyne, R., Horsman, D., Jones, S.J., and Marra, M.A. (2009). Circos: an information aesthetic for comparative genomics. Genome Research *19*, 1639–1645.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nature Methods *9*, 357–359.

Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. Genome Biology *10*, R25.

Letunic, I., and Bork, P. (2016). Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. Nucleic Acids Res. *44*, W242-245.

Li, W., and Godzik, A. (2006). Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. Bioinformatics *22*, 1658–1659.

Marçais, G., and Kingsford, C. (2011). A fast, lock-free approach for efficient parallel counting of occurrences of k-mers. Bioinformatics (Oxford, England) *27*, 764–770.

Martin, M. (2013). Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal *17*, 10–12.

Pop, M., Kosack, D.S., and Salzberg, S.L. (2004). Hierarchical scaffolding with Bambus. Genome Research *14*, 149–159.

Pryszcz, L.P., and Gabaldón, T. (2016). Redundans: an assembly pipeline for highly heterozygous genomes. Nucleic Acids Research *44*, e113.

Raymond, O., Gouzy, J., Just, J., Badouin, H., Verdenaud, M., Lemainque, A., Vergne, P., Moja, S., Choisne, N., Pont, C., et al. (2018). The Rosa genome provides new insights into the domestication of modern roses. Nature Genetics *50*, 772–777.

Shulaev, V., Sargent, D.J., Crowhurst, R.N., Mockler, T.C., Folkerts, O., Delcher, A.L., Jaiswal, P., Mockaitis, K., Liston, A., Mane, S.P., et al. (2011). The genome of woodland strawberry (Fragaria vesca). Nature Genetics *43*, 109–116.

Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E. V., and Zdobnov, E.M. (2015). BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. Bioinformatics *31*, 3210–3212.

Tang, H., Zhang, X., Miao, C., Zhang, J., Ming, R., Schnable, J.C., Schnable, P.S., Lyons, E., and Lu, J. (2015). ALLMAPS: robust scaffold ordering based on multiple maps. Genome Biology *16*, 3.

VanBuren, R., Bryant, D., Bushakra, J.M., Vining, K.J., Edger, P.P., Rowley, E.R., Priest, H.D., Michael, T.P., Lyons, E., Filichkin, S.A., et al. (2016). The genome of black raspberry (*Rubus occidentalis*). The Plant Journal *87*, 535–547.

VanBuren, R., Wai, C.M., Colle, M., Wang, J., Sullivan, S., Bushakra, J.M., Liachko, I, Vining, K.J., Dossett, M., Finn, C.E., et al. (2018). A near complete, chromosome-scale assembly of the black raspberry (Rubus occidentalis) genome. Gigascience *7*.

Velasco, R., Zharkikh, A., Affourtit, J., Dhingra, A., Cestaro, A., Kalyanaraman, A., Fontana, P., Bhatnagar, S.K., Troggio, M., Pruss, D., et al. (2010). The genome of the domesticated apple (Malus × domestica Borkh.). Nature Genetics *42*, 833–839.

Verde, I., Abbott, A.G., Scalabrin, S., Jung, S., Shu, S., Marroni, F., Zhebentyayeva, T., Dettori, M.T., Grimwood, J., Cattonaro, F., et al. (2013). The high-quality draft genome of peach (Prunus persica) identifies unique patterns of genetic diversity, domestication and genome evolution. Nature Genetics *45*, 487–494.

Walker, B.J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C.A., Zeng, Q., Wortman, J., Young, S.K., et al. (2014). Pilon: An Integrated Tool for Comprehensive Microbial Variant Detection and Genome Assembly Improvement. PLoS ONE *9*, e112963.

Wang, Y., Tang, H., DeBarry, J.D., Tan, X., Li, J., Wang, X., Lee, T., Jin, H., Marler, B., Guo, H., et al. (2012). MCScanX: a toolkit for detection and evolutionary analysis of gene synteny and collinearity. Nucleic Acids Res *40*, e49.

Ward, J.A., Bhangoo, J., Fernández-Fernández, F., Moore, P., Swanson, J., Viola, R., Velasco, R., Bassil, N., Weber, C.A., and Sargent, D.J. (2013). Saturated linkage map construction in Rubus idaeus using genotyping by sequencing and genome-independent imputation. BMC Genomics *14*, 2.

Wu, J., Wang, Z., Shi, Z., Zhang, S., Ming, R., Zhu, S., Khan, M.A., Tao, S., Korban, S.S., Wang, H., et al. (2013). The genome of the pear (Pyrus bretschneideri Rehd.). Genome Research *23*, 396–408.

Xiang, Y., Huang, C.-H., Hu, Y., Wen, J., Li, S., Yi, T., Chen, H., Xiang, J., and Ma, H. (2017). Evolution of Rosaceae Fruit Types Based on Nuclear Phylogeny in the Context of Geological Times and Genome Duplication. Molecular Biology and Evolution *34*, 262–281.

Zimin, A. V., Marçais, G., Puiu, D., Roberts, M., Salzberg, S.L., and Yorke, J.A. (2013). The MaSuRCA genome assembler. Bioinformatics *29*, 2669–2677.

Table 1. Statistics of genome and transcriptome assemblies. Single (S), Duplicated (D), Fragmented (F) and Missing (M) single-copy orthologs are reported alongside the BUSCO completeness score.

Total length	300,259,977 bp
Scaffold N50	638,152 bp
Contig N50	250,294 bp
Smallest Scaffold	501 bp
Largest Scaffold	4,458,320 bp
N's	174,429 bp (.000582%)
Sequence GC's	37.9%
% Repeats	43.35%
Busco Completeness Score (Genome)	95.3% (S:86.1%, D:9.2%), F:1.5%, M:3.2%
Number of Annotated Protein Coding Genes	35,566
Busco Completeness Score (Transcriptome)	97.2% (S:92.9%,D:4.3%), F:1.1%, M:1.7%

Figure 1. The correlation between physical map and the linkage maps of seven chromosomes.
For each chromosome, the left figure illustrates the connections between physical positions on the assembled pseudomolecule and the two flanking linkage maps colored in orange and teal respectively. The orange coloring represents the tulmaneen linkage map whereas the teal represents the heritage linkage map (Ward et al., 2013). On the right is the scatter plot with dots representing the physical position on the chromosome (*x* axis) versus the map position (*y* axis). Rho (ρ) is the Pearson correlation coefficient (right panel). Each panel represents distinct chromosome.

Figure 2. The distribution of shared gene families among nine *Rosaceae* species and *Arabidopsis thaliana.*

(A)The left panel describes the phylogeny among the species. The branch length distances represent substitutions per site. The right panel is an UpSet plot (Conway et al., 2017): an alternative representation of a venn diagram with intersections (shared genes) greater than 100. The species described in each intersection is represented by the dotted lines, the size of the intersection is described by the bar chart above. (B) Circos plots (Krzywinski et al., 2009) displaying macrosynteny between the genomes of *Rubus idaeus* and *Rubus occidentalis*. (C) Macrosynteny between *Rubus idaeus* and *Fragaria vesca*. (D) Macrosynteny between *Rubus idaeus* and *Prunus persica*. For B to D, each connecting line represents an orthologous gene pair and the right half of each circle consists of the seven *Rubus ideaus* chromosomes colored by the spectral order in the rainbow.

Supplemental Tables

Supplementary Table 1. Summary statistics of DNA sequence data for *Rubus idaeus* genome assembly

	Mean Read length	Read count	Total base, bp
Illumina PE	150	249,081,860	37,455,877,274
PacBio	9,879	1,305,619	8,007,129,543

Supplementary Table 2. Summary statistics of RNA-seq data for *Rubus idaeus* fruit tissues.

Sample	Number of Reads	% of Uniquely Mapped Reads	% of reads mapped to multiple loci	Total % reads mapped
Ovule-0-26	60144925	87.94%	7.13%	95.07%
Ovule-0-41	57979785	89.56%	7.24%	96.80%
Ovule-0-7	66004490	89.29%	7.30%	96.59%
Receptacle-0-17	64293938	89.79%	7.16%	96.95%
Receptacle-0-27	61401941	89.54%	7.18%	96.72%
Receptacle-0-41	68480278	88.20%	7.18%	95.38%
Receptacle-12-13	67769659	89.39%	6.32%	95.71%
Receptacle-12-1	54088666	91.41%	6.22%	97.63%
Receptacle-12-4	50260693	90.69%	6.70%	97.39%
Seed-12-13	53332584	84.84%	8.07%	92.91%
Seed-12-1	55781661	89.18%	8.47%	97.65%
Seed-12-7	62967294	89.73%	7.99%	97.72%
Wall-0-24	65304690	89.32%	7.27%	96.59%
Wall-0-7	59200984	89.64%	7.51%	97.15%
Wall-0-13	71863354	89.68%	7.34%	97.02%
Wall-12-13	68284797	83.95%	8.43%	92.38%
Wall-12-1	70217618	88.17%	8.27%	96.44%
Wall-12-4	61592260	89.07%	9.06%	98.13%

*Sample names are "Tissue-Stage-unique ID of the specific sample". The two stages are 0 DPA and 12 DPA. The tissues are Ovule, Seed, Receptacle, and Wall (ovary wall).

Supplemental Figures

Supplementary Figure 1. Bimodal K-mer distribution of *Rubus idaeus* (variety Joan J.) genome 31-mer distribution of *Rubus idaeus* genome obtained, using jellyfish, from 150-bp paired-end whole genome sequencing data.

Supplemental Figure 2. **Genome assembly pipeline.**
Flowchart represents all steps of the genome assembly process upstream of anchoring to the
linkage map.