Environmental DNA (eDNA) metabarcoding of pond water as a tool to

survey conservation and management priority mammals

Lynsey R. Harper[1*], Lori Lawson Handley[1], Angus I. Carpenter[2], Gill Murray-Dickson[3], Cristina

Di Muri[1], Callum J. Macgregor[4], Thomas W. Logan[1], Alan Law[5], Thomas Breithaupt[1], Daniel S.

Read[6], Allan D. McDevitt[7], and Bernd Hänfling[1]

[1] Department of Biological and Marine Sciences, University of Hull, Hull, HU6 7RX, UK

[2] Wildwood Trust, Canterbury Rd, Herne Common, Herne Bay, CT6 7LQ, UK

[3] The Royal Zoological Society of Scotland, Edinburgh Zoo, 134 Corstorphine Road, Edinburgh, EH12 6TS, UK

[4] Department of Biology, University of York, Wentworth Way, York, YO10 5DD, UK

[5] Biological and Environmental Sciences, University of Stirling, Stirling, FK9 4LA, UK

[6] Centre for Ecology & Hydrology (CEH), Benson Lane, Crowmarsh Gifford, Wallingford, Oxfordshire, OX10 8BB, UK

[7] Ecosystems and Environment Research Centre, School of Environment and Life Sciences, University of Salford,

Salford, M5 4WT, UK

*Corresponding author: lynsey.harper2@gmail.com

Lynsey Harper, Department of Biological and Marine Sciences, University of Hull, Hull, HU6 7RX,

UK

Abstract

1. Environmental DNA (eDNA) metabarcoding is largely used to survey aquatic communities, but can also provide data on terrestrial taxa utilising aquatic habitats. However, the entry, dispersal, and detection of terrestrial species' DNA within waterbodies is understudied.

2. We evaluated eDNA metabarcoding of pond water for monitoring semi-aquatic, ground-dwelling, and arboreal mammals, and examined spatiotemporal variation in mammal eDNA signals using experiments in captive and wild conditions.

3. We selected nine focal species of conservation and management concern: European water vole, European otter, Eurasian beaver, European hedgehog, European badger, red deer, Eurasian lynx, red squirrel, and European pine marten. We hypothesised that eDNA signals (i.e. proportional read counts) would be stronger for semi-aquatic than terrestrial species, and at sites where mammals exhibited behaviours (e.g. swimming, urination). We tested this by sampling waterbodies in enclosures of captive focal species at specific sites where behaviours had been observed ('directed' sampling) and at equidistant intervals along the shoreline ('stratified' sampling). We then surveyed natural ponds ($N = 6$) where focal species were present using stratified water sampling, camera traps, and field signs. eDNA samples were metabarcoded using vertebrate-specific primers.

4. All focal species were detected in captivity. eDNA signal strength did not differ between directed and stratified samples across or within species, between species lifestyles (i.e. semi-aquatic, ground-dwelling, arboreal), or according to behaviours. Therefore, eDNA was evenly distributed within artificial waterbodies. Conversely, eDNA was unevenly distributed

in natural ponds. eDNA metabarcoding, camera trapping, and field signs detected beaver,

red deer, and roe deer. Badger and red fox were recorded with cameras and field signs, but

not eDNA metabarcoding. However, eDNA metabarcoding detected small mammals missed

by cameras and field signs, e.g. water vole. Terrestrial mammal eDNA signals were weaker

and detected in fewer samples than semi-aquatic mammal eDNA signals.

5. eDNA metabarcoding has potential for inclusion in mammal monitoring schemes by

enabling large-scale, multi-species distribution assessment for priority and difficult to survey

species, and could provide early indication of range expansions or contractions. However,

eDNA surveys need high spatiotemporal resolution and metabarcoding biases require

further investigation before this tool is routinely implemented.

Key-words: camera traps, environmental DNA (eDNA), field signs, lentic, monitoring, semi-

aquatic mammals, terrestrial mammals

1. Introduction

Mammals are among the most threatened vertebrate groups globally, with at least 25% of species at risk of extinction due to harvesting (i.e. food, medicine, materials), habitat degradation/loss (e.g. agriculture), non-native species (i.e. competition, hybridisation, disease transmission) or perception as pests (Visconti et al., 2011). The paucity of data for mammals prevents robust estimation of range expansions or declines and population trends (Bland, Collen, Orme, & Bielby, 2015). Most species lack long-term, systematic monitoring, with survey efforts biased towards rare species while widespread species receive less attention (Massimino, Harris, & Gillings, 2018). Consequently, there is a need for more effective and evidence-based strategies for mammal conservation and management (Battersby & Tracking Mammals Partnership, 2005; Mathews et al., 2018).

Many mammals are nocturnal and elusive thus monitoring often requires non-invasive, observational monitoring tools, such as camera traps and field signs (Caravaggi et al., 2018; Harris & Yalden, 2004; McShea, Forrester, Costello, He, & Kays, 2016; Sadlier, Webbon, Baker, & Harris, 2004). Camera trapping is cost-efficient, standardised, reproducible, and produces data suited to site occupancy modelling. However, camera traps can only survey a fraction of large, heterogeneous landscapes, trap placement can substantially influence species detection probabilities and community insights, and traps often miss small species (Burton et al., 2015; Caravaggi et al., 2018; Glen, Cockburn, Nichols, Ekanayake, & Warburton, 2013; Ishige et al., 2017; McShea et al., 2016; Stat et al., 2019). Field sign surveys are inexpensive, but resource-intensive (i.e. time, volunteers) for broad geographic coverage (Sadlier et al., 2004). Some

species also have similar footprints and scat, increasing the potential for misidentification

(Franklin et al., 2019; Harris & Yalden, 2004). Mammal survey methods are usually species-

specific, and multiple methods are necessary for large-scale, multi-species monitoring schemes

(Massimino et al., 2018).

Environmental DNA (eDNA) analysis is a recognised tool for rapid, non-invasive, cost-

efficient biodiversity assessment (Deiner et al., 2017). Organisms transfer their genetic material

to their environment via secretions, excretions, gametes, blood, or decomposition, which can

be isolated from environmental samples (Thomsen & Willerslev, 2015). Targeted (i.e. single-

species) eDNA analysis, using PCR, quantitative PCR (qPCR), or droplet digital PCR of eDNA

samples, is increasingly applied to survey marine (Baker, Steel, Nieukirk, & Klinck, 2018; Hunter

et al., 2018; Parsons, Everett, Dahlheim, & Park, 2018), freshwater (Lugg, Griffiths, van Rooyen,

Weeks, & Tingley, 2017; Padgett-Stewart et al., 2016; Thomsen et al., 2012), and terrestrial

(Aylward, Johnson, Sullivan, Perry, & Louis, 2018; Franklin et al., 2019; Rodgers & Mock, 2015;

Williams, Huyvaert, Vercauteren, Davis, & Piaggio, 2018) mammals. Alternatively, eDNA

metabarcoding can screen entire communities using PCR combined with high-throughput

sequencing (Deiner et al., 2017; Thomsen & Willerslev, 2015), but eDNA metabarcoding

assessments of mammal communities are less common (Klymus, Richter, Thompson, & Hinck,

2017; Ushio et al., 2017). Mammal assemblages have been obtained by metabarcoding

invertebrate blood meals (e.g. Tessler et al., 2018) and salt licks (Ishige et al., 2017) in tropical

habitats, but samples from the physical environment have tremendous potential to reveal

mammal biodiversity over broad spatiotemporal scales (Ushio et al., 2017).

In aquatic ecosystems, eDNA metabarcoding has predominantly been applied to

characterise fish (e.g. Evans et al., 2017; Hänfling et al., 2016; Lawson Handley et al., 2018;

Valentini et al., 2016) and amphibian (e.g. Bálint et al., 2018; Valentini et al., 2016)

communities. However, mammals also leave eDNA signatures in water that metabarcoding can

detect (Harper et al., 2018a; Klymus et al., 2017; Port et al., 2016; Ushio et al., 2017). Ponds in

particular act as stepping stones for semi-aquatic and terrestrial mammals by providing

drinking, foraging, dispersive, and reproductive opportunities (Klymus et al., 2017). Therefore,

samples from these waterbodies could provide information on biodiversity present in the wider

environment (Deiner et al., 2017; Harper et al., 2018a).

Despite evidence for eDNA deposition by semi-aquatic and terrestrial mammals in

freshwater systems (Harper et al., 2018a; Klymus et al., 2017; Lugg et al., 2017; Padgett-Stewart

et al., 2016; Rodgers & Mock, 2015; Thomsen et al., 2012; Ushio et al., 2017; Williams et al.,

2018), little is known about the influence of mammal behaviour on the distribution and

strength of the eDNA signal left behind. Drinking is a major source of eDNA deposition in water

due to the release of saliva, but mammals may also swim, wallow, urinate or defecate in water

(Rodgers & Mock, 2015; Ushio et al., 2017; Williams et al., 2018). Furthermore, arboreal

mammals may be expected to use ponds less than semi-aquatic and ground-dwelling species,

non-territorial mammals may visit ponds less than territorial species, and species living in

groups may leave more DNA in water than solitary species (Williams et al., 2018).

In this study, we conducted two experiments to evaluate eDNA metabarcoding of pond

water as a tool for monitoring semi-aquatic, ground-dwelling, and arboreal mammals of

conservation or management concern. The first experiment, carried out on nine focal species

housed at two wildlife parks, was designed to examine the role of sampling strategy, mammal

lifestyle, and mammal behaviour on eDNA detection and signal strength (i.e. read counts). Our

hypotheses for Experiment 1 were as follows: (1) eDNA would be unevenly distributed thus

directed sampling would yield stronger eDNA signals (i.e. higher read counts) for mammals than

stratified sampling; (2) semi-aquatic mammals would have stronger eDNA signals than ground-

dwelling or arboreal mammals; and (3) mammal behaviours involving water contact (e.g.

swimming, drinking) would be associated with stronger eDNA signals. The second experiment

aimed to validate eDNA metabarcoding against camera trapping and field sign searches at

natural ponds within sites where focal species were present. For Experiment 2, our hypotheses

were: (1) eDNA metabarcoding would detect more mammals than camera trapping or field

signs; (2) semi-aquatic mammals would be readily detected and their eDNA evenly distributed

in ponds in comparison to terrestrial mammals; and (3) temporal sampling would reveal that

terrestrial mammal eDNA is detectable for short periods in comparison to fully aquatic

vertebrates.

2. Materials and methods

2.1 Study species

We studied nine mammal species that are the focus of UK conservation or management

(Battersby & Tracking Mammals Partnership, 2005; Mathews et al., 2018): European water vole

(*Arvicola amphibius*), European otter (*Lutra lutra*), Eurasian beaver (*Castor fiber*), European

hedgehog (*Erinaceus europaeus*), European badger (*Meles meles*), red deer (*Cervus elaphus*),

Eurasian lynx (*Lynx lynx*), red squirrel (*Sciurus vulgaris*), and European pine marten (*Martes*

martes). Water vole, otter, red squirrel, pine marten and hedgehog are UK Biodiversity Action

Plan species (Joint Nature Conservation Committee, 2018). Lynx is not currently present in the

UK, but reintroduction trials have been proposed (Lynx UK Trust, 2018). Water vole, otter, and

beaver are semi-aquatic, red squirrel and pine marten are arboreal, and the other species are

ground-dwelling. Badger and red deer live in groups whereas the other species are solitary,

except when courting, mating, and rearing young.

2.2 Experiment 1: eDNA detection and signal strength in artificial systems

We performed eDNA metabarcoding for focal species under artificial conditions at two wildlife

parks to investigate whether mammal eDNA signal strength (i.e. read counts) depends on

sampling strategy, species lifestyle (semi-aquatic, ground-dwelling, arboreal), and behaviour.

Behavioural observation and eDNA sampling were conducted between 18[th] - 21[st] September

2017 at Wildwood Trust (WT), Kent, England, and 10[th] - 11[th] October 2017 at Royal Zoological

Society of Scotland (RZSS) Highland Wildlife Park (HWP), Kingussie, Scotland. Sixteen categories

of behaviour were defined based on potential contact with waterbodies and species lifestyle,

and the frequency and duration of behaviours recorded (Table S1). The number of individuals in

each enclosure was recorded alongside waterbody size (Table 1). Beaver, lynx, red deer, and

red squirrel were present at both wildlife parks, whereas other species were only present at

WT. Each species was observed for one hour on two separate occasions with exceptions.

Badger and beaver were nocturnal and required overnight observation using camera traps (Bushnell Trophy Cam Aggressor). Camera traps (one per enclosure) were positioned perpendicular to the ground at a height of 1 m and distance of 2 m from the pond shoreline to capture water and shoreline in the field of view. Cameras were set to take 30 s videos (1920 x 1080) when triggered (30 s interval between triggers) at high sensitivity. Behavioural observation was not undertaken for WT water voles as animals were under quarantine or HWP red squirrels as individuals were wild and widely distributed.

Water samples were collected from enclosures within 3 hrs of the second behavioural observation period. Up to six directed or stratified samples were collected, but sample number varied by species due to waterbody size as well as type and frequency of behaviours observed (Tables S1, S2). Enclosure drinking containers were also sampled and classed as 'other' samples. There were no artificial waterbodies in the WT water vole, red squirrel, or hedgehog enclosures, thus bathing and drinking bowls were sampled. The HWP beaver enclosure was empty for 24 hrs before sampling as animals were moved to RZSS Edinburgh Zoo (EZ), Scotland, for quarantine before wild release. Water from the empty enclosure and the EZ beaver quarantine enclosure was sampled. A sample was also collected from a water bath situated in the HWP woods to capture any wild red squirrels present and classed as 'other'.

Directed samples (2 L surface water taken approximately where behaviours were observed) were collected before stratified samples (2 L surface water [8 x 250 ml pooled subsamples] taken at equidistant points around the waterbody perimeter where access permitted) to minimise disturbance to the water column and cross-contamination risk. All samples were collected using sterile Gosselin™ HDPE plastic bottles (Fisher Scientific UK Ltd,

UK) and disposable gloves. A field blank (1 L molecular grade water [MGW]) was taken into

each species enclosure and opened. Samples (n = 80) collected from WT and HWP were

transported alongside field blanks (n = 13) in sterile coolboxes with ice packs to the University

of Kent (UoK) and EZ respectively, where ice was added to coolboxes.

Samples and blanks were vacuum-filtered within 6 hrs of collection in a UoK wet

laboratory that housed exotic amphibians (Table S3) and within 24 hrs of collection in a EZ staff

room. All surfaces and equipment were sterilised before, during, and after set-up in temporary

work areas. Surfaces and vacuum pumps were wiped with 10% v/v chlorine-based commercial

bleach (Elliott Hygiene Ltd, UK) solution. Non-electrical equipment was immersed in 10% bleach

solution for 10 minutes, followed by 5% v/v MicroSol detergent (Anachem, UK), and rinsed with

purified water. Up to 500 ml of each 2 L sample was vacuum-filtered through sterile 0.45 µm

mixed cellulose ester membrane filters with pads (47 mm diameter; Whatman, GE Healthcare,

UK) using Nalgene™ filtration units. One hour was allowed for each sample to filter and a

second filter used if clogging occurred. A filtration blank (1 L MGW) was processed during each

filtration round (n = 12), and equipment sterilised after each filtration round. After 500 ml had

filtered or one hour had passed, filters were removed from pads using sterile tweezers, placed

in sterile 47 mm petri dishes (Fisher Scientific UK Ltd, UK), sealed with parafilm (Sigma-Aldrich

Company Ltd, UK), and stored at -20 °C. The total water volume filtered per sample was

recorded for downstream analysis (Table S2; Fig. S1).

2.3 Experiment 2: eDNA detection and signal strength in natural systems

We performed eDNA metabarcoding for focal species at natural ponds in conjunction with conventional surveys to validate this molecular approach for mammal identification and to investigate spatiotemporal variation in mammal eDNA signals. At three sites where focal species were present based on cumulative survey data from landowners, reserve managers, researchers, and volunteers, we selected two ponds (range 293-5056 m^2, average 1471 m^2) within <4 km of each other. The Bamff Estate (BE), Alyth, Scotland, was selected for beaver, otter, badger, red deer, and red squirrel, but roe deer (*Capreolus capreolus*) and red fox (*Vulpes vulpes*) were also present. Otter, water vole, and badger were also present at Tophill Low Nature Reserve (TLNR), Driffield, East Yorkshire, where American mink (*Neovison vison*), stoat (*Mustela erminea*), weasel (*Mustela nivalis*), rabbit (*Oryctolagus cuniculus*), brown hare (*Lepus europaeus*), fox, roe deer, and grey squirrel (*Sciurus carolinensis*) additionally occur. We selected Thorne Moors (TM), Doncaster, South Yorkshire, for red deer and badger, but stoat, weasel, red fox, roe deer, and Reeve's muntjac (*Muntiacus reevesi*) were also present. Camera traps (Bushnell Trophy Cam Aggressor) were deployed at TM (one per pond) and BE (three per pond) one week prior to eDNA sampling and collected once sampling was completed. At TLNR, camera traps (two at one pond and three at the other pond) were deployed one day before a 5-day period of eDNA sampling and collected one week after sampling was completed. Camera traps were positioned perpendicular to the ground at a height of 1 m and distance of 0.3-1 m from the pond shoreline to capture water and shoreline in the field of view. Cameras were set to take three photographs (5 megapixel) when triggered (3 s interval between triggers) at high

sensitivity.

Ten stratified samples (2 L surface water [8 x 250 ml pooled subsamples] taken at

equidistant points around the pond perimeter where access permitted) were collected from the

shoreline of each pond and a field blank (1 L MGW) included using the same procedures as

Experiment 1. TM ponds were sampled on 17th April 2018, and BE ponds on 20th April 2018.

TLNR ponds were sampled every 24 hrs between 23rd - 27th April 2018 to investigate

spatiotemporal variation in mammal eDNA signals. TM and TLNR samples were transported on

ice in sterile coolboxes with ice packs to the University of Hull (UoH) eDNA facility, which is

devoted to pre-PCR processes with separate rooms for filtration, DNA extraction and PCR

preparation of environmental samples, and stored at 4 °C. BE samples were transported in

sterile coolboxes with ice packs to BE accommodation. Surfaces and equipment were sterilised

before, during, and after set-up in local accommodation as in Experiment 1. Samples (n = 140)

and field blanks (n = 14) were vacuum-filtered within 4 hrs of collection as outlined in

Experiment 1 with minor modifications to maximise detection probability as follows. The full 2 L

of each sample was vacuum-filtered where possible, two filters were used for each sample, and

duplicate filters were stored in one petri dish at -20 °C. A filtration blank (1 L MGW) was

processed during each filtration round (n = 21). The total water volume filtered per sample was

recorded (Table S4).

2.4 DNA extraction

DNA was extracted within 2 weeks of filtration at the UoH eDNA facility using the water variant of mu-DNA, a DNA extraction protocol tailored for complex environmental samples (Sellers, Di Muri, Gómez, & Hänfling, 2018). Duplicate filters from samples in Experiment 1 were lysed independently and the lysate from each loaded onto one spin column. Due to a larger sample size in Experiment 2, duplicate filters from samples were co-extracted by placing both filters in a single tube for bead milling. An extraction blank, consisting only of extraction buffers, was included for each round of DNA extraction (n = 17). Eluted DNA (100 µl) was stored at -20 °C until PCR amplification.

2.5 eDNA metabarcoding

Our eDNA metabarcoding workflow is fully described in Supporting Information. Briefly, we performed nested metabarcoding using a two-step PCR protocol, where Multiplex Identification (MID) tags were included in the first and second PCR for sample identification (Kitson et al., 2019). In the first PCR, eDNA was amplified in triplicate with published 12S ribosomal RNA (rRNA) primers 12S-V5-F (5'-ACTGGGATTAGATACCCC-3') and 12S-V5-R (5'-TAGAACAGGCTCCTCTAG-3') (Riaz et al., 2011) that were validated *in silico* for UK vertebrates (Harper et al., 2018a, b). PCR positive controls (two per PCR plate; n = 16) were exotic cichlid (*Maylandia zebra*) DNA (0.05 ng/µl), and PCR negative controls (two per PCR plate; n = 16) were MGW (Fisher Scientific UK Ltd, UK). PCR products were pooled to create sub-libraries (Fig. S2)

and purified. The second PCR bound pre-adapters, MID tags, and Illumina adapters to the sub-

libraries. PCR products were purified, quantified, and pooled, following which the library was

purified, quantified by qPCR, and fragment size (330 bp) and removal of secondary product

verified. The library (220 eDNA samples, 27 field blanks, 33 filtration blanks, 17 extraction

blanks, 16 PCR negative controls, and 16 PCR positive controls) was sequenced on an Illumina

MiSeq®. Raw sequence reads were demultiplexed using a custom Python script, following which

metaBEAT   (metaBarcoding   and   Environmental   Analysis   Tool)   v0.97.11

(https://github.com/HullUni-bioinformatics/metaBEAT) was used for quality trimming, merging,

chimera removal, clustering, and taxonomic assignment of sequences against our UK vertebrate

reference database (Harper et al., 2018a, b) which contains sequences for all UK mammals

including focal species.

2.6 Data analysis

Analyses were performed in R v.3.4.3 (R Core Team, 2017). The total unrefined read counts (i.e.

raw taxonomically assigned reads) per sample were calculated and retained for downstream

analyses. Assignments were corrected and merged, taxon-specific sequence thresholds (i.e. the

maximum sequence frequency of each taxon in PCR positive controls) used to mitigate cross-

contamination and false positives (Fig. S3), and remnant contaminants and higher taxonomic

assignments removed with exceptions. The genus *Anas* (Dabbling ducks) was not removed due

to potential for hybridisation reducing confidence in species-level assignments, and the genera

Emberiza (Buntings) and *Larus* (White-headed gulls) were not removed as reference sequences

were missing for several common species. Consequently, taxonomic assignments in the refined

dataset (i.e. reads left after assignments were corrected, merged, removed, and had sequence

thresholds applied) were predominantly of species resolution and considered true positives.

Dataset refinement is fully described in Supporting Information: Appendix 1. We subsetted the

data by experiment to generate separate datasets for eDNA samples from artificial waterbodies

(Experiment 1) and natural ponds (Experiment 2). Proportional read counts for each species

were calculated from the total unrefined read counts per sample. Our proportional read count

data were not normally distributed (Shapiro–Wilk normality test: $W = 0.915$, $P < 0.001$), thus we

used a Mann-Whitney U test to compare the median proportional read count of stratified and

directed samples across species.

We employed binomial Generalized Linear Mixed Models (GLMMs) with the logit link

function from the development version of the R package glmmTMB (Brooks et al., 2017) for the

following tests. First, we compared the eDNA signals from stratified and directed samples for

each mammal species, with higher proportional read counts expected from directed than

stratified samples. A hierarchical model including sample type nested within species (fixed) and

wildlife park (random) as effects was used. We tested the hypothesis that species lifestyle

would influence mammal eDNA signals, with higher proportional read counts anticipated for

semi-aquatic than ground-dwelling or arboreal species. This model included species lifestyle

(fixed) and species nested within wildlife park (random) as effects. Using the directed samples,

we tested the hypothesis that mammal behaviour influences their eDNA signals. We expected

that behaviour(s) involving water contact would be associated with higher proportional read

counts. We used two hierarchical models including species nested within wildlife park as a

random effect to test the influence of behaviour(s) on proportional read counts. These

modelled specific (e.g. swimming, drinking) or generic (i.e. water contact versus no water

contact) behaviour(s) as a fixed effect. We did not have enough data on mammal behavioural

frequencies or duration to test for an effect on proportional read counts. We assessed model fit

using diagnostic plots and performed validation checks to ensure model assumptions were met

where possible and absence of overdispersion (Zuur, Ieno, Walker, Saveliev, & Smith, 2009).

Predictions for each model were obtained using the *predict* function and 95% confidence

intervals were calculated from the standard error of the predictions.

For Experiment 2, we tested two hypotheses relating to spatiotemporal variation in

mammal eDNA signals at natural ponds. We qualitatively compared mammal presence-absence

records generated by eDNA metabarcoding, camera trapping, and field signs, with eDNA

metabarcoding expected to identify more species than other survey methods. TLNR samples

spanned a 5-day period, where 10 samples were taken from each pond at the same locations

every 24 hrs. For the method performance comparison with BE and TM ponds, which were

sampled once each at 10 locations, the proportional read count data for TLNR samples were

averaged to condense 50 samples to 10 samples per pond. To test the hypothesis that eDNA

signals from aquatic species would be more evenly distributed and persist longer than those

from terrestrial mammals, the unaveraged data from TLNR ponds were used. We qualitatively

assessed change in proportional read counts for identified species over the 5-day period using

tile plots and heat maps. All figures were produced using the R package ggplot2 v3.0.0

(Wickham, 2016).

3. Results

3.1 eDNA metabarcoding

The sequencing run generated 47,713,656 raw sequence reads, of which 37,590,828 remained

following trimming, merging, and length filter application. After removal of chimeras and

redundancy via clustering, the library contained 21,127,061 sequences (average read count of

64,215 per sample including controls), of which 16,787,750 (79.46%) were assigned a

taxonomic rank. Environmental contamination was observed in the field blanks (badger,

beaver, lynx, pine marten, red squirrel, and water vole) as well as environmental and/or

laboratory contamination in the filtration and extraction blanks (human [*Homo sapiens*] and

cichlid). PCR negative controls were contaminated with human, cichlid, beaver, and pine

marten as well as non-focal species (Fig. S4). After threshold application, contaminants

remaining in eDNA samples included Gentoo penguin (*Pygoscelis papua*), reindeer (*Rangifer*

tarandus), cichlid, and human. The refined dataset contained 62 vertebrate species, including

six amphibians, 10 fish, 22 birds, and 24 mammals (Table S6).

3.2 Experiment 1: eDNA detection and signal strength in artificial systems

All nine focal species were detected in captivity. Seven species (red deer and hedgehog

excluded) were detected in all water samples taken from their respective enclosures. Red deer

were not detected in 2 of 5 stratified samples from HWP, and hedgehog was not detected in 1

of 2 WT drinking bowl samples (Fig. 1). 'Other' samples (neither directed nor stratified) were

excluded from further comparisons, thus hedgehog, red squirrel, and water vole were omitted

in downstream analyses.

Across species, stratified samples (0.406) had a higher median proportional read count

than directed samples (0.373), but this difference was not significant (Mann-Whitney U test: U

= 1181.5, P = 0.829). Within species, sample type also had no effect (χ^2_6 = 0.364, P = 0.999) on

proportional read counts (GLMM: θ = 0.168, χ^2_{53} = 8.915, P = 1.000, pseudo-R^2 = 39.21%).

Proportional read counts for directed and stratified samples belonging to each species did not

significantly differ. Otter proportional read counts were lower than other species, but not

significantly so (Fig 2a). Similarly, species lifestyle had no influence (χ^2_2 = 0.655, P = 0.721) on

proportional read counts (GLMM: θ = 0.213, χ^2_{61} = 13.002, P = 1.000, pseudo-R^2 = 11.85%).

Proportional read counts for semi-aquatic, ground-dwelling, and arboreal species did not

significantly differ (Fig. 2b).

Specific behaviours exhibited by species had no influence (χ^2_{11} = 1.369, P = 0.999) on

proportional read counts (GLMM: θ = 0.355, χ^2_{31} = 11.013, P = 0.999, pseudo-R^2 = 9.17%),

where proportional read counts did not differ across species' behaviours (Fig. 3a). Likewise,

generic behaviour (i.e. water contact versus no water contact) had no effect on proportional

read counts (χ^2_{11} = 0.002, P = 0.964). Proportional read counts did not differ between

behaviour that involved or lacked water contact (Fig. 3b, GLMM: θ = 0.217, χ^2_{41} = 8.897, P =

1.000, pseudo-R^2 = 8.50%).

3.3 Experiment 2: eDNA detection and signal strength in natural systems

At natural ponds, eDNA metabarcoding, camera trapping, and field signs all detected beaver, red deer, and roe deer. Camera traps (Fig. 4) and field signs recorded red fox and badger when eDNA metabarcoding did not (Fig. 5). However, eDNA metabarcoding revealed small mammal species not caught on camera, including water vole, water shrew (*Neomys fodiens*), bank vole (*Myodes glareolus*), common shrew (*Sorex araneus*), brown rat (*Rattus norvegicus*), rabbit, grey squirrel, and common pipistrelle (*Pipistrellus pipistrellus*). Mice or vole footprints were observed at BE Pond 1, but species could not be determined. Fig. 5 summarises mammals recorded by different methods at each site with reference to cumulative survey data. Notably, only beaver presence was captured at the same ponds by all methods. Although methods shared species at site level, species were not always detected at the same pond. Detection rates for species captured by at least one survey method are summarised in Table S7.

Sampling of natural ponds revealed spatial patterns in eDNA detection and signal strength. eDNA from non-domestic terrestrial mammals (i.e. mammals excluding dog [*Canis lupus familiaris*], pig [*Sus scrofa domesticus*], sheep [*Ovis aries*] and cow [*Bos taurus*]) was unevenly dispersed compared with semi-aquatic mammals (Fig. S5). Semi-aquatic beaver and water vole were detected in at least 90% and 60% respectively of water samples (*N* = 10) collected from single ponds, whereas water shrew was only detected in 10% of samples. Conversely, non-domestic terrestrial mammals were consistently detected in less than 20% of water samples collected from a pond and left relatively weak eDNA signals. Overall, beaver was

the most consistently detected mammal with the highest proportional read counts. However,

the strongest and most evenly distributed signals belonged to amphibians, particularly common

frog (*Rana temporaria*) and great crested newt (*Triturus cristatus*) (Fig. S5).

TLNR samples collected over a 5-day period (D01-05) revealed that mammal detection

was highly dependent on the spatial and temporal resolution of eDNA metabarcoding surveys

(Fig. S6). Mammal eDNA signals in pond water were ephemeral, often disappearing within 24-

48 hrs of initial detection, as opposed to amphibians that were detected for multiple days and

whose eDNA signal increased in strength. The majority of semi-aquatic or terrestrial mammals

were only detected in a single sample on each day.

4. Discussion

We have demonstrated that eDNA metabarcoding is a potential monitoring tool for

conservation and management priority mammals, but species detection rates are highly

variable. We used two experiments to validate this molecular approach and gained new insights

that will inform the development and application of eDNA metabarcoding for mammals.

Sampling strategy, mammal lifestyle, and mammal behaviour had little influence on eDNA

detection and signal strength in captivity, but all played vital roles in natural ponds. Although

mammals were detected from pond water, their eDNA signals were temporary and weak in

comparison to amphibians. Nonetheless, this would suggest eDNA is representative of

contemporary and local mammal diversity in a given area.

4.1 Experimental insights

In Experiment 1, all nine focal species were detected in captivity, and seven of those species

were detected in all water samples taken from their respective enclosures. This demonstrates

that our method can successfully detect a wide range of mammal species from pond and

drinking water. Perhaps surprisingly, we found that none of sampling strategy, mammal

lifestyle or mammal behaviour influenced eDNA detectability and signal strength in enclosures.

We found no significant differences in eDNA detection or signal strength between stratified and

directed samples collected from artificial waterbodies across or within species. There were no

differences in eDNA signal strength between semi-aquatic, ground-dwelling, and arboreal

species. No relationships between proportional read counts and specific or generic behaviour(s)

were found. This included behaviours associated with eDNA deposition, e.g. swimming,

drinking, urination, and defecation (Rodgers & Mock, 2015; Ushio et al., 2017; Williams et al.,

2018). Enclosures were permanently occupied and artificial waterbodies likely saturated with

eDNA, which possibly masked behavioural signals. Our sample size was constrained as most

focal species were housed by WT only. Modest replication may have limited experimental

power, preventing patterns being detected statistically. Nonetheless, our results have

demonstrated that mammal contact with water enables eDNA deposition and detection.

Unsurprisingly, given the nature of wild mammal interactions with natural systems

versus those in captivity, results from Experiment 2 highlighted the challenges of mammal

eDNA detection. We detected 17 mammals using three monitoring tools, comparable to the 17

mammals expected from cumulative survey data despite some differences. Field signs and

camera trapping performed better than eDNA metabarcoding for red fox and badger detection,

but only eDNA metabarcoding identified water vole and other small mammals missed on

camera or with ambiguous field signs, e.g. mice, voles, shrews. Importantly, the deployment

period, height, and positioning of camera traps may have influenced small mammal detection

by this method (Caravaggi et al., 2018; Glen et al., 2013). Ishige et al. (2017) achieved

comparable mammal detection at salt licks with eDNA metabarcoding and camera trapping, but

species presence was inconsistent between salt licks surveyed. Currently, there are no

published studies comparing eDNA metabarcoding of water to camera trapping for mammal

identification. However, Stat et al. (2019) found highest marine fish species richness using both

baited remote underwater video systems and eDNA metabarcoding, but each method

identified unique species.

Notably, no survey method captured the semi-aquatic otter despite known presence at

study sites and successful detection in eDNA metabarcoding studies of UK ponds (Harper et al.,

2018a) and lakes (Hänfling et al., 2017) as well as Experiment 1 here; although, the eDNA signal

for otter was weaker than other semi-aquatic mammals studied. eDNA metabarcoding may

have performed poorly for otter, badger, and red fox due to species' ecologies. These mammals

are wide-ranging (Gaughran et al., 2018; Thomsen et al., 2012) and may not readily release DNA

in water. Otters often spraint on grass or rock substrata outside water and use latrines

associated with caves and dens (Ruiz-Olmo & Gosálbez, 1997). As terrestrial mammals, red fox

and badger must drink from or enter ponds for eDNA deposition to occur (Rodgers & Mock,

2015; Ushio et al., 2017; Williams et al., 2018). Otter, badger, and red fox detection may require

greater spatiotemporal resolution of eDNA sampling. This is reinforced by other eDNA

metabarcoding studies where mammal detection was highly variable across sites surveyed

(Ishige et al., 2017; Klymus et al., 2017; Ushio et al., 2017). Alternatively, false negatives may be

symptomatic of metabarcoding bias (section 4.2).

eDNA from other semi-aquatic mammals (e.g. beaver, water vole) was evenly

distributed, being found in most or all samples collected on fine spatial scales within natural

ponds, whereas terrestrial mammal (e.g. red and roe deer) eDNA was highly localised and often

detected in less than 20% of samples. Mammal eDNA signals varied temporally, being

detectable for two consecutive days maximum. Depending on the species, mammal eDNA may

be spatially and temporally clumped in lentic ecosystems due to the nature and frequency of

contact that individuals have with water. Unless behaviours involving prolonged water contact

(e.g. swimming, wallowing) are exhibited, non-domestic mammals may only be detected at

drinking sites (Klymus et al., 2017; Ushio et al., 2017; Williams et al., 2018). In contrast,

domestic mammals (e.g. cow, sheep, pig) may have elevated detection rates in ponds due to

high occurrence of these waterbodies in agricultural landscapes as well as eDNA transport by

rainfall and run-off (Staley et al., 2018). eDNA detection and persistence are further influenced

by group size, where eDNA from multiple individuals endures for longer time periods in water

than eDNA from single individuals (Williams et al., 2018). Detailed investigations incorporating

biotic (e.g. population size, sex, body mass, behaviour) and abiotic (e.g. temperature, pH,

ultraviolet light, rainfall) factors are needed to understand the longevity of mammal eDNA

signals in aquatic ecosystems (Parsons et al., 2018; Rodgers & Mock, 2015; Williams et al.,

2018).

Both of our experiments have shown that sampling strategy drastically influences

mammal eDNA detection probability. Mammal eDNA was evenly distributed in closed, artificial

waterbodies, but locally distributed in open, natural ponds. In captivity, mammal species were

housed individually (excluding HWP red deer) with a drinking container(s) and/or small

waterbody (range 0.01-162 m^2, mean 27.4 m^2) in their enclosure. Some enclosures housed

more individuals of a species than others, thereby increasing eDNA deposition and detection

probability (Williams et al., 2018). Wild mammals have an array of freshwater habitats at their

disposal and can hold vast territories. Therefore, rates of pond visitation and eDNA deposition

are more random and irregular (Klymus et al., 2017; Ushio et al., 2017) which may lead to

between-sample variation (Williams et al., 2018).

4.2 Technical pitfalls of eDNA metabarcoding for mammal monitoring

eDNA metabarcoding has strong potential for inclusion in mammal monitoring schemes

(section 4.3), but suffers from biases that may result in false negatives or false positives. Our

process controls identified low-level contamination at all stages of metabarcoding, but

contaminants primarily occurred in field and PCR negative controls (Supporting Information:

Appendix 2). We applied taxon-specific sequence thresholds to our data to mitigate false

positives as in Harper *et al*. (2018a, b). Remnant contaminants were cichlid (laboratory), Gentoo

penguin (environment), reindeer (environment), and human (environment/laboratory). Gentoo

penguin is housed at EZ and was identified from beaver quarantine enclosure water. The WT

red squirrel and reindeer enclosures are in close proximity. DNA transport by wildlife (e.g.

waterfowl [Hänfling et al., 2016]) and park staff or visitors may explain this environmental

contamination. Human DNA was present across process controls corresponding to artificial and

natural waterbodies. Human DNA may be amplified and sequenced instead of focal species,

potentially resulting in false negative detections for rare and/or less abundant species. Human

DNA blocking primers can prevent this bias, but may impair PCR amplification efficiency

(Klymus et al., 2017; Port et al., 2016; Ushio et al., 2017; Valentini et al., 2016).

In our study, eDNA metabarcoding produced false negatives for otter, badger, and red

fox at natural ponds. DNA from purely aquatic and more abundant species may have

overwhelmed otter, badger, and red fox DNA during amplification and sequencing, i.e. species-

masking (Kelly, Port, Yamahara, & Crowder, 2014; Klymus et al., 2017). More biological and

technical replication could help mitigate amplification bias and improve species detection

probabilities (Evans et al., 2017; Lawson Handley et al., 2018; Stat et al., 2019; Valentini et al.,

2016), but potential for primer bias remains. We selected a 12S rRNA primer pair designed to

amplify vertebrate DNA (Riaz et al., 2011) for metabarcoding. There were no primer

mismatches with otter, badger or red fox reference sequences. These species also amplified *in*

silico with up to three mismatches between each primer and reference sequence allowed.

However, other species may have been preferentially amplified *in vitro* due to greater primer

binding affinity (Evans et al., 2017; Kelly et al., 2014; Klymus et al., 2017; Stat et al., 2019).

Metabarcoding primers targeting mammals (Ushio et al., 2017), or multi-marker (e.g. 12S, 16S,

COI) investigations (Evans et al., 2017; Hänfling et al., 2016; Kelly et al., 2014; Klymus et al.,

2017) may improve mammal detection in systems with competition from aquatic species. Yet,

fish and amphibians also had higher qPCR detection than otter (Thomsen et al., 2012). A

comparison of metabarcoding and qPCR would confirm whether poor amplification efficiency

for otter is due to technical bias or species ecology, and whether eDNA metabarcoding can

reliably monitor otter alongside the wider mammal community. Similar comparisons have been

made for the great crested newt (Harper et al., 2018b) and wood turtle (*Glyptemys insculpta*)

(Lacoursière-Roussel, Dubois, Normandeau, & Bernatchez, 2016).

4.3 Scope of eDNA metabarcoding for mammal monitoring

The most recent assessment of UK mammal populations emphasised the lack of data and

systematic monitoring for many species (Mathews et al., 2018). Distribution and occupancy

data are poor for most species, with ongoing survey effort biased toward rare species. Surveys

are also heavily reliant on citizen science and casual records (Massimino et al., 2018). Tools that

can provide standardised, systematic monitoring of UK mammal populations are needed

(Mathews et al., 2018). Despite issues inherent to metabarcoding for biodiversity monitoring

(Deiner et al., 2017), this tool has enormous potential to enhance mammal monitoring,

conservation, and management. eDNA metabarcoding generates distribution data for multiple

species, whether rare, invasive, or abundant, and could be a powerful tool for tracking species

in conflict with one another, e.g. water vole, American mink, and otter (Bonesi & Macdonald,

2004) or red squirrel, grey squirrel, and pine marten (Sheehy, Sutherland, O'Reilly, & Lambin,

2018). Different sample types (e.g. water, soil, snow, salt licks, feeding traces, faeces, hair, and

blood meals) may also offer new insights to mammal biodiversity in a given area (Aylward et al.,

2018; Franklin et al., 2019; Ishige et al., 2017; Tessler et al., 2018; Ushio et al., 2017).

eDNA metabarcoding can rapidly survey multitudes of aquatic sites at landscape-scale

where camera traps might be resource-intensive, cost-inefficient, and susceptible to theft or

damage (Glen et al., 2013; Stat et al., 2019; Ushio et al., 2017). Field signs can be employed at

comparable spatial scales to eDNA metabarcoding but depend on volunteer time and skill

(McShea et al., 2016). eDNA metabarcoding could provide accurate data for species

misidentified from field signs, e.g. mice and voles, otter and mink (Franklin et al., 2019; Harris &

Yalden, 2004). However, both camera traps and field signs recorded species that eDNA

metabarcoding did not. Therefore, eDNA metabarcoding is complementary and should be

incorporated into, not replace, existing monitoring schemes. This tool could be most effective in

mammal monitoring if deployed at the edges of known species distributions, in areas where

species presence is unknown, and in areas with isolated species records (Mathews et al., 2018).

Mammal eDNA detection probabilities from natural ponds will likely be high when areas

with dense populations are studied. However, rigorous sampling strategies are required to

track mammals in areas that are sparsely populated by individuals. Multiple ponds must be

sampled repeatedly, and samples taken at multiple locations within ponds. Ideally, samples and

subsamples would not be pooled to allow inferences about site occupancy to be made.

Importantly, we sampled natural ponds in spring but sampling in other seasons may produce

different results. For example, Lawson Handley et al. (2018) found high repeatability in eDNA

metabarcoding detection and abundance estimates across seasons, but strong spatial

differences reflective of species ecology. We recommend that researchers and practitioners

using eDNA metabarcoding for mammal monitoring channel efforts into extensive sampling of

numerous waterbodies in a given area over larger timescales. This will account for differential

mammal visitation rates and maximise eDNA detection probabilities.

Data accessibility

Acknowledgements

The authors declare that they have no conflicts of interest in presenting this work for

publication. This work was funded by the University of Hull, and D.S.R was supported by the

Natural Environment Research Council award number NE/R016429/1 as part of the UK-SCAPE

programme delivering National Capability. We would like to thank Helen Zenn (RZSS Edinburgh

Zoo) for feedback on the study design, and Richard Griffiths (DICE, University of Kent) for

support with filtration of water samples from Wildwood Trust. We are grateful to staff at

Wildwood Trust and RZSS Highland Wildlife Park for assisting with eDNA sampling from animal

enclosures. We thank Richard Hampshire and volunteers at Tophill Low Nature Reserve for

assisting with camera trap deployment. Tim Kohler (Natural England) and Paul Ramsay kindly

gave permission to sample at Thorne Moors and the Bamff Estate respectively.

Author contributions

L.R.H, B.H, and L.L.H conceived and designed the study. A.I.C and G.M-D coordinated sampling

at Wildwood Trust and RZSS Highland Wildlife Park respectively. L.R.H, C.D.M, C.J.M, and T.L

collected and filtered water samples. A.L, T.L, and T.B helped select natural ponds to be

surveyed using eDNA, camera trapping, and field signs, and provided camera traps for the

study. L.R.H, A.L, and T.L deployed camera traps, which were then collected and footage

analysed by L.R.H. L.R.H processed samples in the laboratory with advice from C.D.M and A.M.

D.S.R sequenced the final library. L.R.H completed bioinformatic processing of samples, and

subsequent data analysis. L.R.H wrote the manuscript, which all authors contributed critically to

drafts of and gave final approval for publication.

References

Aylward, M. L., Johnson, S. E., Sullivan, A. P., Perry, G. H., & Louis, E. E. (2018). A novel environmental DNA (eDNA) sampling method for aye-ayes from their feeding traces. *bioRxiv, 272153*. https://doi.org/10.1101/272153

Baker, C. S., Steel, D., Nieukirk, S., & Klinck, H. (2018). Environmental DNA (eDNA) From the Wake of the Whales: Droplet Digital PCR for Detection and Species Identification. *Frontiers in Marine Science*, *5*, 133.

Bálint, M., Nowak, C., Márton, O., Pauls, S. U., Wittwer, C., Aramayo, J. L., Schulze, A., Chambert, T., Cocchiararo, B. & Jansen, M. (2018). Accuracy, limitations and cost efficiency of eDNA-based community survey in tropical frogs. *Molecular Ecology Resources*, *18*, 1415–1426. https://doi.org/10.1111/1755-0998.12934

Battersby, J. & Tracking Mammals Partnership (2005) *UK Mammals: Species Status and Population Trends. First Report by the Tracking Mammals Partnership*. Joint Nature Conservation Committee /Tracking Mammals Partnership, Peterborough.

Bland, L. M., Collen, B., Orme, C. D. L., & Bielby, J. (2015). Predicting the conservation status of data-deficient species: Predicting Extinction Risk. *Conservation Biology*, *29*, 250–259. https://doi.org/10.1111/cobi12372

Bonesi, L., & Macdonald, D. W. (2004). Differential habitat use promotes sustainable coexistence between the specialist otter and the generalist mink. *Oikos* , *106*, 509–519. https://doi.org/10.1111/j.0030-1299.2004.13034.x

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Machler, M. & Bolker, B. M. (2017). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal*, *9*, 378–400. https://doi.org/10.32614/RJ-2017-066

Burton, A. C., Neilson, E., Moreira, D., Ladle, A., Steenweg, R., Fisher, J. T., Bayne, E. & Boutin, S. (2015). REVIEW: Wildlife camera trapping: a review and recommendations for linking surveys to ecological processes. *Journal of Applied Ecology*, *52*, 675–685. https://doi.org/10.1111/1365-2664.12432

Caravaggi, A., Gatta, M., Vallely, M. -C., Hogg, K., Freeman, M., Fadaei, E., Dick, J. T. A., Montgomery, W. I., Reid, N. & Tosh, D. G. (2018). Seasonal and predator-prey effects on circadian activity of free-ranging mammals revealed by camera traps. *PeerJ*, *6*, e5827. https://doi.org/10.7717/peerj.5827

Deiner, K., Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., Creer, S., Bista, I., Lodge, D. M., de Vere, N., Pfrender, M. E. & Bernatchez, L. (2017). Environmental DNA metabarcoding: transforming how we survey animal and plant communities. *Molecular Ecology*, *26*, 5872–5895. https://doi.org/10.1111/mec.14350

Evans, N. T., Li, Y., Renshaw, M. A., Olds, B. P., Deiner, K., Turner, C. R., Jerde, C. L., Lodge, D. M.,
Lamberti, G. A. & Pfrender, M. E (2017). Fish community assessment with eDNA
metabarcoding: effects of sampling design and bioinformatic filtering. *Canadian Journal of*
Fisheries and Aquatic Sciences, *74*, 1362–1374. https://doi.org/10.1139/cjfas-2016-0306

Franklin, T. W., McKelvey, K. S., Golding, J. D., Mason, D. H., Dysthe, J. C., Pilgrim, K. L., Squires,
J. R., Aubry, K. B., Long, R. A., Greaves, S. E., Raley, C. M., Jackson, S., MacKay, P., Lisbon, J.,
Sauder, J. D., Pruss, M. T., Heffington, D. & Schwartz, M. K (2019). Using environmental DNA
methods to improve winter surveys for rare carnivores: DNA from snow and improved
noninvasive techniques. *Biological Conservation*, *229*, 50–58.
https://doi.org/10.1016/j.biocon.2018.11.006

Gaughran, A., Kelly, D. J., MacWhite, T., Mullen, E., Maher, P., Good, M., & Marples, N. M.
(2018). Super-ranging. A new ranging strategy in European badgers. *PLoS ONE*, *13*,
e0191818. https://doi.org/10.1371/journal.pone.0191818

Glen, A. S., Cockburn, S., Nichols, M., Ekanayake, J., & Warburton, B. (2013). Optimising camera
traps for monitoring small mammals. *PLoS ONE*, *8*, e67940.
https://doi.org/10.1371/journal.pone.0067940

Hänfling, B., Lawson Handley, L., Lunt, D., Shum, P., Winfield, I., & Read, D. S. (2017). *A Review*
of Recent Advances in Genetic Methods to Identify Improvements in CAMERAS Partners
Monitoring Activities. University of Hull, Hull, UK.

Hänfling, B., Lawson Handley, L., Read, D. S., Hahn, C., Li, J., Nichols, P., Blackman, R. C., Oliver,
A. & Winfield, I. J. (2016). Environmental DNA metabarcoding of lake fish communities
reflects long-term data from established survey methods. *Molecular Ecology*, *25*, 3101–3119.
https://doi.org/10.1111/mec.13660

Harper, L. R., Handley, L. L., Hahn, C., Boonham, N., Rees, H. C., Lewis, E., Adams, I. P.,
Brotherton, P., Phillips, S. & Hänfling, B. (2018a) Ground-truthing environmental DNA (eDNA)
metabarcoding for ecological hypothesis testing at the pondscape. *bioRxiv*, *278309*.
https://doi.org/10.1101/278309

Harper, L. R., Lawson Handley, L., Hahn, C., Boonham, N., Rees, H. C., Gough, K. C., Lewis, E.,
Adams, I. P., Brotherton, P., Phillips, S. & Hänfling, B. (2018b). Needle in a haystack? A
comparison of eDNA metabarcoding and targeted qPCR for detection of the great crested
newt (*Triturus cristatus*). *Ecology and Evolution*, *8*, 6330–6441.
https://doi.org/10.1002/ece3.4013

Harris, S., & Yalden, D. W. (2004). An integrated monitoring programme for terrestrial mammals
in Britain. *Mammal Review*, *34*, 157–167. https://doi.org/10.1046/j.0305-1838.2003.00030.x

Hunter, M. E., Meigs-Friend, G., Ferrante, J. A., Takoukam Kamla, A., Dorazio, R. M., Keith-
Diagne, L., Luna, F., Lanyon, J. M. & Reid, J. P. (2018). Surveys of environmental DNA (eDNA):
a new approach to estimate occurrence in Vulnerable manatee populations. *Endangered*
Species Research, *35*, 101–111. https://doi.org/10.3354/esr00880

Ishige, T., Miya, M., Ushio, M., Sado, T., Ushioda, M., Maebashi, K., Yonechi, R., Lagan, P. &
Matsubayashi, H. (2017). Tropical-forest mammals as detected by environmental DNA at
natural saltlicks in Borneo. *Biological Conservation*, *210*, 281–285.
https://doi.org/10.1016/j.biocon.2017.04.023

Joint Nature Conservation Committee (2018) UK BAP priority terrestrial mammal species.
Accessed on 1st December 2018. http://jncc.defra.gov.uk/page-5170

Kelly, R. P., Port, J. A., Yamahara, K. M., & Crowder, L. B. (2014). Using environmental DNA to
census marine fishes in a large mesocosm. *PLoS ONE*, *9*, e86175.
https://doi.org/10.1371/journal.pone.0086175

Kitson, J. J. N., Hahn, C., Sands, R. J., Straw, N. A., Evans, D. M., & Lunt, D. H. (2019). Detecting
host-parasitoid interactions in an invasive Lepidopteran using nested tagging DNA-
metabarcoding. *Molecular Ecology*, *28*, 471–483. https://doi.org/10.1111/mec.14518

Klymus, K. E., Richter, C. A., Thompson, N., & Hinck, J. E. (2017). Metabarcoding of
Environmental DNA Samples to Explore the Use of Uranium Mine Containment Ponds as a
Water Source for Wildlife. *Diversity*, *9*, 54. https://doi.org/10.3390/d9040054

Lacoursière-Roussel, A., Dubois, Y., Normandeau, E., & Bernatchez, L. (2016). Improving
herpetological surveys in eastern North America using the environmental DNA method.
Genome, *59*, 991–1007. https://doi.org/10.1139/gen-2015-0218

Lawson Handley, L. J., Read, D., Winfield, I., Kimbell, H., Johnson, H., Li, J., Hahn, C., Blackman,
R., Wilcox, R., Donnelly, R., Szitenberg, A. & Hänfling, B. (2018) Temporal and spatial
variation in distribution of fish environmental DNA in England's largest lake. *bioRxiv*, *376400*.
https://doi.org/10.1101/376400

Lugg, W. H., Griffiths, J., van Rooyen, A. R., Weeks, A. R., & Tingley, R. (2017). Optimal survey
designs for environmental DNA sampling. *Methods in Ecology and Evolution*, *9*, 1049–1059.
https://doi.org/10.1111/2041-210X.12951

Lynx UK Trust (2018) Accessed on 1st December 2018. http://www.lynxuk.org/

Massimino, D., Harris, S. J., & Gillings, S. (2018). Evaluating spatiotemporal trends in terrestrial
mammal abundance using data collected during bird surveys. *Biological Conservation*, *226*,
153–167. https://doi.org/10.1016/j.biocon.2018.07.026

Mathews, F., Kubasiewicz, L. M., Gurnell, J., Harrower, C. A., McDonald, R. A. & Shore, R. F.
(2018) *A Review of the Population and Conservation Status of British Mammals: Technical*
Summary. Natural England, Peterborough.

McShea, W. J., Forrester, T., Costello, R., He, Z., & Kays, R. (2016). Volunteer-run cameras as
distributed sensors for macrosystem mammal research. *Landscape Ecology*, *31*, 55–66.
https://doi.org/10.1007/s10980-015-0262-9

Padgett-Stewart, T. M., Wilcox, T. M., Carim, K. J., McKelvey, K. S., Young, M. K., & Schwartz, M.
K. (2016). An eDNA assay for river otter detection: a tool for surveying a semi-aquatic
mammal. *Conservation Genetics Resources*, *8*, 5–7. https://doi.org/10.1007/s12386-015-
0511-s

Parsons, K. M., Everett, M., Dahlheim, M., & Park, L. (2018). Water, water everywhere:
environmental DNA can unlock population structure in elusive marine species. *Royal Society*
Open Science, *5*, 180537. https://doi.org/10.1098/rsos.180537

Port, J. A., O'Donnell, J. L., Romero-Maraccini, O. C., Leary, P. R., Litvin, S. Y., Nickols, K. J.,
Yamahara, K. M. & Kelly, R. P. (2016). Assessing vertebrate biodiversity in a kelp forest
ecosystem using environmental DNA. *Molecular Ecology*, *25*, 527–541.
https://doi.org/10.1111/mec.13481

Riaz, T., Shehzad, W., Viari, A., Pompanon, F., Taberlet, P., & Coissac, E. (2011). ecoPrimers:
inference of new DNA barcode markers from whole genome sequence analysis. *Nucleic Acids*
Research, *39*, e145. https://doi.org/10.1093/nar/gkr732

Rodgers, T. W., & Mock, K. E. (2015). Drinking water as a source of environmental DNA for the
detection of terrestrial wildlife species. *Conservation Genetics Resources*, *7*, 693–696.
https://doi.org/10.1007/s12686-015-0478-7

Ruiz-Olmo, J., & Gosálbez, J. (1997). Observations on the sprainting behaviour of the otter *Lutra*
lutra in the NE Spain. *Acta Theriologica*, *42*, 259–270.

Sadlier, L. M. J., Webbon, C. C., Baker, P. J., & Harris, S. (2004). Methods of monitoring red foxes
Vulpes vulpes and badgers Meles meles: are field signs the answer? *Mammal Review*, *34*,
75–98. https://doi.org/10.1046/j.0305-1838.2003.00029.x

Sellers, G. S., Di Muri, C., Gómez, A., & Hänfling, B. (2018). Mu-DNA: a modular universal DNA
extraction method adaptable for a wide range of sample types. *Metabarcoding and*
Metagenomics, *2*, e24556. https://doi.org/10.3897/mbmg.2.24556

Sheehy, E., Sutherland, C., O'Reilly, C. & Lambin, X. (2018) The enemy of my enemy is my friend:
native pine marten recovery reverses the decline of the red squirrel by suppressing grey
squirrel populations. *Proceedings of the Royal Society of London B*, *285*, 20172603.
https://doi.org/10.1098/rspb.2017.2603

Staley, Z. R., Chuong, J. D., Hill, S. J., Grabuski, J., Shokralla, S., Hajibabaei, M., & Edge, T. A.
(2018). Fecal source tracking and eDNA profiling in an urban creek following an extreme rain
event. *Scientific Reports*, *8*, 14390. https://doi.org/10.1038/s41598-018-32680-z

Stat, M., John, J., DiBattista, J. D., Newman, S. J., Bunce, M. & Harvey, E. S. (2019) Combined use
of eDNA metabarcoding and video surveillance for the assessment of fish biodiversity.
Conservation Biology, *33*, 196–205. https://doi.org/10.1111/cobi.13183

Tessler, M., Weiskopf, S. R., Berniker, L., Hersch, R., Mccarthy, K. P., Yu, D. W., & Siddall, M. E.
(2018). Bloodlines: mammals, leeches, and conservation in southern Asia. *Systematics and*
Biodiversity, *16*, 488–496. https://doi.org/10.1080/14772000.2018.1433729

Thomsen, P. F., Iversen, L. L., Wiuf, C., Rasmussen, M., Gilbert, M. T. P., Orlando, L., &
Willerslev, E. (2012). Monitoring endangered freshwater biodiversity using environmental
DNA. *Molecular Ecology*, *21*, 2565–2573. https://doi.org/10.1111/j.1365-294X.2011.05418.x

Thomsen, P. F., & Willerslev, E. (2015). Environmental DNA–an emerging tool in conservation
for monitoring past and present biodiversity. *Biological Conservation*, *183*, 4–18.
https://doi.org/10.1016/j.biocon.2014.11.019

Ushio, M., Fukuda, H., Inoue, T., Makoto, K., Kishida, O., Sato, K., Murata, K., Nikaido, M., Sado,
T., Sato, Y., Takeshita, M., Iwasaki, W., Yamanaka, H., Kondoh, M. & Miya, M. (2017).
Environmental DNA enables detection of terrestrial mammals from forest pond water.
Molecular Ecology Resources, *17*(6), e63–e75. https://doi.org/10.1111/1755-0998.12690

Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P. F., Bellemain, E.,
Besnard, A., Coissac, E., Boyer, F., Gaboriaud, C., Jean, P., Poulet, N., Roset, N., Copp, G. H.,
Geniez, P., Pont, D., Argillier, C., Baudoin, J. -M., Peroux, T., Crivelli, A. J., Olivier, A.,
Acqueberge, M., Le Brun, M., Møller, P. R., Willerslev, E. & Dejean, T. (2016). Next-
generation monitoring of aquatic biodiversity using environmental DNA metabarcoding.
Molecular Ecology, *25*, 929–942. https://doi.org/10.1111/mec.13428

Visconti, P., Pressey, R. L., Giorgini, D., Maiorano, L., Bakkenes, M., Boitani, L., Alkemade, R.,
Falcucci, A., Chiozza, F. & Rondinini, C. (2011). Future hotspots of terrestrial mammal loss.
Philosophical Transactions of the Royal Society B, *366*, 2693–2702.
https://doi.org/10.1098/rstb.2011.0105

Wickham, H. (2016). *ggplot2: elegant graphics for data analysis*. Springer, New York, USA.
http://ggplot2.org

Williams, K. E., Huyvaert, K. P., Vercauteren, K. C., Davis, A. J., & Piaggio, A. J. (2018). Detection
and persistence of environmental DNA from an invasive, terrestrial mammal. *Ecology and*
Evolution, *8*, 688–695. https://doi.org/10.1002/ece3.3698

Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A. & Smith, G. M. (2009) *Mixed effects models*
and extensions in ecology with R. Springer, New York, USA.

Table 1. Summary of focal species studied at wildlife parks and their lifestyle. The number of

individuals present and waterbody size in enclosures is provided.

Site	Species	Lifestyle	Enclosure	Number of individuals	Waterbody size (m^2)
Wildwood Trust	European otter (*Lutra lutra*)	Semi-aquatic	1	2	162
	European water vole (*Arvicola amphibius*)	Semi-aquatic	1	4	0.09
			2	1	0.09
	European beaver (*Castor fiber*)	Semi-aquatic	1	2	100
			2	1	100
	European hedgehog (*Erinaceus europaeus*)	Ground-dwelling	1	1	0.04
			2	2	0.04
	European badger (*Meles meles*)	Ground-dwelling	1	4	1.73
	Red deer (*Cervus elaphus*)	Ground-dwelling	1	8	100
	Eurasian lynx (*Lynx lynx*)	Ground-dwelling	1	2	2
	Red squirrel (*Sciurus vulgaris*)	Arboreal	1	2	0.01
			2	3	0.01
			3	3	0.01
			4	2	0.01
	European pine marten (*Martes martes*)	Arboreal	1	1	2
			2	1	0.375
Highland Wildlife Park	Red squirrel (*Sciurus vulgaris*)	Arboreal	NA	NA	0.25
	Eurasian lynx (*Lynx lynx*)	Ground-dwelling	1	8	2
	European beaver (*Castor fiber*)	Semi-aquatic	1	2	50
	Red deer (*Cervus elaphus*)	Ground-dwelling	1	30	NA

Figure 1. Heatmap showing proportional read counts for eDNA samples. Each square

800 represents a sample taken from a wildlife park enclosure. Directed (DIR01-DIR06) and stratified

801 (STR01-STR06) samples were collected from artificial waterbodies. Samples were collected from

802 drinking containers (E1, E2, E3, E4, BOWL, BUCK), water vole (QUAR1, QUAR2) and beaver

803 (ZOO) quarantine enclosures, and a water bath (BATH) in RZSS Highland Wildlife Park woods.

Figure 2. Relationships predicted by the binomial GLMMs between proportional read counts

and sample type nested within species **(a)** or species lifestyle **(b)**. The observed data (coloured

points) are displayed against the predicted relationships (black points with error bars) for each

species **(a)** or species lifestyle **(b)**. Points are shaped by sample type **(a)** or wildlife park **(b)**, and

coloured by species lifestyle. Error bars represent the standard error around the predicted

means.

Figure 3. Boxplots showing the mean proportional read counts for specific **(a)** and generic **(b)** behaviour(s) exhibited by focal species. Boxes show 25th, 50th, and 75th percentiles, and whiskers show 5th and 95th percentiles. Points are coloured by species lifestyle, and each point in **(a)** represents a directed sample sized by frequency of behaviour. The behaviour 'none' for beaver represents occurrences of beaver in water but out of view of camera traps.

Figure 4. Exemplar camera trap photos taken at natural ponds where focal species were present. Red deer was recorded at Thorne Moors **(a)**, roe deer **(b)** and red fox **(c)** were recorded at Tophill Low Nature Reserve, and beaver was recorded at the Bamff Estate **(d)**.

Figure 5. Tile plot showing species presence-absence at individual pond and site-level as indicated by field signs, camera trapping, and eDNA metabarcoding at sites with focal species presence confirmed by cumulative survey data.