Environmental DNA (eDNA) metabarcoding of pond water as a tool to

survey conservation and management priority mammals

Lynsey R. Harper[a*], Lori Lawson Handley[a], Angus I. Carpenter[b], Muhammad Ghazali[c], Cristina

Di Muri[a], Callum J. Macgregor[d], Thomas W. Logan[a], Alan Law[e], Thomas Breithaupt[a], Daniel S.

Read[f], Allan D. McDevitt[g], and Bernd Hänfling[a]

[a] Department of Biological and Marine Sciences, University of Hull, Hull, HU6 7RX, UK

[b] Wildwood Trust, Canterbury Rd, Herne Common, Herne Bay, CT6 7LQ, UK

[c] The Royal Zoological Society of Scotland, Edinburgh Zoo, 134 Corstorphine Road, Edinburgh, EH12 6TS, UK

[d] Department of Biology, University of York, Wentworth Way, York, YO10 5DD, UK

[e] Biological and Environmental Sciences, University of Stirling, Stirling, FK9 4LA, UK

[f] Centre for Ecology & Hydrology (CEH), Benson Lane, Crowmarsh Gifford, Wallingford, Oxfordshire, OX10 8BB, UK

[g] Ecosystems and Environment Research Centre, School of Science, Engineering and Environment, University of

Salford, Salford, M5 4WT, UK

*Corresponding author: lynsey.harper2@gmail.com

Lynsey Harper, Department of Biological and Marine Sciences, University of Hull, Hull, HU6 7RX,

UK

Abstract

Environmental DNA (eDNA) metabarcoding can identify terrestrial taxa utilising aquatic habitats

alongside aquatic communities, but terrestrial species' eDNA dynamics are understudied. We

evaluated eDNA metabarcoding for monitoring semi-aquatic and terrestrial mammals,

specifically nine species of conservation or management concern, and examined

spatiotemporal variation in mammal eDNA signals. We hypothesised eDNA signals would be

stronger for semi-aquatic than terrestrial mammals, and at sites where individuals exhibited

behaviours. In captivity, we sampled waterbodies at points where behaviours were observed

('directed' sampling) and at equidistant intervals along the shoreline ('stratified' sampling). We

surveyed natural ponds (N = 6) where focal species were present using stratified water

sampling, camera traps, and field signs. eDNA samples were metabarcoded using vertebrate-

specific primers. All focal species were detected in captivity. eDNA signal strength did not differ

between directed and stratified samples across or within species, between semi-aquatic or

terrestrial species, or according to behaviours. eDNA was evenly distributed in artificial

waterbodies, but unevenly distributed in natural ponds. Survey methods deployed at natural

ponds shared three species detections. Metabarcoding missed badger and red fox recorded by

cameras and field signs, but detected small mammals these tools overlooked, e.g. water vole.

Terrestrial mammal eDNA signals were weaker and detected less frequently than semi-aquatic

mammal eDNA signals. eDNA metabarcoding could enhance mammal monitoring through

large-scale, multi-species distribution assessment for priority and difficult to survey species, and

provide early indication of range expansions or contractions. However, eDNA surveys need high

44   spatiotemporal resolution and metabarcoding biases require further investigation before

45   routine implementation.

47   **Key-words:** camera traps, field signs, lentic, monitoring, semi-aquatic mammals, terrestrial

48   mammals

1. Introduction

Mammals are a highly threatened taxon, with 25% of species at risk of extinction globally due to harvesting, habitat degradation/loss, non-native species or perception as pests (Visconti et al., 2011). Most species lack long-term, systematic monitoring, with survey efforts biased towards rare species (Massimino, Harris, & Gillings, 2018). Data deficiency prevents robust estimation of mammalian range expansions/declines and population trends (Bland, Collen, Orme, & Bielby, 2015). Therefore, effective and evidence-based strategies for mammal conservation and management are urgently needed (Mathews et al., 2018).

Many mammals are nocturnal and elusive thus monitoring requires non-invasive, observational methods such as camera traps and field signs, e.g. footprints, scat (Caravaggi et al., 2018; Harris & Yalden, 2004; Kinoshita et al., 2019; Sadlier, Webbon, Baker, & Harris, 2004). Camera trapping is cost-efficient, standardised, reproducible, and produces data suited to site occupancy modelling, but only surveys a fraction of large, heterogeneous landscapes. Trap placement can substantially influence species detection probabilities, and traps often miss small species (Burton et al., 2015; Caravaggi et al., 2018; Ishige et al., 2017; Leempoel, Hebert, & Hadly, 2019). Field sign surveys are inexpensive, but resource-intensive for broad geographic coverage (Kinoshita et al., 2019; Sadlier et al., 2004). Species can have similar footprints and scat, increasing the potential for misidentification (Franklin et al., 2019; Harris & Yalden, 2004). Mammal survey methods can be species-specific, thus multiple methods are necessary for large-scale, multi-species monitoring schemes (Massimino et al., 2018; Sales et al., 2019).

Environmental DNA (eDNA) analysis is a recognised tool for rapid, non-invasive, cost-

efficient biodiversity assessment across aquatic and terrestrial ecosystems (Deiner et al., 2017).

Organisms transfer genetic material to their environment via secretions, excretions, gametes,

blood, or decomposition, which can be isolated from environmental samples (Thomsen &

Willerslev, 2015). Studies using eDNA analysis to target specific semi-aquatic and terrestrial

mammals have employed PCR or quantitative PCR (qPCR) (e.g. Franklin et al., 2019; Lugg,

Griffiths, van Rooyen, Weeks, & Tingley, 2017; Rodgers & Mock, 2015; Thomsen et al., 2012;

Williams, Huyvaert, Vercauteren, Davis, & Piaggio, 2018). eDNA metabarcoding can screen

entire communities using PCR combined with high-throughput sequencing (Deiner et al., 2017;

Thomsen & Willerslev, 2015), but mammalian assessments are uncommon (Klymus, Richter,

Thompson, & Hinck, 2017; Kinoshita et al., 2019; Leempoel et al., 2019; Sales et al., 2019; Ushio

et al., 2017). Tropical mammal assemblages have been obtained by metabarcoding invertebrate

blood meals (e.g. Tessler et al., 2018) and salt licks (Ishige et al., 2017), but samples from the

physical environment have tremendous potential to reveal mammal biodiversity over broad

spatiotemporal scales (Sales et al., 2019; Ushio et al., 2017).

In aquatic ecosystems, eDNA metabarcoding has predominantly been applied to

characterise fish (e.g. Evans et al., 2017; Hänfling et al., 2016; Lawson Handley et al., 2018;

Valentini et al., 2016) and amphibian (e.g. Bálint et al., 2018; Valentini et al., 2016)

communities. However, mammals also leave eDNA signatures in water that metabarcoding can

detect (Harper et al., 2019; Klymus et al., 2017; Sales et al., 2019; Ushio et al., 2017). Ponds in

particular provide drinking, foraging, dispersive, and reproductive opportunities for semi-

aquatic and terrestrial mammals (Klymus et al., 2017). Samples from these waterbodies could

uncover biodiversity present in the wider environment (Deiner et al., 2017; Harper et al., 2019).

Drinking is a major source of eDNA deposition due to the release of saliva, but mammals may

also swim, wallow, urinate or defecate in water (Rodgers & Mock, 2015; Ushio et al., 2017;

Williams et al., 2018). Furthermore, arboreal mammals may use ponds less than semi-aquatic

and ground-dwelling species, non-territorial mammals may visit ponds less than territorial

species, and group-living species may deposit more eDNA than solitary species (Williams et al.,

2018). Despite evidence for eDNA deposition by semi-aquatic and terrestrial mammals in

freshwater ecosystems, little is known about the influence of mammal behaviour on the

distribution and strength of the eDNA signal left behind (defined here as proportional read

counts).

In this study, we conducted two experiments under artificial and natural conditions to

evaluate eDNA metabarcoding of pond water as a tool for monitoring semi-aquatic, ground-

dwelling, and arboreal mammals of conservation or management concern. The first

experiment, carried out on nine focal species housed at two wildlife parks, examined the role of

sampling strategy, mammal lifestyle, and mammal behaviour on eDNA detection and signal

strength under artificial conditions. Mammal eDNA detection is expected from enclosure water

that is frequently used by individuals for drinking, swimming and bathing. We hypothesised

that: (1) eDNA would be unevenly distributed, thus directed sampling would yield stronger

eDNA signals (i.e. higher proportional read counts) for mammals than stratified sampling; (2)

semi-aquatic mammals would have stronger eDNA signals than ground-dwelling or arboreal

mammals; and (3) mammal behaviours involving water contact would generate stronger eDNA

signals. The second experiment validated eDNA metabarcoding against camera trapping and

field sign searches for mammal identification at natural ponds, and investigated spatiotemporal

variation in mammal eDNA signals. Mammal eDNA detection is unpredictable at natural

waterbodies that can be extensive, subject to environmental fluctuations, and used rarely or

not at all by individuals. We hypothesised that: (1) eDNA metabarcoding would detect more

mammals than camera trapping or field signs; (2) semi-aquatic mammals would be readily

detected and their eDNA evenly distributed in ponds in comparison to terrestrial mammals; and

(3) temporal sampling would reveal that terrestrial mammal eDNA is detectable for short

periods in comparison to fully aquatic vertebrates.

2. Materials and methods

2.1 Study species

We studied nine mammal species that are the focus of European conservation or management

(Mathews et al., 2018): European water vole (*Arvicola amphibius*), European otter (*Lutra lutra*),

Eurasian beaver (*Castor fiber*), European hedgehog (*Erinaceus europaeus*), European badger

(*Meles meles*), red deer (*Cervus elaphus*), Eurasian lynx (*Lynx lynx*), red squirrel (*Sciurus*

vulgaris), and European pine marten (*Martes martes*). Water vole, otter, red squirrel, pine

marten and hedgehog are UK Biodiversity Action Plan species (Joint Nature Conservation

Committee, 2018). Water vole, otter, and beaver are semi-aquatic, red squirrel and pine

marten are arboreal, and the other species are ground-dwelling. Badger and red deer live in

groups whereas the other species are predominantly solitary.

2.2 Experiment 1: eDNA detection and signal strength in artificial systems

Behavioural observation and eDNA sampling were conducted between 18[th] – 21[st] September

2017 at Wildwood Trust (WT), Kent, England, and 10[th] – 11[th] October 2017 at Royal Zoological

Society of Scotland (RZSS) Highland Wildlife Park (HWP), Kingussie, Scotland. Sixteen categories

of behaviour were defined based on potential contact with waterbodies and species lifestyle,

and the frequency and duration of behaviours recorded (Table 1, Appendix A: Table A1). The

number of individuals in each enclosure was recorded alongside waterbody size (Table 2).

Beaver, lynx, red deer, and red squirrel were present at both wildlife parks, whereas other

captive species were only present at WT. Each species was observed for one hour on two

separate occasions except nocturnal mammals (badger and beaver), which were observed

overnight using camera traps (Bushnell Trophy Cam Standard, Bushnell Corporation, KS, USA).

One camera trap per enclosure was positioned perpendicular to the ground (1 m height, 2 m

from shoreline) to capture water and shoreline. Cameras took 30 s videos (1920 x 1080) when

triggered (30 s interval between triggers) at high sensitivity. Behavioural observation was not

undertaken for WT water voles as animals were under quarantine or HWP red squirrels as

individuals were wild. Photos of waterbodies in animal enclosures are provided in Appendix B.

Water samples were collected from enclosures within 3 hrs of the second behavioural

observation period. Up to six directed or stratified samples were collected, but sample number

varied by species according to waterbody size and observed behaviours (Tables A1, A2).

Enclosure drinking containers were also sampled and classed as 'other' samples. Bathing and

drinking bowls were sampled where enclosures contained no artificial waterbodies (WT water

vole, red squirrel, and hedgehog). The HWP beaver enclosure was empty for 24 hrs before

sampling. Water was sampled from a RZSS Edinburgh Zoo (EZ) enclosure containing beavers

and classed as 'other'. A sample was collected from a water bath in the HWP woods to capture

wild red squirrels and classed as 'other'.

Directed samples (2 L surface water taken approximately where behaviours were

observed) were collected before stratified samples (2 L surface water [8 x 250 ml pooled

subsamples] taken at equidistant points [access permitting] around the waterbody perimeter)

to minimise disturbance to the water column and cross-contamination risk. Samples were

collected using sterile Gosselin™ HDPE plastic bottles (Fisher Scientific UK Ltd, UK) and

disposable gloves. A field blank (1 L molecular grade water [MGW]) was taken into each species

enclosure, opened, and closed before artificial water sources were sampled. Samples (n = 80)

collected from WT and HWP were transported alongside field blanks (n = 13) in sterile

coolboxes with ice packs to the University of Kent (UoK) and EZ respectively, where ice was

added to coolboxes.

Samples and blanks were vacuum-filtered within 6 hrs of collection in a UoK wet

laboratory and within 24 hrs of collection in an EZ staff room. Surfaces and equipment were

sterilised before, during, and after set-up in temporary work areas. Surfaces and vacuum

pumps were wiped with 10% v/v chlorine-based commercial bleach (Elliott Hygiene Ltd, UK)

solution. Non-electrical equipment was immersed in 10% bleach solution for 10 minutes,

followed by 5% v/v MicroSol detergent (Anachem, UK), and rinsed with purified water. Up to

500 ml of each 2 L sample was vacuum-filtered through sterile 0.45 μm mixed cellulose ester

membrane filters with pads (47 mm diameter; Whatman, GE Healthcare, UK) using Nalgene™

filtration units. One hour was allowed for each sample to filter and a second filter used if

clogging occurred. A filtration blank (1 L MGW) was processed during each filtration round ($n =$

12), and equipment sterilised after each filtration round. After 500 ml had filtered or one hour

had passed, filters were removed from pads using sterile tweezers, placed in sterile 47 mm

petri dishes (Fisher Scientific UK Ltd, UK), sealed with parafilm (Sigma-Aldrich Company Ltd,

UK), and stored at -20 °C. The total water volume filtered per sample was recorded for

downstream analysis (Table A2; Fig. A1).

2.3 Experiment 2: eDNA detection and signal strength in natural systems

At three sites where focal species were present based on cumulative survey data, we selected

two ponds (range 293-5056 m^2, average 1471 m^2) within 4 km of each other. The Bamff Estate

(BE), Alyth, Scotland, was selected for beaver, otter, badger, red deer, and red squirrel, but roe

deer (*Capreolus capreolus*) and red fox (*Vulpes vulpes*) were also present. Otter, water vole, and

badger were present at Tophill Low Nature Reserve (TLNR), Driffield, East Yorkshire, alongside

American mink (*Neovison vison*), stoat (*Mustela erminea*), weasel (*Mustela nivalis*), rabbit

(*Oryctolagus cuniculus*), brown hare (*Lepus europaeus*), red fox, roe deer, and grey squirrel

(*Sciurus carolinensis*). We selected Thorne Moors (TM), Doncaster, South Yorkshire, for red deer

and badger, but stoat, weasel, red fox, roe deer, and Reeve's muntjac (*Muntiacus reevesi*) were

also present. Camera traps (Bushnell Trophy Cam Standard/Aggressor, Bushnell Corporation,

KS, USA) were deployed at TM (one per pond) and BE (three per pond) one week prior to eDNA

sampling and collected once sampling was completed. At TLNR, camera traps (two to three per

pond) were deployed one day before a 5-day period of eDNA sampling and collected one week

after sampling was completed. Camera traps were positioned perpendicular to the ground (1 m

height, 0.3-1 m from shoreline) to capture water and shoreline. Cameras took three

photographs (5 megapixel) when triggered (3 s interval between triggers) at high sensitivity.

209   Ten stratified samples were collected from the shoreline of each pond (TM: 17[th] April

2018; BE: 20[th] April 2018; TLNR: 23[rd] – 27[th] April 2018) and a field blank (1 L MGW) included as

in Experiment 1. TLNR ponds were sampled every 24 hrs over 5 days to investigate

spatiotemporal variation in mammal eDNA signals. TM and TLNR samples were transported on

ice in sterile coolboxes to the University of Hull (UoH) eDNA facility, and stored at 4 °C. BE

samples were transported in sterile coolboxes with ice packs to BE accommodation. Surfaces

and equipment were sterilised before, during, and after set-up as in Experiment 1. Samples (n =

140) and field blanks (n = 14) were vacuum-filtered within 4 hrs of collection as in Experiment 1

with minor modifications to maximise detection probability as follows. The full 2 L of each

sample was vacuum-filtered where possible, two filters were used for each sample, and

duplicate filters were stored in one petri dish at -20 °C. A filtration blank (1 L MGW) was

processed during each filtration round (n = 21). The total water volume filtered per sample was

recorded (Table A3).

2.4 DNA extraction

DNA was extracted within 2 weeks of filtration at the UoH eDNA facility using the Mu-DNA water protocol (Sellers, Di Muri, Gómez, & Hänfling, 2018). The full protocol is available at: https://doi.org/10.17504/protocols.io.qn9dvh6. Duplicate filters from samples in Experiment 1 were lysed independently and the lysate from each loaded onto one spin column. As more samples were collected in Experiment 2, duplicate filters were co-extracted by placing both in a single tube for bead milling. An extraction blank, consisting only of extraction buffers, was included for each round of DNA extraction (n = 17). Eluted DNA (100 µl) was stored at -20 °C until PCR amplification.

2.5 eDNA metabarcoding

Our eDNA metabarcoding workflow is fully described in Appendix A. Briefly, we performed nested metabarcoding using a two-step PCR protocol, where Multiplex Identification (MID) tags were included in the first and second PCR for sample identification (Kitson et al., 2019). The first PCR amplified eDNA in triplicate with published 12S ribosomal RNA (rRNA) primers 12S-V5-F (5'-ACTGGGATTAGATACCCC-3') and 12S-V5-R (5'-TAGAACAGGCTCCTCTAG-3') (Riaz et al., 2011). Harper et al. (2018) validated these primers *in silico* for UK vertebrates, and found 91/112 mammal species listed on the Natural History Museum Checklist of Mammalia v1 (subspecies excluded) could be distinguished. Nine indistinguishable species lacked reference sequences, whereas 12 had reference sequences but did not amplify. PCR positive controls (two per PCR

plate; *n* = 16) were exotic cichlid (*Maylandia zebra*) DNA (0.05 ng/µl), and PCR negative controls

(two per PCR plate; *n* = 16) were MGW (Fisher Scientific UK Ltd, UK). PCR products were pooled

to create sub-libraries (Fig. A2) and purified with Mag-BIND® RxnPure Plus magnetic beads

(Omega Bio-tek Inc, GA, USA), following the double size selection protocol established by

Bronner et al. (2009). Ratios of 0.9x and 0.15x magnetic beads to 100 µL of each sub-library

were used. Eluted DNA (30 µL) was stored at -20 °C until the second PCR could be performed.

The second PCR bound pre-adapters, MID tags, and Illumina adapters to the sub-libraries. PCR

products were purified with Mag-BIND® RxnPure Plus magnetic beads (Omega Bio-tek Inc, GA,

USA), following the double size selection protocol established by Bronner et al. (2009). Ratios of

0.7x and 0.15x magnetic beads to 50 µL of each sub-library were used. Eluted DNA (30 µL) was

stored at 4 °C until quantification and normalisation. The library was purified again, quantified

by qPCR using the NEBNext® Library Quant Kit for Illumina® (New England Biolabs® Inc., MA,

USA), and fragment size (330 bp) and removal of secondary product verified using an Agilent

2200 TapeStation and High Sensitivity D1000 ScreenTape (Agilent Technologies, CA, USA). The

library (220 eDNA samples, 27 field blanks, 33 filtration blanks, 17 extraction blanks, 16 PCR

negative controls, and 16 PCR positive controls) was sequenced on an Illumina MiSeq® using a

MiSeq Reagent Kit v3 (600-cycle) (Illumina, Inc, CA, USA). Raw sequence reads were

demultiplexed using a custom Python script. metaBEAT v0.97.11 (https://github.com/HullUni-

bioinformatics/metaBEAT) was used for quality trimming, merging, chimera removal, clustering,

and taxonomic assignment of sequences against our UK vertebrate reference database (Harper

et al., 2018) which contains sequences for 103 UK mammals. Taxonomic assignment used a

lowest common ancestor approach based on the top 10% BLAST matches for any query that

matched a reference sequence across more than 80% of its length at minimum identity of 98%.

2.6 Data analysis

Analyses were performed in R v.3.4.3 (R Core Team, 2017). The total unrefined read counts (i.e.

raw taxonomically assigned reads) per sample were calculated and retained for downstream

analyses. Assignments were corrected: family and genera containing a single UK species were

reassigned to that species, species were reassigned to domestic subspecies, and

misassignments were corrected, e.g. *Lynx pardinus* and *Lynx lynx*. Manual reassignment

duplicated some metaBEAT assignments thus the read count data for these assignments were

merged. Taxon-specific sequence thresholds (i.e. maximum sequence frequency of each taxon

in PCR positive controls) were used to mitigate cross-contamination and false positives (Table

A4, Fig. A3), and remnant contaminants and higher taxonomic assignments removed excluding

the following genera. *Anas* (Dabbling ducks) was retained because potential for hybridisation

reduced confidence in species-level assignments, and *Emberiza* (Buntings) and *Larus* (White-

headed gulls) were retained because reference sequences were missing for several common

species. Dataset refinement is fully described in Appendix A. Taxonomic assignments remaining

in the refined dataset were predominantly of species resolution and considered true positives.

We split the refined dataset by Experiment 1 (artificial waterbodies) and Experiment 2 (natural

ponds). Proportional read counts for each species were calculated from the total unrefined

read counts per sample. Our proportional read count data were not normally distributed

(Shapiro–Wilk normality test: $W = 0.915$, $P < 0.001$), thus we used a Mann-Whitney U test to

compare the median proportional read count of stratified and directed samples across species.

We employed binomial Generalized Linear Mixed-effects Models (GLMMs) with the logit

link function using the package glmmTMB (development version; Brooks et al., 2017) for the

following tests. First, we compared the eDNA signals from stratified and directed samples for

each mammal species using a hierarchical model including sample type nested within species

(fixed) and wildlife park (random) as effects. We tested the influence of species lifestyle on

mammal eDNA signals using a model with species lifestyle (fixed) and species nested within

wildlife park (random) as effects. Using directed samples, we tested the influence of behaviour

on mammal eDNA signals using two hierarchical models, including species nested within

wildlife park (random) and specific (e.g. swimming, drinking) or generic (i.e. water contact

versus no water contact) behaviour(s) respectively (fixed) as effects. We assessed model fit

using diagnostic plots and performed validation checks to ensure model assumptions were met

and overdispersion was absent (Zuur, Ieno, Walker, Saveliev, & Smith, 2009).

For Experiment 2, we qualitatively compared mammal presence-absence records

generated by eDNA metabarcoding, camera trapping, and field signs. TLNR ponds were

sampled every 24 hrs for 5 days, thus proportional read counts were averaged across days for

comparison to BE and TM ponds (sampled once each). We qualitatively compared the

distribution and persistence of eDNA signals between semi-aquatic and terrestrial mammals

using tile plots and heat maps of the unaveraged proportional read counts for identified species

at TLNR over the 5-day period. All figures were produced using the package ggplot2 v3.0.0

(Wickham, 2016).

3. Results

3.1 eDNA metabarcoding

The sequencing run generated 47,713,656 raw sequence reads, of which 37,590,828 remained following trimming, merging, and length filter application. After removal of chimeras and redundancy via clustering, the library contained 21,127,061 sequences (average read count of 64,215 per sample including controls), of which 16,787,750 (79.46%) were assigned a taxonomic rank. Contamination (Fig. A4) was observed in the field blanks (badger, beaver, lynx, pine marten, red squirrel, and water vole) as well as in the filtration and extraction blanks (human [*Homo sapiens*] and cichlid). PCR negative controls were contaminated to different extents with human, cichlid, beaver, and pine marten as well as non-focal species. After threshold application, contaminants remaining in eDNA samples included Gentoo penguin (*Pygoscelis papua*), reindeer (*Rangifer tarandus*), cichlid, and human. The refined dataset contained 59 vertebrate species, including six amphibians, 10 fish, 19 birds, and 24 mammals (Table A5).

3.2 Experiment 1: eDNA detection and signal strength in artificial systems

All nine focal species were detected in captivity, of which seven were detected in all water

samples taken from their respective enclosures. HWP red deer were not detected in 2 of 5

stratified samples, and WT hedgehog was not detected in 1 of 2 drinking bowl samples (Fig. 1).

'Other' samples (neither directed nor stratified) were excluded from further comparisons, thus

hedgehog, red squirrel, and water vole were omitted in downstream analyses. Across species,

stratified samples (0.406) had a higher median proportional read count than directed samples

(0.373), but this difference was not significant (Mann-Whitney U test: U = 1181.5, P = 0.829).

Proportional read counts for directed and stratified samples did not significantly differ (χ^2_6 =

0.364, P = 0.999) within species either (Fig. 2a; GLMM: θ = 0.168, χ^2_{53} = 8.915, P = 1.000,

pseudo-R^2 = 39.21%). Otter proportional read counts were lower than other species, but not

significantly so. Similarly, species lifestyle (semi-aquatic, ground-dwelling, arboreal) did not

influence (χ^2_2 = 0.655, P = 0.721) proportional read counts (Fig. 2b; GLMM: θ = 0.213, χ^2_{61} =

13.002, P = 1.000, pseudo-R^2 = 11.85%). Proportional read counts did not differ (χ^2_{11} = 1.369, P

= 0.999) according to specific behaviours exhibited by species (Fig. 3a; GLMM: θ = 0.355, χ^2_{31} =

11.013, P = 0.999, pseudo-R^2 = 9.17%). Likewise, generic behaviour (i.e. water contact versus no

water contact) did not influence (χ^2_{11} = 0.002, P = 0.964) proportional read counts (Fig. 3b;

GLMM: θ = 0.217, χ^2_{41} = 8.897, P = 1.000, pseudo-R^2 = 8.50%).

3.3 Experiment 2: eDNA detection and signal strength in natural systems

At natural ponds, eDNA metabarcoding, camera trapping, and field signs all detected beaver,

red deer, and roe deer. Camera traps (Fig. 4) and field signs recorded red fox and badger when

eDNA metabarcoding did not (Fig. 5). However, eDNA metabarcoding revealed small mammals

missed by cameras and field signs, including water vole, water shrew (*Neomys fodiens*), bank

vole (*Myodes glareolus*), common shrew (*Sorex araneus*), brown rat (*Rattus norvegicus*), rabbit,

grey squirrel, and common pipistrelle (*Pipistrellus pipistrellus*). We observed mice or vole

footprints at BE Pond 1, but could not ascertain species. Fig. 5 summarises mammals recorded

by different methods at each site with reference to cumulative survey data. Notably, only

beaver was found at the same ponds by all methods. Although methods shared species at site

level, species were not always detected at the same pond. Detection rates for species captured

by at least one survey method are summarised in Table A6.

Sampling of natural ponds revealed spatial patterns in eDNA detection and signal

strength. eDNA from non-domestic terrestrial mammals (i.e. mammals excluding dog [*Canis*

lupus familiaris], pig [*Sus scrofa domesticus*], sheep [*Ovis aries*] and cow [*Bos taurus*]) was

unevenly dispersed compared with semi-aquatic mammals (Fig. A5). Semi-aquatic beaver and

water vole were detected in at least 90% and 60% respectively of water samples (*n* = 10)

collected from single ponds, albeit water shrew was only detected in 10% of samples. Non-

domestic terrestrial mammals were routinely detected in <20% of water samples collected from

a pond and left relatively weak eDNA signals. Overall, beaver was the most consistently

detected mammal with the highest proportional read counts. However, the strongest and most

evenly distributed signals belonged to amphibians, particularly common frog (*Rana temporaria*)

and great crested newt (*Triturus cristatus*) (Fig. A5).

TLNR samples collected over a 5-day period (D01-05) revealed that mammal detection

heavily depends on the spatial and temporal resolution of eDNA metabarcoding surveys (Fig.

A6). Mammal eDNA signals in pond water were ephemeral, often disappearing within 24-48 hrs

of initial detection, as opposed to amphibians that were detected for multiple days and whose

eDNA signal increased in strength. The majority of semi-aquatic or terrestrial mammals were

only detected in a single sample on each day.

4. Discussion

We have demonstrated the potential of eDNA metabarcoding for monitoring conservation and

management priority mammals, but species detection rates are variable. Our experiments have

validated this molecular approach and provided new insights that will inform the development

and application of mammal eDNA metabarcoding. Sampling strategy, mammal lifestyle, and

mammal behaviour did not influence eDNA detection and signal strength in captivity, but all

played vital roles in natural ponds. Although semi-aquatic and terrestrial mammals were

detected from pond water, their eDNA signals were temporary and weak in comparison to

aquatic amphibians and fishes. Nonetheless, this suggests that eDNA is representative of

contemporary and local mammal diversity.

4.1 Influence of sampling strategy and mammal behaviour on eDNA detection

In Experiment 1, all nine focal species were detected in captivity, and seven were detected in all

water samples taken from their respective enclosures. This demonstrates that our method can

successfully detect a variety of mammals from pond and drinking water. Surprisingly, we found

that neither sampling strategy nor mammal lifestyle nor mammal behaviour influenced eDNA

detectability and signal strength in captivity. This included behaviours associated with eDNA

deposition, e.g. swimming, drinking, urination, and defecation (Rodgers & Mock, 2015; Ushio et

al., 2017; Williams et al., 2018). Enclosures were permanently occupied and artificial

waterbodies likely saturated with eDNA, which possibly masked behavioural signals. Modest

replication may have limited experimental power, preventing patterns being detected

statistically. Nonetheless, our results show that mammal contact with water enables eDNA

deposition and detection.

Unsurprisingly, given the nature of wild mammal interactions with natural systems

versus those in captivity, Experiment 2 results highlight the challenges of mammal eDNA

detection. We recorded 17 mammals using three monitoring tools, comparable to the 17

mammals expected from cumulative survey data despite discordance. Field signs and camera

trapping detected red fox and badger where eDNA metabarcoding did not, but eDNA

metabarcoding identified water vole and other small mammals missed on camera or with

ambiguous field signs, i.e. mice, voles, shrews. Importantly, camera trap deployment period,

height, and positioning may have influenced small mammal detection by this method

(Caravaggi et al., 2018). Ishige et al. (2017) achieved comparable mammal detection at salt licks

with eDNA metabarcoding and camera trapping, but species presence was inconsistent

between salt licks surveyed. Using multi-species occupancy modelling for three mammal

species, Sales et al. (2019) observed water-based eDNA metabarcoding provided comparable

detection probabilities to conventional survey methods and actually outperformed camera

trapping. Similarly, Leempoel et al. (2019) found soil-based eDNA metabarcoding identified the

same mammals as camera trapping as well as small mammals rarely seen on camera, albeit the

methods differed between sites. Our own results echo all three studies, where despite some

inconsistencies, eDNA metabarcoding enhanced species inventories and identified smaller,

cryptic taxa.

Notably, no survey method captured semi-aquatic otter despite presence at study sites

and successful detection in eDNA metabarcoding studies of UK ponds (Harper et al., 2019),

lakes (Hänfling et al., 2017), and rivers/streams (Sales et al., 2019). Captive otter also had a

weaker eDNA signal than other semi-aquatic mammals studied here. Lower eDNA detection

rates for otter, badger, and red fox may stem from species' ecologies (Sales et al., 2019). These

mammals are wide-ranging (Gaughran et al., 2018; Thomsen et al., 2012) and may not readily

release DNA in water. Otters often spraint on grass or rock substrata outside water and use

latrines associated with caves and dens (Ruiz-Olmo & Gosálbez, 1997). As terrestrial mammals,

red fox and badger must drink from or enter ponds for eDNA deposition to occur (Rodgers &

Mock, 2015; Ushio et al., 2017; Williams et al., 2018). Otter, badger, and red fox detection may

require greater spatiotemporal resolution of eDNA sampling. This is reinforced by other eDNA

metabarcoding studies where mammal detection was highly variable across sites surveyed

(Ishige et al., 2017; Klymus et al., 2017; Leempoel et al., 2019; Sales et al., 2019; Ushio et al.,

2017). False negatives may instead be symptomatic of metabarcoding bias, but this is unlikely in

our study (section 4.2).

eDNA from other semi-aquatic mammals was evenly distributed, being found in most or

all samples collected on fine spatial scales within natural ponds, whereas terrestrial mammal

eDNA was highly localised and detected in few (<20%) samples. Mammal eDNA signals varied

temporally, being detectable for two consecutive days maximum. Depending on the species,

mammal eDNA may be spatially and temporally clumped in lentic ecosystems due to the nature

and frequency of water contact. Unless non-domestic mammals exhibit behaviours involving

prolonged water contact (e.g. swimming, wallowing), they may only be detected at drinking

sites (Klymus et al., 2017; Ushio et al., 2017; Williams et al., 2018). Conversely, domestic

mammals may have elevated detection rates in ponds due to high occurrence of these

waterbodies in agricultural landscapes as well as eDNA transport by rainfall and run-off (Staley

et al., 2018). eDNA detection and persistence are further influenced by group size, where eDNA

from multiple individuals endures for longer periods in water than eDNA from single individuals

(Williams et al., 2018). Detailed investigations incorporating biotic (e.g. population size, body

mass, behaviour) and abiotic (e.g. temperature, pH, rainfall) factors are needed to understand

the longevity of mammal eDNA signals in aquatic ecosystems (Rodgers & Mock, 2015; Sales et

al., 2019; Williams et al., 2018).

Our two experiments have shown that sampling strategy influences mammal eDNA

detection. Mammal eDNA was evenly distributed in closed, artificial waterbodies, but locally

distributed in open, natural ponds. Captive mammal enclosures contained one species

(excluding HWP red deer) and a drinking container(s) and/or small waterbody (range 0.01-162

m^2, mean 27.4 m^2). Some enclosures housed more individuals of a species than others, thereby

increasing eDNA deposition and detection probability (Williams et al., 2018). Wild mammals

have an array of freshwater habitats at their disposal and can hold vast territories. Therefore,

rates of pond visitation and eDNA deposition are more irregular (Klymus et al., 2017; Ushio et

al., 2017), possibly leading to between-sample variation (Williams et al., 2018).

4.2 Accounting for false positives and false negatives in metabarcoding

eDNA metabarcoding has potential for inclusion in mammal monitoring schemes (section 4.3),

but like existing monitoring tools, may produce false negatives or false positives. Our process

controls identified low-level contamination at all stages of metabarcoding, but primarily during

sampling or PCR (Appendix A). We applied taxon-specific sequence thresholds to our data to

mitigate false positives as in Harper et al. (2019). Remnant contaminants were cichlid

(laboratory), Gentoo penguin (environment), reindeer (environment), and human

(environment/laboratory). Gentoo penguin is housed at EZ and was identified from EZ beaver

enclosure water. The WT red squirrel and reindeer enclosures are in close proximity. DNA

transport by wildlife (e.g. waterfowl [Hänfling et al., 2016]) and park staff/visitors may explain

this environmental contamination. Human DNA was present across process controls

corresponding to artificial and natural waterbodies. Human DNA may be amplified and

sequenced instead of focal species, potentially resulting in false negative detections for rare

and/or less abundant species. Human DNA blocking primers can prevent this bias, but may

impair PCR amplification efficiency (Klymus et al., 2017; Ushio et al., 2017; Valentini et al.,

2016). Sequence thresholds are one method of accounting for contamination in metabarcoding

datasets, but this is a topic that warrants deeper investigation aimed at researching and

refining standardised methods for false positive identification and mitigation, e.g. the R

package microDecon (McKnight et al., 2019).

In our study, eDNA metabarcoding produced false negatives for otter, badger, and red

fox at natural ponds. We selected a 12S metabarcode designed to amplify vertebrate DNA (Riaz

et al., 2011). One of four fox reference sequences (NCBI Accession: KF387633.1) possessed one

mismatch to the forward primer, and one of three otter reference sequences (NCBI Accession:

EF672696.1) possessed one mismatch to the reverse primer. These mismatches did not occur

within the first or last four bases of either primer sequence, and there were no primer

mismatches with the badger reference sequences (Harper et al., 2018). Therefore, amplification

bias was not responsible for these false negatives. DNA from aquatic and more abundant

species may have overwhelmed otter, badger, and red fox DNA during amplification and

sequencing, i.e. species-masking (Kelly, Port, Yamahara, & Crowder, 2014; Klymus et al., 2017).

Metabarcoding primers targeting mammals (Ushio et al., 2017) or multi-marker (e.g. 12S, 16S,

COI) investigations (Evans et al., 2017; Hänfling et al., 2016; Kelly et al., 2014; Klymus et al.,

2017) may improve mammal detection in systems with competition from non-target aquatic

species. Similarly, more biological and technical replication may improve species detection

probabilities (Evans et al., 2017; Lawson Handley et al., 2019; Sales et al., 2019; Valentini et al.,

2016). Importantly, otter also had lower qPCR detection than amphibians and fish (Thomsen et

al., 2012). A metabarcoding and qPCR comparison (e.g. Harper et al., 2018; Lacoursière-Roussel,

Dubois, Normandeau, & Bernatchez, 2016) would confirm whether poor amplification

efficiency for otter arises from technical bias or species ecology, and whether eDNA

metabarcoding can reliably monitor otter alongside the wider mammalian community.

4.3 Scope of eDNA metabarcoding for mammal monitoring

4.3 Scope of eDNA metabarcoding for mammal monitoring

Mammal population assessments are hindered by lack of data and systematic monitoring for

many species (Mathews et al., 2018). Distribution and occupancy data are poor for most

species, with ongoing survey effort biased toward rare species. Surveys heavily rely on citizen

science and casual records (Massimino et al., 2018). Tools that provide standardised, systematic

monitoring of mammal populations are needed (Mathews et al., 2018). Despite issues inherent

to metabarcoding for biodiversity monitoring (Deiner et al., 2017), this tool has enormous

potential to enhance mammal monitoring, conservation, and management. eDNA

metabarcoding generates distribution data for multiple species, whether rare, invasive, or

abundant, and could track conflicting species simultaneously, e.g. water vole, American mink,

and otter (Bonesi & Macdonald, 2004) or red squirrel, grey squirrel, and pine marten (Sheehy,

Sutherland, O'Reilly, & Lambin, 2018).

eDNA metabarcoding can rapidly survey multitudes of aquatic sites at landscape-scale

where camera traps might be resource-intensive, cost-inefficient, and susceptible to

theft/damage (Ushio et al., 2017). Field signs require volunteer time and skill (Sadlier et al.,

2004) to be employed at comparable spatial scales to eDNA metabarcoding which could

provide accurate data for species misidentified from field signs, e.g. mice and voles, otter and

mink (Franklin et al., 2019; Harris & Yalden, 2004). However, camera traps and field signs both

recorded species that eDNA metabarcoding missed. Therefore, eDNA metabarcoding is

complementary and should be incorporated into, not replace, existing monitoring schemes

(Leempoel et al., 2019; Sales et al., 2019). This tool could be most effective in mammal

monitoring if deployed at the edges of known species distributions, in areas where species

presence is unknown, and in areas with isolated species records (Mathews et al., 2018).

4.4 Recommendations for mammal survey using eDNA metabarcoding

Water-based eDNA metabarcoding shows great promise for mammal monitoring encompassing

conservation and management priority species (Sales et al., 2019). However, there are factors

to be considered when designing and conducting mammal eDNA surveys that may not be

problematic for surveys of fishes or amphibians. Mammal eDNA detection probabilities from

natural ponds will likely be high when areas with dense populations are studied, but rigorous

sampling strategies will be required to track mammals in areas sparsely populated by

individuals. Multiple ponds must be sampled repeatedly, and samples taken at multiple

locations within ponds without pooling to enable site occupancy inferences. Importantly, we

sampled natural ponds in spring but sampling in other seasons may produce different results,

reflective of species' ecologies (Lawson Handley et al., 2019). To account for differential

mammal visitation rates and maximise eDNA detection probabilities, we recommend that

researchers and practitioners using eDNA metabarcoding for mammal monitoring channel their

efforts into extensive sampling of numerous waterbodies in a given area over prolonged

timescales. Water-based eDNA appears to be indicative of contemporary mammal presence,

with most mammal eDNA signals lost within 1-2 days. Therefore, eDNA metabarcoding could

provide valuable mammalian community "snapshots" that may not be obtained with other

survey methods (Ushio et al., 2017). Different sample types (e.g. water, soil, snow, salt licks,

feeding traces, faeces, hair, and blood meals) may also offer new insights to mammal

biodiversity (Franklin et al., 2019; Ishige et al., 2017; Kinoshita et al., 2019; Leempoel et al.,

2019; Sales et al., 2019; Tessler et al., 2018; Ushio et al., 2017).

Data accessibility

Raw sequence reads have been archived on the NCBI Sequence Read Archive (Study:

SRP164740; BioProject: PRJNA495011; BioSamples: SAMN10195928 - SAMN10196255; SRA

accessions: SRR7986451 - SRR7986778). Jupyter notebooks, R scripts and corresponding data

are archived online (https://doi.org/10.5281/zenodo.2561415).

Acknowledgements

The authors declare that they have no conflicts of interest in presenting this work for

publication. This work was funded by the University of Hull, and D.S.R was supported by the

Natural Environment Research Council award number NE/R016429/1 as part of the UK-SCAPE

programme delivering National Capability. We would like to thank Gill Murray-Dickson

(National Museums Scotland) and Helen Senn (RZSS Edinburgh Zoo) for feedback on the study

design, and Richard Griffiths (DICE, University of Kent) for support with filtration of water

samples from Wildwood Trust. We are grateful to staff at Wildwood Trust and RZSS Highland

Wildlife Park for assisting with eDNA sampling from animal enclosures. We thank Richard

Hampshire and volunteers at Tophill Low Nature Reserve for assisting with camera trap

deployment. Tim Kohler (Natural England) and Paul Ramsay kindly gave permission to sample at

Thorne Moors and the Bamff Estate respectively.

Author contributions

L.R.H, B.H, and L.L.H conceived and designed the study. A.I.C and M.G coordinated sampling at

Wildwood Trust and RZSS Highland Wildlife Park respectively. L.R.H, C.D.M, C.J.M, and T.L

collected and filtered water samples. A.L, T.L, and T.B helped select natural ponds to be

surveyed using eDNA, camera trapping, and field signs, and provided camera traps for the

study. L.R.H, A.L, and T.L deployed camera traps, which were then collected and footage

analysed by L.R.H. L.R.H processed samples in the laboratory with advice from C.D.M and A.M.

D.S.R sequenced the final library. L.R.H completed bioinformatic processing of samples, and

subsequent data analysis. L.R.H wrote the manuscript, which all authors contributed critically to

drafts of and gave final approval for publication.

References

Alberdi, A., Aizpurua, O., Gilbert, M. T. P., & Bohmann, K. (2018) Scrutinizing key steps for reliable metabarcoding of environmental samples. *Methods in Ecology and Evolution*, *9*, 134–147.

Bálint, M., Nowak, C., Márton, O., Pauls, S. U., Wittwer, C., Aramayo, J. L., Schulze, A., Chambert, T., Cocchiararo, B., & Jansen, M. (2018). Accuracy, limitations and cost efficiency of eDNA-based community survey in tropical frogs. *Molecular Ecology Resources*, *18*, 1415–1426. https://doi.org/10.1111/1755-0998.12934

Bland, L. M., Collen, B., Orme, C. D. L., & Bielby, J. (2015). Predicting the conservation status of data-deficient species: Predicting Extinction Risk. *Conservation Biology*, *29*, 250–259. https://doi.org/10.1111/cobi12372

Bonesi, L., & Macdonald, D. W. (2004). Differential habitat use promotes sustainable coexistence between the specialist otter and the generalist mink. *Oikos* , *106*, 509–519. https://doi.org/10.1111/j.0030-1299.2004.13034.x

Bronner, I. F., Quail, M. A., Turner, D. J. & Swerdlow, H. (2009) Improved Protocols for Illumina Sequencing. *Current Protocols in Human Genetics*, *18*, 18.2.1–18.2.42.

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Machler, M., & Bolker, B. M. (2017). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal*, *9*, 378–400. https://doi.org/10.32614/RJ-2017-066

Burton, A. C., Neilson, E., Moreira, D., Ladle, A., Steenweg, R., Fisher, J. T., Bayne, E., & Boutin, S. (2015). REVIEW: Wildlife camera trapping: a review and recommendations for linking surveys to ecological processes. *Journal of Applied Ecology*, *52*, 675–685. https://doi.org/10.1111/1365-2664.12432

Caravaggi, A., Gatta, M., Vallely, M. -C., Hogg, K., Freeman, M., Fadaei, E., Dick, J. T. A., Montgomery, W. I., Reid, N., & Tosh, D. G. (2018). Seasonal and predator-prey effects on circadian activity of free-ranging mammals revealed by camera traps. *PeerJ*, *6*, e5827. https://doi.org/10.7717/peerj.5827

Deiner, K., Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., Creer, S., Bista, I., Lodge, D. M., de Vere, N., Pfrender, M. E., & Bernatchez, L. (2017). Environmental DNA metabarcoding: transforming how we survey animal and plant communities. *Molecular Ecology*, *26*, 5872–5895. https://doi.org/10.1111/mec.14350

Evans, N. T., Li, Y., Renshaw, M. A., Olds, B. P., Deiner, K., Turner, C. R., Jerde, C. L., Lodge, D. M., Lamberti, G. A., & Pfrender, M. E (2017). Fish community assessment with eDNA metabarcoding: effects of sampling design and bioinformatic filtering. *Canadian Journal of Fisheries and Aquatic Sciences*, *74*, 1362–1374. https://doi.org/10.1139/cjfas-2016-0306

Franklin, T. W., McKelvey, K. S., Golding, J. D., Mason, D. H., Dysthe, J. C., Pilgrim, K. L., Squires,
J. R., Aubry, K. B., Long, R. A., Greaves, S. E., Raley, C. M., Jackson, S., MacKay, P., Lisbon, J.,
Sauder, J. D., Pruss, M. T., Heffington, D., & Schwartz, M. K (2019). Using environmental DNA
methods to improve winter surveys for rare carnivores: DNA from snow and improved
noninvasive techniques. *Biological Conservation*, *229*, 50–58.
https://doi.org/10.1016/j.biocon.2018.11.006

Gaughran, A., Kelly, D. J., MacWhite, T., Mullen, E., Maher, P., Good, M., & Marples, N. M.
(2018). Super-ranging. A new ranging strategy in European badgers. *PLoS ONE*, *13*,
e0191818. https://doi.org/10.1371/journal.pone.0191818

Hänfling, B., Lawson Handley, L., Read, D. S., Hahn, C., Li, J., Nichols, P., Blackman, R. C., Oliver,
A., & Winfield, I. J. (2016). Environmental DNA metabarcoding of lake fish communities
reflects long-term data from established survey methods. *Molecular Ecology*, *25*, 3101–3119.
https://doi.org/10.1111/mec.13660

Hänfling, B., Lawson Handley, L., Lunt, D., Shum, P., Winfield, I., & Read, D. S. (2017). *A Review*
of Recent Advances in Genetic Methods to Identify Improvements in CAMERAS Partners
Monitoring Activities. University of Hull, Hull, UK.

Harper, L. R., Lawson Handley, L., Hahn, C., Boonham, N., Rees, H. C., Gough, K. C., Lewis, E.,
Adams, I. P., Brotherton, P., Phillips, S. & Hänfling, B. (2018). Needle in a haystack? A
comparison of eDNA metabarcoding and targeted qPCR for detection of the great crested
newt (*Triturus cristatus*). *Ecology and Evolution*, *8*, 6330–6441.
https://doi.org/10.1002/ece3.4013

Harper, L. R., Lawson Handley, L., Hahn, C., Boonham, N., Rees, H. C., Lewis, E., Adams, I. P.,
Brotherton, P., Phillips, S., & Hänfling, B. (2019) Testing ecological hypotheses at the
pondscape with environmental DNA metabarcoding: a case study on a threatened
amphibian. *bioRxiv*, 278309. https://doi.org/10.1101/278309

Harris, S., & Yalden, D. W. (2004). An integrated monitoring programme for terrestrial mammals
in Britain. *Mammal Review*, *34*, 157–167. https://doi.org/10.1046/j.0305-1838.2003.00030.x

Ishige, T., Miya, M., Ushio, M., Sado, T., Ushioda, M., Maebashi, K., Yonechi, R., Lagan, P., &
Matsubayashi, H. (2017). Tropical-forest mammals as detected by environmental DNA at
natural saltlicks in Borneo. *Biological Conservation*, *210*, 281–285.
https://doi.org/10.1016/j.biocon.2017.04.023

Joint Nature Conservation Committee (2018) UK BAP priority terrestrial mammal species.
Accessed on 1st December 2018. http://jncc.defra.gov.uk/page-5170

Kelly, R. P., Port, J. A., Yamahara, K. M., & Crowder, L. B. (2014). Using environmental DNA to
census marine fishes in a large mesocosm. *PLoS ONE*, *9*, e86175.
https://doi.org/10.1371/journal.pone.0086175

Kinoshita, G., Yonezawa, S., Murakami, S., & Isagi, Y. (2019) Environmental DNA Collected from
Snow Tracks is Useful for Identification of Mammalian Species. *Zoological Science*, *36*, 198–
207. http://dx.doi.org/10.2108/zs180172

Kitson, J. J. N., Hahn, C., Sands, R. J., Straw, N. A., Evans, D. M., & Lunt, D. H. (2019). Detecting
host-parasitoid interactions in an invasive Lepidopteran using nested tagging DNA-
metabarcoding. *Molecular Ecology*, *28*, 471–483. https://doi.org/10.1111/mec.14518

Klymus, K. E., Richter, C. A., Thompson, N., & Hinck, J. E. (2017). Metabarcoding of
Environmental DNA Samples to Explore the Use of Uranium Mine Containment Ponds as a
Water Source for Wildlife. *Diversity*, *9*, 54. https://doi.org/10.3390/d9040054

Lacoursière-Roussel, A., Dubois, Y., Normandeau, E., & Bernatchez, L. (2016). Improving
herpetological surveys in eastern North America using the environmental DNA method.
Genome, *59*, 991–1007. https://doi.org/10.1139/gen-2015-0218

Lawson Handley, L., Read, D.S., Winfield, I.J., Kimbell, H., Johnson, H., Li, J., Hahn, C., Blackman,
R., Wilcox, R., Donnelly, R., Szitenberg, A., & Hänfling, B. (2019) Temporal and spatial
variation in distribution of fish environmental DNA in England's largest lake. *Environmental*
DNA, **12**, e0176343. https://doi.org/10.1002/edn3.5

Leempoel, K., Hebert, T., & Hadly, E. A. (2019) A comparison of eDNA to camera trapping for
assessment of terrestrial mammal diversity. *bioRxiv*, 634022.
https://doi.org/10.1101/634022

Lugg, W. H., Griffiths, J., van Rooyen, A. R., Weeks, A. R., & Tingley, R. (2017). Optimal survey
designs for environmental DNA sampling. *Methods in Ecology and Evolution*, *9*, 1049–1059.
https://doi.org/10.1111/2041-210X.12951

Massimino, D., Harris, S. J., & Gillings, S. (2018). Evaluating spatiotemporal trends in terrestrial
mammal abundance using data collected during bird surveys. *Biological Conservation*, *226*,
153–167. https://doi.org/10.1016/j.biocon.2018.07.026

Mathews, F., Kubasiewicz, L. M., Gurnell, J., Harrower, C. A., McDonald, R. A., & Shore, R. F.
(2018) *A Review of the Population and Conservation Status of British Mammals: Technical*
Summary. Natural England, Peterborough.

McKnight, D. T., Huerlimann, R., Bower, D. S., Schwarzkopf, L., Alford, R. A., & Zenger, K. R.
(2019) microDecon: A highly accurate read-subtraction tool for the post-sequencing removal
of contamination in metabarcoding studies. *Environmental DNA*, *1*, 14–25.
https://doi.org/10.1002/edn3.11

Riaz, T., Shehzad, W., Viari, A., Pompanon, F., Taberlet, P., & Coissac, E. (2011). ecoPrimers:
inference of new DNA barcode markers from whole genome sequence analysis. *Nucleic Acids*
Research, *39*, e145. https://doi.org/10.1093/nar/gkr732

Rodgers, T. W., & Mock, K. E. (2015). Drinking water as a source of environmental DNA for the
detection of terrestrial wildlife species. *Conservation Genetics Resources*, *7*, 693–696.
https://doi.org/10.1007/s12686-015-0478-7

Ruiz-Olmo, J., & Gosálbez, J. (1997). Observations on the sprainting behaviour of the otter *Lutra*
lutra in the NE Spain. *Acta Theriologica*, *42*, 259–270.
Sadlier, L. M. J., Webbon, C. C., Baker, P. J., & Harris, S. (2004). Methods of monitoring red foxes
Vulpes vulpes and badgers Meles meles: are field signs the answer? *Mammal Review*, *34*,
75–98. https://doi.org/10.1046/j.0305-1838.2003.00029.x
Sales, N. G., McKenzie, M. B., Drake, J., Harper, L. R., Browett, S. S., Coscia, I., Wangensteen, O.
S., Baillie, C., Bryce, E., Dawson, D. A., Ochu, E., Hänfling, B., Lawson Handley, L., Mariani, S.,
Lambin, X., Sutherland, C., & McDevitt, A. D. (2019) Fishing for mammals: landscape-level
monitoring of terrestrial and semi-aquatic communities using eDNA from lotic ecosystems.
bioRxiv, 629758. https://doi.org/10.1101/629758
Sellers, G. S., Di Muri, C., Gómez, A., & Hänfling, B. (2018). Mu-DNA: a modular universal DNA
extraction method adaptable for a wide range of sample types. *Metabarcoding and*
Metagenomics, *2*, e24556. https://doi.org/10.3897/mbmg.2.24556
Sheehy, E., Sutherland, C., O'Reilly, C., & Lambin, X. (2018) The enemy of my enemy is my
friend: native pine marten recovery reverses the decline of the red squirrel by suppressing
grey squirrel populations. *Proceedings of the Royal Society of London B*, *285*, 20172603.
https://doi.org/10.1098/rspb.2017.2603
Staley, Z. R., Chuong, J. D., Hill, S. J., Grabuski, J., Shokralla, S., Hajibabaei, M., & Edge, T. A.
(2018). Fecal source tracking and eDNA profiling in an urban creek following an extreme rain
event. *Scientific Reports*, *8*, 14390. https://doi.org/10.1038/s41598-018-32680-z
Tessler, M., Weiskopf, S. R., Berniker, L., Hersch, R., Mccarthy, K. P., Yu, D. W., & Siddall, M. E.
(2018). Bloodlines: mammals, leeches, and conservation in southern Asia. *Systematics and*
Biodiversity, *16*, 488–496. https://doi.org/10.1080/14772000.2018.1433729
Thomsen, P. F., Iversen, L. L., Wiuf, C., Rasmussen, M., Gilbert, M. T. P., Orlando, L., &
Willerslev, E. (2012). Monitoring endangered freshwater biodiversity using environmental
DNA. *Molecular Ecology*, *21*, 2565–2573. https://doi.org/10.1111/j.1365-294X.2011.05418.x
Thomsen, P. F., & Willerslev, E. (2015). Environmental DNA–an emerging tool in conservation
for monitoring past and present biodiversity. *Biological Conservation*, *183*, 4–18.
https://doi.org/10.1016/j.biocon.2014.11.019
Ushio, M., Fukuda, H., Inoue, T., Makoto, K., Kishida, O., Sato, K., Murata, K., Nikaido, M., Sado,
T., Sato, Y., Takeshita, M., Iwasaki, W., Yamanaka, H., Kondoh, M., & Miya, M. (2017).
Environmental DNA enables detection of terrestrial mammals from forest pond water.
Molecular Ecology Resources, *17*(6), e63–e75. https://doi.org/10.1111/1755-0998.12690

Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P. F., Bellemain, E.,
Besnard, A., Coissac, E., Boyer, F., Gaboriaud, C., Jean, P., Poulet, N., Roset, N., Copp, G. H.,
Geniez, P., Pont, D., Argillier, C., Baudoin, J. -M., Peroux, T., Crivelli, A. J., Olivier, A.,
Acqueberge, M., Le Brun, M., Møller, P. R., Willerslev, E., & Dejean, T. (2016). Next-
generation monitoring of aquatic biodiversity using environmental DNA metabarcoding.
Molecular Ecology, *25*, 929–942. https://doi.org/10.1111/mec.13428

Visconti, P., Pressey, R. L., Giorgini, D., Maiorano, L., Bakkenes, M., Boitani, L., Alkemade, R.,
Falcucci, A., Chiozza, F., & Rondinini, C. (2011). Future hotspots of terrestrial mammal loss.
Philosophical Transactions of the Royal Society B, *366*, 2693–2702.
https://doi.org/10.1098/rstb.2011.0105

Wickham, H. (2016). *ggplot2: elegant graphics for data analysis*. Springer, New York, USA.
http://ggplot2.org

Williams, K. E., Huyvaert, K. P., Vercauteren, K. C., Davis, A. J., & Piaggio, A. J. (2018). Detection
and persistence of environmental DNA from an invasive, terrestrial mammal. *Ecology and*
Evolution, *8*, 688–695. https://doi.org/10.1002/ece3.3698

Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009) *Mixed effects models*
and extensions in ecology with R. Springer, New York, USA.

Table 1. Ethogram used to catalogue mammal behaviours that occur in or near artificial

waterbodies in captive enclosures. Importantly, this ethogram was designed to catalogue

mammal behaviours potentially leading to eDNA deposition. Therefore, it may not be

comparable to ethograms typically used in study of captive animals.

Behaviour	Definition
Swimming	Mammal completely submerged in and moving through waterbody using limbs
Bodypart in water	Mammal partially submerged in waterbody, e.g. foot or tail in water
Drinking	Water taken into mouth and swallowed by mammal
Feeding	Food taken into mouth and swallowed by mammal in or near waterbody, e.g. otter and fish
Scratching	Bodypart or external object in enclosure used by mammal to relieve itch near waterbody
Urinating/scent-marking	Liquid excretion passed by mammal in or near waterbody
Pooing	Solid excretion passed by mammal in or near waterbody
Sniffing	Air visibly drawn through nose of mammal to detect a smell around waterbody, possibly involving contact with water
Standing	Mammal motionless in or near waterbody
Walking	Mammal moving around waterbody at a regular pace by lifting and setting down each foot in turn, never having both feet off the ground at once
Running	Mammal moving around waterbody at a speed faster than a walk, never having both or all the feet on the ground at the same time
Vocalising	Mammal producing sound while in or near waterbody
Grooming	Mammal cleaning fur or skin with its tongue while in or near waterbody
Resting	Mammal lying down or sitting in or near waterbody
Other	Behaviour exhibited in or near waterbody that does not conform to other categories, e.g. chasing tail
Not visible	Mammal moved to part of enclosure not visible to the observer

759   **Table 2.** Summary of focal species studied at wildlife parks and their lifestyle. The number of

760   individuals present and waterbody size in enclosures is provided.

Site	Species	Lifestyle	Enclosure	Number of individuals	Waterbody size (m^2)
Wildwood Trust	European otter (*Lutra lutra*)	Semi-aquatic	1	2	162
	European water vole (*Arvicola amphibius*)	Semi-aquatic	1	4	0.09
			2	1	0.09
	European beaver (*Castor fiber*)	Semi-aquatic	1	2	100
			2	1	100
	European hedgehog (*Erinaceus europaeus*)	Ground-dwelling	1	1	0.04
			2	2	0.04
	European badger (*Meles meles*)	Ground-dwelling	1	4	1.73
	Red deer (*Cervus elaphus*)	Ground-dwelling	1	8	100
	Eurasian lynx (*Lynx lynx*)	Ground-dwelling	1	2	2
	Red squirrel (*Sciurus vulgaris*)	Arboreal	1	2	0.01
			2	3	0.01
			3	3	0.01
			4	2	0.01
	European pine marten (*Martes martes*)	Arboreal	1	1	2
			2	1	0.375
Highland Wildlife Park	Red squirrel (*Sciurus vulgaris*)	Arboreal	NA	NA	0.25
	Eurasian lynx (*Lynx lynx*)	Ground-dwelling	1	8	2
	European beaver (*Castor fiber*)	Semi-aquatic	1	2	50
	Red deer (*Cervus elaphus*)	Ground-dwelling	1	30	NA

Figure 1. Heatmap showing proportional read counts for eDNA samples from Experiment 1. Each square represents a sample taken from a wildlife park enclosure. Directed (DIR01-DIR06) and stratified (STR01-STR06) samples were collected from artificial waterbodies. Samples were collected from drinking containers (E1, E2, E3, E4, BOWL, BUCK), water vole (QUAR1, QUAR2) and RZSS Edinburgh Zoo beaver (ZOO) enclosures, and a water bath (BATH) in RZSS Highland Wildlife Park woods. Samples containing a 0 in the corresponding cell had proportional read counts greater than two decimal places. Samples with proportional read counts of 0 are represented by empty cells.

Figure 2. Relationships predicted by the binomial GLMMs between proportional read counts and sample type nested within species **(a)** or species lifestyle **(b)** for Experiment 1. The observed data (coloured points) are displayed against the predicted relationships (black points with error bars) for each species **(a)** or species lifestyle **(b)**. Points are shaped by sample type **(a)** or wildlife park **(b)**, and coloured by species lifestyle. Error bars represent the standard error around the predicted means.

Figure 3. Boxplots showing the mean proportional read counts for specific **(a)** and generic **(b)** behaviour(s) exhibited by focal species in Experiment 1. Boxes show 25th, 50th, and 75th percentiles, and whiskers show 5th and 95th percentiles. Points are coloured by species lifestyle, and each point in **(a)** represents a directed sample sized by frequency of behaviour. The behaviour 'none' for beaver represents occurrences of beaver in water but out of view of camera traps.

Figure 4. Exemplar camera trap photos taken at natural ponds where focal species were present in Experiment 2. Red deer was recorded at Thorne Moors **(a)**, roe deer **(b)** and red fox **(c)** were recorded at Tophill Low Nature Reserve, and beaver was recorded at the Bamff Estate **(d)**.

Figure 5. Tile plot showing species presence-absence at individual pond and site-level as indicated by field signs, camera trapping, and eDNA metabarcoding in Experiment 2. Surveys were performed at sites where focal species presence was confirmed by cumulative survey data.