Environmental DNA (eDNA) metabarcoding of pond water as a tool to

survey conservation and management priority mammals

Lynsey R. Harper[1*], Lori Lawson Handley[1], Angus I. Carpenter[2], Gill Murray-Dickson[3], Cristina

Di Muri[1], Callum J. Macgregor[4], Thomas W. Logan[1], Alan Law[5], Thomas Breithaupt[1], Daniel S.

Read[6], Allan D. McDevitt[7], and Bernd Hänfling[1]

[1] Department of Biological and Marine Sciences, University of Hull, Hull, HU6 7RX, UK

[2] Wildwood Trust, Canterbury Rd, Herne Common, Herne Bay, CT6 7LQ, UK

[3] The Royal Zoological Society of Scotland, Edinburgh Zoo, 134 Corstorphine Road, Edinburgh, EH12 6TS, UK

[4] Department of Biology, University of York, Wentworth Way, York, YO10 5DD, UK

[5] Biological and Environmental Sciences, University of Stirling, Stirling, FK9 4LA, UK

[6] Centre for Ecology & Hydrology (CEH), Benson Lane, Crowmarsh Gifford, Wallingford, Oxfordshire, OX10 8BB, UK

[7] Ecosystems and Environment Research Centre, School of Environment and Life Sciences, University of Salford,

Salford, M5 4WT, UK

*Corresponding author: lynsey.harper2@gmail.com

Lynsey Harper, Department of Biological and Marine Sciences, University of Hull, Hull, HU6 7RX,

UK

Abstract

1. Environmental DNA (eDNA) metabarcoding is largely used to survey aquatic communities, but can also provide data on terrestrial taxa utilising aquatic habitats. However, the entry, dispersal, and detection of terrestrial species' DNA within waterbodies is understudied.

2. We evaluated eDNA metabarcoding of pond water for monitoring semi-aquatic, ground-dwelling, and arboreal mammals, and examined spatiotemporal variation in mammal eDNA signals using experiments in captive and wild conditions.

3. We selected nine focal species of conservation and management concern: European water vole, European otter, Eurasian beaver, European hedgehog, European badger, red deer, Eurasian lynx, red squirrel, and European pine marten. We hypothesised that eDNA signals (i.e. proportional read counts) would be stronger for semi-aquatic than terrestrial species, and at sites where mammals exhibited behaviours (e.g. swimming, urination). We tested this by sampling waterbodies in enclosures of captive focal species at specific sites where behaviours had been observed ('directed' sampling) and at equidistant intervals along the shoreline ('stratified' sampling). We then surveyed natural ponds (N = 6) where focal species were present using stratified water sampling, camera traps, and field signs. eDNA samples were metabarcoded using vertebrate-specific primers.

4. All focal species were detected in captivity. eDNA signal strength did not differ between directed and stratified samples across or within species, between species lifestyles (i.e. semi-aquatic, ground-dwelling, arboreal), or according to behaviours. Therefore, eDNA was evenly distributed within artificial waterbodies. Conversely, eDNA was unevenly distributed

in natural ponds. eDNA metabarcoding, camera trapping, and field signs detected beaver,

red deer, and roe deer. Badger and red fox were recorded with cameras and field signs, but

not eDNA metabarcoding. However, eDNA metabarcoding detected small mammals missed

by cameras and field signs, e.g. water vole. Terrestrial mammal eDNA signals were weaker

and detected in fewer samples than semi-aquatic mammal eDNA signals.

5. eDNA metabarcoding has potential for inclusion in mammal monitoring schemes by

enabling large-scale, multi-species distribution assessment for priority and difficult to survey

species, and could provide early indication of range expansions or contractions. However,

eDNA surveys need high spatiotemporal resolution and metabarcoding biases require

further investigation before this tool is routinely implemented.

1. Introduction

Mammals are among the most threatened vertebrate groups globally, with at least 25% of species at risk of extinction due to harvesting, habitat degradation/loss, non-native species or perception as pests (Visconti et al., 2011). Lack of data prevents robust estimation of mammalian range expansions or declines and population trends (Bland, Collen, Orme, & Bielby, 2015). Most species lack long-term, systematic monitoring, with survey efforts biased towards rare species (Massimino, Harris, & Gillings, 2018). Consequently, there is a need for more effective and evidence-based strategies for mammal conservation and management (Battersby & Tracking Mammals Partnership, 2005; Mathews et al., 2018).

Many mammals are nocturnal and elusive thus monitoring often requires non-invasive, observational methods, such as camera traps and field signs (Caravaggi et al., 2018; Harris & Yalden, 2004; Sadlier, Webbon, Baker, & Harris, 2004). Camera trapping is cost-efficient, standardised, reproducible, and produces data suited to site occupancy modelling. However, camera traps can only survey a fraction of large, heterogeneous landscapes, trap placement can substantially influence species detection probabilities and community insights, and traps often miss small species (Burton et al., 2015; Caravaggi et al., 2018; Ishige et al., 2017; Stat et al., 2019). Field sign surveys are inexpensive, but resource-intensive (i.e. time, volunteers) for broad geographic coverage (Sadlier et al., 2004). Some species also have similar footprints and scat, increasing the potential for misidentification (Franklin et al., 2019; Harris & Yalden, 2004). Mammal survey methods are usually species-specific, and multiple methods are necessary for large-scale, multi-species monitoring schemes (Massimino et al., 2018).

Environmental DNA (eDNA) analysis is a recognised tool for rapid, non-invasive, cost-

efficient biodiversity assessment across a wide range of aquatic and terrestrial habitats (Deiner

et al., 2017). Organisms transfer genetic material to their environment via secretions,

excretions, gametes, blood, or decomposition, which can be isolated from environmental

samples (Thomsen & Willerslev, 2015). Studies using eDNA analysis for semi-aquatic and

terrestrial mammal detection have largely employed PCR or quantitative PCR (qPCR) to target

single species (Aylward, Johnson, Sullivan, Perry, & Louis, 2018; Franklin et al., 2019; Lugg,

Griffiths, van Rooyen, Weeks, & Tingley, 2017; Padgett-Stewart et al., 2016; Rodgers & Mock,

2015; Thomsen et al., 2012; Williams, Huyvaert, Vercauteren, Davis, & Piaggio, 2018).

Alternatively, eDNA metabarcoding can screen entire communities using PCR combined with

high-throughput sequencing (Deiner et al., 2017; Thomsen & Willerslev, 2015), but such

assessments of mammal communities are less common (Klymus, Richter, Thompson, & Hinck,

2017; Ushio et al., 2017). Mammal assemblages have been obtained by metabarcoding

invertebrate blood meals (e.g. Tessler et al., 2018) and salt licks (Ishige et al., 2017) in tropical

habitats, but samples from the physical environment have tremendous potential to reveal

mammal biodiversity over broad spatiotemporal scales (Ushio et al., 2017).

In aquatic ecosystems, eDNA metabarcoding has predominantly been applied to

characterise fish (e.g. Evans et al., 2017; Hänfling et al., 2016; Lawson Handley et al., 2018;

Valentini et al., 2016) and amphibian (e.g. Bálint et al., 2018; Valentini et al., 2016)

communities. However, mammals also leave eDNA signatures in water that metabarcoding can

detect (Harper et al., 2018a; Klymus et al., 2017; Ushio et al., 2017). Ponds in particular provide

drinking, foraging, dispersive, and reproductive opportunities for semi-aquatic and terrestrial

mammals (Klymus et al., 2017). Therefore, samples from these waterbodies could uncover

biodiversity present in the wider environment (Deiner et al., 2017; Harper et al., 2018a).

Drinking is a major source of eDNA deposition in water due to the release of saliva, but

mammals may also swim, wallow, urinate or defecate in water (Rodgers & Mock, 2015; Ushio et

al., 2017; Williams et al., 2018). Furthermore, arboreal mammals may use ponds less than semi-

aquatic and ground-dwelling species, non-territorial mammals may visit ponds less than

territorial species, and group-living species may leave more DNA in water than solitary species

(Williams et al., 2018). Despite evidence for eDNA deposition by semi-aquatic and terrestrial

mammals in freshwater systems, little is known about the influence of mammal behaviour on

the distribution and strength of the eDNA signal left behind.

In this study, we conducted two experiments to evaluate eDNA metabarcoding of pond

water as a tool for monitoring semi-aquatic, ground-dwelling, and arboreal mammals of

conservation or management concern. The first experiment, carried out on nine focal species

housed at two wildlife parks, examined the role of sampling strategy, mammal lifestyle, and

mammal behaviour on eDNA detection and signal strength. We hypothesised that: (1) eDNA

would be unevenly distributed, thus directed sampling would yield stronger eDNA signals (i.e.

higher read counts) for mammals than stratified sampling; (2) semi-aquatic mammals would

have stronger eDNA signals than ground-dwelling or arboreal mammals; and (3) mammal

behaviours involving water contact (e.g. swimming, drinking) would be associated with stronger

eDNA signals. The second experiment validated eDNA metabarcoding against camera trapping

and field sign searches at natural ponds within sites where focal species were present. We

hypothesised that: (1) eDNA metabarcoding would detect more mammals than camera

trapping or field signs; (2) semi-aquatic mammals would be readily detected and their eDNA

evenly distributed in ponds in comparison to terrestrial mammals; and (3) temporal sampling

would reveal that terrestrial mammal eDNA is detectable for short periods in comparison to

fully aquatic vertebrates.

130   2. Materials and methods

132   2.1 Study species

We studied nine mammal species that are the focus of European conservation or management

(Battersby & Tracking Mammals Partnership, 2005; Mathews et al., 2018): European water vole

(*Arvicola amphibius*), European otter (*Lutra lutra*), Eurasian beaver (*Castor fiber*), European

hedgehog (*Erinaceus europaeus*), European badger (*Meles meles*), red deer (*Cervus elaphus*),

Eurasian lynx (*Lynx lynx*), red squirrel (*Sciurus vulgaris*), and European pine marten (*Martes*

martes). Water vole, otter, red squirrel, pine marten and hedgehog are UK Biodiversity Action

Plan species (Joint Nature Conservation Committee, 2018). Water vole, otter, and beaver are

semi-aquatic, red squirrel and pine marten are arboreal, and the other species are ground-

dwelling. Badger and red deer live in groups whereas the other species are predominantly

solitary.

2.2 Experiment 1: eDNA detection and signal strength in artificial systems

We performed eDNA metabarcoding for focal species under artificial conditions at two wildlife parks to investigate whether mammal eDNA signal strength (i.e. read counts) depends on sampling strategy, species lifestyle (semi-aquatic, ground-dwelling, arboreal), and behaviour. Behavioural observation and eDNA sampling were conducted between 18/09/2017 – 21/09/2017 at Wildwood Trust (WT), Kent, England, and 10/10/2017 – 11/10/2017 at Royal Zoological Society of Scotland (RZSS) Highland Wildlife Park (HWP), Kingussie, Scotland. Sixteen categories of behaviour were defined based on potential contact with waterbodies and species lifestyle, and the frequency and duration of behaviours recorded (Table S1). The number of individuals in each enclosure was recorded alongside waterbody size (Table 1). Beaver, lynx, red deer, and red squirrel were present at both wildlife parks, whereas other captive species were only present at WT. Each species was observed for one hour on two separate occasions except nocturnal mammals (badger and beaver), which were observed overnight using camera traps (Bushnell Trophy Cam Aggressor). One camera trap per enclosure was positioned perpendicular to the ground (1 m height, 2 m from pond shoreline) to capture water and shoreline. Cameras took 30 s videos (1920 x 1080) when triggered (30 s interval between triggers) at high sensitivity. Behavioural observation was not undertaken for WT water voles as animals were under quarantine or HWP red squirrels as individuals were wild.

Water samples were collected from enclosures within 3 hrs of the second behavioural observation period. Up to six directed or stratified samples were collected, but sample number varied by species according to waterbody size and observed behaviours (Tables S1, S2).

Enclosure drinking containers were also sampled and classed as 'other' samples. Bathing and

drinking bowls were sampled where enclosures contained no artificial waterbodies (WT water

vole, red squirrel, and hedgehog). The HWP beaver enclosure was empty for 24 hrs before

sampling. Water was sampled from an enclosure containing beavers at RZSS Edinburgh Zoo (EZ)

and classed as 'other'. A sample was collected from a water bath in the HWP woods to capture

wild red squirrels and classed as 'other'.

Directed samples (2 L surface water taken approximately where behaviours were

observed) were collected before stratified samples (2 L surface water [8 x 250 ml pooled

subsamples] taken at equidistant points [access permitting] around the waterbody perimeter)

to minimise disturbance to the water column and cross-contamination risk. Samples were

collected using sterile Gosselin™ HDPE plastic bottles (Fisher Scientific UK Ltd, UK) and

disposable gloves. A field blank (1 L molecular grade water [MGW]) was taken into each species

enclosure and opened. Samples (n = 80) collected from WT and HWP were transported

alongside field blanks (n = 13) in sterile coolboxes with ice packs to the University of Kent (UoK)

and EZ respectively, where ice was added to coolboxes.

Samples and blanks were vacuum-filtered within 6 hrs of collection in a UoK wet

laboratory and within 24 hrs of collection in an EZ staff room. Surfaces and equipment were

sterilised before, during, and after set-up in temporary work areas. Surfaces and vacuum

pumps were wiped with 10% v/v chlorine-based commercial bleach (Elliott Hygiene Ltd, UK)

solution. Non-electrical equipment was immersed in 10% bleach solution for 10 minutes,

followed by 5% v/v MicroSol detergent (Anachem, UK), and rinsed with purified water. Up to

500 ml of each 2 L sample was vacuum-filtered through sterile 0.45 µm mixed cellulose ester

membrane filters with pads (47 mm diameter; Whatman, GE Healthcare, UK) using Nalgene™

filtration units. One hour was allowed for each sample to filter and a second filter used if

clogging occurred. A filtration blank (1 L MGW) was processed during each filtration round (n =

12), and equipment sterilised after each filtration round. After 500 ml had filtered or one hour

had passed, filters were removed from pads using sterile tweezers, placed in sterile 47 mm

petri dishes (Fisher Scientific UK Ltd, UK), sealed with parafilm (Sigma-Aldrich Company Ltd,

UK), and stored at -20 °C. The total water volume filtered per sample was recorded for

downstream analysis (Table S2; Fig. S1).

2.3 Experiment 2: eDNA detection and signal strength in natural systems

We performed eDNA metabarcoding for focal species at natural ponds in conjunction with

conventional surveys to validate this molecular approach for mammal identification and to

investigate spatiotemporal variation in mammal eDNA signals. At three sites where focal

species were present based on cumulative survey data, we selected two ponds (range 293-5056

m^2, average 1471 m^2) within 4 km of each other. The Bamff Estate (BE), Alyth, Scotland, was

selected for beaver, otter, badger, red deer, and red squirrel, but roe deer (*Capreolus*

capreolus) and red fox (*Vulpes vulpes*) were also present. Otter, water vole, and badger were

present at Tophill Low Nature Reserve (TLNR), Driffield, East Yorkshire, alongside American

mink (*Neovison vison*), stoat (*Mustela erminea*), weasel (*Mustela nivalis*), rabbit (*Oryctolagus*

cuniculus), brown hare (*Lepus europaeus*), red fox, roe deer, and grey squirrel (*Sciurus*

carolinensis). We selected Thorne Moors (TM), Doncaster, South Yorkshire, for red deer and

badger, but stoat, weasel, red fox, roe deer, and Reeve's muntjac (*Muntiacus reevesi*) were also

present. Camera traps (Bushnell Trophy Cam Aggressor) were deployed at TM (one per pond)

and BE (three per pond) one week prior to eDNA sampling and collected once sampling was

completed. At TLNR, camera traps (two to three per pond) were deployed one day before a 5-

day period of eDNA sampling and collected one week after sampling was completed. Camera

traps were positioned perpendicular to the ground (1 m height, 0.3-1 m from pond shoreline)

to capture water and shoreline. Cameras took three photographs (5 megapixel) when triggered

(3 s interval between triggers) at high sensitivity.

Ten stratified samples were collected from the shoreline of each pond (TM: 17/04/2018;

BE: 20/04/2018; TLNR: 23/04/2018 -27/04/2018) and a field blank (1 L MGW) included as in

Experiment 1. TLNR ponds were sampled every 24 hrs over 5 days to investigate spatiotemporal

variation in mammal eDNA signals. TM and TLNR samples were transported on ice in sterile

coolboxes to the University of Hull (UoH) eDNA facility, and stored at 4 °C. BE samples were

transported in sterile coolboxes with ice packs to BE accommodation. Surfaces and equipment

were sterilised before, during, and after set-up as in Experiment 1. Samples (*n* = 140) and field

blanks (*n* = 14) were vacuum-filtered within 4 hrs of collection as in Experiment 1 with minor

modifications to maximise detection probability as follows. The full 2 L of each sample was

vacuum-filtered where possible, two filters were used for each sample, and duplicate filters

were stored in one petri dish at -20 °C. A filtration blank (1 L MGW) was processed during each

filtration round (*n* = 21). The total water volume filtered per sample was recorded (Table S3).

2.4 DNA extraction

DNA was extracted within 2 weeks of filtration at the UoH eDNA facility using the mu-DNA water protocol (Sellers, Di Muri, Gómez, & Hänfling, 2018). Duplicate filters from samples in Experiment 1 were lysed independently and the lysate from each loaded onto one spin column. As more samples were collected in Experiment 2, duplicate filters were co-extracted by placing both in a single tube for bead milling. An extraction blank, consisting only of extraction buffers, was included for each round of DNA extraction (n = 17). Eluted DNA (100 µl) was stored at -20 °C until PCR amplification.

2.5 eDNA metabarcoding

Our eDNA metabarcoding workflow is fully described in Supporting Information. Briefly, we performed nested metabarcoding using a two-step PCR protocol, where Multiplex Identification (MID) tags were included in the first and second PCR for sample identification (Kitson et al., 2019). In the first PCR, eDNA was amplified in triplicate with published 12S ribosomal RNA (rRNA) primers 12S-V5-F (5'-ACTGGGATTAGATACCCC-3') and 12S-V5-R (5'-TAGAACAGGCTCCTCTAG-3') (Riaz et al., 2011) that were validated *in silico* for UK vertebrates (Harper et al., 2018a, b). The *in silico* analysis found that 91/112 mammal species listed on the Natural History Museum Checklist of Mammalia v1 (subspecies excluded) can be distinguished using this 12S metabarcode. Nine indistinguishable species lacked reference sequences, whereas the other 12 had reference sequences but did not amplify. PCR positive controls (two

per PCR plate; *n* = 16) were exotic cichlid (*Maylandia zebra*) DNA (0.05 ng/μl), and PCR negative

controls (two per PCR plate; *n* = 16) were MGW (Fisher Scientific UK Ltd, UK). PCR products

were pooled to create sub-libraries (Fig. S2) and purified. The second PCR bound pre-adapters,

MID tags, and Illumina adapters to the sub-libraries. PCR products were purified, quantified,

and pooled. The library was then purified, quantified by qPCR, and fragment size (330 bp) and

removal of secondary product verified. The library (220 eDNA samples, 27 field blanks, 33

filtration blanks, 17 extraction blanks, 16 PCR negative controls, and 16 PCR positive controls)

was sequenced on an Illumina MiSeq®. Raw sequence reads were demultiplexed using a custom

Python script, then metaBEAT v0.97.11 (https://github.com/HullUni-bioinformatics/metaBEAT)

was used for quality trimming, merging, chimera removal, clustering, and taxonomic

assignment of sequences against our UK vertebrate reference database (Harper et al., 2018a, b)

which contains sequences for 103 UK mammals.

2.6 Data analysis

Analyses were performed in R v.3.4.3 (R Core Team, 2017). The total unrefined read counts (i.e.

raw taxonomically assigned reads) per sample were calculated and retained for downstream

analyses. Assignments were corrected and merged, taxon-specific sequence thresholds (i.e.

maximum sequence frequency of each taxon in PCR positive controls [Table S4]) used to

mitigate cross-contamination and false positives (Fig. S3), and remnant contaminants and

higher taxonomic assignments removed excluding the following genera. *Anas* (Dabbling ducks)

was retained because potential for hybridisation reduced confidence in species-level

assignments, and *Emberiza* (Buntings) and *Larus* (White-headed gulls) were retained because

reference sequences were missing for several common species. Dataset refinement is fully

described in Supporting Information: Appendix 1. Taxonomic assignments remaining in the

refined dataset were predominantly of species resolution and considered true positives. We

split the refined dataset by Experiment 1 (artificial waterbodies) and Experiment 2 (natural

ponds). Proportional read counts for each species were calculated from the total unrefined

read counts per sample. Our proportional read count data were not normally distributed

(Shapiro–Wilk normality test: W = 0.915, P < 0.001), thus we used a Mann-Whitney U test to

compare the median proportional read count of stratified and directed samples across species.

We employed binomial Generalized Linear Mixed-effects Models (GLMMs) with the logit

link function using the package glmmTMB (development version; Brooks et al., 2017) for the

following tests. First, we compared the eDNA signals from stratified and directed samples for

each mammal species using a hierarchical model including sample type nested within species

(fixed) and wildlife park (random) as effects. We tested the influence of species lifestyle on

mammal eDNA signals using a model with species lifestyle (fixed) and species nested within

wildlife park (random) as effects. Using directed samples, we tested the influence of behaviour

on mammal eDNA signals using two hierarchical models, including species nested within

wildlife park (random) and specific (e.g. swimming, drinking) or generic (i.e. water contact

versus no water contact) behaviour(s) respectively (fixed) as effects. We assessed model fit

using diagnostic plots and performed validation checks to ensure model assumptions were met

and overdispersion was absent (Zuur, Ieno, Walker, Saveliev, & Smith, 2009).

For Experiment 2, we qualitatively compared mammal presence-absence records

generated by eDNA metabarcoding, camera trapping, and field signs. TLNR ponds were

sampled every 24 hrs for 5 days, thus proportional read counts were averaged across days for

comparison to BE and TM ponds (sampled once each). We qualitatively compared the

distribution and persistence of eDNA signals between semi-aquatic and terrestrial mammals

using tile plots and heat maps of the unaveraged proportional read counts for identified species

at TLNR over the 5-day period. All figures were produced using the package ggplot2 v3.0.0

(Wickham, 2016).

3. Results

3.1 eDNA metabarcoding

The sequencing run generated 47,713,656 raw sequence reads, of which 37,590,828 remained

following trimming, merging, and length filter application. After removal of chimeras and

redundancy via clustering, the library contained 21,127,061 sequences (average read count of

64,215 per sample including controls), of which 16,787,750 (79.46%) were assigned a

taxonomic rank. Contamination (Fig. S4) was observed in the field blanks (badger, beaver, lynx,

pine marten, red squirrel, and water vole) as well as in the filtration and extraction blanks

(human [*Homo sapiens*] and cichlid). PCR negative controls were contaminated to different

extents with human, cichlid, beaver, and pine marten as well as non-focal species. After

threshold application, contaminants remaining in eDNA samples included Gentoo penguin

(*Pygoscelis papua*), reindeer (*Rangifer tarandus*), cichlid, and human. The refined dataset

contained 59 vertebrate species, including six amphibians, 10 fish, 19 birds, and 24 mammals

(Table S5).

3.2 Experiment 1: eDNA detection and signal strength in artificial systems

All nine focal species were detected in captivity. Seven species (red deer and hedgehog

excluded) were detected in all water samples taken from their respective enclosures. Red deer

were not detected in 2 of 5 stratified samples from HWP, and hedgehog was not detected in 1

of 2 WT drinking bowl samples (Fig. 1). 'Other' samples (neither directed nor stratified) were

excluded from further comparisons, thus hedgehog, red squirrel, and water vole were omitted

in downstream analyses.

Across species, stratified samples (0.406) had a higher median proportional read count

than directed samples (0.373), but this difference was not significant (Mann-Whitney U test: U

= 1181.5, P = 0.829). Within species, sample type also had no effect (χ^2_6 = 0.364, P = 0.999) on

proportional read counts (GLMM: θ = 0.168, χ^2_{53} = 8.915, P = 1.000, pseudo-R^2 = 39.21%).

Proportional read counts for directed and stratified samples belonging to each species did not

significantly differ. Otter proportional read counts were lower than other species, but not

significantly so (Fig 2a). Similarly, species lifestyle had no influence (χ^2_2 = 0.655, P = 0.721) on

proportional read counts (GLMM: θ = 0.213, χ^2_{61} = 13.002, P = 1.000, pseudo-R^2 = 11.85%).

Proportional read counts for semi-aquatic, ground-dwelling, and arboreal species did not

significantly differ (Fig. 2b).

Specific behaviours exhibited by species had no influence (χ^2_{11} = 1.369, P = 0.999) on

proportional read counts (GLMM: θ = 0.355, χ^2_{31} = 11.013, P = 0.999, pseudo-R^2 = 9.17%),

where proportional read counts did not differ across species' behaviours (Fig. 3a). Likewise,

generic behaviour (i.e. water contact versus no water contact) had no effect on proportional

read counts (χ^2_{11} = 0.002, P = 0.964). Proportional read counts did not differ between

behaviour that involved or lacked water contact (Fig. 3b, GLMM: θ = 0.217, χ^2_{41} = 8.897, P =

1.000, pseudo-R^2 = 8.50%).

3.3 Experiment 2: eDNA detection and signal strength in natural systems

At natural ponds, eDNA metabarcoding, camera trapping, and field signs all detected beaver,

red deer, and roe deer. Camera traps (Fig. 4) and field signs recorded red fox and badger when

eDNA metabarcoding did not (Fig. 5). However, eDNA metabarcoding revealed small mammal

species not caught on camera, including water vole, water shrew (*Neomys fodiens*), bank vole

(*Myodes glareolus*), common shrew (*Sorex araneus*), brown rat (*Rattus norvegicus*), rabbit, grey

squirrel, and common pipistrelle (*Pipistrellus pipistrellus*). Mice or vole footprints were

observed at BE Pond 1, but species could not be determined. Fig. 5 summarises mammals

recorded by different methods at each site with reference to cumulative survey data. Notably,

only beaver presence was captured at the same ponds by all methods. Although methods

shared species at site level, species were not always detected at the same pond. Detection

rates for species captured by at least one survey method are summarised in Table S6.

Sampling of natural ponds revealed spatial patterns in eDNA detection and signal

strength. eDNA from non-domestic terrestrial mammals (i.e. mammals excluding dog [*Canis*

lupus familiaris], pig [*Sus scrofa domesticus*], sheep [*Ovis aries*] and cow [*Bos taurus*]) was

unevenly dispersed compared with semi-aquatic mammals (Fig. S5). Semi-aquatic beaver and

water vole were detected in at least 90% and 60% respectively of water samples (*n* = 10)

collected from single ponds, whereas water shrew was only detected in 10% of samples.

Conversely, non-domestic terrestrial mammals were consistently detected in less than 20% of

water samples collected from a pond and left relatively weak eDNA signals. Overall, beaver was

the most consistently detected mammal with the highest proportional read counts. However,

the strongest and most evenly distributed signals belonged to amphibians, particularly common

frog (*Rana temporaria*) and great crested newt (*Triturus cristatus*) (Fig. S5).

TLNR samples collected over a 5-day period (D01-05) revealed that mammal detection

was highly dependent on the spatial and temporal resolution of eDNA metabarcoding surveys

(Fig. S6). Mammal eDNA signals in pond water were ephemeral, often disappearing within 24-

48 hrs of initial detection, as opposed to amphibians that were detected for multiple days and

whose eDNA signal increased in strength. The majority of semi-aquatic or terrestrial mammals

were only detected in a single sample on each day.

4. Discussion

We have demonstrated that eDNA metabarcoding is a potential monitoring tool for

conservation and management priority mammals, but species detection rates are variable. We

used two experiments to validate this molecular approach and gained new insights that will

inform the development and application of eDNA metabarcoding for mammals. Sampling

strategy, mammal lifestyle, and mammal behaviour had little influence on eDNA detection and

signal strength in captivity, but all played vital roles in natural ponds. Although mammals were

detected from pond water, their eDNA signals were temporary and weak in comparison to

amphibians and fish. Nonetheless, this would suggest eDNA is representative of contemporary

and local mammal diversity in a given area.

4.1 Experimental insights

In Experiment 1, all nine focal species were detected in captivity, of which seven were detected

in all water samples taken from their respective enclosures. This demonstrates that our method

can successfully detect a wide range of mammal species from pond and drinking water. Perhaps

surprisingly, we found that neither sampling strategy, mammal lifestyle or mammal behaviour

influenced eDNA detectability and signal strength in enclosures. This included behaviours

associated with eDNA deposition, e.g. swimming, drinking, urination, and defecation (Rodgers

& Mock, 2015; Ushio et al., 2017; Williams et al., 2018). Enclosures were permanently occupied

and artificial waterbodies likely saturated with eDNA, which possibly masked behavioural

signals. Modest replication may also have limited experimental power, preventing patterns

being detected statistically. Nonetheless, our results have demonstrated that mammal contact

with water enables eDNA deposition and detection.

Unsurprisingly, given the nature of wild mammal interactions with natural systems

versus those in captivity, results from Experiment 2 highlighted the challenges of mammal

eDNA detection. We detected 17 mammals using three monitoring tools, comparable to the 17

mammals expected from cumulative survey data despite some differences. Field signs and

camera trapping performed better than eDNA metabarcoding for red fox and badger detection,

but only eDNA metabarcoding identified water vole and other small mammals missed on

camera or with ambiguous field signs, e.g. mice, voles, shrews. Importantly, the deployment

period, height, and positioning of camera traps may have influenced small mammal detection

by this method (Caravaggi et al., 2018). Ishige et al. (2017) achieved comparable mammal

detection at salt licks with eDNA metabarcoding and camera trapping, but species presence was

inconsistent between salt licks surveyed. Currently, there are no published studies comparing

eDNA metabarcoding of water to camera trapping for mammal identification. However, our

results reflect those of Stat et al. (2019) who found highest marine fish species richness using

both baited remote underwater video systems and eDNA metabarcoding, but each method

identified unique species.

Notably, no survey method captured the semi-aquatic otter despite known presence at

study sites and successful detection in eDNA metabarcoding studies of UK ponds (Harper et al.,

2018a) and lakes (Hänfling et al., 2017). Furthermore, the eDNA signal for captive otter was

weaker than that of other semi-aquatic mammals studied here. Lower eDNA detection rates for

otter, badger, and red fox may stem from species' ecologies. These mammals are wide-ranging

(Gaughran et al., 2018; Thomsen et al., 2012) and may not readily release DNA in water. Otters

often spraint on grass or rock substrata outside water and use latrines associated with caves

and dens (Ruiz-Olmo & Gosálbez, 1997). As terrestrial mammals, red fox and badger must drink

from or enter ponds for eDNA deposition to occur (Rodgers & Mock, 2015; Ushio et al., 2017;

Williams et al., 2018). Otter, badger, and red fox detection may require greater spatiotemporal

resolution of eDNA sampling. This is reinforced by other eDNA metabarcoding studies where

mammal detection was highly variable across sites surveyed (Ishige et al., 2017; Klymus et al.,

2017; Ushio et al., 2017). False negatives may instead be symptomatic of metabarcoding bias,

but this is unlikely in our study (section 4.2).

eDNA from other semi-aquatic mammals (e.g. beaver, water vole) was evenly

distributed, being found in most or all samples collected on fine spatial scales within natural

ponds, whereas terrestrial mammal (e.g. red and roe deer) eDNA was highly localised and often

detected in less than 20% of samples. Mammal eDNA signals varied temporally, being

detectable for two consecutive days maximum. Depending on the species, mammal eDNA may

be spatially and temporally clumped in lentic ecosystems due to the nature and frequency of

contact that individuals have with water. Unless behaviours involving prolonged water contact

(e.g. swimming, wallowing) are exhibited, non-domestic mammals may only be detected at

drinking sites (Klymus et al., 2017; Ushio et al., 2017; Williams et al., 2018). In contrast,

domestic mammals (e.g. cow, sheep, pig) may have elevated detection rates in ponds due to

high occurrence of these waterbodies in agricultural landscapes as well as eDNA transport by

rainfall and run-off (Staley et al., 2018). eDNA detection and persistence are further influenced

by group size, where eDNA from multiple individuals endures for longer time periods in water

than eDNA from single individuals (Williams et al., 2018). Detailed investigations incorporating

biotic (e.g. population size, sex, body mass, behaviour) and abiotic (e.g. temperature, pH,

ultraviolet light, rainfall) factors are needed to understand the longevity of mammal eDNA

signals in aquatic ecosystems (Rodgers & Mock, 2015; Williams et al., 2018).

Both of our experiments have shown that sampling strategy drastically influences

mammal eDNA detection probability. Mammal eDNA was evenly distributed in closed, artificial

waterbodies, but locally distributed in open, natural ponds. In captivity, mammal species were

housed individually (excluding HWP red deer) with a drinking container(s) and/or small

waterbody (range 0.01-162 m^2, mean 27.4 m^2) in their enclosure. Some enclosures housed

more individuals of a species than others, thereby increasing eDNA deposition and detection

probability (Williams et al., 2018). Wild mammals have an array of freshwater habitats at their

disposal and can hold vast territories. Therefore, rates of pond visitation and eDNA deposition

are more irregular (Klymus et al., 2017; Ushio et al., 2017) which may lead to between-sample

variation (Williams et al., 2018).

4.2 Technical pitfalls of eDNA metabarcoding for mammal monitoring

eDNA metabarcoding has strong potential for inclusion in mammal monitoring schemes

(section 4.3), but like existing monitoring tools, may produce false negatives or false positives.

Our process controls identified low-level contamination at all stages of metabarcoding, but

contaminants primarily occurred in field and PCR negative controls (Supporting Information:

Appendix 2). We applied taxon-specific sequence thresholds to our data to mitigate false

positives as in Harper *et al*. (2018a, b). Remnant contaminants were cichlid (laboratory), Gentoo

penguin (environment), reindeer (environment), and human (environment/laboratory). Gentoo

penguin is housed at EZ and was identified from EZ beaver enclosure water. The WT red squirrel

and reindeer enclosures are in close proximity. DNA transport by wildlife (e.g. waterfowl

[Hänfling et al., 2016]) and park staff or visitors may explain this environmental contamination.

Human DNA was present across process controls corresponding to artificial and natural

waterbodies. Human DNA may be amplified and sequenced instead of focal species, potentially

resulting in false negative detections for rare and/or less abundant species. Human DNA

blocking primers can prevent this bias, but may impair PCR amplification efficiency (Klymus et

al., 2017; Ushio et al., 2017; Valentini et al., 2016).

In our study, eDNA metabarcoding produced false negatives for otter, badger, and red

fox at natural ponds. We selected a 12S metabarcode designed to amplify vertebrate DNA (Riaz

et al., 2011). There were no primer mismatches with otter, badger or red fox reference

sequences (Harper et al., 2018a, b), thus amplification bias was not responsible for these false

negatives. DNA from purely aquatic and more abundant species may have overwhelmed otter,

badger, and red fox DNA during amplification and sequencing, i.e. species-masking (Kelly, Port,

Yamahara, & Crowder, 2014; Klymus et al., 2017). Metabarcoding primers targeting mammals

(Ushio et al., 2017), or multi-marker (e.g. 12S, 16S, COI) investigations (Evans et al., 2017;

Hänfling et al., 2016; Kelly et al., 2014; Klymus et al., 2017) may improve mammal detection in

systems with competition from aquatic species. Similarly, more biological and technical

replication may improve species detection probabilities (Evans et al., 2017; Lawson Handley et

al., 2018; Stat et al., 2019; Valentini et al., 2016). Importantly, fish and amphibians also had

higher qPCR detection than otter (Thomsen et al., 2012). A comparison of metabarcoding and

qPCR (e.g. Harper et al., 2018b; Lacoursière-Roussel, Dubois, Normandeau, & Bernatchez, 2016)

would confirm whether poor amplification efficiency for otter is due to technical bias or species

ecology, and whether eDNA metabarcoding can reliably monitor otter alongside the wider

mammal community.

4.3 Scope of eDNA metabarcoding for mammal monitoring

Assessments of UK mammal populations are hindered by lack of data and systematic

monitoring for many species (Mathews et al., 2018). Distribution and occupancy data are poor

for most species, with ongoing survey effort biased toward rare species. Surveys are also

heavily reliant on citizen science and casual records (Massimino et al., 2018). Tools that can

provide standardised, systematic monitoring of UK mammal populations are needed (Mathews

et al., 2018). Despite issues inherent to metabarcoding for biodiversity monitoring (Deiner et

al., 2017), this tool has enormous potential to enhance mammal monitoring, conservation, and

management. eDNA metabarcoding generates distribution data for multiple species, whether

rare, invasive, or abundant, and could be a powerful tool for tracking species in conflict with

one another, e.g. water vole, American mink, and otter (Bonesi & Macdonald, 2004) or red

squirrel, grey squirrel, and pine marten (Sheehy, Sutherland, O'Reilly, & Lambin, 2018).

Different sample types (e.g. water, soil, snow, salt licks, feeding traces, faeces, hair, and blood

meals) may also offer new insights to mammal biodiversity in a given area (Aylward et al., 2018;

Franklin et al., 2019; Ishige et al., 2017; Tessler et al., 2018; Ushio et al., 2017).

eDNA metabarcoding can rapidly survey multitudes of aquatic sites at landscape-scale

where camera traps might be resource-intensive, cost-inefficient, and susceptible to

theft/damage (Stat et al., 2019; Ushio et al., 2017). Field signs require volunteer time and skill

to be employed at comparable spatial scales to eDNA metabarcoding (Sadlier et al., 2004).

eDNA metabarcoding could provide accurate data for species misidentified from field signs, e.g.

mice and voles, otter and mink (Franklin et al., 2019; Harris & Yalden, 2004). However, both

camera traps and field signs recorded species that eDNA metabarcoding did not. Therefore,

eDNA metabarcoding is complementary and should be incorporated into, not replace, existing

monitoring schemes. This tool could be most effective in mammal monitoring if deployed at the

edges of known species distributions, in areas where species presence is unknown, and in areas

with isolated species records (Mathews et al., 2018).

Mammal eDNA detection probabilities from natural ponds will likely be high when areas

with dense populations are studied. However, rigorous sampling strategies are required to

track mammals in areas that are sparsely populated by individuals. Multiple ponds must be

sampled repeatedly, and samples taken at multiple locations within ponds without pooling so

that inferences about site occupancy can be made. Importantly, we sampled natural ponds in

spring but sampling in other seasons may produce different results, reflective of species'

ecologies (Lawson Handley et al., 2018). We recommend that researchers and practitioners

using eDNA metabarcoding for mammal monitoring channel efforts into extensive sampling of

numerous waterbodies in a given area over larger timescales. This will account for differential

mammal visitation rates and maximise eDNA detection probabilities.

Data accessibility

Raw sequence reads have been archived on the NCBI Sequence Read Archive (Study: SRP164740; BioProject: PRJNA495011; BioSamples: SAMN10195928 - SAMN10196255; SRA accessions: SRR7986451 - SRR7986778). Jupyter notebooks, R scripts and corresponding data are archived online (https://doi.org/10.5281/zenodo.2561415).

Acknowledgements

The authors declare that they have no conflicts of interest in presenting this work for publication. This work was funded by the University of Hull, and D.S.R was supported by the Natural Environment Research Council award number NE/R016429/1 as part of the UK-SCAPE programme delivering National Capability. We would like to thank Helen Senn (RZSS Edinburgh Zoo) for feedback on the study design, and Richard Griffiths (DICE, University of Kent) for support with filtration of water samples from Wildwood Trust. We are grateful to staff at Wildwood Trust and RZSS Highland Wildlife Park for assisting with eDNA sampling from animal enclosures. We thank Richard Hampshire and volunteers at Tophill Low Nature Reserve for assisting with camera trap deployment. Tim Kohler (Natural England) and Paul Ramsay kindly gave permission to sample at Thorne Moors and the Bamff Estate respectively.

Author contributions

L.R.H, B.H, and L.L.H conceived and designed the study. A.I.C and G.M-D coordinated sampling at Wildwood Trust and RZSS Highland Wildlife Park respectively. L.R.H, C.D.M, C.J.M, and T.L collected and filtered water samples. A.L, T.L, and T.B helped select natural ponds to be surveyed using eDNA, camera trapping, and field signs, and provided camera traps for the study. L.R.H, A.L, and T.L deployed camera traps, which were then collected and footage analysed by L.R.H. L.R.H processed samples in the laboratory with advice from C.D.M and A.M. D.S.R sequenced the final library. L.R.H completed bioinformatic processing of samples, and subsequent data analysis. L.R.H wrote the manuscript, which all authors contributed critically to drafts of and gave final approval for publication.

References

Aylward, M. L., Johnson, S. E., Sullivan, A. P., Perry, G. H., & Louis, E. E. (2018). A novel environmental DNA (eDNA) sampling method for aye-ayes from their feeding traces. *bioRxiv, 272153*. https://doi.org/10.1101/272153

Bálint, M., Nowak, C., Márton, O., Pauls, S. U., Wittwer, C., Aramayo, J. L., Schulze, A., Chambert, T., Cocchiararo, B. & Jansen, M. (2018). Accuracy, limitations and cost efficiency of eDNA-based community survey in tropical frogs. *Molecular Ecology Resources*, *18*, 1415–1426. https://doi.org/10.1111/1755-0998.12934

Battersby, J. & Tracking Mammals Partnership (2005) *UK Mammals: Species Status and Population Trends. First Report by the Tracking Mammals Partnership*. Joint Nature Conservation Committee /Tracking Mammals Partnership, Peterborough.

Bland, L. M., Collen, B., Orme, C. D. L., & Bielby, J. (2015). Predicting the conservation status of
data-deficient species: Predicting Extinction Risk. *Conservation Biology*, *29*, 250–259.
https://doi.org/10.1111/cobi12372

Bonesi, L., & Macdonald, D. W. (2004). Differential habitat use promotes sustainable
coexistence between the specialist otter and the generalist mink. *Oikos* , *106*, 509–
519. https://doi.org/10.1111/j.0030-1299.2004.13034.x

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug,
H. J., Machler, M. & Bolker, B. M. (2017). glmmTMB Balances Speed and Flexibility Among
Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal*, *9*, 378–400.
https://doi.org/10.32614/RJ-2017-066

Burton, A. C., Neilson, E., Moreira, D., Ladle, A., Steenweg, R., Fisher, J. T., Bayne, E. & Boutin, S.
(2015). REVIEW: Wildlife camera trapping: a review and recommendations for linking
surveys to ecological processes. *Journal of Applied Ecology*, *52*, 675–
685. https://doi.org/10.1111/1365-2664.12432

Caravaggi, A., Gatta, M., Vallely, M. -C., Hogg, K., Freeman, M., Fadaei, E., Dick, J. T. A.,
Montgomery, W. I., Reid, N. & Tosh, D. G. (2018). Seasonal and predator-prey effects on
circadian activity of free-ranging mammals revealed by camera traps. *PeerJ*, *6*, e5827.
https://doi.org/10.7717/peerj.5827

Deiner, K., Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., Creer, S.,
Bista, I., Lodge, D. M., de Vere, N., Pfrender, M. E. & Bernatchez, L. (2017). Environmental
DNA metabarcoding: transforming how we survey animal and plant communities. *Molecular*
Ecology, *26*, 5872–5895. https://doi.org/10.1111/mec.14350

Evans, N. T., Li, Y., Renshaw, M. A., Olds, B. P., Deiner, K., Turner, C. R., Jerde, C. L., Lodge, D. M.,
Lamberti, G. A. & Pfrender, M. E (2017). Fish community assessment with eDNA
metabarcoding: effects of sampling design and bioinformatic filtering. *Canadian Journal of*
Fisheries and Aquatic Sciences, *74*, 1362–1374. https://doi.org/10.1139/cjfas-2016-0306

Franklin, T. W., McKelvey, K. S., Golding, J. D., Mason, D. H., Dysthe, J. C., Pilgrim, K. L., Squires,
J. R., Aubry, K. B., Long, R. A., Greaves, S. E., Raley, C. M., Jackson, S., MacKay, P., Lisbon, J.,
Sauder, J. D., Pruss, M. T., Heffington, D. & Schwartz, M. K (2019). Using environmental DNA
methods to improve winter surveys for rare carnivores: DNA from snow and improved
noninvasive techniques. *Biological Conservation*, *229*, 50–58.
https://doi.org/10.1016/j.biocon.2018.11.006

Gaughran, A., Kelly, D. J., MacWhite, T., Mullen, E., Maher, P., Good, M., & Marples, N. M.
(2018). Super-ranging. A new ranging strategy in European badgers. *PLoS ONE*, *13*,
e0191818. https://doi.org/10.1371/journal.pone.0191818

Hänfling, B., Lawson Handley, L., Lunt, D., Shum, P., Winfield, I., & Read, D. S. (2017). *A Review*
of Recent Advances in Genetic Methods to Identify Improvements in CAMERAS Partners
Monitoring Activities. University of Hull, Hull, UK.

Hänfling, B., Lawson Handley, L., Read, D. S., Hahn, C., Li, J., Nichols, P., Blackman, R. C., Oliver,
A. & Winfield, I. J. (2016). Environmental DNA metabarcoding of lake fish communities
reflects long-term data from established survey methods. *Molecular Ecology*, *25*, 3101–3119.
https://doi.org/10.1111/mec.13660

Harper, L. R., Handley, L. L., Hahn, C., Boonham, N., Rees, H. C., Lewis, E., Adams, I. P.,
Brotherton, P., Phillips, S. & Hänfling, B. (2018a) Ground-truthing environmental DNA (eDNA)
metabarcoding for ecological hypothesis testing at the pondscape. *bioRxiv*, *278309*.
https://doi.org/10.1101/278309

Harper, L. R., Lawson Handley, L., Hahn, C., Boonham, N., Rees, H. C., Gough, K. C., Lewis, E.,
Adams, I. P., Brotherton, P., Phillips, S. & Hänfling, B. (2018b). Needle in a haystack? A
comparison of eDNA metabarcoding and targeted qPCR for detection of the great crested
newt (*Triturus cristatus*). *Ecology and Evolution*, *8*, 6330–6441.
https://doi.org/10.1002/ece3.4013

Harris, S., & Yalden, D. W. (2004). An integrated monitoring programme for terrestrial mammals
in Britain. *Mammal Review*, *34*, 157–167. https://doi.org/10.1046/j.0305-1838.2003.00030.x

Ishige, T., Miya, M., Ushio, M., Sado, T., Ushioda, M., Maebashi, K., Yonechi, R., Lagan, P. &
Matsubayashi, H. (2017). Tropical-forest mammals as detected by environmental DNA at
natural saltlicks in Borneo. *Biological Conservation*, *210*, 281–285.
https://doi.org/10.1016/j.biocon.2017.04.023

Joint Nature Conservation Committee (2018) UK BAP priority terrestrial mammal species.
Accessed on 1[st] December 2018. http://jncc.defra.gov.uk/page-5170

Kelly, R. P., Port, J. A., Yamahara, K. M., & Crowder, L. B. (2014). Using environmental DNA to
census marine fishes in a large mesocosm. *PLoS ONE*, *9*, e86175.
https://doi.org/10.1371/journal.pone.0086175

Kitson, J. J. N., Hahn, C., Sands, R. J., Straw, N. A., Evans, D. M., & Lunt, D. H. (2019). Detecting
host-parasitoid interactions in an invasive Lepidopteran using nested tagging DNA-
metabarcoding. *Molecular Ecology*, *28*, 471–483. https://doi.org/10.1111/mec.14518

Klymus, K. E., Richter, C. A., Thompson, N., & Hinck, J. E. (2017). Metabarcoding of
Environmental DNA Samples to Explore the Use of Uranium Mine Containment Ponds as a
Water Source for Wildlife. *Diversity*, *9*, 54. https://doi.org/10.3390/d9040054

Lacoursière-Roussel, A., Dubois, Y., Normandeau, E., & Bernatchez, L. (2016). Improving
herpetological surveys in eastern North America using the environmental DNA method.
Genome, *59*, 991–1007. https://doi.org/10.1139/gen-2015-0218

Lawson Handley, L. J., Read, D., Winfield, I., Kimbell, H., Johnson, H., Li, J., Hahn, C., Blackman,
R., Wilcox, R., Donnelly, R., Szitenberg, A. & Hänfling, B. (2018) Temporal and spatial
variation in distribution of fish environmental DNA in England's largest lake. *bioRxiv*, *376400*.
https://doi.org/10.1101/376400

Lugg, W. H., Griffiths, J., van Rooyen, A. R., Weeks, A. R., & Tingley, R. (2017). Optimal survey
designs for environmental DNA sampling. *Methods in Ecology and Evolution*, *9*, 1049–1059.
https://doi.org/10.1111/2041-210X.12951

Massimino, D., Harris, S. J., & Gillings, S. (2018). Evaluating spatiotemporal trends in terrestrial
mammal abundance using data collected during bird surveys. *Biological Conservation*, *226*,
153–167. https://doi.org/10.1016/j.biocon.2018.07.026

Mathews, F., Kubasiewicz, L. M., Gurnell, J., Harrower, C. A., McDonald, R. A. & Shore, R. F.
(2018) *A Review of the Population and Conservation Status of British Mammals: Technical*
Summary. Natural England, Peterborough.

Padgett-Stewart, T. M., Wilcox, T. M., Carim, K. J., McKelvey, K. S., Young, M. K., & Schwartz, M.
K. (2016). An eDNA assay for river otter detection: a tool for surveying a semi-aquatic
mammal. *Conservation Genetics Resources*, *8*, 5–7. https://doi.org/10.1007/s12386-015-
0511-s

Riaz, T., Shehzad, W., Viari, A., Pompanon, F., Taberlet, P., & Coissac, E. (2011). ecoPrimers:
inference of new DNA barcode markers from whole genome sequence analysis. *Nucleic Acids*
Research, *39*, e145. https://doi.org/10.1093/nar/gkr732

Rodgers, T. W., & Mock, K. E. (2015). Drinking water as a source of environmental DNA for the
detection of terrestrial wildlife species. *Conservation Genetics Resources*, *7*, 693–696.
https://doi.org/10.1007/s12686-015-0478-7

Ruiz-Olmo, J., & Gosálbez, J. (1997). Observations on the sprainting behaviour of the otter *Lutra*
lutra in the NE Spain. *Acta Theriologica*, *42*, 259–270.

Sadlier, L. M. J., Webbon, C. C., Baker, P. J., & Harris, S. (2004). Methods of monitoring red foxes
Vulpes vulpes and badgers Meles meles: are field signs the answer? *Mammal Review*, *34*,
75–98. https://doi.org/10.1046/j.0305-1838.2003.00029.x

Sellers, G. S., Di Muri, C., Gómez, A., & Hänfling, B. (2018). Mu-DNA: a modular universal DNA
extraction method adaptable for a wide range of sample types. *Metabarcoding and*
Metagenomics, *2*, e24556. https://doi.org/10.3897/mbmg.2.24556

Sheehy, E., Sutherland, C., O'Reilly, C. & Lambin, X. (2018) The enemy of my enemy is my friend:
native pine marten recovery reverses the decline of the red squirrel by suppressing grey
squirrel populations. *Proceedings of the Royal Society of London B*, *285*, 20172603.
https://doi.org/10.1098/rspb.2017.2603

Staley, Z. R., Chuong, J. D., Hill, S. J., Grabuski, J., Shokralla, S., Hajibabaei, M., & Edge, T. A.
(2018). Fecal source tracking and eDNA profiling in an urban creek following an extreme rain
event. *Scientific Reports*, *8*, 14390. https://doi.org/10.1038/s41598-018-32680-z

Stat, M., John, J., DiBattista, J. D., Newman, S. J., Bunce, M. & Harvey, E. S. (2019) Combined use
of eDNA metabarcoding and video surveillance for the assessment of fish biodiversity.
Conservation Biology, *33*, 196–205. https://doi.org/10.1111/cobi.13183

Tessler, M., Weiskopf, S. R., Berniker, L., Hersch, R., Mccarthy, K. P., Yu, D. W., & Siddall, M. E.
(2018). Bloodlines: mammals, leeches, and conservation in southern Asia. *Systematics and*
Biodiversity, *16*, 488–496. https://doi.org/10.1080/14772000.2018.1433729

Thomsen, P. F., Iversen, L. L., Wiuf, C., Rasmussen, M., Gilbert, M. T. P., Orlando, L., &
Willerslev, E. (2012). Monitoring endangered freshwater biodiversity using environmental
DNA. *Molecular Ecology*, *21*, 2565–2573. https://doi.org/10.1111/j.1365-294X.2011.05418.x

Thomsen, P. F., & Willerslev, E. (2015). Environmental DNA–an emerging tool in conservation
for monitoring past and present biodiversity. *Biological Conservation*, *183*, 4–18.
https://doi.org/10.1016/j.biocon.2014.11.019

Ushio, M., Fukuda, H., Inoue, T., Makoto, K., Kishida, O., Sato, K., Murata, K., Nikaido, M., Sado,
T., Sato, Y., Takeshita, M., Iwasaki, W., Yamanaka, H., Kondoh, M. & Miya, M. (2017).
Environmental DNA enables detection of terrestrial mammals from forest pond water.
Molecular Ecology Resources, *17*(6), e63–e75. https://doi.org/10.1111/1755-0998.12690

Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P. F., Bellemain, E.,
Besnard, A., Coissac, E., Boyer, F., Gaboriaud, C., Jean, P., Poulet, N., Roset, N., Copp, G. H.,
Geniez, P., Pont, D., Argillier, C., Baudoin, J. -M., Peroux, T., Crivelli, A. J., Olivier, A.,
Acqueberge, M., Le Brun, M., Møller, P. R., Willerslev, E. & Dejean, T. (2016). Next-
generation monitoring of aquatic biodiversity using environmental DNA metabarcoding.
Molecular Ecology, *25*, 929–942. https://doi.org/10.1111/mec.13428

Visconti, P., Pressey, R. L., Giorgini, D., Maiorano, L., Bakkenes, M., Boitani, L., Alkemade, R.,
Falcucci, A., Chiozza, F. & Rondinini, C. (2011). Future hotspots of terrestrial mammal loss.
Philosophical Transactions of the Royal Society B, *366*, 2693–2702.
https://doi.org/10.1098/rstb.2011.0105

Wickham, H. (2016). *ggplot2: elegant graphics for data analysis*. Springer, New York, USA.
http://ggplot2.org

Williams, K. E., Huyvaert, K. P., Vercauteren, K. C., Davis, A. J., & Piaggio, A. J. (2018). Detection
and persistence of environmental DNA from an invasive, terrestrial mammal. *Ecology and*
Evolution, *8*, 688–695. https://doi.org/10.1002/ece3.3698

Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A. & Smith, G. M. (2009) *Mixed effects models*
and extensions in ecology with R. Springer, New York, USA.

Table 1. Summary of focal species studied at wildlife parks and their lifestyle. The number of

individuals present and waterbody size in enclosures is provided.

Site	Species	Lifestyle	Enclosure	Number of individuals	Waterbody size (m^2)
Wildwood Trust	European otter (*Lutra lutra*)	Semi-aquatic	1	2	162
	European water vole (*Arvicola amphibius*)	Semi-aquatic	1	4	0.09
			2	1	0.09
	European beaver (*Castor fiber*)	Semi-aquatic	1	2	100
			2	1	100
	European hedgehog (*Erinaceus europaeus*)	Ground-dwelling	1	1	0.04
			2	2	0.04
	European badger (*Meles meles*)	Ground-dwelling	1	4	1.73
	Red deer (*Cervus elaphus*)	Ground-dwelling	1	8	100
	Eurasian lynx (*Lynx lynx*)	Ground-dwelling	1	2	2
	Red squirrel (*Sciurus vulgaris*)	Arboreal	1	2	0.01
			2	3	0.01
			3	3	0.01
			4	2	0.01
	European pine marten (*Martes martes*)	Arboreal	1	1	2
			2	1	0.375
Highland Wildlife Park	Red squirrel (*Sciurus vulgaris*)	Arboreal	NA	NA	0.25
	Eurasian lynx (*Lynx lynx*)	Ground-dwelling	1	8	2
	European beaver (*Castor fiber*)	Semi-aquatic	1	2	50
	Red deer (*Cervus elaphus*)	Ground-dwelling	1	30	NA

Figure 1. Heatmap showing proportional read counts for eDNA samples. Each square represents a sample taken from a wildlife park enclosure. Directed (DIR01-DIR06) and stratified (STR01-STR06) samples were collected from artificial waterbodies. Samples were collected from drinking containers (E1, E2, E3, E4, BOWL, BUCK), water vole (QUAR1, QUAR2) and RZSS Edinburgh Zoo beaver (ZOO) enclosures, and a water bath (BATH) in RZSS Highland Wildlife Park woods.

Figure 2. Relationships predicted by the binomial GLMMs between proportional read counts

and sample type nested within species **(a)** or species lifestyle **(b)**. The observed data (coloured

points) are displayed against the predicted relationships (black points with error bars) for each

species **(a)** or species lifestyle **(b)**. Points are shaped by sample type **(a)** or wildlife park **(b)**, and

coloured by species lifestyle. Error bars represent the standard error around the predicted

means.

Figure 3. Boxplots showing the mean proportional read counts for specific **(a)** and generic **(b)** behaviour(s) exhibited by focal species. Boxes show 25th, 50th, and 75th percentiles, and whiskers show 5th and 95th percentiles. Points are coloured by species lifestyle, and each point in **(a)** represents a directed sample sized by frequency of behaviour. The behaviour 'none' for beaver represents occurrences of beaver in water but out of view of camera traps.

757 **Figure 4.** Exemplar camera trap photos taken at natural ponds where focal species were

758 present. Red deer was recorded at Thorne Moors **(a)**, roe deer **(b)** and red fox **(c)** were

759 recorded at Tophill Low Nature Reserve, and beaver was recorded at the Bamff Estate **(d)**.

Figure 5. Tile plot showing species presence-absence at individual pond and site-level as indicated by field signs, camera trapping, and eDNA metabarcoding at sites with focal species presence confirmed by cumulative survey data.