The Out of East Asia model versus the African Eve model of modern human origins in light of ancient mtDNA findings

Ye Zhang and Shi Huang*

Center for Medical Genetics, Hunan Key Laboratory of Medical Genetics, School of

Life Sciences, Xiangya Medical School, Central South University, 110 Xiangya Road,

Changsha, Hunan 410078, P.R. China

*Corresponding author: Shi Huang

Keywords: mtDNA, ancient DNAs, Out of East Asia model, African Eve model, Out of Africa model, MGD theory, neutral theory, molecular clock

Abstract

The first molecular model of modern human origins published in 1983 had the

mtDNA phylogenetic tree rooted in Asia. This model was subsequently overlooked

and superseded by the African Eve model in 1987 that was premised on the

unrealistic infinite site assumption and the now failed molecular clock hypothesis. We

have recently developed a new framework of molecular evolution, the maximum

genetic diversity (MGD) hypothesis, which has in turn led us to discover a new model

of modern human origins with the roots of uniparental DNAs placed in East Asia.

While the African mtDNA Eve model has haplotype N as ancestral to R, our Asia

model places R as the ancestor of all. We here examined ancient mtDNAs from the

literature focusing on the relationship between N and R. The data showed that all

three oldest mtDNAs were R with the 45000 year old Ust'-Ishim a basal type and the

two ~40000 year old samples sub-branch of R. Among the numerous mtDNAs of

39500-30000 year old, most were R subtype U and only two were N samples, the

39500 year old Oase1 and the 34425 year old Salkhit. These N types are basal and

hence likely close to the root of N. These ancient DNA findings suggest that basal R is

~5000 years older than basal N, thereby confirming the East Asia model and

invalidating the African Eve model.

Introduction

Investigation into the question of human origins has a long history and two

competing models termed "Multiregional" and the "Recent Out-of-Africa" hypothesis

have been proposed [1]. In the Multiregional model [2-4], recent human evolution is seen

as the product of the early and middle Pleistocene radiation of *Homo erectus* from

Africa. Thereafter, local differentiation led to the establishment of regional populations

which evolved to produce anatomically modern humans (AMH) in different regions of

the world. *Homo* has been a single species since the genus first appeared in the fossil

record ~2.3-2.8 million years ago. The model has ample evidence from fossils and

Paleolithic cultural remains but consistent molecular evidence has been lacking until

very recently [5].

The single origin Recent Out of Africa model assumes that there was a relatively

recent common ancestral population for *Homo sapiens* which already showed most if

not all of the anatomical features shared by present day people. This population

originated in Africa ~200 ky ago, followed by an initiation of African regional

differentiation, subsequent radiation from Africa, and final establishment of modern

regional characteristics outside Africa [1,6]. These modern Africans replaced the archaic

Homo in Eurasia with limited genetic mixing [7-11]. Support for this model comes from

the African location of the earliest fossils of some AMH features [12,13] and the molecular

clock and neutral theory interpretation of the greater genetic diversity in Africans [6].

However, completely modern humans older than 20000 years have yet to be found in

Africa but did exist in Daoxian South China, although the Daoxian site has no

skeletons other than teeth, leaving it uncertain whether features other than teeth are

also completely modern [14,15].

While long overlooked, there is in fact a third model based on mtDNA analyses. In

1983, researchers derived the first mtDNA phylogenetic tree and rooted the tree in

Asia [16]. Unfortunately, this Asia model was superseded 4 years later by the African

Eve model without any valid reasons [6]. The African Eve model assumes the molecular

clock while the Asia model not. Given that the universal molecular clock is widely

acknowledged to be unreal [17-22], a reexamination of these models is clearly warranted.

Furthermore, the Africa model requires the neutral theory and the infinite site

assumption, which, while largely sound as a null model and a framework for

pre-saturation evolutionary processes, are not *a priori* valid and have met with great

difficulties as an explanatory framework for most molecular evolutionary phenomena

[23-27]. Obviously, inferring human origins by using genetic diversity data must wait until

one has a complete understanding of what genetic diversity means.

We have in recent years examined the longstanding genetic diversity riddle and

developed a more complete evolutionary account, the maximum genetic diversity

(MGD) hypothesis, that has been productive in addressing both evolutionary and

biomedical problems [21,28-36]. The MGD theory has solved the two major puzzles of

genetic diversity, the genetic equidistance phenomenon and the much narrower range

of genetic diversity relative to the large variation in population size [27,28]. The genetic

equidistance result of Margoliash in 1963 is in fact the first and best evidence for MGD

rather than linear distance as mis-interpreted by the molecular clock and in turn the

neutral theory [19-21,28,34,37,38]. The MGD regards the genome to be mostly functional and

under stabilizing selection. We have shown that genetic distances or diversities today

are mostly at optimum saturation no longer related to time of evolution. However, the

molecular clock and neutral theory misread reality of genetic diversity as still related to

time of evolution.

Based on the MGD theory, we have developed the "slow clock" method that only

uses slow evolving DNAs for demographic inferences because they are more likely to

be neutral and away from the saturation phase of evolution [5,34,39,40]. Our analyses

have led to a new scheme of modern human origins largely consistent with the

multiregional model in terms of autosomes where the first split among major human

groups is placed at ~1.9 million years ago [5]. Different from the multiregional model,

however, our new scheme has solved the origin problem of the modern uniparental

DNAs that obviously can only have originated from a single region as the coalescence

time of the modern mtDNA lineages is much shorter than 1.9 million years and as

archaic humans, such as Neanderthals, Denisovans, and Heidelbergensis, clearly

carried distinct lineages [41]. Our model placed the roots of both mtDNA and Y

chromosome in East Asia [5], which independently confirms the 1983 finding on the

Asia origin of mtDNA [16]. Our model furthermore specifically places the least

differentiated haplotype R0 or R* as the ancestor of all mtDNA haplotypes, which is in

contrast to the African Eve model that puts R downstream of haplotype N (Figure 1).

The R0 haplotype is most common today in the Southern Chinese group in the 1000

genomes project, implicating the origin of the modern mtDNA lineage in Southern

China [5]. Our mtDNA tree was derived from analyzing present day haplotypes without

needing to use ancient samples. Therefore, its validity can be independently tested by

using ancient DNAs.

Figure 1. mtDNA phylogenetic tree of the Out of East Asia model. Only major

braches and representative mutations are shown. Slow evolving sites (altering protein

and RNA sequences) in black and fast sites in red. Also shown are several

representative ancient DNAs that were placed based on their variant types. It should

be noted that tree topology was built without making use of any ancient DNAs.

The Africa model assumes the infinite site assumption to classify haplotype

relationships based on inference of ancestral and derived alleles with the allele

identical to that of chimpanzees as the ancestral. However, the infinite site model is in

fact unrealistic because there is overwhelming evidence of mutation saturation as

explained by the MGD theory, which makes the inference of ancestral alleles

meaningless. Actually, the Africa Eve phylogenetic tree cannot even meet the

minimum standard of a scientific model, i.e., self-consistency, as it contains huge

number of back mutations (reversions to an ancestral allele) that violate the infinite

site assumption required to build the Africa model in the first place. These mutations

can be found in the official mtDNA tree, phylotree (http://www.phylotree.org/), and are

designated by an exclamation mark (!) following the position numbers in phylotree.

The total number of these back mutations in phylotree is 1180 per our manual

counting. For example, of the 5 defining mutations for haplotype N, one at site 15301

is a back mutation.

Another premise (inferred from the molecular clock) for the Africa model is that

the original ancestor group should today show highest genetic diversity. However, as

the MGD theory explains, reality is that different groups may show different maximum

genetic diversity and human groups today are at saturation level of genetic diversity

for the fast evolving DNAs [5]. Africans do have the highest genetic diversity but only in

fast changing variants and deleterious variants such as stopcodon gain/loss variants[5].

They in fact do have the highest genetic diversity in slow evolving neutral variants [5].

Unlike the Africa model, our Asia model, as well as that of Johnson et al [16],

however, is based on the common sense that the original type should be the least

differentiated. The original ancestor should mostly carry major alleles at SNP sites

because as the ancestor population grew in size only a small fraction of the group

should carry alternative mutant alleles (mutations are rare events). Indeed, the 45000

year old Ust'-Ishim's mtDNA had fewer SNPs to begin with and the one he did have

are all common alleles in today's population (>67%) except one site 16150 [10]. The

Asia model also regards mutations in mtDNA to be functional, which has ample

evidential support [26,33,42,43], and classifies haplotypes based on sharing of alleles with

more weight on the slow evolving sites (altering protein or RNA sequences) [5]. Hence,

the Asia model is inherently more sound and self-consistent than the Africa model due

to stronger theoretical foundations and far more realistic assumptions. Nonetheless,

direct experimental test of these models is required to be absolutely certain. As our

mtDNA tree was not derived by making use of ancient DNAs, it can be independently

tested by using ancient samples. Recent discoveries in ancient DNAs have made

such test possible and we here found the Asia model strongly supported by the

ancient mtDNAs findings from the recent literature.

Results and discussion

The relationship between N and R is the most obvious difference between the

Africa and the Asia model. Of the three oldest mtDNAs, all turned out to be R, the

45000 year old Ust'-Ishim from Siberia [10], the 40328 year old Tianyuan from Northern

China [44], and the 39805 year old Fumane2 from Riparo Bombrini Italy [45]. Tianyuan

was haplotype R subtype B as reported. Fumane2 was closest to extant HV or H

haplotype (KP34013 and KF523402) per our analyses of the NCBI database. Only the

oldest R sample Ust'-Ishim was found to be basal, consistent with expectations. All

three had the expected mutations within R haplogroup (73G, 263G, 750G, 1438G,

2706G, 4769G, 7028T, 8860G, 11719A, 14766T and 15326G). But Ust'-Ishim had

only 1 extra mutation (16150T), Tianyuan had 6 extra mutations (a 9 bp deletion, 5348,

5836, 11257, 16189, 16293), and Fumane2 had 3 (5585A, 16129A and 16304C).

These extra mutations are mostly minor or rare alleles in present day people,

indicating that Ust'-Ishim is more basal and less differentiated than the other two

younger samples.

Among relatively younger samples of 39500-30000 years old, two N samples

were found, the 39500 year old Oase1 from Romania [9] and the 34425 year old Salkhit

from Mongolia [46]. In contrast to the rarity of N, numerous samples of R subtype U

were found in Europe as recently summarized by Deviese et al [46], indicating that R*

basal samples may only be expected near the time when the root of R lived, which

should be close to 45000 years ago as marked by the R* sample Ust'-Ishim. If R is

indeed the root of modern mtDNA, then the modern lineage should not be much older

than 45000 years old. In fact, while partially modern skeletons can be found in many

sites in Africa and Asia at 310000-80000 years ago [14,15,47-53], no human skeletons that

were fully modern for all key skeletal features have been found prior to 50000 years

ago [54]. As the demise of Neanderthals is thought to be associated with the arrival of

modern humans and occurred at ~40000 years ago [45], one can infer that the

appearance of fully modern humans may not be much earlier than 40000 years ago. If

it is much earlier, one would have the difficulty to explain the long lag before the

demise of Neanderthals.

Both the N samples were found to be basal N* and not directly ancestral to any

present day haplogroups [46]. Oase1 had all N defining mutations except 8701G and

9540C and 6 extra mutations (3205A, 3462T, 4232C, 7158G, 8749C, 11016A). Salkhit

had all N defining mutations and 10 extra mutations (4113A, 5492C, 8155A, 9456G

slow, 9509C, 10364A, 14578T, 15077A slow, 16169T, 16092Gap). These extra

mutations are also mostly minor or rare alleles in present day people. So, both of

these N* samples appear to be far more differentiated than the R* type of Ust'-Ishim.

As these two N* samples had ages of 39500-34425 years and no N subtypes were

found in the numerous mtDNA samples from 40000-30000 years ago, which is in

contrast to the abundantly found R subtype U, one can infer that the basal N*

ancestor probably lived at 39500-34425 years ago and cannot be much earlier than

39500 years ago or older than R*. It is therefore highly unlikely to detect the presence

of N in future samples older than 45000 years or the oldest R.

These two samples all have private mutations that are rarely found in today's

samples, indicating that those rare private mutations may represent adaptive changes

related to adaptation to ancient environments and may not be used to classify these

two samples as non-basal N. We note that Deviese et al [46] used the molecular clock

method to calculate a split time between Salkhit and basal N* to be 19000 years and

thus considered the appearance of N* basal type to be at ~50000 years ago. However,

such calculation may not be realistic as it did not consider the adaptive non-neutral

roles of the ancient variants. Variations in mtDNA are mostly adaptive or under natural

selection. Although distance among haplotypes may show some meaningful

correlation with time of separation, such correlation are all derived by using present

day haplotypes where the effects of variations due to adaption to environments from

different eras are minimized. But comparing ancient DNA with present day haplotypes

would represent a very different picture, because some variants in ancient DNAs may

represent adaptive changes specific to the ancient environment. Those variants could

be changed by today's environments rather than by neutral drift. Therefore, one

cannot use the molecular clock derived from present day samples to calculate the

split time between an ancient sample such as Salkhit and other haplotypes of either

today's or ancient. In addition, if basal N* is indeed 50000 years old, it would

contradict other findings. For example, one should find within the time window of

40000-30000 years ago differentiated N subtypes. And yet, only non-differentiated N

were in fact found (Oase1 and Salkhit) that do not belong to any known present day

haplotypes under N. In contrast, for a basal haplotype of truly 45000 years old such as

R* , nearly all R samples found for the time window of 40000-30000 years ago were

differentiated R subtypes such as U and B. Taken together, the combined data do not

support placing basal N* at >40000 years ago.

There were two M samples aged 34000-35000 years, Goyet Q116-1 and Goyet

Q376-3 from Belgium, found among samples older than 30000 years, and one M

sample of 28000 years old from La Rochette France [55,56]. As M is downstream of R

and N in the Asia model but is parallel to N in the Africa model, the rarity of M relative

to R and N in samples older than 35000 years is consistent with the Asia model but

not the Africa model. Also, if the Africa tree is true where N and M are sister branches,

the rarity of M in samples older than 35000 years ago further suggest that basal N*

could not be 50000 years old.

That R* haplotype was found to be 5000 year older than N* haplotype as

discussed above is consistent with the 2 SNP difference between R and N (Figure 1)

and the mtDNA mutation rate of 2.67×10^{-8} substitution per site per year (5000 x 16500

x 2.67×10^{-8} = 2.2 substitution) as recently used by others [46]. This suggests that the

ages of the oldest R* and N* samples found so far were likely to be close to the real

ages for the R and N ancestors. If the Africa model is true, the age of the original

African L0 ancestor of all modern humans would be 120000 years old (35 SNPs

separate L0 and N). However, as there are no completely modern humans found in

Africa that are more than 20000 years old, the classification of L0 lineage as the

original ancestor may be unrealistic. It is however placed as one of the last evolved in

the Asia model, where its highly differentiated feature is a result of hybridization of

AMH with archaic humans and coevolution of mtDNA with the admixed nuclear

genome [5]. It should be noted that the calculations here using molecular clock were all

dealing with present day haplotypes within the same environmental or time period.

Thus, effects of variants adaptive to environments of different eras are non-issues.

The original mtDNA type should leave more descendants today than the subtypes

because mutations are rare events and occur only to rare individuals in a large

population. We next asked whether the R haplotype is more popular today than other

types by examining the data collected by the mitomap database

(https://www.mitomap.org/MITOMAP). There are 64531 R samples in the database as

inferred by using the R defining SNP 16223C, representing 53.6% of the whole

collection. While the database may not be free of sampling bias and not all 16223C

carrying samples are R due to independent mutations (again violating the infinite site

assumption), one can nonetheless infer that it is highly likely for the R group to be

larger than any other haplogroup as it is found larger than all the other haplogroups

combined.

In contrast to the Asia model, the Africa model requires rare individuals with new

mutations to have a reproductive advantage over ancestral populations, hence

accounting for why the R group is much larger than any of its ancestor group, or why

L0 is smaller than the mutant group L1'2'3'4'5'6, why L1 is smaller than the mutant

group L2'3'4'5'6, why L5 is smaller than the mutant group L2'3'4'6, why L2 is smaller

than the mutant group L3'4'6, why L6 is smaller than the mutant group L3'4, and why

L4 is smaller than the mutant group L3. Such repeated reproductive advantages that

must be granted to each round of new mutations leading to a new haplotype are in

plain contradiction to the neutral mutation premise required to build the Africa model in

the first place.

In summary, the Out of East Asia mtDNA model is inherently more sound and

self-consistent than the African Eve model due to stronger theoretical foundations and

far more realistic assumptions. The original R and N are likely to have ages close to

those of the so far found oldest R* and N* samples, which showed that R* is older

than N* by 5000 years. These results plainly invalidate the African Eve model and

confirm the Out of Asia model. We expect our conclusion here to be further

strengthened by future ancient DNA studies.

Methods:

Sequence alignment and mismatches were performed using blastn at the NCBI

database. The number of back mutations in the mtDNA tree was inferred by counting

the exclamation marks (!) that designate the back mutations in the official mtDNA tree

phylotree (http://www.phylotree.org/). The number of R samples in the mitomap

database (https://www.mitomap.org/MITOMAP) were calculated by counting the

number of haplotypes having the R defining SNP 16223C.

Acknowledgments:

Supported by the National Natural Science Foundation of China grant 81171880,

the National Basic Research Program of China grant 2011CB51001 (S.H.).

Additional Information:

Competing Interests

The authors declare that they have no competing interests.

Author contributions

Y.Z. and S.H. devised the project, analyzed the data and wrote the manuscript.

References:

1   Stringer, C. B. & Andrews, P. Genetic and fossil evidence for the origin of modern humans. *Science* **239**, 1263-1268 (1988).

2   Thorne, A. G. & Wolpoff, M. H. Regional continuity in Australasian Pleistocene
hominid evolution. *Am J Phys Anthropol* **55**, 337-349 (1981).
3   Wolpoff, M. H., Wu, X. Z. & Thorne, A. G. *Modern homo sapiens origins: a general*
theory of hominid evolution involving the fossil evidence from east Asia.   411-483
(Alan R. Liss, 1984).
4   Wu, X. On the origin of modern humans in China. *Quaternary International* **117**,
131-140 (2004).
5   Yuan, D. *et al.* Modern human origins: multiregional evolution of autosomes and East
Asia origin of Y and mtDNA. *bioRxiv*, doi: https://doi.org/10.1101/106864 (2017).
6   Cann, R. L., Stoneking, A. C. & Wilson, A. C. Mitochondrial DNA and human evolution.
Nature **325**, 31-36 (1987).
7   Green, R. E., Krause, J., et. & al. A draft sequence of the Neandertal Genome. *Science*
328, 710-722 (2010).
8   Meyer, M. *et al.* A High-Coverage Genome Sequence from an Archaic Denisovan
Individual. *Science*, doi:10.1126/science.1224344 (2012).
9   Fu, Q. *et al.* An early modern human from Romania with a recent Neanderthal
ancestor. *Nature* **524**, 216-219, doi:10.1038/nature14558 (2015).
10   Fu, Q. *et al.* Genome sequence of a 45,000-year-old modern human from western
Siberia. *Nature* **514**, 445-449, doi:10.1038/nature13810 (2014).
11   Vernot, B. & Akey, J. M. Resurrecting surviving Neandertal lineages from modern
human genomes. *Science* **343**, 1017-1021, doi:10.1126/science.1245938 (2014).
12   Hublin, J. J. *et al.* New fossils from Jebel Irhoud, Morocco and the pan-African origin
of Homo sapiens. *Nature* **546**, 289-292, doi:10.1038/nature22336 (2017).
13   White, T. D. *et al.* Pleistocene Homo sapiens from Middle Awash, Ethiopia. *Nature*
423, 742-747 (2003).
14   Scerri, E. M. L. *et al.* Did Our Species Evolve in Subdivided Populations across Africa,
and   Why   Does   It   Matter?   *Trends   Ecol   Evol*   **33**,   582-594,
doi:10.1016/j.tree.2018.05.005 (2018).
15   Liu, W. *et al.* The earliest unequivocally modern humans in southern China. *Nature*
526, 696-699, doi:10.1038/nature15696 (2015).
16   Johnson, M. J., Wallace, D. C., Ferris, S. D., Rattazzi, M. C. & Cavalli-Sforza, L. L.
Radiation of human mitochondria DNA types analyzed by restriction endonuclease
cleavage patterns. *J Mol Evol* **19**, 255-271 (1983).
17   Ayala, F. J. Molecular clock mirages. *BioEssays* **21**, 71-75 (1999).
18   Ayala, F. J. On the virtues and pitfalls of the molecular evolutionary clock. *J Hered* **77**,
226-235 (1986).
19   Hu, T. *et al.* The genetic equidistance result, misreading by the molecular clock and
neutral theory and reinterpretation nearly half of a century later. *Sci China Life Sci* **56**,
254-261 (2013).
20   Huang, S. The genetic equidistance result of molecular evolution is independent of
mutation rates. *J. Comp. Sci. Syst. Biol.* **1**, 092-102 (2008).
21   Huang, S. The overlap feature of the genetic equidistance result, a fundamental
biological phenomenon overlooked for nearly half of a century. *Biological Theory* **5**,
40-52 (2010).

22 Kumar, S. Molecular clocks: four decades of evolution. *Nat Rev Genet* **6**, 654-662
(2005).
23 Hahn, M. W. Toward a selection theory of molecular evolution. *Evolution* **62**, 255-265,
doi:10.1111/j.1558-5646.2007.00308.x (2008).
24 Kreitman, M. The neutral theory is dead. Long live the neutral theory. *Bioessays* **18**,
678-683; discussion 683, doi:10.1002/bies.950180812 (1996).
25 Ohta, T. & Gillespie, J. H. Development of Neutral and Nearly Neutral Theories.
Theoretical population biology **49**, 128-142 (1996).
26 Ballard, J. W. & Kreitman, M. Is mitochondrial DNA a strictly neutral marker? *Trends
Ecol Evol* **10**, 485-488 (1995).
27 Leffler, E. M. *et al.* Revisiting an old riddle: what determines genetic diversity levels
within species? *PLoS Biol* **10**, e1001388, doi:10.1371/journal.pbio.1001388 (2012).
28 Huang, S. New thoughts on an old riddle: What determines genetic diversity within
and between species? *Genomics* **108**, 3-10, doi:10.1016/j.ygeno.2016.01.008 (2016).
29 Zhu, Z., Lu, Q., Wang, J. & Huang, S. Collective effects of common SNPs in foraging
decisions in Caenorhabditis elegans and an integrative method of identification of
candidate genes. *Sci. Rep.*, doi:10.1038/srep16904 (2015).
30 Zhu, Z., Yuan, D., Luo, D., Lu, X. & Huang, S. Enrichment of Minor Alleles of Common
SNPs and Improved Risk Prediction for Parkinson's Disease. *PLoS ONE* **10**, e0133421,
doi:10.1371/journal.pone.0133421 (2015).
31 Zhu, Z. *et al.* Collective effects of SNPs on transgenerational inheritance in
Caenorhabditis   elegans   and   budding   yeast. *Genomics*   **106**,   23-29,
doi:10.1016/j.ygeno.2015.04.002 (2015).
32 Biswas, K., Chakraborty, S., Podder, S. & Ghosh, T. C. Insights into the dN/dS ratio
heterogeneity between brain specific genes and widely expressed genes in species of
different complexity. *Genomics* **108**, 11-17, doi:10.1016/j.ygeno.2016.04.004 (2016).
33 Zhu, Z., Lu, Q., Zeng, F., Wang, J. & Huang, S. Compatibility between mitochondrial
and nuclear genomes correlates wtih quantitative trait of lifespan in Caenorhabditis
elegans. *Sci. Rep.*, doi:10.1038/srep17303 (2015).
34 Huang, S. Primate phylogeny: molecular evidence for a pongid clade excluding
humans and a prosimian clade containing tarsiers. *Sci China Life Sci* **55**, 709-725
(2012).
35 Huang, S. Inverse relationship between genetic diversity and epigenetic complexity.
Preprint available at Nature Precedings doi.org/10.1038/npre.2009.1751.1032
(2009).
36 Huang, S. *Histone methylation and the initiation of cancer, Cancer Epigenetics*.   (CRC
Press, 2008).
37 Margoliash, E. Primary structure and evolution of cytochrome c. *Proc. Natl. Acad. Sci.*
50, 672-679 (1963).
38 Luo, D. & Huang, S. The genetic equidistance phenomenon at the proteomic level.
Genomics **108**, 25-30, doi:10.1016/j.ygeno.2016.03.002 (2016).
39 Yuan, D. & Huang, S. On the peopling of the Americas: molecular evidence for the
Paleoamerican   and   the   Solutrean   models. *bioRxiv* **bioRxiv 130989; doi:**
https://doi.org/10.1101/130989 (2017).

40 Zhang, Y., Lei, X., Chen, H., Zhou, H. & Huang, S. Ancient DNAs and the Neolithic
Chinese super-grandfather Y haplotypes. *bioRxiv* **doi:**
https://doi.org/10.1101/487918 (2018).

41 Meyer, M. *et al.* A mitochondrial genome sequence of a hominin from Sima de los
Huesos. *Nature* **505**, 403-406, doi:10.1038/nature12788 (2014).

42 Teske, P. R. *et al.* Mitochondrial DNA is unsuitable to test for isolation by distance.
Scientific reports **8**, 8448, doi:10.1038/s41598-018-25138-9 (2018).

43 Towarnicki, S. G. & Ballard, J. W. O. Mitotype Interacts With Diet to Influence
Longevity, Fitness, and Mitochondrial Functions in Adult Female Drosophila.
Frontiers in genetics **9**, 593, doi:10.3389/fgene.2018.00593 (2018).

44 Fu, Q. *et al.* DNA analysis of an early modern human from Tianyuan Cave, China. *Proc*
Natl Acad Sci U S A **110**, 2223-2227, doi:10.1073/pnas.1221359110 (2013).

45 Benazzi, S. *et al.* Archaeology. The makers of the Protoaurignacian and implications
for Neandertal extinction. *Science* **348**, 793-796, doi:10.1126/science.aaa2773
(2015).

46 Deviese, T. *et al.* Compound-specific radiocarbon dating and mitochondrial DNA
analysis of the Pleistocene hominin from Salkhit Mongolia. *Nature communications*
10, 274, doi:10.1038/s41467-018-08018-8 (2019).

47 Hershkovitz, I. *et al.* The earliest modern humans outside Africa. *Science* **359**,
456-459, doi:10.1126/science.aap8369 (2018).

48 Groucutt, H. S. *et al.* Homo sapiens in Arabia by 85,000 years ago. *Nature ecology &*
evolution **2**, 800-809, doi:10.1038/s41559-018-0518-2 (2018).

49 Athreya, S. & Wu, X. A multivariate assessment of the Dali hominin cranium from
China: Morphological affinities and implications for Pleistocene evolution in East Asia.
Am J Phys Anthropol **164**, 679-701, doi:10.1002/ajpa.23305 (2017).

50 Zhao, L. *et al.* New discovery of human fossils and associated mammal faunas in Bijie
Guizhou. *Acta Anthropologica Sinica* **35**, 24-35 (2016).

51 Curnoe, D., Ji, X., Shajin, H., Tacon, P. S. C. & Li, Y. Dental remains from Longtanshan
cave 1 (Yunnan, China), and the initial presence of anatomically modern humans in
East Asia. *Quaternary International* **400**, 180-186 (2016).

52 Bae, C. J. *et al.* Modern human teeth from Late Pleistocene Luna Cave (Guangxi,
China). *Quaternary International* **354**, 169-183 (2014).

53 Li, Z. Y. *et al.* Late Pleistocene archaic human crania from Xuchang, China. *Science*
355, 969-972, doi:10.1126/science.aal2482 (2017).

54 Neubauer, S., Hublin, J. J. & Gunz, P. The evolution of modern human brain shape.
Science advances **4**, eaao5961, doi:10.1126/sciadv.aao5961 (2018).

55 Fu, Q. *et al.* The genetic history of Ice Age Europe. *Nature* **534**, 200-205,
doi:10.1038/nature17993 (2016).

56 Posth, C. *et al.* Pleistocene Mitochondrial Genomes Suggest a Single Major Dispersal
of Non-Africans and a Late Glacial Population Turnover in Europe. *Curr Biol* **26**,
827-833, doi:10.1016/j.cub.2016.01.037 (2016).

423