GPCRs show widespread differential mRNA expression and frequent mutation and copy number variation in solid tumors

Krishna Sriram[1], Kevin Moyung[1], Ross Corriden[1], Hannah Carter[2], Paul A. Insel[1, 2 *]
Departments of Pharmacology[1] and Medicine[2]
University of California, San Diego

*: Corresponding Author

Contact Information:

Email: pinsel@ucsd.edu
Phone: 858 534 2298

Mailing Address:
9500 Gilman Drive,
BSB 3076, Mail Code 0636
La Jolla, CA, 92093

Abstract:
G protein-coupled receptors (GPCRs) are the most widely targeted gene family for FDA-approved drugs. To assess possible roles for GPCRs in cancer, we analyzed The Cancer Genome Atlas (TCGA) data for mRNA expression, mutations, and copy number variation (CNV) in 20 categories/45 sub-types of solid tumors and quantified differential expression of GPCRs by comparing tumors against normal tissue from the GTEx database. GPCRs are over-represented among coding genes with elevated expression in solid tumors. This analysis reveals that most tumor types differentially express >50 GPCRs, including many targets for approved drugs, hitherto largely unrecognized as targets of interest in cancer. GPCR mRNA signatures characterize specific tumor types and correlate with expression of cancer-related pathways. Tumor GPCR mRNA signatures have prognostic relevance for survival and correlate with expression of numerous cancer-related genes and pathways. GPCR expression in tumors is largely independent of staging/grading/metastasis/driver mutations. GPCRs expressed in cancer cell lines largely parallels GPCR expression in tumors. Certain GPCRs are frequently mutated and appear to be hotspots, serving as bellwethers of accumulated genomic damage. CNV of GPCRs is common but does not generally correlate with mRNA expression. Our results suggest a previously under-appreciated role for GPCRs in cancer, perhaps as functional oncogenes, biomarkers, surface antigens and pharmacological targets.

List of abbreviations/acronyms
CAFs: Cancer associated fibroblasts; CCLE: Cancer Cell Line Encyclopedia; CNV/CNA: Copy number variation/ amplification; DE: Differential Expression; GPCR: G protein-coupled receptor; GTEx: Gene Tissue Expression Project [1] ; GtoPdb: IUPHAR/BPS Guide to Pharmacology [2]; MDS: Multi-dimensional Scaling; Nmut: Number of genes with somatic non-silent mutations per tumor genome; OE: Overexpression; TCGA: The Cancer Genome Atlas; Table 1 lists abbreviated cancer names.

Introduction

G protein-coupled receptors (GPCRs), the largest family of cell-surface receptors (>800 in the human genome), mediate the signaling of a wide variety of ligands, including hormones, neurotransmitters, proteases, lipids, and peptides. GPCRs regulate many functions (e.g., metabolism, migration, proliferation) and interactions of cells with their environment. GPCRs are also the largest family of targets for approved drugs [3,4], interacting with ~35% of FDA approved drugs, but are infrequently targeted in tumors other than endocrine cancers, even though a role for GPCRs has been implicated in features of the malignant phenotype [5,6]. One reason for their limited use is the notion that GPCRs are rarely mutated in cancer [7,8] although mutations occur in heterotrimeric GTP binding (G) proteins that GPCRs activate [8] and GPCRs regulate pathways, such as Wnt, MAPK and PI3K signaling, with mutations in cancer [9]. The biological relevance of GPCRs for the malignant phenotype and their high druggability imply that GPCRs might be an under-explored class of contributors to and targets in cancer.

To define the landscape of GPCRs in cancer, we undertook an integrated analysis of Differential Expression (DE), mutations, and CNV of GPCRs in 20 types of solid tumors (**Table 1, Supplementary Tables 1, 2**). Using RNA-seq data from TCGA and the GTEx database [1], we performed DE analysis of GPCRs in tumors compared to normal tissue, respectively, an analysis facilitated by the TOIL recompute project [10]. We studied GPCRs annotated by GtoPdb [2], including endoGPCRs (which respond to endogenous agonists) and taste receptors but not olfactory GPCRs for which such annotations are unavailable (**Supplement 2**). Our findings identify many differentially expressed GPCRs in solid tumors and corresponding cancer cell lines but a less important role for mutations and CNV. GPCRs with DE predict survival and are associated with expression of oncogenes and tumorigenic pathways. Overall, these results reveal a largely under-appreciated potential of GPCRs as contributors to cancer biology and as potential therapeutic targets. Our results from DE (N = 6224 individual tumors), mutation (N = 5103), and CNV analyses (N = 7545) are available as a resource at *insellab.github.io*.

FIGURE 1

Figure 1. Heatmaps of GPCR expression and differential expression (DE) in solid tumors

(**A**) For all 45 tumor subtypes, a heatmap showing the Log2 Fold-Change of GPCR expression in tumors compared to normal tissue (positive values indicate higher expression in tumors), with hierarchical clustering of GPCR genes to reveal patterns of DE.

(**B**) The number of GPCRs that show significant (FDR < 0.05) changes in expression compared to normal tissue among tumor types tested with large numbers of replicates (**Table 1**) and that correspond to the most lethal types of cancer.

(**C-D**) The GPCRs that most frequently (i.e., in most tumor types) show increases (**C**) or decreases (**D**) in expression among the 45 tumor subtypes

Figure 2. Solid tumors have large numbers of differentially expressed GPCRs compared to normal tissue

(**A-B**) The 30 highest expressed GPCRs in PDAC (**A**) or primary SKCM (**B**) and their corresponding expression in normal pancreatic (**A**) or skin (**B**) tissue.

(**C-D**) The 30 GPCRs with the highest fold-increase in expression in tumors compared to normal tissue for (**C**) PDAC and (**D**) Primary SKCM, sorted by fold-increase in tumors compared to normal tissue.

(**E-F**) For two highly expressed GPCRs in (**A-D**) as examples, the median expression of (**E**) GPRC5A and (**F**) GPR143 in all tumor types tested and corresponding normal tissue, normalized in CPM, allowing for comparison between tissue/tumor types. A lookup file that enables generation of similar plots (as well as upper and lower quartiles of expression) for any GPCR can be found at *https://insellab.github.io/gpcr_tcga_exp*.

FIGURE 2

Results

Differential expression (DE) of GPCRs in solid tumors compared to normal tissues

We focused on GPCRs with both substantial DE and magnitude of expression in solid tumors, i.e., 1) > 2-fold increase/decrease in DE in tumors compared to normal tissue, 2) FDR < 0.05 and 3) median expression in tumors > 1 TPM. We used the latter threshold for median expression in order to identify GPCRs that may be useful as therapeutic targets, for which higher expression is preferable. For DE analysis, we divided the 20 TCGA tumor types into 45 tumor subtypes (**Table 1**), based on histological classification of tumors in TCGA metadata. We found that different tumor subtypes within the same TCGA tumor classification have distinct GPCR expression, e.g., subtypes of breast cancer (BRCA), thyroid cancer (THCA) and esophageal cancer (ESCA) (**Figs S1A-F**).

Figure 1A shows a heatmap with fold changes (where statistically significant) for mRNA expression of all GPCR genes in solid tumors, relative to their corresponding normal tissue. Hierarchical clustering was performed on the GPCR genes, revealing three main clusters of GPCRs: a) those with frequent increases in expression across multiple tumor types; b) those with relatively little change; and c) GPCRs frequently reduced in expression in solid tumors compared to normal tissue. A phylogenetic tree identifying the GPCRs in each cluster is shown in **Fig S1**. Among the most lethal forms of cancer (in terms of annual deaths), **Figure 1B** shows that >25 GPCRs have >2-fold increased expression relative to normal tissue (with median expression in tumors > 1 TPM), whereas >20 GPCRs have significantly down-regulated expression, while remaining expressed > 1 TPM in solid tumors. Thus, large numbers of GPCRs that are expressed in solid tumors show DE, including tumor types that are most lethal. **Figure 1C** shows the 20 GPCRs that have >2-fold increased expression (with median expression >1 TPM) in the largest number of tumor types. **Figure 1D** shows the same, but for GPCRs frequently reduced >2-fold in expression, but which are still detected at >1 TPM in those tumors. Among the GPCRs with frequently increased expression are receptors likely expressed in the tumor cells themselves (e.g., GPRC5A, [11,12]) and expressed in the tumor microenvironment, such as in fibroblasts (e.g., F2R, [13]) and immune cells (e.g., FPR3, CCR1, CCR5).

Figure S3A-C shows DE for pancreatic ductal adenocarcinoma (PDAC) tumors (as an example) compared to normal pancreas. An MDS plot (**Fig S3A**) reveals clusters for tumors and normal tissue, implying distinct transcriptomic profiles. The more diffuse cluster of PDAC samples likely reflects their heterogeneity. Smear and volcano plots (**Figs S3B-C**) reveal many genes (>5000) with high, statistically significant DE (FDR <<0.05). **Figures S3D-F** show examples of genes (other than GPCRs) with high overexpression that prior studies implicated as having a role in PDAC. Multiple other tumor types also show expression of genes relevant to the malignant phenotype, cluster separately from their respective normal tissues, and have a large number of genes with DE, thus supporting the validity of our analysis.

Many GPCRs show DE in tumors, including those from each GPCR class: A (rhodopsin-like), B (secretin-like), C (metabotropic glutamate and others), frizzled and adhesion GPCRs. The highest expressed GPCRs in PDAC tumors (as an example, this finding is generalizable to other tumor types) are generally overexpressed compared to normal tissue and include orphan receptors (e.g., *GPRC5A* and *ADGRF4*/*GPR115*) and GPCRs with known agonists (e.g., *GPR68*) (**Fig 2A, C**). *GPRC5A*, the most highly expressed GPCR in PDAC, is 50-fold higher expressed; 95% of PDAC samples have >8-fold higher median *GPRC5A* expression than in normal pancreas (**Fig S3I**). Within a tumor type, a large majority of individual tumors express such overexpressed GPCRs at far higher levels than corresponding normal tissue (**Fig 3A**, e.g., GPRC5A); a subset of GPCRs are expressed in >90% of PDAC tumors at abundances greater than in any normal pancreas sample (**Fig 2F**). As discussed in subsequent sections, **Figures 3B, D-F** (and **Figs 7A-E**) indicate that GPCR expression is relatively consistent, ubiquitous among members of a given tumor cohort and largely independent of patient characteristics (e.g. tumor grade, pathological T and Sex, **Figs 3D-F**) that one might use to stratify patients. In addition, the GPCRs expressed couple to all major classes of Gα G protein signaling mechanisms (**Fig 3C**).

Similar to PDAC, numerous GPCRs are highly, consistently overexpressed in other tumor types. For instance, in skin cutaneous melanoma (SKCM) (**Fig 2B, D**); *ADGRG1/GPR56*, *GPR143* and *EDNRB* are highly overexpressed and highly expressed in magnitude compared to normal skin in >90% of melanoma samples (**Figs S3G-I**). In general, such highly overexpressed GPCRs are expressed in the vast majority, typically >90%, of samples within a tumor subtype.

Figure 2E shows the median expression of GPRC5A, the highest expressed GPCR in PDAC (and highly differentially expressed, **Figs 2A, C**) and highly expressed in many adenocarcinomas. **Figure 2F** shows this analysis across all tumor types and normal tissue for GPR143, the highest up-regulated GPCR in SKCM. For many GPCRs, we observe similar patterns of expression with pronounced up-regulation in tumors, as shown in **Figure 1**. To facilitate exploration of these data, we provide a spreadsheet-based tool (downloadable at *https://insellab.github.io/gpcr_tcga_exp*) wherein users can generate similar plots (along with upper and lower quartiles of expression) as in **Figure 2E-F** for any GPCR gene of interest. Of note, overexpression of certain GPCRs tends to be more prevalent within specific tumor types/subtypes than are common mutations. For example, *KRAS* and *TP53* are the most frequently mutated genes in PDAC (>70% and >60% of TCGA samples, respectively) but increased expression of multiple GPCRs occurs with greater frequency (**Fig 3B**). Each GPCR shown in **Figure 3B** shows statistically elevated expression in tumors compared to normal tissue, with FDRs << 0.05. The magnitude of DE and corresponding FDR for each GPCR shown are provided in **Supplement 3** and the project website.

Figure 3. Increased expression of certain GPCRs in PDAC is more frequent than that of KRAS and TP53 mutations, is independent of tumor grade, pathological T, and patient sex and includes GPCRs that couple to each of the 4 major G protein families.
(**A**) The expression of GPRC5A in all PDAC samples and normal pancreas tissues analyzed.
(**B**) Frequency of 2-fold increase and percent of TCGA-PDAC samples with higher maximal expression compared to normal pancreas of the indicated GPCRs with comparison of the frequency of mutations of KRAS and TP53, the most frequent somatic, non-silent mutations in PDAC tumors in TCGA.
(**C**) Changes in GPCR expression in PDAC alter the GPCR repertoire that couple to different G proteins.
(**D-F**) Tumor grade (D), Pathological T (E) and patient sex (F) do not impact on GPCR expression. The 30 highest expressed GPCRs in PDAC tumors are shown for each case; no statistically significant differences occur between the groups.

Figure 4. Grouping GPCRs and tumor types based on patterns of GPCR expression
(**A**) A phylogenetic tree showing the hierarchical clustering of GPCRs from Figure 1A reveals subsets of GPCRs that are either high- or low-expressed in solid tumors.
(**B**) Hierarchical clustering of types of solid tumors, based on their expression (in CPM) of GPCRs reveals clusters of tumor types, characterized by expression of particular GPCRs.

FIGURE 3

FIGURE 4

Magnitude of Expression (Log 2 TPM)

A

All GPCR genes

GPCRs that are generally low expressed in tumors, except for a few cases

GPCRs with very low expression in tumors

B

GPCRs with intermediate expression and for some tumors, prominently elevated relative to normal tissue

GPCRs widely expressed in multiple tumor types.

FIGURE 5

Figure 5. Hierarchical clustering of types of solid tumors, based on their expression (in CPM) of GPCRs reveals clusters of tumor types with expression of various GPCRs at characteristic levels of expression.

1 *A resource for exploring GPCR expression in tumors and normal tissue*
2
We compiled a list of GPCRs overexpressed in solid tumors with fold-changes and FDR along with expression in
TPM (for median expression and within-group comparisons of different genes) and CPM (for inter-group
comparisons of the same gene). The analysis revealed that 35 of 45 tumor types/subtypes show increased
expression of >30 GPCRs; 203 GPCRs are overexpressed in at least one type of cancer (**Supplementary Table
4; Supplement 3**), including 47 orphan GPCRs and >15 GPCRs that couple to each of the major G protein
classes. Increased expression of 130 GPCRs occurs in ≥4 tumor subtypes (**Supplement 3**). A subset of GPCRs
is overexpressed in many tumors, e.g., *FPR3* in 38 of the 45 tumor categories. **Supplementary Table 4** lists
other examples along with GPCRs that have reduced expression compared to normal tissue. Importantly, of the
203 GPCRs with increased expression in one or more tumors, 77 are targets for approved drugs. These include
ADORA2B, *CCR5* and *F2R*, which are overexpressed in 27, 27 and 26 tumor subtypes, respectively. **Table S4**
and **Supplement 3** provide further details regarding such druggable GPCRs.
14
Data generated and mined in this study, including DE analysis, re-normalized GPCR expression data, expression
of every GPCR in every individual tumor sample analyzed, accompanying CNV data and mutation data for
GPCRs are provided in the online resource at *https://insellab.github.io.* This resource also includes plots of GPCR
expression in tumors and corresponding normal tissue (such as those shown in **Figure 2A** and **2C)**, MDS scaling
plots showing the extent to which tumor and normal samples cluster together based on overall gene expression,
the aforementioned lookup tool to plot GPCR expression and information on GPCR expression in cancer cell
lines. In addition, high-resolution images of the heatmaps and phylogenetic trees in **Figures 1A, 4A-B** and **S1** are
available for download. Thus, while we have shown examples of GPCR expression and DE in certain tumor types
in this text, the resource website provides similar information for all GPCRs in all tumor types.
24
25 *Patterns of GPCR expression across solid tumors*
26
Figure 4A shows a heatmap that plots median GPCR expression in TPM across all tumor types, for all non-
olfactory GPCRs and reveals that GPCRs can be divided into four groups: a) those widely expressed at high
levels in all tumor types (often at >10 TPM) b) those with intermediate expression more broadly (~1 TPM), but
with high expression in specific tumor types c) GPCRs with generally very low expression (~O(0.1) TPM or less)
in most tumor types and d) GPCRs that are generally not detected in solid tumors. A phylogenetic (**Figure 4B**)
tree shows the identities of these groups of GPCRs. In general, many of the GPCRs in **Figure 4** that are widely
expressed in cancer are also widely differentially expressed (**Figure 1)**.
34
We also performed hierarchical clustering on tumor types to explore whether GPCR expression is distinct in
different subsets of solid tumors. **Figure 5** shows a phylogenetic tree of solid tumors, based on hierarchical
clustering of median GPCR expression in CPM (thereby allowing for comparisons between groups) in each tumor
type. Certain types of tumors cluster together, in a manner that corresponds more generally to their biology. For
example, nearly all adenocarcinomas cluster together and express a common set of GPCRs (**Figure 5)**. Similar
findings are observed for the other clusters. All tumor types within each group had a median expression of these
GPCRs ≥ 10 TPM, implying a common GPCR profile for adenocarcinomas compared to squamous cell
carcinomas or other groups highlighted. Such widely expressed GPCRs may be potential drug targets. Several of
these GPCRs commonly appear across multiple groups/clusters, but at different levels of expression. For
example, GPRC5A is widely expressed in both adenocarcinomas and squamous cell carcinomas but is typically
higher expressed in adenocarcinomas.
46
47 *GPCR expression is associated with cancer-related pathways and with survival: PDAC as an example*
48
49 A subset of GPCRs in PDAC is highly overexpressed and prominently expressed in tumors and in PDAC cells
50 (*vide infra*). Combining expression (normalized to median expression in PDAC) of 5 of the most highly DE GPCRs
51 (ADGRF1, ADGRF4, GPRC5A, HRH1 and LPAR5) yields a composite 'marker' whose expression positively

correlates with a subset of ~1200 genes (Benjamini-Hochberg adjusted p-values < 0.001) (**Fig S5A**). A reconstruction of the resulting network of genes using STRING [14] is shown in **Figure 6A**.

We conducted further analyses related to PDAC, including with GSEA [15] of the sets of negatively and positively associated genes (with respect to the five gene marker described above), pre-ranked/weighted by their FDRs and found an enrichment of a number of KEGG [16] pathways relevant to cancer (**Fig 6B**). The set of positively associated genes shows similar associations with cancer-related pathways when analyzed via GO [17,18] and Enrichr [19]. Enrichr also identifies, based on the Jensen compartment database [20], an enrichment of vesicle and exosome-related gene products among the set of positively correlated genes (**Fig 6C**). Network-analysis of the genes positively associated with the composite GPCR marker via STRING [14] provides an intuitive picture of this gene set (**Fig 6A**). Two 'clusters' of genes and pathways are evident; those associated with KRAS (including KRAS itself) and related processes (e.g., focal adhesion pathways) and a second group associated with regulation of cell cycle, cell division and differentiation. Expression of highly overexpressed GPCRs is positively correlated with one another and with expression of KRAS, implicating this GPCR subset as a PDAC signature. Leading edge analysis of the GSEA results confirmed that KRAS and other oncogenes are common elements in multiple enriched gene sets associated with this GPCR signature (**Fig 6D**). Survival analysis indicates that GPCR expression has prognostic relevance: patients with above-median expression of the five GPCRs had a ~200-day shorter survival compared to those with less than the median expression (**Fig 6F; Fig S3A)**.

We tested whether we could identify associations between GPCR expression in PDAC tumors and expression of markers for cell types commonly found in these tumors, thereby perhaps serving as an indicator for the cell types in the tumor microenvironment that express certain GPCRs. The cell types and markers we explored were: Tumor epithelial cells (E-Cadherin as a marker), cancer-associated fibroblasts (CAFs, Collagen1A1), endothelial cells (*VWF*, Von Willebrand Factor), the immune compartment (*CD45*), T-Cells (*CD3G*), myeloid cells (*CD33*) and macrophages (*CD14*). Pearson correlations were calculated between expression of each GPCR and of these markers, from which p-values were then calculated, followed by adjusted p-values using the Benjamini-Hochberg method so as to identify GPCRs associated with markers for each cell type. **Figure 6E** lists the most strongly correlated GPCRs for each cell type. For CAFs and cancer cells, these data appear in excellent agreement with prior experimental results. For example, *GPR68*/*OGR1* is strongly associated with CAFs in our analysis, consistent with evidence that it is a novel functional receptor in PDAC-derived CAFs [13]. Similarly, epithelial-enriched GPCRs (**Figure 6E)** are expressed in cancer cells [21] and as shown below, in cancer cell lines.

PDAC tumors have a subset of highly overexpressed (and highly expressed in terms of magnitude) chemokine receptors (CCR6, CCR7, CXCR3 and CXCR4) not expressed in PDAC cancer cells but likely associated with immune cells and their activation. Genes that correlated with expression of these GPCRs are involved with immune-associated processes, especially T-cell and B-cell related pathways (**Fig S4B**). Combined expression of these GPCRs is a positive predictor of survival (**Fig 6G**). The observation that GPCR expression may be a marker for survival is not unique to PDAC (**Fig 9D-F**): In SKCM, ESCA and LIHC, expression of individual GPCRs is associated with survival but combinations of such GPCRs are even better predictors of survival. GPCR expression may thus serve as a prognostic indicator in multiple tumor types.

The finding that GPCRs with high expression and DE in tumors show an association with tumorigenic pathways appears generalizable. For example, GPCRs that are highly expressed/overexpressed in other adenocarcinomas (**Figure 5**) are positively correlated with expression of genes from pathways similar to ones shown in **Figure 6A-C** for PDAC; i.e., focal adhesion, cell motility, cell cycle and division. By contrast, GPCR expression of solid tumor types from different branches of the cancer/GPCR phylogenetic tree shows an association with different pathways from those observed in adenocarcinomas. For example, GPR143, EDNRB and ADGRG1 are highly expressed and differentially expressed in SKCM and are adverse indicators of survival (**Figs S2F-H and Fig 9D**). Pathways enriched among genes that correlate with expression of the GPCRs are ones implicated in metastatic SKCM (**Figs S5 A-C**), such as transferrin transport [22], melanosome organization [23] and insulin receptor signaling

[24]. In general, highly expressed GPCRs in solid tumors show a positive correlation between GPCR expression and expression of tumorigenic pathways, implicating these GPCRs as potentially "functional oncogenes".

Functionality of overexpressed GPCRs

Evidence for functional roles in cancer cells of GPCRs that are highly expressed and overexpressed in solid tumors and cancer cells include findings for *PAR1/F2R* in breast cancer [25], gastric cancer [26], colon cancer [27] and melanoma [28] cells and for *PAR2/F2RL1* in melanoma [28], breast [29] and colon cancer cells [30]. Higher *PAR2* expression in ovarian cancer predicts poorer prognosis [31]. *EDNRB,* which is highly overexpressed in SKCM, promotes migration and transformation of melanocytes and melanoma cells and inhibition of *EDNRB* is pro-apoptotic [32,33]. *GPR143* promotes migration [34] and chemotherapeutic resistance [35] of melanoma cells. *GPR160* and *GPRC5A*, two frequently overexpressed GPCRs, are orphan receptors that influence the malignant phenotype. Knockdown of *GPR160* in prostate cancer cells increases apoptosis and growth arrest [36]. It has been suggested that *GPRC5A* is an oncogene that promotes proliferation, migration and colony formation of PDAC cells [11,12]. GPCRs with increased expression may thus be functional in cancer cells and activated by endogenous agonists or have constitutive activity that regulates signaling via heterotrimeric G proteins and/or β-arrestin [4]. At least certain of the many overexpressed GPCRs may thus serve as phenotypic drivers.

Incorporating omics analysis similar to what is presented here, our laboratory has recently shown [13] that *GPR68* (a proton-sensing GPCR) is highly overexpressed in PDAC tumors, in particular, in CAFs. We validated these data at the protein level and discovered that *GPR68* mediates symbiotic crosstalk between CAFs and PDAC cells and contributes to the tumor phenotype. Such findings provide an example as to how such omics data can identify overexpressed GPCRs with relevance to cancer cells themselves and to other cells in the tumor microenvironment.

Driver mutations, patient sex and stage/grade of tumors does not impact on GPCR expression

GPCR expression and DE is largely independent of tumor stage and grade. **Figure 3D** shows the similarity in GPCR expression for Grades 1 to 3 (G1 to G3) PDAC tumors. Median expression of GPCRs was also similar in PDAC tumors with different pathological T (**Fig 3E**). Similarly, Stage I and Stage IIIA BRCA IDC HR+ (Hormone Receptor positive) tumors have comparable GPCR expression and DE (**Fig 7D**).

GPCR expression appears largely independent of driver mutations, such as in BRCA HR+ IDC tumors with either *PI3KA* or *P53* mutations (**Fig 7A-C**); both groups have similar GPCR expression and DE of the same GPCRs compared to normal breast tissue. Similar results occur for LUAD and STAD that have or lack P53 mutations. Increased GPCR expression in solid tumors may thus not depend on specific driver mutations. Presence of highly overexpressed GPCRs may be a more ubiquitous feature of tumors than the presence of specific driver mutations, as exemplified by PDAC (**Fig 3B**) and in other tumor types with DE of numerous GPCRs (**Table 1**).

Moreover, GPCR expression also appears to be independent of a patient's sex. **Figure 3F** shows this for PDAC as a representative example, with the 30 highest expressed GPCRs in males and females. This finding appears to be generalizable to other tumor types as well. Thus, the elevated expression of highly expressed GPCRs in tumors appears to occur in nearly every patient within a tumor type and is independent of attributes such as sex, tumor progression and the mutations present. As discussed below, GPCR expression tends to be independent of CNV. Each tumor type thus expresses a repertoire of GPCRs that is largely, conserved among all patients with that type of cancer. This finding suggests the potential relevance of such highly expressed GPCRs as cancer drug targets.

GPCR expression is likely to be similar in metastatic and primary tumors

The SKCM gene expression dataset in TCGA has the most replicates of metastases. Primary and metastatic SKCM show similar expression and DE of GPCRs (e.g., *GPR143, EDNRB* and other highly expressed GPCRs) even though major differences occur in overall gene expression between primary and metastatic SKCM (**Fig 7E, F; S3F-H**). We found similar results for GPCR expression with primary and metastatic BRCA and THCA tumors and for recurrent and primary ovarian tumors (**Figs S6A-F**), though the number of replicates for each is small (<10). As new databases with large numbers of metastatic and primary tumor become available, we anticipate extending this analysis to strengthen this conclusion.

GPCRs highly expressed in tumors are highly expressed in cancer cells

We assessed RNA-seq data for GPCR expression in cancer cell lines from the EBI portal generated via the iRAP analysis pipeline [37] for cell lines in CCLE [38] and from Genentech [39]. The use of a different analysis pipeline than that used for TCGA data does not allow direct statistical comparisons of the datasets but confirms that most GPCRs in TCGA tumors are present in cancer cells and vice versa. We also mined RNA-seq for primary melanoma cells [40] and PDAC cells [41] from the NCBI GEO database. The data from these sources (**Methods, Section 1**) allow an approximate comparison with data for tumors. **Supplement 3** shows GPCR expression in cancer cell lines.

As an example, GPCRs with the highest median expression in TCGA PDAC tumors are generally highly expressed in PAAD (pancreatic adenocarcinoma, most likely PDAC) cell lines and patient-derived PDAC cells [41] (**Fig 7G**). A few exceptions exist, perhaps from effects of cell culture or expression by non-cancer cells in tumors. Even so, highly expressed GPCRs in PAAD cells are highly expressed in PDAC tumors (**Fig 7H**), findings that also occur in other tumors, such as SKCM (**Fig 7I, J**). Thus, most highly expressed GPCRs in tumors are also highly expressed in cancer cells and vice versa.

Most overexpressed GPCRs are rarely mutated

The most frequently mutated GPCRs in solid tumors are rarely overexpressed (**Figs 8A-B, 10H**) and conversely, highly overexpressed GPCRs in solid tumors are rarely mutated (**Fig 8A-B, Supplementary Table 5**). In SKCM, which has the highest mutation burden among TCGA tumor types, the most highly overexpressed GPCRs (*GPR143*, *EDNRB* and *GPR56*) are mutated in <2% of SKCM tumors whereas frequently mutated GPCRs (e.g., *GPR98,* mutated in nearly 40% of tumors) typically have low expression. The most frequently overexpressed GPCRs across all tumors (e.g., *FPR3*; **Table S5**) are mutated in <1% of all tumors surveyed, compared to frequently mutated GPCRs, e.g., *GPR98, GPR112*, which are mutated in >5% of all TCGA tumors surveyed. Thus, the frequency of GPCR mutations and likelihood of overexpression do not correlate (**Fig 8B**). The majority of GPCRs overexpressed in >20 tumor types/subtypes are mutated in < 50 samples out of > 5000 TCGA samples surveyed. Further, as discussed in following sections on GPCR mutation, mutations to these GPCRs are predicted to have no functional impact and are not enriched significantly for mutations at specific sites; thus, overexpressed GPCRs in tumors are not expected to be altered in their function by mutations.

Figure 6. GPCRs expression correlates with cancer-related pathways and predicts survival in PDAC

(**A**) Network analysis via STRING [14] of the genes positively correlated with expression of GPCRs highly expressed in PDAC, with FDR < 0.001, revealing distinct clusters of genes associated with specific cancer-related pathways.

(**B**) KEGG [16] gene sets with positive enrichment among genes most significantly positively or negatively correlated with expression of the identified subset of overexpressed GPCRs, based on GSEA pre-ranked analysis [15].

(**C**) Enrichr [19] analysis of the genes positively correlated in (**A**), with FDR < 0.001, based on their enrichment in cellular compartments, from the Jensen compartment database [20].

(**D**) Leading-edge analysis of GSEA results from (**B**), showing numerous cancer-related genes, including KRAS, which are common to multiple enriched KEGG genes sets.

(**E**) GPCRs in PDAC show a positive, statistically significant association (FDR < 0.001) between their expression and that of markers for various cell types as shown. The top 10 GPCRs associated with each cell type are highlighted, along with the strength/significance of these correlations.

(**F, G**) Kaplan-Meier curves for the impact of combined, normalized expression of subsets of GPCRs on PDAC patients. Total number of patients = 141; 59 patients were censored due to inadequate follow-up. (**F**) Impact of highly expressed GPCRs on median survival: 652 days (if below median expression) and 470 days (if above median expression) and for (**G**) Impact of highly expressed immune-related GPCRs on median survival: 460 days (below median expression) and 603 days (above median expression).

FIGURE 6

FIGURE 7

Figure 7. GPCR expression and presence of driver mutations and the similarity in GPCR expression of primary tumors, metastases and cancer cells derived from the tumors.

(**A**) Correlation of median expression of GPCRs in P53 mutant and PI3K mutant HR-positive BRCA IDC tumors

(**B**) Median expression of the 25 highest expressed GPCRs in P53-mutated tumors compared to expression in PI3K mutant HR-positive BRCA IDC tumors

(**C**) Fold-changes of GPCRs in P53 mutant and PI3K mutation HR-positive BRCA IDC tumors compared to normal breast tissue.

(**D**) Fold-changes of GPCRs in Stage 1 and Stage 3 ER-positive BRCA IDC tumors over normal breast tissue.

(**E**) Expression of the 25 highest expressed GPCRs and (inset) correlation of median GPCR expression between primary and metastatic SKCM.

(**F**) Gene expression of primary and distant metastatic SKCM tumors cluster differently and have large numbers of DE genes (**Supplement 3**)

(**G**) Median expression of highest expressed GPCRs in PDAC tumors compared to cancer cells (CCLE [38], n = 33; Genentech [39], n = 16; Witkiewicz et al., [41], n = 72).

(**H**) Median expression of highest expressed GPCRs in CCLE PDAC cell lines compared to other cell lines (as in panel **G**), primary cells and PDAC tumors.

(**I**) Median expression of highest expressed GPCRs in SKCM tumors compared to cancer cells (CCLE [38], n = 45; Genentech [39], n = 44; Müller et al., [40], n = 29).

(**J**) Median expression of highest expressed GPCRs in CCLE SKCM cell lines compared to other cell lines (as in panel **I**), primary cells and SKCM tumors.

Figure 8. Frequent overexpression of GPCRs in tumors does not correlate with frequency of mutation

(**A**). The number of tumors with GPCR mutations and of tumor types/subtypes in which the same GPCR is overexpressed for the 10 most frequently mutated GPCRs in TCGA tumors surveyed and the 10 most frequently overexpressed GPCRs.

(**B**) The frequency of overexpression compared to the frequency of mutations.

FIGURE 8

The predicted association of expressed GPCRs with Gα G proteins

Supplement 2 shows GPCRs annotated in the IUPHAR / BPS Guide to Pharmacology GPCR database [2], their signal transduction via G protein heterotrimers and if they are orphan GPCRs. Tissues and tumors typically express >150 GPCRs (at detection thresholds >0.1 TPM) that couple to the major types of G proteins (Gs, Gi/o, Gq/11, G12/13), most frequently Gi/Go and Gq/G11 (**Figs S6A-B**). We calculated the abundance of GPCRs that couple to each G protein by summing median GPCR expression (TPM), thereby yielding an expression 'repertoire' for each signaling mechanism (**Figs S6C-D**). Gs-coupled GPCRs typically account for the smallest GPCR expression repertoire for which such coupling is known. **Supplement 3** provides GPCR expression and G protein linkage data for normal tissues and solid tumors. Summing expression (TPM) of all GPCRs provides an estimate of the proportion of GPCRs among total mRNA (**Figs S6E-F**).

The GPCR expression repertoire varies among tissues in terms of total expression, number and identities of GPCRs. Tumors typically have a different GPCR repertoire than normal tissue, with increased or decreased expression of many GPCRs (**Fig 1C**, **Figs S6E-F**). Total GPCR expression and the number of GPCRs above expression thresholds increases in certain tumors (e.g., PDAC) but decreases in others (e.g., LIHC and SKCM) compared to normal tissue. Tumors also differ from normal tissue with respect to the abundance of GPCRs that couple to different G proteins (**Fig 1I, Figs S6C-D)**, suggesting changes in signaling. For example, Gs-coupled GPCR expression decreases in many tumors (e.g., PDAC, **Fig 3C**), implying decreases in cAMP signaling.

GPCRs as potential therapeutic targets in cancer

Among the >200 GPCRs overexpressed in at least one of the 45 tumor subtypes, 77 are targets for drugs approved by the FDA and/or EMA. (**Supplement 3**). Among these GPCR drug targets, >50% are overexpressed in 4 or more tumor subtypes (**Fig 9A**) and 15 GPCRs are increased in expression in 10 or more tumor subtypes. These results highlight the potential of GPCRs as targets in cancers and importantly, for the possible repurposing of drugs approved for other indications. Among the 77 GPCRs with increased expression in tumors, nearly two thirds link to either Gs- or Gi-coupled signaling and thus are predicted to regulate cAMP formation (**Fig 9B**).

Of the solid tumor types we analyzed, lung, colon, pancreatic, breast, and prostate cancers account for the largest annual number of deaths in the U.S. (https://www.cancer.gov/types/common-cancers). **Figure 9C** shows currently druggable GPCRs with increased expression in subtypes of those tumor types. Approved drugs target at least 10 GPCRs that have increased expression in those tumors. Hierarchical clustering of GPCR expression in different tumor types (based on their median expression of all GPCRs) reveals that GPCR expression distinguishes tumor types into groups that are consistent with other molecular/physiological traits (**Fig 5**). As discussed in sections above, GPCR expression appears to characterize categories of tumors and certain GPCRs may be targets across tumor classes/families.

As a protein-coding family of genes, GPCRs are disproportionately enriched among overexpressed genes in solid tumors, when compared to all protein-coding genes. Evidence for this was obtained as follows: In each tumor type indicated (**Fig 9G-H**), the ratio of number of coding genes with increased expression (above a prescribed threshold) over the total number of differentially expressed genes present was computed for a) GPCRs only; b) all coding genes. Fischer's exact test was used to verify if overrepresentation of GPCRs among genes with increased expression is significant ($p < 0.05$). Data shown are for coding genes with >4 fold increased expression in solid tumors, i.e., highlighting genes with drastic increases in expression. We found that many tumor types have 2-fold or greater overrepresentation of GPCRs among coding genes with large increases in expression.

Moreover, extending from the data shown in fig (**6F-G**), we find that many GPCRs expressed by solid tumors may be prognostic markers. GPCR mRNA expression is associated with differences in survival in multiple tumor types (**Fig 10**). Using GPCR expression normalized in CPM, we performed survival analysis using the modified Peto-Peto test for every GPCR in tumor types where sufficient numbers of TCGA replicates were available (typically

>60 replicates, allowing for >30 samples in each group for survival analysis). We omitted tumor types such as TGCT (Testicular cancer) for which almost no fatalities were recorded in the metadata. For certain tumor types with large numbers of replicates (e.g. OV, LUAD, LSQC etc), we have also provided alternative survival analysis on the related website omitting patients who are >75 years old at time of diagnosis, to reduce the risk of other sources of mortality confounding the analysis. This analysis yielded many GPCRs with a statistically significant impact on survival of each tumor type. Combined expression, i.e. a mean-normalized sum of expression, of GPCRs that individually predict survival and/or are highly differentially expressed yield stronger composite markers with prognostic relevance. **Figure 10A-C** shows examples of such combined markers in LSQC (**A**), LUAD (**B**), SKCM (**C**), which demonstrate strong adverse effects on survival. Certain tumor types, in particular KIRP, KIRC, LSQC and LUAD, express individual GPCRs that have statistically significant associations with survival (**Fig 10D**).

For two tumor types as examples (ESCA Adenocarcinoma and SKCM distant metastases, **Fig 10E-F**), we show the difference in mean survival times (in days) between patients with high expression (above median) and low expression (below median); negative values imply that elevated expression of these GPCRs is associated with adverse survival rates. GPCR expression is associated with both negative and positive survival effects, depending on the tumor type and GPCR. Several GPCRs are associated with differences in survival of >1 year. Some GPCRs have low median expression (<1 TPM) in the tumor population in general (e.g., taste receptors *TAS2R14* and *TAS2R20* in ESCA) but may be expressed in subsets of these populations, wherein they may contribute to differences in survival. Our analysis focusses on higher expressed GPCRs but we provide (at the accompanying website) expression and DE data for all GPCRs regardless of expression level, since in certain tumors very low-expressed GPCRs may have disease relevance in subsets of patients within a cancer cohort.

In total, we found that 301 GPCRs show statistically significant evidence (p <0.05) of an association with survival, in at least one tumor type/subtype (among 20 where such analysis was feasible). A subset of 32 GPCRs was significantly associated with survival in 5 or more of these tumor types/subtypes. All associations of GPCRs with survival in the tumor types tested are available at the accompanying website.

Figure 9. Solid tumor-expressed GPCRs with potential as drug targets
(**A**) The number of GPCRs that are targets for approved drugs and have increased expression in 1-3, 4-9 or ≥10 tumor subtypes.
(**B**) The linkage to G proteins of the 77 GPCRs targeted by approved drugs and with increased expression in at least 1 tumor subtype. Note: multiple GPCRs couple to more than one G protein.
(**C**) The number of GPCRs targeted by approved drugs that show increased expression in lung, colon, pancreatic, breast, and prostate cancers, the leading causes of cancer deaths in the U.S.
(**D**) Overrepresentation of GPCRs among genes with >4-fold elevated expression (FDR < 0.05) for the indicated tumor types/subtypes with p-value calculated via Fischer's exact test
(**E**) The magnitude of overrepresentation (relative enrichment) of GPCRs corresponding to the p-value in (**H**)

FIGURE 9

Figure 10. GPCR expression is associated with survival in many types of solid tumors
(**A-C**) Kaplan-Meier survival curves in the indicated tumor types, for weighted, combined expression of the
indicated GPCRs
(**D**) In the indicated tumor types, the number of GPCRs with significant (p < 0.05) association with survival
(**E-F**) As examples, the impact on mean survival of GPCRs in (**E**) ESCA adenocarcinoma and (**F**) SKCM (distant
metastases)
7
8

FIGURE 10

Somatic mutations of GPCRs in solid tumors

Analysis of 5103 TCGA samples in 20 tumor types (**Table S1**; 21 tumor types if one divides ESCA into esophageal adenocarcinoma and squamous cell carcinomas) revealed many GPCRs with frequent non-silent mutations (**Fig 11A, S8A**), including a more frequently mutated subset (**Fig 11A, inset**). *GPR98*/*ADGRV1,* the most frequently mutated GPCR, occurs in >8% of TCGA samples. Tumor types with high mutation burdens have a high frequency of GPCR mutations (**Fig 11B**). SKCM has the highest frequency: ~40% SKCM tumors have *GPR98* mutations (**Fig 11C, H**). Approximately 65% of tumors have ≥1 non-silent GPCR mutation. Certain GPCRs are mutated in >10% of specific tumor types (**Fig 11C**).

Nmut, the number of genes with somatic non-silent mutations per tumor genome, and the number of mutated GPCRs scale linearly in individual tumors (**Figs 11D, S8E-G**). Frequently mutated GPCRs (e.g., *GPR98*, *GPR112*, *and BAI3*) are more likely to be mutated as Nmut increases (**Fig 11E,** SKCM as an example). The relationship between Nmut and likelihood of GPR98 mutation is similar in SKCM and other cancers (**Fig 11F**); this is also observed for other frequently mutated GPCRs. Hence, the likelihood of a GPCR being mutated appears to depend on the accumulation of genome damage and to be independent of the mechanisms for the mutations. The relationship between GPCR mutation rates and Nmut is identical in BLCA, LUAD and SKCM, although the factors driving DNA damage and oncogenesis are likely different. Mutations of certain GPCRs, such as GPR98, may thus serve as a bellwether for genome-wide DNA damage.

Missense mutations and in-frame deletions are the most frequent non-silent mutations in GPCR genes (**Fig S8C-D, Table S3**). Mutations in frequently mutated GPCRs occur at many sites (**Fig S9A**), which contrasts with the smaller number of such sites in common oncogenes, e.g., *KRAS* [9]. (**Fig S9B**). Certain GPCR genes (e.g., *GPR98*) may be in genomic regions vulnerable to dysregulation of DNA damage and repair and belong to a subset of mutated genes; *GPR98* mutations frequently occur alongside other frequently mutated genes such as *TTN* and *MUC16* (**Fig S10A-G**). *GPR98* is among the 25 most frequently mutated genes in all tumor types surveyed; its mutational frequency is similar to that of genes (e.g., *BAGE2* [**Fig S8B]**) that are mutational hotspots [42]. As the GPCR with the largest gene length (~19,000 bp), *GPR98* has more mutational events. Compared with other very long genes, e.g., genes > 10,000 bp, *GPR98* belongs to a subset of ~20 genes with high mutational frequencies (**Fig 12A**), implicating *GPR98* as a hotspot for both silent and non-silent somatic mutations. *GPR98* has a >4-fold increased density of mutational events (normalized for gene length) compared to the average of these very long genes. One obtains a similar result by assessing the density of somatic mutations across all genes, irrespective of length (**Fig 12B**). GPR98, GPR112 and BAI3 are among the top 5% of genes in number of mutations per unit gene length, highlighting these genes as chromosomal regions with higher than normal rates of somatic mutation.

Survival analysis of metastatic SKCM samples was performed in order to evaluate the impact on tumors of somatic non-silent mutations to GPR98, GPR112, or other frequently mutated GPCRs. Analysis of Kaplan-Meier survival curves using the modified Peto-Peto method reveals that presence of somatic mutations in these GPCRs does not have a statistically significant impact on survival. **Fig 12C-D** shows this for GPR98 and GPR112, the two most frequently mutated GPCRs in SKCM. We find the same result in other tumor types as well and thus conclude that somatic non-silent mutations to GPCRs have no impact on patient survival.

Most mutated GPCR genes have low levels of mRNA expression (**Fig 8; 11G [SKCM as an example])** so may not be functionally relevant but certain GPCR genes (e.g., *CELSR1* and *LPHN2*/*ADGRL2*) are frequently mutated and moderately/highly expressed. Several such GPCRs are orphan receptors (without known physiologic agonists or roles), for which it is unclear if they impact on cell function. As cell-surface receptors, frequently mutated, well expressed GPCRs may represent neo-antigens. For SKCM, which has the most GPCR mutations among tumors types surveyed, DE analysis of primary melanomas and distant metastases that have or lack GPCR mutations (e.g., *GPR98* and *LPHN2*) revealed little evidence that these mutations alter the tumor transcriptome, implying that such GPCR mutations are likely passenger, rather than driver, mutations (**Fig 11H**;

Fig S8C, D). Conversely, previous work has suggested that for known oncogenes (e.g., for TP53, [43]), there are often widespread transcriptomic changes associated with specific mutations. We found similar behavior for other tumors (e.g., BLCA) that have frequent GPCR mutations.

As a further approach, we evaluated GPCR mutations, predicting the likelihood of functional consequences and site-specific enrichment of the mutations via MutSig 2CV v3.1 (*gdac.broadinstitute.org*). The majority of GPCRs frequently mutated (**Fig 11-I**, SKCM as example) show non-silent mutations that are non-significant in terms of enrichment (compared to the background mutation rate of silent mutations over the same regions) for individual mutation sites. These mutations are not predicted to be functional (calculated from estimations of functional impact of mutations based on whether mutated regions are highly evolutionarily conserved) by MutSig 2CV, consistent with the idea that the frequent GPCR mutations are likely passenger and not driver mutations. MutSig 2CV results for all GPCRs in all tumor types are available at the accompanying website.

FIGURE 11

Figure 11: Somatic non-silent mutations of GPCRs
(**A**) Frequency of GPCR mutations in the TCGA cohort (n = 5103). Inset: 20 most frequently mutated GPCRs.
(**B**) The average number of all genes (red) and GPCRs (blue) with somatic, non-silent mutations per tumor
genome for the TCGA tumor types surveyed
(**C**) The number of mutated GPCRs per tumor for several types of solid tumors in TCGA (black), the proportion of
samples in each tumor type with at least one mutated GPCR (orange) and proportion with non-silent mutations for
the most commonly mutated GPCR (gray) for each tumor type.
(**D**) GPCR mutation frequency is linearly related to Nmut in SKCM.
(**E**) Probability of GPCR mutation as Nmut increases in SKCM for the 10 most frequently mutated GPCRs
(**F**) Normalized probability of GPR98 mutation as Nmut increases in SKCM, and the same for several cancers
with high mutational burden and combined for BLCA, LUAD, LSQC, COAD, and SKCM.
(**G**) The 20 most frequently mutated GPCRs in primary SKCM, with frequency of mutation and median (and upper
and lower quartile) expression in TPM
(**H**) Expression (CPM) of the 5000 most abundant genes in SKCM correlate closely in primary SKCM tumors that
have or lack *GPR98* mutations.
(**I**) MutSig 2CV v3.1 scores in SKCM obtained from https://gdac.broadinstitute.org/, showing the q-values for the
significance of mutation scores for each annotated coding gene (blue); GPCR (black) and for those GPCRs, the
number of mutation events (red) among the SKCM cohort.
19
20 **Figure 12: Mutated GPCRs show high density of mutations but do not impact survival**
21 (**A-B**) Number of somatic (silent and non-silent) mutations per unit of gene length (total length of all exons per
22 genes) for (**A**) all genes and (**B**) genes >10 kb in length in SKCM.
23 (**C-D**) Survival analysis in metastatic SKCM for (**C**) GPR112 and (**D**) GPR98 mutations, with p-values calculated
24 using the Peto & Peto modification of the Gehan-Wilcoxon test.
25

FIGURE 12

Figure 13. Copy number variations (CNVs) of GPCRs in solid tumors
(**A**) The number of solid tumors with CNV for each GPCR across all TCGA samples (plotted in descending order of frequency for high-level amplification and homozygous deletions; see text for definition of high- and low-level amplification)
(**B**) The same as in (**A**) for low-level amplification and single-copy deletions
(**C**) In SKCM tumors (n = 367), the distribution of high-level amplification of GPCRs (n = 390 genes) compared to that of all protein coding genes (n = 24,776)
(**D**) The total number of homozygous deletions, single copy deletions, low-level and high-level amplifications for all GPCRs combined in the 7545 TCGA tumors surveyed for CNV
(**E**) The most frequently amplified GPCR for each TCGA tumor type and the proportion of samples with High and Low-level amplification for that GPCR
(**F**) Proportion of samples of various tumors types with High and Low-level amplification of *GPR160*, the most frequently amplified GPCR overall)
(**G**) Ovarian cancer samples with and without High-level amplification of GPR160, with the median for each group indicated. The difference between groups was not statistically significant)
(**H**) The risk ratio of elevated GPR160 expression (above the median value for OV) when GPR160 also shows high-level amplification; amplification of GPR160 increases the likelihood that GPR160 expression is elevated
(**I**) For the 30 GPCRs in OV with the highest fold-increase in expression relative to normal ovarian tissue (with FDR < 0.05 and median expression in OV > 1TPM), the corresponding number (of 579) OV tumors with high-level amplification of those same GPCRs
(**J**) The same as (**I**), but comparing fold-increase against low-level amplification

FIGURE 13

Copy-number variation (CNV) of GPCRs in solid tumors

CNV of certain GPCRs occurs frequently in TCGA solid tumor samples (**Fig 13A-B; Table S1**), with some GPCRs (e.g., *GPR160*) amplified in >5% of all TCGA samples surveyed. CNV data were obtained as GISTIC 2.0 [44] thresholded data, wherein values of -2, -1, 0, 1 and 2 respectively denote homozygous deletion, single copy deletion, diploid copy number, low level amplification (i.e. increase of 0.1 to 0.9 of copy number, expressed as a log2 ratio) and high-level amplification (amplification of >0.9 of the log2 ratio, i.e. >1.7 extra copies in a diploid cell) [45]. The distribution of amplification events among GPCRs parallels that of all genes (**Fig 13C,** SKCM as an example) but a subset of GPCRs is disproportionally amplified (**Fig 13D-F, and Supplement 3**). Most GPCRs have infrequent amplification (in 2% of tumors or less). Amplification does not predict high expression or overexpression of GPCRs (**Fig 13G-J**; OV as an example): frequently amplified GPCRs in tumors often have limited mRNA expression in those tumors while most highly expressed, overexpressed GPCRs are not amplified.

Single-copy/heterozygous deletions of GPCRs are widespread, whereas homozygous deletions are rare (**Fig 13A, D**). GPCR genes with single-copy deletions are generally not significantly expressed in tumors or normal tissues, implying that such deletions lack functional effects but exceptions exist. *PTH1R*, frequently deleted in KIRC (~77% samples have single-copy deletions), has ~10-fold reduced expression compared to normal kidney tissue. Similarly, *ADRA1A* is frequently deleted and has reduced expression in hepatocellular and prostate adenocarcinomas.

Figure 13E shows the identity/frequency of amplification of the most frequently amplified GPCR in each tumor type. Several cancers (e.g., OV and LSQC) have a high level of amplification of specific GPCRs in >25% and low-level amplification in >40% of samples. **Figure 13F** shows the CNA frequency of *GPR160*, the most frequently amplified GPCR overall, among all tumors surveyed. Except for *GPR160* and *FZD6*, most GPCRs with frequent amplification are rarely overexpressed (**Supplement 2**). CNA alone thus does not generally predict increased mRNA expression in tumors compared to normal tissue and highly expressed GPCRs in tumors are typically not amplified. We tested for correlation between high-level amplification and increased mRNA expression. **Figure 13G** shows GPR160 expression in OV tumors with and without high-level amplification; CNA is not a prerequisite for high mRNA expression and the small difference in median expression is not statistically significant, based on DE analysis via edgeR. However, tumors with amplification of GPR160 show a higher likelihood (~33%, **Fig 13H**) of expressing GPR160 at levels above the median for OV. We did not observe statistically significant risk ratios relating GPCR expression with amplification for other frequently amplified GPCRs. We conclude that CNA and GPCR mRNA expression are generally poorly correlated; hence, examination of amplified GPCR genes does not predict which GPCRs are highly and/or differentially expressed in a tumor (**Fig 13I, J**).

Supplements 2 and 3 provide, respectively, the frequency of GPCR CNV and changes in expression of GPCRs in the tumors surveyed. The widespread CNV of certain GPCRs suggests that they (and/or neighboring genes that vary along with these GPCRs) contribute to the malignant phenotype and may be markers for malignancy.

Discussion and Conclusions

In this study, we identified mutations, CNVs and alterations in mRNA expression of GPCRs in a range of solid tumors. The results reveal a broad landscape of changes, suggesting a functional role for this gene superfamily in such tumors and possible therapeutic utility of GPCRs in a large number of solid tumors.

Mutations of certain GPCRs have been implicated in cancer [8], but a comprehensive analysis of GPCR amplification, expression and DE has been lacking. The largest public datasets of normal (GTEx) [1] and cancer tissues (TCGA) provide RNA-seq data analyzed/normalized differently, making it difficult to compare datasets. For many tumors types, few replicates of 'normal' TCGA tissue are available and samples from matched non-tumor tissue from patients may not be representative of normal tissue (**Supp. Methods, Section 1**). The TOIL project enabled comparison of TCGA and GTEx data with analysis of both RNA-seq datasets via the same pipeline.

Our analysis identified frequently mutated GPCRs (e.g., *GPR98/ADGRV1* and *GPR112/ADGRG4*) in multiple cancers, especially melanoma (SKCM). GPCR mutations appear to reflect accumulation of DNA damage and mutations across the genome and may be tumor markers for this process. Multiple prior studies (e.g. [8,46]) have suggested that GPCRs that are frequently mutated (e.g., certain adhesion GPCRs) in tumors should be further studied for their role as drivers of tumor progression and as targets for novel therapeutics. Several observations lead us to question this idea:

1.   The absence of enrichment of mutations in specific sites/regions on frequently mutated GPCRs
2.   The analysis from MutSig2CV indicating the likelihood that such mutations are not enriched at specific residues at rates statistically elevated above the background mutation rate, i.e. the rate of non-synonymous mutations is not statistically elevated, adjusted for the rate of synonymous mutations.
3.   The general low expression of such mutated GPCRs at the mRNA level
4.   The lack of impact of GPCR mutations on survival
5.   The lack of effect of GPCR mutations on gene expression

While GPCRs are of interest in cancer due to their high mRNA expression, using their frequency of mutation as a rationale to choose particular GPCRs as potentially functionally important or possible therapeutic targets is likely a flawed approach—a conclusion that contrasts with previous suggestions about mutated GPCRs. Our data also raise a cautionary point about mutations: frequency of mutation of a gene is insufficient evidence to suggest its importance as a potential oncogene or therapeutic target without additional analyses such as those noted above.

While we conclude that frequently mutated GPCRs are unlikely to be targets for therapeutic intervention in solid tumors, such GPCR genes may be useful for studies related to oncogenesis. For example, GPR98 is frequently mutated in a number of tumor types and is one of several genes with a high frequency of somatic mutations per kilobase of coding gene length, suggesting that the GPR98 gene lies on a chromosomal region particularly vulnerable to accumulation of DNA damage. Mutation to this gene seems to scale with mutational burden, in a manner that is independent of tumor type: in tumors such as melanoma and bladder cancer, in which DNA damage from environmental factors may occur via different mechanisms, the accumulation of damage at the GPR98 locus in response to genome-wide damage appears to occur at a constant rate. This rate is elevated relative to the rest of the coding genome in terms of mutation rate per kilobase. Understanding how and why this occurs may provide additional insight into the mechanisms that drive DNA damage.

Known driver mutations do not appear to influence GPCR expression in tumors but we excluded rare mutations. In order to ensure large numbers of replicates and high statistical significance, we analyzed tumors with high-frequency mutations (e.g., *P53* or *KRAS*) [9]. Highly expressed GPCRs are widely expressed among replicates of specific tumor types and are more prevalent in tumors than are common driver mutations. The overrepresentation of GPCRs among protein-coding genes with increased expression in solid tumors supports the hypothesis that the elevated expression of specific GPCRs is a hitherto underappreciated feature of solid tumors.

The finding that many GPCRs show altered mRNA expression in tumors at rates higher than occurs for coding genes in general raises the question: what causes GPCR-specific changes in expression? Regulation of GPCR mRNA expression is poorly understood; hence, exploration of potential mechanisms for such regulation/dysregulation is of interest. Studies of GPCRs with DE in tumors may shed light on such mechanisms and perhaps also have relevance for other disease settings with altered GPCR expression.

Given the widespread changes in copy number that occur in cancer, especially copy number amplification, which might explain the increased expression of particular GPCRs in tumors, we tested but failed to find that CNV can generally explain altered GPCR mRNA expression or DE. CNV is not stochastically distributed among the GPCR family, certain GPCRs are more frequently amplified (e.g., *GPR160*) or deleted (e.g., *PTH1R* in KIRC). Amplified and deleted GPCRs may have potential as biomarkers [47]. Our findings with respect to CNV and GPCRs raise a broader concern and cautionary note: changes in copy number of certain genes should not be taken as evidence of dysregulation of the same genes, unless one obtains evidence for corresponding changes at the mRNA level.

Numerous solid tumors have increased mRNA expression of large numbers of mostly non-mutated GPCRs: 72 GPCRs are overexpressed in >10 tumor subtypes, implying that common mechanisms may regulate GPCR expression in such tumors. Highly overexpressed GPCRs are potential candidates as drug targets. Of note, 77 such overexpressed GPCRs are targets of approved drugs that have the potential to be repurposed to treat tumors. The similarity of GPCR expression in primary tumors and metastases supports such therapeutic potential.

The data reveal that clusters of GPCRs may be prognostic indicators for survival and provide a molecular signature of the malignant phenotype. GPCRs whose expression adversely or positively predicts survival are candidates for antagonists or agonists, respectively as novel cancer drugs. Hierarchical clustering of tumor types based on GPCR expression identifies groups of tumors consistent with other molecular/phenotypic features of these tumors. Thus, the tumor GPCRome appears to be predictive of the broader molecular landscape of tumors.

Do GPCR mRNA data predict protein expression? Direct quantification of GPCR proteins is challenging, due to their generally low abundance and paucity of well-validated antibodies. However, mRNA expression of GPCRs, especially highly expressed GPCRs, generally predicts the presence of functionally active GPCRs in human and animal cells [48–52]. In contrast to earlier ideas, recent evidence supports the view that mRNA expression broadly predicts protein expression [53–57] (**Supplemental Note 1**). As an example, GPRC5a protein and mRNA abundance are concordant (**Supplemental Note 2; Fig S11**). GPCR detection via mass-spectrometry has been challenging; proteomics data (e.g., [58] indicate that at present, few GPCRs are detectable by such methods, likely due to the low abundance of GPCR proteins. As noted in **Results**, functional evidence is available for numerous GPCRs with DE in solid tumors. As cell-surface proteins enriched in tumors and cancer cells, certain GPCRs may represent novel tumor-associated proteins that might be targeted for diagnosis and/or treatment.

As a caveat to those ideas regarding mRNA and proteins expression, most analyses on their concordance cited above have used model organisms. Such concordance may be less evident in native mammalian cells, particularly in certain cell and tissue types, as observed in [54]. It is unclear whether GPCR mRNA expression predicts protein expression and functional effects in native cells/tissues. However, our recent work on Gq-coupled GPCRs in pancreatic cancer cells indicates a concentration-response relationship between GPCR mRNA expression and function, such that high GPCR expression corresponds to strong signaling and functional responses [59]. Previous work on GPCR mRNA expression in native cells has shown robust functional effects and/or protein expression in a range of settings, including cancer, for GPCRs identified as highly expressed, based on omics methods (reviewed in [50]). We thus suggest that highly expressed GPCRs (e.g. those expressed at ≥ 5-10 TPM from RNA-seq data) are very likely to be functional.

GPCR mutations, CNV and DE thus occur at a high frequency in solid tumors. Therefore, this receptor super-family may have unappreciated functional roles in such tumors, especially since GPCR expression appears to be

largely independent of tumor grade/stage and mutations. Our results imply that new insights may derive from further studies of GPCRs regarding mechanisms of gene expression and phenotype in solid tumors and perhaps other cancers. Of particular and perhaps rapid, translational importance is the potential of GPCR-targeted drugs, including FDA/EMA-approved drugs that might be repurposed as therapeutics for a variety of solid tumors.

Acknowledgements
This work was supported by the American Society of Pharmacology and Experimental Therapeutics (ASPET) David Lehr Award and Padres Pedal the Cause #PTC2017.

Results are in part based upon data from the Cancer Genome Atlas (TCGA) Research Network: http://cancergenome.nih.gov/.

Author Contributions
KS and PAI conceived the project. KS designed and implemented the analysis approach and co-wrote the manuscript. PAI directed the project, interpreted results and co-wrote the manuscript. KM implemented analysis methods and maintained datasets/databases and the project website. RC co-wrote and edited the manuscript. HC interpreted results, designed analysis approach and co-wrote the manuscript.

Supplemental Information
Supplement 1: Supplementary Figures 1-13, Supplementary Tables 1-5 and Supplementary Notes 1-3.
Supplement 2: Data file with GPCRs annotated by GtoPdb plus GPCR mutations and CNV
Supplement 3: Data file with GPCR expression in normal tissue, tumors and cancer cells and GPCRs with DE
Additional downloadable content: *insellab.github.io*

Table 1: Tumors surveyed for Differential Expression (DE) analysis. TCGA cancer type and sub-classification, if applicable, for solid tumors with distinct histological classification are shown, along with the number of replicates and GPCRs with increased or decreased expression for each type of tumor.

	Cancer Type	Histology/Subtype	Replicates	# GPCRs ↑	# GPCRs ↓
1	Adrenocortical Cancer (ACC)	Adrenocortical carcinoma - Usual Type	73	22	4
2	Bladder Cancer (BLCA)	Papillary bladder cancer (BLCA_P)	130	26	12
		Non-papillary bladder cancer (BLCA_NP)	267	36	14
3	Breast Cancer (BRCA)	Infiltrating ductal carcinoma (IDC), Her2 positive (BRCA_IDC_Her2+)	48	46	25
		IDC, Hormone Receptor positive (BRCA_IDC_HR+)	431	49	29
		IDC, Triple positive (BRCA_IDC_3pl+)	54	51	25
		IDC, Triple negative (BRCA_IDC_3pl-)	109	51	24
		Invasive Lobular Carcinoma (ILC), Hormone R positive (BRCA_Lob_HR+)	57	50	31
4	Cervical Cancer (CESC)	Cervical Squamous Cell Carcinoma (CESC_CervSq)	252	40	18
		Endocervical Adenocarcinoma of the Usual Type (CESC_ECAD)	21	38	23
		Mucinous Adenocarcinoma of Endocervical Type (CESC_Muc)	17	38	24
5	Colon Cancer (COAD)	Colon Adenocarcinoma in the sigmoid colon (COAD_Sig)	71	41	28
		Colon Adenocarcinoma in the transverse colon (COAD_Trans)	22	31	23
6	Esophageal Cancer (ESCA)	Esophagus Adenocarcinoma (ESCA AD)	89	59	17
		Esophagus Squamous Cell Carcinoma (ESCA SQC)	92	43	9
7	Kidney Papillary Cell Carcinoma (KIRP)	-	288	28	14
8	Kidney Clear Cell Carcinoma (KIRC)	-	523	65	11
9	Kidney Chromophobe (KICH)	-	66	29	10
10	Liver Cancer (LIHC)	Liver Hepatocellular Carcinoma (LIHC)	360	11	7
11	Lung adenocarcinoma (LUAD)	Lung Papillary Adenocarcinoma (LUAD_Pap)	23	22	33
		Lung Bronchioloalveolar Carcinoma Non-Mucinous (LUAD_BCNM)	19	34	36
		Lung Adenocarcinoma- Not Otherwise Specified (LUAD_NOS)	308	33	33
		Lung Adenocarcinoma – Mixed (LUAD_Mixed)	105	29	31
		Lung Acinar Adenocarcinoma (LUAD_Acinar)	18	27	36
12	Lung squamous cell carcinoma (LSQC)	Lung Squamous Cell Carcinoma- Not Otherwise Specified (LSQC_NOS)	468	34	31
		Lung Basaloid Squamous Cell Carcinoma (LSQC_Basal)	14	38	39
13	Skin Cutaneous Melanoma (SKCM)	Primary melanomas (SKCM_Primary)	100	34	18
		Distant metastases (SKCM_DMet)	68	41	11
14	Ovarian Cancer (OV)	Ovarian Serous Cystadenocarcinoma (OV)	418	57	11
15	Pancreatic Cancer (PAAD)	Pancreatic Ductal Adenocarcinoma (PDAC)	147	68	11
16	Prostate Cancer (PRAD)	Prostate Adenocarcinoma Acinar Type (PRAD)	475	27	25
17	Stomach Cancer (STAD)	Stomach, Adenocarcinoma, Diffuse Type (STAD_Diff)	68	53	12
		Stomach, Adenocarcinoma, Not Otherwise Specified (STAD_NOS)	154	48	14
		Stomach, Intestinal Adenocarcinoma, Mucinous Type (STAD_Muc)	19	55	15
		Stomach, Intestinal Adenocarcinoma, Not Otherwise Specified (STAD_IntNOS)	73	40	16
		Stomach, Intestinal Adenocarcinoma, Tubular Type (STAD_IntTub)	76	39	12
		Stomach Adenocarcinoma, Signet Ring Type (STAD_Sig)	12	37	10
18	Testicular Cancer (TGCT)	Seminoma (TGCT_Sem)	72	73	16
		Non-seminoma (TGCT_NonSem)	65	76	9
19	Thyroid Cancer (THCA)	Thyroid Papillary Carcinoma - Classical/usual (THCA_Usual)	358	33	15
		Thyroid Papillary Carcinoma - Follicular (>= 99% follicular patterned) (THCA_fol)	101	17	6
		Thyroid Papillary Carcinoma - Tall Cell (>= 50% tall cell features) (THCA_TC)	36	33	17
20	Uterine Carcinosarcoma (UCS)	Uterine Carcinosarcoma/Malignant Mixed Mullerian Tumor (MMMT): (UCS_NOS)	24	43	19
		Uterine Carcinosarcoma/ MMMT: Heterologous Type (UCS_Het)	20	44	22
		Uterine Carcinosarcoma/MMMT: Homologous Type (UCS_Homo)	13	43	20

Methods

Software used and availability of data

RESOURCE	SOURCE	LOCATION OF DATA
Software and Algorithms		
R v 3.3.2	**The Comprehensive R Archive Network**	**https://cran.r-project.org/**
edgeR	**Bioconductor**	**http://bioconductor.org/packages/edgeR**
Deposited Data		
Numbers of GPCR mutations in each tumor type	**This paper**	**Supplement 2**
Occurrence of CNV for GPCRs in each tumor type	**This paper**	**Supplement 2**
GPCR expression and G protein linkages in tumors and healthy tissue	**This paper**	**Supplement 1**
GPCRs with DE in tumors	**This paper**	**Supplement 1**
GPCR expression in cancer cells	**This paper**	**Supplement 1**
GPCRs annotated by IUPHAR	**IUPHAR/BJP**	**Supplement 2**
Analyzed TCGA mutation data	**TCGA/UCSC Xena**	*https://insellab.github.io/gpcr_mutations*
Analyzed TCGA CNV data	**TCGA/UCSC Xena**	*https://insellab.github.io/gpcr_cnv*
Analyzed TCGA RNAseq data	**TCGA/UCSC Xena**	*https://insellab.github.io/gpcr_tcga_exp*
Analyzed GTEX RNAseq data	**GTEX/UCSC Xena**	*https://insellab.github.io/gpcr_gtex_exp*
Analyzed RNAseq data from cancer cells	**Various cited sources**	*https://insellab.github.io/gpcr_cells_exp*

1. Contact for Resource Sharing

A website has been created for sharing all data at *https://insellab.github.io/*. Links to data files will be posted on this website following peer review. All data provided will be open access following peer review; information about how to cite data generated from this study is available at *https://insellab.github.io/*.

Contact for access to data:
Krishna Sriram
Department of Pharmacology, UC San Diego.
ksriram@ucsd.edu
Tel: 858-534-2298

2. Details of Methods

2.1. Differential Expression (DE) Analysis:

Gene expression for the GTEX and TCGA datasets, assayed via RNAseq, was downloaded from the UCSC Xena Portal (xena.ucsc.edu). For DE analysis, RSEM expected counts were obtained, which were computed via the TOIL pipeline, described in [10] and available at the Xena Portal by the authors of the TOIL project (*https://xenabrowser.net/datapages/?host=https://toil.xenahubs.net*).

The analyzed data from the TOIL project were generated as follows. Merged FASTQ files were adapter-trimmed via CUTADAPT, followed by alignment via STAR [60]. Gene expression was then quantified using RSEM [61]. The HG38 reference genome, with Gencode V23 annotations, was used in the TOIL analysis. For this study, both RSEM estimated counts (for DE analysis) and RSEM TPMs (for evaluating magnitudes of expression) were used.

Files were accessed from (*https://xenabrowser.net/datapages/?dataset=tcga_gene_expected_count&host=https://toil.xenahubs.net*) for TCGA expected counts (version 2016-09-01) and (*https://xenabrowser.net/datapages/?dataset=gtex_gene_expected_count&host=https://toil.xenahubs.net*) for GTEX expected counts (version 2016-05-19).

Expression in TPMs for GPCRs was queried via *https://xenabrowser.net/heatmap/* for both TCGA and GTEX.

Following the download of gene expression data, corresponding files on sample phenotype were obtained from the relevant links hosted at *https://xenabrowser.net/datapages/?host=https://tcga.xenahubs.net* for TCGA and at *https://xenabrowser.net/datapages/?cohort=GTEX* for GTEX samples, respectively. Samples were then grouped for DE analysis based on attributes such as tissue type and tumor type.

We used a table of estimated counts from tumor and normal tissue as input for analysis in edgeR [62], yielding normalized abundance in CPM (using TMM normalization) and DE, showing magnitude of fold change and statistical significance estimated by FDR (False Discovery Rate). We estimated fold-changes of gene expression in tumors compared to normal tissue via an exact test. Genes that changed with FDR <0.05 were considered statistically significant, however we focused attention on GPCRs that besides low FDRs are also expressed at >1 TPM in tumors, as high expressed GPCRs are likely of greater interest. The 20 TCGA tumor types were divided into 45 subtypes/categories (**Table 1**) based on histological classification. Different tumor subtypes show distinct GPCR expression, e.g., Classical vs. Follicular Thyroid Cancer (THCA), Triple negative vs. Her2 Breast Cancer Cancer, Infiltrating Ductal Carcinoma (BRCA IDC), and Esophageal (ESCA) squamous cell carcinoma vs. Adenocarcinoma (**Fig S1A-F**), hence requiring this subdivision into tumor subtypes.

In addition to the standard TMM approach in edgeR, we tested upper-quartile normalization before conducting DE analysis in edgeR. The two methods yielded nearly identical results (**Fig S11C, D).** Log2 fold-changes for all genes and GPCRs show that expression changes calculated by both methods are closely correlated, with nearly identical magnitude. We also evaluated DE via EBseq [63]. EBseq and edgeR yielded very similar results (**Fig S11A, B**), in particular for GPCRs, implying that assumptions implicit in the DE analysis via edgeR/TMM normalization do not skew or bias the results. As an empirical test, we verified that GPCRs that show large fold-changes between tumor and normal samples also show large differences in normalized gene expression in TPM, e.g., *EDNRB*, *GPR143* and *ADGRG1* in SKCM (**Fig S2F-H**).

2.2. Database of normalized GPCR expression in tumors and normal tissue

For GPCR expression in tissue, expression as TPM (Transcripts Per Million, a normalization of gene abundance that corrects for effective length of genes) and CPM (Counts Per Million, number of times a gene is encountered per million reads, hence normalization for library size without length normalization) are provided in **Supplement 3**.

Data in TPM are provided for assessing the relative abundance of members of a gene family, such as GPCRs, *within* an individual sample or a set of biological replicates while data in CPM are provided for comparing a specific gene *between* multiple groups of dissimilar samples, where length normalization is problematic because

dissimilar data sets may be normalized differently. We provide GPCR expression in both formats to facilitate
different approaches for analysis.

This resource thus enables an estimate of GPCR abundance in more rigorous terms than comparing FPKM/TPM
values for a particular gene across different tissue types. Further, this approach provides normalized gene
expression estimates (in TPM and CPM) using the same units and analysis methods for both normal and cancer
tissue, thus allowing for direct comparison.

Gene abundances in CPM were calculated via EdgeR. In several cases the same normal tissue dataset was used
to compare multiple tumors (e.g., GTEX Kidney data was used for comparison with KIRP, KICH and KIRC). In
these cases, the calculated CPM values for normal tissue from each analysis were (as expected) very similar but
not equal, as EdgeR TMM normalization yields slightly different normalization factors in each case. The CPM
values presented for these tissues are thus average values (e.g., for CPMs in normal Kidney tissue, the values
provided are the average of the data obtained from comparisons of normal kidney with KIRP, KICH and KIRC
respectively). The normal tissue types where this was performed are breast, lung and kidney.
14
15 **2.3. GPCR mutation and copy number analysis**

16 For each cancer type, tables of somatic, non-silent mutations (gene-level) and somatic mutations (SNPs and
17 small INDELs) and Gene-level GISTIC2 thresholded copy number variation were obtained using
18 *https://xenabrowser.net/datapages/?host=https://tcga.xenahubs.net* and links within*.
19
20 For mutation data, we used results obtained via the Broad Automated Pipeline, where available. In other cases,
21 we used data from the Baylor College of Medicine sequencing center. The source of the mutation data is
22 indicated on the respective downloadable files and **Table S1**. In all cases, the HG19 reference genome was used
23 for calling mutations. Mutation data for genes coding for GPCRs were extracted as part of the present study; all
24 GPCR mutation data are available for download as supplemental material.
25
26 TCGA copy number estimates were obtained using Affymetrix SNP 6.0 arrays. The data were analyzed via
27 GISTIC 2.0 [44] to obtain gene-level estimates of copy number variation. The resulting 'thresholded' GISTIC 2.0
28 data yields values of -2,-1, 0, 1, 2, indicating homozygous/2 copy deletion, heterozygous/single copy deletions, no
29 change, low level amplification, and high level amplification, respectively. Copy number variation for GPCR genes
30 in each tumor type was extracted and is available as downloadable material.
31
32 **2.4. Which genes are included in this analysis?**
33
We evaluated all GPCRs annotated by IUPHAR/*British Journal of Pharmacology* [2], accessible via
(*http://www.guidetopharmacology.org/GRAC/ReceptorFamiliesForward?type=GPCR*). This list primarily focuses
on endoGPCRs (GPCRs natively expressed in peripheral tissue, possess endogenous ligands and receptors
primarily used as drug targets). The IUPHAR list includes taste and vision GPCRs but not olfactory receptors. We
excluded several GPCRs annotated by IUPHAR but which are thought to be pseudogenes. The list of GPCRs in
this analysis is provided in **Supplement 2**.
40
41 For these annotated GPCRs, we included information about their linkages to G proteins and their status as
42 orphans or not. For non-orphans, an example of an endogenous ligand is provided. These data are almost
43 entirely based on information available at the IUPHAR website cited above. In a few cases where such
44 information is not provided by IUPHAR, we have used other literature sources for annotation.
45
46 **2.5 GPCR expression in cancer cells mined from other sources**
47
48 GPCR expression in a range of cancer cell lines was queried via the EBI Expression Atlas
49 (*https://www.ebi.ac.uk/gxa/home*) for cell line profiles part of CCLE [38] as well as by Genentech [39], profiled via
50 RNA-seq. These data were analyzed as part of the EBI Expression Atlas via the iRAP bioinformatics analysis

pipeline, described in detail in [37] wherein gene expression was computed in FPKM (Fragments Per Kilobase of exon, per Million reads), a length-normalized expression abundance estimate analogous to TPM units, used for TOIL TCGA data. Precisely, statistically relevant comparisons between gene abundances in TPM and FPKM are not feasible; however, empirical comparisons between such data sets are possible. In general, genes with high abundances in TPM or FPKM will be highly expressed relative to other genes within sets of samples; hence, our comparison of CCLE and other cell-based data vs. TOIL TCGA data serves as an empirical confirmation of the fact that GPCRs highly expressed in TCGA tumors are also present in cancer cells.

Normalized gene expression in cancer cells from other sources [40,41,64] were obtained via NCBI GEO, wherein analyzed RNA-seq data with quantification of gene expression were provided. As with the data from EBI above, such data allowed for empirical comparisons vs. TCGA TOIL data to confirm the presence of GPCRs in cancer cells, which were also detected in tumors.

3. Quantification and Statistical Analysis

DE analysis was performed in the R software environment via EdgeR [62], as discussed above in section 2.1. We used the following criteria to evaluate GPCRs with significant DE:

1. $FDR < 0.05$. In the majority of cases, genes with a high fold change also showed FDRs $<< 0.05$.
2. Magnitude of fold change > 2 fold (increase or decrease).
3. Magnitude of expression > 1 TPM median expression in tumors, as calculated by RSEM in the TOIL pipeline, discussed in section 2 above. We focused on genes with significant DE and high expression because our primary goal was to identify GPCRs that may be drug targets and/or biomarkers.

For compilation and distribution of data we assembled data files primarily in Microsoft Excel, with files stored in .xlsb format.

Plots of normalized expression in tumors and normal tissue, whether in TPM or CPM, show median expression for respective cohorts, along with upper and lower quartiles, as indicated in figure legends where applicable.

The numbers of replicates in each sample group/category of normal tissue and tumors are provided in **Tables S1 and S2 and Table 1**, respectively. Tumor types with 10 or more replicates were considered for DE analysis. A small number of samples (<< 1% of TCGA samples studied) were excluded because they were duplicated in downloaded databases from TOIL and showed discrepancies between expression in these downloaded data vs expression data queried via the visualization tool on the TOIL website. These discrepant samples are provided on a downloadable list at *insellab.github.io*.

DE results presented in this text are from comparisons between TCGA and GTEX samples, but we also include in our MDS analysis and in all downloadable counts files, data for TCGA-matched "normal" samples taken from tissue adjacent to tumors of TCGA patients. In general, normal TCGA and GTEX samples cluster closer together than do TCGA tumors and GTEX normal samples (**Figures S1G-J**).

The overlap, however, is not exact. In several cases, we found differences between TCGA normal and GTEX samples. It is unclear if these differences result from biological or technical factors; prior data show that tumors impact surrounding "normal" tissue and can also induce global changes [65–68]. Hence, we have not used batch-correction methods to account for these variations between TCGA normal and GTEX tissues. In general, DE of GPCRs is similar whether TCGA normal tissue or GTEX tissue is compared to TCGA tumor samples (e.g., **Figures S11E-F**) suggesting that such differences are unlikely to impact upon the general conclusions of this study.

For TCGA data, some recent efforts (e.g., *http://bioinformatics.mdanderson.org/tcgambatch/*) have been made to
account for batch effects, though peer-reviewed studies have not yet established best practices for batch
correction of TCGA data. Many TCGA datasets for individual tumor types contain numerous batches (with
batches defined in terms of factors such as sequencing runs, or location of tissue collection), with small numbers
of replicates in each batch. Given this, it is unclear if such batch corrections account for technical variation among
batches or merely suppress biological variation, especially with the known heterogeneity among tumor samples.
In light of this, we present all data from TCGA and GTEX without correcting for batch effects.
8
In nearly all cases, DE of GPCRs we highlight have large fold changes with high statistical significance (i.e., FDR
<< 0.05), such that minor technical variations ought not substantially impact our key findings. Moreover, the fact
that TCGA tumors and GTEX normal tissues form distinct, separated clusters (and hence show a high degree of
DE) is unlikely to be due to technical factors. In several cases (e.g., KICH matched normal vs GTEX kidney
samples; **Fig S1G**), TCGA matched normal and GTEX normal tissues are in fact highly similar, whereas in other
cases they are not (e.g., PRAD, **Fig S1J**). This suggests that technical factors between the two studies do not
consistently skew/bias the two data sets vs. each other.
16
Analysis to identify dependence of GPCR expression on patient characteristics, such as sex or tumor stage was
done as follows: GPCR expression for tumor samples normalized in CPM, output from edgeR was used to
evaluate whether grouping samples together based on a specified attribute (e.g., sex) resulted in a statistically
significant relative risk for that attribute tested for results in elevated GPCR expression in one group compared to
the other. We tested this using Fisher's exact test in R, yielding calculations of risk ratio and p-values, with the
terms in the contingency table for Fisher's test being the number of samples in each group with GPCR expression
either above or below the population median. P-values were then adjusted for multiple testing using the p_adjust
function in R, utilizing the Benjamini–Hochberg method. These adjusted p-values thus indicate whether any
GPCRs have expression that is significantly associated with a given patient attribute. Specific attributes tested for
were: sex, tumor stage, grade and pathological T (depending on availability of metadata for each tumor type and
the number of available replicates). This method was also used to evaluate the significance of associations
between expression of GPCRs and presence of specific driver mutations (e.g. presence or absence of mutations
to TP53 or KRAS) and association between GPCR mRNA expression and the thresholded GISTIC 2.0 CNV call.
30
Survival analysis was performed in R using gene expression data normalized in CPM (i.e., units appropriate for
comparisons <u>between samples</u>) via the "survival" and "survminer" packages. For each tumor type analyzed,
samples were divided into two groups based on median expression of the gene being tested, and differences in
survival between the two groups were calculated. The modified Peto-Peto (mPP) method was used to estimate
the statistical significance of differences in survival rates. We noted cases where the effects of GPCR expression
on survival were related/coupled (hence, the presence of composite markers of survival noted in the text). Thus,
these statistical tests were not independent; given a lack of understanding about the nature of such dependencies
(which to our knowledge were hitherto unknown), we did not adjust p-values for multiple testing. We also note
that a) due to our use of relevant units for normalizing the gene expression data prior to performing survival
analysis (CPM instead of units such as FPKM or RPKM) and b) the subdivision of broad TCGA categories into
appropriate sub-types of tumors with consistent histological classification (e.g. dividing ESCA into
adenocarcinomas and squamous cell carcinomas and analyzing each separately) our analysis yields associations
of genes with survival that may differ in some cases from other sources that also provide survival analysis of
TCGA data. We selected a significance threshold of $p < 0.05$, which is relatively lenient for survival analysis, as
our initial priority was to minimize false negatives; we anticipate these analyses will need subsequent validation
efforts in specific tumor types, with larger numbers of patients.
47
48　**4. Data and Software Availability**
49　All data generated in this project are hosted at *https://insellab.github.io/*. These data are all open access. Relevant
50　software such as R and edgeR are also freely available, refer key resources table.
51

1   References

1.   Lonsdale J, Thomas J, Salvatore M, Phillips R, Lo E, Shad S, et al. The Genotype-Tissue Expression (GTEx) project. Nat Genet. 2013;45: 580. doi:10.1038/ng.2653 https://www.nature.com/articles/ng.2653#supplementary-information

2.   Alexander SP, Davenport AP, Kelly E, Marrion N, Peters JA, Benson HE, et al. The Concise Guide to PHARMACOLOGY 2015/16: G protein-coupled receptors. Br J Pharmacol. 2015/12/10. School of Biomedical Sciences, University of Nottingham Medical School, Nottingham, NG7 2UH, UK. Clinical Pharmacology Unit, University of Cambridge, Cambridge, CB2 0QQ, UK. School of Physiology and Pharmacology, University of Bristol, Bristol, BS8 1TD, U; 2015;172: 5744–5869. doi:10.1111/bph.13348

3.   Sriram K IP. G protein-coupled receptors as targets for approved drugs: How many targets and how many drugs? Mol Pharmacol. 2018;Apr 1;93(4.

4.   Wacker D, Stevens RC, Roth BL. How Ligands Illuminate GPCR Molecular Pharmacology. Cell. 2017/07/29. Department of Pharmacology and Division of Chemical Biology and Medicinal Chemistry, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, NC 27514, USA. Departments of Biological Sciences and Chemistry, Bridge Institute, University; 2017;170: 414–427. doi:10.1016/j.cell.2017.07.009

5.   Dorsam RT, Gutkind JS. G-protein-coupled receptors and cancer. Nat Rev Cancer. 2007;7: 79–94. Available: http://dx.doi.org/10.1038/nrc2069

6.   Hanahan D, Weinberg RA. The hallmarks of cancer. Cell. 2000/01/27. Department of Biochemistry, Hormone Research Institute, University of California at San Francisco, 94143, USA.; 2000;100: 57–70.

7.   Bar-Shavit R, Maoz M, Kancharla A, Nag JK, Agranovich D, Grisaru-Granovsky S, et al. G Protein-Coupled Receptors in Cancer. Int J Mol Sci. 2016/08/17. Sharett Institute of Oncology, Hadassah-Hebrew University Medical Center, Jerusalem 91120, Israel. Rachelbar@ekmd.huji.ac.il. Sharett Institute of Oncology, Hadassah-Hebrew University Medical Center, Jerusalem 91120, Israel. Myriamm@hadassah.org.il. Share; 2016;17. doi:10.3390/ijms17081320

8.   O'Hayre M, Vazquez-Prado J, Kufareva I, Stawiski EW, Handel TM, Seshagiri S, et al. The emerging mutational landscape of G proteins and G-protein-coupled receptors in cancer. Nat Rev Cancer. 2013;13: 412–424. doi:10.1038/nrc3521

9.   Kandoth C, McLellan MD, Vandin F, Ye K, Niu B, Lu C, et al. Mutational landscape and significance across 12 major cancer types. Nature. 2013;502: 333–339. doi:10.1038/nature12634 http://www.nature.com/nature/journal/v502/n7471/abs/nature12634.html#supplementary-information

10. Vivian J, Rao AA, Nothaft FA, Ketchum C, Armstrong J, Novak A, et al. Toil enables reproducible, open source, big biomedical data analyses. Nat Biotechnol. 2017/04/12. Computational Genomics Lab, UC Santa Cruz Genomics Institute, University of California Santa Cruz, Santa Cruz, California, USA. AMP Lab, University of California Berkeley, Berkeley, California, USA. UC Berkeley ASPIRE Lab, Berkeley, Berkeley, California, ; 2017;35: 314–316. doi:10.1038/nbt.3772

11. Zhou H, Rigoutsos I. The emerging roles of GPRC5A in diseases. Oncoscience. 2015/01/27. Computational Medicine Center, Jefferson Alumni Hall, Thomas Jefferson University, Philadelphia, PA.; 2014;1: 765–776. doi:10.18632/oncoscience.104

12. Zhou H, Telonis AG, Jing Y, Xia NL, Biederman L, Jimbo M, et al. GPRC5A is a potential oncogene in pancreatic ductal adenocarcinoma cells that is upregulated by gemcitabine with help from HuR. Cell Death Dis. 2016/07/16. Computational Medicine Center, Sidney Kimmel Medical College, Thomas Jefferson University, 1020 Locust Street Philadelphia, PA 19107, USA. Department of Neuroscience and The Farber

Institute for Neuroscience, Thomas Jefferson University, 900 Walnut Street; 2016;7: e2294. doi:10.1038/cddis.2016.169

13.   Wiley SZ, Sriram K, Liang W, Chang SE, French R, McCann T, et al. GPR68, a proton-sensing GPCR, mediates interaction of cancer-associated fibroblasts and cancer cells. Faseb j. 2017/11/03. Department of Pharmacology, University of California, San Diego, La Jolla, California, USA. The Key Laboratory of Cardiovascular Remodeling and Function Research, Qilu Hospital of Shandong University, Jinan, China. Moores Cancer Center, University of Cali; 2018;32: 1170–1183. doi:10.1096/fj.201700834R

14.   Szklarczyk D, Morris JH, Cook H, Kuhn M, Wyder S, Simonovic M, et al. The STRING database in 2017: quality-controlled protein-protein association networks, made broadly accessible. Nucleic Acids Res. 2016/12/08. Institute of Molecular Life Sciences and Swiss Institute of Bioinformatics, University of Zurich, 8057 Zurich, Switzerland. Resource on Biocomputing, Visualization, and Informatics, University of California, San Francisco, CA 94158-2517, USA. Novo Nordisk; 2017;45: D362-d368. doi:10.1093/nar/gkw937

15.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A. 2005/10/04. Broad Institute of Massachusetts Institute of Technology and Harvard, 320 Charles Street, Cambridge, MA 02141, USA.; 2005;102: 15545–15550. doi:10.1073/pnas.0506580102

16.   Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids Res. 1999/12/11. Institute for Chemical Research, Kyoto University, Uji, Kyoto 611-0011, Japan. kanehisa@kuicr.kyoto-u.ac.jp; 2000;28: 27–30.

17.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. Nat Genet. 2000/05/10. Department of Genetics, Stanford University School of Medicine, California, USA. cherry@stanford.edu; 2000;25: 25–29. doi:10.1038/75556

18.   Consortium TGO. Expansion of the Gene Ontology knowledgebase and resources. Nucleic Acids Res. 2016/12/03. 2017;45: D331–D338. doi:10.1093/nar/gkw1108

19.   Kuleshov M V, Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, et al. Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. Nucleic Acids Res. 2016/05/05. Department of Pharmacology and Systems Therapeutics, BD2K-LINCS Data Coordination and Integration Center, Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place Box 1215, New York, NY 10029, USA. Fluid Physics and Transport Processes Branch, N; 2016;44: W90-7. doi:10.1093/nar/gkw377

20.   Binder JX, Pletscher-Frankild S, Tsafou K, Stolte C, O'Donoghue SI, Schneider R, et al. COMPARTMENTS: unification and visualization of protein subcellular localization evidence. Database (Oxford). 2014/02/28. Structural and Computational Biology Unit, European Molecular Biology Laboratory (EMBL), 69117 Heidelberg, Germany, Bioinformatics Core Facility, Luxembourg Centre for Systems Biomedicine (LCSB), University of Luxembourg, 4362 Esch-sur-Alzette, Luxembourg; 2014;2014: bau012. doi:10.1093/database/bau012

21.   Insel PA, Sriram K, Wiley SZ, Wilderman A, Katakia T, McCann T, Yokouchi H, Zhang L, Corriden R, Liu D FM. GPCRomics: GPCR expression in cancer cells and tumors identifies new, potential biomarkers and therapeutic targets. Front Pharmacol. 2018;9.

22.   Richardson DR, Baker E. The uptake of iron and transferrin by the human malignant melanoma cell. Biochim Biophys Acta. 1990/06/12. Department of Physiology, University of Western Australia, Nedlands.; 1990;1053: 1–12.

23. Chen KG, Leapman RD, Zhang G, Lai B, Valencia JC, Cardarelli CO, et al. Influence of melanosome dynamics on melanoma drug sensitivity. J Natl Cancer Inst. 2009/08/26. Laboratory of Cell Biology, National Cancer Institute, National Institutes of Health, Bldg 37, Rm 2108, Bethesda, MD 20892, USA.; 2009;101: 1259–1271. doi:10.1093/jnci/djp259

24. Stracke ML, Engel JD, Wilson LW, Rechler MM, Liotta LA, Schiffmann E. The type I insulin-like growth factor receptor is a motility receptor in human melanoma cells. J Biol Chem. 1989/12/25. Laboratory of Pathology, National Cancer Institute, Bethesda, Maryland 20892.; 1989;264: 21544–21549.

25. Boire A, Covic L, Agarwal A, Jacques S, Sherifi S, Kuliopulos A. PAR1 is a matrix metalloprotease-1 receptor that promotes invasion and tumorigenesis of breast cancer cells. Cell. 2005/02/15. Molecular Oncology Research Institute, Tufts-New England Medical Center, Boston, Massachusetts 02111, USA.; 2005;120: 303–313. doi:10.1016/j.cell.2004.12.018

26. Fujimoto D, Hirono Y, Goi T, Katayama K, Matsukawa S, Yamaguchi A. The activation of Proteinase-Activated Receptor-1 (PAR1) mediates gastric cancer cell proliferation and invasion. BMC Cancer. 2010/08/21. First Department of Surgery, Faculty of Medicine and Division of Bioresearch Laboratories, University of Fukui, Fukui 910-1193, Japan. rui5218@yahoo.co.jp; 2010;10: 443. doi:10.1186/1471-2407-10-443

27. Darmoul D, Gratio V, Devaud H, Peiretti F, Laburthe M. Activation of proteinase-activated receptor 1 promotes human colon cancer cell proliferation through epidermal growth factor receptor transactivation. Mol Cancer Res. 2004/09/24. Neuroendocrinologie et Biologie Cellulaire Digestives, Institut National de la Sante et de la Recherche Medicale U410, Faculte de Medecine Xavier Bichat, 75018 Paris, France. darmoul@bichat.inserm.fr; 2004;2: 514–522.

28. Shi X, Gangadharan B, Brass LF, Ruf W, Mueller BM. Protease-activated receptors (PAR1 and PAR2) contribute to tumor cell motility and metastasis. Mol Cancer Res. 2004/07/29. Cancer Biology Division, La Jolla Institute for Molecular Medicine, 4570 Executive Drive, San Diego, CA 92121, USA.; 2004;2: 395–402.

29. Su S, Li Y, Luo Y, Sheng Y, Su Y, Padia RN, et al. Proteinase-activated receptor 2 expression in breast cancer and its role in breast cancer cell migration. Oncogene. 2009/06/23. Research Center for Traditional Chinese Medicine Complexity System, Shanghai University of Traditional Chinese Medicine, Shanghai, China.; 2009;28: 3047–3057. doi:10.1038/onc.2009.163

30. Darmoul D, Marie JC, Devaud H, Gratio V, Laburthe M. Initiation of human colon cancer cell proliferation by trypsin acting at protease-activated receptor-2. Br J Cancer. 2001/09/05. Neuroendocrinologie et Biologie Cellulaire Digestives, Institut National de la Sante et de la Recherche Medicale, INSERM U410, Faculte de Medecine Xavier Bichat, Paris, 75018, France.; 2001;85: 772–779. doi:10.1054/bjoc.2001.1976

31. Jahan I, Fujimoto J, Alam SM, Sato E, Sakaguchi H, Tamaya T. Role of protease activated receptor-2 in tumor advancement of ovarian cancers. Ann Oncol. 2007/09/01. Department of Obstetrics and Gynecology, Gifu University School of Medicine, 1-1 Yanagido, Gifu City 501-1194, Japan.; 2007;18: 1506–1512. doi:10.1093/annonc/mdm190

32. Lahav R, Suva ML, Rimoldi D, Patterson PH, Stamenkovic I. Endothelin receptor B inhibition triggers apoptosis and enhances angiogenesis in melanomas. Cancer Res. 2004/12/18. Division of Experimental Pathology, Institut Universitaire de Pathologie Universite de Lausanne, Lausanne, Switzerland. ronit.lahav@melcure.com; 2004;64: 8945–8953. doi:10.1158/0008-5472.can-04-1510

33. Lahav R. Endothelin receptor B is required for the expansion of melanocyte precursors and malignant melanoma. Int J Dev Biol. 2005/05/21. Institut Universitaire de Pathologie, Lausanne, Switzerland. ronit.lahav@melcure.com; 2005;49: 173–180. doi:10.1387/ijdb.041951rl

34. Bai J, Xie X, Lei Y, An G, He L, Lv X. Ocular albinism type 1-induced melanoma cell migration is mediated through the RAS/RAF/MEK/ERK signaling pathway. Mol Med Rep. 2014/04/17. Department of Medical Oncology, Shaanxi Provincial People's Hospital, The Third Affiliated Hospital of The School of Medicine, Xi'an Jiaotong University, Xi'an, Shaanxi 710068, P.R. China. Key Laboratory of Resource Biology and Biotechnology in Western Ch; 2014;10: 491–495. doi:10.3892/mmr.2014.2154

35. Hertzman Johansson C, Azimi A, Frostvik Stolt M, Shojaee S, Wiberg H, Grafstrom E, et al. Association of MITF and other melanosome-related proteins with chemoresistance in melanoma tumors and cell lines. Melanoma Res. 2013/08/08. Department of Oncology-Pathology, Karolinska Institutet, Karolinska University Hospital, Stockholm, Sweden. carolina.johansson@ki.se; 2013;23: 360–365. doi:10.1097/CMR.0b013e328362f9cd

36. Zhou C, Dai X, Chen Y, Shen Y, Lei S, Xiao T, et al. G protein-coupled receptor GPR160 is associated with apoptosis and cell cycle arrest of prostate cancer cells. Oncotarget. 2016/02/13. The National Center for Drug Screening and The CAS Key Laboratory of Receptor Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences (CAS), Shanghai 201203, China. The State Key Laboratory of Drug Research, Shanghai Institute of Mater; 2016;7: 12823–12839. doi:10.18632/oncotarget.7313

37. Fonseca NA, Petryszak R, Marioni J, Brazma A. iRAP - an integrated RNA-seq Analysis Pipeline. bioRxiv. 2014; Available: http://biorxiv.org/content/early/2014/06/06/005991.abstract

38. Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, et al. The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. Nature. 2012;483: 307–603. doi:http://www.nature.com/nature/journal/v483/n7391/abs/nature11003.html#supplementary-information

39. Klijn C, Durinck S, Stawiski EW, Haverty PM, Jiang Z, Liu H, et al. A comprehensive transcriptional portrait of human cancer cell lines. Nat Biotech. 2015;33: 306–312. doi:10.1038/nbt.3080 http://www.nature.com/nbt/journal/v33/n3/abs/nbt.3080.html#supplementary-information

40. Müller J, Krijgsman O, Tsoi J, Robert L, Hugo W, Song C, et al. Low MITF/AXL ratio predicts early resistance to multiple targeted drugs in melanoma. Nat Commun. 2014;5: 5712. doi:10.1038/ncomms6712 https://www.nature.com/articles/ncomms6712#supplementary-information

41. Witkiewicz AK, Balaji U, Eslinger C, McMillan E, Conway W, Posner B, et al. Integrated Patient-Derived Models Delineate Individualized Therapeutic Vulnerabilities of Pancreatic Cancer. Cell Rep. 2016/08/09. McDermott Center for Human Growth and Development, University of Texas Southwestern Medical Center, 6000 Harry Hines Boulevard, Dallas, TX 75390, USA; Simmons Cancer Center, University of Texas Southwestern Medical Center, 6000 Harry Hines Boulevard, Dall; 2016;16: 2017–2031. doi:10.1016/j.celrep.2016.07.023

42. Ruault M, van der Bruggen P, Brun ME, Boyle S, Roizes G, De Sario A. New BAGE (B melanoma antigen) genes mapping to the juxtacentromeric regions of human chromosomes 13 and 21 have a cancer/testis expression profile. Eur J Hum Genet. 2002/12/04. Institut de Genetique Humaine, CNRS UPR 1142, 141, rue de la Cardonille, 34396 Montpellier, France.; 2002;10: 833–840. doi:10.1038/sj.ejhg.5200891

43. Miller LD, Smeds J, George J, Vega VB, Vergara L, Ploner A, Pawitan Y, Hall P, Klaar S, Liu ET BJ. An expression signature for p53 status in human breast cancer predicts mutation status, transcriptional effects, and patient survival. Proc Natl Acad Sci. 2005;Sep 20;102.

44. Mermel CH, Schumacher SE, Hill B, Meyerson ML, Beroukhim R, Getz G. GISTIC2.0 facilitates sensitive and confident localization of the targets of focal somatic copy-number alteration in human cancers. Genome Biol. 2011/04/30. Cancer Program, The Broad Institute of MIT and Harvard, 7 Cambridge Center, Cambridge, MA 02142, USA.; 2011;12: R41. doi:10.1186/gb-2011-12-4-r41

45.   Beroukhim R, Getz G, Nghiemphu L, Barretina J, Hsueh T, Linhart D, et al. Assessing the significance of chromosomal aberrations in cancer: methodology and application to glioma. Proc Natl Acad Sci U S A. 2007/12/14. Broad Institute, Massachusetts Institute of Technology and Harvard University, 7 Cambridge Center, Cambridge, MA 02142, USA.; 2007;104: 20007–20012. doi:10.1073/pnas.0710052104

46.   Purcell RH HR. Adhesion G Protein–Coupled Receptors as Drug Targets. Annual review of pharmacology and toxicology. Annu Rev Pharmacol Toxicol. 2018;Jan 6;58:4.

47.   Vang Nielsen K, Ejlertsen B, Møller S, Trøst Jørgensen J, Knoop A, Knudsen H, et al. The value of TOP2A gene copy number variation as a biomarker in breast cancer: Update of DBCG trial 89D. Acta Oncol (Madr). 2008;47: 725–734. doi:10.1080/02841860801995396

48.   Snead AN, Insel PA. Defining the cellular repertoire of GPCRs identifies a profibrotic role for the most highly expressed receptor, protease-activated receptor 1, in cardiac fibroblasts. FASEB J. 2012/08/04. Department of Pharmacology, University of California at San Diego, La Jolla, California 92093, USA.; 2012;26: 4540–4547. doi:10.1096/fj.12-213496

49.   Berger M, Scheel DW, Macias H, Miyatsuka T, Kim H, Hoang P, et al. Galphai/o-coupled receptor signaling restricts pancreatic beta-cell expansion. Proc Natl Acad Sci U S A. 2015/02/20. Departments of Psychiatry and Center for Neurobiology and Psychiatry, Diabetes Center, Hormone Research Institute, and. Diabetes Center, Hormone Research Institute, and. Diabetes Center, Hormone Research Institute, and Medicine. Departments of Psychiatry ; 2015;112: 2888–2893. doi:10.1073/pnas.1319378112

50.   Insel PA, Wilderman A, Zambon AC, Snead AN, Murray F, Aroonsakool N, et al. G Protein-Coupled Receptor (GPCR) Expression in Native Cells: "Novel" endoGPCRs as Physiologic Regulators and Therapeutic Targets. Mol Pharmacol. 2015/03/05. Departments of Pharmacology (P.A.I., A.W., A.C.Z., A.N.S., N.A., D.S.M., S.Z., T.M., L.Z., K.S., A.M.C., A.V.M., R.M.L., A.C.O., R.C.) and Medicine (P.A.I., F.M.), University of California, San Diego, La Jolla, California pinsel@ucsd.edu. Departments of P; 2015;88: 181–187. doi:10.1124/mol.115.098129

51.   Koyama H, Iwakura H, Dote K, Bando M, Hosoda H, Ariyasu H, et al. Comprehensive Profiling of GPCR Expression in Ghrelin-Producing Cells. Endocrinology. 2015/12/17. Medical Innovation Center (H.I., K.D., M.B., T.K., C.S., K.H., K.K., K.N.) and Departments of Diabetes, Endocrinology, and Nutrition (H.K.) and Human Health Sciences (K.H.), Kyoto University Graduate School of Medicine, Kyoto 606-8507, Japan; National Cer; 2016;157: 692–704. doi:10.1210/en.2015-1784

52.   Regard JB, Sato IT, Coughlin SR. Anatomical profiling of G protein-coupled receptor expression. Cell. 2008/11/06. Cardiovascular Research Institute, University of California, San Francisco, San Francisco, CA 94158, USA. regardj@mail.nih.gov; 2008;135: 561–571. doi:10.1016/j.cell.2008.08.040

53.   Csardi G, Franks A, Choi DS, Airoldi EM, Drummond DA. Accounting for experimental noise reveals that mRNA levels, amplified by post-transcriptional processes, largely determine steady-state protein levels in yeast. PLoS Genet. 2015/05/08. Dept. of Statistics, Harvard University, Cambridge, Massachusetts, United States of America. Dept. of Statistics, Harvard University, Cambridge, Massachusetts, United States of America,; The Broad Institute of Harvard & MIT, Cambridge, Massachusetts, Unit; 2015;11: e1005206. doi:10.1371/journal.pgen.1005206

54.   Edfors F, Danielsson F, Hallstrom BM, Kall L, Lundberg E, Ponten F, et al. Gene-specific correlation of RNA and protein levels in human cells and tissues. Mol Syst Biol. 2016/12/13. Science for Life Laboratory, KTH - Royal Institute of Technology, Stockholm, Sweden. Department of Immunology, Genetics and Pathology, Rudbeck Laboratory, Uppsala University, Uppsala, Sweden. Science for Life Laboratory, KTH - Royal Institute of Technolog; 2016;12: 883. doi:10.15252/msb.20167144

55. Guimaraes JC, Rocha M, Arkin AP. Transcript level and sequence determinants of protein abundance and noise in Escherichia coli. Nucleic Acids Res. 2014/02/11. Department of Bioengineering, University of California, Berkeley, CA 94720, USA, California Institute for Quantitative Biosciences, University of California, Berkeley, CA 94720, USA, Computer Science and Technology Center, School of Engineering, Universit; 2014;42: 4791–4799. doi:10.1093/nar/gku126

56. Koussounadis A, Langdon SP, Um IH, Harrison DJ, Smith VA. Relationship between differentially expressed mRNA and mRNA-protein correlations in a xenograft model system. Sci Rep. 2015/06/09. School of Biology, University of St Andrews, St Andrews, Fife, KY16 9TH, UK. Division of Pathology, University of Edinburgh, Edinburgh, EH4 2XU, UK. School of Medicine, University of St Andrews, St Andrews, Fife, KY16 9TF, UK.; 2015;5: 10775. doi:10.1038/srep10775

57. Li JJ, Bickel PJ, Biggin MD. System wide analyses have underestimated protein abundances and the importance of transcription in mammals. PeerJ. 2014/04/02. Department of Statistics, University of California , Berkeley, CA , USA ; Departments of Statistics and Human Genetics, University of California , Los Angeles, CA , USA. Department of Statistics, University of California , Berkeley, CA , USA. Genomics Div; 2014;2: e270. doi:10.7717/peerj.270

58. Lapek Jr. JD, Greninger P, Morris R, Amzallag A, Pruteanu-Malinici I, Benes CH, et al. Detection of dysregulated protein-association networks by high-throughput proteomics predicts cancer vulnerabilities. Nat Biotechnol. 2017/09/12. Massachusetts General Hospital Cancer Center and Department of Medicine, Harvard Medical School, Charlestown, Massachusetts, USA.; 2017;35: 983–989. doi:10.1038/nbt.3955

59. Sriram K, Wiley SZ, Moyung K, Gorr MW, Marucut J, French RP, Lowy AM IP. High-content detection of GPCR mRNA expression: Implications for GPCR biology, translational studies and drug discovery. ACS Pharmacol Transl Sci. 2019; *Submitted, Under Review*.

60. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2012/10/30. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, USA. dobin@cshl.edu; 2013;29: 15–21. doi:10.1093/bioinformatics/bts635

61. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC Bioinformatics. 2011/08/06. Department of Computer Sciences, University of Wisconsin-Madison, Madison, WI, USA.; 2011;12: 323. doi:10.1186/1471-2105-12-323

62. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics. 2009/11/17. Cancer Program, Garvan Institute of Medical Research, 384 Victoria Street, Darlinghurst, NSW 2010, Australia. mrobinson@wehi.edu.au; 2010;26: 139–140. doi:10.1093/bioinformatics/btp616

63. Leng N, Dawson JA, Thomson JA, Ruotti V, Rissman AI, Smits BM, et al. EBSeq: an empirical Bayes hierarchical model for inference in RNA-seq experiments. Bioinformatics. 2013/02/23. Department of Statistics, University of Wisconsin, Madison, WI 53706, USA.; 2013;29: 1035–1043. doi:10.1093/bioinformatics/btt087

64. Reemann P, Reimann E, Ilmjarv S, Porosaar O, Silm H, Jaks V, et al. Melanocytes in the skin--comparative whole transcriptome analysis of main skin cell types. PLoS One. 2014/12/30. Department of Physiology, Institute of Biomedicine and Translational Medicine, University of Tartu, Tartu, Estonia. Core Facility of Clinical Genomics, Department of Pathophysiology, Department of Biomedicine, Institute of Biomedicine and Translational Me; 2014;9: e115717. doi:10.1371/journal.pone.0115717

65. Acharyya S, Ladner KJ, Nelsen LL, Damrauer J, Reiser PJ, Swoap S, et al. Cancer cachexia is regulated by selective targeting of skeletal muscle gene products. J Clin Invest. 2004/08/03. Division of Human

Cancer Genetics, Department of Molecular Virology, Immunology and Medical Genetics, The Ohio State University, Columbus 43210, USA.; 2004;114: 370–378. doi:10.1172/jci20174

66. Egeblad M, Nakasone ES, Werb Z. Tumors as organs: complex tissues that interface with the entire organism. Dev Cell. 2010/07/16. Cold Spring Harbor Laboratory, 1 Bungtown Road, Cold Spring Harbor, NY 11724, USA. egeblad@cshl.edu; 2010;18: 884–901. doi:10.1016/j.devcel.2010.05.012

67. Gabrilovich DI, Nagaraj S. Myeloid-derived suppressor cells as regulators of the immune system. Nat Rev Immunol. 2009/02/07. Department of Oncologic Sciences, H Lee Moffitt Cancer Center and Research Institute, University of South Florida, Tampa, Florida 33612, USA. dmitry.gabrilovich@moffitt.org; 2009;9: 162–174. doi:10.1038/nri2506

68. Redon CE, Dickey JS, Nakamura AJ, Kareva IG, Naf D, Nowsheen S, et al. Tumors induce complex DNA damage in distant proliferative tissues in vivo. Proc Natl Acad Sci U S A. 2010/09/22. Laboratory of Molecular Pharmacology, Center for Cancer Research, National Cancer Institute, Bethesda, MD 20892, USA.; 2010;107: 17992–17997. doi:10.1073/pnas.1008260107

69. Peshkin L, Wühr M, Pearl E, Haas W, Freeman Jr RM, Gerhart JC, Klein AM, Horb M, Gygi SP, Kirschner MW. On the relationship of protein and mRNA dynamics in vertebrate embryonic development. Developmental cell. 2015 Nov 9;35(3):383-94.

70. Schwanhausser B, Busse D, Li N, Dittmar G, Schuchhardt J, Wolf J, et al. Global quantification of mammalian gene expression control. Nature. 2011/05/20. Max Delbruck Center for Molecular Medicine, Robert-Rossle-Str. 10, D-13092 Berlin, Germany.; 2011;473: 337–342. doi:10.1038/nature10098