Plantarflexor Torque and Work is Positively Correlated with Medial Gastrocnemius

Fascicle Length in Healthy Young Adults

John F Drazan[1], Todd J Hullfish[1], Josh R Baxter[1]

[1]Department of Orthopedic Surgery, University of Pennsylvania, Philadelphia, Pennsylvania,

United States

Corresponding Author:

Josh R Baxter[1]

Human Motion Lab, 3737 Market Street, Philadelphia, Pennsylvania, 19104, United States

Email address: josh.baxter@uphs.upenn.edu

Key words:

Word count: 2628

Submission type: Original Research

ABSTRACT

Ankle kinetics are critical for ambulatory function in elite athletes, the elderly, and many
patient populations. Despite the robust findings linking plantarflexor muscle structure to gross
function with these populations, the link between plantarflexor fascicle length and ankle kinetics
has not been established in the literature. In this study, we determined the relationship between
muscle structure and peak torque and total work produced by the plantarflexors during maximal
effort contractions. We measured resting fascicle length and pennation angle of the medial
gastrocnemius using ultrasound in healthy adult subjects (n=12). Subjects performed
contractions on a dynamometer during isokinetic and isometric conditions. Longer fascicles
were positively correlated with higher peak torque and total work ($R^2 > 0.41$, $p < 0.013$) across
all isokinetic velocities. Higher pennation angles were negatively correlated with peak torque
and total work ($R^2 > 0.296$, $p < 0.067$). None of these correlations were significant in isometric
conditions. This provides experimental evidence demonstrating the link between plantarflexor
muscle structure and ankle kinetics in healthy young adults which has hitherto not been
experimentally demonstrated in existing literature.

INTRODUCTION

Torque generation about the ankle plays a crucial role human locomotion and is largely
determined by the structure of the plantarflexor muscles. Small variations in plantarflexor
structure explain functional differences in athletes (Abe et al., 2000; Kearns et al., 2000), the
elderly (Thom et al., 2007), and patients with Achilles tendon injuries (Hullfish et al., 2018).
Sprinters benefit from reduced mechanical advantage and longer muscle fascicle to accelerate
at the start of a sprint race (Abe et al., 2000; Lee and Piazza, 2009). Even amongst highly
trained sprinters, faster athletes have longer plantarflexor fascicles than their slower counter
parts (Abe et al., 2001; Kumagai et al., 2000). Conversely, elderly adults with limited mobility
benefit from increased plantarflexor leverage to generate the torques required to walk quickly
(Lee and Piazza, 2012). Achilles tendon ruptures stimulate rapid remodeling of the plantarflexor
muscles (Hullfish et al., 2018) that permanently reduce muscle shortening capacity and heel
raise function (Baxter et al., 2018).

Plantarflexor function is governed by active muscle shortening that acts about the ankle
joint to extend the foot. Plantarflexion power is better explained by joint leverage than muscle
mass, which suggests that muscle shortening demands are stronger determinants of
plantarflexor power than isometric muscle force (Nagano et al., 2007). Several critical
musculoskeletal parameters have been linked to plantarflexor torque and work capacity.
Plantarflexor muscle volume and moment arm are both correlates of peak isometric torque
(Baxter and Piazza, 2014). The effects of fascicle length on isokinetic muscle function have
been demonstrated in isolated muscle preparations (Lieber, 1997) and expanded to
plantarflexor function using computational musculoskeletal models (Baxter et al., 2019). The
force-velocity properties of skeletal muscle were first described by A.V. Hill in his seminal work
on muscle function (Hill, 1938). However, these foundational properties of skeletal muscle
mechanics relating fascicle length to force-velocity have not yet been demonstrated in the
physiologic context of the human plantarflexors (Ema et al., 2016).

The purpose of this study was to determine the relationship between muscle structural
parameters, which govern fascicle shortening speed, and plantarflexor function in healthy young

adults. Given that longer and less pennate fascicles increase the potential for total muscle
shortening velocity, we hypothesized longer fascicles be correlated with increased plantarflexor
torque and work and that increased pennation would be correlated with decreased plantarflexor
torque and work during isokinetic contractions. To test this hypothesis, we quantified the fascicle
length and pennation angle in the medial gastrocnemius using ultrasound imaging and
measured plantarflexion torque and work in maximal isometric and isokinetic conditions at three
rates of ankle rotation on an isokinetic dynamometer. Based on our previous computational
modeling (Baxter et al., 2019) , we hypothesized that fascicle length would be the strongest
correlate with plantarflexor kinetics, while pennation angle would be less strongly correlated.

METHODS

Plantarflexor muscle structure and function was quantified in 12 healthy young adults (6
male, age: 25 ± 4.54, BMI: 23.1 ± 4.48) who provided written informed consent in this IRB
approved by the University of Pennsylvania IRB (828374). Subjects were recreationally active
and had no reported history of Achilles tendon injury in either leg. We acquired measurements
of plantarflexor structure and function of the right leg with subjects positioned prone on an
isokinetic dynamometer (System 4, Biodex, Shirley, NY). Each subject was positioned on a
treatment table that was rigidly secured to the dynamometer while wearing standardized lab
shoes. After their foot was secured to the foot plate with the medial malleolus aligned with the
spindle of the dynamometer, each subject selected their ankle range of motion by fully
dorsiflexing and plantarflexing their ankle. Once the subject specific range of motion was set,
the investigator set the ankle neutral position, which was recorded for post-processing. To
ensure consistency between subjects, all experimental procedures were performed by a single
investigator.

To quantify medial gastrocnemius structure, we acquired ultrasound images of the
medial gastrocnemius throughout the entire passive range of motion of each subject. We
positioned the 7.5 MHz, 6 cm probe (LV7.5/60/128Z-2, SmartUs, TELEMED) over the mid-
substance of the muscle belly and secured it to the leg using a custom-made cast and strap
system. Ultrasound images were acquired between 30 and 60 Hz while each subject's ankle
was moved through its passive range of motion moving from plantarflexion to dorsiflexion at a
rate of 10 degrees/s. We quantified resting fascicle length, pennation angle, and muscle
thickness with the ankle at the resting angle. This resting angle was set to 16 degrees based on
values of average resting ankle position across 42 patients reported in the literature (Zellers et
al., 2018). We manually identified the superficial and deep aponeuroses and a muscle fascicle
by selecting two points on each structure to form a line (Fig. 1). This manual approach of
fascicle measurement has been demonstrated to be repeatable within the same observer
(Drazan et al., 2019). We calculated fascicle length as the distance between the superficial and
deep attachments of the fascicle and pennation angle as the angle between the fascicle and the
deep aponeurosis. We averaged each of these measurements of structure across three passive
range of motion trials to determine the measures of resting muscle structure we used in our
analyses.

To determine plantarflexor work and peak torque, subjects performed maximal voluntary
plantarflexor contractions on an isokinetic dynamometer in maximal isometric contractions at
neutral ankle angle and isokinetic contractions at three rotational speeds. The subject's right
foot was secured to a foot plate with the medial malleolus of the ankle aligned with the
dynamometer's spindle. First, we measured peak isometric plantarflexor torque with the ankle at

neutral with subjects in prone position. Next, peak isokinetic plantarflexion contractions were
performed throughout the entire range of motion at three speeds: slow (30 degrees per second),
medium (120 degrees per second), and fast (210 degrees per second). At the start of each of
these trials, we confirmed that the subject positioned their ankle in their peak dorsi-flexion angle.
We provided verbal encouragement (McNair et al., 1996) as well as visual feedback to ensure
that subjects maximally contract their plantarflexors during each condition. Subjects continued
to perform maximal plantarflexion contractions for each test condition until the peak torque was
consistent for two consecutive trials. We calculated peak torque for all of the contraction
conditions and total work for each of the three isokinetic conditions and selected peak torque
and maximal work values at each condition to test our research question.

To test our hypothesis that fascicle length would be positively and strongly correlated
with plantarflexor kinetics and that pennation angle would be negatively and moderately
correlated with plantarflexor kinetics, we performed univariate linear regression to determine the
relationship between measures of medial gastrocnemius structure with plantarflexor torque and
work at each contraction condition. We calculated the effect size of these regression analyses
using the coefficient of determination (R^2), which is an index of the correlation strength ranging
between 0 and 1 where values between 0 and 0.04 indicate negligible correlation, 0.04 and 0.25
indicate weak correlation, 0.25 and 0.64 indicate moderate correlation, and 0.64 and 1 indicate
strong correlation (Morton et al., 2005). We set an *a priori* alpha level of 0.05 and performed all
statistical analysis using scientific computing software (MATLAB, MathWorks, Natick, MA).

Because muscle structure and anthropometry varied in our subjects, we quantified
muscle thickness and lower-leg length to determine if these factors accounted for some of the
variability in plantarflexor kinetics. We calculated muscle thickness as the product of the resting
fascicle length and the sine of the pennation angle. We quantified lower-leg length by measuring
the distance between reflective markers placed on the lateral malleolus and the proximal head
of the fibula that was measured using a 12 camera motion capture system (Raptor Series,
Motion Analysis Corporation, Rohnert Park, CA). To evaluate the effect of patient stature on
resting muscle structure, we linearly regressed fascicle length, pennation angle, and muscle
thickness against body mass and leg length. Additionally, muscle thickness was linearly
regressed against fascicle length and pennation angle.

RESULTS

Fascicle length was positively and moderately correlated with plantarflexor work (Fig. 2A) and
peak torque (Fig. 3A) during isokinetic plantarflexion contractions. More than half of the
variability in plantarflexor work ($R^2 = 0.599$, $p = 0.003$) and peak torque ($R^2 = 0.521$, $p = 0.008$)
during 30°/s isokinetic contractions was explained by resting fascicle length. However, the
correlation between resting fascicle length and plantarflexor work and peak torque decreased
during faster isokinetic contractions at 120°/s ($0.413 > R^2 > 0.415$, $p < 0.024$) and 210°/s (0.477
$> R^2 > 0.4494$, $p < 0.013$). Fascicle length had the weakest correlation with peak torque during
isometric conditions ($R^2 = 0.325$, $p = 0.053$). Subjects generated less torque and did less work
as rotational velocity increased (Table 1).

Pennation angle was negatively and moderately correlated with plantarflexor work ($0.255 > R^2$
> 0.39, $p < 0.052$) and peak torque ($0.296 > R^2 > 0.417$, $p < 0.067$ during isokinetic
plantarflexion contractions (Fig. 2B and Fig. 3B). However, these correlations had a smaller
effect size than resting fascicle length for each test condition and only reached statistical

significance during 30°/s (p = 0.047) and 210 °/s (p = 0.023) conditions for measurements of
peak torque. Peak isometric torque was not explained by resting pennation angle (R^2 = 0.09, p =
0.345).

Muscle thickness was moderately and significantly correlated with fascicle length (R^2 = 0.536, p
< 0.001) but not correlated with pennation angle (R^2 = 0.064, p = 0.084). Despite the positive
correlation between resting muscle thickness and fascicle length, resting muscle thickness was
not significantly correlated with any isokinetic testing condition (Table 2). Muscle thickness was
positively and weakly correlated with peak torque at isometric max (R^2 = 0.161). However, this
correlation did not reach statistical significance (p = 0.197).

Muscle structure was weakly correlated with subject stature (Table 3). Fascicle length was
weakly correlated with leg length (R^2 = 0.084, p = 0.046) and body mass (R^2 = 0.132, p = 0.011).
Pennation angle was weakly correlated with body mass (R^2 = 0.098, p = 0.030) but not leg
length. Muscle thickness was not correlated with either leg length or body mass.

DISCUSSION

In this study, we established the relationship between plantarflexor structure and function in a
cohort of healthy young adults. Our findings support our hypothesis that resting fascicle length is
positively correlated with plantarflexor function while pennation angle has a smaller, negative
correlation with plantarflexor function. To determine if plantarflexor function was simply
explained by muscle thickness, we regressed muscle thickness with plantarflexor kinetics and
found no effect on plantarflexor function in any condition. Although muscle kinetics have been
shown to be positively affected by increased fascicle length in isolated muscle experiments
(Lieber, 1997); to our knowledge, this is the first study to link natural variation in gastrocnemius
fascicle length with plantarflexor kinetics in healthy adults (Ema et al., 2016).

Our measurements of medial gastrocnemius structure and plantarflexor kinetics compare
favorably with previous reports in the literature. We decided to measure resting muscle structure
at 16 degrees plantarflexion to approximate muscle-tendon slack (Zellers et al., 2018). Other
measurements of gastrocnemius structure were acquired with the ankle either under load or
neutrally aligned which may explain slightly longer (4-9 mm) and less pennate (6-11 degrees)
muscle fascicles (Baxter and Piazza, 2014; Kubo et al., 2003; Thom et al., 2007) than those we
measured in this current study. Similarly, our measurements of pennation at slack are similar to
earlier measurements in the medial gastrocnemius at the same position (Hoang et al., 2007).
Our measurements of plantarflexion torque and work capacity compares well with the literature.
Our values for torque and work done at 30 and 120 degrees per second compares well with
previously reported values (Randhawa and Wakeling, 2013; Woodson et al., 1995) and our
values for maximal isometric torque compares well with studies with similar subject positioning
(Arampatzis et al., 2006). A previous study reported higher values for torque generation,
however in this case, the subjects were all male and were seated rather than positioned in
prone (Baxter and Piazza, 2014).

As hypothesized, fascicle length and pennation angle, which govern absolute muscle shortening
velocity (Lieber and Ward, 2011), have greater effects on isokinetic plantarflexor function than
muscle thickness. The results are intuitive within our understanding of the force-velocity
properties of muscle. While fascicles of different lengths undergo the same absolute contractile
velocity during a given isokinetic contraction, longer fascicles have additional sarcomeres in
series which reduce relative shortening velocity. This enables longer fascicles to operate at

slower velocities on the force-velocity curve, increasing force production at all isokinetic speeds
(Fig. 4A) (Lieber and Fridén, 2000). Similarly, the force-velocity effects are also affected by
variation in pennation angle. Greater muscle pennation increases the fascicle shortening
demands for a given muscle shortening contraction. Thus, more pennate muscles generate less
force and do less work in isokinetic conditions (Fig. 4B). Our experimental findings that fascicle
length explained more variation in plantarflexor kinetics than pennation angle agrees with the
theoretical framework of the force-velocity properties of muscle and our recent computational
model (Baxter et al., 2019).

Our results are notable because while the force-velocity properties of skeletal muscle have been
well defined in isolated muscle preparations in animal models, the effects of natural variability in
muscle fascicle lengths on kinetic function have not been reported in humans. Force-length
characteristics have been demonstrated in humans in vivo (Maganaris, 2001). Despite evidence
of variations in muscle structure between different populations of human subjects (Abe et al.,
2000; Hullfish et al., 2018; Kearns et al., 2000; Kubo et al., 2003; Lee and Piazza, 2009), a
recent review of the literature on muscle structure and function concluded that there were no
studies reporting the relationship between fascicle length and muscle shortening velocity in
humans (Ema et al., 2016). Another study found that elbow extensor velocity under no load was
positively correlated with muscle volume and pennation angle but not muscle length (Wakahara
et al., 2013). However, this previous study on elbow extensor kinetics did not correlate muscle
structure with maximal isokinetic contractions under load and also did not directly measurement
fascicle length.

This study was affected by several limitations. Despite having a small sample size, our study
results are in strong agreement with both basic principles of skeletal muscle mechanics (Lieber
and Fridén, 2000) and computational simulations (Baxter et al., 2019). We did not confirm
maximal effort during contraction using electromyography. Instead, we provided subjects verbal
encouragement and real-time visual feedback and had each subject repeat each contraction
condition until their peak torques were consistent between consecutive trials. We used
ultrasound to measure fascicle length and pennation angle but did not quantify muscle volume
or moment arm. Our recent work found that moment arm measured with ultrasound imaging is
not a good surrogate measure of geometric moment arm (Baxter and Piazza, 2018). We used
muscle thickness as a proxy for muscle volume, however this yielded weak and insignificant
correlations. Additionally, we did not quantify the lateral gastrocnemius or soleus muscle. While
the lateral gastrocnemius functions similarly to the medial muscle belly (Héroux et al., 2016), the
soleus muscle is a uniarticulate muscle with different mechanics during plantarflexor
contractions (Bojsen-Møller et al., 2004). However, our findings appear to be robust given that
more than 40% of plantarflexor kinetics during plantarflexion rotations were explained by
variation in fascicle lengths of the medial gastrocnemius muscle.

In conclusion, our study demonstrates the link between resting structure of the medial
gastrocnemius with isokinetic plantarflexion function. These findings may have important
implications on plantarflexor function following muscle remodeling elicited either through injury
or training. However, isokinetic plantarflexor function has limited translation to activities of daily
living, and future work should directly test the link between muscle structure and movement
biomechanics.

ACKNOWLEDGEMENTS

This work was supported by the Thomas B. and Jeannette E. Laws McCabe Fund and NIH grant K12GM081259. We would like to thank R. Mathew and S. Donde for assistance with data collection and processing.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to disclose.

FIGURE CAPTIONS

Figure 1 (1 column). Resting muscle architecture was measured from an ultrasound frame captured when each subject's ankle was in 16° plantarflexion. An examiner identified the two aponeuroses as well as a fascicle within the frame, each of which was approximated as a straight line. Pennation angle was calculated as the angle between the fascicle and the deep aponeurosis and muscle thickness was calculated as the sine of the pennation angle times the length of the fascicle.

Figure 2 (1 column). Plantarflexor work produced at three isokinetic speeds (30°/s – *top row*, 120°/s – *middle row*, and 210°/s – *bottom row*) is positively correlated with (*A*) fascicle length and negatively correlated with (*B*) pennation angle. Fascicle lengths explained more variation in plantarflexor work than pennation angle.

Figure 3 (1 column). Peak plantarflexor torque produced at three isokinetic speeds (30°/s – *second row*, 120°/s – *third row*, and 210°/s – *bottom row*) is positively correlated with (*A*) fascicle length and negatively correlated with (*B*) pennation angle. Conversely, fascicle length explains less variation in peak isometric torque (*top row*) and pennation angle is not correlated with isometric torque. Fascicle lengths explained more variation in plantarflexor work than pennation angle.

Figure 4 (1 column). (*A*) Longer and (*B*) less pennate fascicles decrease relative shortening velocity during a given joint rotation, which results in increased fascicle force. The force-velocity properties of skeletal muscle are more sensitive to changes in fascicle length than pennation angle, which agrees with our experimental findings that ankle torque is more strongly correlated with fascicle length than pennation angle.

269 Table 1. Descriptive data on subjects,
270 muscle structure, and plantarflexion
271 function.

	Mean ± Standard Deviation
Plantarflexor Kinetics	
Peak Isometric Torque (N*m)	116.82 ± 39.13
Peak Isokinetic Torque 30°/s (N*m)	95.12 ± 31.33
Peak Isokinetic Torque 120°/s (N*m)	50.09 ± 23.96
Peak Isokinetic Torque 210°/s (N*m)	35.87 ± 16.95
Total Work 30°/s (J)	53.7 ± 20.01
Total Work 120°/s (J)	33.75 ± 17.31
Total Work 210°/s (J)	26.64 ± 12.94
Musculoskeletal Parameters	
Range of Motion (degrees)	53.70 ± 4.17
Fascicle Length (mm)	48.00 ± 10.02
Pennation Angle (degrees)	29.52 ± 4.06
Muscle Thickness (mm)	23.15 ± 4.15
Leg Length (mm)	362.2 ± 36.88
Mass (kg)	67.26 ± 19.25

277   Table 2. Correlations between muscle thickness and plantarflexion kinetics

	Peak Torque	Total Work
Isometric Max	0.153 (0.209)	-
Isokinetic 30 °/s	0.124 (0.262)	0.172 (0.181)
Isokinetic 120 °/s	0.104 (0.307)	0.117 (0.276)
Isokinetic 210 °/s	0.076 (0.387)	0.089 (0.347)

287   Correlations of determination (R^2) and regression p-values are reported as R^2 (p-value)

289   Table 3. Correlations between muscle structure and subject stature.

	Leg Length	Mass	Muscle Thickness
Fascicle Length	0.084 (0.046)	0.132 (0.011)	0.53 (<0.0001)
Pennation Angle	0.013 (0.445)	0.098 (0.030)	0.063 (0.083)
Muscle Thickness	0.070 (0.069)	0.037 (0.193)	-

299   Correlations of determination (R^2) and regression p-values are reported as R^2 (p-value)

REFERENCES

Abe, T., Fukashiro, S., Harada, Y., Kawamoto, K., 2001. Relationship between sprint performance and muscle fascicle length in female sprinters. J. Physiol. Anthropol. Appl. Human Sci. 20, 141–147.

Abe, T., Kumagai, K., Brechue, W.F., 2000. Fascicle length of leg muscles is greater in sprinters than distance runners. Med. Sci. Sports Exerc. 32, 1125–1129.

Arampatzis, A., Karamanidis, K., Stafilidis, S., Morey-Klapsing, G., DeMonte, G., Brüggemann, G.-P., 2006. Effect of different ankle- and knee-joint positions on gastrocnemius medialis fascicle length and EMG activity during isometric plantar flexion. J. Biomech. 39, 1891–1902. https://doi.org/10.1016/j.jbiomech.2005.05.010

Baxter, J.R., Farber, D.C., Hast, M.W., 2019. Plantarflexor fiber and tendon slack length are strong determinates of simulated single-leg heel raise height. J. Biomech. https://doi.org/10.1016/j.jbiomech.2019.01.035

Baxter, J.R., Hullfish, T.J., Chao, W., 2018. Functional deficits may be explained by plantarflexor remodeling following Achilles tendon rupture repair: Preliminary findings. J. Biomech. 79, 238–242. https://doi.org/10.1016/j.jbiomech.2018.08.016

Baxter, J.R., Piazza, S.J., 2018. Plantarflexor moment arms estimated from tendon excursion in vivo are not strongly correlated with geometric measurements. J. Biomech. 77, 201–205. https://doi.org/10.1016/j.jbiomech.2018.06.020

Baxter, J.R., Piazza, S.J., 2014. Plantar flexor moment arm and muscle volume predict torque-generating capacity in young men. J. Appl. Physiol. 116, 538–44. https://doi.org/10.1152/japplphysiol.01140.2013

Bojsen-Møller, J., Hansen, P., Aagaard, P., Svantesson, U., Kjaer, M., Magnusson, S.P., 2004. Differential displacement of the human soleus and medial gastrocnemius aponeuroses during isometric plantar flexor contractions in vivo. J. Appl. Physiol. 97, 1908–1914. https://doi.org/10.1152/japplphysiol.00084.2004

Drazan, J.F., Hullfish, T.J., Baxter, J.R., n.d. An automatic fascicle tracking algorithm quantifying gastrocnemius architecture during maximal effort contractions. https://doi.org/10.7287/peerj.preprints.27475v1

Ema, R., Akagi, R., Wakahara, T., Kawakami, Y., 2016. Training-induced changes in architecture of human skeletal muscles: Current evidence and unresolved issues. J. Phys. Fit. Sports Med. 5, 37–46. https://doi.org/10.7600/jpfsm.5.37

Héroux, M.E., Stubbs, P.W., Herbert, R.D., 2016. Behavior of human gastrocnemius muscle fascicles during ramped submaximal isometric contractions. Physiol. Rep. 4, n/a-n/a. https://doi.org/10.14814/phy2.12947

Hill, A.V., 1938. The Heat of Shortening and the Dynamic Constants of Muscle. Proc. R. Soc. B Biol. Sci. 126, 136–195. https://doi.org/10.1098/rspb.1938.0050

Hoang, P.D., Herbert, R.D., Todd, G., Gorman, R.B., Gandevia, S.C., 2007. Passive mechanical properties of human gastrocnemius muscle tendon units, muscle fascicles and tendons in vivo. J. Exp. Biol. 210, 4159–4168. https://doi.org/10.1242/jeb.002204

Hullfish, T.J., O'Connor, K.M., Baxter, J.R., 2018. Gastrocnemius fascicles are shorter and more pennate immediately following acute Achilles tendon rupture. BioRxiv Prepr. 445569. https://doi.org/10.1101/445569

Kearns, C.F., Abe, T., Brechue, W.F., 2000. Muscle enlargement in sumo wrestlers includes increased muscle fascicle length. Eur. J. Appl. Physiol. 83, 289–296. https://doi.org/10.1007/s004210000298

Kubo, K., Kanehisa, H., Azuma, K., Ishizu, M., Kuno, S.-Y., Okada, M., Fukunaga, T., 2003. Muscle Architectural Characteristics in Young and Elderly Men and Women. Int. J. Sports Med. 24, 125–130. https://doi.org/10.1055/s-2003-38204

Kumagai, K., Abe, T., Brechue, W.F.W.F., Ryushi, T., Takano, S., Mizuno, M., 2000. Sprint
performance is related to muscle fascicle length in male 100-m sprinters. J. Appl.
Physiol. 88, 811–816. https://doi.org/10.1097/00005768-199905001-01648
Lee, S.S.M., Piazza, S.J., 2012. Correlation between plantarflexor moment arm and preferred
gait velocity in slower elderly men. J. Biomech. 45, 1601–1606.
https://doi.org/10.1016/j.jbiomech.2012.04.005
Lee, S.S.M., Piazza, S.J., 2009. Built for speed: musculoskeletal structure and sprinting ability.
J. Exp. Biol. 212, 3700–3707. https://doi.org/10.1242/jeb.031096
Lieber, R.L., 1997. Muscle fiber length and moment arm coordination during dorsi- and
plantarflexion in the mouse hindlimb. Acta Anat. (Basel) 159, 84–89.
Lieber, R.L., Fridén, J., 2000. Functional and clinical significance of skeletal muscle
architecture. Muscle Nerve 23, 1647–1666.
Lieber, R.L., Ward, S.R., 2011. Skeletal muscle design to meet functional demands. Philos.
Trans. R. Soc. Lond. B. Biol. Sci. 366, 1466–1476.
https://doi.org/10.1098/rstb.2010.0316
Maganaris, C.N.C.C.N., 2001. Force-length characteristics of in vivo human skeletal muscle.
Acta Physiol Scand 172, 279–285.
McNair, P.J., Depledge, J., Brettkelly, M., Stanley, S.N., 1996. Verbal encouragement: effects
on maximum effort voluntary muscle: action. Br. J. Sports Med. 30, 243–245.
https://doi.org/10.1136/bjsm.30.3.243
Morton, R.F., Hebel, J.R., McCarter, R.J., 2005. A Study Guide to Epidemiology and
Biostatistics. Jones & Bartlett Learning.
Nagano, A., Yoshioka, S., Komura, T., Fukashiro, S., 2007. Influence of the Fascicle Length and
Physiological Cross-sectional Area of M. soleus on Ankle Joint Kinetics. Int. J. Sport
Health Sci. 5, 98–104. https://doi.org/10.5432/ijshs.5.98
Randhawa, A., Wakeling, J.M., 2013. Associations between muscle structure and contractile
performance in seniors. Clin. Biomech. 28, 705–711.
https://doi.org/10.1016/j.clinbiomech.2013.04.010
Thom, J.M., Morse, C.I.., Birch, K.M., Narici, M.V., 2007. Influence of muscle architecture on the
torque and power-velocity characteristics of young and elderly men. Eur. J. Appl.
Physiol. 100, 613–619. https://doi.org/10.1007/s00421-007-0481-0
Wakahara, T., Kanehisa, H., Kawakami, Y., Fukunaga, T., Yanai, T., 2013. Relationship
between Muscle Architecture and Joint Performance during Concentric Contractions in
Humans. J. Appl. Biomech. 29, 405–412. https://doi.org/10.1123/jab.29.4.405
Woodson, C., Bandy, W.D., Curis, D., Baldwin, D., 1995. Relationship of Isokinetic Peak Torque
With Work and Power for Ankle Plantar Flexion and Dorsiflexion. J. Orthop. Sports Phys.
Ther. 22, 113–115. https://doi.org/10.2519/jospt.1995.22.3.113
Zellers, J.A., Carmont, M.R., Silbernagel, K.G., 2018. Achilles Tendon Resting Angle Relates to
Tendon Length and Function. Foot Ankle Int. 39, 343–348.
https://doi.org/10.1177/1071100717742372
391

Superficial Aponeurosis

Fascicle Length

Pennation Angle

Deep Aponeurosis

A Fascicle Shortening Speed

B Fascicle Shortening Speed