Surpassing thermodynamic, kinetic, and stability barriers to isomerization catalysis for tagatose biosynthesis.

Abstract:

There are many enzymes that are relevant for making rare and valuable chemicals that while active, are severely limited by thermodynamic, kinetic, or stability issues (e.g. isomerases, lyases, transglycosidase etc.). In this work, we study an enzymatic reaction system − *Lactobacillus sakei* L-arabinose isomerase (LsLAI) for D-galactose to D-tagatose isomerization – that is limited by all three reaction parameters. The enzyme has a low catalytic efficiency for non-natural substrate galactose, has low thermal stability at temperatures $> 40\ °C$, and equilibrium conversion $< 50\%$. After exploring several strategies to overcome these limitations, we finally show that encapsulating the enzyme in a gram-positive bacterium (*Lactobacillus plantarum*) that is chemically permeabilized can enable reactions at high rates, high conversion, and at high temperatures. The modified whole cell system stabilizes the enzyme, differentially partitions substrate and product across the membrane to shift the equilibrium toward product formation enables rapid transport of substrate and product for fast kinetics. In a batch process, this system enables approximately 50 % conversion in 4 h starting with 300 mM galactose (an average productivity of 37 mM/h), and 85 % conversion in 48 h, which are the highest reported for food-safe mesophilic tagatose synthesis. We suggest that such an approach may be invaluable for other enzymatic processes that are similarly kinetically-, thermodynamically-, and/or stability-limited.

D-Tagatose is a rare ketohexose sugar with sweetness similar to that of sucrose. However,

its glycemic index and caloric value is much lower because of low bioavailability, making

it an attractive GRAS (generally regarded as safe) sugar substitute. Recent studies have also

demonstrated that it is anti-hyperglycemic [1] and prebiotic, which promotes gut health [2,3].

Thus, there exists a high demand in food industry for the economical production of rare

sugars, like tagatose, exemplified by the 2016 global artificial sweetener market estimated

to be USD 3.2 billion. This market is expected to expand given the global diabetes crisis

and increasing relevance of prebiotics.

The enzyme L-arabinose isomerase (LAI) that responsible for the reversible isomerization

of the pentose L-arabinose to L-ribulose can also isomerize the hexose D-galactose to D-

tagatose [4,5]. LAI has thus been the enzyme of choice to produce tagatose, although, to date,

few commercial bioprocesses exist. A variety of LAIs from different microorganisms have

been isolated and have reported optimal activity at a range of temperatures and pH [4,6,7].

Some of the limitations of tagatose biosynthesis using LAI that may be hindering

commercial viability are, 1) unfavorable enzymatic kinetics since galactose is not the native

substrate of LAI, 2) low enzyme stability, particularly in the absence of divalent metal ions,

and 3) low equilibrium constant for galactose to tagatose isomerization.

Few previous reports have been successful at engineering enzymatic properties of LAI for

industrial application; often addressing only one of the bottlenecks to productivity. To

address the kinetic issue, several groups have used enzyme engineering methods to enhance

catalytic efficiency of LAI toward galactose and have shown moderate increases in

productivity [8–13]. To counter low-stability issues, many groups have tested the utility of

thermophilic enzymes [12,14–16]. However, most thermophilic enzymes rely on divalent metal

ions (Mn^{2+}, Co^{2+}, Fe^{2+}) for stability[17], and high reaction temperatures ($\geq$ 80 °C) result in

significant caramelization [18], which are all undesirable and must be removed from product,

adding to processing costs. Surface-display [19] or encapsulation in particles or whole-cells

[20–23] can stabilize enzymes [24]. Finally, the thermodynamic limitations of isomerization of

galactose to tagatose are severe and, arguably, the most recalcitrant issue since $\Delta G°_{rxn} \approx$

+1.19 kcal/mol [23], which indicates theoretical maximum equilibrium conversion ~ 14% at

room temperature. Several approaches have been used to overcome this limitation.

Thermophilic enzymes can achieve higher conversions than mesophilic enzymes since the

equilibrium shifts toward tagatose at higher temperatures [6]. Whole-cell biocatalysts with

GRAS organisms (e.g. lactic acid bacteria (LAB) and *E. coli*) that disproportionately

partition substrate and product across their membrane has also been shown to partially

circumvent this thermodynamic limitation while simultaneously enhancing enzyme

stability; albeit at a kinetic penalty imposed by substrate transport limitations [25–27].

Recently, cell permeabilization [27] and sugar transport overexpression [28,29] were

demonstrated as methods to overcome the kinetic penalty imposed by cellular

encapsulation.

There have currently been no studies that look to systematically analyze all three

limitations – kinetic, thermodynamic, and enzyme stability – of the enzymatic

isomerization of galactose to tagatose. This work clearly demonstrates the presence of these

three limitations and provides a novel approach to balance their advantages and limitations.

We use the food-safe engineered probiotic bacterium *Lactobacillus plantarum* as the

expression host due to its increasing relevance to biochemical and biomedical research [30,31].

This approach enabled ~ 50 % conversion of galactose to tagatose in 4 h (productivity of ~

38 mmol tagatose $L^{-1} h^{-1}$) ultimately reaching ~ 85% conversion after 48 h at high galactose

loading (300 mM) in batch culture. This is among the highest conversions and

productivities reported to date for tagatose production using a mesophilic enzyme. Such an

approach is expected to be applicable to other biocatalytic systems where similar trade-offs

between kinetics, thermodynamics, and/or stability pose hurdles to process development.

Results:

Characterizing *L. sakei* LAI (LsLAI) limitations.

At high substrate loading (400mM galactose) and 37 °C, the reported optimal temperature

for this enzyme [32], LsLAI purified from *L. plantarum* exhibited an initial forward reaction

(turnover) rate ($k_{initial}$) of 9.3 ± 0.3 s^{-1}, which is lower than the maximum reaction rate

possible by this enzyme of 17.0 ± 1.3 s^{-1} min^{-1} with its preferred substrate, arabinose, at 400

mM. Increasing the reaction temperature to 50 °C increased the initial reaction rate to 11.2

s^{-1} (Fig. 1a) but was accompanied by rapid enzyme inactivation, which is consistent with

previous reports of thermal instability of this enzyme [32]. The enzyme exhibited first-order

degradation with a half-life ($t_{½}$) of approximately 18.5 h at 37 °C (Fig. 1b) and 1.5 h at 50

°C. Indeed, the enzyme retaining only ~ 6 % activity after 6 h at this elevated temperature.

Additionally, the highly reversible isomerization reaction suffers from thermodynamic

limitations with reported free-enzyme equilibrium conversions < 50 % ($\Delta G°_{rxn} = +1.19$ kcal

mol^{-1}) [23]. Our results indicate similar low conversion of 35 – 40 % after 5 days with pure

enzyme at 37 °C with daily supplementation of fresh enzyme to account for inactivation

(Fig. 1c).

Cell surface display is unsuitable to stabilize LsLAI.

To achieve surface localization, we fused LsLAI with six *L. plantarum* surface proteins,

which have all been previously described as suitable display carriers, at either the C- or N-

terminus (Table S1). The constructs also contained the strong Lp_3050 secretion signal, a

10 amino acid linker containing a thrombin protease cleavage site, and either a C- or N-

terminal His_6 tag for immunofluorescence assays. We characterized surface localization

using flow cytometry and measured the whole-cell isomerization activity. All three of the

tested C-terminal anchor strains (A1, A2, and A3) exhibited > 90 % surface detection (Fig.

2a). However, the measured tagatose production after a 2 h incubation with 200 mM

galactose was only 225 ± 72, 55 ± 39, 91 ± 25 µM tagatose (normalized to individual

reaction cell densities of ~ $OD_{600} = 0.5$), respectively, significantly less than that of

intracellularly expressed (I) LsLAI (538 ± 34 µM) (Fig. 2b). LsLAI was detected on N-

terminal anchor strains A5 and A6 in 28 % and 96 % of the population and produced $286 \pm$

31 and 422 ± 34 µM tagatose respectively. Interestingly, A4 exhibited low surface

detection, comparable to control (WT & IC1) cells while producing the highest titers of

tagatose (533 ± 58 µM) among all the surface display constructs. These data indicate there

is not a significant correlation ($p > 0.05$) between whole-cell activity with the presence of

displayed LsLAI on the *L. plantarum* cell surface (Fig. 2c).

Next, we sought to explain the observation regarding absence of positive correlation

between display level and activity using strain A6. Sodium dodecyl sulfate (SDS) is a

strong anionic surfactant commonly used in laboratory techniques to lyse cells through

membrane disruption and denature proteins by destabilizing non-covalent bonds. We

posited that SDS could be used to extract surface proteins from Gram-positive bacteria

without lysing the cells. We incubated strains IC1 and A6 with either PBS (control) or PBS

containing 0.5 % SDS, to remove surface displayed LsLAI, and then quantified surface

detection and whole-cell activity. As expected, we detected LsLAI only on the surface of

PBS-treated control A6 but not the SDS-treated cells by flow cytometry (Fig. S1a).

Interestingly, whole-cell activity of both IC1 and A6 increased approximately 3- and 5-fold,

respectively, post-SDS-treatment. These data suggest LsLAI is present in both the

intracellular compartment and membrane bound of A6 with the vast majority of measured

whole-cell activity coming from the intracellular fraction. We confirmed the presence of

soluble anchor-LAI fusion (A6-LsLAI) in the intracellular fraction of A6 via Western blot

analysis (Fig. S1b). Additionally, we noticed enhanced whole-cell activity that we attribute

to permeabilization/fluidization of the membrane due to removal of surface proteins. While

this study only focused one of our six explored anchor protein constructs, we suspect that

these results are consistent amongst the other LsLAI surface display constructs. Thus, the

measured whole-cell activity differing between constructs was likely dependent on the

activity of soluble anchor-LsLAI present in the cytoplasm and not due to surface displayed

enzyme.

Finally, to further elucidate the discrepancy between activity of intracellular and surface

displayed LsLAI constructs we tested the enzymes activity of secreted, unanchored

enzyme. Post-induction, we separated whole-cells from culture media and compared the

activity of each fraction. We observed a 6-fold higher in activity in the whole-cell fraction

of the secretion strain compared to wild-type control cells; however, there was no

detectable activity in the supernatant of either strain (Fig. S2a). We confirmed the presence

of LsLAI in the supernatant and soluble cell fraction via Western blot analysis (Fig. S2b).

These data, taken together with our previous experiments, suggest LsLAI is largely in an

inactive state after secretion and that any activity is due to accumulation of active enzyme

in the *L. plantarum* cytoplasm.

Encapsulating LAI improves stability and conversion.

Since surface display failed as a stabilization mechanism, yielding minimally active

enzyme, we evaluated whether whole-cell *L. plantarum* cells encapsulating LsLAI could be

used as a stabilization method instead. Our data shows that such encapsulation within *L.*

plantarum protected the enzyme from thermal deactivation at 50 °C; the enzyme retained ~

85 % of its initial activity after 24 h (Fig. 3a). Interestingly, resting whole-cells at 37 °C

displayed no loss of activity after 24 h; in fact, we observed a slight enhancement in

activity. This observation could be due to changes to cellular morphology or physiology.

Encapsulated LsLAI quickly reached an equilibrium conversion of approximately 60 % in

24 h, overcoming the thermodynamic limitation of < 50 % conversion presented in the pure

enzyme system (Fig. 3b). We believe this happens because the substrate (galactose) and

product (tagatose) differentially partition across the cell membrane. Although initial

reaction rates of encapsulated LsLAI at 37 °C and 50 °C with either galactose or tagatose

were lower than that of their free-enzyme counterparts, encapsulated LsLAI favored

reaction in the forward direction more than in the reverse direction, as determined by initial

rates (Fig. S3). At 50 °C the ratio of forward to reverse initial reaction rate for encapsulated

LsLAI was 1.8 whereas for the pure enzyme, the ratio was more unfavorable at 0.7. While

encapsulation was shown to provide thermal stability and a thermodynamic advantage, it

imposes a kinetic penalty due to transport limitations across the cell membrane. To test this,

we measured initial whole-cell activity under a range of induction levels to determine the

maximum achievable activity and compared the whole-cell activity to that of the cell lysate

at the same conditions. Strain I produced the maximum amount of tagatose (1.5 mM) in 2 h

at inducer concentrations ≥ 5 ng/mL (Fig. 3c), indicating that this represented that the

system is at maximum reaction rate under all these inducer levels. Conversely, lysate of

cells induced at 5 and 25 ng/mL inducer conditions generated approximately 4-fold greater

tagatose than that of the equivalent whole-cell catalysts in the same 2 h interval. This

supports the idea that cellular encapsulation limits reaction kinetics.

Permeabilizated cells overcome all three barriers.

Seeking to overcome the transport-limited whole-cell production of tagatose, we

investigated the use of permeabilization surfactants − Triton X-100 and SDS. Following the

same optimized protocol as previously reported for permeabilization of *L. plantarum* with 1

% Triton X-100 [27], we observed a 4.2 ± 0.1 fold increase in tagatose production as

compared to untreated cells under the tested conditions (Fig. S4a). We assessed the ability

of 0.005 % – 0.5 % SDS to permeabilize encapsulated LsLAI cells and enhance tagatose

production. Our results showed the optimal concentration of SDS that enhanced tagatose

production via cell permeabilization while still retaining LsLAI intracellularly to be 0.01 %

(Fig. S4b). Tagatose production was enhanced 5.7 ± 0.1 fold after permeabilization with

0.01 % SDS, greater than that of treatment with 1% Triton X-100 under the tested

conditions. Additionally, this concentration of SDS is below the critical micellular

concentration (CMC = 0.24 – 0.26 % at 35 – 40 °C) therefore producing minimal frothing

as compared to 1 % Triton X-100, the CMC of which is reported to be 0.05 % [33]. Although

SDS is commonly used to denature proteins, low concentrations of SDS (0.001 – 0.005 %)

have been previously used to permeabilize *E. coli* – albeit less successfully than with

CTAB (cetyl trimethylammonium bromide) and Triton X-100 [34].

To further understand the effect of SDS-treatment on cell permeability, we measured the

fluorescence generation rate using 5-carboxyfluorescein diacetate (cFDA) as a substrate.

cFDA is an uncharged ester that becomes fluorescent upon hydrolysis by abundant and

nonspecific intracellular esterases. Hydrolysis also results in entrapment of the fluorescent

dye in the cytoplasm, unless the membrane has been damaged. The rate of generated

fluorescence serves as a proxy to determine substrate transport rates assuming hydrolysis

rates are higher than transport rate. Wild-type *L. plantarum* treated with 0.005, 0.01, and

0.05 % had 1.6-, 3.0-, and 3.4-fold increase in substrate transport rates compared to

untreated cells (Fig. S5). SDS concentrations > 0.5 % resulted in fluorescence leakage, a

sign of damaged cell membrane that may also be expelling other cytoplasmic content.

These data suggest that the treatment with 0.001 % SDS maintains cellular integrity while

increasing tagatose synthesis rate due to a decreased kinetic penalty presented upon

encapsulation.

Finally, we monitored the tagatose production with different LsLAI preparations at high

loading (OD_{600} = 40 or equivalent 0.6 mg/mL LsLAI) in batch processes starting with 300

mM galactose. Consistent with our initial experiments, the pure enzyme preparation was

limited to 39 ± 2.3 % conversion after 48 h at 37 °C (Fig. 4). Increasing the temperature of

the reaction to 50 °C allowed for high initial production of tagatose; however, the enzyme

denatured quickly reaching a maximum conversion of only 16 ± 2.8 %. Bacterial

encapsulation of LsLAI produced 47 ± 3.3 % tagatose at 37 °C, improvement over the

pure-enzyme system. Encapsulation also provided thermal stability to the enzyme allowing

for sustained activity at 50 °C increasing both the rate of production (46 ± 2.3 % in 6 h) and

equilibrium conversion (83 ± 6.1 %) compared to unmodified whole-cell catalyst at 37 °C.

Rate of production was further increased by modifying the whole-cell catalyst through SDS

permeabilization to achieve 59 ± 3.5 % conversion in 6 h, ultimately reaching a similar

equilibrium conversion of 85 ± 6.7 % as unpermeabilized cells. Thus, permeabilized, cell-

encapsulated LAI overcomes the three reaction limitations by supporting greater thermal

stability, higher equilibrium conversion, and faster reaction rates than possible by a free

enzyme system.

Discussion:

While an increasing number of enzymes are being used for biocatalysis, poor stability

under reaction conditions, and unfavorable kinetic properties toward non-native substrates

can limit their utility toward biosynthetic functions. Further, for certain classes of enzymes

like isomerases, transglycosidases, and lyases, limitations of endergonic or mild exergonic

thermodynamics are additional hurdles to economical processes. D-Tagatose synthesis from

D-galactose using the enzyme LAI (L-arabinose isomerase) suffers from all three of these

limitations. Our results confirm the presence of all three limitations (stability,

thermodynamic, kinetic) during enzymatic conversion of galactose to tagatose using the

acid-tolerant mesophilic LAI from *Lactobacillus sakei* (LsLAI). At low temperatures (37

°C), the enzyme is stable and quickly reaches equilibrium. However, the product titer is low

due to low equilibrium conversion (36 − 39 %) at low temperatures (this study and

previously reported [32]). At elevated temperatures (50 °C), the initial reaction rates are

faster, but the enzyme quickly inactivates and is unable to realize the higher conversion and

product titer possible at this temperature. These limitations of LAI for tagatose synthesis

have been realized previously, and each issue has been addressed individually. While some

successes have been reported, especially by selecting or engineering enzymes with more

desirable properties [4,11,35–38], this is the first study to address and overcome all three issues

simultaneously.

LAI display on *Bacillus subtilis* endospore surface has shown to improve thermostability

while maintaining free enzyme-like kinetic properties [19]. Therefore, this approach addresses

the stability limitations but not the thermodynamic issue. To assess whether cell surface

display on *L. plantarum* could improve stability of LsLAI, we tested six previously

reported anchor proteins and found there was no correlation linking whole-cell activity and

surface display level/accessibility. We showed evidence for presence of cytoplasmic

LsLAI-anchor fusions and concluded that enzymatic activity seen was largely due to this

active fraction and that the enzyme displayed on the surface was largely inactive. When we

tested secretion constructs (i.e., LsLAI with secretion signal but no anchor), we found

similar results, suggesting that LsLAI cannot fold extracellularly upon secretion through

this pathway and that active enzyme accumulated cytoplasmically. This is surprising since

there is no precedence for such an observation in literature. *L. plantarum* utilizes the Sec-

pathway to secrete proteins [39] and it is generally accepted that translocation occurs co-

translationally[40,41]; therefore, for a protein to be active it must fold properly in the

extracellular environment. While we detected LsLAI on the surface using flow cytometry

and Western blot analysis, we could not correlate display level to activity or subsequently

attribute any detectable activity to the displayed protein at all. Our results suggest the need

to be cautious when attributing activity of surface displayed proteins, even if they are

readily detected. There are other reports of surface displayed enzymes or antigens have

shown lack of correlation between surface detection and activity both in-vitro and in-vivo

[42–44]. We suggest future studies to more directly attribute activity to displayed proteins and

not due to presence of active cytoplasmic proteins in whole or lysed cells. We posit that

overexpression of surface directed proteins may yield a "back-up" resulting in translation of

active protein fusions directly in the cytoplasm. Surface display of LsLAI has been shown

before with *B. subtilis* endospores; however, the construct is assembled on the spore

cytoplasmically [19], without the need for any secretion. Our results contradict reports from

another study that claimed secretion of active LsLAI using *Lactococcus lactis* [1]. However,

closer analysis of the study revealed that measured activity was never attributed directly to

culture supernatant. Rather activity was detected either for whole cells or in the elution

fraction post-chromatographic purification, which could have allowed for refolding to

occur. Thus, we posit that cell-surface display is not a viable option to improve the stability

of this enzyme.

Cellular or abiotic encapsulation has shown to improve enzyme stability against thermal,

pH, solvent, and impurity interactions [24,45]. In the context of tagatose production, whole

cells expressing L-AI has also shown to improve conversion by preferentially partitioning

substrate and product across the *E. coli* membrane barrier through active transport [25];

however, the same was not observed in resting *L. lactis* cells [46]. Our results demonstrate

that encapsulation of LAI in *L. plantarum* behaves in a manner similar to that in Gram-

negative *E. coli* but not Gram-positive *L. lactis*. That is, the cell membrane preferentially

allows passage of galactose to the intracellular space relative to tagatose to aid the forward

reaction. A concurrent advantage of the encapsulation was that the enzyme stability was

markedly improved. Specifically, while the free enzyme rapidly inactivates at 50 °C, the

cell-encapsulated enzyme maintains activity for a prolonged period. Indeed, the resistance

to lysis at elevated temperatures is afforded by the thick cell wall of the Gram-positive *L.*

plantarum, as opposed to Gram-negative *E. coli*, which is known to readily lyse at 50 °C [47].

Thus, while many previous studies have demonstrated the benefits of using whole cells to

enhance enzyme stability, this work is the first to show the twofold benefit by combining

improved stability with substrate-product partitioning. While the selective membrane

circumvents the thermodynamic barrier that limits maximum conversion at moderate

temperatures, it imposes a barrier to mass transport – a penalty to reaction kinetics. We

addressed this issue next.

Cell permeabilization using chemical agents like detergents or solvents have been shown to

non-selectively increase substrate exchange while maintaining biological cellular

components [48,49], although the exact mechanism of this phenomenon is poorly understood.

We used detergents to improve reactant/product transport across the *L. plantarum* cell

membrane without altering its selectivity. In our best-case scenario, sub-micellar

concentration of SDS resulted in significant increase in whole cell reaction rates to free-

enzyme-like rates while still allowing the system to circumvent the thermodynamic barrier

to product conversion. Since the conversion was unchanged, we presume that the detergents

were not making pores in the cell membrane. Interestingly, permeabilization increased the

initial reaction rate in the forward direction 3.8-fold, however the rate of the reverse

reaction remained low, indicating that the membrane retained its selectivity (Fig S3).

Studies with cFDA further confirmed that cytoplasmic components were not "leaking" into

the extracellular space and that the cellular integrity was maintained. Thus, substrate and

product transport were still likely facilitated through transporters, as in wild-type, untreated

cells, which is also consistent with the observation that conversion reaches the same

maximum with or without detergent treatment. Since SDS and Triton-X treatment resulted

similar outcomes, we believe there are no specific charge-related effects of these

detergents, either. Adding SDS to a pure enzyme solution did not alter its kinetic properties,

indicating that the detergents do not directly interact with the enzyme in any appreciable

manner. However, our study revealed that treatment with SDS removed extracellular- and

surface-attached proteins, potentially altering the fluidity of the membrane. It is unclear

how removal of surface proteins may further facilitate galactose and tagatose transporter

activity (since transport across the membrane is the rate-limiting step in our system), but it

does seem to the mechanism though which we achieve enhanced kinetics.

Having addressed all three limitations (stability, thermodynamics, kinetics), we found that

L. plantarum whole-cell biocatalyst achieved a conversion from of ~ 50 % in 4 h and ~ 85

% 48 h in batch process at 50 °C starting with 300 mM galactose. Improved stability is

apparent by comparison with the free enzyme system, which quickly inactivates at 50 °C

and achieves only 16 % conversion in 48 h. Enhanced equilibrium conversion compared to

free enzyme system can be seen at both 50 °C and 37 °C, but especially in the latter, where

the free enzyme is more stable but is still able to achieve only 13 % conversion in 4 h and

47 % in 48 h. Finally, improved kinetics can be seen compared to unpermeabilized cells

where initial reaction rates at 0.5 mM/min compared to 1.6 mM/min in permeabilized cells.

This rate is higher than the maximum reaction rates sustainable by the free enzyme for

extendable period (initial rate at 37 °C for free enzyme = 0.3 mM/h) due to the ability to

operate at 50 °C. This is the among the highest reported productivity (averaging ~ 11.4

g/L/h over 48 h) for biocatalytic tagatose production, and the highest conversion reported

for a mesophilic enzyme (~85 %) [19,50]. Although further improvements to conversion may

be possible by including borate in the reaction buffer, demonstrated previously, it would

also increase purification costs [16,26,46]. The productivity and conversion reported here is the

highest obtained for a mesophilic LAI under any conditions Further, this system nearly

matches the conversion obtained in a recent study with a extremophilic LAI at 95 °C and

pH 9.5 that required use of Ni^{2+} and borate [51] but at 50 °C and without use of toxic metals

and salts. Additionally, the system avoids product caramelization, which is known to

happen at high temperatures (≥ 80 °C) [18]. Thus, the system described here demonstrates

how to overcome three major limitations of enzymatic catalysis (stability, thermodynamic,

kinetic) for tagatose production using LAI. We believe that a similar strategy may be

applicable to other biocatalytic systems as well – particularly lyases [52], translgycosidases [53],

and isomerases [17,25,54,55] – where free enzyme performance is poor due to stability,

thermodynamic, and/or kinetic limitations.

Materials and Methods:

Reagents and Enzymes: All enzymes for cloning were purchased from NEB (Beverly, MA). DNA primers were ordered from Eurofins Genomics LLC (Louisville, KY) or GENEWIZ Inc. (Cambridge, MA). Growth media and chemicals were purchased from Amresco (Solon, OH) or RPI Corp (Mount Prospect, IL). High-purity L-arabinose and D-galactose were purchased from Sigma-Aldrich (St. Louis, MO). Plasmid Mini Kit I, PCR Purification and Gel Extraction Kits were obtained from Omega Bio-tek (Norcross, GA). Mouse-anti-His$_6$ primary antibody and goat anti-mouse IgG H&L (Alexa Fluor® 488) were purchased from Thermo Fisher Scientific (Waltham, MA).

Bacterial Strains and Culture Conditions: *Lactobacillus plantarum* WCFS1 (kind gift from Prof. Michiel Kleerebezem, Wageningen University & Research) was cultivated in deMan−Rogosa−Sharpe (MRS) medium at 30 °C without agitation. *E. coli* strain NEB5α (New England Biolabs, Ipswich, MA, USA) was grown in Luria−Bertani (LB) medium at 37 °C with shaking at 225 rpm. When needed, media were supplemented with erythromycin at concentrations of 5 μg/mL and 200 μg/mL for *L. plantarum* and *E. coli*, respectively. Solid media were prepared by adding 1.5 % (w/v) agar to the respective media.

Cloning: DNA amplification was performed with Phusion High-Fidelity DNA Polymerase according to the manufacturer's instructions. The oligonucleotide primers were obtained from GeneWiz (Vienna, Austria). *E. coli* was used as host for plasmid propagation before transformation into *L. plantarum*. The amplified sequences were verified by a commercial

sequencing service (GeneWiz). A list of all strains and plasmids used in this study in Table

1.

Plasmid Construction: Each plasmid was transformed into chemically competent *E. coli*

NEB5α cells according to the manufacturer's instructions. *L. plantarum* was transformed

using a previously described electroporation method [56] with slight modifications. Briefly,

stationary phase cells were subcultured to an $OD_{600} = 0.1$ in MRS and allowed to grow until

an $OD_{600} = 0.85$. Cells were harvested via centrifugation at 4 °C and washed twice in ice-

cold 10 mM $MgCl_2$ and once with 0.5 M sucrose and 10 % glycerol. Cells were

resuspended in 1:50 (v/v) of the same solution and kept on ice prior to transformation. 100

ng of plasmid DNA was diluted into 5 μL of water and mixed with 50 μL electrocompetent

cells immediately before adding to a chilled 1 mM electroporation cuvette. The bacteria

were transformed using a BioRad Pulser at 1300 V with a fixed time constant of 5 ms. 1

mL of warm MRS was added and the cells transferred to a new centrifuge tube and allowed

to recover statically at 37 °C for 3 h before plating on appropriate media containing

selective antibiotic. Positive transformants were confirmed using colony PCR.

Enzyme Purification: *L. plantarum* cells carrying pJRB14Q-LSH were grown and induced

as described previously. Cells were washed in lysis/equilibration buffer (50 mM sodium

phosphate, 300 mM sodium chloride, 15 % glycerol, pH 8.0) and resuspended in lysis

buffer containing 10 mM lysozyme and incubated at 37 °C with shaking for one hour. Lysis

was achieved via sonication and soluble protein collected as described previously.

Purification was achieved via immobilized metal affinity chromatography (IMAC) using an

N-terminal His_6 tag. TALON metal affinity resin (Invitrogen) was used as directed by the

manufacturer. A single elution of protein was achieved in elution buffer containing 500

mM imidazole. Active fractions were pooled, and the buffer exchanged with PBS

containing 1 mM $MgCl_2$ (PBSM) using a Microsep 10 kDa Omega centrifugation spin

column (PALL Corp; Port Washington, NY). Purification was confirmed via SD-PAGE

analysis. Purified enzyme was stored in PBSM with 20 % glycerol at −80 °C prior to use.

Enzyme Analysis: *L. plantarum* wild-type (WT) cells or those carrying plasmids pJRB01Q-

LSH (A1), pJRB02Q-LSH (A2), pJRB03Q-LSH (A3), pJRB04Q-LSH (IC1), pJRB05Q-

LSH (A4), pJRB06Q-LSH (A5), pJRB08Q-LSH (A6) were cultured in MRS media with

antibiotic as needed at 37 °C overnight. Overnight cultures were diluted 1:50 (approximate

OD_{600} = 0.1) in MRS media with antibiotic as needed and grown at 37 °C for 2.5 h. Cells

were then induced with IP-673 (synthesized by Life Technologies; Carlsbad, CA) at a final

concentration of 25 ng/ml and $MgCl_2$ was added to a final concentration of 1 mM to support

enzyme folding. Cultures were incubated at 37 °C for an additional 3 h before harvesting the

cells by centrifugation at $3000 \times g$ for 5 min.

Cell pellets were washed in 1 mL of 20 mM phosphate buffer (PB) pH 6.8 and transferred

to a 1.7 mL centrifuge tube. Cells were washed twice with 1 mL of PB before being

suspended in 200 μL PB buffer. The enzyme activity assay was performed in a 96-well

plate as follows: 20 μL cell suspension was incubated with 0.8 mM $MnCl_2$, 0.8mM $MgCl_2$,

sugar (200 mM galactose, 200 mM glucose, or 20 mM arabinose) and diluted to 200 μL

with PB. The cell density within each well was measured for data normalization at the start

of the reaction. The plates were sealed and incubated at 37 °C for 2 h. Enzyme activity was

measured colorimetrically using the cysteine− carbazole−sulfuric-acid method (CCSAA) [57].

Absorbance at 560 nm was measured after 1 h of incubation at room temperature. Triplicate

technical replicates of each cell suspension were analyzed and averaged due to variability

in the assay itself.

To determine the conversion achieved with our balanced system, *L. plantarum* cells

carrying pJRB14Q-LSH (IC2) and wild-type were grown and induced as described

previously with slight modification. Cells were induced with 25 ng/mL IP-673 for 8 h

instead of 3 h to maximize the cell density of the culture. Cells were harvested via

centrifugation and washed twice in phosphate-buffered saline pH 7.4 with 1.0 mM $MgCl_2$

(PBSM). Cells to be permeabilized were diluted to an $OD_{600} = 3.0$ before incubation with

0.0 1% SDS in PBSM for 30 min at room temperature. The enzyme activity assay was

performed in 5 mL tubes as follows: cell suspensions adjusted to an $OD_{600} = 40$ were

incubated in phosphate-buffered saline (PBS) pH 7.4 with 1.0 mM $MgCl_2$, 300 mM

galactose and adjusted to 5 mL with H_2O.

Initial rates of reaction for both pure-enzyme, unmodified, and modified whole-cell

catalysts were determined. *L. plantarum* cells carrying pJRB14Q-LSH were grown and

induced for 8 h as described previously. Permeabilized cells were prepared as described

previously. The enzyme activity assay was performed in 200 μL PCR tubes as follows:

purified enzyme or cell suspensions were incubated in PBSM, variable substrate

concentration of galactose or tagatose, and adjusted to 100 μl with H_2O. Reactions were

initiated by adding $1/10^{th}$ volume cell suspensions adjusted to an $OD_{600} = 10$ or 2 μg of

purified enzyme to prewarmed reaction mixtures in a PCR block at either 37 or 50 °C.

Reactions proceeded for 6 min before termination to measure initial reaction rates. Cell

suspensions were immediately centrifuged for 1 min using a benchtop quick-spin centrifuge

and the supernatant was transferred to a 96-well 0.2 μm centrifugal filter plate with

collection plate. The remaining solution was filtered via centrifugation and stored at -80C

until analysis. Purified enzyme reactions were terminated by adding $1/10^{th}$ volume 2M

perchloric acid. Prior to HPLC analysis the reactions were neutralized using a sodium

hydroxide PBS buffered solution.

Enzyme stability was measured by incubating either purified enzyme or unmodified whole

cell catalysts in PBSM at 37 or 50 °C statically for the duration of the experiment. At

determined time intervals and aliquot of catalyst was added to PBSM reaction mixture

containing 300mM galactose at the same temperature as the catalyst. Reactions proceeded

for 20 min before termination as described previously. Tagatose production was confirmed

via HPLC analysis.

Equilibrium conversion was measured by in incubating either purified enzyme or

unmodified whole cell catalysts in PBSM at 37 °C. Purified enzyme was incubated with

either 10 mM galactose, 5 mM of galactose and tagatose, or 10 mM tagatose. The reaction

was sampled every 24 h and additional enzyme was added after sampling due to incomplete

reaction due to enzyme degradation. Whole cell catalyst was incubated with either 30 mM

galactose, 15 mM of galactose and tagatose, or 30 mM tagatose.

HPLC Analysis: Agilent Infinity HPLC system (Agilent; Santa Clara, CA) equipped with

a Hi-Plex Ca+ 300 x 7.7mm column with a guard column and detected using Agilent

Infinity 1260 ELSD detector. Aqueous mobile phase was run at 85 °C with a flow rate of

0.6 mL/min. The ELSD detector's evaporation temperature was set at 90 °C, the nebulizer

temperature set to 50 °C, and nitrogen flow rate at 1.6 SLM (standard liter per minute). For

calculation of the reaction product(s) L-arabinose, D-galactose, L-ribulose, and D-tagatose

standards were included in the run.

Flow Cytometry and Immunostaining: *L. plantarum* WT cells or those carrying plasmids

pJRB01QLs-pJRB08QLs were grown and induced as described previously. Cells were

collected by centrifugation at 5000 ×g for 5 min and washed twice in PBS before being

suspended to an $OD_{600} = 1.0$. An aliquot of 200 µL cells was pelleted and was resuspended

in 200 µL PBSA (PBS with 2 % bovine serum albumin). The suspension of cells rested on

ice for 30 min before the cells were pelleted and solution removed. The cells were then

resuspended in 50 uL of PBSA containing 1 µg mouse-anti- His_6 primary antibody. The

cells were incubated for 1h at room temperature with gentle rocking. Cells were collected

and washed thrice with PBSA before being suspended in 50 µL of PBSA containing 0.4 µg

goat anti-mouse IgG H&L (Alexa Fluor® 488). The cells were incubated for one fifteen

minutes on ice and blocked from light. Finally, cells were washed thrice in PBSA and

resuspended in 500ul of PBSA. A 20ul aliquot was diluted 10X in PBSA and transferred to

a flat-bottom 96-well plate for flow cytometric analysis. A positive control sample was

used to calibrate the fluorescence intensity. A negative control sample was used to gate the

level of autofluorescence of the cells and any non-specific secondary antibody binding (Fig

S6). A total of 1×10^5 cell counts per sample plotted on a 'logical' or bi-exponential x-axis.

Data analyzed using FCS Express 6 (De Novo Software, Glendale,CA). Surface detection

level was calculated as the fraction of cells with a measured fluorescence greater than that

of the gated negative control.

Surface Treatment: *L. plantarum* WT cells or those carrying either pJRB04Q-LSH or

pJRB08Q-LSH were grown and induced as previously described. Cells were washed in

PBS twice before further processing. A concentrated cell solution was incubated in PBS

containing either 0 % or 0.5 % SDS for 1 h before being pelleted by centrifugation. The

SDS solution was removed and the cell pellet washed twice with PBS. Surface detection

was measured via immunofluorescence flow cytometry as described previously. Whole-cell

activity was measured via the CCSAA after a 1-hour incubation in 200 mM arabinose and 1

mM $MgCl_2$ in 100 mM sodium acetate buffer pH 5.0 at 37 °C.

Western Blot Analysis: *L. plantarum* WT cells or those carrying either pJRB04Q-LSH or

pJRB08Q-LSH were grown and induced as previously described. A concentrated cell

solution was incubated with 0.05 % SDS in PBS for 30 min at 37 °C before being pelleted.

The SDS buffer solution was removed and stored separately. The cell pellet was then

washed twice with PBS. A fresh cell solution of the same concentration was prepared in

PBS as a control. Both aliquots of cells were then incubated with 10 mg/mL lysozyme for 1

h at 37 °C before being lysed via sonication. Sonication was performed using a Branson

150 system equipped with a microtip probe sonicating samples on ice at 55 % amplitude in

30 s ON 2 min OFF cycles for a total of 5 min of ON time. The insoluble cell fraction

containing membrane components and the insoluble protein fraction was pelleted via

centrifugation at 20,000 ×g for 5 min. The soluble fraction was removed, filtered through

0.4 μm sterile filter, and stored separately. An aliquot of each of the cell fractions for the

three cell types was added to denaturing loading dye and incubated at 98 °C for 15 min. 10

μL of each sample were added to a 4 % - 12 % Bis-Tris PAGE gel in MOPS buffer and run

at 75 V for 15 min followed by 110 V for 1.5 h. After separation the gel was removed from

the cassette and washed twice in deionized water. The proteins were next transferred to a

PVDF membrane for Western blot analysis. Transfer was visually confirmed through the

transfer of the stained protein ladder. The membrane was blocked overnight at room

temperature with 5 % skim milk in Tris buffered saline containing 1 % Tween-20 (TBST).

The membrane was washed twice with TBST before being incubated with 0.2 µg/mL

primary antibody in blocking buffer for 2 h at room temperature. The membrane was

washed twice with TBST before being incubated with 2 ng/mL of the horseradish

peroxidase (HRP) conjugated secondary antibody at room temperature for 1 h. The

membrane was then washed twice with TBST before incubation in the substrate containing

solution (Thermo SuperSignal West) for 15 min at room temperature. The

chemiluminescent signal was captured electronically via a camera set to a 30 min exposure

time.

Cell Permeabilization: *L. plantarum* WT cells or those carrying pJRB14Q-LSH were

grown and induced and washed in PBS as previously described. Cells were resuspended to

an $OD_{600} = 3.0$ in 200 µL of PBS. Samples were either incubated with 1% Triton X-100 [27],

0.1 % SDS, or no treatment at room temperature for 1 h. The solution was removed, and the

cell pellet washed twice with PBS before enzymatic activity of the whole-cells were

measured as described previously using 200 mM of glucose, galactose, or arabinose as

substrates. Activity was quantified via the CCSAA and normalized to cell concentration as

described previously.

The optimal concentration of SDS to maximize whole-cell activity was determined. Cells

were prepared as previously described. Cell lysate from the equivalent cell density was

prepared via lysozyme treatment via sonication as described previously. Permeabilized

samples were either incubated with varying concentrations of SDS, or no treatment at room

temperature for 30 min. The solution was removed, and the cell pellet washed twice with

PBS before enzymatic activity of the whole-cells were measured as described previously

using 200 mM of glucose, or galactose, as substrates. Activity was quantified via the

CCSAA and normalized to cell concentration as described previously.

Cell permeabilization was also tested using carboxyfluorescein diacetate (cFDA) (Sigma-

Aldrich) as substrate. Cell suspensions were diluted to $OD_{600} = 0.5$ in PBS in a black

polystyrene 96-well plate. The reaction was started by adding 10 mM cFDA to the cell

suspensions. Transport of cFDA was estimated by measuring the fluorescent signal released

when cFDA becomes activated upon intracellular cleavage via nonspecific esterases. The

total fluorescence was measured on a plate-reader using an excitation and emission

wavelength of 485 nm and 520 nm, respectively, every 15 s for 30 min. Apparent transport

rates were calculated from the slope of the linear fit to the data.

525
526
527

Figure 1: Free-enzyme LsLAI suffers from poor thermal stability, low thermodynamic equilibrium, and kinetic limitations. a) Initial turnover rates of purified LsLAI compared at medium (green) and high (orange) substrate loading at low (37 °C) and high (50 °C) temperatures. Comparison of the non-native substrate, galactose (GAL), to that of native substrate, arabinose (ARA). b) Loss of activity of purified LsLAI over time incubated at 37 °C (green) or 50 °C (orange). Half-life calculated from first-order decay equation. c, Equilibrium conversion of purified LsLAI starting from 10 mM total substrate with galactose to tagatose ratio at 1:0 (purple), 1:1 (orange), or 0:1 (green). Additional enzyme was added every 24 h to account for thermal inactivation. The data are means from three biological replicates. *** = p < 0.001 (Students t-test).

Figure 2: LsLAI surface display and activity. a) Flow cytometry analysis of *L. plantarum* wild-type cells (orange), cells expressing intracellular His$_6$-tagged LsLAI (orange), or cells expressing LsLAI fused to native surface anchor proteins A1-A6 for display (blue) with 10^5 counts per sample. Data is normalized to counts. b) Comparing the amount of tagatose produced in 2 h (hashed) with the percentage of the population with surface detection intensity above that of wild-type "WT" for intracellularly expressed "IC1" or anchor protein A1-A6 (colored). c) Scatterplot of each replicate measurement correlating surface detection vs. tagatose produced. There is no significant correlation between the display level and activity based on Pearson Product Moment Correlation test (Correlation coefficient = −0.395, p = 0.0564). The data are means from three biological replicates.

Figure 3: Encapsulation of LsLAI improves equilibrium conversion and provides thermal stability, albeit at a kinetic penalty a) Loss of activity of encapsulated LsLAI over time incubated in PBSM pH 7.4 at 37°C (orange) or 50°C (blue). Half-life calculated assuming first-order decay. b) Equilibrium conversion of encapsulated LsLAI starting from 30 mM total substrate: 1:0 (purple), 1:1 (orange), or 0:1 (green) galactose:tagatose respectively. c) Tagatose production of *L. plantarum* wild-type "WT" cells (green), IC2 cells expressing LsLAI intracellularly "I" (purple), or lysate of IC2 cells expressing LsLAI cytoplasmically "LYS" (orange) at different induction conditions. The data are means from three biological replicates.

Initial rate (mM/min)	0.3	0.5	1.6	0.5	0.7
6 h (%)	22	46	59	13	8
48 h (%)	47	82	85	39	16

Figure 4: Modified encapsulation of LsLAI maximizes tagatose production.
Batch tagatose production over time starting from 300 mM galactose. Encapsulated LsLAI at 37 °C is "PBS-37C" (green), encapsulated LsLAI at 50 °C is "PBS-50C" (pink), permeabilized encapsulated LsLAI at 50 °C is "SDS-50C" (blue), purified free-enzyme LsLAI at 37 °C is "PURE-37C" (orange), and purified free-enzyme LsLAI at 50 °C is "PURE-50C" (purple). Inset shows initial reaction rates. Salient feature are summarized in table. The data are means from three biological replicates.

538 **Table 1: Plasmids & Strains**

Plasmid	Description	Reference
pLp_3050Ag85B-E6cwa2	pSIP401 based plasmid containing an oncofetal antigen and Lp_2578 anchor for *L. plantarum* surface display.	58
pLp_1261Ag85B-E6	pSIP401 based plasmid containing invasion and Lp_1261 anchor for *L. plantarum* surface display.	59
pLp_1452Inv	pSIP401 based plasmid containing invasion and Lp_1452 anchor for *L. plantarum* surface display.	59
pSIP411	Lactobacillus inducible plasmid system with broad host SH71 origin.	60
pJRB01Q-LSH "A1"	pSIP401 based plasmid containing Ls-araA, sppQ, Lp_2578, SP_3050, His$_6$ tag	This work
pJRB02Q-LSH "A2"	pSIP401 based plasmid containing Ls-araA, sppQ, Lp_2162, SP_3050, His$_6$ tag	This work
pJRB03Q-LSH "A3"	pSIP401 based plasmid containing Ls-araA, sppQ, Lp_2940, SP_3050, His$_6$ tag	This work
pJRB04Q-LSH "IC1"	pSIP401 based plasmid containing Ls-araA, sppQ, His$_6$ tag	This work
pJRB05Q-LSH "A4"	pSIP401 based plasmid containing Ls-araA, sppQ, Lp_1261, SP_3050, His$_6$ tag	This work
pJRB06Q-LSH "A5"	pSIP401 based plasmid containing Ls-araA, sppQ, Lp_1452, SP_3050, His$_6$ tag	This work
pJRB08Q-LSH "A6"	pSIP401 based plasmid containing Ls-araA, sppQ, Lp_3014, SP_3050, His$_6$ tag	This work
pJRB09Q-LSH "Sec"	pSIP401 based plasmid containing Ls-araA, SP_3050, His$_6$ tag	This work
pJRB14Q-LSH "IC2"	pSIP411 based plasmid containing Ls-araA, sppQ, His$_6$ tag	This work

References:

546   1.   Rhimi, M. *et al.* The secreted l-arabinose isomerase displays anti-hyperglycemic

547   effects in mice. *Microb. Cell Fact.* **14,** 204 (2015).

548   2.   Luecke, K. J. & Bell, L. N. Thermal stability of tagatose in solution. *J. Food Sci.* **75,**

549   C346-51 (2010).

550   3.   Bertelsen, H., Andersen, H. & Tvede, M. Fermentation of D-Tagatose by Human

551   Intestinal Bacteria and Dairy Lactic Acid Bacteria. *Microb. Ecol. Health Dis.* **13,**

552   87–95 (2001).

553   4.   Xu, Z., Li, S., Feng, X., Liang, J. & Xu, H. L-Arabinose isomerase and its use for

554   biotechnological production of rare sugars. *Appl. Microbiol. Biotechnol.* **98,** 8869–

555   8878 (2014).

556   5.   Cheetham, P. S. J. & Wootton, a. N. Bioconversion of d-galactose into d-tagatose.

557   *Enzyme Microb. Technol.* **15,** 105–108 (1993).

558   6.   Xu, W., Zhang, W., Zhang, T., Jiang, B. & Mu, W. l-arabinose isomerases:

559   Characteristics, modification, and application. *Trends Food Sci. Technol.* **78,** 25–33

560   (2018).

561   7.   Xu, Z. *et al.* A novel l-arabinose isomerase from Lactobacillus fermentum

562   CGMCC2921 for d-tagatose production: Gene cloning, purification and

563   characterization. *J. Mol. Catal. B Enzym.* **70,** 1–7 (2011).

8.   Waltman, M. J., Yang, Z. K., Langan, P., Graham, D. E. & Kovalevsky, A.

Engineering acidic Streptomyces rubiginosus D-xylose isomerase by rational

enzyme design. *Protein Eng. Des. Sel.* **27,** 59–64 (2014).

9.   Zheng, Z., Mei, W., Xia, M., He, Q. & Ouyang, J. Rational Design of Bacillus

coagulans NL01 L-Arabinose Isomerase and Use of Its F279I Variant in D-Tagatose

Production. (2017). doi:10.1021/acs.jafc.7b01709

10.   Fan, C. *et al.* Engineering of Alicyclobacillus hesperidum l-Arabinose Isomerase for

Improved Catalytic Activity and Reduced pH Optimum Using Random and Site-

Directed Mutagenesis. *Appl. Biochem. Biotechnol.* (2015). doi:10.1007/s12010-015-

1828-3

11.   Oh, H. J., Kim, H. J. & Oh, D. K. Increase in D-tagatose production rate by site-

directed mutagenesis of L-arabinose isomerase from Geobacillus

thermodenitrificans. *Biotechnol. Lett.* **28,** 145–149 (2006).

12.   Hong, Y.-H., Lee, D.-W., Pyun, Y.-R. & Lee, S. H. Creation of Metal-Independent

Hyperthermophilic l -Arabinose Isomerase by Homologous Recombination. *J. Agric.*

Food Chem. **59,** 12939–12947 (2011).

13.   Prabhu, P., Jeya, M. & Lee, J. K. Probing the molecular determinant for the catalytic

efficiency of L-arabinose isomerase from bacillus licheniformis. *Appl. Environ.*

Microbiol. **76,** 1653–1660 (2010).

14.   Choi, J. M. *et al.* Structure of the thermophilic l-Arabinose isomerase from

Geobacillus kaustophilus reveals metal-mediated intersubunit interactions for

activity and thermostability. *Arch. Biochem. Biophys.* **596,** 51–62 (2016).

15.   Kim, H. J. & Oh, D. K. Purification and characterization of an L-arabinose

isomerase from an isolated strain of Geobacillus thermodenitrificans producing D-

tagatose. *J. Biotechnol.* **120,** 162–173 (2005).

16.   Li, Y., Zhu, Y., Liu, A. & Sun, Y. Identification and characterization of a novel L-

arabinose isomerase from Anoxybacillus flavithermus useful in D-tagatose

production. *Extremophiles* **15,** 441–450 (2011).

17.   Hong, Y. H., Lee, D. W., Pyun, Y. R. & Lee, S. H. Creation of metal-independent

hyperthermophilic l -arabinose isomerase by homologous recombination. *J. Agric.*

Food Chem. **59,** 12939–12947 (2011).

18.   Kwon, S. Y. & Baek, H. H. Effects of Temperature, pH, Organic Acids, and Sulfites

on Tagatose Browning in Solutions during Processing and Storage. *Food Sci.*

Biotechnol **23,** 677–684 (2014).

19.   Liu, Y. *et al.* Efficient Production of d-Tagatose Using a Food-Grade Surface

Display System. *J. Agric. Food Chem.* **62,** 6756–62 (2014).

20.   Zhang, Y.-W., Jeya, M. & Lee, J.-K. Enhanced activity and stability of l-arabinose

isomerase by immobilization on aminopropyl glass. *Appl. Microbiol. Biotechnol.* **89,**

1435–1442 (2011).

21.   Bortone, N. & Fidaleo, M. Immobilization of the recombinant (His)6-tagged l-

arabinose isomerase from Thermotoga maritima on epoxy and cupper-chelate epoxy

supports. *Food Bioprod. Process.* **95,** 155–162 (2015).

22.   Kim, P., Yoon, S.-H., Roh, H.-J. & Choi, J.-H. High Production of D-Tagatose, a

Potential Sugar Substitute, Using Immobilized L-Arabinose Isomerase. *Biotechnol. Prog.* **17,** 208–210 (2001).

23. Hong, Y. H. *et al.* Production of D-tagatose at high temperatures using immobilized Escherichia coli cells expressing L-arabinose isomerase from Thermotoga neapolitana. *Biotechnol. Lett.* **29,** 569–574 (2007).

24. Lin, B. & Tao, Y. Whole-cell biocatalysts by design. *Microb. Cell Fact.* **16,** 106 (2017).

25. Kim, J. H. *et al.* Differential selectivity of the Escherichia coli cell membrane shifts the equilibrium for the enzyme-catalyzed isomerization of galactose to tagatose. *Appl. Environ. Microbiol.* **74,** 2307–2313 (2008).

26. Xu, Z. *et al.* Production of D-tagatose, a functional sweetener, utilizing alginate immobilized Lactobacillus fermentum CGMCC2921 cells. *Appl. Biochem. Biotechnol.* **166,** 961–73 (2012).

27. Jayamuthunagai, J., Srisowmeya, G., Chakravarthy, M. & Gautam, P. d -Tagatose production by permeabilized and immobilized Lactobacillus plantarum using whey permeate. *Bioresour. Technol.* **235,** 250–255 (2017).

28. Reider Apel, A., Ouellet, M., Szmidt-Middleton, H., Keasling, J. D. & Mukhopadhyay, A. Evolved hexose transporter enhances xylose uptake and glucose/xylose co-utilization in Saccharomyces cerevisiae. *Sci. Rep.* **6,** 19512 (2016).

29. Miskovic, L. *et al.* A design–build–test cycle using modeling and experiments

reveals interdependencies between upper glycolysis and xylose uptake in

recombinant S. cerevisiae and improves predictive capabilities of large-scale kinetic

models. *Biotechnol. Biofuels* **10,** 166 (2017).

30.   Bober, J. R., Beisel, C. L. & Nair, N. U. *Synthetic Biology Approaches to Engineer*

Probiotics and Members of the Human Microbiota for Biomedical Applications.

Annual Review of Biomedical Engineering **20,** (2018).

31.   Mays, Z. J. & Nair, N. U. Synthetic biology in probiotic lactic acid bacteria: At the

frontier of living therapeutics. *Curr. Opin. Biotechnol.* **53,** 224–231 (2018).

32.   Rhimi, M. *et al.* The acid tolerant l-arabinose isomerase from the food grade

Lactobacillus sakei 23K is an attractive d-tagatose producer. *Bioresour. Technol.*

101, 9171–9177 (2010).

33.   Mukerjee, P. & Mysels, K. J. Critical Micelle Concentrations of Aqueous Surfactant

Systems. *J. Pharm. Sci.* **61,** 319 (1972).

34.   Tao, F., Zhang, Y., Ma, C. & Xu, P. One-pot bio-synthesis: N-acetyl-d-neuraminic

acid production by a powerful engineered whole-cell catalyst. *Sci. Rep.* **1,** 142

(2011).

35.   Rhimi, M. *et al.* Rational design of Bacillus stearothermophilus US100 l-arabinose

isomerase: Potential applications for d-tagatose production. *Biochimie* **91,** 650–653

(2009).

36.   Hong, Y., Lee, D., Pyun, Y. & Lee, S. H. Creation of Metal-Independent

Hyperthermophilic. *Agric. Food Chem.* **59,** 12939–47 (2011).

37.   Fan, C. *et al.* Engineering of Alicyclobacillus hesperidum l-Arabinose Isomerase for

Improved Catalytic Activity and Reduced pH Optimum Using Random and Site-

Directed Mutagenesis. *Appl. Biochem. Biotechnol.* **177,** 1480–1492 (2015).

38.   Kim, B. J., Hong, S. H., Shin, K. C., Jo, Y. S. & Oh, D. K. Characterization of a

F280N variant of l-arabinose isomerase from Geobacillus thermodenitrificans

identified as a d-galactose isomerase. *Appl. Microbiol. Biotechnol.* 9271–9281

(2014). doi:10.1007/s00253-014-5827-z

39.   Mathiesen, G. *et al.* Genome-wide analysis of signal peptide functionality in

Lactobacillus plantarum WCFS1. *BMC Genomics* **10,** 425 (2009).

40.   Schneewind, O. & Missiakas, D. Sec-secretion and sortase-mediated anchoring of

proteins in Gram-positive bacteria. *Biochim. Biophys. Acta* **1843,** 1687–1697 (2014).

41.   Mori, H. & Ito, K. The Sec protein-translocation pathway. *Trends Microbiol.* **9,** 494–

500 (2001).

42.   Nguyen, H.-M. *et al.* Display of a β-mannanase and a chitosanase on the cell surface

of Lactobacillus plantarum towards the development of whole-cell biocatalysts.

Microb. Cell Fact. **15,** 169 (2016).

43.   Kuczkowska, K. *et al.* Immunogenic properties of *Lactobacillus plantarum*

producing surface-displayed *Mycobacterium tuberculosis* antigens. *Appl. Environ.*

Microbiol. AEM.02782-16 (2016). doi:10.1128/AEM.02782-16

44.   Kuczkowska, K., Mathiesen, G., Eijsink, V. G. H. & Øynebråten, I. Lactobacillus

plantarum displaying CCL3 chemokine in fusion with HIV-1 Gag derived antigen

causes increased recruitment of T cells. *Microb. Cell Fact.* **14,** 169 (2015).

45.   Mohamad, N. R., Marzuki, N. H. C., Buang, N. A., Huyop, F. & Wahab, R. A. An

overview of technologies for immobilization of enzymes and surface analysis

techniques for immobilized enzymes. *Biotechnol. Biotechnol. Equip.* **29,** 205–220

(2015).

46.   Salonen, N., Salonen, K., Leisola, M. & Nyyssölä, A. D-Tagatose production in the

presence of borate by resting Lactococcus lactis cells harboring Bifidobacterium

longum L-arabinose isomerase. *Bioprocess Biosyst. Eng.* **36,** 489–497 (2013).

47.   Russell, A. D. & Harries, D. Damage to Escherichia coli on exposure to moist heat.

Appl. Microbiol. **16,** 1394–9 (1968).

48.   de Carvalho, C. C. C. R. Whole cell biocatalysts: essential workers from Nature to

the industry. *Microb. Biotechnol.* **10,** 250–263 (2017).

49.   Mattei, B., Lira, R. B., Perez, K. R. & Riske, K. A. Membrane permeabilization

induced by Triton X-100: The role of membrane phase state and edge tension. *Chem.*

Phys. Lipids **202,** 28–37 (2017).

50.   Nguyen, T.-K., Hong, M.-G., Chang, P.-S., Lee, B.-H. & Yoo, S.-H. Biochemical

properties of L-arabinose isomerase from Clostridium hylemonae to produce D-

tagatose as a functional sweetener. *PLoS One* **13,** e0196099 (2018).

51.   Chen, H. *et al.* Surface display of the thermophilic lipase Tm1350 on the spore of

Bacillus subtilis by the CotB anchor protein. *Extremophiles* **19,** 799–808 (2015).

52.   Parmeggiani, F., Weise, N. J., Ahmed, S. T. & Turner, N. J. Synthetic and

691   Therapeutic Applications of Ammonia-lyases and Aminomutases. (2017).

692   doi:10.1021/acs.chemrev.6b00824

693   53.   Nair, N. U., Denard, C. A. & Zhao, H. *Engineering of Enzymes for Selective*

694   *Catalysis*. *Current Organic Chemistry* **14,** (2010).

695   54.   Guo, Q. *et al.* Enhanced d -tagatose production by spore surface-displayed l -

696   arabinose isomerase from isolated Lactobacillus brevis PC16 and biotransformation.

697   *Bioresour. Technol.* **247,** 940–946 (2018).

698   55.   Cheng, L., Mu, W. & Jiang, B. Thermostable L-arabinose isomerase from Bacillus

699   stearothermophilus IAM 11001 for D-tagatose production: gene cloning, purification

700   and characterisation. *J. Sci. Food Agric.* **90,** 1327–33 (2010).

701   56.   Alegre, M. T., Rodríguez, M. C. & Mesas, J. M. Transformation of Lactobacillus

702   plantarum by electroporation with in vitro modified plasmid DNA. *FEMS Microbiol.*

703   *Lett.* **241,** 73–7 (2004).

704   57.   Dische, Z. & Borenfreund, E. A new spectrophotometric method for the detection

705   and determination of keto sugars and trioses. *Chem, J Biol* **192,** 583–587 (1951).

706   58.   Fredriksen, L., Mathiesen, G., Sioud, M. & Eijsink, V. G. H. Cell wall anchoring of

707   the 37-kilodalton oncofetal antigen by Lactobacillus plantarum for mucosal cancer

708   vaccine delivery. *Appl. Environ. Microbiol.* **76,** 7359–62 (2010).

709   59.   Fredriksen, L. *et al.* Surface display of N-terminally anchored invasin by

710   Lactobacillus plantarum activates NF-κB in monocytes. *Appl. Environ. Microbiol.*

711   **78,** 5864–71 (2012).

712   60.   Karlskås, I. L. *et al.* Heterologous protein secretion in lactobacilli with modified

713   pSIP vectors. *PLoS One* **9,** e91125 (2014).

714