Surpassing thermodynamic, kinetic, and stability barriers to isomerization catalysis

for tagatose biosynthesis.

Josef R. Bober[1] & Nikhil U. Nair[1]*

[1]Department of Chemical and Biological Engineering, Tuts University, Medford, MA
02155

* corresponding author

Abstract:

There are many enzymes that are relevant for making rare and valuable chemicals that

while active, are severely limited by thermodynamic, kinetic, or stability issues (e.g.

isomerases, lyases, transglycosidases, etc.). In this work, we study an enzymatic reaction

system – *Lactobacillus sakei* L-arabinose isomerase (LsLAI) for D-galactose to D-tagatose

isomerization – that is limited by all three reaction parameters. The enzyme has a low

catalytic efficiency for the non-natural substrate galactose, has low thermal stability at

temperatures $> 40\ °C$, and equilibrium conversion $< 50\%$. After exploring several strategies

to overcome these limitations, we finally show that encapsulating the enzyme in a gram-

positive bacterium (*Lactobacillus plantarum*) that is chemically permeabilized can enable

reactions at high rates, high conversion, and high temperatures. The modified whole cell

system stabilizes the enzyme, differentially partitions substrate and product across the

membrane to shift the equilibrium toward product formation and enables rapid transport of

substrate and product for fast kinetics. In a batch process, this system enables

approximately 50 % conversion in 4 h starting with 300 mM galactose (an average

productivity of 37 mM/h), and 85 % conversion in 48 h, which are the highest reported for

food-safe mesophilic tagatose synthesis. We suggest that such an approach may be

invaluable for other enzymatic processes that are similarly kinetically-, thermodynamically-

, and/or stability-limited.

D-Tagatose is a rare ketohexose sugar with sweetness similar to that of sucrose. However,

its glycemic index and caloric value is much lower because of low bioavailability, making

it an attractive generally regarded as safe (GRAS) sugar substitute. Recent studies have also

demonstrated that it is anti-hyperglycemic [1] and prebiotic, which promotes gut health [2,3].

Thus, there exists a high demand in food industry for the economical production of rare

sugars, like tagatose, exemplified by the 2016 global artificial sweetener market estimated

to be USD 3.2 billion. This market is expected to expand given the global diabetes crisis

and increasing relevance of prebiotics.

The enzyme L-arabinose isomerase (LAI) that responsible for the reversible isomerization

of the pentose L-arabinose to L-ribulose can also isomerize the hexose D-galactose to D-

tagatose [4,5]. LAI has thus been the enzyme of choice to produce tagatose, although, to date,

few commercial bioprocesses exist. A variety of LAIs from different microorganisms have

been isolated and have reported optimal activity at a range of temperatures and pH [4,6,7].

Some of the limitations of tagatose biosynthesis using LAI that may be hindering

commercial viability are, 1) unfavorable enzymatic kinetics since galactose is not the native

substrate of LAI, 2) low enzyme stability, particularly in the absence of divalent metal ions,

and 3) low equilibrium constant for galactose to tagatose isomerization.

Few previous reports have been successful at engineering enzymatic properties of LAI for

industrial application; often addressing only one of the bottlenecks to productivity. To

address the kinetic issue, several groups have used enzyme engineering methods to enhance

catalytic efficiency of LAI toward galactose and have shown moderate increases in

productivity [8–13]. To counter low-stability issues, many groups have tested the utility of

thermophilic enzymes [12,14–16]. However, most thermophilic enzymes rely on divalent metal

ions (Mn^{2+}, Co^{2+}, Fe^{2+}) for stability[17], and high reaction temperatures ($\geq 80\ °C$) result in

significant caramelization [18], which are all undesirable and must be removed from product,

adding to processing costs. Surface-display [19] or encapsulation in particles or whole-cells

[20–23] can stabilize enzymes [24]. Finally, the thermodynamic limitations of isomerization of

galactose to tagatose are severe and, arguably, the most recalcitrant issue since $\Delta G°_{rxn} \approx$

$+1.19$ kcal/mol [23], which indicates theoretical maximum equilibrium conversion ~ 14% at

room temperature. Several approaches have been used to try and overcome this limitation,

including the use of an oxidoreductive pathway rather than isomerization [25]. However, this

method resulted in low productivity, byproduct formation, and the need for a sacrificial

substrate to regenerate cofactors and drive the reaction uphill. Thermophilic LAI enzymes

can achieve higher conversions than their mesophilic counterparts since the equilibrium

shifts toward tagatose at higher temperatures [6]. Whole-cell biocatalysts with GRAS

organisms (e.g. lactic acid bacteria (LAB) and *E. coli*) that disproportionately partition

substrate and product across their membrane has also been shown to partially circumvent

this thermodynamic limitation while simultaneously enhancing enzyme stability; albeit at a

kinetic penalty imposed by substrate transport limitations [26–28]. Recently, cell

permeabilization [28] and sugar transport overexpression [29,30] were demonstrated as methods

to overcome the kinetic penalty imposed by cellular encapsulation.

There have currently been no studies that look to systematically analyze all three

limitations – kinetic, thermodynamic, and enzyme stability – of the enzymatic

isomerization of galactose to tagatose. This work clearly demonstrates the presence of these

three limitations and provides a novel approach to balance their advantages and limitations.

We used the food-safe engineered probiotic bacterium *Lactobacillus plantarum* as the

expression host due to its increasing relevance to biochemical and biomedical research [31,32].

This approach enabled ~ 50 % conversion of galactose to tagatose in 4 h (productivity of ~

38 mmol tagatose L^{-1} h^{-1}) ultimately reaching ~ 85% conversion after 48 h at high galactose

loading (300 mM) in batch culture. This is among the highest conversions and

productivities reported to date for tagatose production using a mesophilic enzyme. Such an

approach is expected to be applicable to other biocatalytic systems where similar trade-offs

between kinetics, thermodynamics, and/or stability pose hurdles to process development.

Results:

Production of tagatose using *L. sakei* LAI (LsLAI) is limited by poor conversion and
low thermal stability.

At high substrate loading (400 mM galactose) and 37 °C, the reported optimal temperature

for this enzyme [33], LsLAI purified from *L. plantarum* exhibited an initial forward reaction

(turnover) rate ($k_{initial}$) of 9.3 ± 0.3 s^{-1}, which is lower than the maximum reaction rate

possible by this enzyme of 17.0 ± 1.3 s^{-1} min^{-1} with its preferred substrate, arabinose, at 400

mM. Increasing the reaction temperature to 50 °C increased the initial reaction rate to 11.2

s^{-1} (Fig. 1a) but was accompanied by rapid enzyme inactivation, which is consistent with

previous reports of thermal instability of this enzyme [33]. The enzyme exhibited first-order

degradation with a half-life ($t_{1/2}$) of approximately 18.5 h at 37 °C (Fig. 1b) and 1.5 h at 50

°C. Indeed, the enzyme retains only ~ 6 % activity after 6 h at this elevated temperature.

Additionally, the highly reversible isomerization reaction suffers from thermodynamic

limitations with reported free-enzyme equilibrium conversions < 50 % (ΔG°_{rxn} = +1.19 kcal

mol^{-1}) [23]. Our results indicate similar low conversion of 35 – 40 % after 5 days with pure

enzyme at 37 °C with daily supplementation of fresh enzyme to account for inactivation

(Fig. 1c).

***L. plantarum* cell surface display is unsuitable to stabilize LsLAI.**

To achieve surface localization, we fused LsLAI with six *L. plantarum* surface proteins,

which have all been previously described as suitable display carriers, at either the C- or N-

terminus (Table S4). The constructs also contained the strong Lp_3050 secretion signal, a

10 amino acid linker containing a thrombin protease cleavage site, and either a C- or N-

terminal His$_6$ tag for immunofluorescence assays. We characterized surface localization

using flow cytometry and measured the whole-cell isomerization activity. All three of the

tested C-terminal anchor strains (A1, A2, and A3) exhibited > 90 % surface detection (Fig.

2a) (see Table S3 for strain descriptions). However, the measured tagatose production after

a 2 h incubation with 200 mM galactose was only 225 ± 72, 55 ± 39, 91 ± 25 µM tagatose

(normalized to individual reaction cell densities of ~ OD$_{600}$ = 0.5), respectively,

significantly less than that of intracellularly expressed (IC1) LsLAI (538 ± 34 µM) (Fig.

2b). LsLAI was detected on N-terminal anchor strains A5 and A6 in 28 % and 96 % of the

population and produced 286 ± 31 and 422 ± 34 µM tagatose respectively. Interestingly,

strain A4 exhibited low surface detection, comparable to control (WT & IC1) cells while

producing the highest titers of tagatose (533 ± 58 µM) among all the surface display

constructs. These data indicate there is not a significant correlation (p > 0.05) between

whole-cell activity with the presence of displayed LsLAI on the *L. plantarum* cell surface

(Fig. 2c).

Next, we sought to explain the observation regarding absence of positive correlation

between display level and activity using strain A6. Sodium dodecyl sulfate (SDS) is a

strong anionic surfactant that is commonly used to lyse cells through membrane disruption

and denature proteins by destabilizing non-covalent bonds. We posited that SDS could be

used to extract surface proteins from Gram-positive bacteria without lysing the cells. We

incubated strains IC1 and A6 with either PBS (control) or PBS containing 0.5 % SDS, to

remove surface displayed LsLAI, and then quantified surface detection and whole-cell

activity. As expected, we detected LsLAI only on the surface of PBS-treated control A6 but

not the SDS-treated cells by flow cytometry (Fig. S1a). Interestingly, whole-cell activity of

both IC1 and A6 increased approximately 3- and 5-fold, respectively, post-SDS-treatment.

These data suggest LsLAI is present in both the intracellular compartment and membrane

bound of strain A6 with the vast majority of measured whole-cell activity coming from the

intracellular fraction. We confirmed the presence of soluble anchor-LAI fusion (Lp_3014-

LsLAI) in the intracellular fraction of A6 via Western blot analysis (Fig. S1b).

Additionally, we noticed enhanced whole-cell activity that we attribute to

permeabilization/fluidization of the membrane/cell wall due to removal of surface proteins.

While this study only focused one of our six explored anchor protein constructs, we suspect

that these results are consistent amongst the other LsLAI surface display constructs. Thus,

the measured whole-cell activity differing between constructs was likely dependent on the

activity of soluble anchor-LsLAI present in the cytoplasm and not due to surface displayed

enzyme.

Finally, to further elucidate the discrepancy between activity of intracellular and surface

displayed LsLAI constructs we tested the enzymes activity of secreted, unanchored

enzyme. Post-induction, we separated whole cells from culture media and compared the

activity of each fraction. We observed a 6-fold higher in activity in the whole-cell fraction

of the secretion strain compared to wild-type control cells; however, there was no

detectable activity in the supernatant of either strain (Fig. S2a). We confirmed the presence

of LsLAI in the supernatant and soluble cell fraction via Western blot analysis (Fig. S2b).

These data, taken together with our previous experiments, suggest LsLAI is largely in an

inactive state after secretion and that any activity is due to accumulation of active enzyme

in the *L. plantarum* cytoplasm.

Cellular encapsulation improves LAI stability and conversion.

Since surface display failed as a stabilization mechanism, yielding minimally active

enzyme, we evaluated whether whole-cell *L. plantarum* cells encapsulating LsLAI via

cytoplasmic expression (referred to as 'encapsulated LAI') could be used as a stabilization

method instead. Our data show that such encapsulation within *L. plantarum* protected the

enzyme from thermal deactivation at 50 °C; the enzyme retained ~ 85 % of its initial

activity after 24 h (Fig. 3a). Interestingly, resting whole cells at 37 °C displayed no loss of

activity after 24 h; in fact, we observed a slight enhancement in activity. This observation

could be due to changes to cellular morphology or physiology. Cell-encapsulated LsLAI

quickly reached an equilibrium conversion of approximately 60 % in 24 h, overcoming the

thermodynamic limitation of < 50 % conversion presented in the pure enzyme system (Fig.

3b). We believe this happens because the substrate (galactose) and product (tagatose)

differentially partition across the cell membrane. Although initial reaction rates of

encapsulated LsLAI at 37 °C and 50 °C with either galactose or tagatose were lower than

that of their free-enzyme counterparts, encapsulated LsLAI favored reaction in the forward

direction more than in the reverse direction, as determined by initial rates (Fig. S3). At 50

°C the ratio of forward to reverse initial reaction rate for encapsulated LsLAI was 1.8

whereas for the pure enzyme, the ratio was more unfavorable at 0.7.

To further demonstrate that the *L. plantarum* cell acts as a selective barrier, we tested the

capacity of other sugars to inhibit the rate controlling step of galactose to tagatose

isomerization. Tagatose production by cell encapsulated LAI was significantly inhibited in

the presence of glucose while the presence of arabinose had no effect (Fig. S4). Conversely,

glucose had minimal inhibition on tagatose production in pure enzyme reactions while

arabinose (the preferred substrate for LAI) significantly inhibited the production of

tagatose. These data suggest that the rate-controlling step in whole cell system and pure

enzyme are completely different – sugar transporters in whole cell systems and passive

diffusion in pure enzyme systems. Further, since glucose is a highly preferred substrate of

L. plantarum, the fact that glucose inhibits whole cell galactose isomerization suggests that

the rate-controlling step is one or more glucose-preferring hexose transporter(s).

While encapsulation was shown to provide thermal stability and a thermodynamic

advantage, it imposes a kinetic penalty due to transport limitations across the cell

membrane. To test this, we measured initial whole-cell activity under a range of induction

levels to determine the maximum achievable activity and compared the whole-cell activity

to that of the cell lysate at the same conditions. Strain "I" produced the maximum amount

of tagatose (1.5 mM) in 2 h at inducer concentrations ≥ 5 ng/mL (Fig. 3c), indicating that

this represented that the system is at maximum reaction rate under all these inducer levels.

Conversely, lysate of the same cells induced at 5 and 25 ng/mL inducer conditions

generated approximately 4-fold greater tagatose than that of the equivalent whole-cell

catalysts in the same 2 h interval. This further supports the idea that cellular encapsulation

changes the rate-limiting step to transport and imposes a ceiling on reaction kinetics.

SDS permeabilized cells overcome all three barriers to tagatose production.

Seeking to overcome the transport-limited whole-cell production of tagatose, we

investigated the use of permeabilization surfactants – Triton X-100 and SDS. Following the

same optimized protocol as previously reported for permeabilization of *L. plantarum* with 1

% Triton X-100 [28], we observed a 4.2 ± 0.1 fold increase in tagatose production as

compared to untreated cells under the tested conditions (Fig. S5a). We assessed the ability

of 0.005 % – 0.5 % SDS to permeabilize cell encapsulated LsLAI and enhance tagatose

production. Our results showed the optimal concentration of SDS that enhanced tagatose

production via cell permeabilization while still retaining LsLAI intracellularly to be 0.01 %

(Fig. S5b). Tagatose production was enhanced 5.7 ± 0.1 fold after permeabilization with

0.01 % SDS, greater than that of treatment with 1% Triton X-100 under the tested

conditions. Additionally, this concentration of SDS is below the critical micellular

concentration (CMC = 0.24 – 0.26 % at 35 – 40 °C) therefore producing minimal frothing

as compared to 1 % Triton X-100, the CMC of which is reported to be 0.05 % [34]. Although

SDS is commonly used to denature proteins, low concentrations of SDS (0.001 – 0.005 %)

has been previously used to permeabilize *E. coli* – albeit less successfully than with CTAB

(cetyl trimethylammonium bromide) and Triton X-100 [35].

To further understand the effect of SDS-treatment on cell permeability, we measured the

fluorescence generation rate using 5-carboxyfluorescein diacetate (cFDA) as a substrate.

cFDA is an uncharged ester that becomes fluorescent upon hydrolysis by abundant and

nonspecific intracellular esterases. Hydrolysis also results in entrapment of the fluorescent

dye in the cytoplasm, unless the membrane has been damaged. The rate of generated

fluorescence serves as a proxy to determine substrate transport rates assuming hydrolysis

rates are higher than transport rate. Wild-type *L. plantarum* treated with 0.005 %, 0.01 %,

and 0.05 % SDS had 1.6-, 3.0-, and 3.4-fold increase in substrate transport rates,

respectively, compared to untreated cells (Fig. S6). SDS concentrations > 0.5 % resulted in

fluorescence leakage, a sign of damaged cell membrane that may also be expelling other

cytoplasmic content. These data suggest that the treatment with 0.01 % SDS maintains

cellular integrity while increasing tagatose synthesis rate due to a decreased kinetic penalty

presented upon encapsulation.

To test whether the enhanced transport could be attributed to changes to the Gram-positive

cell wall composition, we mildly treated *L. plantarum* encapsulated LAI (strain IC2) with

lysozyme, an enzyme which catalyzes the destruction of the peptidoglycan component of

the gram-positive cell wall. These cells mildly displayed a 4.3-fold increase in activity like

cells chemically treated with SDS detergent compared to untreated cells (Fig. S7 and Fig.

S5). These data suggest that the cell wall poses a transport barrier and SDS treatment may

be modifying the structure/rigidity of the cell wall to allow faster transport; albeit further

direct experimentation should be done to confirm this hypothesis. However, transmission

electron microscopy (TEM) of SDS-treated *L. plantarum* cells found no evidence of pore

formation or leakage of cytoplasmic biological components (Fig. S9).

Finally, we monitored the tagatose production with different LsLAI catalyst preparations at

high loading (Strain IC2 OD_{600} = 40 or equivalent 0.6 mg/mL purified LsLAI) in batch

processes starting with 300 mM galactose. Consistent with our initial experiments, the pure

enzyme preparation was limited to 39 ± 2.3 % conversion after 48 h at 37 °C (Fig. 4).

Increasing the temperature of the reaction to 50 °C allowed for high initial production of

tagatose; however, the enzyme denatured quickly reaching a maximum conversion of only

16 ± 2.8 %. Bacterial encapsulation of LsLAI (IC2 + PBS) produced 47 ± 3.3 % tagatose at

37 °C, improvement over the pure-enzyme system. Encapsulation also provided thermal

stability to the enzyme allowing for sustained activity at 50 °C increasing both the rate of

production (46 ± 2.3 % in 6 h) and equilibrium conversion (83 ± 6.1 %) compared to

unmodified whole-cell catalyst at 37 °C. Rate of production was further increased by

modifying the whole-cell catalyst through SDS permeabilization (IC2+SDS) to achieve 59

± 3.5 % conversion in 6 h, ultimately reaching a similar equilibrium conversion of 85 ± 6.7

% as unpermeabilized cells. Thus, permeabilized, cell-encapsulated LAI overcomes the

three reaction limitations by supporting greater thermal stability, higher equilibrium

conversion, and faster reaction rates than possible by a free enzyme system.

To investigate the potential to increase the average productivity of our system, we increased

the substrate level from 300 mM to 600 mM and 1 M galactose and measured the

production of tagatose using permeabilized *L. plantarum* encapsulating *L. sakei* LAI (strain

IC2 + SDS) over the course of 24 h. Higher substrate loading lowered the final conversion

of the system to 47 % and 39 % at 600 mM and 1 M galactose respectively (Fig. S8). We

observed the highest productivity at low conversions, as expected. After 6 hours, the

highest average productivity of our system was 9.3 g/L/h of tagatose starting from 1 M

galactose substrate loading compared to 5.3 g/L/h starting from 300 mM galactose. Based

on these results we conclude that increasing the substrate loading increases average

productivity at the cost of conversion, which is consistent with many previous studies

[19,36,37].

Discussion:

While an increasing number of enzymes are being used for biocatalysis, poor stability

under reaction conditions, and unfavorable kinetic properties toward non-native substrates

can limit their utility toward biosynthetic functions. Further, for certain classes of enzymes

like isomerases, transglycosidases, and lyases, limitations of endergonic or mild exergonic

thermodynamics are additional hurdles to economical processes. D-Tagatose synthesis from

D-galactose using the enzyme LAI (L-arabinose isomerase) suffers from all three of these

limitations. Our results confirm the presence of all three limitations (stability,

thermodynamic, kinetic) during enzymatic conversion of galactose to tagatose using the

acid-tolerant mesophilic LAI from *Lactobacillus sakei* (LsLAI). At low temperatures (37

°C), the enzyme is stable and quickly reaches equilibrium. However, the product titer is low

due to low equilibrium conversion (36 – 39 %) at low temperatures (this study and

previously reported [33]). At elevated temperatures (50 °C), the initial reaction rates are

faster, but the enzyme quickly inactivates and is unable to realize the higher conversion and

product titer possible at this temperature. These limitations of LAI for tagatose synthesis

have been realized previously, and each issue has been addressed individually. While some

successes have been reported, especially by selecting or engineering enzymes with more

desirable properties [4,11,38–41], this is the first study to address and overcome all three issues

simultaneously.

LAI display on *Bacillus subtilis* endospore surface was shown to improve thermostability

while maintaining free enzyme-like kinetic properties [19]. Therefore, this approach addresses

the stability limitations but not the thermodynamic issue. To assess whether cell surface

display on *L. plantarum* could improve stability of LsLAI, we tested six previously

reported anchor proteins and found there was no correlation linking whole-cell activity and

surface display level/accessibility. We showed evidence for presence of cytoplasmic

LsLAI-anchor fusions and concluded that enzymatic activity seen was largely due to this

active fraction and that the enzyme displayed on the surface was largely inactive. When we

tested secretion constructs (i.e., LsLAI with secretion signal but no anchor), we found

similar results, suggesting that LsLAI cannot fold extracellularly upon secretion through

this pathway and that active enzyme accumulated cytoplasmically. This is surprising since

there is no precedence for such an observation in literature. *L. plantarum* utilizes the Sec-

pathway to secrete proteins [42] and it is generally accepted that translocation occurs co-

translationally[43,44]; therefore, for a protein to be active it must fold properly in the

extracellular environment. While we detected LsLAI on the surface using flow cytometry

and Western blot analysis, we could not correlate display level to activity or subsequently

attribute any detectable activity to the displayed protein at all. Our results suggest the need

to be cautious when attributing activity of surface displayed proteins, even if they are

readily detected. There are other reports of surface displayed enzymes or antigens have

shown lack of correlation between surface detection and activity both in-vitro and in-vivo

[45–47]. We suggest future studies to more directly attribute activity to displayed proteins and

not due to presence of active cytoplasmic proteins in whole or lysed cells. We posit that

overexpression of surface directed proteins may yield a "back-up" resulting in translation of

active protein fusions directly in the cytoplasm. Surface display of LsLAI has been shown

before with *B. subtilis* endospores; however, the construct is assembled on the spore

cytoplasmically [19], without the need for any secretion. Our results contradict reports from

another study that claimed secretion of active LsLAI using *Lactococcus lactis* [1]. However,

closer analysis of the study revealed that measured activity was never attributed directly to

culture supernatant. Rather activity was detected either for whole cells or in the elution

fraction post-chromatographic purification, which could have allowed for refolding to

occur. Thus, we posit that cell-surface display is not a viable option to improve the stability

of this enzyme.

Cellular or abiotic encapsulation has shown to improve enzyme stability against thermal,

pH, solvent, and impurity interactions [24,48]. In the context of tagatose production, whole

cells expressing L-AI has also shown to improve conversion by preferentially partitioning

substrate and product across the *E. coli* membrane barrier through active transport [26];

however, the same was not observed in resting *L. lactis* cells [49]. Our results demonstrate

that encapsulation of LAI in *L. plantarum* behaves in a manner similar to that in Gram-

negative *E. coli* but not Gram-positive *L. lactis*. That is, the cell membrane preferentially

allows passage of galactose to the intracellular space relative to tagatose to aid the forward

reaction. A concurrent advantage of the encapsulation was that the enzyme stability was

markedly improved. Specifically, while the free enzyme rapidly inactivates at 50 °C, the

cell-encapsulated enzyme maintains activity for a prolonged period. Indeed, the resistance

to lysis at elevated temperatures is afforded by the thick cell wall of the Gram-positive *L.*

plantarum, as opposed to Gram-negative *E. coli*, which is known to readily lyse at 50 °C [50].

Thus, while many previous studies have demonstrated the benefits of using whole cells to

enhance enzyme stability, this work is the first to show the twofold benefit by combining

improved stability with substrate-product partitioning. While the selective membrane

circumvents the thermodynamic barrier that limits maximum conversion at moderate

temperatures, it imposes a barrier to mass transport – a penalty to reaction kinetics. We

addressed this issue next.

Cell permeabilization using chemical agents like detergents or solvents have been shown to

non-selectively increase substrate exchange while maintaining biological cellular

components [51,52], although the exact mechanism of this phenomenon is poorly understood.

We used detergents to improve reactant/product transport across the *L. plantarum* cell

membrane without altering its selectivity. In our best-case scenario, sub-micellar

concentration of SDS resulted in significant increase in whole cell reaction rates to free-

enzyme-like rates while still allowing the system to circumvent the thermodynamic barrier

to product conversion. Since the conversion was unchanged, we presume that the detergents

were not making pores in the cell membrane. TEM images of SDS-treated and untreated

cells further confirmed that the cell wall and membrane were intact under reaction

conditions. Interestingly, permeabilization increased the initial reaction rate in the forward

direction 3.8-fold, however the rate of the reverse reaction remained low, indicating that the

membrane retained its selectivity (Fig. S3). Studies with cFDA further confirmed that

cytoplasmic components were not "leaking" into the extracellular space and that the

cellular integrity was maintained. Thus, substrate and product transport were still likely

facilitated through transporters, as in untreated cells, which is also consistent with the

observation that glucose inhibits only whole cell reactions (but not that of free enzyme) and

conversion reaches the same maximum with or without detergent treatment. Since SDS and

Triton-X treatment resulted similar outcomes, we believe there are no specific charge-

related effects of these detergents, either. Adding SDS to a pure enzyme solution did not

alter its kinetic properties, indicating that the detergents do not directly interact with the

enzyme in any appreciable manner. However, our study revealed that treatment with SDS

removed extracellular- and surface-attached proteins, potentially altering the fluidity of the

membrane and/or permeability of the cell wall. The latter is supported by the fact mild

lysozyme treatment also increases reaction rates, suggesting that the thick Gram-positive

peptidoglycan poses a transport barrier.

Having addressed all three limitations (stability, thermodynamics, kinetics), we found that

L. plantarum whole-cell biocatalyst achieved a conversion from of ~ 50 % in 4 h and ~ 85

% 48 h in batch process at 50 °C starting with 300 mM galactose. Improved stability is

apparent by comparison with the free enzyme system, which quickly inactivates at 50 °C

and achieves only 16 % conversion in 48 h. Enhanced equilibrium conversion compared to

free enzyme system can be seen at both 50 °C and 37 °C, but especially in the latter, where

the free enzyme is more stable but is still able to achieve only 13 % conversion in 4 h and

47 % in 48 h. Finally, improved kinetics can be seen compared to unpermeabilized cells

where initial reaction rates at 0.5 mM/min compared to 1.6 mM/min in permeabilized cells.

This rate is higher than the maximum reaction rates sustainable by the free enzyme for

extendable period (initial rate at 37 °C for free enzyme = 0.3 mM/h) due to the ability to

operate at 50 °C. This is the among the highest reported average productivity (5.3 g/L/h

after 6 h) for biocatalytic tagatose production, and the highest conversion reported for a

mesophilic enzyme (~85 %) [19,53]. Average productivity of the modified cell encapsulated

LAI batch process could be improved by increasing the initial concentration of galactose

from 300 mM to 1 M, albeit with lower conversion. Although further improvements to

conversion may be possible by including borate in the reaction buffer, demonstrated

previously, it would also increase purification costs [16,27,49]. Further, this system nearly

matches the conversion obtained in a recent study with a extremophilic LAI at 95 °C and

pH 9.5 that required use of Ni^{2+} and borate [54] but at 50 °C and without use of toxic metals

and salts. Additionally, the system avoids product caramelization, which is known to

happen at high temperatures ($\geq$ 80 °C) [18]. Thus, the system described here demonstrates

how to overcome three major limitations of enzymatic catalysis (stability, thermodynamic,

kinetic) for tagatose production using LAI. We believe that a similar strategy may be

applicable to other biocatalytic systems as well – particularly lyases [55], transglycosidases [56],

and isomerases [17,26,36,57] – where free enzyme performance is poor due to stability,

thermodynamic, and/or kinetic limitations.

Statistics and data reproducibility:

383   All the experiments were conducted using biological replicates and were carried on

384   different days as specified to calculate measure of variability between the samples. All the

385   data shown are mean with error bars representing standard deviation.

Acknowledgments:

This work was partly supported by National Institutes of Health (NIH) Grant DP2HD91798

and R03HD090444 to N.U.N. We thank Dr. Michiel Kleerebezem (Wageningen University

& Research, Wageningen, Netherlands) for sharing strain *L. plantarum* WCFS1 and Dr.

Lars Axelsson (Nofima, Tromsø, Norway) and Dr. Geir Mathiesen (Norwegian University

of Life Sciences, Ås, Norway) for sharing plasmids. Electron microscopy was conducted

utilizing the W.M. Keck Foundation Biological Imaging Facility at the Whitehead Institute.

Special thanks to Nicki Watson for performing the electron microscopy imaging.

Author contributions:

397 J.R.B. and N.U.N. planned the experiments, wrote the manuscript, and analyzed the results.

398 J.R.B conducted all the experiments.

Materials and Methods:

Reagents and Enzymes: All enzymes for cloning were purchased from NEB (Beverly,

MA). DNA primers were ordered from Eurofins Genomics LLC (Louisville, KY) or

GENEWIZ Inc. (Cambridge, MA). Growth media and chemicals were purchased from

Amresco (Solon, OH) or RPI Corp (Mount Prospect, IL). High-purity L-arabinose and D-

galactose were purchased from Sigma-Aldrich (St. Louis, MO). Plasmid Mini Kit I, PCR

Purification and Gel Extraction Kits were obtained from Omega Bio-tek (Norcross, GA).

Mouse-anti-His$_6$ primary antibody and goat anti-mouse IgG H&L (Alexa Fluor® 488) were

purchased from Thermo Fisher Scientific (Waltham, MA).

Bacterial Strains and Culture Conditions: *Lactobacillus plantarum* WCFS1 (kind gift

from Prof. Michiel Kleerebezem, Wageningen University & Research) was cultivated in

deMan−Rogosa−Sharpe (MRS) medium at 30 °C without agitation. *E. coli* strain NEB5α

(New England Biolabs, Ipswich, MA, USA) was grown in Luria−Bertani (LB) medium at

37 °C with shaking at 225 rpm. When needed, media were supplemented with

erythromycin at concentrations of 5 μg/mL and 200 μg/mL for *L. plantarum* and *E. coli*,

respectively. Solid media were prepared by adding 1.5 % (w/v) agar to the respective

media.

Plasmid and Strain Construction: DNA amplification was performed with Phusion High-

Fidelity DNA Polymerase according to the manufacturer's instructions. *E. coli* was used as

host for plasmid propagation before transformation into *L. plantarum*. The amplified

sequences were verified by a commercial sequencing service (GeneWiz). A list of all

strains and plasmids used in this study in Table S2 and Table S3.

L. plantarum surface display plasmids were constructed based on the previously published

pSIP401 based plasmid pLp_3050Ag85B-E6cwa2 (Table S2). The LAI gene from *L. sakei*

23K was synthesized by Twist Biosciences. This gene was subsequently cloned using

primer pair oJRB133 and oJRB134 and inserted into pLp_3050Ag85B-E6cwa2 using BfaI

and HindIII restriction sites to create plasmid pJRB01Q-LSH. pJRB02Q-LSH was created

via was the amplification of Lp_2162 from the *L. plantarum* chromosome using primer pair

oJRB10 and oJRB11 and inserted into pJRB01Q-LSH using MluI and HindIII restrictions

sites. pJRB03Q-LSH was created via was the amplification of Lp_2940 from the *L.*

plantarum chromosome using primer pair oJRB20 and oJRB21 and inserted into pJRB01Q-

LSH using MluI and HindIII restrictions sites. pJRB05Q-LSH was created via the

amplification of Lp_1261 from the plasmid pLp_1261Ag85B-E6 (Table S2) using primer

pair oJRB7 and oJRB8 before being overlapped with *L. sakei* LAI and inserted into

pJRB01Q-LSH using SalI and PmlI restriction sites. pJRB06Q-LSH was created via

amplification of Lp_1452 from the *L. plantarum* chromosome using primer pair oJRB14

and oJRB15 and inserted into pJRB05Q-LSH using SalI and PmlI restrictions sites.

pJRB08Q-LSH was created via the amplification of Lp_3014 from the *L. plantarum*

chromosome using primer pair oJRB82 and oJRB83 and inserted into pJRB05Q-LSH using

SalI and PmlI restrictions sites. pJRB04Q-LSH was created via amplification of the *L. sakei*

LAI gene using primer pair oJRB170 and oJRB171 and inserted into pJRB01Q-LSH using

NdeI and HindIII restriction sites. pJRB09Q-LSH was created via amplification of the *L.*

sakei LAI gene using primer pair oJRB304 and oJRB305 and inserted into pJRB01Q-LSH

using SalI and PmlI restriction sites. Plasmid pJRB14Q-LSH was created via amplification

of the *L. sakei* LAI gene and 6x-His tag using primer pair oJRB457 and oJRB458 and

inserted into pSIP411 (Table S2) using NcoI and HindIII restriction sites.

Each plasmid was transformed into chemically competent *E. coli* NEB5α cells according to

the manufacturer's instructions. *L. plantarum* was transformed using a previously described

electroporation method [58] with slight modifications. Briefly, stationary phase cells were

subcultured to an $OD_{600} = 0.1$ in MRS and allowed to grow until an $OD_{600} = 0.85$. Cells

were harvested via centrifugation at 4 °C and washed twice in ice-cold 10 mM $MgCl_2$ and

once with 0.5 M sucrose and 10 % glycerol. Cells were resuspended in 1:50 (v/v) of the

same solution and kept on ice prior to transformation. 100 ng of plasmid DNA was diluted

into 5 μL of water and mixed with 50 μL electrocompetent cells immediately before adding

to a chilled 1 mM electroporation cuvette. The bacteria were transformed using a BioRad

Pulser at 1300 V with a fixed time constant of 5 ms. 1 mL of warm MRS was added and the

cells transferred to a new centrifuge tube and allowed to recover statically at 37 °C for 3 h

before plating on appropriate media containing selective antibiotic. Positive transformants

were confirmed using colony PCR.

Enzyme Purification: *L. plantarum* cells carrying pJRB14Q-LSH were grown and induced

as described previously. Cells were washed in lysis/equilibration buffer (50 mM sodium

phosphate, 300 mM sodium chloride, 15 % glycerol, pH 8.0) and resuspended in lysis

buffer containing 10 mM lysozyme and incubated at 37 °C with shaking for one hour. Lysis

was achieved via sonication and soluble protein collected as described previously.

Purification was achieved via immobilized metal affinity chromatography (IMAC) using an

N-terminal His_6 tag. TALON metal affinity resin (Invitrogen) was used as directed by the

manufacturer. A single elution of protein was achieved in elution buffer containing 500

mM imidazole. Active fractions were pooled, and the buffer exchanged with PBS

containing 1 mM $MgCl_2$ (PBSM) using a Microsep 10 kDa Omega centrifugation spin

column (PALL Corp; Port Washington, NY). Purification was confirmed via SD-PAGE

analysis. Purified enzyme was stored in PBSM with 20 % glycerol at −80 °C prior to use.

Enzyme Analysis: *L. plantarum* wild-type (WT) cells or those carrying plasmids

pJRB01Q-LSH (A1), pJRB02Q-LSH (A2), pJRB03Q-LSH (A3), pJRB04Q-LSH (IC1),

pJRB05Q-LSH (A4), pJRB06Q-LSH (A5), pJRB08Q-LSH (A6) were cultured in MRS

media with antibiotic as needed at 37 °C overnight. Overnight cultures were diluted 1:50

(approximate $OD_{600} = 0.1$) in MRS media with antibiotic as needed and grown at 37 °C for

2.5 h. Cells were then induced with IP-673 (synthesized by Life Technologies; Carlsbad,

CA) at a final concentration of 25 ng/ml and $MgCl_2$ was added to a final concentration of 1

mM to support enzyme folding. Cultures were incubated at 37 °C for an additional 3 h

before harvesting the cells by centrifugation at $3000 \times g$ for 5 min.

Cell pellets were washed in 1 mL of 20 mM phosphate buffer (PB) pH 6.8 and transferred

to a 1.7 mL centrifuge tube. Cells were washed twice with 1 mL of PB before being

suspended in 200 μL PB buffer. The enzyme activity assay was performed in a 96-well

plate as follows: 20 μL cell suspension was incubated with 0.8 mM $MnCl_2$, 0.8mM $MgCl_2$,

sugar (200 mM galactose or 200 mM glucose) and diluted to 200 μL with PB. The cell

density within each well was measured for data normalization at the start of the reaction.

The plates were sealed and incubated at 37 °C for 2 h. Enzyme activity was measured

colorimetrically using the cysteine– carbazole–sulfuric-acid method (CCSAA) [59].

Absorbance at 560 nm was measured after 1 h of incubation at room temperature. Triplicate

technical replicates of each cell suspension were analyzed and averaged due to variability

in the assay itself.

To determine the conversion achieved with our balanced system, *L. plantarum* cells

carrying pJRB14Q-LSH (IC2) and wild-type were grown and induced as described

previously with slight modification. Cells were induced with 25 ng/mL IP-673 for 8 h

instead of 3 h to maximize the cell density of the culture. Cells were harvested via

centrifugation and washed twice in phosphate-buffered saline pH 7.4 with 1.0 mM $MgCl_2$

(PBSM). Cells to be permeabilized were diluted to an $OD_{600} = 3.0$ before incubation with

0.0 1% SDS in PBSM for 30 min at room temperature. The enzyme activity assay was

performed in 5 mL tubes as follows: cell suspensions adjusted to an $OD_{600} = 40$ were

incubated in phosphate-buffered saline (PBS) pH 7.4 with 1.0 mM $MgCl_2$, 300 mM

galactose and adjusted to 5 mL with H_2O. The conversion and productivity at increasing

concentrations of galactose was performed by increasing the substrate concentration to 600

mM or 1 M with the same assay conditions.

Initial rates of reaction for both pure-enzyme, unmodified, and modified whole-cell

catalysts were determined. *L. plantarum* cells carrying pJRB14Q-LSH were grown and

induced for 8 h as described previously. Permeabilized cells were prepared as described

previously. The enzyme activity assay was performed in 200 μL PCR tubes as follows:

purified enzyme or cell suspensions were incubated in PBSM, variable substrate

concentration of galactose or tagatose, and adjusted to 100 μl with H_2O. Reactions were

initiated by adding $1/10^{th}$ volume cell suspensions adjusted to an $OD_{600} = 10$ or 2 μg of

purified enzyme to prewarmed reaction mixtures in a PCR block at either 37 or 50 °C.

Reactions proceeded for 6 min before termination to measure initial reaction rates. Cell

suspensions were immediately centrifuged for 1 min using a benchtop quick-spin centrifuge

and the supernatant was transferred to a 96-well 0.2 μm centrifugal filter plate with

collection plate. The remaining solution was filtered via centrifugation and stored at -80C

until analysis. Purified enzyme reactions were terminated by adding $1/10^{th}$ volume 2M

perchloric acid. Prior to HPLC analysis the reactions were neutralized using a sodium

hydroxide PBS buffered solution. "$k_{initial}$" is defined as the initial reaction rate of each

catalyst at a given, non-saturating substrate concentration and stopped the reaction prior to

10% total conversion.

Enzyme stability was measured by incubating either purified enzyme or unmodified whole

cell catalysts in PBSM at 37 or 50 °C statically for the duration of the experiment. At

determined time intervals and aliquot of catalyst was added to PBSM reaction mixture

containing 300mM galactose at the same temperature as the catalyst. Reactions proceeded

for 20 min before termination as described previously. Tagatose production was confirmed

via HPLC analysis.

Equilibrium conversion was measured by in incubating either purified enzyme or

unmodified whole cell catalysts in PBSM at 37 °C. Purified enzyme was incubated with

either 10 mM galactose, 5 mM of galactose and tagatose, or 10 mM tagatose. The reaction

was sampled every 24 h and additional enzyme was added after sampling due to incomplete

reaction due to enzyme degradation. Whole cell catalyst was incubated with either 30 mM

galactose, 15 mM of galactose and tagatose, or 30 mM tagatose.

The selectivity of the *L. plantarum* cell barrier was tested by comparing the inhibition of

the transport of galactose compared to that of the pure enzyme. *L. plantarum* strain IC2

carrying pJRB14Q-LSH was grown and induced for 8 h as described previously. *L.*

plantarum encapsulating LAI (IC2) or purified LAI (PURE) was added to reaction mixture

containing and incubated at 37 °C. Reactions proceeded for 20 min before termination as

described previously.

HPLC Analysis: Agilent Infinity HPLC system (Agilent; Santa Clara, CA) equipped with

a Hi-Plex Ca+ 300 x 7.7mm column with a guard column and detected using Agilent

Infinity 1260 ELSD detector. The mobile phase was filtered deionized water run at 85 °C

with a flow rate of 0.6 mL/min. The ELSD detector's evaporation temperature was set at 90

°C, the nebulizer temperature set to 50 °C, and nitrogen flow rate at 1.6 SLM (standard liter

per minute). For calculation of the reaction product(s) L-arabinose, D-galactose, L-ribulose,

and D-tagatose standards were included in the run.

Flow Cytometry and Immunostaining: *L. plantarum* WT cells or those carrying plasmids

pJRB01QLs-pJRB08QLs were grown and induced as described previously. Cells were

collected by centrifugation at 5000 ×g for 5 min and washed twice in PBS before being

suspended to an $OD_{600} = 1.0$. An aliquot of 200 μL cells was pelleted and was resuspended

in 200 μL PBSA (PBS with 2 % bovine serum albumin). The suspension of cells rested on

ice for 30 min before the cells were pelleted and solution removed. The cells were then

resuspended in 50 uL of PBSA containing 1 μg mouse-anti- His_6 primary antibody. The

cells were incubated for 1h at room temperature with gentle rocking. Cells were collected

and washed thrice with PBSA before being suspended in 50 μL of PBSA containing 0.4 μg

goat anti-mouse IgG H&L (Alexa Fluor® 488). The cells were incubated for one fifteen

minutes on ice and blocked from light. Finally, cells were washed thrice in PBSA and

resuspended in 500ul of PBSA. A 20ul aliquot was diluted 10X in PBSA and transferred to

a flat-bottom 96-well plate for flow cytometric analysis. A positive control sample was

used to calibrate the fluorescence intensity. A negative control sample was used to gate the

level of autofluorescence of the cells and any non-specific secondary antibody binding (Fig.

S10). A total of 1×10^5 cell counts per sample plotted on a 'logical' or bi-exponential x-

axis. Data analyzed using FCS Express 6 (De Novo Software, Glendale,CA). Surface

detection level was calculated as the fraction of cells with a measured fluorescence greater

than that of the gated negative control.

Surface Treatment: *L. plantarum* WT cells or those carrying either pJRB04Q-LSH or

pJRB08Q-LSH were grown and induced as previously described. Cells were washed in

PBS twice before further processing. A concentrated cell solution was incubated in PBS

containing either 0 % or 0.5 % SDS for 1 h before being pelleted by centrifugation. The

SDS solution was removed and the cell pellet washed twice with PBS. Surface detection

was measured via immunofluorescence flow cytometry as described previously. Whole-cell

activity was measured via the CCSAA after a 1-hour incubation in 200 mM arabinose and 1

mM $MgCl_2$ in 100 mM sodium acetate buffer pH 5.0 at 37 °C.

Western Blot Analysis: *L. plantarum* WT cells or those carrying either pJRB04Q-LSH or

pJRB08Q-LSH were grown and induced as previously described. A concentrated cell

solution was incubated with 0.05 % SDS in PBS for 30 min at 37 °C before being pelleted.

The SDS buffer solution was removed and stored separately. The cell pellet was then

washed twice with PBS. A fresh cell solution of the same concentration was prepared in

PBS as a control. Both aliquots of cells were then incubated with 10 mg/mL lysozyme for 1

h at 37 °C before being lysed via sonication. Sonication was performed using a Branson

150 system equipped with a microtip probe sonicating samples on ice at 55 % amplitude in

30 s ON 2 min OFF cycles for a total of 5 min of ON time. The insoluble cell fraction

containing membrane components and the insoluble protein fraction was pelleted via

centrifugation at 20,000 ×g for 5 min. The soluble fraction was removed, filtered through

0.4 μm sterile filter, and stored separately. An aliquot of each of the cell fractions for the

three cell types was added to denaturing loading dye and incubated at 98 °C for 15 min. 10

μL of each sample were added to a 4 % - 12 % Bis-Tris PAGE gel in MOPS buffer and run

at 75 V for 15 min followed by 110 V for 1.5 h. After separation the gel was removed from

the cassette and washed twice in deionized water. The proteins were next transferred to a

PVDF membrane for Western blot analysis. Transfer was visually confirmed through the

transfer of the stained protein ladder. The membrane was blocked overnight at room

temperature with 5 % skim milk in Tris buffered saline containing 1 % Tween-20 (TBST).

The membrane was washed twice with TBST before being incubated with 0.2 μg/mL

primary antibody in blocking buffer for 2 h at room temperature. The membrane was

washed twice with TBST before being incubated with 2 ng/mL of the horseradish

peroxidase (HRP) conjugated secondary antibody at room temperature for 1 h. The

membrane was then washed twice with TBST before incubation in the substrate containing

solution (Thermo SuperSignal West) for 15 min at room temperature. The

chemiluminescent signal was captured electronically via a camera set to a 30 min exposure

time.

Cell Permeabilization: *L. plantarum* WT cells or those carrying pJRB14Q-LSH (IC2)

were grown and induced and washed in PBS as previously described. Cells were

resuspended to an OD_{600} = 3.0 in 200 μL of PBS. Samples were either incubated with 1%

Triton X-100 [28], 0.1 % SDS, 0.1 μM chicken egg lysozyme, or no treatment at room

temperature for 1 h. The solution was removed, and the cell pellet washed twice with PBS

before enzymatic activity of the whole-cells were measured as described previously using

200 mM galactose in PBS pH 7.4 containing 1 mM $MgCl_2$. Activity was quantified via the

CCSAA and normalized to cell concentration as described previously.

The optimal concentration of SDS to maximize whole-cell activity was determined. Cells

were prepared as previously described. Cell lysate from the equivalent cell density was

prepared via lysozyme treatment via sonication as described previously. Permeabilized

samples were either incubated with varying concentrations of SDS, or no treatment at room

temperature for 30 min. The solution was removed, and the cell pellet washed twice with

PBS before enzymatic activity of the whole-cells were measured as described previously

using 200 mM galactose. Activity was quantified via the CCSAA and normalized to cell

concentration as described previously.

Cell permeabilization was also tested using carboxyfluorescein diacetate (cFDA) (Sigma-

Aldrich) as substrate. Cell suspensions were diluted to OD_{600} = 0.5 in PBS in a black

polystyrene 96-well plate. The reaction was started by adding 10 mM cFDA to the cell

suspensions. Transport of cFDA was estimated by measuring the fluorescent signal released

when cFDA becomes activated upon intracellular cleavage via nonspecific esterases. The

total fluorescence was measured on a plate-reader using an excitation and emission

wavelength of 485 nm and 520 nm, respectively, every 15 s for 30 min. Apparent transport

rates were calculated from the slope of the linear fit to the data.

Electron Microscopy: The cells were fixed in 2.5 % gluteraldehyde, 3 %

paraformaldehyde with 5 % sucrose in 0.1 M sodium cacodylate buffer (pH 7.4), pelleted,

and post fixed in 1 % OsO_4 in veronal-acetate buffer. The cells were stained *en block*

overnight with 0.5 % uranyl acetate in veronal-acetate buffer (pH 6.0), then dehydrated and

embedded in Embed-812 resin. Sections were cut on a Leica EM UC7 ultra microtome with

a Diatome diamond knife at a thickness setting of 50 nm, stained with 2 % uranyl acetate,

and lead citrate. The sections were examined using a FEI Tecnai spirit at 80 KV and

photographed with an AMT ccd camera.

Data availability

Data supporting the findings of this work are available within the paper and its

Supplementary Information files. A reporting summary for this article is available as a

Supplementary Information file. The source data underlying Figs. 1, 2, 3, 4, and

Supplementary Figures are provided as a Source Data file. All other data are available from

the corresponding author upon reasonable request.

Figure 1: Free-enzyme LsLAI suffers from poor thermal stability, low thermodynamic equilibrium, and kinetic limitations. a) Initial turnover rates of purified LsLAI compared at medium (100 mM, green) and high (400 mM, orange) substrate loading at low (37 °C) and high (50 °C) temperatures. Comparison of the non-native substrate, galactose at low (GAL-37C) and high (GAL-50C) temperature, to that of native substrate, arabinose at low temperature (ARA-37C). b) Loss of activity of purified LsLAI over time incubated at 37 °C (green) or 50 °C (orange). Half-life calculated from first-order decay equation. c) Equilibrium conversion of purified LsLAI starting from 10 mM total substrate with galactose to tagatose ratio at 1:0 (10 mM galactose, purple), 1:1 (5 mM each galactose and tagatose, orange), or 0:1 (10 mM tagatose, green) incubated at 37 °C. Additional enzyme was added every 24 h to account for thermal inactivation. The data are means from three biological replicates. *** = p < 0.001 (Students t-test).

Figure 2: LsLAI surface display and activity. a) Flow cytometry analysis of *L. plantarum* wild-type cells (green), cells expressing intracellular His$_6$-tagged LsLAI (orange), or cells expressing LsLAI fused to native surface anchor proteins A1-A6 for display (blue) with 10^5 counts per sample. Data is normalized to number of counts. b) Comparing the amount of tagatose produced from 200 mM galactose in 2 h (hashed) with the positive percentage of the population with surface detection intensity above that of wild-type "WT" for intracellularly expressed "IC1" or anchor protein A1-A6 (colored) based on negative gate. c) Scatterplot of each replicate measurement correlating surface detection vs. tagatose produced. There is no significant correlation between the display level and activity based on Pearson Product Moment Correlation test (Correlation coefficient = −0.395, p = 0.0564). The data are means from three biological replicates.

Figure 3: Encapsulation of LsLAI improves equilibrium conversion and provides thermal stability, albeit at a kinetic penalty a) Loss of activity of encapsulated LsLAI (IC2) over time incubated in PBSM pH 7.4 at 37°C (orange) or 50°C (blue). Half-life calculated assuming first-order decay. b) Equilibrium conversion of encapsulated LsLAI starting from 30 mM total substrate: 1:0 (30 mM galactose, purple), 1:1 (15 mM each, orange), or 0:1 (30 mM tagatose, green) galactose:tagatose respectively. c) Tagatose production from 200 mM galactose of *L. plantarum* wild type "WT" cells (green), IC2 cells expressing LsLAI intracellularly "I" (purple), or crude lysate of the same IC2 cells expressing LsLAI cytoplasmically "LYS" (orange) at different induction concentrations. The data are means from three biological replicates.

Initial rate (mM/min)	0.3	0.5	1.6	0.5	0.7
6 h (%)	22	46	59	13	8
48 h (%)	47	82	85	39	16

Figure 4: Modified encapsulation of LsLAI maximizes tagatose production.
Batch tagatose production over time starting from 300 mM galactose. Encapsulated LsLAI (IC2) was tested for tagatose production as unmodified (PBS) or SDS permeabilized (SDS) compared to purified enzyme at 37 °C or 50 °C to demonstrate its advantages. Encapsulated LsLAI is "PBS-37C" (green), encapsulated LsLAI at 50 °C is "PBS-50C" (pink), permeabilized encapsulated LsLAI at 50 °C is "SDS-50C" (blue), purified free-enzyme LsLAI at 37 °C is "PURE-37C" (orange), and purified free-enzyme LsLAI at 50 °C is "PURE-50C" (purple). Inset plot shows 0 - 1 h data to highlight initial reaction rates of each catalyst for comparison purposes. Salient features are summarized in table. The data are means from three biological replicates.

References:

1.   Rhimi, M. et al. The secreted L-arabinose isomerase displays anti-hyperglycemic

effects in mice. Microb. Cell Fact. 14, 204 (2015).

2.   Luecke, K. J. & Bell, L. N. Thermal stability of tagatose in solution. J. Food Sci. 75,

C346-51 (2010).

3.   Bertelsen, H., Andersen, H. & Tvede, M. Fermentation of D-tagatose by human

intestinal bacteria and dairy lactic acid bacteria. Microb. Ecol. Health Dis. 13, 87–95

(2001).

4.   Xu, Z., Li, S., Feng, X., Liang, J. & Xu, H. L-Arabinose isomerase and its use for

biotechnological production of rare sugars. Appl. Microbiol. Biotechnol. 98, 8869–8878

(2014).

5.   Cheetham, P. S. J. & Wootton, a. N. Bioconversion of D-galactose into D-tagatose.

Enzyme Microb. Technol. 15, 105–108 (1993).

6.   Xu, W., Zhang, W., Zhang, T., Jiang, B. & Mu, W. L-arabinose isomerases:

Characteristics, modification, and application. Trends Food Sci. Technol. 78, 25–33 (2018).

7.   Xu, Z. et al. A novel L-arabinose isomerase from *Lactobacillus fermentum*

CGMCC2921 for D-tagatose production: Gene cloning, purification and characterization. J.

Mol. Catal. B Enzym. 70, 1–7 (2011).

8.   Waltman, M. J., Yang, Z. K., Langan, P., Graham, D. E. & Kovalevsky, A.

Engineering acidic *Streptomyces rubiginosus* D-xylose isomerase by rational enzyme

design. Protein Eng. Des. Sel. 27, 59–64 (2014).

9.   Zheng, Z., Mei, W., Xia, M., He, Q. & Ouyang, J. Rational design of *Bacillus*

coagulans NL01 L-arabinose isomerase and use of its F279I variant in D-tagatose

production. (2017).

10.   Fan, C. et al. Engineering of *Alicyclobacillus hesperidum* L-arabinose isomerase for

improved catalytic activity and reduced pH optimum using random and site-directed

mutagenesis. Appl. Biochem. Biotechnol. (2015).

11.   Oh, H. J., Kim, H. J. & Oh, D. K. Increase in D-tagatose production rate by site-

directed mutagenesis of L-arabinose isomerase from *Geobacillus thermodenitrificans*.

Biotechnol. Lett. 28, 145–149 (2006).

12.   Hong, Y.-H., Lee, D.-W., Pyun, Y.-R. & Lee, S. H. Creation of metal-independent

hyperthermophilic L-arabinose isomerase by homologous recombination. J. Agric. Food

Chem. 59, 12939–12947 (2011).

13.   Prabhu, P., Jeya, M. & Lee, J. K. Probing the molecular determinant for the

catalytic efficiency of L-arabinose isomerase from *Bacillus licheniformis*. Appl. Environ.

Microbiol. 76, 1653–1660 (2010).

14.   Choi, J. M. et al. Structure of the thermophilic L-arabinose isomerase from

Geobacillus kaustophilus reveals metal-mediated intersubunit interactions for activity and

thermostability. Arch. Biochem. Biophys. 596, 51–62 (2016).

15.   Kim, H. J. & Oh, D. K. Purification and characterization of an L-arabinose

isomerase from an isolated strain of *Geobacillus thermodenitrificans* producing D-tagatose.

J. Biotechnol. 120, 162–173 (2005).

16.   Li, Y., Zhu, Y., Liu, A. & Sun, Y. Identification and characterization of a novel L-

arabinose isomerase from *Anoxybacillus flavithermus* useful in D-tagatose production.

Extremophiles 15, 441–450 (2011).

17.   Hong, Y. H., Lee, D. W., Pyun, Y. R. & Lee, S. H. Creation of metal-independent

hyperthermophilic L-arabinose isomerase by homologous recombination. J. Agric. Food

Chem. 59, 12939–12947 (2011).

18.   Kwon, S. Y. & Baek, H. H. Effects of temperature, pH, organic acids, and sulfites

on tagatose browning in solutions during processing and storage. Food Sci. Biotechnol. 23,

677–684 (2014).

19.   Liu, Y. et al. Efficient production of D-Tagatose using a food-grade surface display

system. J. Agric. Food Chem. 62, 6756–62 (2014).

20.   Zhang, Y.-W., Jeya, M. & Lee, J.-K. Enhanced activity and stability of L-arabinose

isomerase by immobilization on aminopropyl glass. Appl. Microbiol. Biotechnol. 89,

1435–1442 (2011).

21.   Bortone, N. & Fidaleo, M. Immobilization of the recombinant (His)6-tagged L-

arabinose isomerase from *Thermotoga maritima* on epoxy and cupper-chelate epoxy

supports. Food Bioprod. Process. 95, 155–162 (2015).

22.   Kim, P., Yoon, S.-H., Roh, H.-J. & Choi, J.-H. High production of D-tagatose, a

potential sugar substitute, using immobilized L-arabinose isomerase. Biotechnol. Prog. 17,

208–210 (2001).

23.   Hong, Y. H. et al. Production of D-tagatose at high temperatures using immobilized

Escherichia coli cells expressing L-arabinose isomerase from *Thermotoga neapolitana*.

Biotechnol. Lett. 29, 569–574 (2007).

24.   Lin, B. & Tao, Y. Whole-cell biocatalysts by design. Microb. Cell Fact. 16, 106

(2017).

25.   Liu, J.-J. et al. Overcoming the thermodynamic equilibrium of an isomerization

reaction through oxidoreductive reactions for biotransformation.

26.   Kim, J. H. et al. Differential selectivity of the *Escherichia coli* cell membrane shifts

the equilibrium for the enzyme-catalyzed isomerization of galactose to tagatose. Appl.

Environ. Microbiol. 74, 2307–2313 (2008).

27.   Xu, Z. et al. Production of D-tagatose, a functional sweetener, utilizing alginate

immobilized *Lactobacillus fermentum* CGMCC2921 cells. Appl. Biochem. Biotechnol.

166, 961–73 (2012).

28.   Jayamuthunagai, J., Srisowmeya, G., Chakravarthy, M. & Gautam, P. D-Tagatose

production by permeabilized and immobilized *Lactobacillus plantarum* using whey

permeate. Bioresour. Technol. 235, 250–255 (2017).

29.   Reider Apel, A., Ouellet, M., Szmidt-Middleton, H., Keasling, J. D. &

Mukhopadhyay, A. Evolved hexose transporter enhances xylose uptake and glucose/xylose

co-utilization in *Saccharomyces cerevisiae*. Sci. Rep. 6, 19512 (2016).

30.   Miskovic, L. et al. A design–build–test cycle using modeling and experiments

reveals interdependencies between upper glycolysis and xylose uptake in recombinant *S.*

cerevisiae and improves predictive capabilities of large-scale kinetic models. Biotechnol.

Biofuels 10, 166 (2017).

31.   Bober, J. R., Beisel, C. L. & Nair, N. U. Synthetic biology approaches to engineer

probiotics and members of the human microbiota for biomedical applications. Annual

Review of Biomedical Engineering 20, (2018).

32.   Mays, Z. J. & Nair, N. U. Synthetic biology in probiotic lactic acid bacteria: At the

frontier of living therapeutics. Curr. Opin. Biotechnol. 53, 224–231 (2018).

33.   Rhimi, M. et al. The acid tolerant L-arabinose isomerase from the food grade

Lactobacillus sakei 23K is an attractive D-tagatose producer. Bioresour. Technol. 101,

9171–9177 (2010).

34.   Mukerjee, P. & Mysels, K. J. Critical micelle concentrations of aqueous surfactant

systems. J. Pharm. Sci. 61, 319 (1972).

35.   Tao, F., Zhang, Y., Ma, C. & Xu, P. One-pot bio-synthesis: N-Acetyl-D-neuraminic

acid production by a powerful engineered whole-cell catalyst. Sci. Rep. 1, 142 (2011).

36.   Guo, Q. et al. Enhanced D-tagatose production by spore surface-displayed L-

arabinose isomerase from isolated *Lactobacillus brevis* PC16 and biotransformation.

Bioresour. Technol. 247, 940–946 (2018).

37.   Mei, W., Wang, L., Zang, Y., Zheng, Z. & Ouyang, J. Characterization of an L-

arabinose isomerase from *Bacillus coagu*lans NL01 and its application for D-tagatose

production. BMC Biotechnol. 16, 1–11 (2016).

38.   Rhimi, M. et al. Rational design of *Bacillus stearothermophilus* US100 L-arabinose

isomerase: Potential applications for D-tagatose production. Biochimie 91, 650–653 (2009).

39.   Hong, Y., Lee, D., Pyun, Y. & Lee, S. H. Creation of metal-independent

hyperthermophilic L-arabinose isomerase by homologous recombination. J. Agric. Food

Chem. 59, 12939–12947 (2011).

40.   Fan, C. et al. Engineering of *Alicyclobacillus hesperidum* L-arabinose isomerase for

improved catalytic activity and reduced pH optimum using random and site-directed

mutagenesis. Appl. Biochem. Biotechnol. 177, 1480–1492 (2015).

41.   Kim, B. J., Hong, S. H., Shin, K. C., Jo, Y. S. & Oh, D. K. Characterization of a

F280N variant of L-arabinose isomerase from *Geobacillus thermodenitrificans* identified as

a D-galactose isomerase. Appl. Microbiol. Biotechnol. 9271–9281 (2014).

doi:10.1007/s00253-014-5827-z

42.   Brurberg, M. et al. Genome-wide analysis of signal peptide functionality in

Lactobacillus plantarum WCFS1. BMC Genomics 10, 425 (2009).

43.   Schneewind, O. & Missiakas, D. Sec-secretion and sortase-mediated anchoring of

proteins in Gram-positive bacteria. Biochim. Biophys. Acta 1843, 1687–1697 (2014).

44.   Mori, H. & Ito, K. The Sec protein-translocation pathway. Trends Microbiol. 9,

494–500 (2001).

45.   Nguyen, H.-M. et al. Display of a β-mannanase and a chitosanase on the cell surface

of *Lactobacillus plantarum* towards the development of whole-cell biocatalysts. Microb.

Cell Fact. 15, 169 (2016).

46.   Kuczkowska, K. et al. Immunogenic properties of *Lactobacillus plantarum*

producing surface-displayed *Mycobacterium tuberculosis* antigens. Appl. Environ.

Microbiol. AEM.02782-16 (2016). doi:10.1128/AEM.02782-16

47.   Kuczkowska, K., Mathiesen, G., Eijsink, V. G. H. & Øynebråten, I. *Lactobacillus*

plantarum displaying CCL3 chemokine in fusion with HIV-1 Gag derived antigen causes

increased recruitment of T cells. Microb. Cell Fact. 14, 1 (2015).

48.   Mohamad, N. R., Marzuki, N. H. C., Buang, N. A., Huyop, F. & Wahab, R. A. An

overview of technologies for immobilization of enzymes and surface analysis techniques

for immobilized enzymes. Biotechnol. Biotechnol. Equip. 29, 205–220 (2015).

49.   Salonen, N., Salonen, K., Leisola, M. & Nyyssölä, A. D-Tagatose production in the

presence of borate by resting *Lactococcus lactis* cells harboring *Bifidobacterium longum* L-

arabinose isomerase. Bioprocess Biosyst. Eng. 36, 489–497 (2013).

50.   Russell, A. D. & Harries, D. Damage to *Escherichia coli* on exposure to moist heat.

Appl. Enviromental Microbiol. 16, 1394 (1968).

51.   de Carvalho, C. C. C. R. Whole cell biocatalysts: Essential workers from nature to

the industry. Microbial Biotechnology 10, 250–263 (2017).

52.   Mattei, B., Lira, R. B., Perez, K. R. & Riske, K. A. Membrane permeabilization

induced by Triton X-100: The role of membrane phase state and edge tension. Chem. Phys.

Lipids 202, 28–37 (2017).

53.   Nguyen, T. K., Hong, M. G., Chang, P. S., Lee, B. H. & Yoo, S. H. Biochemical

properties of L-arabinose isomerase from *Clostridium hylemonae* to produce D-tagatose as a

functional sweetener. PLoS One 13, e0196099 (2018).

54.   Chen, H. et al. Surface display of the thermophilic lipase Tm1350 on the spore of

Bacillus subtilis by the CotB anchor protein. Extremophiles 19, 799–808 (2015).

55.   Parmeggiani, F., Weise, N. J., Ahmed, S. T. & Turner, N. J. Synthetic and

therapeutic applications of ammonia-lyases and aminomutases. Chemical Reviews 118, 73–

118 (2018).

56.   U. Nair, N., A. Denard, C. & Zhao, H. Engineering of enzymes for selective

catalysis. Curr. Org. Chem. 14, 1870–1882 (2010).

57.   Cheng, L., Mu, W. & Jiang, B. Thermostable L-arabinose isomerase from *Bacillus*

stearothermophilus IAM 11001 for D-tagatose production: gene cloning, purification and

characterization. J. Sci. Food Agric. 90, 1327–33 (2010).

58.   Alegre, M. T., Rodríguez, M. C. & Mesas, J. M. Transformation of *Lactobacillus*

plantarum by electroporation with in vitro modified plasmid DNA. FEMS Microbiol. Lett.

241, 73–7 (2004).

59.   Dische, Z. & Borenfreund, E. A new spectrophotometric method for the detection

and determination of keto sugars and trioses. Chem, J Biol 192, 583–587 (1951).