Title: Stable Pom1 clusters form a glucose-modulated concentration gradient that regulates mitotic entry

Authors: Corey A.H. Allard, Hannah E. Opalko, and James B. Moseley*

Author Affiliations: Department of Biochemistry and Cell Biology

The Geisel School of Medicine at Dartmouth, Hanover, NH 03755

*Corresponding author

james.b.moseley@dartmouth.edu

Abstract: Control of cell size requires molecular size sensors that are coupled to the cell cycle. Rod-shaped fission yeast cells divide at a threshold size partly due to Cdr2 kinase, which forms nodes at the medial cell cortex where it inhibits the Cdk1-inhibitor Wee1. Pom1 kinase phosphorylates and inhibits Cdr2, and forms cortical concentration gradients from cell poles. Pom1 inhibits Cdr2 signaling to Wee1 specifically in small cells, but the time and place of their regulatory interactions were unclear. We show that Pom1 forms stable oligomeric clusters that dynamically sample the cell cortex. Binding frequency is patterned into a concentration gradient by the polarity landmarks Tea1 and Tea4. Pom1 clusters colocalize with Cdr2 nodes, forming a glucose-modulated inhibitory threshold against node activation. Our work reveals how Pom1-Cdr2-Wee1 operates in multiprotein clusters at the cortex to promote mitotic entry at a cell size that can be modified by nutrient availability.

Introduction:

Many cell types display a remarkable ability to maintain a constant size during rapid cycles of growth and division (Fantes and Nurse, 1977; Ginzberg et al., 2015; Jorgensen and Tyers, 2004; Lloyd, 2013). Such cell size control is a systems-level property that emerges from the integration of multiple size-dependent signal transduction pathways. Each signaling pathway is comprised of tunable biochemical parameters, including gene-expression and post-translational modifications such as protein phosphorylation (Alberghina et al., 2009). One major challenge in cell size research is to understand the biochemical mechanisms of signal transduction in each pathway, and what makes them size-dependent. These control systems can generate size homogeneity for a given cell type, but cell size is also an adaptable property. For example, nutritional cues and other environmental factors can alter cell size (Fantes and Nurse, 1977; Kelkar and Martin, 2015; Shiozaki, 2009; Yanagida Mitsuhiro et al., 2011). Thus, a second major challenge in cell size research is to understand how size-dependent signaling pathways respond to changes in cell metabolism and stress.

In eukaryotic cells, these signaling pathways lead to regulated activation of the conserved cyclin-dependent kinase Cdk1 (Harashima et al., 2013). Activated Cdk1 triggers mitotic entry and the cascade of events that lead to cell division (Gould and Nurse, 1989; Simanis and Nurse, 1986). The fission yeast *Schizosaccharomyces pombe* is an excellent model system to study size-dependent signaling pathways that regulate Cdk1. Genetic screens performed in past decades have identified many conserved factors that regulate Cdk1, but how these factors form size-dependent signaling pathways remains less clear. Fission yeast cells have a simple geometry that facilitates cell size studies. These cylindrical cells maintain a constant cell width, and grow by linear extension during interphase (Fantes and Nurse, 1977; Moreno et al., 1989). A network of cell polarity proteins positioned at cell tips restricts growth to these sites and maintains proper cell morphology (Chang and Martin, 2009). As a result, cell length doubles in one cell cycle, and many aspects of cell geometry scale with this increase in cell length (Gu and Oliferenko, 2019; Neumann and Nurse, 2007). Recent studies used cell shape mutants to show that fission yeast cells primarily measure surface area, not length or volume, for

56 cell size control (Facchetti et al., 2019; Pan et al., 2014). A critical next step is to
57 understand how signaling pathways that regulate Cdk1 might operate at the cell surface
58 in a size-dependent manner.
59
Cdk1 activity is established by the opposing activities of the inhibitory protein
kinase Wee1, and the counteracting phosphatase Cdc25 (Gautier et al., 1991; Gould and
Nurse, 1989; Kumagai and Dunphy, 1991; Russell and Nurse, 1986, 1987; Strausfeld et
al., 1991). In fission yeast, mutations in Wee1, Cdc25, and their upstream regulators lead
to changes in cell size. Separate mechanisms link cell size with regulation of Wee1 versus
Cdc25. The cellular concentration of Cdc25 increases as cells grow during interphase
(Keifenheim et al., 2017; Moreno et al., 1990). In contrast, the concentration of Wee1
remains constant during interphase, but it is progressively phosphorylated by the
conserved inhibitory kinases Cdr1 and Cdr2 (Aligue et al., 1997; Breeding et al., 1998;
Kanoh and Russell, 1998; Keifenheim et al., 2017; Lucena et al., 2017; Opalko and
Moseley, 2017; Russell and Nurse, 1987; Wu and Russell, 1993; Young and Fantes,
1987). cdr2 mutants fail to divide at a constant surface area, and instead divide according
to cell volume or length (Facchetti et al., 2019). This change suggests that Cdr2-Cdr1-
Wee1 signaling underlies the primary size-sensing pathway that measures cell surface
area, while additional pathways related to volume and length are engaged in its absence.
The localization of Cdr2, Cdr1, and Wee1 support this model **(Figure 1A)**: Cdr2 forms
punctate oligomeric structures called nodes that stably bind to the medial cell cortex, and
recruits Cdr1 to these sites (Akamatsu et al., 2014, 2017; Guzmán-Vendrell et al., 2015;
Martin and Berthelot-Grosjean, 2009; Morrell et al., 2004; Moseley et al., 2009). Wee1
localizes primarily in the nucleus and spindle-pole body, where it encounters Cdk1 to
prevent mitotic entry (Masuda et al., 2011; Moseley et al., 2009; Wu et al., 1996). In
addition, Wee1 transiently visits cortical Cdr1/2 nodes in bursts that lead to inhibition of
Wee1 kinase activity (Allard et al., 2018). The frequency and duration of these Wee1
bursts at Cdr1/2 nodes increase approximately twenty-fold as cells double in size (Allard
et al., 2018), leading to size-dependent inhibition of Wee1 at the cell surface.
85

This size-dependent change in Wee1 bursting dynamics is encoded into the Cdr1-
Cdr2-Wee1 pathway at two proposed levels. First, increased Wee1 bursting depends
upon the doubling in number of Cdr1/2 nodes during one cell cycle, but this 2-fold increase
is smaller than the 20-fold increase in Wee1 bursting (Allard et al., 2018; Deng and
Moseley, 2013; Pan et al., 2014). Second, activation of Cdr2 increases as cells increase
in size (Deng et al., 2014). Cdr2 kinase activity is required for Wee1 localization to nodes,
and is controlled by the upstream kinase Pom1 (Allard et al., 2018; Moseley et al., 2009).
Pom1 directly phosphorylates and inhibits the activation of Cdr2 (Bhatia et al., 2014; Deng
et al., 2014; Kettenbach et al., 2015; Martin and Berthelot-Grosjean, 2009; Moseley et al.,
2009). Pom1 also phosphorylates a separate set of sites on Cdr2 to disrupt
oligomerization and membrane binding (Bhatia et al., 2014; Rincon et al., 2014). Pom1
regulation of Cdr2 increases the frequency and duration of Wee1 bursts at Cdr1/2 nodes,
but only in small cells (Allard et al., 2018). Consistent with this defect, *pom1Δ* cells divide
at a small size due to dysregulation of the Cdr2-Cdr1-Wee1 pathway (Bähler and Pringle,
1998; Bhatia et al., 2014; Martin and Berthelot-Grosjean, 2009; Moseley et al., 2009;
Wood and Nurse, 2013). Taken together, these results suggest a size-dependent
interaction between Pom1 and its substrate Cdr2. However, the location and timing of
Pom1 interactions with Cdr2 have remained poorly defined.
104
Pom1 localizes in a cortical spatial gradient that is enriched at cell tips, with a lower
concentration at the medial cell cortex, where its substrate Cdr2 forms nodes (Bähler and
Pringle, 1998; Bhatia et al., 2014; Martin and Berthelot-Grosjean, 2009; Moseley et al.,
2009; Pan et al., 2014). Thus, the majority of Pom1 and Cdr2 molecules in a cell are
spatially separated **(Figure 1A)**, raising the question of when and where they interact.
Several lines of evidence suggest that the spatial distributions of Pom1 and Cdr2 are
critical for the size-dependent signaling properties of this pathway, and suggest the lateral
cell cortex as the key interface. For example, ectopic targeting of Pom1 to the medial cell
cortex inhibits Cdr2 node formation and Cdr2-dependent cell size signaling (Martin and
Berthelot-Grosjean, 2009). Additionally, the Pom1 gradient is disrupted as part of a
cellular response to glucose deprivation, which leads to increased Pom1 concentration at
the lateral cortex where it delays mitotic entry (Kelkar and Martin, 2015). This result also

demonstrates that Pom1-Cdr2 signaling responds to environmental input to coordinate
cell size with nutrient availability. In this system, Pom1 functions analogously to an
intracellular morphogen, acting as a concentration-dependent and localization-controlled
inhibitor of Cdr2 nodes. However, levels of Pom1 at the medial cell cortex in wild-type
cells growing under steady state conditions are low, constant, and do not vary with cell
size (Bhatia et al., 2014; Pan et al., 2014). How Pom1 provides size-dependent input to
Cdr2 is unclear and requires analysis of their molecular dynamics at the lateral cell cortex.
124
Past studies have led to a model for Pom1 gradient formation driven by the binding
and diffusion of individual Pom1 molecules in the plasma membrane (Hachet et al., 2011;
Hersch et al., 2015; Saunders et al., 2012). Intriguingly, dynamic clusters of Pom1 have
also been observed and are proposed to form by unstable oligomerization of molecules
diffusing on the membrane (Hachet et al., 2011; Saunders et al., 2012). These clusters
are thought to form and decay rapidly, temporarily slowing Pom1 diffusion rates and
reducing noise in the gradient (Saunders et al., 2012). These clusters represent nano-
scale pockets of increased Pom1 concentration, but whether these clusters contribute to
regulation of Cdr2 nodes is unknown. Here, we combine biochemical and microscopy
approaches to show that the Pom1 gradient is comprised of these punctate clusters.
Pom1 clusters are stable oligomers that bind and release the membrane with minimal
lateral diffusion. These clusters bind more frequently at cell tips than at cell sides, resulting
in a concentration gradient. A portion of Pom1 clusters in the medial cell cortex colocalize
with Cdr2 nodes, and the ratio of Pom1 to Cdr2 at the medial cell cortex changes as a
function of cell size. When glucose is restricted, Pom1 clusters redistribute to the medial
cell cortex to prevent dramatic dysregulation of Wee1. Our work reveals that the Pom1-
Cdr2-Wee1 signaling pathway is organized as a series of cortical clusters. The relative
distribution of these clusters at the plasma membrane changes with cell size and glucose
availability, thus relaying cell surface area to the core cell cycle machinery in a nutrient-
controlled manner.
145
146

Results:

The Pom1 gradient is formed by transient cortical clusters

We sought to examine the molecular dynamics of Pom1-Cdr2 signaling at the cell cortex. As a starting point, we imaged Pom1 by TIRF microscopy, which selectively excites fluorophores near the coverslip. Pom1 was tagged at the endogenous locus with the bright and photostable fluorophore mNeonGreen (mNG). Surprisingly, along the lateral cortex of interphase cells, Pom1-mNG was localized almost exclusively in discreet clusters, with no apparent diffuse signal **(Figure 1B)**. In images from a single time-point, Pom1 clusters formed a noisy concentration gradient **(Figure 1B,D)**. Time-averaging produced a smoother gradient dotted by occasional clusters **(Figure 1B,D)**, similar to the gradient observed by confocal microscopy **(Figure 1— figure supplement 1A,B)**. Pom1 clusters were highly dynamic **(Video 1)**. They appeared on the cell sides with a frequency of ~2 $\mu m^{-2}s^{-1}$, and remained bound for less than 2 seconds on average **(Figure 1E,F)**. During this brief cortical attachment, Pom1 clusters exhibited minimal diffusion which lacked directionality, and instead appeared to diffuse randomly in submicron patches of cortex **(Figure 1—figure supplement 2A,B,E)**. Mean squared displacement (MSD) measurements were consistent with free diffusion of Pom1 clusters within submicron corrals **(Figure 1—figure supplement 2A,B,C,D)** (Qian et al., 1991). The measured diffusion coefficient (D=0.134±0.017μm^2/s) was slow for a membrane-associated protein (Knight et al., 2010; Weiß et al., 2013), likely reflecting the large size and/or multiple membrane contacts for oligomeric Pom1 clusters.

We next used "head-on" TIRF microscopy to image Pom1 at cell tips. Similar to cell sides, Pom1-mNG localized at cell tips almost entirely in clusters **(Figure 1C)**. We also used "head-on" confocal imaging to confirm that Pom1 localization in clusters at tips was not an artifact of the TIRF approach **(Figure 1—figure supplement 1C)**; a conclusion further supported by work using high resolution wide-field microscopy (Dodgson et al., 2013). Compared to cell sides, Pom1 clusters appeared twice as frequently at cell tips but with a similar cortical duration, again exhibiting minimal and non-directional diffusion away from their binding site **(Figure 1E,F, Figure 1—figure supplement 2E)**. Fluorescence intensity of individual clusters at cell tips and cell sides

were similar, although we detected a low number of brighter clusters at cell tips, likely
representing multiple diffraction-limited clusters **(Figure 1—figure supplement 1C,D,E)**.
From these data, we conclude that the Pom1 gradient is formed by patterning the
membrane binding frequency of cortical clusters along the long axis of the cell, rather
than by diffusion of Pom1 molecules from the cell tips to the cell side.

***In vitro* analysis shows Pom1 exists in large, stable clusters**

Past work suggested that Pom1 structures assemble and disassemble at the
cortex through oligomerization of individual molecules diffusing in the plane of the
membrane (Saunders et al., 2012). However, our high-speed (20ms/frame) continuous
TIRF videos suggested that Pom1 clusters bind and release the cortex as a pre-formed
unit **(Figure 2—figure supplement 1A)**. Consistent with this possibility, in the rare cases
when we observed cell lysis events during TIRF imaging, Pom1 clusters remained intact
in extruded cytoplasm **(Figure 2A)**. This apparent stability while removed from the plasma
membrane led us to test the existence and properties of Pom1 clusters in detergent cell
extracts. Using TIRF microscopy, we observed clusters of Pom1-mNG in extracts made
from Pom1-mNG cells, but not from untagged wild-type cells **(Figure 2B)**. We next
examined the size of these Pom1 clusters by velocity sucrose gradient sedimentation. By
fractionating detergent cell extracts from *pom1-3xHA* cells, we found that most Pom1
exists in a large 60S complex, consistent with Pom1 clusters observed in cells. Fractions
6-8 containing this peak were dialyzed and centrifuged on a second sucrose gradient.
These pooled fractions again sedimented at 60S, indicating that they represent a stable
complex **(Figure 2C, Figure 2—figure supplement 2A,B,C)**.

Next, we tested the behavior of isolated Pom1 clusters on fluid artificial supported
lipid bilayers (SLBs) **(Figure 2D, Figure 2—figure supplement 1B)**. In this cell free-
system, Pom1-mNG clusters bound and released SLB lipids with strikingly similar
dynamics as in cells **(Figure 2E,F, Video 2)**. MSD measurements for Pom1 clusters on
SLBs revealed a diffusion coefficient similar to that measured in cells
(D_{cell}=0.134±0.017µm^2/s vs D_{SLB}= 0.168±0.018µm^2/s), but the relationship was linear over
the lifetime of most particles **(Figure 2—figure supplement 3A,B,C,D)**. These results

indicate that whereas membrane diffusion of Pom1 clusters in cells is confined within
microdomains, clusters can diffuse without confinement on synthetic bilayers. Dwell times
were increased for the kinase-dead mutant Pom1(K728R)-mNG, which was previously
shown to increase cortical Pom1 levels in cells **(Figure 2—figure supplement 1C).**
(Bähler and Pringle, 1998; Hachet et al., 2011). Pom1 clusters did not colocalize with
Cdr2 nodes in detergent extracts, suggesting they remain as separate structures **(Figure**
2 – figure supplement 1D). Collectively, our TIRF microscopy and *in vitro* analysis of
Pom1 clusters support a model whereby the majority of cellular Pom1 protein is contained
within discreet, highly-stable oligomers that bind and release membranes with kinetics
dictated by their catalytic activity in a pattern that generates a concentration gradient.
219

Formation of clusters could represent an intrinsic property of Pom1 protein, or
alternatively might require additional cellular factors. To distinguish between these
possibilities, we expressed and purified GST-Pom1 from bacteria, and then performed
sucrose gradient centrifugation experiments. Recombinant GST-Pom1 sedimented in a
low molecular weight peak unlike Pom1 clusters from cells **(Figure 2G)**. Remarkably,
purified GST-Pom1 was assembled into a cluster-sized high molecular weight complex
by incubation with wild-type yeast detergent extract **(Figure 2G)**. Complex formation was
also induced by incubation of GST-Pom1 with *pom1Δ* cell extracts, meaning that
additional cellular factors drive this assembly process **(Figure 2G).** Once assembled,
Pom1 clusters are stable oligomeric complexes with intrinsic membrane-binding
properties that can be reconstituted *in vitro*. Additional cellular regulatory proteins are
likely to promote the binding of Pom1 clusters at cell tips, leading to the spatial gradient.
232

Polarity landmarks pattern cortical dynamics of Pom1 clusters to shape the
gradient
The key role of clusters in forming the Pom1 gradient led us to reexamine the role
of cell polarity landmark proteins Tea1 and Tea4. Both of these proteins localize at cell
tips and are required for proper Pom1 gradient formation (Hachet et al., 2011). However,
past studies have shown that Tea1 and Tea4 have distinct mechanistic roles: Tea4 is
required for localization of Pom1 to the cortex, whereas Tea1 is required for enrichment

of cortical Pom1 to the cell tip (Hachet et al., 2011). We first used sucrose gradient
centrifugation to test if either protein is required for assembly of Pom1 into stable
biochemical complexes. The sedimentation pattern of Pom1 clusters isolated from *tea1Δ*
and *tea4Δ* cells in velocity sucrose gradients is unchanged from wild-type cells,
suggesting that neither Tea1 nor Tea4 are required for assembly of Pom1 into stable
biochemical complexes **(Figure 3A)**. Rather, these regulators are likely to act
downstream of cluster assembly.

We next tested the localization and dynamics of Pom1-mNG in *tea1Δ* and *tea4Δ*
cells. We confirmed that Pom1 was absent from the cell cortex in confocal micrographs
of *tea4Δ* cells **(Figure 3—figure supplement 1A,** (Hachet et al., 2011)**)**. Surprisingly, we
did observe Pom1-mNG clusters throughout the cortex of *tea4Δ* cells by TIRF microscopy
(Figure 3B). These binding events occurred with low frequency and short dwell times
(Figure 3C,D, Figure 3—figure supplement 1B,C,D), which likely preclude their
detection by confocal microscopy. In contrast, confocal micrographs of Pom1-mNG in
tea1Δ cells confirmed even distribution around the entire cell cortex **(Figure 3—figure**
supplement 1A, (Hachet et al., 2011)**)**. In TIRF microscopy of *tea1Δ* cells, Pom1-mNG
clusters bound the cell cortex with the same frequency at cell sides and cell tips **(Figure**
3B,C,D, Figure 3—figure supplement 1B,C,D) resulting in enrichment of cortical Pom1
but not a gradient. Thus, Tea1 and Tea4 cooperate to promote localized membrane
binding but not assembly of Pom1 clusters. These spatial cues pattern the frequency and
binding duration of Pom1 cluster cortical interaction along the long axis of the cell to
generate a concentration gradient that emanates from cell tips.

Pom1 clusters interact with Cdr2 nodes at the lateral cortex
It has been unclear when and where Pom1 interacts with its inhibitory target Cdr2,
which localizes in cortical nodes positioned in the cell middle. Both Pom1 and Cdr2 are
enriched at the cortex, and their cytoplasmic concentrations do not change with cell size
(Figure 4—figure supplement 1A,B,C,D,E). Furthermore, Cdr2 nodes and Pom1
clusters do not colocalize in cytoplasmic extracts, suggesting the cytoplasm is not the
location of their interaction **(Figure 2—figure supplement 1D)**. Since Pom1 clusters

appear to bind throughout the medial cell cortex, we used simultaneous two-color TIRF microscopy to test colocalization of Pom1 clusters and Cdr2 nodes in this region of overlap. We observed colocalization between Cdr2 nodes and some Pom1 clusters, as well as Pom1 clusters that bound to the cortex without encountering a Cdr2 node **(Figure 4A)**. Thus, Pom1 clusters bind to the medial cortex in an apparently stochastic pattern that can generate overlap with Cdr2 nodes. These patterns of colocalization were apparent in both static images and time-lapse movies **(Figure 4A,B)**. Similar results were obtained if different fluorophores were used to tag Pom1 and Cdr2, indicating that colocalization is unlikely an artifact of the fluorescent tags **(Figure 4—figure supplement 2A,B)**. The frequency, dwell time, and displacement of Pom1 cortical clusters were unaffected in *cdr2Δ* cells **(Figure 4—figure supplement 3A,B,C)**. These data suggest that Pom1 interacts with its substrate Cdr2 when Pom1 clusters associate with Cdr2 nodes at the medial cell cortex.

Phosphorylation by Pom1 prevents activation of Cdr2 in small cells, thereby contributing to cell size-dependent regulation of Wee1 and mitotic entry (Allard et al., 2018). To place this genetic pathway in the cellular context of Pom1 clusters and Cdr2 nodes, we analyzed how these structures accumulate and colocalize as cells grow. Intriguingly, the total number of both Cdr2 nodes and Pom1 clusters detected along entire cell sides in TIRF images were equivalent for cells of a given size **(Figure 4C,D)**, and both structures show a similar size-dependent doubling. We next restricted our analysis to a 2μm x 2μm square ROI positioned at the cell middle, where Cdr2 nodes concentrate. The local density of Cdr2 nodes in this region increased as a function of cell size, consistent with past studies (Deng and Moseley, 2013; Pan et al., 2014). In contrast, the density of Pom1 clusters in this region was largely independent of cell size **(Figure 4E)**. In both TIRF and confocal images, the concentration of Pom1 protein and Pom1 clusters decreased slightly as cells increase in size, but this trend was dwarfed by the cell size-dependent increase in Cdr2 node density **(Figure 4E, Figure 4—figure supplement 3D)**. Thus, as cells grow larger, the ratio of Pom1 to Cdr2 in the medial cell cortex changes to favor Cdr2 because the densities of these two structures scale differently with cell size **(Figure 4G)**.

We next tested how this density scaling affects colocalization of Pom1 clusters and
Cdr2 nodes as cells grow. Due to the increasing density of Cdr2 nodes, we observed an
increased number of colocalized Pom1 clusters and Cdr2 nodes at cell sides as cells
increase in size **(Figure 4F, Figure 4—figure supplement 3E)**. The increased
colocalization, combined with the constant density of Pom1 clusters, means that there
are less "free" Pom1 clusters as cells increase in size. Perhaps more importantly, the
number of "free" Cdr2 nodes increased as a function of cell size **(Figure 4—figure**
supplement 3F). These free Cdr2 nodes are not occupied by an inhibitory Pom1 cluster,
and thus have increased potential to promote mitotic entry by inhibiting Wee1. In this
manner, Pom1 sets an inhibitory threshold that must be overcome by an increase in Cdr2
node density. The inhibitory threshold decreases slightly as cells grow, and functions
most effectively in small cells, where Pom1 was previously shown to inhibit downstream
signaling to Wee1 (Allard et al., 2018).

Control of Pom1 cluster levels in the medial cell cortex
Our results suggest that Pom1 sets an inhibitory threshold that opposes Cdr2
activation until cells reach an appropriate size for division. We hypothesized that the
strength of the Pom1 inhibitory threshold could be tuned by altering the concentration of
Pom1 cortical clusters in the cell middle. To increase the abundance of Pom1 clusters in
the cell middle, we grew Pom1-mNG cells in low glucose media. In low glucose
conditions, the fission yeast microtubule cytoskeleton depolymerizes, which leads to
Pom1 redistribution to the lateral cell cortex where it delays mitotic entry (Kelkar and
Martin, 2015). In time-lapse TIRF microscopy experiments, we observed increased
numbers of discreet Pom1 clusters at the lateral cell cortex under low glucose conditions
(Figure 5A). These clusters bound to the lateral cortex more frequently than under normal
glucose conditions, but the average dwell time was unaffected **(Figure 5B,C)**. This
increased on-rate leads to accumulation of Pom1 clusters at the medial cell cortex.

We tested how increased numbers of Pom1 clusters at the lateral cell cortex affect
Cdr2 nodes using TIRF microscopy. In low glucose conditions, the number of Cdr2 nodes

measured per cell was largely unchanged. However, we observed a significant reduction in the number of Cdr2 molecules per node, and in the overall concentration of Cdr2 at the medial cortex **(Figure 5D,E,F, Figure 5—figure supplement 1A,B,C)**. Low glucose did not affect Cdr2 nodes in *pom1Δ* cells, so these effects are Pom1-dependent **(Figure 5D,E,F)**. Thus, Pom1 redistribution to the lateral cortex under low glucose is required to induce partial disassembly of each Cdr2 node. We quantified how this redistribution effect scales with glucose levels. As glucose was reduced, Pom1 concentration decreased at cell tips and increased at cell sides **(Figure 6A,C,D)**. Pom1 concentration at the lateral cortex therefore serves as a cellular indicator of glucose availability. In contrast, Cdr2 concentration increased at cell tips and decreased at cell sides **(Figure 6B,C,D)**. The concentrations of Pom1 and Cdr2 at cell sides were nearly perfectly anti-correlated (Pearson's r=-0.9352, *p*=0.0006) **(Figure 6E,F)**. We conclude that the increasing frequency of Pom1 cluster binding at the lateral cell cortex leads to a reduced number of Cdr2 molecules per node.

Pom1 disrupts Wee1 bursting under glucose deprivation

We next examined the downstream effects of altered Pom1-Cdr2 signaling in low glucose by using TIRF microscopy to monitor the previously described bursts of Wee1-mNG localization at Cdr2 nodes (Allard et al., 2018). In wild-type cells, both the frequency and duration of Wee1 bursts increased linearly with cell size, consistent with increased inhibition of Wee1 as cells grow **(Figure 7A,B, Supplemental Table 1, Supplemental Table 2)**. In low glucose, the frequency of Wee1-mNG bursts still scaled with cell size as in normal glucose. However, the duration of each burst was uniformly short and independent of cell size **(Figure 7A,B, Supplemental Table 1, Supplemental Table 2)**. Therefore, glucose modulates how long Wee1 molecules stay at inhibitory Cdr2 nodes. These same properties were seen for Wee1-mNG bursts in kinase-dead *cdr2(E177A)* mutant cells grown in normal glucose **(Figure 7—figure supplement 1A,B, Supplemental Table 1, Supplemental Table 2)**. These results indicate that Cdr2 kinase activity is required to retain Wee1 at nodes, but not for the initial binding event. Further, they demonstrate that low glucose phenocopies the kinase-dead *cdr2(E177A)* mutant, consistent with Pom1-dependent inhibition of Cdr2 kinase activity.

These results suggested that reduced Wee1 bursting in low glucose might require

Pom1 redistribution. We tested this idea by examining the glucose dependency of Wee1

localization to nodes in *pom1Δ* cells. In *pom1Δ* cells growing under high glucose, the

Wee1 burst frequency scales with cell size but Wee1 burst duration is uniformly high and

independent of cell size **(Figure 7A,C, Supplemental Table 1, Supplemental Table 2)**.

Unlike wild type cells, which suppress Wee1 bursts in low glucose, *pom1Δ* cells show a

striking increase in Wee1 bursts under low glucose. This increase is not caused by burst

duration, which was independent of both cell size and glucose concentration in *pom1Δ*

cells **(Figure 7C, Supplemental Table 1, Supplemental Table 2)**. Rather, we measured

increased Wee1-mNG burst frequency in low glucose *pom1Δ* cells **(Figure 7A,C,**

Supplemental Table 1, Supplemental Table 2). The reason for this increase is

unknown, and suggests the existence of additional glucose-regulated pathways that

operate on Cdr2-Wee1 signaling.

Wee1 localization at Cdr2 nodes leads to its inhibitory phosphorylation by Cdr2

and the related kinase Cdr1. To test how our microscopy results connect with Wee1

phosphorylation status, we analyzed phosphorylation-dependent shifts of Wee1 migration

using western blots **(Figure 7D)**. In high glucose, Wee1 migrates as a smear of

phosphorylated isoforms (Allard et al., 2018; Lucena et al., 2017). The upper,

hyperphosphorylated forms are increased in *pom1Δ* cells but absent in *cdr2Δ* cells. This

hyperphosphorylated Wee1 is lost in wild-type cells grown in low glucose, consistent with

reduced bursts of Wee1 localization to nodes. In contrast, Wee1 appears even more

hyperphosphorylated in *pom1Δ* cells grown under low glucose. This result suggests that

Pom1 prevents phospho-inactivation of Wee1 in response to low glucose. These

combined experiments support a model where Pom1 redistribution to the lateral cell

cortex inactivates Cdr2 nodes to relieve inhibition of Wee1 under low glucose conditions.

Discussion:

In this study, we have shown that the Pom1 concentration gradient is generated

by oligomeric Pom1 clusters that rapidly bind and release from the plasma membrane.

These clusters bind more frequently at cell tips versus cell sides in high glucose media,
while their binding rate is increased at cell sides in low glucose media. At the medial cell
cortex, these clusters overlap with their inhibitory target Cdr2, which localizes in static
oligomeric nodes. More Cdr2 nodes are free from Pom1 inhibition as cells grow larger
due to different density scaling of these two structures at the medial cell cortex. Our TIRF-
based colocalization experiments on these structures were limited to short periods of time
to avoid photobleaching, partly caused by the imaging conditions needed to observe
these highly dynamic Pom1 clusters. Nevertheless, Pom1 clusters colocalized with Cdr2
nodes even over these short timescales. Given the high frequency of Pom1 cluster
binding, we expect that many more colocalization events occur between Cdr2 nodes and
Pom1 clusters during the course of a full cell cycle, with the potential for Pom1 clusters
to visit each Cdr2 node multiple times. Thus, these rapid dynamics of Pom1 clusters
interacting with Cdr2 nodes may be integrated in time throughout the cell cycle. It appears
likely that the most critical temporal window for Pom1-Cdr2 interactions occurs during
early G2, as *pom1Δ* show defects in Cdr2-Wee1 signaling specifically in small cells (Allard
et al., 2018). It will be interesting to determine how these rapid interactions in small cells
are integrated into the mitotic entry decision, which occurs later in the cell cycle when
cells are longer.
413
Molecular clusters are widespread in signal transduction
The Pom1-Cdr2-Wee1 signaling pathway appears to function entirely within the
confines of oligomeric protein clusters at the plasma membrane. The development of
imaging technologies with increased signal-to-noise and spatial and temporal resolution
has enabled the discovery of protein clustering as a paradigm in signaling. As a result, a
growing number of signaling events and proteins have been found in similar clustered
structures that are referred to as nanodomains, clusters, nanoclusters, and nodes, among
other names. Such structures have been shown to operate in polarization of the *C.*
elegans zygote, signal transduction nanodomains at the plant plasma membrane, Ras
isoform-specific signaling, bacterial chemotactic receptors, and both B-cell and T-cell
receptors (Bray et al., 1998; Dickinson et al., 2017; Douglass and Vale, 2005; Duke and
Bray, 1999; Falke, 2002; Gronnier et al., 2017; Liu et al., 2016; Maity et al., 2015; Munro,

2017; Rodriguez et al., 2017; Wang et al., 2017; Zhou and Hancock, 2015). These T cell

receptor clusters enrich downstream kinases while excluding phosphatases (Su et al.,

2016), a mechanism that could relate to concentrated clusters of the kinases Pom1 and

Cdr2. In addition, theoretical work has suggested that clustering can reduce noise by

generating reaction bursts (Kalay et al., 2012), which have the capacity to overcome

inhibitory thresholds, such as those imparted by phosphatases, more effectively than a

system driven by gradual accumulation of signal. Insight into the functional role of

clustering for Pom1-Cdr2-Wee1 will require additional *in vivo* analysis combined with *in*

vitro reconstitution. We have shown that Pom1 clusters remain stable through

biochemical enrichment, and isolated clusters exhibit the same membrane-binding

properties *in vitro* as in cells. Past work has demonstrated Cdr2 nodes also remain stable

through biochemical fractionation (Allard et al., 2018), providing the necessary tools for

future dissection of the pathway in a reconstituted system.

A new model for cortical gradient formation

Our work has shown that Pom1 clusters bind to the cell cortex with different

frequencies along the long axis of the cell, resulting in a concentration gradient. Past work

suggested single Pom1 molecules (or small oligomers) bind to the membrane at tips, and

then diffuse away from the tip before dissociating (Hachet et al., 2011; Saunders et al.,

2012). Clusters were proposed to dynamically assemble and disassemble at the cortex

from this diffusing population (Saunders et al., 2012). Our findings suggest a new model

for gradient formation, which is distinct from this previous model in four ways. First, we

found that clusters themselves form a concentration gradient that emanates from the cell

tips. Second, our combined imaging and biochemical approaches demonstrated that

clusters are stable complexes that bind and rapidly release from the membrane. Third,

we did not observe extensive diffusion by complexes bound to the membrane. It remains

possible that smaller, diffusing Pom1 complexes or single molecules were not detected

by our imaging approach, but we favor a model where diffusion is limited to short, non-

directional movements that do not contribute to gradient formation. Fourth, the key step

in gradient formation is a positional system that increases the on-rate for Pom1 clusters

at cell tips versus the cell middle. Concentration gradients are found across vast size

scales in biology, so the mechanism of gradient formation by Pom1 clusters may have
broad implications. It is also important to note that the spatial gradient of on-rates for
Pom1 clusters binding to the cortex may depend on a concentration gradient of specific
lipids in the plasma membrane. In support of this hypothesis, Pom1 binds to
phosphatidylserine *in vitro* (Hachet et al., 2011), and this lipid is enriched at the tips of
growing fission yeast cells (Haupt and Minc, 2017). Thus, further investigation of these
lipids and their localization dynamics could reveal additional layers of this morphogen-like
gradient.

We found that the polarity landmarks Tea1 and Tea4 provide the positional
information for Pom1 clusters to form a concentration gradient. Past work has shown that
Pom1 localizes homogenously throughout the cortex in *tea1Δ* cells, and does not localize
to the cortex in *tea4Δ* cells (Bähler and Pringle, 1998; Hachet et al., 2011). Based on
these and other results, Tea1 was proposed to recruit Tea4 to cell tips, where it promotes
cortical loading of Pom1 molecules. Several results suggest that additional mechanisms
may contribute to gradient formation. For example, we found that Pom1 clusters still bind
to the cortex in *tea4Δ* cells but for very short durations. Past work has also shown that
Tea1 and Tea4 form cortical clusters that co-localize at cell tips, but these Tea1-Tea4
clusters do not co-localize with Pom1 clusters (Dodgson et al., 2013). Thus, Tea1-Tea4
may pattern binding of Pom1 clusters to the cortex through an additional mechanism. For
example, Tea1 and Tea4 have been shown to assemble sterol-enriched lipid domains at
cell tips when cells initiate polarization following exit from starvation (Makushok et al.,
2016). Sterols and phosphatidylserine are thought to form a common membrane domain
in fission yeast, suggesting that these proteins could act through the recruitment and
organization of lipids to generate cortical domains permissive for Pom1 cluster binding
(Haupt and Minc, 2017; Makushok et al., 2016).

Adjustable Pom1 gradients regulate mitotic entry
We propose an integrated mechanism for fission yeast cell size sensing in which
Cdr2 acts as an important sensor of cell size that is dynamically antagonized by cortical
Pom1 **(Figure 8)**. Pom1 clusters bind to the medial cell cortex in a manner that can

overlap with Cdr2 nodes. These dynamic Pom1 clusters and static Cdr2 cortical nodes suggest a system that monitors surface expansion, consistent with the notion that fission yeast cells monitor surface area as the primary determinant of size at division (Facchetti et al., 2019; Pan et al., 2014). As cells grow, the density of Pom1 clusters in the cell middle decreases slightly due to their binding preference at cell tips. In contrast, the density of Cdr2 nodes doubles to overcome this inhibitory Pom1 threshold. The density of Pom1 clusters and the resulting inhibitory threshold can be modulated by glucose availability, consistent with past work (Kelkar and Martin, 2015). As fission yeast cells begin to starve due to depletion of nutrients such as glucose, their length at division progressively decreases and the length of the cell cycle increases (Pluskal et al., 2011; Yanagida Mitsuhiro et al., 2011). The Pom-Cdr2 network has a prominent role in this nutrient modulation, suggesting that it may represent a key physiological role for the pathway (Kelkar and Martin, 2015). We found that the size dependence of Wee1 localization bursts to Cdr2 nodes, which facilitate Wee1 inhibition, was dampened in low glucose conditions. Re-localization of Pom1 clusters to the medial cortex mediates this response. However, for *pom1Δ* mutants in low glucose, Wee1 localization to nodes exceeds even wild type cells in high glucose. This result indicates that additional mechanisms contribute to Wee1 regulation in low glucose, for example Wee1 nuclear transport could be altered under these conditions. Thus, Pom1-Cdr2 signal transduction serves two functions: (1) Under steady-state growth conditions, Pom1 and Cdr2 cooperate to measure cell size and ensure that cells divide only when they have grown large enough. In this context, the Pom1 threshold is only strong enough to inhibit Cdr2 signaling in small cells, where node density is low. In this manner, Pom1 prevents precocious entry into mitosis. (2) Under glucose restriction, Pom1 and Cdr2 cooperate to integrate information about both cell size and glucose availability, and signal that information to the core cell cycle machinery.

The Pom1 gradient as a read-out for cell polarity

More broadly, our study reveals the adjustable nature of the concentration gradient formed by Pom1 clusters. Concentration of Pom1 clusters at cell tips reads out the polarity state of the cell for downstream signaling by Cdr2 and Wee1. Conditions that alter the

polarized growth state of the cell would lead to changes in the gradient of Pom1 clusters,
resulting in altered cell size at division. This dynamic system appears to operate in a
manner that depends on both cell size and nutrient availability. Identification of additional
growth and environmental conditions that alter the relative distributions of Pom1 clusters
and Cdr2 nodes, and determination of how they connect with other cell cycle signaling
pathways, may reveal new mechanisms for nutrient modulation of cell size control.

Integration of the Pom1-Cdr2-Wee1 pathway with other sizing mechanisms

Pom1-Cdr2-Wee1 signaling is cell size dependent but is not the only pathway that
contributes to the overall cell size control network. Intriguingly, a recent study reported
that *cdr2Δ* cells no longer divide based on accumulation of a threshold surface area and
instead revert to a secondary mechanism of cell size control based on volume sensing
(Facchetti et al., 2019). These results demonstrate that cells use different signal
transduction networks to measure different aspects of their size. Whereas the Pom1-Cdr2
network is well suited to measure cortical surface area, size-control systems that measure
other aspects of size would require different signaling logic. Theoretically, mechanisms
for cell volume sensing could depend on proteins whose concentrations do not scale
linearly with cell size. Such mechanisms are thought to underlie cell size control in other
organisms, such as the budding yeast, where growth-dependent dilution of the
transcriptional repressor Whi5 is thought to underlie G_1/S size control (Schmoller et al.,
2015). One possible candidate for cellular volume sensing in fission yeast is Cdc25, which
increases in concentration with cell size (Keifenheim et al., 2017; Moreno et al., 1990).
These mechanisms are attractive candidates for volume sensors because cytoplasmic
concentration is independent of cell geometry. The existence of sizing mechanisms that
measure other aspects of cell size, and function independently of the Pom1-Cdr2 surface
area sensing network, underscores that cell size is a systems level property controlled by
multiple signaling pathways. Thus, it remains important to discover size-dependent
signaling mechanisms within each individual pathway, as a step towards understanding
the integration of multiple pathways into a larger size control system.

Materials and Methods:

 Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
strain, strain background ()	*pom1-yomNeonGreen::hphR*	This Paper	JM4496	
strain, strain background ()	*pom1-3HA::hphR ura4-D18 leu1-32 h+*	PN948	JM797	Paul Nurse Lab
strain, strain background ()	*pom1Δ::natR h-*	Lab Stock	JM966	
strain, strain background ()	*pom1-m2-yomNeonGreen::hphR ura4-D18 leu1-32 h-*	This Paper	JM5412	
strain, strain background ()	*pom1-3HA::hphR tea1Δ::ura4+ ura4-D18*	This Paper	JM5414	
strain, strain background ()	*pom1-3HA::hphR tea4Δ::kanMX6*	This Paper	JM5415	
strain, strain background ()	*pom1-yomNeonGreen::hphR tea1Δ::kanMX6*	This Paper	JM4792	
strain, strain background ()	*pom1-yomNeonGreen::hphR tea4Δ::kanMX6*	This Paper	JM4791	
strain, strain background ()	*pom1-yomNeonGreen::hphR cdr2-tagRFP-t::hphR*	This Paper	JM5373	
strain, strain background ()	*cdr2-yomNeonGreen::hphR h-*	Lab Stock	JM4493	

Category	Description	Source	Identifier	Additional information
strain, strain background ()	*wee1-yomNeonGreen::hphR cdr2∆::natR*	Lab Stock	JM4525	
strain, strain background ()	*cdr2-yomNeonGreen::hphR pom1-tdTomato::natR*	This Study	JM5135	
strain, strain background ()	*cdr2-mEGFP::kanMX6 ura4-D18 leu1-32 ade6-m210 h+*	Lab Stock	JM346	
strain, strain background ()	*cdr2-yomNeonGreen::hphR pom1∆::natR*	This Paper	JM5359	
strain, strain background ()	*wee1-yomNeonGreen::hphR h-*	Lab Stock	JM4495	
strain, strain background ()	*wee1-yomNeonGreen::hphR pom1∆::kanMX6*	Lab Stock	JM4527	
strain, strain background ()	*972 h-*	PN1	JM366	Paul Nurse lab
strain, strain background ()	*cdr2∆::natR ura4-D18 leu1-32 h+*	Lab Stock	JM600	
strain, strain background ()	*wee1-yomNeonGreen::hphR cdr2(E177A)*	Lab Stock	JM4578	
recombinant DNA reagent	pFA6a-yomNeonGreen::hphR	Lab Stock	pJM1344	
recombinant DNA reagent	pFA6a-tdTomato::natR	Lab Stock	pJM248	
recombinant DNA reagent	pFA6a-yomTagRFP-T::hphR	Lab Stock	pJM1221	Derived from Addgene Plasmid 44842

recombinant DNA reagent	pFA6a-mEGFP::kanMX6	Lab Stock	pJM228	
recombinant DNA reagent	pFA6a-3HA::hphR	Lab Stock	pJM216	
antibody	anti-HA (mouse monoclonal)	Covance	MMS-101R	WB (1:3000)
antibody	anti-GST (rabbit polycolonal)	This Paper		WB (1:3000)
chemical compound, drug	1,2-dioleoyl-sn-glycero-3-phospho-L-serine (sodium salt)	Avanti Polar Lipids	840035C	18:1 DOPS or "PS"
chemical compound, drug	1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-N-(lissamine rhodamine B sulfonyl) (ammonium salt)	Avanti Polar Lipids	810150C	18:1 Liss Rhod PE
chemical compound, drug	1,2-dioleoyl-sn-glycero-3-phosphocholine	Avanti Polar Lipids	850375C	18:1 (Δ9-Cis) PC (DOPC) or "PC"

Strain Construction and Media

Standard *S. pombe* media and methods were used (Moreno et al., 1991). Strains used in this study are listed in the Key Resources Table. Gene tagging and deletion were performed using PCR and homologous recombination (Bähler et al., 1998). The mNeonGreen (mNG) sequence was used under license from Allele Biotechnology. Addgene Plasmid 44842 (pFA6a-link-yoTagRFP-T-SpHis5) was a gift from Wendell Lim & Kurt Thorn (Addgene plasmid # 44842; http://n2t.net/addgene:44842 ; RRID:Addgene_44842).

TIRF Microscopy and Analysis

Pom1 clusters and node components were imaged using simultaneous dual-color total internal reflection fluorescence (TIRF) microscopy to limit excitation of fluorophores

to those nearest to coverslip. Imaging was performed on a commercially available TIRF microscope (Micro Video Instruments) composed of a Nikon Eclipse Ti microscope base equipped with a 100x Nikon Apo TIRF NA 1.49 objective and a two-camera imaging adaptor (Tu-CAM, Andor Technlogy) containing a dichroic and polarization filters (Semrock FF580-FDi01-25x36, FF02-525/40-25, FF01-640/40-25) to split red and green signal between two aligned Andor iXon electron-multiplied CCD cameras (Andor Technology). Red/green beam alignment was performed prior to imaging using a TetraSpeck Fluorescent Microsphere size kit (Thermofisher).

Standard #1.5 glass coverslips were RCA cleaned before use to remove fluorescent debris. Cells were grown in EMM4S, and washed into fresh EMM4S immediately before imaging to remove auto-fluorescent debris resulting from overnight culture. Cells were imaged in EMM4S media on glass slides at ambient temperature. Individual slides were used for no more than five minutes to prevent cells from exhausting nutrients or oxygen. Agar pads were not used due to increased background fluorescence. Image analysis and processing was performed using ImageJ2 (NIH).

"Head-On" Tip Imaging

"Head-On" or "Tip" imaging was performed using a protocol modified from (Dodgson et al., 2013). Custom micro-well coverslips or Ibidi Sticky-Slide VI channels (Ibidi, Martinsried, Germany) were coated with BS-I lectin (1mg/mL in water) by incubation for 30 minutes and washed 3X with water to remove non-adherent lectin. Cells were then added to the imaging chambers and allowed to settle to the cover-slip bottom where they adhere to the lectin. Cells oriented with the long axis perpendicular to the coverslip were identified using brightfield prior to fluorescence imaging.

Spinning Disc Microscopy and Analysis

Spinning-disc confocal imaging was performed using a commercially available system (Micro Video Instruments, Avon, MA) featuring a Nikon Eclipse Ti base equipped with an Andor CSU-W1 two-camera spinning disc module, dual Zyla sCMOS cameras (Andor, South Windsor, CT) an Andor ILE laser module, and a Nikon 100X Plan Apo λ

1.45 oil immersion objective. Cells were imaged in EMM4S media on glass slides at
ambient temperature unless otherwise noted.

Airyscan Super-Resolution Microscopy and Analysis

To achieve maximum resolution and sensitivity, fluorescence intensity of Pom1
clusters on both cell sides and cell tips was measured using a Zeiss Airyscan microscope
(Figure 1—figure supplement 1C,D,E), composed of a Zeiss LSM-880 laser scanning
confocal microscope (Zeiss, Oberkochen, Germany) equipped with 100X alpha Plan-
Apochromat/NA 1.46 Oil DIC M27 Elyra objective, Airyscan super-resolution module and
GaAsP Detectors, and Zen Blue acquisition software using the Super-resolution mode
with pin-hole size of 1.2 airy-units to prioritize resolution. Z-volumes of 16 slices with
0.19µm spacing for high spatial resolution in all dimensions were centered on the cell
cortex closest to the coverslip. Airyscan images were processed in Zeiss Zen Blue
software, and quantification was performed on sum projections of Airyscan reconstructed
stacks.

Single Particle Tracking

Image analysis and processing was performed using ImageJ2 (NIH). Cdr2 node
numbers and Pom1/Wee1 clusters number, frequency and binding duration were
quantified using the Trackmate plugin (Tinevez et al., 2017) to analyze TIRF microscopy
movies. Due to variable fluorescence intensity in different TIRF fields and images,
thresholding parameters were determined separately for each image, and accuracy was
confirmed by visual inspection to ensure that only nodes/clusters were counted and that
no nodes/clusters were omitted. For Wee1 burst and Pom1 cluster tracking, Particle
Diameter was set to 0.3 microns (approximate XY resolution), with Maximum Gap Linking
set to 2 frames, and Linking Range for particle tracking was set to 0.15 microns. Lookup
table was adjusted to fire for some images to emphasize signal intensities. For Pom1
cluster tracking, analysis was restricted to 2µm x 2µm square ROIs positioned at the cell
middle or tip.

Colocalization Analysis

Colocalization analysis of Cdr2 nodes and Pom1 clusters as in **Figure 4F,G** and
Figure 4—figure supplement 2E,F was performed using two-color simultaneous TIRF
microscopy images as in **Figure 4A**. Analysis was restricted to $2\mu m^2$ square ROIs
positioned at the cell middle. The spot detection algorithm in the Trackmate plugin
(Tinevez et al., 2017) for ImageJ2 was used to assign 0.3 µm diameter circular ROIs to
Cdr2 nodes (cyan) and Pom1 clusters (magenta) in images as in **Figure 4F**. Pom1 and
Cdr2 structures were counted as colocalized if the centroids of their respective ROIs were
spaced <150nm apart (yellow ROI pairs in **Figure 4F**.
635

Mean Squared Displacement
Mean Squared Displacement analysis and calculation of diffusion coefficients were
performed using the MATLAB (MathWorks, Natick, MA) class @msdanalyzer developed
in (Tarantino et al., 2014). Briefly, single-particle traces of high-speed (100ms acquisition)
TIRFM videos were generated using the Trackmate plugin for ImageJ2 as described
above and were then exported to MATLAB for further analysis. @msdanalyzer was then
used to plot and calculate MSD for each particle trace, plot the MSD curves, compute and
fit the mean MSD curve, and compute the diffusion coefficient.
644

Pom1 Purification
The full length Pom1 sequence was purified as follows: Pom1 sequence was
subcloned into pGEX6P1 vector using the Xho1 restriction site and expressed as a GST-
fusion protein in *Escherichia coli* strain BL21(DE3). Transformants were cultured to log-
phase at 37˚C, followed by a shift to 25˚C for 30 minutes. Expression was then induced
by addition of 1-thio-β-D-galactopyranoside to 200µM, followed by growth for an
additional 3 hours at 25˚C. Cells were harvested by centrifugation and lysed by passing
them twice through a French press in lysis buffer (1xPBS, 100mM NaCl, 1mM DTT, and
complete EDTA-free protease inhibitor tablets (1 tablet/50mL Buffer) (Roche, Basel,
Switzerland)). Following lysis, Triton-X 100 was added to 1% V/V. Lysates were then
cleared by centrifugation for 10 minutes at 12,000 x g at 4˚C in a Sorval SS-34 fixed-angle
rotor (Thermo Scientific, Waltham, MA). The supernatant was then incubated with
glutathione-agarose (Sigma Aldrich, St. Louis, MO) for 2h at 4˚C. 20 mM glutathione (pH

8.0) was used to elute purified protein, and the eluate was dialyzed overnight at 4°C into

10mM Tris-HCl, pH8 + 5% V/V glycerol.

Preparation of Yeast Extracts

Fission yeast detergent extracts were prepared by growing 1.5L cells to mid-log

phase (OD ~0.3), and then washed twice with 50mL Node Isolation Buffer (NIB – 50mM

HEPES, 100mM KCl, 1mM $MgCl_2$, 1mM EDTA, pH 7.5) (Allard et al., 2018). Next, we

resuspended the pellet in an equal volume of 2X NIB (W/V) containing a

protease/phosphatase inhibitor cocktail (10µL/mL 200x PI, 50µL/mL 1M β-glycerol

phosphate, 50µL/mL 1M NaF, 2.5 µL/mL 200mM PMSF, 1mM DTT), and snap froze the

resuspension as pellets by pipetting drop-wise into a liquid nitrogen bath. Then, yeast

pellets were ground using liquid-nitrogen chilled coffee-grinders for 2 min, and collected

into chilled falcon tubes and stored at -80°C. 1.5g of frozen yeast powder was then thawed

on ice, and Triton X-100 was added to a final concentration of 1%, and the extracts were

mixed by gentle pipetting. Extracts were then centrifuged at 4°C for 10min at 20,000 x g

to yield a low speed supernatant, which were then used for subsequent experiments.

Sucrose Gradient Ultracentrifugation

Discontinuous sucrose gradients were prepared in 14 x 89mm Ultra Clear

Ultracentrifuge tubes (Beckman Coulter, Brea, CA) by layering 5 – 23% (top to bottom)

sucrose in NIB + 1% Triton X-100 in 0.37mL steps of 2% sucrose increments. 700µL of

yeast extract or recombinant protein diluted in NIB Buffer was then added to the top of

the gradient. Sucrose gradients were centrifuged at 100k x g (50kRPM) in a Beckman L8-

M ultracentrifuge for 2hr at 4°C in a chilled SW60ti swinging bucket rotor. 0.5mL gradient

fractions were collected from the top by hand, vortexed, and 100µL of each fraction was

mixed 2:1 in 3X SDS-PAGE sample buffer (65 mM Tris pH 6.8, 3% SDS, 10% glycerol,

10% 2-mercaptoethanol, 50 mM NaF, 50 mM β-glycerophosphate, 1 mM sodium

orthovanadate) and boiled for 5 min. Samples were then subjected to SDS-PAGE and

western blotting or Coomassie staining. To calculate S-values of sedimentation peaks

from western blot signal intensities, mean band intensities were measured using Image

Studio Lite (LICOR, Lincoln, NE), and Gaussian Curves were fit to these values. The peak

of the Gaussian for each sedimentation peak of each protein was used to assign the
known or interpolated S-value.

Western Blotting

For western blots, cells were lysed in 150 µl of 3X SDS-PAGE sample buffer with
glass beads in a Mini-beadbeater-16 (Biospec, Bartlesville, OK) for 2 minutes. Gels were
run at a constant 20 mAmps until 75kDa marker was at the bottom of the gel. Blots were
probed with anti-HA (Covance, Princeton, NJ) or homemade anti-GST. For monitoring
Wee1 phosphorylation, samples were run on an SDS-PAGE gel containing 6%
acrylamide and 0.02% bisacrylamide, and then probed with a homemade anti-Wee1
antibody (Allard et al., 2018).

Supported Lipid Bilayers

To prepare supported lipid bilayers, we first prepared small unilamellar vesicles
(SUVs) composed of three lipids: (1) DOPC (18:1 (Delta9) Cis PC - 1,2 Dioleoyl-sn-
glycero-3-phosphocholine), (2) DOPS (18:1 PS - 1,2-Dioleoyl-sn-Glycero-3-[Phospho-L-
Serine]), and (3) fluorescent 18:1 Liss Rhod PE (1,2-dioleoyl-sn-glycero-3-
phosphoethanolamine-N-(lissamine rhodamine B sulfonyl). All lipids were dissolved in
chloroform (Avanti, Alabaster, AL). 2µmol total lipid with the desired molar ratios were
mixed in glass vials with PTFE coated caps, which had been RCA cleaned and rinsed
three times with chloroform before use. Excess chloroform was evaporated in a vacuum
desiccator for 1 hour. Lipids were resuspended in 400µL SUV Buffer (50mM HEPES,
100mM KCl, 1mM EDTA, 1mM $MgCl_2$, pH7.5) for 5mM lipid mixture stocks. The lipid
mixtures were vortexed until cloudy, and then transferred to 1.5mL Eppendorf tubes and
subjected to ten freeze-thaw cycles using a liquid nitrogen bath and 32˚C hot plate, with
2 minutes of sonication in a sonicating water bath following each thaw cycle. Stocks were
stored at -80˚C in 20µL aliquots.

Supported lipid bilayers were made by adding 10µL of SUVs to RCA cleaned
custom microwells built on 22x40mm #1.5 coverslips. Chambers were then incubated at
37°C for one hour to induce vesicle fusion, and unincorporated vesicles were rinsed away

using five rinses with 100µL SUV buffer. Yeast extract was added to these chambers by
removing 90µL of buffer, leaving just enough to cover the chamber bottom, and then
adding extract.

Statistics and Line Fitting

All plotting and statistical analysis was performed using GraphPad Prism
8.0.2(263) (GraphPad Software, San Diego, CA), except for MSD analysis, which was
performed using MATLAB as described above. Analysis of statistical significance was
performed using unpaired Student's T-tests for comparison of two data sets where
appropriate or using single-factor ANOVA with Tukey's multiple comparison test for
comparison of >2 data sets. The specific test used for each experiment and the results
of that test are listed in the figure legends, with one exception: Analysis of the data
presented in **Figure 3C,D** is provided in an ANOVA table in **Figure 3—figure**
supplement 1.

In general, line fitting was performed using linear regression by least-squares with
analysis of residuals to ensure fit. Where linear regression did not fit the biological
question, we compared of multiple models (linear regression vs non-linear regression,
and the coefficient of determination (R^2) was considered to determine best-fit. When
considering the regression model in **Figure 6C,D**, visualization of the non-log scale plots
revealed unambiguous exponential relationships between glucose concentration and the
concentration of either Pom1 or Cdr2, reflecting saturable binding. In **Figure 7B,C**, insets
indicate whether the slopes (b_1) and Y-intercepts (b_0) of the two linear regressions are
significantly different at the 95% confidence level.   Details of the linear regressions
presented in **Figure 7B,C** and **Figure 7—figure supplement 1A,B** are provided in
Supplemental Table 1 and **Supplemental Table 2**. To summarize the results of a given
test, statistical significance is indicated within plots by the convention: (****) indicates $p <$
0.0001, (***) indicates $p < 0.001$, (**) indicates $p < 0.01$, (*) indicates $p < 0.05$, n.s.
indicated $p > 0.05$.

Figure Legends:

Figure 1: **The Pom1 gradient is formed by time-averaging of clusters that transiently bind the cortex. (A)** Schematic of the Cdr2-Pom1-Wee1 signal transduction pathway and its coarse cellular localization.   **(B)** Individual frame (top panel) and sum projection (bottom panel) of a high-speed TIRF microscopy movie. Movie was continuous 15s time-lapse acquisition of 200ms exposures. Scale bar 1µm. Blue arrows mark position of line scans performed for data in panel (D). **(C)** Pom1-mNG also forms clusters at the cell tip. Image is a sum projection of three consecutive 200ms exposures from continuous time-lapse TIRF movie of the cell tip as depicted in the cartoon diagram. Scale bar 1µm. **(D)** Line scans of fluorescence intensity along the long axis of the snap shot (blue line, left Y axis) and projection images (red line, right Y axis) in panel (B). Note that time-averaging of Pom1 clusters smoothens the concentration gradient. **(E)** Comparison of Pom1 cluster binding frequency at the cell tip or side (****, $p=<0.0001$, n=10 cells, 42-170 traces/cell). **(F)** Comparison of cortical dwell time of individual Pom1-mNG clusters (n.s., $p=0.6747$, $n=10$ cells, 42-170 traces/cell) at the cell tip or side. For (E-F), each data point represents a single cell mean, and line and error bars represent mean and standard deviation of all cells. Statistical significance was tested using a Student's T-test.

Figure 1—figure supplement 1: **Analysis of the Pom1 gradient by confocal microscopy**. **(A)** Pom1-mNG clusters are apparent at the lateral cell cortex of cells in individual frames (200ms) of spinning-disc microscopy movies (top panel) but become averaged out in a projection of the entire movie (bottom panel). Sum projection is of a continuous 15s time-lapse acquisition of 200ms exposures at the top cell focal plane. Scale bar 1µm. **(B)** Line scans of fluorescence intensity along the long axis of the cell in the snap shot (blue line, left Y axis) and projection image (red line, right Y axis) at positions marked by blue arrow heads in panel (A). Note that time-averaging of Pom1 clusters smoothens the concentration gradient. **(C)** Representative images of Airyscan super-resolution micrographs of Pom1-mNG at cell tip or side. Scale bar, 1µm. **(D)** Fluorescence intensities of all single clusters resolved in Airyscan images as in panel (C). Data points represent individual cluster values, error bars are the mean and standard

deviation (*n*=87 tip, 158 side). **(E)** Histogram of the fluorescence intensity of individual

Pom1 clusters at the cell tip and cell side. Clusters were binned by fluorescence intensity

and displayed as a population frequency (*n*=48 tip, 158 side).

<u>Figure 1—figure supplement 2</u>: **Analysis of Pom1 cluster diffusion (A)** Pom1-mNG

clusters positioned at the cell middle, in a 2x2µm ROI as used for particle tracking. **(B)**

Single-particle diffusion traces of Pom1-mNG clusters show limited lateral diffusion in

cells. **(C)** Mean squared displacement (MSD) curves calculated from traces of individual

Pom1-mNG cluster diffusion as in panel (B). **(D)** Mean MSD curves generated from

averaging of 439 single-particle traces, coming from 6 different cells. Linear regression

of the initial linear segment (red line) of the curve provides the diffusion coefficient

$D=0.134\pm0.017\mu m^2/s$. Black line represents the weighted mean of all traces, and grey

lines indicate standard deviation. **(E)** Average life-time displacement of individual clusters

(***, *p*=0.0002, *n*=10 cells, 42-170 traces/cell). Line and error represent mean and

standard deviation. Statistical significance was tested using a Student's T-test.

<u>Figure 2</u>: **Pom1 clusters are stable structures that can be isolated *in vitro*. (A)** TIRF

microscopy image of Pom1-mNG clusters in the extruded cytoplasm (white dotted line)

of a lysed cell (yellow dotted line). Scale bar 5µm. **(B)** TIRF microscopy images of cell

extracts prepared from wild-type (no tag) or Pom1-mNG cells. Images are 50 frame sum

projections of continuous 200ms time-lapse exposures. The two images were contrasted

equally. Scale bars 1µm. **(C)** Cytoplasmic extracts of *pom1-3HA* cells were subjected to

velocity sucrose gradient sedimentation, and fractions were probed against the HA tag

(upper blot). Fraction 1 corresponds to the top of the gradient and contains smaller

structures; fraction 9 corresponds to bottom of the gradient and contains larger structures.

Fractions 6-8 were pooled, sucrose was removed by dialysis, and then the sample was

subjected to a second identical round of sucrose gradient sedimentation and western

blotting of the resulting fractions (lower blot). **(D)** TIRF microscopy of Pom1-mNG clusters

from cytoplasmic extracts on supported lipid bilayers. Scale bar 1µm. Left panel is single

time point image. Right panel is kymograph taken from a line scan of time-lapse TIRF

experiment. **(E)** Quantification of binding duration of Pom1-mNG clusters on supported

lipid bilayers imaged by TIRF microscopy as in panel (D). Values are compared to cellular

measurements of Pom1 clusters on cell sides (n.s., *p*=0.05954, n= 713 *in vivo*, 421 *in*

vitro). **(F)** Quantification of life-time displacement of Pom1-mNG clusters diffusing on

supported lipid bilayers imaged by TIRF microscopy as in panel (D). Values are compared

to cellular measurements of Pom1 clusters on cell sides (****, *p*<0.0001, n= 713 *in vivo*,

421 *in vitro*). For (E) and (F), statistical significance was tested using a Student's T-test.

(G) Purified GST-Pom1 was subjected to sucrose gradient sedimentation and the

fractions were probed against the GST tag (upper blot). Purified GST-Pom1 was also

added to wild-type or *pom1Δ* cell extracts and incubated for 1hr at 4°C in the presence of

ATP before velocity sucrose sedimentation and western blotting (bottom blots).

Figure 2—figure supplement 1: **Controls and supporting *in vitro* analysis of Pom1**

clusters. (A) Fluorescence intensity trace of a single Pom1-mNG cluster binding to the

cell cortex, measured by high speed TIRF microscopy (~50 frames per second with

continuous acquisition). Rapid intensity changes indicating binding and unbinding are

marked by green upward and magenta downward pointing arrows respectively. **(B)** FRAP

analysis of supported lipid bilayers. Upper panel: A circular ROI (teal dashed line) was

used to bleach fluorescent PE in the bilayers. Lower panel: Kymograph at white dashed

line in upper panel of fluorescence recovery as monitored by TIRF microscopy, indicating

fluidity. **(C)** Quantification and comparison of binding duration of either wild-type or

kinase-dead Pom1-mNG clusters to supported lipid bilayers in TIRF microscopy movies

(***p=<0.0001, *n*=1526 wild-type, 1993 kinase-dead). Statistical significance was tested

with a Student's T-test. **(D)** TIRF microscopy images of cell extracts prepared from cells

co-expressing Pom1-mNG and Cdr2-tagRFP-t. Pom1 clusters are shown in green and

Cdr2 nodes are shown in magenta. Images are 3 frame sum projections of continuous

200ms time-lapse exposures. Scale bar 1µm.

Figure 2—figure supplement 2: **Quantification of Pom1 sedimentation in velocity**

sucrose gradients and size standards. (A) Quantification of Pom1-3HA band

intensities from western blot of yeast extracts in main **(Figure 2C)**. **(B)** Size standards for

velocity sucrose gradients, as in main (Figure 2C,G). Data are quantified from single

Coomassie-stained gels or using spectrophotometry data. **(C)** Plot of S-value versus
sedimentation peak. Line is the linear regression of the three size standards, with the S-
value for Pom1 peaks interpolated.

Figure 2—figure supplement 3: **Analysis of Pom1 cluster diffusion on supported lipid**
bilayers (A) Pom1-mNG clusters diffusing on supported lipid bilayers imaged by high-
speed (100ms acquisition) TIRF microscopy. **(B)** Single-particle diffusion traces of Pom1-
mNG clusters on supported lipid bilayers. **(C)** Mean squared displacement (MSD) curves
calculated from traces of individual Pom1-mNG cluster diffusion as in panel (B). **(D)** Mean
MSD curves generated from averaging of 421 single-particle traces. Weighted linear
regression of the curve (red line) provides the diffusion coefficient D= $0.168\pm0.018\mu m^2$/s.
Note that the weighted regression best fits the initial region of the curve because few
clusters bind longer than 10s. Black line represents weighted mean of all traces, grey
lines indicate standard deviation, and error bars represent weighted standard deviation.

Figure 3: **Polarity landmarks pattern cortical dynamics of Pom1 clusters to shape**
the gradient. (A) Western blots of sucrose gradient fractions after sedimentation analysis
as in Figure 2C of extracts prepared from *tea1Δ* or *tea4Δ* cells expressing Pom1-3HA. **(B)**
TIRF micrographs of Pom1-mNG localization at the tip or side cortex in *tea1Δ* or *tea4Δ*
cells. Scale bar 1μm. **(C)** The mean binding frequency of Pom1-mNG clusters at the tips
and sides of wild-type, *tea1Δ* or *tea4Δ* cells (*n*=10 cells, 7-170 traces/cell). **(D)** The mean
dwell time of Pom1-mNG clusters at the tips and sides of wild-type, *tea1Δ* or *tea4Δ* cells.
For panels C-D, wild-type data are replotted from main **(Figure 1)**. Each data point
represents a single cell mean, and line and error bars represent mean and standard
deviation of all cells (*n*=10 cells, 7-170 traces/cell). Comparisons are 1-way ANOVA with
Tukey's multiple comparisons tests. (****) indicates *p* < 0.0001, (n.s.) indicates *p* > 0.05.
See Figure 3—figure supplement 1D for all results of statistical analysis.

Figure 3—figure supplement 1: **Supporting analysis of Pom1 clusters in polarity**
mutants. (A) Confocal micrographs of Pom1-mNG in *tea1Δ* or *tea4Δ* cells. Images are
from the middle focal plane; scale bar 5μm. **(B)** Quantification of cortical binding dwell

time of individual Pom1-mNG clusters at the tips or sides of representative wild-type,

tea1Δ or *tea4Δ* cells (*n*=44-129 traces). **(C)** Quantification of total lifetime displacement of

individual Pom1-mNG clusters at the tips or sides of representative wild-type, *tea1Δ* or

tea4Δ cells. For B-C, line and error represent single cell mean and standard deviation

(*n*=44-129 traces). **(D)**   Results of ANOVA and Tukey's multiple comparison tests

performed in main (Figure 3C,D).

Figure 4: **Size-scaling of Pom1 clusters and Cdr2 nodes at the lateral cell cortex. (A)**

A subset of Pom1 clusters colocalize with Cdr2 nodes. Panels are TIRF microscopy

images of cells expressing Pom1-mNG and Cdr2-tagRFP-t. Yellow dashed brackets

outline the ROI of the lower zoomed panels. Orange arrows point to Pom1 clusters

colocalized with Cdr2 nodes. Scale bar is 1µm**. (B)** Kymographs generated using the ROI

indicated in panel A. Cyan arrows indicate prominent Pom1 clusters, magenta arrows

indicate prominent Cdr2 nodes, and orange arrows indicate colocalization. **(C)**

Localization of Cdr2-mEGFP (upper panels) or Pom1-mNG (lower panels) in

representative cells of increasing size, imaged by TIRF microscopy. Scale bar is 1µm.

(D) Total number of Cdr2-mEGFP nodes (blue circles) or Pom1-mNG clusters (red

squares) measured per cell. Quantification is limited to clusters detected and resolvable

in the TIRF illumination field. The slopes of the corresponding linear regressions are not

significantly different (*p*=0.3757, *n*>40 cells). **(E)** Density of Cdr2-mEGFP nodes (blue

circles) or Pom1-mNG clusters (red squares) in 2x2µm square ROIs at the cell middle,

counted using TIRF microscopy. The slopes of the corresponding linear regressions are

significantly different (*p*<0.0001, *n*>40 cells). **(F)** Example of colocalization analysis for

Pom1 clusters (magenta ROI) and Cdr2 nodes (cyan ROI) (left panels). Colocalized

structures are marked with overlapping yellow ROIs, whereas non-colocalizing structures

are marked with grey ROIs (right panel, grey circles). **(G)** Ratio of free Cdr2 nodes to free

Pom1 clusters, plotted as a function of cell size. The slope of the linear regression is

positive and significantly non-zero (*p*=0.0002, R^2=0.33, *n*=36 cells).

Figure 4—figure supplement 1: **Quantification of Pom1 and Cdr2 concentration in**

different cellular regions. (A) Schematic showing the ROI locations used to measure

the concentration of Pom1-mNG or Cdr2-mNG in the cytoplasm, cortex, tip cortex, or side

cortex. Images are spinning disc confocal micrographs of middle cell focal planes. Scale

bars 1µm. **(B)** Quantification of Cdr2 concentration (average fluorescence per pixel) at

different cellular ROIs as in panel (A). Each dot represents a measurement from a

different cell, numbers and the line and error represent the mean and standard deviation.

(C) Quantification of Pom1 concentration (average fluorescence per pixel) at different

cellular ROIs as in panel A. Each dot represents a measurement from a different cell,

numbers and the line and error represent the mean and standard deviation. **(D)**

Quantification of the cytoplasmic concentration (average fluorescence per pixel) of Cdr2-

mNG as a function of cell length, using a cytoplasmic ROI as in panel (A). The slope of

the linear regression does not significantly deviate from zero ($b_1 \neq 0$, (n.s.) p=0.6449). Each

point indicates a measurement from a single cell. **(E)** Quantification of the cytoplasmic

concentration (average fluorescence) of Pom1-mNG as a function of cell length, using a

cytoplasmic ROI as in panel (A). The slope of the linear regression does not significantly

deviate from zero ($b_1 \neq 0$, (n.s.) p=0.4926). Each point indicates a measurement from a

single cell.

<u>Figure 4—figure supplement 2:</u> **Colocalization of Cdr2 nodes and Pom1 clusters**

using alternative fluorophore pair. (A) A subset of Pom1 clusters colocalize with Cdr2

nodes. Images are dual-channel simultaneously-acquired TIRF microscopy images of

cells expressing Pom1-tdTomato and Cdr2-mNeonGreen. Yellow dashed brackets

outline the ROI of the lower zoomed panels. White arrows point to Pom1 clusters

colocalized with Cdr2 nodes. Scale bar is 1µm**. (B)** Kymographs generated using the ROI

drawn on the merged image in panel A. Cyan arrows indicate prominent Cdr2 nodes,

magenta arrows indicate prominent Pom1 clusters, and orange arrows adjacent to the

merged kymograph indicate colocalization.

<u>Figure 4—figure supplement 3:</u> **Supporting analysis of Cdr2 nodes and Pom1 clusters**

at the medial cell cortex. (A) Comparison of Pom1 cluster binding frequency on the cell

sides of wild-type and *cdr2Δ* cells (n.s., *p*=0.1076, *n*>10 cells, 25-42 traces/cell). **(B)**

Comparison of Pom1 cluster binding duration on the cell sides of wild-type and *cdr2Δ*

cells (n.s., *p*=0.371, *n*>10 cells, 25-42 traces/cell). **(C)** Comparison of Pom1 cluster

displacement on the cell sides of wild-type and *cdr2Δ* cells (n.s., *p*=0.8191, *n*>10 cells,

25-42 traces/cell). In A-C, significance was tested using a Student's T-test. **(D)** Schematic

and quantification of Pom1 concentration at the lateral cell cortex measured by confocal

microscopy, using 2μm line scans at the cell middle from images such as Figure 4—figure

supplement 1A. The slope of the linear regression is negative, but not significantly non-

zero (*p*=0.2603, R^2=0.03, n=45 cells). **(E)** Plot of the number of colocalized structures as

a function of cell size. The slope of the linear regression is significantly non-zero

(*p*=0.0006, R^2=0.3, *n*=36 cells). **(F)** Plot of free, non-colocalized Cdr2 nodes (blue circles)

and free Pom1 clusters (red squares). The slope of the linear regression of #Pom1

clusters versus cell length is negative, and significantly non-zero (*p*=0.001, R^2=0.27, *n*=36

cells). The slope of the linear regression of #Cdr2 nodes versus cell length is positive,

and significantly non-zero (*p*=0.021, R^2=0.15, *n*=36 cells).

Figure 5: **Redistribution of Pom1 clusters to the lateral cortex disrupts Cdr2 nodes**

during glucose restriction. (A) Localization of Pom1-mNG in cells grown in either high

(2%) or low (0.03%) glucose media. Images were collected using TIRF microscopy. Scale

bar, 5μm. **(B)** Comparison of the binding frequency of Pom1 clusters at the lateral cell

cortex in cells grown under high and low glucose (***, *p*=0.0003, *n*=10 cells, 32-113

traces/cell). Statistical significance was tested using a Student's T-test. **(C)** Comparison

of the binding dwell time of Pom1 clusters at the lateral cell cortex in cells grown under

high and low glucose (n.s., *p*=0.6833, *n*=10 cells, 32-113 traces/cell). Statistical

significance was tested using a Student's T-test. **(D)** Localization of Cdr2-mNG in wild-

type or *pom1Δ* cells grown in either normal (2%) or low (0.03%) glucose media. Images

were collected using TIRF microscopy. Scale bar, 1μm. **(E)** Comparison of the

fluorescence intensity of individual Cdr2-mNG nodes in wild-type or *pom1Δ* cells grown

in either normal (2%) or low (0.03%) glucose media. Low glucose induces partial node

disassembly in wild-type cells (****, *p*<0.0001, *n*=92-169 nodes from >5 cells) but not in

pom1Δ cells (n.s., *p*=0.4015, *n*=113-156 nodes from >5 cells). Measurements were taken

from Airyscan Super-Resolution confocal micrographs. **(F)** Quantification and comparison

of the total number of Cdr2-mNG nodes visible in TIRF micrographs of wild-type or *pom1Δ*

cells grown in either normal (2%) or low (0.03%) glucose media. There is no significant

difference in any condition (*p*>0.05, *n*=28-33 cells). Statistical significance was tested

using a one-way ANOVA.

Figure 5—figure supplement 1: **Redistribution of Pom1 clusters to the lateral cortex**

disrupts Cdr2 nodes during glucose restriction, supporting data. (A) Analysis of

Pom1 and Cdr2 redistribution by spinning-disc confocal microscopy, as in (Kelkar and

Martin, 2015). Images are confocal micrographs of cells co-expressing Pom1-tdTomato

and Cdr2-mNG in high (2%) or low (0.03%) glucose media. Scale bar, 5μm. **(B)** Cortical

fluorescence intensity of Cdr2 and Pom1 was measured using line scans drawn around

the circumference of the representative cell marked with an arrowhead in the 2% glucose

portion of panel A. **(C)** Cortical fluorescence intensity of Cdr2 and Pom1 was measured

using line scans drawn around the circumference of the representative cell marked with

an arrowhead in the 0.03% glucose portion of panel A.

Figure 6: **Pom1 and Cdr2 concentrations at the lateral cell cortex depend on glucose**

availability and are anti-correlated. (A-B) Localization of Pom1-mNG (panel A) and

Cdr2-mNG (panel B) in confocal micrographs at middle cell focal planes. Regions where

line-scans were used to measure fluorescence intensity are marked at the tip and side

(orange lines). Scale bar, 1μm. **(C)** Concentration of Pom1 (red squares) or Cdr2 (blue

circles) at cell sides in a range of glucose concentrations, measured using 'Side ROI' as

in panels A and B (*n*>10 cells/concentration). **(D)** Concentration of Pom1 (red squares)

or Cdr2 (blue circles) at cell tips in a range of glucose concentrations, measured using

'Tip ROI' as in panels A and B (*n*>10 cells/concentration). **(E)** Correlation of Cdr2 vs Pom1

concentrations at cell sides in each glucose concentration from panel C. Concentrations

are anticorrelated across all tested media glucose concentrations (*p*=0.0006, Pearson r=-

0.9352, *n*=8 concentrations). **(F)** Correlation of Cdr2 vs Pom1 concentrations at cell tips

in each glucose concentration from panel D. Concentrations show weak anticorrelation

across the tested media glucose concentrations, but the correlation is not statistically

significant (*p*=0.09, Pearson r=-0.65, *n*=8 concentrations).

Figure 7: **Pom1 redistribution under glucose restriction disrupts Wee1 regulation at**
cortical nodes. (A) Localization of Wee1-mNG in wild-type or *pom1Δ* cells grown in
either high (2%) or low (0.03%) glucose media. Images were collected using TIRF
microscopy. Scale bar, 5μm. **(B)** Quantification of Wee1 bursting kinetics in wild-type cells
grown under high (2%, blue circles) and low (0.03%, red squares) glucose conditions.
The top panel is a plot of the total number of Wee1 bursts as a function of cell length,
counted in single time point TIRF micrographs. The middle panel is a plot of the frequency
of Wee1 bursts as a function of cell length. The bottom panel is a plot of Wee1 burst
duration as a function of cell length. **(C)** Quantification of Wee1 bursting kinetics in *pom1Δ*
cells grown under normal (2%, green circles) and low (0.03%, magenta squares) glucose
conditions, as in panel B. For (B-C), insets represent whether the best fit values of the
slope (b_1) and/or Y-intercept (b_0) of the two linear regressions are different at the 95%
confidence level. See Supplemental Table 1 and Supplemental Table 2 for parameters of
linear regressions and full statistical comparisons. **(D)** Western blot of Wee1 in wild-type*,*
pom1Δ, and *cdr2Δ* cells grown in high (2%) or low (0.03%) glucose media showing
phosphorylation dependent band-shifts.

1015

Figure 7—figure supplement 1: **Wee1 bursting in low glucose phenocopies a kinase-**
dead Cdr2 mutant. (A) Plot of the frequency of Wee1 bursts as a function of cell length
in wild-type cells grown in high glucose (blue circles) and low glucose (red squares), or
kinase-dead *cdr2(E177A)* mutant cells grown in high glucose (green triangles). **(B)** Plot
of Wee1 burst duration as a function of cell length in wild-type cells grown in low glucose
(red squares) and kinase-dead *cdr2(E177A)* mutant cells grown in high glucose (green
triangles). For (A-B), insets represent whether the best fit values of the slope (b_1) and/or
Y-intercept (b_0) of the linear regressions are different at the 95% confidence level. See
Supplemental Table 1 and Supplemental Table 2 for parameters of linear regressions
and full statistical comparisons.

1026

1027   Figure 8: **A model for the glucose-modulated control of Wee1 bursting by Pom1 and**
1028   **Cdr2.** See text for discussion.

1029

Supplemental Table 1: **Parameters of linear regressions in Figure 7B,C and Figure 7—figure supplement 1A,B.** For each data set, the table includes: The slope of the linear regression (b_1) and its 95% confidence intervals; whether the slope of the linear regression significantly deviates from zero at the 95% confidence interval; the Y-intercept of the linear regression (b_0) and its 95% confidence intervals; the coefficient of determination of the linear regression (R^2); the *n* of the data set.

Supplemental Table 2: **Summary of the statistical comparisons of the linear regressions in Figure 7B,C and Figure 7—figure supplement 1A,B.** For each comparison, the table includes: Whether the slopes of the linear regressions (b_1) are significantly different from each other at the 95% confidence level; whether the Y-intercepts of the linear regressions (b_0) are significantly different from each other at the 95% confidence level. If the slopes of the linear regressions differed significantly, the difference in Y-intercepts was not tested.

Video 1: **Pom1-mNG forms clusters that transiently bind the cell cortex.** TIRF microscopy of a cell expressing Pom1-mNG from the endogenous locus. Frames are 200ms exposures from continuous acquisition. Scale bar is 1µm.

Video 2: **Pom1-mNG clusters on synthetic supported lipid bilayers.** TIRF microscopy of Pom1-mNG clusters in cytoplasmic extracts, binding and releasing artificial supported lipid bilayers composed of phosphatidylcholine and phosphatidylserine. Frames are 200ms exposures with 200ms interval between consecutive frames. Scale bar is 1µm.

Acknowledgements:

We thank members of the Moseley lab and Erik Griffin for comments on the manuscript. B. Wickner, C. Barlowe and the Norris Cotton Cancer Center Molecular Interactions & Imaging Core at Dartmouth for sharing equipment. A. Lavanway, Z. Svindrych and the Life Sciences Light Microscopy Facility for assistance with microscopy. A. Orr for assistance with ultracentrifugation. Y. Wu for recombinant Pom1 proteins. This work was supported by grants from the American Cancer Society (RSG-15-140-01-CCG) and the

NIH (R01 GM099774) to J.B.M; and an NIH Training Grant (T32GM008704) to C.A.H.A. Shared imaging resources were supported by grants from the NIH (1S10OD018046 and P20GM113132).

References:

Akamatsu, M., Berro, J., Pu, K.-M., Tebbs, I.R., and Pollard, T.D. (2014). Cytokinetic nodes in fission yeast arise from two distinct types of nodes that merge during interphase. J Cell Biol *204*, 977–988.

Akamatsu, M., Lin, Y., Bewersdorf, J., and Pollard, T.D. (2017). Analysis of interphase node proteins in fission yeast by quantitative and superresolution fluorescence microscopy. MBoC *28*, 3203–3214.

Alberghina, L., Höfer, T., and Vanoni, M. (2009). Molecular networks and system-level properties. J. Biotechnol. *144*, 224–233.

Aligue, R., Wu, L., and Russell, P. (1997). Regulation of Schizosaccharomyces pombe Wee1 Tyrosine Kinase. J. Biol. Chem. *272*, 13320–13325.

Allard, C.A.H., Opalko, H.E., Liu, K.-W., Medoh, U., and Moseley, J.B. (2018). Cell size–dependent regulation of Wee1 localization by Cdr2 cortical nodes. J Cell Biol *217*, 1589–1599.

Bähler, J., and Nurse, P. (2001). Fission yeast Pom1p kinase activity is cell cycle regulated and essential for cellular symmetry during growth and division. The EMBO Journal *20*, 1064–1073.

Bähler, J., and Pringle, J.R. (1998). Pom1p, a fission yeast protein kinase that provides positional information for both polarized growth and cytokinesis. Genes Dev *12*, 1356–1370.

Bähler, J., Wu, J.-Q., Longtine, M.S., Shah, N.G., Iii, A.M., Steever, A.B., Wach, A., Philippsen, P., and Pringle, J.R. (1998). Heterologous modules for efficient and versatile PCR-based gene targeting in Schizosaccharomyces pombe. Yeast *14*, 943–951.

Bhatia, P., Hachet, O., Hersch, M., Rincon, S.A., Berthelot-Grosjean, M., Dalessi, S., Basterra, L., Bergmann, S., Paoletti, A., and Martin, S.G. (2014). Distinct levels in Pom1 gradients limit Cdr2 activity and localization to time and position division. Cell Cycle *13*, 538–552.

Bray, D., Levin, M.D., and Morton-Firth, C.J. (1998). Receptor clustering as a cellular mechanism to control sensitivity. Nature *393*, 85–88.

Breeding, C.S., Hudson, J., Balasubramanian, M.K., Hemmingsen, S.M., Young, P.G., and Gould, K.L. (1998). The cdr2(+) gene encodes a regulator of G2/M progression and cytokinesis in Schizosaccharomyces pombe. Mol. Biol. Cell *9*, 3399–3415.

Chang, F., and Martin, S.G. (2009). Shaping Fission Yeast with Microtubules. Cold Spring Harb Perspect Biol *1*, a001347.

Coleman, T.R., Tang, Z., and Dunphy, W.G. (1993). Negative regulation of the weel protein kinase by direct action of the nim1/cdr1 mitotic inducer. Cell *72*, 919–929.

Deng, L., and Moseley, J.B. (2013). Compartmentalized nodes control mitotic entry signaling in fission yeast. Mol. Biol. Cell *24*, 1872–1881.

Deng, L., Baldissard, S., Kettenbach, A.N., Gerber, S.A., and Moseley, J.B. (2014). Dueling kinases regulate cell size at division through the SAD kinase Cdr2. Curr. Biol. *24*, 428–433.

1121   Dickinson, D.J., Schwager, F., Pintard, L., Gotta, M., and Goldstein, B. (2017). A Single-
1122   Cell Biochemistry Approach Reveals PAR Complex Dynamics during Cell Polarization.
1123   Dev. Cell *42*, 416-434.e11.

1124

1125   Dodgson, J., Chessel, A., Yamamoto, M., Vaggi, F., Cox, S., Rosten, E., Albrecht, D.,
1126   Geymonat, M., Csikasz-Nagy, A., Sato, M., et al. (2013). Spatial segregation of polarity
1127   factors into distinct cortical clusters is required for cell polarity control. Nature
1128   Communications *4*, 1834.

1129

1130   Dolznig, H., Grebien, F., Sauer, T., Beug, H., and Müllner, E.W. (2004). Evidence for a
1131   size-sensing mechanism in animal cells. Nat. Cell Biol. *6*, 899–905.

1132

1133   Douglass, A.D., and Vale, R.D. (2005). Single-molecule microscopy reveals plasma
1134   membrane microdomains created by protein-protein networks that exclude or trap
1135   signaling molecules in T cells. Cell *121*, 937–950.

1136

1137   Duke, T.A., and Bray, D. (1999). Heightened sensitivity of a lattice of membrane
1138   receptors. Proc. Natl. Acad. Sci. U.S.A. *96*, 10104–10108.

1139

1140   Facchetti, G., Chang, F., and Howard, M. (2017). Controlling cell size through sizer
1141   mechanisms. Current Opinion in Systems Biology *5*, 86–92.

1142

1143   Facchetti, G., Knapp, B., Flor-Parra, I., Chang, F., and Howard, M. (2019).
1144   Reprogramming Cdr2-Dependent Geometry-Based Cell Size Control in Fission Yeast.
1145   Current Biology *29*, 350-358.e4.

1146

1147   Falke, J.J. (2002). Cooperativity between bacterial chemotaxis receptors. Proc. Natl.
1148   Acad. Sci. U.S.A. *99*, 6530–6532.

1149

1150   Fantes, P.A. (1977). Control of cell size and cycle time in Schizosaccharomyces pombe.
1151   J. Cell. Sci. *24*, 51–67.

Fantes, P., and Nurse, P. (1977). Control of cell size at division in fission yeast by a growth-modulated size control over nuclear division. Exp. Cell Res. *107*, 377–386.

Feilotter, H., Nurse, P., and Young, P.G. (1991). Genetic and Molecular Analysis of Cdr1/Nim1 in Schizosaccharomyces Pombe. Genetics *127*, 309–318.

Gautier, J., Solomon, M.J., Booher, R.N., Bazan, J.F., and Kirschner, M.W. (1991). cdc25 is a specific tyrosine phosphatase that directly activates p34cdc2. Cell *67*, 197–211.

Ginzberg, M.B., Kafri, R., and Kirschner, M. (2015). Cell biology. On being the right (cell) size. Science *348*, 1245075.

Gould, K.L., and Nurse, P. (1989). Tyrosine phosphorylation of the fission yeast cdc2 + protein kinase regulates entry into mitosis. Nature *342*, 39.

Gronnier, J., Crowet, J.-M., Habenstein, B., Nasir, M.N., Bayle, V., Hosy, E., Platre, M.P., Gouguet, P., Raffaele, S., Martinez, D., et al. (2017). Structural basis for plant plasma membrane protein dynamics and organization into functional nanodomains. Elife *6*.

Gu, Y., and Oliferenko, S. (2019). Cellular geometry scaling ensures robust division site positioning. Nat Commun *10*, 268.

Guzmán-Vendrell, M., Rincon, S.A., Dingli, F., Loew, D., and Paoletti, A. (2015). Molecular control of the Wee1 regulatory pathway by the SAD kinase Cdr2. J. Cell. Sci. *128*, 2842–2853.

Hachet, O., Berthelot-Grosjean, M., Kokkoris, K., Vincenzetti, V., Moosbrugger, J., and Martin, S.G. (2011). A phosphorylation cycle shapes gradients of the DYRK family kinase Pom1 at the plasma membrane. Cell *145*, 1116–1128.

1183 Harashima, H., Dissmeyer, N., and Schnittger, A. (2013). Cell cycle control across the
1184 eukaryotic kingdom. Trends Cell Biol. *23*, 345–356.

1185

1186 Haupt, A., and Minc, N. (2017). Gradients of phosphatidylserine contribute to plasma
1187 membrane charge localization and cell polarity in fission yeast. Mol Biol Cell *28*, 210–220.

1188

1189 Hersch, M., Hachet, O., Dalessi, S., Ullal, P., Bhatia, P., Bergmann, S., and Martin, S.G.
1190 (2015). Pom1 gradient buffering through intermolecular auto-phosphorylation. Mol. Syst.
1191 Biol. *11*, 818.

1192

1193 Jorgensen, P., and Tyers, M. (2004). How Cells Coordinate Growth and Division. Current
1194 Biology *14*, R1014–R1027.

1195

1196 Kalay, Z., Fujiwara, T.K., and Kusumi, A. (2012). Confining domains lead to reaction
1197 bursts: reaction kinetics in the plasma membrane. PLoS ONE *7*, e32948.

1198

1199 Kanoh, J., and Russell, P. (1998). The Protein Kinase Cdr2, Related to Nim1/Cdr1 Mitotic
1200 Inducer, Regulates the Onset of Mitosis in Fission Yeast. Mol Biol Cell *9*, 3321–3334.

1201

1202 Keifenheim, D., Sun, X.-M., D'Souza, E., Ohira, M.J., Magner, M., Mayhew, M.B.,
1203 Marguerat, S., and Rhind, N. (2017). Size-Dependent Expression of the Mitotic Activator
1204 Cdc25 Suggests a Mechanism of Size Control in Fission Yeast. Curr. Biol. *27*, 1491-
1205 1497.e4.

1206

1207 Kelkar, M., and Martin, S.G. (2015). PKA antagonizes CLASP-dependent microtubule
1208 stabilization to re-localize Pom1 and buffer cell size upon glucose limitation. Nature
1209 Communications *6*, 8445.

1210

1211 Kettenbach, A.N., Deng, L., Wu, Y., Baldissard, S., Adamo, M.E., Gerber, S.A., and
1212 Moseley, J.B. (2015). Quantitative phosphoproteomics reveals pathways for coordination

of cell growth and division by the conserved fission yeast kinase pom1. Mol. Cell Proteomics *14*, 1275–1287.

Knight, J.D., Lerner, M.G., Marcano-Velázquez, J.G., Pastor, R.W., and Falke, J.J. (2010). Single Molecule Diffusion of Membrane-Bound Proteins: Window into Lipid Contacts and Bilayer Dynamics. Biophys J *99*, 2879–2887.

Kumagai, A., and Dunphy, W.G. (1991). The cdc25 protein controls tyrosine dephosphorylation of the cdc2 protein in a cell-free system. Cell *64*, 903–914.

Lee, S., Lim, W.A., and Thorn, K.S. (2013). Improved blue, green, and red fluorescent protein tagging vectors for S. cerevisiae. PLoS ONE *8*, e67902.

Liu, W., Wang, H., and Xu, C. (2016). Antigen Receptor Nanoclusters: Small Units with Big Functions. Trends Immunol. *37*, 680–689.

Lloyd, A.C. (2013). The Regulation of Cell Size. Cell *154*, 1194–1205.

Lucena, R., Alcaide-Gavilán, M., Anastasia, S.D., and Kellogg, D.R. (2017). Wee1 and Cdc25 are controlled by conserved PP2A-dependent mechanisms in fission yeast. Cell Cycle *16*, 428–435.

Maity, P.C., Blount, A., Jumaa, H., Ronneberger, O., Lillemeier, B.F., and Reth, M. (2015). B cell antigen receptors of the IgM and IgD classes are clustered in different protein islands that are altered during B cell activation. Sci Signal *8*, ra93.

Makushok, T., Alves, P., Huisman, S.M., Kijowski, A.R., and Brunner, D. (2016). Sterol-Rich Membrane Domains Define Fission Yeast Cell Polarity. Cell *165*, 1182–1196.

Martin, S.G., and Berthelot-Grosjean, M. (2009). Polar gradients of the DYRK-family kinase Pom1 couple cell length with the cell cycle. Nature *459*, 852–856.

1244

1245   Masuda, H., Fong, C.S., Ohtsuki, C., Haraguchi, T., and Hiraoka, Y. (2011).

1246   Spatiotemporal regulations of Wee1 at the G2/M transition. Mol. Biol. Cell *22*, 555–569.

1247

1248   Moreno, S., Hayles, J., and Nurse, P. (1989). Regulation of p34cdc2 protein kinase during

1249   mitosis. Cell *58*, 361–372.

1250

1251   Moreno, S., Nurse, P., and Russell, P. (1990). Regulation of mitosis by cyclic

1252   accumulation of p80cdc25 mitotic inducer in fission yeast. Nature *344*, 549–552.

1253

1254   Moreno, S., Klar, A., and Nurse, P. (1991). Molecular genetic analysis of fission yeast

1255   Schizosaccharomyces pombe. Meth. Enzymol. *194*, 795–823.

1256

1257   Morrell, J.L., Nichols, C.B., and Gould, K.L. (2004). The GIN4 family kinase, Cdr2p, acts

1258   independently of septins in fission yeast. Journal of Cell Science *117*, 5293–5302.

1259

1260   Moseley, J.B., Mayeux, A., Paoletti, A., and Nurse, P. (2009). A spatial gradient

1261   coordinates cell size and mitotic entry in fission yeast. Nature *459*, 857–860.

1262

1263   Munro, E. (2017). Protein Clustering Shapes Polarity Protein Gradients. Dev. Cell *42*,

1264   309–311.

1265

1266   Neumann, F.R., and Nurse, P. (2007). Nuclear size control in fission yeast. J Cell Biol

1267   *179*, 593–600.

1268

1269   Opalko, H.E., and Moseley, J.B. (2017). Dynamic regulation of Cdr1 localization and

1270   phosphorylation during osmotic stress. J. Biol. Chem. jbc.M117.793034.

1271

1272   Pan, K.Z., Saunders, T.E., Flor-Parra, I., Howard, M., and Chang, F. (2014). Cortical

1273   regulation of cell size by a sizer cdr2p. ELife *3*.

1274

Parker, L.L., Walter, S.A., Young, P.G., and Piwnica-Worms, H. (1993). Phosphorylation and inactivation of the mitotic inhibitor Weel by the nim1/cdr1 kinase. Nature *363*, 736.

Pluskal, T., Hayashi, T., Saitoh, S., Fujisawa, A., and Yanagida, M. (2011). Specific biomarkers for stochastic division patterns and starvation-induced quiescence under limited glucose levels in fission yeast. FEBS J. *278*, 1299–1315.

Qian, H., Sheetz, M.P., and Elson, E.L. (1991). Single particle tracking. Analysis of diffusion and flow in two-dimensional systems. Biophysical Journal *60*, 910–921.

Rincon, S.A., Bhatia, P., Bicho, C., Guzman-Vendrell, M., Fraisier, V., Borek, W.E., Alves, F. de L., Dingli, F., Loew, D., Rappsilber, J., et al. (2014). Pom1 regulates the assembly of Cdr2–Mid1 cortical nodes for robust spatial control of cytokinesis. J Cell Biol *206*, 61–77.

Rodriguez, J., Peglion, F., Martin, J., Hubatsch, L., Reich, J., Hirani, N., Gubieda, A.G., Roffey, J., Fernandes, A.R., St Johnston, D., et al. (2017). aPKC Cycles between Functionally Distinct PAR Protein Assemblies to Drive Cell Polarity. Dev. Cell *42*, 400-415.e9.

Russell, P., and Nurse, P. (1986). cdc25+ functions as an inducer in the mitotic control of fission yeast. Cell *45*, 145–153.

Russell, P., and Nurse, P. (1987). Negative regulation of mitosis by wee1+, a gene encoding a protein kinase homolog. Cell *49*, 559–567.

Saunders, T.E., Pan, K.Z., Angel, A., Guan, Y., Shah, J.V., Howard, M., and Chang, F. (2012). Noise reduction in the intracellular pom1p gradient by a dynamic clustering mechanism. Dev. Cell *22*, 558–572.

Sbalzarini, I.F., and Koumoutsakos, P. (2005). Feature point tracking and trajectory analysis for video imaging in cell biology. J. Struct. Biol. *151*, 182–195.

Schmoller, K.M., Turner, J.J., Kõivomägi, M., and Skotheim, J.M. (2015). Dilution of the cell cycle inhibitor Whi5 controls budding-yeast cell size. Nature *526*, 268–272.

Shiozaki, K. (2009). Nutrition-Minded Cell Cycle. Sci. Signal. *2*, pe74–pe74.

Simanis, V., and Nurse, P. (1986). The cell cycle control gene cdc2+ of fission yeast encodes a protein kinase potentially regulated by phosphorylation. Cell *45*, 261–268.

Spira, F., Mueller, N.S., Beck, G., von Olshausen, P., Beig, J., and Wedlich-Söldner, R. (2012). Patchwork organization of the yeast plasma membrane into numerous coexisting domains. Nature Cell Biology *14*, 640–648.

Strausfeld, U., Labbé, J.C., Fesquet, D., Cavadore, J.C., Picard, A., Sadhu, K., Russell, P., and Dorée, M. (1991). Dephosphorylation and activation of a p34cdc2/cyclin B complex in vitro by human CDC25 protein. Nature *351*, 242–245.

Su, X., Ditlev, J.A., Hui, E., Xing, W., Banjade, S., Okrut, J., King, D.S., Taunton, J., Rosen, M.K., and Vale, R.D. (2016). Phase separation of signaling molecules promotes T cell receptor signal transduction. Science *352*, 595–599.

Tarantino, N., Tinevez, J.-Y., Crowell, E.F., Boisson, B., Henriques, R., Mhlanga, M., Agou, F., Israël, A., and Laplantine, E. (2014). TNF and IL-1 exhibit distinct ubiquitin requirements for inducing NEMO–IKK supramolecular structures. J Cell Biol *204*, 231–245.

Tinevez, J.-Y., Perry, N., Schindelin, J., Hoopes, G.M., Reynolds, G.D., Laplantine, E., Bednarek, S.Y., Shorte, S.L., and Eliceiri, K.W. (2017). TrackMate: An open and extensible platform for single-particle tracking. Methods *115*, 80–90.

Wang, S.-C., Low, T.Y.F., Nishimura, Y., Gole, L., Yu, W., and Motegi, F. (2017). Cortical forces and CDC-42 control clustering of PAR proteins for Caenorhabditis elegans embryonic polarization. Nat. Cell Biol. *19*, 988–995.

Weiß, K., Neef, A., Van, Q., Kramer, S., Gregor, I., and Enderlein, J. (2013). Quantifying the Diffusion of Membrane Proteins and Peptides in Black Lipid Membranes with 2-Focus Fluorescence Correlation Spectroscopy. Biophys J *105*, 455–462.

Wood, E., and Nurse, P. (2013). Pom1 and cell size homeostasis in fission yeast. Cell Cycle *12*, 3228–3236.

Wu, J.-Q., and Pollard, T.D. (2005). Counting cytokinesis proteins globally and locally in fission yeast. Science *310*, 310–314.

Wu, L., and Russell, P. (1993). Nim1 kinase promotes mitosis by inactivating Wee1 tyrosine kinase. Nature *363*, 738–741.

Wu, L., Shiozaki, K., Aligue, R., and Russell, P. (1996). Spatial organization of the Nim1-Wee1-Cdc2 mitotic control network in Schizosaccharomyces pombe. Mol. Biol. Cell 7, 1749–1758.

Yanagida Mitsuhiro, Ikai Nobuyasu, Shimanuki Mizuki, and Sajiki Kenichi (2011). Nutrient limitations alter cell division control and chromosome segregation through growth-related kinases and phosphatases. Philosophical Transactions of the Royal Society B: Biological Sciences *366*, 3508–3520.

Young, P.G., and Fantes, P.A. (1987). Schizosaccharomyces pombe mutants affected in their division response to starvation. Journal of Cell Science *88*, 295–304.

1366 Zhou, Y., and Hancock, J.F. (2015). Ras nanoclusters: Versatile lipid-based signaling

1367 platforms. Biochim. Biophys. Acta *1853*, 841–849.

1368

Figure 1

Figure 1 – figure supplement 1

A. Pom1-mNG TIRFM

2µm

2µm

B. Single-Cell Particle Tracks

C. Single Particle Traces

D. $D = 0.134µm^2/s$
$R^2 = 0.960$

E.

TIP   SIDE

***

Figure 1 – figure supplement 2

Figure 2

A. Single Puncta Trace
TIRF Microscopy -20ms/frame

B. 0.01% Liss-Rhod-PE
75% PC 25%PS

Bleach ROI

Time (s) 0 → 40

C.

D. Cell Extract - TIRFM

Cdr2-tagRFP-t Pom1-mNG Merge

Figure 2 – figure supplement 1

Figure 2 – figure supplement 2

A. Pom1-mNG on SLB - TIRFM

2µm

2µm

B. Single-Cell Particle Tracks

C.

D.

D = 0.168µm²/s
R² = 0.972

Figure 2 – figure supplement 3

Figure 3

A.

Pom1-mNG

tea1Δ

tea4Δ

Confocal Microscopy

B. **Single Cell Data**

Dwell Time (s)

WT TIP, WT SIDE, *tea1Δ* TIP, *tea1Δ* SIDE, *tea4Δ* TIP, *tea4Δ* TIP

C. **Single Cell Data**

Displacement (μm)

WT TIP, WT SIDE, *tea1Δ* TIP, *tea1Δ* SIDE, *tea4Δ* TIP, *tea4Δ* TIP

D.

ANOVA summary of **FREQUENCY**	
F	46.32
P value	<0.0001
P value summary	****
Significant diff. among means (P < 0.05)?	Yes
R square	0.8081

ANOVA summary of **DWELL TIME**	
F	10.34
P value	<0.0001
P value summary	****
Significant diff. among means (P < 0.05)?	Yes
R square	0.4846

Tukey's Multiple Comparisons Summary of **FREQUENCY** Tukey's Multiple Comparisons Summary of **DWELL TIME**

-	Mean Diff.	Significant?	Summary	Adjusted P Value	Mean Diff.	Significant?	Summary	Adjusted P Value
WT TIP vs. WT SIDE	1.891	Yes	****	<0.0001	0.05557	No	ns	0.9972
WT TIP vs. tea1Δ TIP	0.98	Yes	**	0.0013	0.5207	Yes	***	0.0009
WT TIP vs. tea1Δ SIDE	1.485	Yes	****	<0.0001	0.2832	No	ns	0.1718
WT TIP vs. tea4Δ TIP	2.763	Yes	****	<0.0001	0.5577	Yes	***	0.0003
WT TIP vs. tea4Δ SIDE	2.983	Yes	****	<0.0001	0.6484	Yes	****	<0.0001
WT SIDE vs. tea1Δ TIP	-0.9114	Yes	**	0.0032	0.4651	Yes	**	0.0039
WT SIDE vs. tea1Δ SIDE	-0.4065	No	ns	0.4816	0.2277	No	ns	0.3927
WT SIDE vs. tea4Δ TIP	0.8714	Yes	**	0.0055	0.5022	Yes	**	0.0015
WT SIDE vs. tea4Δ SIDE	1.091	Yes	***	0.0003	0.5929	Yes	***	0.0001
tea1Δ TIP vs. tea1Δ SIDE	0.5049	No	ns	0.2449	-0.2375	No	ns	0.3458
tea1Δ TIP vs. tea4Δ TIP	1.783	Yes	****	<0.0001	0.03705	No	ns	0.9996
tea1Δ TIP vs. tea4Δ SIDE	2.003	Yes	****	<0.0001	0.1277	No	ns	0.8947
tea1Δ SIDE vs. tea4Δ TIP	1.278	Yes	****	<0.0001	0.2745	No	ns	0.1989
tea1Δ SIDE vs. tea4Δ SIDE	1.498	Yes	****	<0.0001	0.3652	Yes	*	0.034
tea4Δ TIP vs. tea4Δ SIDE	0.22	No	ns	0.9327	0.09068	No	ns	0.9741

Figure 3 – figure supplement 1

Figure 4

Figure 4 – figure supplement 1

A.

Pom1
-tdTomato

Cdr2
-mNG

Merge

B.

Cdr2-mNG

Pom1-tdTomato

Merge

0s   5s   10s   15s

Figure 4 – figure supplement 2

Figure 4 – figure supplement 3

Figure 5

A. Confocal Microscopy

Pom1-tdTomato   Cdr2-mNG   Merge

2% Glucose

0.03% Glucose

B.

2% Glucose

Pom1 Intensity (A.U.) / Cdr2 Intensity (A.U.) vs Distance from Tip (μm)

C.

0.03% Glucose

Pom1 Intensity (A.U.) / Cdr2 Intensity (A.U.) vs Distance from Tip (μm)

Figure 5 – figure supplement 1

Figure 6

Figure 7

Figure 7 – figure supplement 1

CELL TIP

CELL SIDE

Pom1 Puncta   Cdr1/2 Node   Wee1

Small Cell

Large Cell

Glucose Starvation

Figure 8