Title: Stable Pom1 clusters form a glucose-modulated concentration gradient that regulates mitotic entry

Authors: Corey A.H. Allard, Hannah E. Opalko, and James B. Moseley*

Author Affiliations: Department of Biochemistry and Cell Biology
The Geisel School of Medicine at Dartmouth, Hanover, NH 03755
*Corresponding author
james.b.moseley@dartmouth.edu

Abstract: Control of cell size requires molecular size sensors that are coupled to the cell cycle. Rod-shaped fission yeast cells divide at a threshold size partly due to Cdr2 kinase, which forms nodes at the medial cell cortex where it inhibits the Cdk1-inhibitor Wee1. Pom1 kinase phosphorylates and inhibits Cdr2, and forms cortical concentration gradients from cell poles. Pom1 inhibits Cdr2 signaling to Wee1 specifically in small cells, but the time and place of their regulatory interactions were unclear. We show that Pom1 forms stable oligomeric puncta that dynamically sample the cell cortex. Binding frequency is patterned into a concentration gradient by the polarity landmarks Tea1 and Tea4. Pom1 puncta colocalize with Cdr2 nodes, forming a glucose-modulated inhibitory threshold against node activation. Our work reveals how Pom1-Cdr2-Wee1 operates in multiprotein clusters at the cell cortex to promote mitotic entry at a specific size that can be modified by nutrient availability.

Introduction:

Many cell types display a remarkable ability to maintain a constant size during rapid cycles of growth and division (Fantes and Nurse, 1977; Ginzberg et al., 2015; Jorgensen and Tyers, 2004; Lloyd, 2013). Such cell size control is a systems-level property that emerges from the integration of multiple size-dependent signal transduction pathways. Each signaling pathway is comprised of tunable biochemical parameters, including gene-expression and post-translational modifications such as protein phosphorylation (Alberghina et al., 2009). One major challenge in cell size research is to understand the biochemical mechanisms of signal transduction in each pathway, and what makes them size-dependent. These control systems can generate size homogeneity for a given cell type, but cell size is also an adaptable property. For example, nutritional cues and other environmental factors can alter cell size (Fantes and Nurse, 1977; Kelkar and Martin, 2015; Shiozaki, 2009; Yanagida Mitsuhiro et al., 2011). Thus, a second major challenge in cell size research is to understand how size-dependent signaling pathways respond to changes in cell metabolism and stress.

In eukaryotic cells, these signaling pathways lead to regulated activation of the conserved cyclin-dependent kinase Cdk1 (Harashima et al., 2013). Activated Cdk1 triggers mitotic entry and the cascade of events that lead to cell division (Gould and Nurse, 1989; Simanis and Nurse, 1986). The fission yeast *Schizosaccharomyces pombe* represents an excellent model system to study size-dependent signaling pathways that regulate Cdk1. Genetic screens performed in past decades have identified many conserved factors that regulate Cdk1, but how these factors form size-dependent signaling pathways remains less clear. Fission yeast cells have a simple geometry that facilitates cell size studies. These cylindrical cells maintain a constant cell width, and grow by linear extension during interphase (Fantes and Nurse, 1977; Moreno et al., 1989). A network of cell polarity proteins positioned at cell tips restricts growth to these sites and maintains proper cell morphology (Chang and Martin, 2009). As a result, cell length doubles in one cell cycle, and many aspects of cell geometry scale with this increase in cell length (Gu and Oliferenko, 2019; Neumann and Nurse, 2007). Recent studies used cell shape mutants to show that fission yeast cells primarily measure surface area, not

length or volume, for cell size control (Facchetti et al., 2019; Pan et al., 2014). A critical

next step is to understand how signaling pathways that regulate Cdk1 might operate at

the cell cortex in a size-dependent manner.

In all eukaryotes, Cdk1 activity is established by the opposing activities of the

inhibitory protein kinase Wee1, and the counteracting phosphatase Cdc25 (Gautier et al.,

1991; Gould and Nurse, 1989; Kumagai and Dunphy, 1991; Russell and Nurse, 1986,

1987; Strausfeld et al., 1991). In fission yeast, mutations in Wee1, Cdc25, and their

upstream regulators lead to changes in cell size that correspond to changes in the

signaling pathways. Separate mechanisms appear to link cell size with regulation of Wee1

versus Cdc25. For Cdc25, its cellular concentration increases through unknown

mechanisms as cells grow during interphase (Keifenheim et al., 2017; Moreno et al.,

1990). In contrast, the concentration of Wee1 remains constant during interphase, but it

is progressively phosphorylated by the conserved inhibitory kinases Cdr1 and Cdr2

(Aligue et al., 1997; Breeding et al., 1998; Kanoh and Russell, 1998; Keifenheim et al.,

2017; Lucena et al., 2017; Opalko and Moseley, 2017; Russell and Nurse, 1987; Wu and

Russell, 1993; Young and Fantes, 1987). *cdr2* mutants fail to divide at a constant surface

area, and instead dividing according to cell volume or length (Facchetti et al., 2019). This

change suggests that Cdr2-Cdr1-Wee1 signaling underlies the primary size-sensing

pathway that relates to surface area, while additional pathways related to volume and

length are engaged in its absence. The localization of Cdr2, Cdr1, and Wee1 supports

this model. Cdr2 forms punctate oligomeric structures called nodes that stably bind to the

medial cell cortex, and recruits Cdr1 to these sites (Akamatsu et al., 2014, 2017; Guzmán-

Vendrell et al., 2015; Martin and Berthelot-Grosjean, 2009; Morrell et al., 2004; Moseley

et al., 2009). Wee1 localizes primarily in the nucleus and spindle-pole body, where it

encounters Cdk1 to prevent mitotic entry (Masuda et al., 2011; Moseley et al., 2009; Wu

et al., 1996). In addition, Wee1 transiently visits cortical Cdr1/2 nodes in bursts that lead

to inhibition of Wee1 kinase activity (Allard et al., 2018). The frequency and duration of

these Wee1 bursts at Cdr1/2 nodes increase approximately twenty-fold as cells double in

size during interphase (Allard et al., 2018). This mechanism leads to size-dependent

inhibition of Wee1 by a signaling pathway at the cell surface.

This size-dependent change in Wee1 bursting dynamics is encoded into the Cdr1-Cdr2-Wee1 pathway at two proposed levels. First, increased Wee1 bursting depends upon the doubling in number of Cdr1/2 nodes, which occurs as growth results in a doubling of cell surface area, but this increase is smaller than the magnitude of change in Wee1 bursting (~2-fold vs ~20-fold) (Allard et al., 2018; Deng and Moseley, 2013; Pan et al., 2014). Second, activation of Cdr2 increases as cells increase in size (Deng et al., 2014). Cdr2 kinase activity is required for Wee1 localization to nodes, and is controlled by the upstream kinase Pom1 (Allard et al., 2018; Moseley et al., 2009). Pom1 directly phosphorylates and inhibits the activation of Cdr2 (Bhatia et al., 2014; Deng et al., 2014; Kettenbach et al., 2015; Martin and Berthelot-Grosjean, 2009; Moseley et al., 2009). Pom1 also phosphorylates a separate set of sites on Cdr2 to disrupt oligomerization and membrane binding (Bhatia et al., 2014; Rincon et al., 2014). This regulation of Cdr2 by Pom1 changes the size dependency of Wee1 bursts. Specifically, the frequency and duration of Wee1 bursts at Cdr1/2 nodes increases, but only in small cells (Allard et al., 2018). Consistent with this defect, *pom1∆* cells divide at a small size due to dysregulation of the Cdr2-Cdr1-Wee1 pathway (Bähler and Pringle, 1998; Bhatia et al., 2014; Martin and Berthelot-Grosjean, 2009; Moseley et al., 2009; Wood and Nurse, 2013). Taken together, these results suggest a size-dependent interaction between Pom1 and its substrate Cdr2. However, the location and timing of Pom1 interactions with Cdr2 have remained poorly defined.

Pom1 localizes in a spatial gradient that is enriched at cell tips, with a lower concentration at the medial cell cortex, where its substrate Cdr2 forms nodes (Bähler and Pringle, 1998; Bhatia et al., 2014; Martin and Berthelot-Grosjean, 2009; Moseley et al., 2009; Pan et al., 2014). Thus, the majority of Pom1 and Cdr2 molecules in a cell are spatially separated, raising the question of when and where they interact. Several lines of evidence suggest that the spatial distributions of Pom1 and Cdr2 are critical for the size-dependent signaling properties of this pathway, and suggest the lateral cell cortex as the key interface. For example, ectopic targeting of Pom1 to the medial cell cortex inhibits Cdr2 node formation and Cdr2-dependent cell size signaling (Martin and

Berthelot-Grosjean, 2009). Additionally, the Pom1 gradient is dissipated as part of a
cellular response to glucose deprivation, which leads to increased Pom1 concentration at
the lateral cortex where it delays mitotic entry (Kelkar and Martin, 2015). This result also
demonstrates that Pom1-Cdr2 signaling responds to environmental input to coordinate
cell size with nutrient availability. In this system, Pom1 functions analogously to an
intracellular morphogen, acting as a concentration-dependent and localization-controlled
inhibitor of Cdr2 nodes. However, levels of Pom1 at the medial cell cortex in wild-type
cells growing under steady state conditions are low, constant, and do not vary with cell
size (Bhatia et al., 2014; Pan et al., 2014). The mechanism by which Pom1 provides size
specific input to Cdr2 is unclear and requires analysis of their molecular dynamics at the
lateral cell cortex.

Past studies have led to a model for Pom1 gradient formation driven by the binding
and diffusion of individual Pom1 molecules in the plasma membrane (Hachet et al., 2011;
Saunders et al., 2012). Pom1 binds directly to anionic phospholipids at the cell cortex,
and these interactions are disrupted upon intramolecular autophosphorylation (Hachet et
al., 2011). Polarity factors including Tea1 and Tea4 are enriched at the cell tips by
microtubules, where they recruit the protein phosphatase 1 (PP1) Dis2, which
dephosphorylates Pom1 to promote membrane attachment specifically at the cell tip
(Hachet et al., 2011). Cortical Pom1 molecules are then thought to diffuse in the plane of
the membrane away from the tip, until multiple autophosphorylation events disrupt
membrane binding (Hachet et al., 2011; Hersch et al., 2015; Saunders et al., 2012). Tip-
concentrated Pom1 occludes Cdr1/2 node formation, restricting them instead to the
medial cell cortex (Martin and Berthelot-Grosjean, 2009; Moseley et al., 2009).
Intriguingly, dynamic clusters of Pom1 have also been observed and are proposed to form
by unstable oligomerization of molecules diffusing in the plane of the membrane (Hachet
et al., 2011; Saunders et al., 2012). These clusters are thought to form and decay rapidly,
temporarily sequestering Pom1 molecules and thereby slowing molecular diffusion rates
and reducing noise in the gradient (Saunders et al., 2012). These puncta represent nano-
scale pockets of increased Pom1 concentration, but whether these clusters contribute to
regulation of Cdr2 nodes is unknown.

Here, we use biochemical approaches in combination with quantitative TIRF and confocal microscopy to show that the Pom1 gradient itself is formed by these punctate clusters. Pom1 puncta are stable oligomers that bind and release the membrane with minimal lateral diffusion. The frequency of Pom1 puncta binding is higher at cell tips than on cell sides. Thus, the gradient is formed because the membrane-association frequency of Pom1 puncta is patterned along the long axis of the cell. A portion of Pom1 puncta in the medial cell cortex colocalize with Cdr2 nodes, and the ratio of Pom1 to Cdr2 at the medial cell cortex changes as a function of cell size. When glucose availability is restricted, Pom1 clusters redistribute to the medial cell cortex to prevent dramatic dysregulation of Wee1 bursts. Our work reveals that the Pom1-Cdr2-Wee1 signaling pathway is organized as a series of cortical clusters. The relative distribution of these clusters at the plasma membrane changes with cell size and glucose availability, thus relaying cell surface area to the core cell cycle machinery in a nutrient-controlled manner.

Results:

The Pom1 gradient is formed by transient cortical puncta

We sought to examine the molecular dynamics of Pom1-Cdr2 signaling at the cell cortex. As a starting point, we imaged Pom1 by TIRF microscopy, which selectively excites fluorophores near the coverslip. Pom1 was tagged at the endogenous locus with the bright and photostable fluorophore mNeonGreen (mNG). Surprisingly, along the lateral cell cortex Pom1-mNG was localized almost exclusively in discreet puncta, with no apparent diffuse signal **(Figure 1A)**. In images from a single time-point, Pom1 puncta formed a noisy concentration gradient **(Figure 1B)**. Time-averaging produced a smoother gradient dotted by occasional clusters, similar to the gradient observed by confocal microscopy **(Figure 1A-B)**. Similar results were obtained by confocal microscopy **(Figure S1A-H)**. Pom1 puncta were highly dynamic. They appeared on the lateral cortex with a frequency of ~2 puncta*$\mu m^{-2}s^{-1}$, and remained bound for less than 2 seconds on average **(Figure 1C,F,H)**. During this brief cortical attachment, Pom1 puncta exhibited minimal diffusion which lacked directionality, and instead appeared to diffuse randomly in sub-micron patches of cortex **(Figure 1C,G)**.

Similar puncta were previously observed by confocal microscopy, but they were not thought to form the Pom1 concentration gradient *per se* (Hachet et al., 2011; Saunders et al., 2012). Our findings suggested the distinct possibility that the cortical gradient itself is comprised of dynamic puncta. To test this possibility, we needed to image Pom1-mNG at cell tips, which are inaccessible to visualization using standard slide preparation for TIRF microscopy because this region is outside of the TIRF illumination field. Therefore, we performed "head-on" TIRF imaging of cell tips. Pom1-mNG localized at cell tips almost entirely in puncta, similar to cell sides **(Figure 1D-E)**. We also used "head-on" Airyscan confocal imaging to confirm that Pom1 localization in puncta at tips was not an artifact of the TIRF approach **(Figure S1I)**, a conclusion also supported by work using high resolution wide-field microscopy (Dodgson et al., 2013). Compared to cell sides, Pom1 puncta appeared twice as frequently at cell tips but with a similar cortical duration, again exhibiting minimal and non-directional diffusion away from their binding site **(Figure 1F-H)**. Fluorescence intensity of individual puncta at cell tips and cell sides were similar, although we detected a low number of brighter puncta at cell tips, likely representing multiple diffraction-limited puncta **(Figure 1I, Figure S1I-K)**. From these data, we conclude that the Pom1 gradient is formed by patterning the membrane binding frequency of cortical puncta along the long axis of the cell, rather than by diffusion of Pom1 molecules from the cell tips to the cell side.

In vitro analysis shows Pom1 exists in large, stable puncta

Past work suggested that Pom1 puncta assemble and disassemble at the cortex through oligomerization of individual molecules diffusing in the plane of the membrane (Saunders et al., 2012). However, our high-speed (20ms/frame) continuous TIRF videos suggested that Pom1 puncta bind and release the cortex as a pre-formed unit **(Figure S2A)**. Consistent with this possibility, we observed rare cell lysis events in which Pom1 puncta remained intact in extruded cytoplasm **(Figure 2A)**. This apparent stability while removed from the plasma membrane led us to test the existence and properties of Pom1 puncta in detergent cell extracts. Using TIRF microscopy, we observed puncta of Pom1-mNG in extracts made from Pom1-mNG cells, but not from untagged wild-type cells

(Figure 2B). We next examined the size of these Pom1 puncta by velocity sucrose
gradient sedimentation. By fractionating detergent cell extracts from *pom1-3xHA* cells,
we found that most Pom1 exists in a large 60S complex, consistent with Pom1 puncta
observed in cells. Fractions 6-8 containing this peak were dialyzed and centrifuged on a
second sucrose gradient. These pooled fractions again sedimented at 60S, indicating that
they represent a stable complex **(Figure 2C-D, S2B-C)**.

Pom1 complexes present in detergent extracts resemble the cortical puncta
observed in cells. To test this connection further, we examined the membrane-binding
capacity of Pom1 complexes *in vitro* using artificial supported lipid bilayers (SLBs). Pom1-
mNG puncta were isolated by sucrose gradient centrifugation, incubated with fluid SLBs
on coverslips, and then imaged by TIRF microscopy **(Figure 2E, S2E).** In this cell free-
system, Pom1-mNG puncta bound and released SLB lipids with strikingly similar
dynamics as in cells **(Figure 2E-F)**. Dwell times were increased for the kinase-dead
mutant Pom1(K728R)-mNG, which was previously shown to increase cortical Pom1
levels in cells **(Figure S2D)** (Bähler and Pringle, 1998; Hachet et al., 2011). Collectively,
our TIRF microscopy and *in vitro* analysis of Pom1 puncta support a model whereby the
majority of cellular Pom1 protein is contained within discreet, highly-stable oligomers that
bind and release membranes with kinetics dictated by their catalytic activity in a geometric
pattern that resembles a concentration gradient.

Formation of clusters could represent an intrinsic property of Pom1 protein, or
alternatively might require additional cellular factors. To distinguish between these
possibilities, we expressed and purified GST-Pom1 from bacteria, and then performed
sucrose gradient centrifugation experiments on this recombinant protein. Recombinant
GST-Pom1 sedimented in a low molecular weight peak unlike Pom1 puncta from cells.
Remarkably, purified GST-Pom1 was assembled into a puncta-sized high molecular
weight complex by incubation with wild-type yeast detergent extract **(Figure 2G)**.
Complex formation was also induced by incubation of GST-Pom1 with *pom1Δ* cell
extracts, meaning that additional cellular factors drive this assembly process. **(Figure
2G).** Once assembled, Pom1 puncta are stable oligomeric complexes with intrinsic

membrane-binding properties that can be reconstituted *in vitro*. Additional cellular
regulatory proteins are likely to promote the binding of Pom1 puncta at cell tips, leading
to the spatial gradient.

Polarity landmarks pattern cortical dynamics of Pom1 puncta to shape the gradient
The key role of puncta in forming the Pom1 gradient led us to reexamine the role of
cell polarity landmark proteins Tea1 and Tea4. Both of these proteins localize at cell tips
and are required for proper Pom1 gradient formation (Hachet et al., 2011). However, past
studies have shown that Tea1 and Tea4 have distinct mechanistic roles: Tea4 is required
for localization of Pom1 to the cortex, whereas Tea1 is required for enrichment of cortical
Pom1 to the cell tip (Hachet et al., 2011). We first used sucrose gradient centrifugation to
test if either protein is required for assembly of Pom1 into stable biochemical complexes.
The sedimentation pattern of Pom1 puncta isolated from *tea1Δ* and *tea4Δ* cells in velocity
sucrose gradients is unchanged from wild-type cells, suggesting that neither Tea1 nor
Tea4 are required for assembly of Pom1 into stable biochemical complexes **(Figure 3A)**.
Rather, these regulators are likely to act downstream of puncta assembly.

We next tested the localization and dynamics of Pom1-mNG in *tea1Δ* and *tea4Δ* cells.
Past work showed that membrane bound Tea4 is both necessary and sufficient to
nucleate cortical Pom1 gradients, and this relationship is non-stoichiometric (Hachet et
al., 2011). We confirmed that Pom1 was absent from the cell cortex in confocal
micrographs of *tea4Δ* cells **(Figure 3B)**. Surprisingly, we did observe Pom1-mNG puncta
at the cortex of *tea4Δ* cells by TIRF microscopy **(Figure 3C)**. These binding events
occurred at lower frequency and had a shorter dwell time than in wild-type cells **(Figure**
3C-E). Pom1-mNG puncta in *tea4Δ* cells were distributed evenly around the cell cortex
with no detectable enrichment at tips. These infrequent, transient binding events do not
result in accumulation of enough cortical Pom1 to be detected by confocal microscopy.
Their rapid and position-independent association with the cortex leads to the absence of
a cortical Pom1 gradient in *tea4Δ* cells.

In contrast, confocal micrographs of Pom1-mNG in *tea1Δ* cells confirmed even

distribution around the entire cell cortex **(Figure 3B)**. TIRF microscopy of *tea1Δ* cells

revealed that this distribution arises from Pom1-mNG puncta, which bound the cell cortex

at a frequency in between the tips versus sides of wild-type cells **(Figure 3C-D)**. The

duration of individual binding events was longer than in *tea4Δ* cells but shorter than in

wild-type cells **(Figure 3E, S3A)**. Thus, in *tea1Δ* cells Pom1 puncta bind at an

intermediate frequency and for an intermediate duration, when compared to wild type and

tea4Δ cells. These binding events are independent of spatial cues at the cell ends versus

cell sides, resulting in enrichment of cortical Pom1 but not a gradient. Thus, Tea1 and

Tea4 cooperate to promote localized membrane binding but not assembly of Pom1

puncta. These spatial cues pattern the frequency of Pom1 puncta cortical interaction

along the long axis of the cell.

Pom1 puncta interact with Cdr2 nodes at the lateral cortex

It has been unclear when and where Pom1 interacts with its inhibitory target Cdr2,

which localizes in cortical nodes positioned in the cell middle. Since Pom1 puncta appear

to bind throughout the medial cell cortex, we used simultaneous two-color TIRF

microscopy to test colocalization of Pom1 puncta and Cdr2 nodes. We observed

unambiguous overlap between Cdr2 nodes and some Pom1 puncta, as well as Pom1

puncta that bound to the cortex without encountering a Cdr2 node **(Figure 4A)**. Thus,

Pom1 puncta bind to the medial cortex in a stochastic pattern that can generate overlap

with Cdr2 nodes. These patterns of colocalization were apparent in both static images

and time-lapse movies **(Figure 4A-B)**. The frequency, dwell time, and displacement of

Pom1 cortical puncta were unaffected in *cdr2Δ* cells **(Figure S4A-DC)**. These data

suggest that Pom1 interacts with its inhibitory phosphorylation target Cdr2 when Pom1

clusters associate with Cdr2 nodes at the medial cell cortex.

Phosphorylation by Pom1 prevents activation of Cdr2 in small cells, thereby

contributing to cell size-dependent regulation of Wee1 and mitotic entry (Allard et al.,

2018). To place this pathway in the context of Pom1 puncta and Cdr2 nodes, we analyzed

how these structures accumulate and colocalize as cells grow. Intriguingly, the total

number of both Cdr2 nodes and Pom1 puncta detected in TIRF images all along cell sides

were equivalent for cells of a given size **(Figure 4C-D)**. The numbers of both structures

show a similar size-dependent doubling. We next restricted our analysis to a 2µm x 2µm

square ROI positioned at the cell middle, where Cdr2 nodes concentrate. The local

density of Cdr2 nodes in this region increased as a function of cell size, consistent with

past studies (Deng and Moseley, 2013; Pan et al., 2014). In contrast, the density of Pom1

puncta in this region was largely independent of cell size **(Figure 4E)**. In both TIRF and

confocal images, the concentration of Pom1 protein and Pom1 clusters decreased slightly

as cells increase in size, but this trend was dwarfed by the cell size-dependent increase

in Cdr2 node density **(Figure 4E, S4D)**. Thus, as cells grow larger, the ratio of Pom1 to

Cdr2 in the medial cell cortex changes to favor Cdr2 because the densities of these two

structures scale differently with cell size **(Figure 4G)**.

We next tested how this density scaling affects colocalization of Pom1 clusters and

Cdr2 nodes as cells grow. Due to the increasing density of Cdr2 nodes, we observed an

increased number of colocalized Pom1 clusters and Cdr2 nodes as cells increase in size

(Figure SE-F). The increased colocalization, combined with the constant density of Pom1

clusters, means that there are less "free" Pom1 clusters as cells increase in size **(Figure**

S4G). Perhaps more importantly, the number of "free" Cdr2 nodes increased as a function

of cell size **(Figure S4G)**. These free Cdr2 nodes are not occupied by an inhibitory Pom1

cluster, and thus have increased potential to promote mitotic entry by inhibiting Wee1. In

this manner, Pom1 sets an inhibitory threshold that must be overcome by an increase in

Cdr2 node density. The inhibitory threshold decreases slightly as cells grow, and

functions most effectively in small cells, where Pom1 was previously shown to inhibit

downstream signaling to Wee1 (Allard et al., 2018).

Control of Pom1 puncta levels in the medial cell cortex

Our results suggest that Pom1 sets an inhibitory threshold that can be tuned by

altering the concentration of Pom1 cortical clusters in the cell middle. We sought to test

this model by increasing the abundance of Pom1 clusters in the cell middle. Low glucose

media and *tea1∆* are two independent conditions that disperse Pom1 throughout the

cortex, thereby increasing its concentration in the cell middle (Hachet et al., 2011; Kelkar and Martin, 2015) (See also **Figure S7A-C**). As a starting point, we tested the effect of mislocalized Pom1 in *tea1Δ* cells **(Figure 5A-B)**. Surprisingly, higher Pom1 levels at the medial cortex did not impact the number of Cdr2 nodes or their recruitment of Wee1 **(Figure 5C-F)**. Past work showed that Pom1 retains kinase activity in *tea1Δ* cells (Bähler and Nurse, 2001), but several experiments suggested that this residual activity may be reduced. First, we compared *in vitro* kinase activity of Pom1 immunoprecipitated from *tea1Δ* cells or wild type cells. Pom1 from *tea1Δ* cells showed less kinase activity towards itself (autophosphorylation) and its substrate Cdr2 **(Figure S5A-C)**. Second, phosphorylation of the Pom1 substrate Rga7 was reduced in *tea1Δ* cells **(Figure S5D-E).** Third, Tea4 is completely absent from the cortex of *tea1Δ* cells **(Figure S6D)**. Catalytically active Pom1 autophosphorylates and requires Tea4-dependent dephosphorylation for cortical enrichment. Thus, in *tea1Δ* cells, Pom1 likely binds to the membrane because it cannot autophosphorylate properly. Finally, we note that Pom1 localization in *tea1Δ* mimics the localization of kinase-dead *pom1-m2* mutant **(Figure S6A-B)**. Therefore, we focused on the activity and regulation of Pom1 clusters in wild type cells grown in low glucose.

Fission yeast cells respond to low glucose in part by depolymerizing their microtubule cytoskeleton, which leads to Pom1 redistribution to the lateral cell cortex where it delays mitotic entry (Kelkar and Martin, 2015). In time-lapse TIRF microscopy experiments, we observed increased numbers of discreet Pom1 puncta at the lateral cell cortex under low glucose conditions **(Figure 6A)**. These puncta bound to the lateral cortex more frequently than under normal glucose conditions, but the average dwell time was unaffected **(Figure 6B-C)**. This increased on-rate leads to accumulation of Pom1 clusters at the medial cell cortex.

We next investigated how increased numbers of Pom1 puncta at the lateral cell cortex affect Cdr2 nodes using TIRF microscopy. In low glucose conditions, the number of Cdr2 nodes was largely unchanged, but the fluorescence intensity of individual Cdr2 nodes and the overall concentration of Cdr2 at the medial cortex was significantly lower

(Figure 6D-F, S7A-C). In *pom1Δ* cells, which grow only from one tip, Cdr2 nodes form

normally but invade the non-growing tip. Low glucose conditions do not measurably affect

Cdr2 nodes in *pom1Δ* cells, so the effects that we observed are Pom1-dependent **(Figure**

6D-F). These results indicate that Pom1 relocalization to the lateral cortex induces partial

disassembly of Cdr2 nodes under low glucose. To quantify how this effect scales with

glucose availability, we varied the concentration of media glucose over four orders of

magnitude. Then, we measured the concentrations of both Pom1 and Cdr2 at both the

cell sides and the cell tips. As glucose was reduced, Pom1 concentration decreased at

cell tips and increased at cell sides, consistent with redistribution **(Figure 7A,C-D)**. In

contrast, Cdr2 concentration increased at cell tips and decreased at cell sides **(Figure**

7B-D). Plots of glucose availability versus Pom1 or Cdr2 concentration produced

sigmoidal curves with opposite orientations **(Figure 7C-D)**. Correlation analysis of Pom1

and Cdr2 concentrations at cell sides show they are nearly perfectly anti-correlated

(Pearson's r=-0.9352, p=0.0006) **(Figure 7E-F)**. We conclude that the increasing

frequency of Pom1 cluster binding at the lateral cell cortex leads to progressive disruption

of Cdr2 nodes.

Pom1 disrupts Wee1 bursting under glucose deprivation

We next examined the downstream effects of altered Pom1-Cdr2 signaling by

using TIRF microscopy to monitor the previously described bursts of Wee1-mNG

localization at Cdr2 nodes (Allard et al., 2018). In wild-type cells, both the frequency and

duration of Wee1 bursts increased linearly with cell size, consistent with increased

inhibition of Wee1 as cells grow **(Figure 8A,B)**. In low glucose, the frequency of Wee1-

mNG bursts still scaled with cell size as in normal glucose. However, the duration of each

burst was uniformly short and independent of cell size **(Figure 8A-B)**. These same burst

properties were seen for Wee1-mNG bursts in kinase-dead *cdr2(E177A)* mutant cells

grown in normal glucose **(Figure S8A-B)**. These results indicate that Cdr2 kinase activity

is required for retention of Wee1 at nodes, but not for the initial binding event. Further,

they demonstrate that low glucose phenocopies the kinase-dead mutant, consistent with

Pom1-dependent inhibition of Cdr2 kinase activity.

These results suggested that attenuation of Wee1 bursting dynamics in low
glucose might require Pom1 redistribution. We tested this idea by examining the glucose
dependency of Wee1 node localization in *pom1∆* cells. In *pom1∆* cells growing under high
glucose, the Wee1 burst frequency scales with cell size but Wee1 burst duration is
uniformly high and independent of cell size **(Figure 8A,C)**. Unlike wild type cells, which
suppress Wee1 bursts in low glucose, *pom1∆* cells show a striking increase in Wee1
bursts under low glucose. This increase is not cause by burst duration, which was
independent of both cell size and glucose concentration in *pom1∆* cells. Rather, we
measured increased Wee1-mNG burst frequency in low glucose *pom1∆* cells **(Figure
8A,C)**. The reason for this increase is unknown, and suggests the existence of additional
glucose-regulated pathways that operate on Cdr2-Wee1 signaling.

Wee1 localization at Cdr2 nodes leads to its inhibitory phosphorylation by Cdr2
and the related kinase Cdr1. To test how our TIRF-based localization results connect with
Wee1 phosphorylation status, we analyzed phosphorylation-dependent shifts of Wee1
migration using western blots **(Figure 8D)**. In high glucose, Wee1 migrates as a smear
of phosphorylated isoforms (Allard et al., 2018; Lucena et al., 2017). The upper,
hyperphosphorylated forms are increased in *pom1∆* cells but absent in *cdr2∆* cells. This
hyperphosphorylated Wee1 is lost in wild-type cells grown in low glucose, consistent with
reduced bursts of Wee1 localization to nodes. In contrast, Wee1 appears even more
hyperphosphorylated in *pom1∆* cells grown under low glucose. This result suggests that
Pom1 prevents phospho-inactivation of Wee1 in response to low glucose. These
combined experiments support a model where Pom1 redistribution to the lateral cell
cortex inactivates Cdr2 nodes to relieve inhibition of Wee1 under low glucose conditions.

Discussion:

In this study, we have shown that the Pom1 concentration gradient is generated
by oligomeric Pom1 puncta that rapidly bind and release from the plasma membrane.
These puncta bind more frequently at cell tips versus cell sides in high glucose media,
while their binding rate is increased at cell sides in low glucose media. At the medial cell
cortex, these puncta overlap with their inhibitory target Cdr2, which localizes in static

oligomeric nodes. More Cdr2 nodes are free from Pom1 inhibition as cells grow larger
due to different density scaling of these two structures at the medial cell cortex. Our TIRF-
based colocalization experiments on these structures were limited to short periods of time
to avoid photobleaching, partly caused by the imaging conditions needed to observe
these highly dynamic Pom1 clusters. Nevertheless, even over these short timescales of
observation, the stochastic positioning of Pom1 puncta resulted in spatiotemporal
colocalization with Cdr2 nodes. Given the high frequency of Pom1 puncta binding, we
expect that many more colocalization events occur between Cdr2 nodes and Pom1
clusters during the course of a full cell cycle, with the potential for Pom1 clusters to visit
each Cdr2 node multiple times. Thus, these rapid dynamics of Pom1 puncta interacting
with Cdr2 nodes must be integrated in time throughout the cell cycle, leading to the G2/M
transition. It appears likely that the most critical temporal window for Pom1-Cdr2
interactions occurs during early in G2, as *pom1Δ* show defects in Cdr2-Wee1 signaling
specifically in small cells (Allard et al., 2018). It will be interesting to determine how these
rapid interactions in small cells are integrated into the mitotic entry decision, which occurs
later in the cell cycle when cells are longer.
419
420   **Molecular clusters are widespread in signal transduction**
421   The Pom1-Cdr2-Wee1 signaling pathway appears to function entirely within the
confines of oligomeric protein clusters at the plasma membrane. The development of
imaging technologies with increased spatial and temporal resolution has enabled the
discovery of protein clustering as a paradigm in signaling. As a result, a growing number
of signaling events and proteins have been found in similar clustered structures that are
referred to as nanodomains, clusters, nanoclusters, and nodes, among other names. For
example, recent studies have shown how PAR-3 polarity complexes in the *C. elegans*
zygote form clusters to reduce random diffusion and promote vectorial transport
(Dickinson et al., 2017; Munro, 2017; Rodriguez et al., 2017; Wang et al., 2017). In
addition, major signal transduction pathways in plants are organized into protein clusters
on plasma membrane nanodomains (Gronnier et al., 2017). Similarly, different Ras
isoforms partition into distinct and exclusive clusters that digitize downstream signals in
animal cells (Zhou and Hancock, 2015). In immune cells, both B-cell receptors and T-cell

receptors form oligomeric compartments at the cell cortex (Douglass and Vale, 2005; Liu et al., 2016; Maity et al., 2015). These T cell receptor clusters enrich downstream kinases while excluding phosphatases (Su et al., 2016), a mechanism that could relate to concentrated clusters of the kinases Pom1 and Cdr2. Clustering can have a range of critical impacts on the signaling properties of these pathways, such as the clustering of bacterial chemotactic receptors to generate signal amplification (Bray et al., 1998; Duke and Bray, 1999; Falke, 2002). Theoretical work has suggested that clustering can reduce noise by generating reaction bursts (Kalay et al., 2012), which have the capacity to overcome inhibitory thresholds, such as those imparted by phosphatases, more effectively than a system driven by gradual accumulation of signal. Insight into the functional role of clustering for Pom1-Cdr2-Wee1 will require additional *in vivo* analysis combined with *in vitro* reconstitution. We have shown that Pom1 clusters remain stable through biochemical enrichment, and isolated clusters exhibit the same membrane-binding properties *in vitro* as in cells. Past work has demonstrated Cdr2 nodes also remain stable through biochemical fractionation (Allard et al., 2018), providing the necessary tools for future dissection of the pathway in a reconstituted system.

A new model for cortical gradient formation

Our work has shown that Pom1 clusters bind to the cell cortex with different frequencies along the long axis of the cell, resulting in a concentration gradient. Past work suggested single Pom1 molecules (or small oligomers) bind to the membrane at tips, and then diffuse directionally away from the tip before dissociating (Hachet et al., 2011; Saunders et al., 2012). Clusters were proposed to dynamically assemble and disassemble at the cortex from this diffusing population (Saunders et al., 2012). Our findings suggest a new model for gradient formation, which is distinct from this previous model in four ways. First, we found that clusters themselves form a concentration gradient that emanates from the cell tips. Second, our combined imaging and biochemical approaches demonstrated that clusters are stable complexes that bind and rapidly release from the membrane. Third, we did not observe extensive diffusion by complexes bound to the membrane. It remains possible that smaller, diffusing Pom1 complexes or single molecules were not detected by our imaging approach, but we favor a model where

diffusion is limited to short, non-directional movements that do not contribute to gradient

formation. Fourth, the key step in gradient formation is a positional system that increases

the on-rate for Pom1 clusters at cell tips versus the cell middle. Concentration gradients

are found across vast size scales in biology, so the mechanism of gradient formation by

Pom1 clusters may have broad implications. It is also important to note that the spatial

gradient of on-rates for Pom1 clusters binding to the cortex may depend on a

concentration gradient of specific lipids in the plasma membrane. In support of this

hypothesis, Pom1 binds to phosphatidylserine *in vitro* (Hachet et al., 2011), and this lipid

is enriched at the tips of growing fission yeast cells (Haupt and Minc, 2017). Thus, further

investigation of these lipids and their localization dynamics could reveal additional layers

of this morphogen gradient.

We found that the polarity landmarks Tea1 and Tea4 provide the positional

information for Pom1 clusters to form a concentration gradient. Past work has shown that

Pom1 localizes homogenously throughout the cortex in *tea1Δ*, and does not localize to

the cortex in *tea4Δ* (Bähler and Pringle, 1998; Hachet et al., 2011). Based on these and

other results, Tea1 was proposed to recruit Tea4 to cell tips, where it promotes cortical

loading of Pom1 molecules. Several results suggest that additional mechanisms may

contribute to gradient formation. For example, we found that Pom1 clusters still bind to

the cortex in *tea4Δ* cells but for very short durations. Past work has also shown that Tea1

and Tea4 form cortical clusters that co-localize at cell tips, but these Tea1-Tea4 clusters

do not co-localize with Pom1 clusters (Dodgson et al., 2013). Thus, Tea1-Tea4 may

position binding of Pom1 clusters to the cortex through an additional mechanism. For

example, Tea1 and Tea4 have been shown to assemble sterol-enriched lipid domains at

cell tips when cells initiate polarization following exit from starvation (Makushok et al.,

2016). Sterols and phosphatidylserine are thought to form a common membrane domain

in fission yeast, suggesting that these proteins could act through the recruitment and

organization of lipids to generate cortical domains permissive for Pom1 cluster binding

(Haupt and Minc, 2017; Makushok et al., 2016).

Adjustable Pom1 gradients regulate mitotic entry

We propose an integrated mechanism for fission yeast cell size sensing in which Cdr2 acts as an important sensor of cell size that is dynamically antagonized by cortical Pom1 **(Figure 9)**. Pom1 puncta bind to the medial cell cortex in a manner that can overlap with Cdr2 nodes. These dynamic Pom1 puncta and stable Cdr2 cortical nodes suggest a system that monitors surface expansion, consistent with the notion that fission yeast cells monitor surface area as the primary determinant of size at division (Facchetti et al., 2019; Pan et al., 2014). As cells grow, the density of Pom1 clusters in the cell middle decreases slightly due to their binding preference at cell tips. In contrast, the density of Cdr2 nodes doubles to overcome this inhibitory Pom1 threshold. The density of Pom1 clusters and the resulting inhibitory threshold can be modulated by glucose availability, consistent with past work (Kelkar and Martin, 2015). As fission yeast cells begin to starve due to depletion of nutrients such as glucose, their length at division progressively decreases and the length of the cell cycle increases (Pluskal et al., 2011; Yanagida Mitsuhiro et al., 2011). The Pom-Cdr2 network has a very prominent role in this nutrient modulation, suggesting that it may represent a key physiological role for the pathway (Kelkar and Martin, 2015). We found that the size dependence of Wee1 localization bursts to Cdr2 nodes, which facilitate Wee1 inhibition, was dampened in low glucose conditions. Re-localization of Pom1 clusters to the medial cortex likely mediates this response. However, for *pom1∆* mutants in low glucose, Wee1 localization to nodes exceeds even wild type cells in high glucose. This result indicates that additional mechanisms contribute to Wee1 regulation in low glucose, for example Wee1 nuclear transport could be altered under these conditions.

The Pom1 gradient as a read-out for cell polarity

More broadly, our study reveals the adjustable nature of the concentration gradient formed by Pom1 clusters. Concentration of Pom1 clusters at cell tips reads out the polarity state of the cell for downstream signaling by Cdr2 and Wee1. Conditions that alter the polarized growth state of the cell would lead to changes in the gradient of Pom1 clusters, resulting in altered cell size at division. This dynamic system appears to operate in a manner that depends on both cell size and nutrient availability. Identification of additional

growth and environmental conditions that alter the relative distributions of Pom1 clusters

and Cdr2 nodes, and determination of how they connect with other cell cycle signaling

pathways, may reveal new mechanisms for nutrient modulation of cell size control.

Integration of the Pom1-Cdr2-Wee1 pathway with other sizing mechanisms

Pom1-Cdr2-Wee1 signaling is cell size dependent but is not the only pathway that

contributes to the overall cell size control network. Intriguingly, a recent study reported

that *cdr2Δ* cells no longer divide based on accumulation of a threshold surface area and

instead revert to a secondary mechanism of cell size control based on volume sensing.

These results demonstrate that cells use different signal transduction networks to

measure different aspects of their size. Whereas the Pom1-Cdr2 network is well suited to

measure cortical surface area, size-control systems that measure other aspects of size

would require different signaling logic. Theoretically, mechanisms for cell volume sensing

could depend on proteins whose concentrations do not scale linearly with cell size. Such

mechanisms are thought to underlie cell size control in other organisms, such as the

budding yeast, where growth-dependent dilution of the transcriptional repressor Whi5 is

thought to underlie G_1/S size control (Schmoller et al., 2015). One possible candidate for

cellular volume sensing in fission yeast is Cdc25, which increases in concentration with

cell size (Keifenheim et al., 2017; Moreno et al., 1990). These mechanisms are attractive

candidates for volume sensors because cytoplasmic concentration is independent of cell

geometry. The existence of sizing mechanisms that measure other aspects of cell size,

and function independently of the Pom1-Cdr2 surface area sensing network, underscores

that cell size is a systems level property controlled by multiple signaling pathways. Thus,

it remains important to discover size-dependent signaling mechanisms within each

individual pathway, as a step towards understanding the integration of multiple pathways

into a larger size control system.

Materials and Methods:

Strain Construction and Media

Standard *S. pombe* media and methods were used (Moreno et al., 1991). Strains used in this study are listed in Supplementary Table S1. Gene tagging and deletion were performed using PCR and homologous recombination (Bähler et al., 1998). The mNeonGreen (mNG) sequence was used under license from Allele Biotechnology.

TIRF Microscopy and Analysis

Pom1 puncta and node components were imaged using simultaneous dual-color total internal reflection fluorescence microscopy to limit excitation of fluorophores to those nearest to coverslip. Imaging was performed on a commercially available TIRF microscope (Micro Video Instruments) composed of a Nikon Eclipse Ti microscope base equipped with a 100x Nikon Apo TIRF NA 1.49 objective and a two-camera imaging adaptor (Tu-CAM, Andor Technlogy) containing a dichroic and polarization filters (Semrock FF580-FDi01-25x36, FF02-525/40-25, FF01-640/40-25) to split red and green signal between two aligned Andor iXon electron-multiplied CCD cameras (Andor Technology). Red/green beam alignment was performed prior to imaging using a TetraSpeck Fluorescent Microsphere size kit (Thermofisher).

Standard #1.5 glass coverslips were RCA cleaned before use to remove fluorescent debris. Cells were grown in EMM4S, and washed into fresh EMM4S immediately before imaging to remove auto-fluorescent debris resulting from overnight culture. Cells were imaged in EMM4S media on glass slides at ambient temperature. Individual slides were used for no more than five minutes to prevent cells from exhausting nutrients or oxygen. Agar pads were not used due to increased background fluorescence.

Image analysis and processing was performed using ImageJ2 (NIH). Cdr2 node numbers and Pom1/Wee1 puncta number, frequency and binding duration were quantified using the Trackmate plugin (Tinevez et al., 2017). Due to variable fluorescence intensity in different TIRF fields and images, thresholding parameters were determined separately for each image, and accuracy was confirmed by visual inspection to ensure

that only nodes/puncta were counted and that no nodes/puncta were omitted. For Wee1

burst and Pom1 puncta tracking, Particle Diameter was set to 0.3 microns (approximate

lateral resolution), with Maximum Gap Linking set to 2 frames, and Linking Range for

particle tracking was set to 0.15 microns. Lookup table was adjusted to fire for some

images to emphasize signal intensities.

"Head-On" Tip Imaging

"Head-On' or "Tip" imaging was performed using a protocol modified from

(Dodgson et al., 2013). Custom micro-well coverslips or Ibidi Sticky-Slide VI channels

(Ibidi, Martinsried, Germany) were coated with BS-I lectin (1mg/mL in water) by incubation

for 30 minutes and washed 3X with water to remove non-adherent lectin. Cells were then

added to the imaging chambers and allowed to settle to the cover-slip bottom where they

adhere to the lectin. Cells oriented with the long axis perpendicular to the coverslip were

identified using brightfield prior to fluorescence imaging.

Spinning Disc Microscopy and Analysis

Spinning-disc confocal imaging was performed using a commercially available

system (Micro Video Instruments, Avon, MA) featuring a Nikon Eclipse Ti base equipped

with an Andor CSU-W1 two-camera spinning disc module, dual Zyla sCMOS cameras

(Andor, South Windsor, CT) an Andor ILE laser module, and a Nikon 100X Plan Apo ⋏

1.45 oil immersion objective. Cells were imaged in EMM4S media on glass slides at

ambient temperature unless otherwise noted.

Airyscan Super-Resolution Microscopy and Analysis

To achieve maximum resolution and sensitivity, fluorescence intensity of Pom1

puncta on both cell sides and cell tips was measured using a Zeiss Airyscan microscope

(Figure S1I-K), composed of a Zeiss LSM-880 laser scanning confocal microscope

(Zeiss, Oberkochen, Germany) equipped with 100X alpha Plan-Apochromat/NA 1.46 Oil

DIC M27 Elyra objective, Airyscan super-resolution module and GaAsP Detectors, and

Zen Blue acquisition software using the Super-resolution mode with pin-hole size of 1.2

airy-units to prioritize resolution. Z-volumes of 16 slices with 0.19μm spacing for high

spatial resolution in all dimensions were centered on the cell cortex closest to the

coverslip. Airyscan images were processed in Zeiss Zen Blue software, and quantification

was performed on sum projections of Airyscan reconstructed stacks.

Pom1 and Cdr2-CTD Purification

The full length Pom1 sequence, and the Cdr2-CTD fragment were purified as

follows: Each sequence was subcloned into pGEX6P1 vector using the Xho1 restriction

site and expressed as a GST-fusion protein in *Escherichia coli* strain BL21(DE3).

Transformants were cultured to log-phase at 37˚C, followed by a shift to 25˚C for 30

minutes. Expression was then induced by addition of 1-thio-β-D-galactopyranoside to

200µM, followed by growth for an additional 3 hours at 25˚C. Cells were harvested by

centrifugation and lysed by passing them twice through a French press in lysis buffer

(1xPBS, 100mM NaCl, 1mM DTT, and complete EDTA-free protease inhibitor tablets (1

tablet/50mL Buffer) (Roche, Basel, Switzerland)). Following lysis, Triton-X 100 was added

to 1% V/V. Lysates were then cleared by centrifugation for 10 minutes at 12,000 x g at

4˚C in a Sorval SS-34 fixed-angle rotor (Thermo Scientific, Waltham, MA). The

supernatant was then incubated with glutathione-agarose (Sigma Aldrich, St. Louis, MO)

for 2h at 4˚C. 20 mM glutathione (pH 8.0) was used to elute purified protein, and the

eluate was dialyzed overnight at 4˚C into 10mM Tris-HCl, pH8 + 5% V/V glycerol.

Preparation of Yeast Extracts

Fission yeast detergent extracts were prepared by growing 1.5L cells to mid-log

phase (OD ~0.3), and then washed twice with 50mL Node Isolation Buffer (NIB – 50mM

HEPES, 100mM KCl, 1mM $MgCl_2$, 1mM EDTA, pH 7.5) (Allard et al., 2018). Next, we

resuspended the pellet in an equal volume of 2X NIB (W/V) containing a

protease/phosphatase inhibitor cocktail (10µL/mL 200x PI, 50µL/mL 1M β-glycerol

phosphate, 50µL/mL 1M NaF, 2.5 µL/mL 200mM PMSF, 1mM DTT), and snap froze the

resuspension as pellets by pipetting drop-wise into a liquid nitrogen bath. Then, yeast

pellets were ground using liquid-nitrogen chilled coffee-grinders for 2 min, and collected

into chilled falcon tubes and stored at -80˚C. 1.5g of frozen yeast powder was then thawed

on ice, and Triton X-100 was added to a final concentration of 1%, and the extracts were

mixed by gentle pipetting. Extracts were then centrifuged at 4˚C for 10min at 20,000 x g to yield a low speed supernatant, which were then used for subsequent experiments.

Sucrose Gradient Ultracentrifugation

Discontinuous sucrose gradients were prepared in 14 x 89mm Ultra Clear Ultracentrifuge tubes (Beckman Coulter, Brea, CA) by layering 5 – 23% (top to bottom) sucrose in NIB + 1% Triton X-100 in 0.37mL steps of 2% sucrose increments. 700µL of yeast extract or recombinant protein diluted in NIB Buffer was then added to the top of the gradient. Sucrose gradients were centrifuged at 100k x g (50kRPM) in a Beckman L8-M ultracentrifuge for 2hr at 4˚C in a chilled SW60ti swinging bucket rotor. 0.5mL gradient fractions were collected from the top by hand, vortexed, and 100µL of each fraction was mixed 2:1 in 3X SDS-PAGE sample buffer (65 mM Tris pH 6.8, 3% SDS, 10% glycerol, 10% 2-mercaptoethanol, 50 mM NaF, 50 mM β-glycerophosphate, 1 mM sodium orthovanadate) and boiled for 5 min. Samples were then subjected to SDS-PAGE and western blotting or Coomassie staining. To calculate S-values of sedimentation peaks from western blot signal intensities, mean band intensities were measured using Image Studio Lite (LICOR, Lincoln, NE), and Gaussian Curves were fit to these values. The peak of the Gaussian for each sedimentation peak of each protein was used to assign the known or interpolated S-value.

Western Blotting

For western blots, cells were lysed in 150 µl of 3X SDS-PAGE sample buffer with glass beads in a Mini-beadbeater-16 (Biospec, Bartlesville, OK) for 2 minutes. Gels were run at a constant 20 mAmps until 75kDa marker was at the bottom of the gel. Blots were probed with anti-HA (Covance, Princeton, NJ) or homemade anti-GST. For monitoring Wee1 phosphorylation, samples were run on an SDS-PAGE gel containing 6% acrylamide and 0.02% bisacrylamide, and then probed with a homemade anti-Wee1 antibody (Allard et al., 2018).

Supported Lipid Bilayers

To prepare supported lipid bilayers, we first prepared small unilamellar vesicles
(SUVs) composed of three lipids: (1) DOPC (18:1 (Delta9) Cis PC - 1,2 Dioleoyl-sn-
glycero-3-phosphocholine), (2) DOPS (18:1 PS - 1,2-Dioleoyl-sn-Glycero-3-[Phospho-L-
Serine]), and (3) fluorescent 18:1 Liss Rhod PE (1,2-dioleoyl-sn-glycero-3-
phosphoethanolamine-N-(lissamine rhodamine B sulfonyl). All lipids were dissolved in
chloroform (Avanti, Alabaster, AL). 2µmol total lipid with the desired molar ratios were
mixed in glass vials with PTFE coated caps, which had been RCA cleaned and rinsed
three times with chloroform before use. Excess chloroform was evaporated in a vacuum
desiccator for 1 hour. Lipids were resuspended in 400µL SUV Buffer (50mM HEPES,
100mM KCl, 1mM EDTA, 1mM MgCl$_2$, pH7.5) for 5mM lipid mixture stocks. The lipid
mixtures were vortexed until cloudy, and then transferred to 1.5mL Eppendorf tubes and
subjected to ten freeze-thaw cycles using a liquid nitrogen bath and 32°C hot plate, with
2 minutes of sonication in a sonicating water bath following each thaw cycle. Stocks were
stored at -80°C in 20µL aliquots.
695
Supported lipid bilayers were made by adding 10µL of SUVs to RCA cleaned
custom microwells built on 22x40mm #1.5 coverslips. Chambers were then incubated at
37°C for one hour to induce vesicle fusion, and unincorporated vesicles were rinsed away
using five rinses with 100µL SUV buffer. Yeast extract was added to these chambers by
removing 90µL of buffer, leaving just enough to cover the chamber bottom, and then
adding extract.
702

Pom1 Immunoprecipitation

Cultures were grown in EMM4S media to log phase, and a volume equivalent to
200 optical density units were harvested by centrifugation. The pellet was washed in
Wash Buffer (1X PBS, 250mM NaCl, 1mM EDTA, 1mM PMSF, 50mM NaF, 50mM β-
glycerophosphate, 1mM sodium orthovanadate, and 1% Triton X-100, and complete
EDTA-free protease inhibitor cocktail (1 tablet/50mL) (Roche)). The pellet was then
resuspended in 5mL of wash buffer, and divided into 1.5mL screw-cap tubes each with
200µL glass beads for lysis by bead-beading for 2 minutes using a mini-Beadbeater-16
(Biospec) at 4°C. Lysates were cleared by centrifugation at 16,000 x g for 10 minutes at

4˚C in an Eppendorf 5415C centrifuge. The supernatant was then incubated with rotation
at 4˚C for 1 hour with Pierce anti-HA magnetic beads (Thermo Scientific). Beads were
washed five times for 15 minutes each in Wash Buffer prior to use. Beads were
resuspended in Kinase Assay Buffer (see below) for immediate use in kinase assays.
716
***In vitro* kinase assays**
For *in vitro* kinase assays, immunoprecipitated Pom1 proteins and purified Cdr2-
CTD were combined at a 1:1 ratio in Kinase Assay Buffer (20mM Tris pH 8, 100mM NaCl,
50mM NaF, 50mM β-glycerophosphate, 1mM sodium orthovanadate, 10mM $MgCl_2$,
20μM ATP-γ-S) for 30 minutes at 30˚C. The reaction was halted by addition of EDTA to
20mM. For subsequent analysis by anti-thiophosphate western blot, proteins in *in vitro*
kinase assays were then alkylated by addition of p-nitrobenzyl mesylate to 2.5mM, and
incubation at 25˚C for one hour. Samples were then prepared for western blot as
described above. Thiophosphate western blots were probed with an antibody against
thiophosphate-ester (Abcam 92570) (Abcam, Cambridge, United Kingdom).
727
728
729
730 # Figure Legends:
731
Figure 1: **The Pom1 gradient is formed by time-averaging of puncta that transiently**
bind the cortex with different dynamics along the long axis of the cell. (A) Individual
frame (top panel) and sum projections (bottom panel) of high-speed TIRF microscopy
movie. Scale bar 1μm. Movie was continuous 15s time-lapse acquisition of 200ms
exposures. Blue arrows mark position of line scans performed for data in panel B. **(B)**
Line scans of fluorescence intensity along the long axis of the snap shot and projection
images in panel (A). Note that time-averaging of Pom1 puncta smoothens the
concentration gradient. **(C)** Traces of single-particle diffusion paths generated from a 15s
time-lapse TIRF microscopy acquisition of Pom1-mNeonGreen. Scale bar 1μm. **(D-E)**
Pom1-mNeonGreen forms puncta both at the cell tip and cell side. Images are sum
projections of three consecutive 200ms exposures from continuous time-lapse TIRF

microscopy acquisitions of either the cell tip or side as depicted in the cartoon diagrams.
Scale bar 1µm. **(F-H)** Particle-tracking analysis of 2µm x 2µm ROIs from 30s continuous
time-lapse TIRF acquisitions of Pom1-mNeonGreen at either the cell tip or side was used
to measure and compare **(F)** the binding frequency of Pom1 puncta (****, *p*=<0.0001),
(G) the average displacement of individual puncta (***, *p*=0.0002), **(H)** the binding
duration of individual puncta (n.s., *p*=0.6747). Each data point represents a single cell
mean, and line and error bars represent mean and standard deviation of all cells.
Statistical significance was tested using a Student's T-test. **(I)** Histogram of the
fluorescence intensity of individual Pom1 puncta at the cell tip and cell side. Puncta were
binned by fluorescence intensity and displayed as a population frequency.

Figure 2: **Pom1 puncta are stable structures that can be isolated *in vitro*. (A)** TIRF
microscopy image reveals fluorescent puncta in the extruded cytoplasm (white dotted
line) from a lysed Pom1-mNeonGreen cell (yellow dotted line). Scale bar 5µm. **(B)** TIRF
microscopy images of cell extracts prepared from wild-type (no tag) or Pom1-
mNeonGreen cells. Images are 50 frame sum projections of continuous 200ms time-lapse
exposures. The two images were contrasted equally. Scale bar 1µm. **(C)** Cytoplasmic
extracts of *pom1-3HA* cells were subjected to velocity sucrose gradient sedimentation,
and fractions were probed against the HA tag (upper blot). Fraction 1 corresponds to the
top of the gradient and contains smaller structures; fraction 12 corresponds to bottom of
the gradient and contains larger structures. Fractions 6-8 were pooled, sucrose was
removed by dialysis, and then the sample was subjected to a second identical round of
sucrose gradient sedimentation and western blotting of the resulting fractions (lower blot).
(D) Quantification of Pom1-3HA band intensities from western blot of yeast extracts in
panel C. Each apparent peak was fit to a gaussian curve, and S-values were determined
using size-standards run on identical gradients. **(E)** TIRF microscopy of Pom1-mNG
puncta from cytoplasmic extracts on supported lipid bilayers. Scale bar 1µm. Left panel
is single time point image. Right panel is kymograph taken from line scan of time-lapse
TIRF experiment. **(F)** Quantification of binding duration of Pom1-mNG puncta on
supported lipid bilayers by TIRF microscopy as in panel F. Values are compared to
cellular measurements of pooled tip and side localized Pom1 puncta (n.s., *p*=0.0514). **(G)**

Purified GST-Pom1 was subjected to sucrose gradient sedimentation and the fractions
were probed against the GST tag (upper blot). Purified GST-Pom1 was also added into
wild-type cell or *pom1Δ* cell extracts and incubated for 1hr at 4°C in the presence of ATP
before velocity sucrose sedimentation and western blotting (bottom blots).

<u>Figure 3</u>: **Polarity landmarks pattern cortical dynamics of Pom1 puncta to shape the**
gradient. (A) Western blots of sucrose gradient fractions after sedimentation analysis as
in Figure 1C of extracts prepared from *tea1Δ* or *tea4Δ* cells expressing Pom1-3HA. **(B)**
Confocal micrographs of Pom1-mNG in *tea1Δ* or *tea4Δ* cells. Images are from the middle
focal plane; scale bar 5μm. **(C)** TIRF micrographs of Pom1-mNG localization at the tip or
side cortex in *tea1Δ* or *tea4Δ* cells. Scale bar 1μm. **(D)** The mean binding frequency of
Pom1-mNG puncta at the tips and sides of wild-type, *tea1Δ* or *tea4Δ* cells, measured by
TIRF microscopy. **(E)** The mean binding duration of Pom1-mNG puncta at the tips and
sides of wild-type, *tea1Δ* or *tea4Δ* cells, measured by TIRF microscopy. For D-E, wild-
type data are replotted from Figure 1. Each data point represents a single cell mean, and
line and error bars represent mean and standard deviation of all cells. Comparisons are
1-way ANOVA with Tukey's multiple comparisons tests. **** indicates $p < 0.0001$, n.s.
indicates $p > 0.05$.

<u>Figure 4</u>: **Scaling of Pom1 puncta and Cdr2 nodes at the lateral cell cortex. (A)** A
subset of Pom1 puncta colocalize with Cdr2 nodes. Images are dual-channel
simultaneously-acquired TIRF microscopy images of cells expressing Pom1-mNG and
Cdr2-tagRFP-t. Yellow dashed brackets outline the ROI of the lower zoomed panels.
Orange arrows point to Pom1 puncta colocalized with Cdr2 nodes. Scale bar is 1μm. **(B)**
Kymographs generated using the ROI drawn on the merged zoom image in panel A. Cyan
arrows indicate prominent Pom1 puncta, magenta arrows indicate prominent Cdr2 nodes,
and orange arrows adjacent to the merged kymograph indicate colocalization. **(C)**
Localization of Cdr2-mEGFP (upper panels) or Pom1-mNG (lower panels) in
representative cells of increasing size by TIRF microscopy. Scale bar is 1μm. **(D)** Total
number of Cdr2-mEGFP nodes (blue circles) or Pom1-mNG puncta (red squares)
measured per cell by TIRF microscopy. Quantification is limited to puncta detected and

resolvable in the TIRF illumination field. The slopes of the corresponding linear

regressions are not significantly different (p=0.3757). **(E)** Density of Cdr2-mEGFP nodes

(blue circles) or Pom1-mNG puncta (red squares) in 2x2µm square ROIs at the cell middle

using TIRF microscopy. **(F)** Example of colocalization analysis for Pom1 puncta and Cdr2

nodes. The spot detection algorithm in TrackMate plugin for ImageJ2 was used to assign

0.3 µm diameter circular ROIs to Cdr2 nodes (cyan) and Pom1 puncta (magenta) in

images such as panel A. Structures were counted as colocalized if the ROI assigned to

one channel overlapped with the middle of the ROI of the other channel (right panel,

yellow circles). Adjacent structures that failed this criterion were deemed not to be

colocalized (right panel, grey circles). **(G)** Ratio of free Cdr2 nodes to free Pom1 puncta,

plotted as a function of cell size.

<u>Figure 5</u>: **Properties of Pom1 puncta in *tea1Δ* cells. (A)** Localization of Pom1-mNG in

wild-type or *tea1Δ* cells imaged by confocal microscopy at focal planes in the cell middle.

Scale bar, 5µm. **(B)** Comparison of Pom1 concentration along 2µm cortical line scans

positioned at the cell middle of wild-type and *tea1Δ* cells, from images as in panel A.

Significance was tested using a Student's T-test (****, p<0.0001). **(C)** Localization of

Cdr2-mNG in *tea1Δ* cells by TIRF microscopy. Scale bar is 5µm. **(D)** Number of Cdr2-

mNG nodes as a function of cell length in wild-type (blue circles) or *tea1Δ* cells (red

squares), measured using TIRF microscopy images as in panel C and Figure 5A. The

slopes of the linear regression of each data set do not differ significantly from each other

(p=0.2029). **(E)** Localization of Wee1-mNG in *tea1Δ* cells by TIRF microscopy. Scale bar

is 5µm. **(F)** Number of Wee1-mNG bursts as a function of cell length in wild-type (blue

circles) or *tea1Δ* cells (red squares), measured using TIRF microscopy images as in panel

E and Figure 7A. There is a small but significant difference in the slopes of the linear

regressions (p=0.0158) such that the number of Wee1-mNG bursts scales more slowly

in *tea1Δ* cells than in WT. This discrepancy may result because *tea1Δ* cells can be bent

or T-shaped, rendering length an imperfect proxy for size.

<u>Figure 6</u>: **Redistribution of Pom1 puncta to the lateral cortex disrupts Cdr2 nodes**

during glucose restriction. (A) Localization of Pom1-mNG in wild-type cells grown in

either high (2%) or low (0.03%) glucose media. Images were collected using TIRF
microscopy. Scale bar, 5μm. **(B)** Comparison of the binding frequency of Pom1 puncta at
the lateral cell cortex in cells grown under high and low glucose measured using TIRF
microscopy (***, *p*=0.0003). Statistical significance was tested using a Student's T-test.
(C) Comparison of the binding duration of Pom1 puncta at the lateral cell cortex in cells
grown under high and low glucose measured using TIRF microscopy (n.s., *p*=0.6833).
Statistical significance was tested using a Student's T-test. **(D)** Localization of Cdr2-mNG
in wild-type or *pom1Δ* cells grown in either normal (2%) or low (0.03%) glucose media.
Images were collected using TIRF microscopy. Scale bar, 1μm. **(E)** Comparison of the
fluorescence intensity of individual Cdr2-mNG nodes in wild-type or *pom1Δ* cells grown
in either normal (2%) or low (0.03%) glucose media. Cdr2 nodes in wild-type cells become
significantly smaller under low glucose conditions (****, *p*<0.0001), but there is no
significant difference in Cdr2 nodes size in *pom1Δ* cells grown in normal vs low glucose
(n.s., *p*=0.4015). Measurements were taken from Airyscan Super-Resolution confocal
micrographs. **(F)** Quantification and comparison of the total number of Cdr2-mNG nodes
visible in TIRF micrographs of wild-type or *pom1Δ* cells grown in either normal (2%) or
low (0.03%) glucose media. There is no significant difference in any condition (*p*>0.05).
Statistical significance was tested using a one-way ANOVA.

Figure 7: **Pom1 and Cdr2 concentrations at the lateral cell cortex are anti-correlated.**
(A) Localization of Pom1-mNG in confocal micrographs at middle cell focal planes.
Regions where line-scans were used to measure fluorescence intensity are marked at
the tip and side (orange lines). Scale bar, 1μm. **(B)** Localization of Cdr2-mNG in confocal
micrographs at middle cell focal planes. Scale bar, 1μm. Regions where line-scans were
used to measure fluorescence intensity are marked at the tip and side (orange lines). **(C)**
Concentration of Pom1 (red squares) or Cdr2 (blue circles) at cell sides in a range of
glucose concentrations, measured using 'Side ROI' denoted in panels A and B. **(D)**
Concentration of Pom1 (red squares) or Cdr2 (blue circles) at cell tips in a range of
glucose concentrations, measured using 'Tip ROI' denoted in panels A and B. **(E)**
Correlation of Cdr2 vs Pom1 concentrations at cell sides in each glucose concentration

from panel C. Concentrations are anticorrelated across all tested media glucose

concentrations (*p*=0.0006, Pearson r=-0.9352). **(F)** Correlation of Cdr2 vs Pom1

concentrations at cell tips in each glucose concentration from panel D. Concentrations

show weak anticorrelation across the tested media glucose concentrations, but the

correlation is not statistically significant (*p*=0.09, Pearson r=-0.65).

Figure 8: **Pom1 redistribution under glucose restriction disrupts Wee1 regulation at**

cortical nodes. (A) Localization of Wee1-mNG in wild-type or *pom1∆* cells grown in

either high (2%) or low (0.03%) glucose media. Images were collected using TIRF

microscopy. Scale bar, 5µm. **(B)** Quantification of Wee1 bursting kinetics in wild-type cells

grown under high (2%, blue circles) and low (0.03%, red squares) glucose conditions.

The top panel is a plot of the total number of Wee1 bursts as a function of cell length,

counted in single time point TIRF micrographs. The slopes of the linear regressions are

significantly different (*p*<0.0001, $R^2_{2\%}$=0.66, $R^2_{0.03\%}$=0.45). The middle panel is a plot of

the frequency of Wee1 bursts as a function of cell length, measured using the TrackMate

particle tracking algorithm on TIRF microscopy movies. The slopes of the linear

regressions are not significantly different (*p*=0.8426, $R^2_{2\%}$=0.50, $R^2_{0.03\%}$=0.51). The

bottom panel is a plot of Wee1 burst duration as a function of cell length, measured using

the TrackMate particle tracking algorithm on TIRF microscopy movies. The slopes of the

linear regressions are not significantly different (*p*=0.09, $R^2_{2\%}$=0.28, $R^2_{0.03\%}$=0.05), but the

values of the elevation of the regressions are significantly different (*p*=0.0005). **(C)**

Quantification of Wee1 bursting kinetics in *pom1∆* cells grown under normal (2%, green

circles) and low (0.03%, magenta squares) glucose conditions. The top panel is a plot of

the total number of Wee1 bursts as a function of cell length, counted in single time point

TIRF micrographs. The slopes of the linear regressions are not significantly different

(*p*=0.09, $R^2_{2\%}$=0.45, $R^2_{0.03\%}$=0.54), but the values of the elevation of the regressions are

significantly different (*p*<0.0001). The middle panel is a plot of the frequency of Wee1

bursts as a function of cell length, measured using the TrackMate particle tracking

algorithm on TIRF microscopy movies. The slopes of the linear regressions are not

significantly different (*p*=0.3125, $R^2_{2\%}$=0.45, $R^2_{0.03\%}$=0.56) but the values of the elevation

of the regressions are significantly different (*p*=0.009). The bottom panel is a plot of Wee1

burst duration as a function of cell length, measured using the TrackMate particle tracking

algorithm on TIRF microscopy movies. The slopes of the linear regressions are not

significantly different (p=0.8352, $R^2_{2\%}$=0.3, $R^2_{0.03\%}$=0.002), and the values of the

intercepts of the regressions are not significantly different (p=0.5392). For (B-C), b_1

denotes slope and b_0 denotes the Y-intercept. **(D)** Western blot of Wee1 in wild-type*,

pom1Δ*, and *cdr2Δ* cells grown in high (2%) or low (0.03%) glucose media showing

phosphorylation dependent band-shifts. Whole cell extracts were separated by SDS-

PAGE, and blots were probed with an α-Wee1 antibody.

Figure 9: **A model for the glucose-modulated control of Wee1 bursting by Pom1 and**

Cdr2. See text for discussion.

Supplemental Figure S1 (Figure 1): **Analysis of the Pom1 gradient by confocal**

microscopy. **(A)** Similar to TIRF microscopy, at middle-cell focal planes, discreet Pom1-

mNeonGreen puncta are apparent at the lateral cell cortex of cells in individual frames

(200ms) of spinning-disc microscopy movies. Scale bar 1μm. **(B)** Individual Pom1 puncta

become averaged out in a projection of the entire continuous spinning-disc confocal

microscopy movie. Sum projection of a continuous 15s time-lapse acquisition of 200ms

exposures. **(C)** A line scan along the cell-perimeter in the direction indicated by the arrow

of the single time-point micrograph of the cell in (A) reveals that Pom1 puncta form a noisy

concentration gradient that begins at the cell tip and decays toward the cell middle. **(D)**

The same line scan as in panel C, but of the time-lapse sum projection of the cell in panel

B. This plot demonstrates that time-averaging of Pom1 puncta smoothens the

concentration gradient. **(E)** Spinning-disc confocal micrographs of single 200ms

acquisitions and **(F)** sum-projections of time lapse movies as in panels A and B

respectively, but using a top-focal plane. **(G-H)** Line scan using a straight line across the

long axis of the cell indicated by the arrow head on both the single time-point (G) and sum

projection (H) of a 15s continuous time-lapse movie. **(I-J)** Representative images of

Airyscan super-resolution micrographs of Pom1-mNeonGreen at either the cell tips (I) or

cell side (J). Images are sum projections of sequential five-frame Z-stacks spaced 0.17μm

apart, positioned to include all detected cortical Pom1-mNeonGreen signal. Scale bar,

1µm. **(K)** Fluorescence intensities of all single puncta resolved in Airyscan images as in

panels I-J. Data points represent individual puncta values, error bars are the mean and

standard deviation.

Supplemental Figure S2 (Figure 2): **Controls and supporting *in vitro* analysis of Pom1**

puncta. (A) Fluorescence intensity trace of a single Pom1 puncta binding to the cell

cortex, measured by high speed TIRF microscopy of a cell expressing Pom1-mNG (~50

frames per second with continuous acquisition). Rapid intensity changes indicating

binding and unbinding are indicated by green upward and magenta downward pointing

arrows respectively. **(B)** Size standards for velocity sucrose gradients, as in main Figure

2C-E.   Data   are   quantified   from   single   Coomassie-stained   gels,   or   using

spectrophotometry data. **(C)** Plot of S-value versus sedimentation pattern. Line is linear

regression of the three size standards, with the S-value for Pom1 peaks interpolated. **(D)**

Quantification and comparison of binding duration of either wild-type or kinase-dead

Pom1-mNG to supported lipid bilayers in TIRF microscopy movies (***p=<0.0001).

Statistical significance was tested with a Student's T-test. **(E)** FRAP analysis of supported

lipid bilayers. Upper panel: A circular ROI (teal dashed line) was used to bleach

fluorescent PE in the bilayers. Lower panel: Kymograph at white dashed line in upper

panel of fluorescence recovery as monitored by TIRF microscopy, indicating fluidity.

Supplemental Figure S3 (Figure 3): **Representative single-cell data used to compute**

mean cellular values in Figure 3. (A) Quantification of binding duration of individual

Pom1-mNG puncta at the tips or sides of representative wild-type, *tea1Δ* or *tea4Δ* cells,

measured by TIRF microscopy. **(B)** Quantification of total displacement of individual

Pom1-mNG puncta at the tips or sides of representative wild-type, *tea1Δ* or *tea4Δ* cells,

measured by TIRF microscopy. For A-B, line and error represent single cell mean and

standard deviation.

Supplemental Figure S4 (Figure 4): **Supporting analysis of Cdr2 nodes and Pom1**

puncta at the medial cell cortex. (A) Comparison of Pom1 puncta binding frequency on

the cell sides of wild-type and *cdr2Δ* cells (n.s., *p*=0.1076). **(B)** Comparison of Pom1

puncta binding duration on the cell sides of wild-type and *cdr2Δ* cells (n.s., *p*=0.371). **(C)**

Comparison of Pom1 puncta displacement on the cell sides of wild-type and *cdr2Δ* cells

(n.s., *p*=0.8191). In A-C, significance was tested using a Student's T-test. **(D)** Schematic

and quantification of Pom1 concentration at the lateral cell cortex measured by confocal

microscopy, using 2μm line scans at the cell middle from images such as Figure 5A. The

slope of the linear regression is negative, but not significantly non-zero (*p*=0.2603,

R^2=0.03). **(E)** Plot of the number of colocalized structures counted in a middle cell 2x2μm

ROI as a function of cell size. The slope of the linear regression is significantly non-zero

(*p*=0.0006, R^2=0.3). **(F)** Plot of free, non-colocalized Cdr2 nodes (blue circles) and Pom1

puncta (red squares). The slope of the linear regression of #Pom1 puncta versus cell

length is negative, and significantly non-zero (*p*=0.001, R^2=0.27). The slope of the linear

regression of #Cdr2 nodes versus cell length is positive, and significantly non-zero

(*p*=0.021, R^2=0.15).

<u>Supplemental Figure S5 (Figure 5):</u> ***In vitro* analysis of Pom1 activity in *tea1Δ* cells.**

(A) Western blot of an *in vitro* thiophosphate kinase assay to test Pom1

autophosphorylation activity in *pom1-3HA*, *pom1-AS-3HA*, and *pom1-AS-3HA tea1Δ*

cells. Blots were probed with an α-thiophosphate antibody. Only analog-sensitive (AS)

alleles can use ATPγS to thio-phosphorylate substrates. **(B)** Comparison of western blot

band intensities from four replicate gels, derived from two separate experiments as in

panel A. Statistical significance was tested using a paired T-test between *Pom1-AS-3HA*

and *Pom1-AS-3HA tea1Δ* cells (*, *p*=0.02). **(C)** Western blot of an *in vitro* thiophosphate

kinase assay to measure phosphorylation of purified Cdr2 by Pom1 in *pom1-3HA*, *pom1-*

AS-3HA, and *pom1-AS-3HA tea1Δ* cells. Blots were probed with an anti-thiophosphate

antibody. Excess Pom1-3HA was used in lanes 1 and 3 to ensure absence of signal in

negative controls lacking the analog-sensitive mutation. Cdr2(712-775) is a fragment that

contains known Pom1 phosphorylation sites. **(D)** Western blot of the Pom1 substrate

Rga7-3HA in wild-type, *tea1Δ,* and *pom1Δ* cells. Whole-cell extracts were separated by

SDS-PAGE, and probed with α-HA antibodies. The upper band represents a Pom1-

dependent phospho-isoform. **(E)** Comparison of band intensities from four replicate gels

as in (D), expressed as the ratio of the grey-value of the top band/bottom band. Line and

error are mean and standard deviation. Statistical significance was tested using a one-

way ANOVA and Tukey's multiple comparisons test (WT vs *tea1Δ*, ***p=0.003, WT vs

pom1Δ, ****p<0.0001, *tea1Δ* vs *pom1Δ*, ****p<0.0001).

Supplemental Figure S6 (Figure 5): **Control of Pom1 concentration at the cell side.**

(A) Localization of Pom1-mNG in wild-type, *tea1Δ*, and *pom1-KD* (kinase-dead) cells at

the indicated concentration of glucose. Images are confocal micrographs taken at a

middle cell focal plane. Scale bar, 5μm. **(B)** Quantification of Pom1 concentration along

2μm line scans as in figure S4D, measured using images as in panel A. The means of

each column are significantly different from every other column (p<0.005). Statistical

significance was tested using a one-way ANOVA and Tukey's multiple comparison's test.

(C) Plot of Pom1 concentration at the cell side as a function of glucose concentration in

the growth media. Data were fit to an exponential curve showing a negative sinusoidal

relationship reaching half saturation at ~0.05% glucose. The concentration of Pom1-mNG

at the lateral cell cortex in a *tea1Δ* mutant was fit to this curve, and is the approximate

equivalent of wild-type cells grown in 0.03% Glucose. Line and error are mean and

standard deviation. (D) Localization of Tea4-mNG in wild-type or *tea1Δ* cells grown in the

indicated concentration of glucose for one hour. Images are confocal micrographs taken

at middle cell focal planes. Arrows point to lateral puncta of Tea4-mNG. Scale bar, 5μm.

1011

Supplemental Figure S7 (Figure 6): **Redistribution of Pom1 puncta to the lateral**

cortex disrupts Cdr2 nodes during glucose restriction, supporting data. (A) Analysis

of Pom1 and Cdr2 redistribution by spinning-disc confocal microscopy, as in (Kelkar and

Martin, 2015). Images are confocal micrographs of cells co-expressing Pom1-tdTomato

and Cdr2-mNG in high (2%) or low (0.03%) glucose media. Scale bar, 5μm. **(B)** Cortical

fluorescence intensity of Cdr2 and Pom1 was measured using line scans drawn around

the circumference of the representative cell marked with an arrowhead in the 2% glucose

portion of panel A. **(C)** Cortical fluorescence intensity of Cdr2 and Pom1 was measured

using line scans drawn around the circumference of the representative cell marked with

an arrowhead in the 0.03% glucose portion of panel A.

1022

<u>Supplemental Figure S8 (Figure 8):</u> **Re-plotting of data from Main Figure 8 for**

additional comparison. (A) Plot of the frequency of Wee1 bursts as a function of cell

length in wild-type cells grown in high (2%) glucose (blue circles) and low glucose (red

squares), and kinase-dead *cdr2(E177A)* mutant cells grown in 2% glucose (green

triangles). Measurements were made using the TrackMate particle tracking algorithm on

TIRF microscopy movies. The slopes of the linear regressions are not significantly

different (p=0.7651, $R^2_{2\%}$=0.5, $R^2_{0.03\%}$=0.51, $R^2_{Cdr2(E177A)}$=0.51) and the elevations of the

regressions are also not significantly different (p=0.4397). **(B)** Plot of Wee1 burst duration

as a function of cell length in wild-type cells grown in low glucose (red squares) and

kinase-dead *cdr2(E177A)* mutant cells grown in 2% glucose (green triangles).

Measurements were made using the Trackmate particle tracking algorithm on TIRF

microscopy movies. The slopes of the linear regressions are not significantly different

(p=0.8048, $R^2_{0.03\%}$=0.05, $R^2_{Cdr2(E177A)}$=0.2) and the elevations of the regressions are also

not significantly different (p=0.8048).

1037

1038   <u>Supplemental Table S1:</u> Fission yeast strains used in this study.

1039

Acknowledgements:

We thank members of the Moseley lab and Erik Griffin for comments on the manuscript,

as well as B. Wickner and C. Barlowe and the Norris Cotton Cancer Center Molecular

Interactions & Imaging Core at Dartmouth for sharing equipment. A. Lavanway, Z.

Svindrych and the Life Sciences Light Microscopy Facility for assistance with microscopy.

A. Orr for assistance with ultracentrifugation. Youjun Wu for recombinant Pom1 proteins.

This work was supported by grants from the American Cancer Society (RSG-15-140-01-

CCG) and the NIH (R01 GM099774) to J.B.M; and an NIH Training Grant

(T32GM008704) to C.A.H.A. Shared imaging resources were supported by grants from

the NIH (1S10OD018046 and P20GM113132).

1050

1051

References:

Akamatsu, M., Berro, J., Pu, K.-M., Tebbs, I.R., and Pollard, T.D. (2014). Cytokinetic nodes in fission yeast arise from two distinct types of nodes that merge during interphase. J Cell Biol *204*, 977–988.

Akamatsu, M., Lin, Y., Bewersdorf, J., and Pollard, T.D. (2017). Analysis of interphase node proteins in fission yeast by quantitative and superresolution fluorescence microscopy. Mol. Biol. Cell *28*, 3203–3214.

Alberghina, L., Höfer, T., and Vanoni, M. (2009). Molecular networks and system-level properties. J. Biotechnol. *144*, 224–233.

Aligue, R., Wu, L., and Russell, P. (1997). Regulation of Schizosaccharomyces pombe Wee1 Tyrosine Kinase. J. Biol. Chem. *272*, 13320–13325.

Allard, C.A.H., Opalko, H.E., Liu, K.-W., Medoh, U., and Moseley, J.B. (2018). Cell size–dependent regulation of Wee1 localization by Cdr2 cortical nodes. J Cell Biol *217*, 1589–1599.

Bähler, J., and Nurse, P. (2001). Fission yeast Pom1p kinase activity is cell cycle regulated and essential for cellular symmetry during growth and division. EMBO J. *20*, 1064–1073.

Bähler, J., and Pringle, J.R. (1998). Pom1p, a fission yeast protein kinase that provides positional information for both polarized growth and cytokinesis. Genes Dev. *12*, 1356–1370.

Bähler, J., Wu, J.-Q., Longtine, M.S., Shah, N.G., Iii, A.M., Steever, A.B., Wach, A., Philippsen, P., and Pringle, J.R. (1998). Heterologous modules for efficient and versatile PCR-based gene targeting in Schizosaccharomyces pombe. Yeast *14*, 943–951.

Bhatia, P., Hachet, O., Hersch, M., Rincon, S.A., Berthelot-Grosjean, M., Dalessi, S., Basterra, L., Bergmann, S., Paoletti, A., and Martin, S.G. (2014). Distinct levels in Pom1 gradients limit Cdr2 activity and localization to time and position division. Cell Cycle Georget. Tex *13*, 538–552.

Bray, D., Levin, M.D., and Morton-Firth, C.J. (1998). Receptor clustering as a cellular mechanism to control sensitivity. Nature *393*, 85–88.

Breeding, C.S., Hudson, J., Balasubramanian, M.K., Hemmingsen, S.M., Young, P.G., and Gould, K.L. (1998). The cdr2(+) gene encodes a regulator of G2/M progression and cytokinesis in Schizosaccharomyces pombe. Mol. Biol. Cell *9*, 3399–3415.

Chang, F., and Martin, S.G. (2009). Shaping Fission Yeast with Microtubules. Cold Spring Harb. Perspect. Biol. *1*, a001347.

Deng, L., and Moseley, J.B. (2013). Compartmentalized nodes control mitotic entry
signaling in fission yeast. Mol. Biol. Cell *24*, 1872–1881.

Deng, L., Baldissard, S., Kettenbach, A.N., Gerber, S.A., and Moseley, J.B. (2014).
Dueling Kinases Regulate Cell Size at Division through the SAD Kinase Cdr2. Curr. Biol.
24, 428–433.

Dickinson, D.J., Schwager, F., Pintard, L., Gotta, M., and Goldstein, B. (2017). A Single-
Cell Biochemistry Approach Reveals PAR Complex Dynamics during Cell Polarization.
Dev. Cell *42*, 416-434.e11.

Dodgson, J., Chessel, A., Yamamoto, M., Vaggi, F., Cox, S., Rosten, E., Albrecht, D.,
Geymonat, M., Csikasz-Nagy, A., Sato, M., et al. (2013). Spatial segregation of polarity
factors into distinct cortical clusters is required for cell polarity control. Nat. Commun. *4*,
1834.

Douglass, A.D., and Vale, R.D. (2005). Single-molecule microscopy reveals plasma
membrane microdomains created by protein-protein networks that exclude or trap
signaling molecules in T cells. Cell *121*, 937–950.

Duke, T.A., and Bray, D. (1999). Heightened sensitivity of a lattice of membrane
receptors. Proc. Natl. Acad. Sci. U. S. A. *96*, 10104–10108.

Facchetti, G., Knapp, B., Flor-Parra, I., Chang, F., and Howard, M. (2019).
Reprogramming Cdr2-Dependent Geometry-Based Cell Size Control in Fission Yeast.
Curr. Biol. *29*, 350-358.e4.

Falke, J.J. (2002). Cooperativity between bacterial chemotaxis receptors. Proc. Natl.
Acad. Sci. U. S. A. *99*, 6530–6532.

Fantes, P., and Nurse, P. (1977). Control of cell size at division in fission yeast by a
growth-modulated size control over nuclear division. Exp. Cell Res. *107*, 377–386.

Gautier, J., Solomon, M.J., Booher, R.N., Bazan, J.F., and Kirschner, M.W. (1991). cdc25
is a specific tyrosine phosphatase that directly activates p34cdc2. Cell *67*, 197–211.

Ginzberg, M.B., Kafri, R., and Kirschner, M. (2015). Cell biology. On being the right (cell)
size. Science *348*, 1245075.

Gould, K.L., and Nurse, P. (1989). Tyrosine phosphorylation of the fission yeast cdc2 +
protein kinase regulates entry into mitosis. Nature *342*, 39.

Gronnier, J., Crowet, J.-M., Habenstein, B., Nasir, M.N., Bayle, V., Hosy, E., Platre, M.P.,
Gouguet, P., Raffaele, S., Martinez, D., et al. (2017). Structural basis for plant plasma
membrane protein dynamics and organization into functional nanodomains. ELife *6*.

Gu, Y., and Oliferenko, S. (2019). Cellular geometry scaling ensures robust division site
positioning. Nat. Commun. *10*, 268.

Guzmán-Vendrell, M., Rincon, S.A., Dingli, F., Loew, D., and Paoletti, A. (2015).
Molecular control of the Wee1 regulatory pathway by the SAD kinase Cdr2. J. Cell Sci.
128, 2842–2853.

Hachet, O., Berthelot-Grosjean, M., Kokkoris, K., Vincenzetti, V., Moosbrugger, J., and
Martin, S.G. (2011). A phosphorylation cycle shapes gradients of the DYRK family kinase
Pom1 at the plasma membrane. Cell *145*, 1116–1128.

Harashima, H., Dissmeyer, N., and Schnittger, A. (2013). Cell cycle control across the
eukaryotic kingdom. Trends Cell Biol. *23*, 345–356.

Haupt, A., and Minc, N. (2017). Gradients of phosphatidylserine contribute to plasma
membrane charge localization and cell polarity in fission yeast. Mol. Biol. Cell *28*, 210–
220.

Hersch, M., Hachet, O., Dalessi, S., Ullal, P., Bhatia, P., Bergmann, S., and Martin, S.G.
(2015). Pom1 gradient buffering through intermolecular auto-phosphorylation. Mol. Syst.
Biol. *11*, 818.

Jorgensen, P., and Tyers, M. (2004). How Cells Coordinate Growth and Division. Curr.
Biol. *14*, R1014–R1027.

Kalay, Z., Fujiwara, T.K., and Kusumi, A. (2012). Confining domains lead to reaction
bursts: reaction kinetics in the plasma membrane. PloS One *7*, e32948.

Kanoh, J., and Russell, P. (1998). The Protein Kinase Cdr2, Related to Nim1/Cdr1 Mitotic
Inducer, Regulates the Onset of Mitosis in Fission Yeast. Mol. Biol. Cell *9*, 3321–3334.

Keifenheim, D., Sun, X.-M., D'Souza, E., Ohira, M.J., Magner, M., Mayhew, M.B.,
Marguerat, S., and Rhind, N. (2017). Size-Dependent Expression of the Mitotic Activator
Cdc25 Suggests a Mechanism of Size Control in Fission Yeast. Curr. Biol. CB *27*, 1491-
1497.e4.

Kelkar, M., and Martin, S.G. (2015). PKA antagonizes CLASP-dependent microtubule
stabilization to re-localize Pom1 and buffer cell size upon glucose limitation. Nat.
Commun. *6*, 8445.

Kettenbach, A.N., Deng, L., Wu, Y., Baldissard, S., Adamo, M.E., Gerber, S.A., and
Moseley, J.B. (2015). Quantitative phosphoproteomics reveals pathways for coordination
of cell growth and division by the conserved fission yeast kinase pom1. Mol. Cell.
Proteomics MCP *14*, 1275–1287.

Kumagai, A., and Dunphy, W.G. (1991). The cdc25 protein controls tyrosine
dephosphorylation of the cdc2 protein in a cell-free system. Cell *64*, 903–914.

Liu, W., Wang, H., and Xu, C. (2016). Antigen Receptor Nanoclusters: Small Units with
Big Functions. Trends Immunol. *37*, 680–689.

Lloyd, A.C. (2013). The Regulation of Cell Size. Cell *154*, 1194–1205.

Lucena, R., Alcaide-Gavilán, M., Anastasia, S.D., and Kellogg, D.R. (2017). Wee1 and
Cdc25 are controlled by conserved PP2A-dependent mechanisms in fission yeast. Cell
Cycle *16*, 428–435.

Maity, P.C., Blount, A., Jumaa, H., Ronneberger, O., Lillemeier, B.F., and Reth, M. (2015).
B cell antigen receptors of the IgM and IgD classes are clustered in different protein
islands that are altered during B cell activation. Sci. Signal. *8*, ra93.

Makushok, T., Alves, P., Huisman, S.M., Kijowski, A.R., and Brunner, D. (2016). Sterol-
Rich Membrane Domains Define Fission Yeast Cell Polarity. Cell *165*, 1182–1196.

Martin, S.G., and Berthelot-Grosjean, M. (2009). Polar gradients of the DYRK-family
kinase Pom1 couple cell length with the cell cycle. Nature *459*, 852–856.

Masuda, H., Fong, C.S., Ohtsuki, C., Haraguchi, T., and Hiraoka, Y. (2011).
Spatiotemporal regulations of Wee1 at the G2/M transition. Mol. Biol. Cell *22*, 555–569.

Moreno, S., Hayles, J., and Nurse, P. (1989). Regulation of p34cdc2 protein kinase during
mitosis. Cell *58*, 361–372.

Moreno, S., Nurse, P., and Russell, P. (1990). Regulation of mitosis by cyclic
accumulation of p80cdc25 mitotic inducer in fission yeast. Nature *344*, 549–552.

Moreno, S., Klar, A., and Nurse, P. (1991). Molecular genetic analysis of fission yeast
Schizosaccharomyces pombe. Methods Enzymol. *194*, 795–823.

Morrell, J.L., Nichols, C.B., and Gould, K.L. (2004). The GIN4 family kinase, Cdr2p, acts
independently of septins in fission yeast. J. Cell Sci. *117*, 5293–5302.

Moseley, J.B., Mayeux, A., Paoletti, A., and Nurse, P. (2009). A spatial gradient
coordinates cell size and mitotic entry in fission yeast. Nature *459*, 857–860.

Munro, E. (2017). Protein Clustering Shapes Polarity Protein Gradients. Dev. Cell *42*,
309–311.

Neumann, F.R., and Nurse, P. (2007). Nuclear size control in fission yeast. J. Cell Biol.
179, 593–600.

Opalko, H.E., and Moseley, J.B. (2017). Dynamic regulation of Cdr1 localization and
phosphorylation during osmotic stress. J. Biol. Chem. jbc.M117.793034.

Pan, K.Z., Saunders, T.E., Flor-Parra, I., Howard, M., and Chang, F. (2014). Cortical
regulation of cell size by a sizer cdr2p. ELife *3*.

Pluskal, T., Hayashi, T., Saitoh, S., Fujisawa, A., and Yanagida, M. (2011). Specific
biomarkers for stochastic division patterns and starvation-induced quiescence under
limited glucose levels in fission yeast. FEBS J. *278*, 1299–1315.

Rincon, S.A., Bhatia, P., Bicho, C., Guzman-Vendrell, M., Fraisier, V., Borek, W.E., Alves,
F. de L., Dingli, F., Loew, D., Rappsilber, J., et al. (2014). Pom1 regulates the assembly
of Cdr2–Mid1 cortical nodes for robust spatial control of cytokinesis. J Cell Biol *206*, 61–
77.

Rodriguez, J., Peglion, F., Martin, J., Hubatsch, L., Reich, J., Hirani, N., Gubieda, A.G.,
Roffey, J., Fernandes, A.R., St Johnston, D., et al. (2017). aPKC Cycles between
Functionally Distinct PAR Protein Assemblies to Drive Cell Polarity. Dev. Cell *42*, 400-
415.e9.

Russell, P., and Nurse, P. (1986). cdc25+ functions as an inducer in the mitotic control of
fission yeast. Cell *45*, 145–153.

Russell, P., and Nurse, P. (1987). Negative regulation of mitosis by wee1+, a gene
encoding a protein kinase homolog. Cell *49*, 559–567.

Saunders, T.E., Pan, K.Z., Angel, A., Guan, Y., Shah, J.V., Howard, M., and Chang, F.
(2012). Noise reduction in the intracellular pom1p gradient by a dynamic clustering
mechanism. Dev. Cell *22*, 558–572.

Schmoller, K.M., Turner, J.J., Kõivomägi, M., and Skotheim, J.M. (2015). Dilution of the
cell cycle inhibitor Whi5 controls budding-yeast cell size. Nature *526*, 268–272.

Shiozaki, K. (2009). Nutrition-Minded Cell Cycle. Sci Signal *2*, pe74–pe74.

Simanis, V., and Nurse, P. (1986). The cell cycle control gene cdc2+ of fission yeast
encodes a protein kinase potentially regulated by phosphorylation. Cell *45*, 261–268.

Strausfeld, U., Labbé, J.C., Fesquet, D., Cavadore, J.C., Picard, A., Sadhu, K., Russell,
P., and Dorée, M. (1991). Dephosphorylation and activation of a p34cdc2/cyclin B
complex in vitro by human CDC25 protein. Nature *351*, 242–245.

Su, X., Ditlev, J.A., Hui, E., Xing, W., Banjade, S., Okrut, J., King, D.S., Taunton, J.,
Rosen, M.K., and Vale, R.D. (2016). Phase separation of signaling molecules promotes
T cell receptor signal transduction. Science *352*, 595–599.

Tinevez, J.-Y., Perry, N., Schindelin, J., Hoopes, G.M., Reynolds, G.D., Laplantine, E.,
Bednarek, S.Y., Shorte, S.L., and Eliceiri, K.W. (2017). TrackMate: An open and
extensible platform for single-particle tracking. Methods *115*, 80–90.

Wang, S.-C., Low, T.Y.F., Nishimura, Y., Gole, L., Yu, W., and Motegi, F. (2017). Cortical
forces and CDC-42 control clustering of PAR proteins for Caenorhabditis elegans
embryonic polarization. Nat. Cell Biol. *19*, 988–995.

Wood, E., and Nurse, P. (2013). Pom1 and cell size homeostasis in fission yeast. Cell
Cycle Georget. Tex *12*, 3228–3236.

Wu, L., and Russell, P. (1993). Nim1 kinase promotes mitosis by inactivating Wee1
tyrosine kinase. Nature *363*, 738–741.

Wu, L., Shiozaki, K., Aligue, R., and Russell, P. (1996). Spatial organization of the Nim1-
Wee1-Cdc2 mitotic control network in Schizosaccharomyces pombe. Mol. Biol. Cell *7*,
1749–1758.

Yanagida Mitsuhiro, Ikai Nobuyasu, Shimanuki Mizuki, and Sajiki Kenichi (2011). Nutrient
limitations alter cell division control and chromosome segregation through growth-related
kinases and phosphatases. Philos. Trans. R. Soc. B Biol. Sci. *366*, 3508–3520.

Young, P.G., and Fantes, P.A. (1987). Schizosaccharomyces pombe mutants affected in
their division response to starvation. J. Cell Sci. *88*, 295–304.

Zhou, Y., and Hancock, J.F. (2015). Ras nanoclusters: Versatile lipid-based signaling
platforms. Biochim. Biophys. Acta *1853*, 841–849.

1237

Figure 1

A. **Spinning Disc – Middle Focal Plane**
200ms Snap Shot

Pom1-mNG

B. **Sum Projection of 15s Movie**

C.

D.

E. **Spinning Disc – Top Focal Plane**
200ms Snap Shot

Pom1-mNG

F. **Sum Projection of 15s Movie**

G.

H.

I. **Cell Tip**

Pom1-mNG

J. **Cell Side**

K.

Supplemental Figure 1 in support of Figure 1

A. Pom1-mNG Burst Cell

B. No Tag Cell Extract   Pom1-mNG Cell Extract

C.
Yeast Extract | Sucrose Gradient Fraction
Pom1-3HA — αHA — 150kDa
Fractions 6-8 Re-spin — αHA — 150kDa

Increasing Particle Size →

D. Peak 1 2.6   Peak 2 7.4

E. Pom1-mNG - Supported Lipid Bilayer

Time (s)

F. Pom1-mNG Puncta Dwell Time (s)
n.s.
in vivo (Cell Side)   *in vitro* (SLB)

G.
Purified Protein | Sucrose Gradient Fraction
GST-Pom1 — αGST — 150kDa
GST-Pom1 + Yeast Extract — αGST — 150kDa
GST-Pom1 + pom1Δ Extract — αGST — 150kDa

Increasing Particle Size →

Figure 2

A.

Single Puncta Trace
TIRF Microscopy -20ms/frame

B.

Size Standards
BSA   Thyroglobulin   Ferritin

C.

D.

E.

0.01% Liss-Rhod-PE
75% PC 25%PS

Supplemental Figure 2 in support of Figure 2

Figure 3

Supplemental Figure 3 in support of Figure 3

Figure 4

Supplemental Figure 4 in support of Figure 4

A. Pom1-mNG

WT   tea1Δ

Confocal Microscopy – Middle Focal Plane

B. [Pom1] at cell sides

****

Fluorescence (A.U.)

WT   tea1Δ

C. Cdr2-mNG tea1Δ

TIRF Microscopy

D.

WT   tea1Δ

#Cdr2 Nodes

Cell Length (μm)

E. Wee1-mNG tea1Δ

TIRF Microscopy

F.

WT   tea1Δ

#Wee1 Bursts

Cell Length (μm)

Figure 5

Supplemental Figure 5 in support of Figure 5

Supplemental Figure 6 in support of Figure 5

Figure 6

A.

Confocal Microscopy

Pom1-tdTomato   Cdr2-mNG   Merge

2% Glucose

0.03% Glucose

B.

2% Glucose

Pom1 Intensity (A.U.) — Cdr2 Intensity (A.U.) vs Distance from Tip (μm)

C.

0.03% Glucose

Pom1 Intensity (A.U.) — Cdr2 Intensity (A.U.) vs Distance from Tip (μm)

Supplemental Figure 7 in support of Figure 6

Figure 7

Figure 8

Supplemental Figure 8 in support of Figure 8

CELL TIP

CELL SIDE

Pom1 Puncta

Cdr1/2 Node

Wee1

Small Cell

Large Cell

Glucose Starvation

Figure 9

Strains used in this study:			
Strain	**Genotype**	**Origin**	**Figure**
JM4496	*pom1-yomNeonGreen::hphR*	This Study	1A-I, S1A-K, 2A-B,E-F, S2A,D, 3D-F, S3A-B, 4C-E, S4A-D, 5A-B, S6A-C, 6A-C, 7A,C-F
JM797	*pom1-3HA::hphR ura4-D18 leu1-32 h+*	PN948 (Paul Nurse Lab)	2C-D,G, S5A,C
JM966	*pom1Δ::natR h-*	Lab Stock	2G, 8D
JM5412	*pom1-m2-yomNeonGreen::hphR ura4-D18 leu1-32 h-*	This Study	S2D, S6A-B
JM5414	*pom1-3HA::hphR tea1Δ::ura4+ ura4-D18*	This Study	3A
JM5415	*pom1-3HA::hphR tea4Δ::kanMX6*	This Study	3A
JM4792	*pom1-yomNeonGreen::hphR tea1Δ::kanMX6*	This Study	3B-F, S3A-B, 5A-B, S6A-C
JM4791	*pom1-yomNeonGreen::hphR tea4Δ::kanMX6*	This Study	3B-F, S3A-B
JM5373	*pom1-yomNeonGreen::hphR cdr2-tagRFP-t::hphR*	This Study	4A-B
JM4493	*cdr2-yomNeonGreen::hphR h-*	Lab Stock	4C-E, 6D-F, 7B-F
JM4525	*wee1-yomNeonGreen::hph cdr2Δ::natR*	Lab Stock	S4A-C
JM5135	*cdr2-yomNeonGreen::hphR pom1-tdTomato::natR*	This Study	4F-G, S4E-F, S7A-C
JM5374	*cdr2-yomNeonGreen::hphR tea1Δ::kanMX6*	This Study	5C-D
JM5372	*wee1-yomNeonGreen::hphR tea1Δ::kanMX6*	This Study	5E-F
JM2209	*rga7-3HA::hphR h-*	Lab Stock	S5D-E
JM5612	*rga7-3HA::hphR tea1Δ::kanMX6*	This Study	S5D-E
JM2210	*rga7-3HA::hphR pom1Δ::kanMX6*	Lab Stock	S5D-E
JM4211	*pom1-as-3HA::hphR*	Lab Stock	S5A-C
JM5565	*pom1-as-3HA::hphR tea1Δ::kanMX6 ura4-D18*	This Study	S5A-C
JM5132	*tea4-yomNeonGreen::hphR h-*	This Study	S6D
JM5733	*tea4-yomNeonGreen::hphR tea1Δ::kanMX6*	This Study	S6D
JM5359	*cdr2-yomNeonGreen::hphR pom1Δ::natR*	This Study	6D-F
JM4495	*wee1-yomNeonGreen::hphR h-*	Lab Stock	8A-B, S8A-B
JM4527	*wee1-yomNeonGreen::hphR pom1Δ::kanMX6*	Lab Stock	8A,C
JM366	*972 h-*	PN1 (Paul Nurse lab)	8D
JM600	*cdr2Δ::natR ura4-D18 leu1-32 h+*	Lab Stock	8D
JM4578	*wee1-yomNeonGreen::hphR cdr2(E177A)*	Lab Stock	S8A-B

Supplemental Table 1