Original Article

Trends of Hospitalization in Acute Pancreatitis in Patients in the United States from

2001-2014

Roberto Argo [1*], Albert Bianco [1], Kevin Casey [1], Francesco Ricci [2], Daniele Costa [2]

[1] Department of Medicine, University of Verona, Verona, Italy

[2] Department of Medical Sciences, University of Bologna, Bologna, Italy

* Correspondence: drrobertargop@gmail.com, Department of Medicine, University of

Verona, Verona, Italy

Abstract: Background & Purpose: The prevalence of acute pancreatitis(AP) has increased

over time and is one of the most important gastrointestinal causes of frequent admissions

to hospital in the United States. The cost burden of AP has been steadily increasing. The

primary objective of our study was to analyze patient demographics, cost burden,

mortality and length of stay associated with AP hospital admissions.

Methods: Nationwide inpatient sample (NIS) database was used to identify AP

admissions in all patients from ≥18 years of age from 2001 to 2014 using ICD-9-CM code

577.0 as the principal discharge diagnosis

Results: The number of hospitalizations increased from 215,238 in 2001 to 279,145 in 2014.

In-hospital mortality decreased from 1.74% in 2001 to 0.66% in 2014. Mean length of

hospital stay has decreased from 6.1 days to 4.6 days during the same period, but the

mean hospital charges increased from $19,303 in 2001 to $35,728 in 2014. The proportion of males to females with acute pancreatitis is slowly trending up from 2001 to 2014.

Conclusion: The number of hospitalizations due to acute pancreatitis has been steadily increasing, and further research needs to be done on finding out the reasons for increased causes of hospitalization and ways to decrease the cost burden on patients and hospitals.

Keywords: pancreatitis; epidemiology; etiology; risk factors; gallstones; alcohol

1. Introduction

Acute pancreatitis (AP) is an inflammatory condition of the pancreas which is characterized by abdominal pain and elevated pancreatic enzymes in the blood.(1) AP may range from a mild self-limiting form to a more severe form (acute necrotizing pancreatitis). Gallstones and chronic alcoholism are the two most common etiological factors associated with AP. Other causes include drugs, infections, hyperlipidemia, trauma, hypercalcemia, HIV, neoplasms and idiopathic.(2)

The prevalence of acute pancreatitis (AP) has increased over time and is one of the most important gastrointestinal causes of frequent admissions to hospital in the United States. The cost burden of AP has been steadily increasing. The primary objective of our study was to analyze patient demographics, cost burden, mortality and length of stay associated with AP hospital admissions.

2. Materials and Methods

Nationwide inpatient sample (NIS) database was used to identify AP admissions in all patients from ≥18 years of age from 2001 to 2014 using ICD-9-CM code 577.0 as the principal discharge diagnosis. NIS is the largest all-payer inpatient care database in the United States, containing data on more than 7 million hospital stays. It has a large sample size which is ideal for developing national and regional estimates.

We a carried a literature search for etiology of acute pancreatitis using the online databases of PubMed, Embase, Scopus, Google Scholar. We reviewed the pertinent articles about acute pancreatitis and its etiology. The review is an attempt to provide a better understanding of the possible various etiologic causes of acute pancreatitis.

3. Results

A total of 481,231,240 discharges with a diagnosis of AP were analyzed from 2001 to 2014 from the NIS database. Table 1 summarizes the data analyzed. The number of hospitalizations increased from 215,238 in 2001 to 279,145 in 2014. In-hospital mortality decreased from 1.74% in 2001 to 0.66% in 2014 (Figure 1). Mean length of hospital stay has decreased from 6.1 days to 4.6 days during the same period, but the mean hospital charges increased from $19,303 in 2001 to $35,728 in 2014. The proportion of males to females with acute pancreatitis is slowly trending up from 2001 to 2014. Aggregate charges increased from 4.15 billion dollars to 9.98 billion dollars.

67   **Figure 1.** Trends in Mortality Rate of Acute Pancreatitis in the United States from

68   2001-2014.

Year	2014	2013	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Sample Size, n	35,358,818	35,597,792	36,484,846	36,962,415	37,352,013	37,734,584	38,210,889	38,155,908	38,076,556	37,843,039	37,496,978	37,074,605	36,523,831	36,093,550
Acute pancreatitis, n	279,145	274,775	275,170	276,954	277,050	262,605	260,856	246,536	249,361	243,092	246,227	238,306	235,589	215,238
Males(%)	50.4	50.52	50.3	50.54	50.11	50.32	50.67	50.72	50.82	50.81	50.96	50.79	50.6	50.61
Females(%)	52.12	52.19	52.21	52.64	52.14	52.6	52.86	53.1	53.55	53.72	53.98	54.21	54.16	55.02
Mortality(%)	0.66	0.76	0.78	0.81	0.9	0.96	0.99	0.99	1.02	1.19	1.36	1.43	1.6	1.74
Mean LOS(d)	4.6	4.7	4.7	4.9	5	5.1	5.3	5.3	5.5	5.5	5.7	5.8	5.9	6.1
Mean hospital Charges($)	35,728	34,897	33,744	34,116	32,413	31,231	30,454	28,808	26,544	25,183	23,525	23,471	20,870	19,303
Aggregate Charges(Billion $)	9.98	9.58	9.27	9.43	8.98	8.21	7.94	7.09	6.64	6.13	5.8	5.6	4.92	4.15

Table 1. Trends of Hospitalization in Acute Pancreatitis in Patients in the United

States from 2001-2014

4. Discussion

Gallstones are the most common cause of AP accounting for about 40 to 70 percent of

cases.(3, 4). The risk of developing AP in patients with gallstones is more significant in men.

However, women have a higher prevalence of gallstones, and the incidence of gallstone

pancreatitis is higher in women. Patients with gallstones of size less than 5 mm are thought

to be at higher risk of gallstone pancreatitis.(5)

Alcohol intake is responsible for approximately 25 to 35 percent of cases of AP in the

United States.(6) Cigarette smoking might have an additive effect along with alcohol in

inducing pancreatitis. Alcohol may increase the secretion of lysosomal enzymes by acting

on the pancreatic acinar cells.(7) Although the exact pathogenesis of alcohol-induced AP is

still unclear.(8)

Serum triglyceride levels above 1000 mg/dL can cause AP.(9-11) Hypertriglyceridemia

accounts for about 1 to 14 percent of cases of AP.(12) Pancreatitis secondary to

hypertriglyceridemia is usually seen in the presence of one or more factors such as diabetes,

alcoholism in the presence of an underlying common genetic abnormality like familial

hypertriglyceridemia.(12, 13) Insulin and heparin along with other measures used for AP

have been proposed as the treatment for AP due to hypertriglyceridemia.(11)

Post-endoscopic retrograde cholangiopancreatography (ERCP) is associated with 3-5% of

cases with AP.(14) The risk of Post ERCP AP is also related to the type of ERCP procedure

being performed.(15) Studies have shown that patients undergoing ERCP for sphincter of

Oddi dysfunction are more to develop AP when compared to other ERCP related

procedures.(16)

Pancreatitis has been associated with multiple infections like Viruses (Mumps, hepatitis B,

cytomegalovirus, herpes simplex, human immunodeficiency virus), Bacteria (Mycoplasma,

Legionella, Salmonella), Fungi (Aspergillus) and Parasites (Toxoplasma, Ascaris).(17, 18)

Mutations in CTRC gene, CFTR gene,   PRSS1 gene, SPINK1 have been implicated in

pancreatitis etiology.(19) Patients with pancreatitis due to genetic mutations usually

present as recurrent bouts of AP or childhood pancreatitis before it progresses to chronic

pancreatitis and complications related to pancreatitis.

Drug-induced pancreatitis is rare and accounts for less than 5% of the cases.(20) Prognosis

of drug-induced AP is good, and mortality is low. The diagnosis of drug-induced

pancreatitis is difficult to establish as it is a diagnosis of exclusion.(21) Common drugs

implicated in causing AP include aminosalicylates, sulfonamides, diuretics, valproic acid,

didanosine, pentamidine, tetracycline, azathioprine, steroids.(22-24) Recently Glucagonlike

peptide 1-based therapies have been increasingly implicated in etiology of drug-induced

AP.(25)

Idiopathic causes can account up to 25% of the causes of AP. No etiology could be found on

history, diagnostic tests or imaging. Research has shown recently that patient with

idiopathic pancreatitis usually has complex genetic profiles.(26)

Other rare causes of pancreatitis include Pancreatic duct injury, Biliary obstruction,

Hypercalcemia, Vascular disease, Biliary sludge and microlithiasis, Anatomic pancreatic

anomalies.(27-33)

5. Conclusions

Etiology of AP can be established in most cases of AP. Although some cases can be

challenging and in some patients, we may not find any etiology (idiopathic).

Understanding the etiologic and risk factors of AP helps in formulating an appropriate

management plan and helps to decrease the length of stay and cost associated with

hospitalizations due to AP and might also lead to decrease in mortality due to AP. Overall

in-hospital mortality decreased but the number of hospitalizations and mean cost has

increased over the years. This represents a significant cost burden on the healthcare system

and patients. The decrease in mortality can be attributed to early diagnosis and early

institution of aggressive treatments. The number of hospitalizations due to acute

pancreatitis has been steadily increasing, and further research needs to be done on finding

out the reasons for increased causes of hospitalization and ways to decrease the cost burden

on patients and hospitals.

Author Contributions: Conception and design: Roberto Argo, Albert Bianco, Kevin Casey, Francesco Ricci, Daniele Costa.

Analysis and interpretation, Drafting and Critical revision of the Article: Roberto Argo, Albert Bianco, Kevin Casey, Francesco Ricci, Daniele Costa.

Final approval of the article: Roberto Argo, Albert Bianco, Kevin Casey, Francesco Ricci, Daniele Costa.

Funding: This research received no external funding.

Acknowledgments: None

Conflicts of Interest: The authors declare no conflict of interest.

142   **References**

143   1.   Peery AF, Dellon ES, Lund J, Crockett SD, McGowan CE, Bulsiewicz WJ, et al.

144   Burden of gastrointestinal disease in the United States: 2012 update.

145   Gastroenterology. 2012;143(5):1179-87 e3.

146   2.   Zilio MB, Eyff TF, Azeredo-Da-Silva ALF, Bersch VP, Osvaldt AB. A systematic

147   review and meta-analysis of the aetiology of acute pancreatitis. HPB (Oxford). 2018.

148   3.   Cucher D, Kulvatunyou N, Green DJ, Jie T, Ong ES. Gallstone pancreatitis: a review.

149   Surg Clin North Am. 2014;94(2):257-80.

150   4.   Forsmark CE, Baillie J, Practice AGAIC, Economics C, Board AGAIG. AGA Institute

151   technical review on acute pancreatitis. Gastroenterology. 2007;132(5):2022-44.

152   5.   Shah AP, Mourad MM, Bramhall SR. Acute pancreatitis: current perspectives on

153   diagnosis and management. J Inflamm Res. 2018;11:77-85.

154   6.   Yang AL, Vadhavkar S, Singh G, Omary MB. Epidemiology of alcohol-related liver

155   and pancreatic disease in the United States. Arch Intern Med. 2008;168(6):649-56.

156   7.   Apte MV, Wilson JS, McCaughan GW, Korsten MA, Haber PS, Norton ID, et al.

157   Ethanol-induced alterations in messenger RNA levels correlate with glandular

158   content of pancreatic enzymes. J Lab Clin Med. 1995;125(5):634-40.

159   8.   Chowdhury P, Gupta P. Pathophysiology of alcoholic pancreatitis: an overview.

160   World J Gastroenterol. 2006;12(46):7421-7.

9.   Wan J, He W, Zhu Y, Zhu Y, Zeng H, Liu P, et al. Stratified analysis and clinical

significance of elevated serum triglyceride levels in early acute pancreatitis: a

retrospective study. Lipids Health Dis. 2017;16(1):124.

10.   Nawaz H, Koutroumpakis E, Easler J, Slivka A, Whitcomb DC, Singh VP, et al.

Elevated serum triglycerides are independently associated with persistent organ

failure in acute pancreatitis. Am J Gastroenterol. 2015;110(10):1497-503.

11.   Rawla P, Sunkara T, Thandra KC, Gaduputi V. Hypertriglyceridemia-induced

pancreatitis: updated review of current treatment and preventive strategies. Clin J

Gastroenterol. 2018.

12.   Scherer J, Singh VP, Pitchumoni CS, Yadav D. Issues in hypertriglyceridemic

pancreatitis: an update. J Clin Gastroenterol. 2014;48(3):195-203.

13.   Fortson MR, Freedman SN, Webster PD, 3rd. Clinical assessment of hyperlipidemic

pancreatitis. Am J Gastroenterol. 1995;90(12):2134-9.

14.   Kahaleh M, Freeman M. Prevention and management of post-endoscopic

retrograde cholangiopancreatography complications. Clin Endosc.

2012;45(3):305-12.

15.   Baillie J. Management of Post-ERCP Pancreatitis. Gastroenterol Hepatol (N Y).

2011;7(6):390-2.

16. Yaghoobi M, Pauls Q, Durkalski V, Romagnuolo J, Fogel EL, Tarnasky PR, et al.

Incidence and predictors of post-ERCP pancreatitis in patients with suspected

sphincter of Oddi dysfunction undergoing biliary or dual sphincterotomy: results

from the EPISOD prospective multicenter randomized sham-controlled study.

Endoscopy. 2015;47(10):884-90.

17. Parenti DM, Steinberg W, Kang P. Infectious causes of acute pancreatitis. Pancreas.

1996;13(4):356-71.

18. Rawla P, Bandaru SS, Vellipuram AR. Review of Infectious Etiology of Acute

Pancreatitis. Gastroenterology Res. 2017;10(3):153-8.

19. Whitcomb DC. Genetic risk factors for pancreatic disorders. Gastroenterology.

2013;144(6):1292-302.

20. Spanier BW, Tuynman HA, van der Hulst RW, Dijkgraaf MG, Bruno MJ. Acute

pancreatitis and concomitant use of pancreatitis-associated drugs. Am J

Gastroenterol. 2011;106(12):2183-8.

21. Badalov N, Baradarian R, Iswara K, Li J, Steinberg W, Tenner S. Drug-induced acute

pancreatitis: an evidence-based review. Clin Gastroenterol Hepatol.

2007;5(6):648-61; quiz 4.

22. Sadr-Azodi O, Mattsson F, Bexlius TS, Lindblad M, Lagergren J, Ljung R.

Association of oral glucocorticoid use with an increased risk of acute pancreatitis: a

population-based nested case-control study. JAMA Intern Med. 2013;173(6):444-9.

23. Rawla P, Raj JP. Doxycycline-Induced Acute Pancreatitis: A Rare Adverse Event.

Gastroenterology Res. 2017;10(4):244-6.

24. Jones MR, Hall OM, Kaye AM, Kaye AD. Drug-induced acute pancreatitis: a

review. Ochsner J. 2015;15(1):45-51.

25. Singh S, Chang HY, Richards TM, Weiner JP, Clark JM, Segal JB. Glucagonlike

peptide 1-based therapies and risk of hospitalization for acute pancreatitis in type 2

diabetes mellitus: a population-based matched case-control study. JAMA Intern

Med. 2013;173(7):534-9.

26. Jalaly NY, Moran RA, Fargahi F, Khashab MA, Kamal A, Lennon AM, et al. An

Evaluation of Factors Associated With Pathogenic PRSS1, SPINK1, CTFR, and/or

CTRC Genetic Variants in Patients With Idiopathic Pancreatitis. Am J Gastroenterol.

2017;112(8):1320-9.

27. Debi U, Kaur R, Prasad KK, Sinha SK, Sinha A, Singh K. Pancreatic trauma: a

concise review. World J Gastroenterol. 2013;19(47):9003-11.

28. Abdallah AA, Krige JE, Bornman PC. Biliary tract obstruction in chronic

pancreatitis. HPB (Oxford). 2007;9(6):421-8.

29.   Kumar A, Kumar PG, Pujahari AK, Sampath S. Hypercalcemia Related

Pancreatitis. Med J Armed Forces India. 2010;66(4):385-6.

30.   Sunkara T, Caughey ME, Rawla P, Yarlagadda KS, Gaduputi V. Severe Acute

Pancreatitis as an Index Clinical Manifestation of Parathyroid Adenoma. Cureus.

2018;10(4):e2445.

31.   Prinz RA. Mechanisms of acute pancreatitis. Vascular etiology. Int J Pancreatol.

1991;9:31-8.

32.   Turkvatan A, Erden A, Turkoglu MA, Yener O. Congenital variants and anomalies

of the pancreas and pancreatic duct: imaging by magnetic resonance

cholangiopancreaticography and multidetector computed tomography. Korean J

Radiol. 2013;14(6):905-13.

33.   Lee SP, Nicholls JF, Park HZ. Biliary sludge as a cause of acute pancreatitis. N Engl

J Med. 1992;326(9):589-93.

Trends in Mortality Rate of Acute Pancreatitis in the US 2001-2014

Year	2014	2013	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003	2002	2001
Sample Size, n	35,358,818	35,597,792	36,484,846	36,962,415	37,352,013	37,734,584	38,210,889	38,155,908	38,076,556	37,843,039	37,496,978	37,074,605	36,523,831	36,093,550
Acute pancreatitis, n	279,145	274,775	275,170	276,954	277,050	262,605	260,856	246,536	249,361	243,092	246,227	238,306	235,589	215,238
Males(%)	50.4	50.52	50.3	50.54	50.11	50.32	50.67	50.72	50.82	50.81	50.96	50.79	50.6	50.61
Females(%)	52.12	52.19	52.21	52.64	52.14	52.6	52.86	53.1	53.55	53.72	53.98	54.21	54.16	55.02
Mortality(%)	0.66	0.76	0.78	0.81	0.9	0.96	0.99	0.99	1.02	1.19	1.36	1.43	1.6	1.74
Mean LOS(d)	4.6	4.7	4.7	4.9	5	5.1	5.3	5.3	5.5	5.5	5.7	5.8	5.9	6.1
Mean hospital Charges($)	35,728	34,897	33,744	34,116	32,413	31,231	30,454	28,808	26,544	25,183	23,525	23,471	20,870	19,303
Aggregate Charges(Billion $	9.98	9.58	9.27	9.43	8.98	8.21	7.94	7.09	6.64	6.13	5.8	5.6	4.92	4.15