Aversive learning strengthens episodic memory in both adolescents and adults

Alexandra O. Cohen[1], Nicholas G. Matese[1], Anastasia Filimontseva[1], Xinxu Shen[1], Tracey C. Shi[2], Ethan Livne[3], Catherine A. Hartley[1]

[1]Department of Psychology, New York University
[2]Columbia University Irving Medical Center
[3]Weizmann Institute of Science

Abstract

Adolescence is often filled with positive and negative emotional experiences that may change

how individuals remember and respond to stimuli in their environment. In adults, aversive

events can both enhance memory for associated stimuli as well as generalize to enhance

memory for unreinforced but conceptually related stimuli. The present study tested whether

learned aversive associations similarly lead to better memory and generalization across a

category of stimuli in adolescents. Participants completed an olfactory Pavlovian category

conditioning task in which trial-unique exemplars from one of two categories were partially

reinforced with an aversive odor. Participants then returned 24-hours later to complete a

surprise recognition memory test. We found better corrected recognition memory for the

reinforced versus the unreinforced category of stimuli in both adults and adolescents. Further

analysis revealed that enhanced recognition memory was driven specifically by better memory

for the reinforced exemplars. Autonomic arousal during learning was also related to subsequent

memory. These findings build on previous work in adolescent and adult humans and rodents

showing comparable acquisition of aversive Pavlovian conditioned responses across age

groups and demonstrate that memory for stimuli with an acquired aversive association is

enhanced in both adults and adolescents.

Introduction

Emotional experiences shape the information that we remember. Emotional events, particularly those that are negative, have been widely shown to enhance episodic memory in human adults (Cahill & McGaugh, 1998; Labar & Cabeza, 2006; Yonelinas & Ritchey, 2015). However, studies examining whether emotion similarly facilitates episodic memory at earlier developmental stages have yielded mixed results. Studies of autobiographical memories for emotional and neutral events in children and adolescents suggest that emotional life events are remembered more frequently and in greater detail (Bauer et al., 2017; Fivush, Hazzard, McDermott Sales, Sarfati, & Brown, 2003). In contrast, several studies assessing children's subsequent memory for images depicting intrinsically emotional stimuli have shown similar memory for negative and neutral images (Cordon, Melinder, Goodman, & Edelstein, 2013; Leventon, Stevens, & Bauer, 2014). There is some evidence to suggest that emotional information enhances memory in adolescents. Adolescents show enhanced memory for fearful faces relative to neutral faces (Pinabiaux et al., 2013) and their recall for emotional images is similar to that of adults (Vasa et al., 2011). In an attempt to reconcile these findings in more limited age-ranged samples, a recent study examined subsequent memory for negative, neutral, and positive images in 8- to 30-year-olds and showed similar emotional memory enhancement effects across ages (Stenson, Leventon, & Bauer, 2019). Taken, together, these studies suggest that across development, memory for emotional experiences may be better than memory for neutral experiences. Still, the extant research has focused on memory for events from one's own life or for intrinsically emotional stimuli. It remains unclear whether emotional learning, in which a neutral stimulus associated with an emotional experience acquires affective significance, similarly enhances subsequent memory in adolescents and adults.

Studies of emotional learning commonly model the acquisition of emotional associations through Pavlovian learning, in which a previously neutral conditioned stimulus acquires emotional salience through pairing with an intrinsically arousing positive or negative

unconditioned stimulus (LeDoux, 2000; Maren, 2001). However, in real world situations, learned

emotional associations are often more complex. For example, if someone is bitten by a dog,

they may go on to develop a negative association not only with the particular dog that bit them,

but with all dogs, or with animals more generally. The generalization and persistence in memory

of learned aversive associations are core features of anxiety disorders (American Psychiatric

Association, 2013; Dymond, Dunsmoor, Vervliet, Roche, & Hermans, 2015). Characterizing how

these cognitive processes develop is particularly important given the typical emergence and

peak in prevalence of anxiety disorders during adolescence (Kessler, Berglund, Demler, Jin, &

Walters, 2005). In children and adolescents, stimuli that are visually similar to an aversive

conditioned stimulus elicit more negative subjective emotion ratings and heightened

psychophysiological measures of arousal, suggesting a generalization of negative affective

value (Glenn et al., 2012; Michalska et al., 2016; Schiele et al., 2016). However, whether

aversive learning generalizes more broadly to facilitate subsequent memory for similar stimuli in

adolescents has yet to be investigated.

A recently developed "category conditioning" paradigm enables measurement of both

learned affective responses and their generalization to conceptually similar stimuli, as well as

the degree to which the strength and generalization of subsequent memory is influenced by

emotionally salient events. In this paradigm, trial-unique exemplars from one conceptual

category are partially reinforced with an intrinsically positive or negative stimulus, while

exemplars from another conceptual category are never reinforced (Dunsmoor, Kragel, Martin, &

La Bar, 2014; Dunsmoor, Martin, & LaBar, 2012; Patil, Murty, Dunsmoor, Phelps, & Davachi,

2017). In adults, emotional associations formed via category conditioning can generalize across

the conceptual category and lead to enhanced memory for the reinforced category of exemplars

(e.g. Dunsmoor et al., 2014, 2012; Dunsmoor, Murty, Davachi, & Phelps, 2015; Kroes,

Dunsmoor, Lin, Evans, & Phelps, 2017; Patil et al., 2017). Here we leverage this paradigm to

examine in both adults and adolescents whether emotional memory is facilitated for stimuli with

an aversive association, whether such a memory benefit generalizes to non-reinforced

exemplars within a category, and whether psychophysiological signatures of aversive learning

also generalize to conceptually similar stimuli.

In the present study, 60 participants ages 13- to 25-years-old completed a novel

olfactory Pavlovian category conditioning task, followed by a surprise recognition memory test

24-hours later (Figure 1). Our category conditioning paradigm used aversive odor as an

unconditioned stimulus, rather than mild electrical shock, as aversive odors have been

successfully used in conditioning paradigms in human (Gottfried, O'Doherty, & Dolan, 2002) and

non-human primates (U. Livneh & Paz, 2012; Uri Livneh & Paz, 2010, 2012) and can be

ethically administered in developmental populations without risk of physical harm. We chose to

administer the memory test a day after learning due to convergent evidence from previous

studies suggesting that emotional memory enhancement effects emerge with time, after at least

several hours (Yonelinas & Ritchey, 2015). The skin conductance response (SCR) to each

stimulus was collected during the category conditioning task to serve as a psychophysiological

measure of learning. Additionally, following the recognition memory test, we collected measures

of self-reported anxiety and intolerance of uncertainty, which we hypothesized might relate to

individual differences in emotional memory enhancement effects. Our primary aim was to test

whether acquired aversive associations, using odor as a reinforcer, enhance memory in

adolescents, similarly to adults, and whether these aversive associations generalize across a

category. We hypothesized that adolescents and adults would show similar facilitation of

memory for the aversively reinforced stimuli, but that adolescents might show greater

generalization of aversive associations across a category relative to adults, which might confer

heightened vulnerability to anxiety during this developmental stage.

Results

DAY 1:
Odorant Selection & Conditioning

6 s + breath

6 s

DAY 2:
Recognition Memory Test

1: Definitely Old 2: Maybe Old
3: Maybe New 4: Definitely new

1: Definitely Old 2: Maybe Old
3: Maybe New 4: Definitely new

1: Definitely Old 2: Maybe Old
3: Maybe New 4: Definitely new

1: Definitely Old 2: Maybe Old
3: Maybe New 4: Definitely new

Figure 1. Experimental design. Participants first completed an odor selection procedure (see Methods) to determine which odorant would be used as the unconditioned stimulus (US). Immediately afterwards, participants underwent aversive olfactory Pavlovian category conditioning, using a breath-triggered paradigm. Participants returned 24-hours later and completed a surprise, self-paced recognition memory test.

Recognition Memory

In line with previous category conditioning studies (Dunsmoor et al., 2014, 2012, 2015), we first examined corrected recognition memory (hit minus false alarms) for stimuli from the reinforced (CS+) versus unreinforced (CS-) category by continuous age (Figure 2A) and controlling for which category (objects or places) served as the reinforced category. We found a significant effect of stimulus type ($F(1,58) = 8.91$, $p = 0.004$), such that subjects showed better corrected recognition memory for the CS+ stimuli than the CS- stimuli. There was no significant effect of age ($F(1,57) = 0.31$, $p = 0.58$), no age by stimulus type interaction ($F(1,58) = 0.33$, $p = 0.57$), and no effect of which category was reinforced ($F(1,57) = 0.58$, $p = 0.45$).

Figure 2. Similar effects of aversive learning on recognition memory and skin conductance response across age. Across age, corrected recognition memory is better for items from the CS+ versus CS- category (A), driven by better recognition memory for the reinforced items (CS+US) (B). There was a trend towards higher skin conductance in response to CS+ items relative to CS- items (C). Participants separated by age group (Teen: 13-17, Adult: 18-25) for visualization purposes only. The corresponding statistical analyses treat age as a continuous variable. Different colored dots represent individual participants. Error bars are s.e.m. ** $p < .01$, ~ $p < .1$

To assess whether the memory benefit conferred by learned aversive associations generalized to unreinforced exemplars within the same conceptual category, we next examined hit rate by stimulus type (CS+US, CS+, and CS-) and by continuous age (Figure 2B), controlling for the reinforced category. We found a significant effect of stimulus type ($F(2,116) = 14.95$, $p < 0.0001$), but no significant effects of age ($F(1,57) = 0.69$, $p = 0.41$), no age by stimulus type interaction ($F(2,116) = 1.14$, $p = 0.32$), and no effect of which category was reinforced ($F(1,57) = 0.01$, $p = 0.93$). Post-hoc t-tests ($\alpha = .0167$, adjusted for multiple comparisons) revealed that the main effect of stimulus type was driven by better memory for the CS+US stimuli relative to the unreinforced CS+ stimuli ($t(116.59) = 3.55$, $p < 0.001$) and the CS- stimuli ($t(117.99) = 3.73$, $p < 0.001$). There was no significant difference between memory for the unreinforced CS+ stimuli relative to the CS- stimuli ($t(116.79) = 0.02$, $p = 0.99$). We also examined trial-wise memory accuracy by stimulus type (CS+US, CS+, and CS-) and by continuous age, controlling for both the reinforced category and the order of presentation of the stimuli during learning. We again found a significant effect of stimulus type ($\chi^2(2) = 43.58$, $p < .0001$) and no significant effects of

age (χ^2(1) = 0.01, *p* = .94), no age by stimulus type interaction (χ^2(2) = 3.52, *p* = .17), and no

effect of which category was reinforced (χ^2(1) = 0.33, *p* =.56). There was a significant primacy

effect of stimulus presentation order (χ^2(2) = 77.46, *p* < .0001), such that stimuli presented near

the beginning of learning were better remembered than those presented near the end. These

results suggest that improved corrected recognition memory for the CS+ category of stimuli was

driven specifically by enhanced memory for images paired with an aversive odor and was not

due to generalization of memory facilitation to non-reinforced images in the same category.

However, given numerous previous findings of such a generalization effect in category

conditioning studies conducted in adults, we also conducted a post-hoc exploratory analysis in

which we used an ordinal model (Burkner & Vuorre, 2018) to examine the influence of stimulus

type on subsequent memory, using all four confidence levels of memory responses (1 =

Definitely Old, 2 = Maybe Old, 3 = Maybe New, and 4 = Definitely New). This analysis

suggested that among the high-confidence hit stimuli, there is evidence for generalization of

memory facilitation to unreinforced CS+ stimuli (see Supplemental Materials, Tables S1 & S2)

The pattern of results reported above remained consistent when participants who were

excluded for not showing a variable skin-conductance signal during conditioning were included

in the analyses (see Supplemental Materials, Figure S1).

Psychophysiological Measure of Learning

To test for acquisition of category conditioning across adolescents and adults, we

examined average skin conductance responses for stimuli from the reinforced (CS+) versus

unreinforced (CS-) category by continuous age (Figure 2C). We found a marginal effect of

stimulus type (*F*(1,58) = 3.03, *p* = 0.09), such that subjects showed a trend towards higher skin

conductance for the CS+ stimuli relative to the CS- stimuli. There was no significant effect of

age (*F*(1,58) = 1.50, *p* = 0.23) or an age by stimulus type interaction (*F*(1,58) = 0.59, *p* = 0.45).

Psychophysiology-Recognition Memory Relationships

To gain a better understanding of the large degree of individual variability in recognition

memory, we explored relationships between psychophysiological responses during learning and

subsequent memory. First, we examined the relationship between subjects' overall hit rate and

their averaged skin conductance in response to the cues. We found a positive relationship (r(58)

= 0.26, p = 0.041), such that individuals with larger conditioned responses showed better

memory overall (Figure 3A). Further analyses (α = .025, adjusted for multiple comparisons)

revealed that this pattern was evident in the adult subjects (r(28) = 0.42, p = 0.021) but not the

adolescent subjects (r(28) = 0.06, p = 0.76; Figure 3B). This suggests that increased

psychophysiological arousal during encoding of images is related to better performance, on

average, and that this relationship is driven by the adult participants.

We next investigated the relationship between trial-evoked skin conductance responses

to cues, irrespective of the stimulus type and experienced outcome, and subsequent memory.

We first examined whether trial-evoked psychophysiological responses to the cue predicted

subsequent memory, including continuous age as a regressor of interest. We found no

significant effects of skin conductance in response to the cue (χ^2(1) = 0.01, p = .91), age (χ^2(1)

= 0.34, p = .56), and no interaction between skin conductance in response to the cue and age

(χ^2(1) = 0.003, p = .95). We also examined whether trial-evoked skin conductance responses at

the time of the outcome predicted subsequent memory, including continuous age as a regressor

of interest. We found a significant main effect (χ^2(1) = 14.11, p < .001), such that larger skin

conductance responses at the time of the outcome were associated with better memory (Figure

3C). There was no statistically significant effect of age (χ^2(1) = 0.22, p = .64), or interaction

between outcome SCR and age (χ^2(1) = 0.09, p = .77). Thus, trial-evoked SCRs at the time of

outcome were predictive of subsequent memory whereas trial-evoked SCRs to the cue itself

were not.

Figure 3. Psychophysiological arousal during learning relates to memory 24-hours later.
Participants' average skin conductance response to cue presentation was positively correlated
with their overall recognition memory performance (A). This relationship is observed in the adult
participants but not in the adolescents (B). While trial-evoked responses to the cue did not
predict subsequent memory for that item, higher responses at the time of the outcome were
predictive of better item memory (C).
232

Recognition Memory-Individual Difference Measure Relationships

To examine how individual differences in memory enhancement and generalization

might relate to participants' state or trait anxiety, and intolerance of uncertainty, we first

computed memory bias scores for CS+US (CS+US hit rate – CS- hit rate) and unreinforced CS+

(CS+ hit rate – CS- hit rate) stimuli. Consistent with our previously reported findings, linear

regressions revealed no relationships between CS+US memory bias and age ($F(1,58) = 0.19$, p

$= 0.67$) or CS+ memory bias and age ($F(1,58) = 1.78$, $p = 0.19$).

We next examined the relationships between the State Trait Anxiety Inventory (STAI)

state and trait scales (Spielberger, Gorsuch, & Lushene, 1988) and these memory bias

measures ($\alpha = .0125$, adjusted for multiple comparisons). Linear regressions revealed no

significant relationships between age and STAI state ($F(1,57) = 0.39$, $p = 0.53$), STAI trait

($F(1,57) = 1.39$, $p = 0.24$), or IUS ($F(1,57) = 0.77$, $p = 0.38$). We did not find statistically

significant relationships between the STAI state measure and either memory bias index

(CS+US memory bias, $r(57) = -0.07$, $p = 0.58$; CS+ bias, $r(57) = -0.10$, $p = 0.44$) or the STAI trait

and CS+US memory bias ($r(57) = -0.19$, $p = 0.15$). However, we did observe a negative

correlation between the STAI trait and CS+ memory bias ($r(57) = -0.35$, $p = 0.006$), such that

individuals with lower STAI trait scores showed a stronger CS+ memory bias than those with

high STAI trait scores. A follow-up analysis ($\alpha = .017$, adjusted for multiple comparisons) was

conducted to determine whether this result was due to differences in recognition memory for the

unreinforced CS+ stimuli or the CS- stimuli. We also examined the relationship between

recognition memory for the CS+US stimuli and STAI trait for completeness. We found that

neither CS+US hit rate ($r(57) = 0.09$, $p = 0.48$) nor CS+ hit rate ($r(57) = -0.11$, $p = 0.40$)

correlated with STAI trait scores. However, we observed a positive correlation between CS- hit

rate and STAI ($r(57) = 0.31$, $p = 0.016$), such that individuals with higher trait anxiety showed

better memory for the CS- stimuli (Figure 4A). Further analyses ($\alpha = .025$, adjusted for multiple

comparisons) revealed that this relationship was statistically significant in the adolescent

subjects ($r(27) = 0.49$, $p = 0.007$) but not the adult subjects ($r(28) = 0.17$, $p = 0.36$; Figure 4B).

These results indicate that higher self-reported anxiety was related to better for stimuli from the

non-reinforced category, but was not related to memory for stimuli from the reinforced category.

Figure 4. Better memory for the stimuli from the unreinforced category is associated with higher
trait anxiety. While there was no significant relationship between recognition memory for the
CS+ or CS+US stimuli and trait anxiety, there was a positive relationship between memory for
the CS- stimuli and trait anxiety (A). Follow-up analysis revealed that this positive relationship
was more pronounced in adolescents than adults (B).

We also examined the relationships between Intolerance of Uncertainty Scale (IUS)

(Freeston, Rhéaume, Letarte, Dugas, & Ladouceur, 1994) and the memory bias measures ($\alpha =$

.025, adjusted for multiple comparisons). We did not observe statistically significant correlations

between either memory bias index and IUS (CS+US memory bias, $r(57) = -0.06$, $p = 0.67$; CS+

bias, $r(57) = -0.21$, $p = 0.11$).

The pattern of results reported above remained consistent when participants who were

excluded for not showing a variable skin-conductance signal during conditioning were included

in the analyses (see Supplemental Materials, Figure S2).

Discussion

The present study employed a novel olfactory variant of a Pavlovian category

conditioning task to test whether aversive learning leads to similar memory enhancement and

generalization across a conceptual category in adults and adolescents. By using trial-unique

stimuli as "tags" for each learning trial, we show that aversive learning leads to better episodic

memory for trials associated with an aversive event in both adolescents and adults. The age

invariance of this effect is consistent with previous observations that adolescent and adult

humans and rodents exhibit equivalent acquisition of aversive Pavlovian conditioning using

simple stimuli (Deal, Erickson, Shiers, & Burman, 2016; Kim, Li, & Richardson, 2011; Pattwell et

al., 2012; Rudy, 1993). Our finding extends this literature by showing similar memory

improvements in adolescents and adults for items with an acquired aversive association.

While few studies have examined the neural mechanisms underlying emotional

facilitation of episodic memory prior to adulthood, our findings are consistent with evidence of

the early development of this circuitry. Multiple memory systems, centered on the amygdala for

emotional memory and the hippocampus for episodic and declarative memory, are proposed to

interact to facilitate memory of emotional events (Mcdonald, Devan, & Hong, 2004; Phelps,

2004). Under the "emotional binding" account of episodic memory, the amygdala binds

emotional information to an item, communicating with both the perirhinal cortex and the

hippocampus to modulate encoding, storage, and recollection of these memories (Yonelinas &

Ritchey, 2015). Evidence from rodent studies suggests that signatures of a functional emotional

memory system emerge early in development (Stanton, 2000), indicating that emotional

memory enhancement effects should be present during childhood and adolescence.

While memory for items directly associated with an aversive odor was facilitated across

age, unreinforced exemplars from the same category as the odor-paired stimuli were not better

remembered in either adults or adolescents. This result does not replicate previous studies

showing that emotional associations generalize across a category and lead to enhanced

memory for the reinforced category of exemplars in adults (Dunsmoor et al., 2014, 2012, 2015;

Kroes et al., 2017; Patil et al., 2017). However, we did find some evidence of increased correct

high-confidence memory judgements for both odor-paired stimuli and unreinforced stimuli of the

same category, relative to stimuli that were never reinforced. This indicates a possible

interaction between metacognitive ability and memory generalization effects, such that

generalization of memory facilitation is primarily observed when examining strong memories.

Another reason that we may not have fully replicated prior studies is our use of a different

primary reinforcer. Previous work has shown that the intensity of the aversive stimulus is related

to the degree of generalization (Dunsmoor, Kroes, Braren, & Phelps, 2017), suggesting that

olfactory reinforcers may not be potent enough to induce widespread generalization effects.

Further studies comparing aversive learning across different modality reinforcers (e.g. shock

versus noise versus odor) and manipulating the duration and intensity of reinforcement will be

necessary to determine the effectiveness of odor conditioning in producing generalization

effects.

In this study, we used cue-evoked skin conductance response as a psychophysiological

measure of emotional learning. Moreover, in this category condition paradigm, the measure of

anticipatory arousal also provides a measure of the degree to which learned aversive

associations generalize across a conceptual category. Skin conductance responses showed a

trend towards increased anticipatory arousal for the reinforced category of stimuli across

participants. While this marginal increase in anticipatory arousal indicates some degree of

learning of the association between the partially reinforced category and a potential aversive

outcome, evidence for emotional learning in our study was weak. In the current experiment, we

used skin conductance response as a psychophysiological index of learning due to the

prevalence of this measure in the human conditioning literature (Bradley, Miccoli, Escrig, &

Lang, 2008; Hamm & Stark, 1993; LaBar, Gatenby, Gore, LeDoux, & Phelps, 1998; LANG,

GREENWALD, BRADLEY, & HAMM, 1993). Other psychophysiological measures of learning,

such as pupillometry (Leuchs, Schneider, & Spoormaker, 2018) and breathing measures (Uri

Livneh & Paz, 2010) should also be examined to determine whether they might provide more

robust indices of learning dynamics during olfactory conditioning.

In order to probe individual variability in aversive learning and memory, we examined

relationships between skin conductance responses during learning and subsequent memory.

Previous studies in adults have shown that autonomic arousal during encoding is associated

with better memory (Bradley, Greenwald, Petry, & Lang, 1992; Kensinger EA & Corkin S, 2004;

Kleinsmith & Kaplan, 1963). In both humans and rodents (Glascher, Adolphs, Gläscher, &

Adolphs, 2003; Mather, Clewett, Sakaki, & Harley, 2016; Reis & LeDoux, 1987; Reyes,

Carvalho, Vakharia, & Van Bockstaele, 2011; Roozendaal, Luyten, de Voogd, & Hermans,

2016), the amygdala can modulate noradrenergic autonomic arousal responses to aversive

stimuli, providing a putative mechanism through which emotion might influence memory.

Consistent with this prior work, we found that individuals showing higher anticipatory arousal in

response to cues on average, throughout the task, also showed better memory overall. This

result was driven primarily by the adult participants and not the adolescents. However, in

accordance with previous findings (de Voogd, Fernández, & Hermans, 2016), trial-evoked

anticipatory arousal to the cue did not predict subsequent memory for the corresponding trial.

We instead found that trial-evoked responses at the time of the outcome predicted memory 24-

hours later. These data suggest that while increased anticipatory arousal during learning may

foster a general memory benefit, unlearned autonomic arousal reactions to individual stimuli

predict whether or not that stimulus will be remembered at a later time.

Finally, we examined how individual difference measures related to subsequent

memory. Unexpectedly, we found positive relationship between recognition memory for CS-

stimuli and trait anxiety, such that individuals with higher trait anxiety showed better memory for

items from the category that was never reinforced. A follow-up analysis suggested that this

correlation was largely driven by adolescent participants. This surprising finding supports the

idea that overgeneralization of aversive experiences to dissimilar stimuli is a defining feature of

anxiety (Dymond et al., 2015) and that this overgeneralization may be more readily observable

during adolescence, the period of development in which anxiety disorders often first emerge

(Kessler et al., 2005; Lee et al., 2014). However, it is noteworthy that trait anxiety was not

correlated with memory for the specific items associated with aversive events (CS+US stimuli)

or the unreinforced items from that same category (CS+ stimuli). Given the exploratory nature of

these results, replication and further investigation of the relationship between generalization,

overgeneralization, and trait anxiety in adolescents and adults is warranted.

In summary, the current study demonstrated that aversive learning enhances episodic

memory in both adolescents and adults, particularly for items directly associated with an

aversive odor. We found that autonomic arousal during learning was related to later memory.

Specifically, unlearned arousal responses to outcomes during encoding were predictive of

subsequent memory for individual stimuli. These results indicate that aversive odors are

sufficiently evocative to induce memory enhancements in both adolescents and adults. While

further refinement of olfactory conditioning methods for use in developmental populations is

necessary, this study suggests that aversive and appetitive odors might be fruitfully utilized to

study emotional learning and memory processes across development.

Materials and Methods

Participants

Sixty participants between the ages of 13 and 25 years (mean age = 18.69, 30 female)

were included in analyses. A target sample size of n = 60, including 30 adolescents and 30

adults, was determined based on group sample sizes previously reported in category

conditioning studies (Dunsmoor et al., 2014, 2012, 2015). Data from 28 additional participants

(mean age = 19.45, 18 female) were excluded from primary analyses for not showing a variable

skin-conductance signal (defined as fewer than four scorable trials) during conditioning. Data

from 22 additional participants were excluded from analyses due to the discovery of a software

bug that yielded inconsistencies in timing and delivery of the aversive reinforcers. Three

additional participants were excluded from analyses due to failure to return for the second

session of the study. All participants were volunteers from a community sample of New York

City. Of the 60 participants included in primary analyses, 45% of participants self-identified as

Caucasian/White, 15% as African American, 25% as Asian, and 15% as mixed race.

Additionally, 16.67% of the sample identified as Hispanic. Of the 28 participants who did not

exhibit a variable skin conductance signal (but were included in the supplemental analyses of

the memory data), 32.14% of participants self-identified as Caucasian/White, 21.43% as African

American, 39.29% as Asian, and 7.14% as mixed race. Additionally, 7.14% of the sample

identified as Hispanic. Participants were screened for difficulties seeing without corrective

lenses (as the nasal mask precluded the simultaneous use of glasses; contact lenses were

permitted), history of psychiatric diagnoses, use of psychoactive medication or beta blockers, or

difficulties breathing. Participants provided informed written consent or assent per research

procedures approved by New York University's Institutional Review Board. Parents or guardians

of teenagers under age eighteen also provided written consent on behalf of the teenager, prior

to their participation in the study. All participants were compensated $30 for their participation in

two approximately 1-hour sessions scheduled 24-hours apart.

Olfactory Pavlovian Category Conditioning Paradigm

A category conditioning paradigm used in previous studies in healthy adults (Dunsmoor

et al., 2014, 2012) was adapted for use with a custom built olfactometer, allowing for odorants to

serve as the unconditioned stimulus (US). The breath-triggered conditioning paradigm consisted

of four 12-trial blocks, where each trial was a unique exemplar from one of two conceptual

categories. Over the course of conditioning, subjects viewed 24 unique objects and 24 unique

scenes (Konkle & Caramazza, 2013). Each stimulus category (object or scene) was randomly

assigned to serve as the reinforced conditioned stimulus category (CS+) for half the participants

and the unreinforced conditioned stimulus category (CS-) for the other half of participants.

Within each 12-trial block, half of the trials were exemplars from the CS+ category and half were

from the CS- category. The CS- trials were never paired with an odor and the CS+ trials were

reinforced 50% of the time (three CS+US, three CS+, and six CS- trials per block, resulting in 12

trials total). Trial order was pseudorandomized such that no more than two reinforced (CS+US)

trials and no more than three exemplars from the same category appeared in a row. Four

different trial orders and two possible assignments the reinforced category of stimuli resulted in

eight different versions of the task that were administered to participants.

During conditioning, participants passively viewed images presented on the screen via

Psychtoolbox-3 in Matlab R2015b while breathing through a nasal mask connected to the

olfactometer. They were instructed simply to notice any associations between the pictures and

smells. Clean air was continuously circulated through the mask and participants' nasal breathing

was measured via pressure sensors in the olfactometer and processed in real-time using

LabVIEW 2016 Version 16.0f5 (64-bit). The paradigm used breath-triggered stimulus

presentation to insure that odor delivery was timed to a participants' inhalation. Each trial began

with a fixation cross presented for a fixed interval of six seconds. On the subsequent inhale after

the fixation (variable duration), a trial-unique exemplar appeared on the screen for a fixed

interval of six seconds. After a one-second buffer to ensure separation of respiratory cycles (Uri

Livneh & Paz, 2010), the participant's next inhale while the image was still on the screen

triggered an olfactometer "shoot" event for two seconds. For CS- and unreinforced CS+ trials,

this shoot event consisted of continued release of clean air and for the CS+US trials, the

aversive odor was delivered. If the participant did not inhale while the image was on the screen,

an olfactometer shoot event was not triggered and the participant only experienced clean air. If

this occurred on a CS+US trial, this trial was reclassified as a CS+ image without reinforcement

during data processing and in subsequent analyses. Twelve of the 60 participants missed at

least one odor shoot event. Nine of the 12 missed a single shoot event, leading to a 45.83%

reinforcement rate, two missed two shoot events, leading to a 41.67% reinforcement rate, and

one participant missed three shoot events, leading to a 37.5% reinforcement rate.

Odorant Selection

At the beginning of the first session, participants underwent an odor selection procedure

to identify the odorant to be used as the aversive reinforcer in the category conditioning

paradigm. This procedure was designed to take into account individual differences in whether

an odorant is considered to be aversive, mirroring calibration procedures that are typically

performed in aversive learning studies using mild electrical shock as the aversive reinforcer

(e.g. Dunsmoor et al., 2014; Dunsmoor, Mitroff, & LaBar, 2009). Each odorant was rated on

valence and arousal using a modified version of the Self-Assessment Manikin (SAM) (Bradley,

M. M. & Lang, P. J., 1994). A suite of eight different aversive odorants, supplied by DreamAir

perfumers, were first administered to participants using Whispi air puff canisters (Scentovation,

Novia Products, LLC). Three of the odorants were the following chemical compounds: isovaleric

aldehyde 10% diluted in isopropyl myristate, dimethyl acetate undecadienol, and Ozonil^{TM}

(tridec2-ene nitrile). Five of the odorants were proprietary DreamAir odorant blends ("Bad smell

3", "Horse hair", "Fumier", "Frog 3", and "Fear 45l"). Participants were asked to rate the valence

of the smell on a scale from one to nine in which a one represented a bad smell, labeled 'Don't

Like' on the scale, and a nine represented a good smell, labeled 'Like' on the scale. Immediately

following the valence rating, the arousal rating measured the perceived strength of the smell on

a scale of one to nine in which one represented a 'Weak', unnoticeable odor and a nine

represented a 'Strong', noticeable odor.

452   Each smell was presented and ranked on these scales three times, and the average

scores for each odorant were computed to determine the four most aversive odors, as indexed

by ratings of lowest valence and highest strength. The four most aversive odors were then

presented through the nasal mask via inhale-triggered odor release delivered using the

olfactometer, which allowed the participants to experience the odors as they would during

conditioning. Participants were asked to rate the four odorants three times each on a scale from

one to five, where one indicated the smell was bad and five indicated that the smell was so bad

that the participant would not be able to handle smelling it several times during the conditioning

task. Ratings were averaged and the odorant with the highest average of a score of four or

below, meaning that the odor never received a rating of five, was used in the conditioning task.

Surprise Recognition Memory Test

463   Participants returned 24-hours later for a surprise recognition memory test presented via

MATLAB's Psychtoolbox-3. The self-paced memory test included the 24 CS+ and 24 CS-

category exemplars from day one, as well as 24 new objects and 24 new scenes, for a total of

96 images. Images used in the task on day 1 and as new images on day 2 were

counterbalanced across participants. Participants rated whether each picture was new or old on

a four-point scale: 1 = Definitely Old, 2 = Maybe Old, 3 = Maybe New, and 4 = Definitely New.

Consistent with previous studies, responses were collapsed across new versus old ratings. We

examined corrected recognition memory, which is a difference score between hits, old images

correctly identified as old, and false alarms, new images incorrectly identified as old.

Additionally, we examined hit rate for the CS+US, CS+, and CS- images to look for

generalization across the reinforced category of exemplars.

Psychophysiological Data Acquisition & Analysis

Skin conductance data was recording during the conditioning paradigm using a BIOPAC

MP-100 System (Goleta, CA). Pre-gelled SCR electrodes were placed on the hypothenar

eminence of the palm (Dunsmoor et al., 2015) of the non-dominant hand and the phasic skin

conductance response (SCR) to each CS onset and outcome timepoint (US or no US) were

scored using AcqKnowledge 3.9 software (BIOPAC Systems). SCR data were low-pass filtered

and smoothed. SCR scores were based on the window 0.5 seconds after stimulus onset to 0.5

seconds after shoot onset and outcome response scores were based on the window 0.5

seconds after shoot onset to 0.5 seconds after shoot offset. The trough-to-peak difference of the

first waveform (in μSiem) (Dunsmoor et al., 2015; Hermans et al., 2017) beginning within these

windows was measured. Using MATLAB R2016a, distributions were normalized using square

root transformation of the raw SCR magnitudes, and then divided by the maximum response

(across all cue and outcome responses) to enable between-subject comparison. Any trial

without a shoot event was considered missing for analyses that examined SCR at outcome.

Self-Report Measures

Following the recognition memory test on day two, participants completed several self-

report measures via Qualtrics surveys. Participants completed the State Trait Anxiety Inventory

(STAI) state and trait scales (Spielberger et al., 1988), the Intolerance of Uncertainty Scale

(IUS) for adults or the IUS-C for teenagers (Freeston et al., 1994), and a free response question

asking whether the subject noticed anything about the types of images that were paired with

smells.

Analysis Approach

Data processing was completed in MATLAB R2016 and all statistical analyses were

conducted in R version 3.5.1 (R Core team, 2016). Mixed-effects models were run using the

'lme4' package (version 1.1-17) lmer (for analyzing recognition memory and average skin

conductance response) and glmer (for trial-wise analyses) functions (Bates D, Maechler M,

Bolker B, & Walker S, 2015). Numeric variables included as regressors in the model were z-

scored across all participants. Each model included a random intercept for each participant.

Statistics were reported from analysis of variance (Type III using Satterthwaite's method)

performed on lmer models and analysis of deviance (Type III Wald chi-square tests) performed

on glmer models. Welch two sample t-tests were performed for post-hoc analyses of recognition

memory data and Pearson product-moment correlations were computed for all reported

correlations. Where applicable, statistical significance thresholds (alphas) adjusted for multiple

comparisons are reported in the Results section.

Acknowledgements

We thank DreamAir, LLC for providing us with the odorants used in this study, Noam Sobel and

his engineering team for building our olfactometer, and the Frueauff Foundation for a generous

equipment grant that funded its construction. This work was supported by a Klingenstein-

Simons Fellowship in Neuroscience, a Jacobs Foundation Research Fellowship, and a National

Science Foundation CAREER Award Grant No. 1654393 (to C.A.H.) as well as a National

Science Foundation SBE Postdoctoral Research Fellowship Grant No. 1714321 (to A.O.C).

References

American Psychiatric Association. (2013). *Diagnostic and Statistical Manual DSM 5. Diagnostic*

and Statistical Manual of Mental Disorders, 5th Edition.

https://doi.org/10.1176/appi.books.9780890425596.744053

Bates D, Maechler M, Bolker B, & Walker S. (2015). lme4. *Journal of Statistical Software*, *67*(1),

1–48. https://doi.org/10.18637/jss.v067.i01

Bauer, P. J., Stark, E. N., Ackil, J. K., Larkina, M., Merrill, N., & Fivush, R. (2017). The

recollective qualities of adolescents' and adults' narratives about a long-ago tornado.

Memory, *25*(3), 412–424. https://doi.org/10.1080/09658211.2016.1180396

Bradley, M. M., & Lang, P. J. (1994). Measuring emotion: the self-assessment manikin and the

semantic differential. *Journal of Behavior Therapy and Experimental Psychiatry*, *25*(1), 49–

59. https://doi.org/10.1016/0005-7916(94)90063-9

Bradley, M. M., Greenwald, M. K., Petry, M. C., & Lang, P. J. (1992). Remembering Pictures:

Pleasure and Arousal in Memory. *Journal of Experimental Psychology: Learning, Memory,*

and Cognition, *18*(2), 379–390. https://doi.org/10.1037/0278-7393.18.2.379

Bradley, M. M., Miccoli, L., Escrig, M. A., & Lang, P. J. (2008). The pupil as a measure of

emotional arousal and autonomic activation. *Psychophysiology*, *45*(4), 602–607.

https://doi.org/10.1111/j.1469-8986.2008.00654.x

Burkner, P.-C., & Vuorre, M. (2018). Ordinal Regression Models in Psychology: A Tutorial.

PsyArXiv, 1–17. https://doi.org/10.31234/osf.io/x8swp

Cahill, L., & McGaugh, J. L. (1998). Mechanisms of emotional arousal and lasting declarative

memory. *Trends in Neurosciences*, *21*(7), 294–299. Retrieved from

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&l

ist_uids=9683321

Cordon, I. M., Melinder, A. M. D., Goodman, G. S., & Edelstein, R. S. (2013). Children's and

adults' memory for emotional pictures: Examining age-related patterns using the

Developmental Affective Photo System. *Journal of Experimental Child Psychology*, *114*(2),

339–356. https://doi.org/10.1016/j.jecp.2012.08.004

Deal, A. L., Erickson, K. J., Shiers, S. I., & Burman, M. A. (2016). Limbic system development

underlies the emergence of classical fear conditioning during the third and fourth weeks of

life in the rat. *Behavioral Neuroscience*, *130*(2), 212–230.

https://doi.org/10.1037/bne0000130

de Voogd, L. D., Fernández, G., & Hermans, E. J. (2016). Disentangling the roles of arousal and

amygdala activation in emotional declarative memory. *Social Cognitive and Affective*

Neuroscience, *11*(9), 1471–1480. https://doi.org/10.1093/scan/nsw055

Dunsmoor, J. E., Kragel, P. A., Martin, A., & La Bar, K. S. (2014). Aversive learning modulates

cortical representations of object categories. *Cerebral Cortex*, *24*(11), 2859–2872.

https://doi.org/10.1093/cercor/bht138

Dunsmoor, J. E., Kroes, M. C. W., Braren, S. H., & Phelps, E. A. (2017). Threat intensity widens

fear generalization gradients. *Behavioral Neuroscience*, *131*(2), 168–175.

https://doi.org/10.1037/bne0000186

Dunsmoor, J. E., Martin, A., & LaBar, K. S. (2012). Role of conceptual knowledge in learning

and retention of conditioned fear. *Biological Psychology*, *89*(2), 300–305.

https://doi.org/10.1016/j.biopsycho.2011.11.002

Dunsmoor, J. E., Mitroff, S. R., & LaBar, K. S. (2009). Generalization of conditioned fear along a

dimension of increasing fear intensity. *Learning and Memory*, *16*(7), 460–469.

https://doi.org/10.1101/lm.1431609

Dunsmoor, J. E., Murty, V. P., Davachi, L., & Phelps, E. A. (2015). Emotional learning

selectively and retroactively strengthens memories for related events. *Nature*, *520*(7547),

345–348. https://doi.org/10.1038/nature14106

Dymond, S., Dunsmoor, J. E., Vervliet, B., Roche, B., & Hermans, D. (2015). Fear

Generalization in Humans: Systematic Review and Implications for Anxiety Disorder

Research. *Behavior Therapy*, *46*(5), 561–582. https://doi.org/10.1016/j.beth.2014.10.001

Fivush, R., Hazzard, A., McDermott Sales, J., Sarfati, D., & Brown, T. (2003). Creating

coherence out of Chaos? Children's narratives of emotionally positive and negative events.

Applied Cognitive Psychology, *17*(1), 1–19. https://doi.org/10.1002/acp.854

Freeston, M. H., Rhéaume, J., Letarte, H., Dugas, M. J., & Ladouceur, R. (1994). Why do

people worry? *Personality and Individual Differences*, *17*(6), 791–802.

https://doi.org/10.1016/0191-8869(94)90048-5

Glascher, J., Adolphs, R., Gläscher, J., & Adolphs, R. (2003). Processing of the arousal of

subliminal and supraliminal emotional stimuli by the human amygdala. *J Neurosci*,

23(1529–2401 (Electronic)), 10274–10282. https://doi.org/23/32/10274 [pii]

Glenn, C. R., Klein, D. N., Lissek, S., Britton, J. C., Pine, D. S., & Hajcak, G. (2012). The

development of fear learning and generalization in 8-13 year-olds. *Developmental*

Psychobiology, *54*(7), 675–684. https://doi.org/10.1002/dev.20616

Gottfried, J. A., O'Doherty, J., & Dolan, R. J. (2002). Appetitive and aversive olfactory learning in humans studied using event-related functional magnetic resonance imaging. *The Journal of Neuroscience*, *22*(24), 10829–10837. https://doi.org/22/24/10829 [pii]

Hamm, A. O., & Stark, R. (1993). Sensitization and aversive conditioning: Effects on the startle reflex and electrodermal responding. *Integrative Physiological and Behavioral Science*, *28*(2), 171–176. https://doi.org/10.1007/BF02691223

Hermans, E. J., Kanen, J. W., Tambini, A., Fernández, G., Davachi, L., & Phelps, E. A. (2017). Persistence of Amygdala-Hippocampal Connectivity and Multi-Voxel Correlation Structures During Awake Rest After Fear Learning Predicts Long-Term Expression of Fear. *Cerebral Cortex*, *27*(5), 3028–3041. https://doi.org/10.1093/cercor/bhw145

Kensinger EA, & Corkin S. (2004). Two routes to emotional memory: distinct neural processes for valence and arousal. *Proceedings of the National Academy of Sciences of the United States of America*, *101*(9), 3310–5. https://doi.org/10.1073/pnas.0306408101

Kessler, R. C., Berglund, P., Demler, O., Jin, R., & Walters, E. E. (2005). Lifetime Prevalence and Age-of-Onset Distributions of DSM-IV Disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, *62*(6), 593–602. Retrieved from http://search.ebscohost.com/login.aspx?direct=true&db=psyh&AN=2005-06168-003&site=ehost-live&scope=site

Kim, J. H., Li, S., & Richardson, R. (2011). Immunohistochemical analyses of long-term extinction of conditioned fear in adolescent rats. *Cerebral Cortex*, *21*(3), 530–538.

Kleinsmith, L. J., & Kaplan, S. (1963). Paired-associate learning as a function of arousal and interpolated interval. *Journal of Experimental Psychology*, *65*(2), 190–193. https://doi.org/10.1037/h0040288

Konkle, T., & Caramazza, A. (2013). Tripartite Organization of the Ventral Stream by Animacy and Object Size. *Journal of Neuroscience*, *33*(25), 10235–10242.

605  https://doi.org/10.1523/JNEUROSCI.0983-13.2013

606 Kroes, M. C. W., Dunsmoor, J. E., Lin, Q., Evans, M., & Phelps, E. A. (2017). A reminder before

607  extinction strengthens episodic memory via reconsolidation but fails to disrupt generalized

608  threat responses. *Scientific Reports*, *7*(1). https://doi.org/10.1038/s41598-017-10682-7

609 Labar, K. S., & Cabeza, R. (2006). Cognitive neuroscience of emotional memory. *Nature*

610  *Reviews Neuroscience*, *7*, 54–64. https://doi.org/10.1038/nrn1825

611 LaBar, K. S., Gatenby, J. C., Gore, J. C., LeDoux, J. E., & Phelps, E. A. (1998). Human

612  amygdala activation during conditioned fear acquisition and extinction: a mixed-trial fMRI

613  study. *Neuron*, *20*(5), 937–945. Retrieved from

614  http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&l

615  ist_uids=9620698

616 LANG, P. J., GREENWALD, M. K., BRADLEY, M. M., & HAMM, A. O. (1993). Looking at

617  pictures: Affective, facial, visceral, and behavioral reactions. *Psychophysiology*, *30*(3),

618  261–273. https://doi.org/10.1111/j.1469-8986.1993.tb03352.x

619 LeDoux, J. E. (2000). Emotion circuits in the brain. *Annu Rev Neurosci*, *23*, 155–184.

620 Lee, F. S., Heimer, H., Giedd, J. N., Lein, E. S., Šestan, N., Weinberger, D. R., & Casey, B. J.

621  (2014). Adolescent mental health— Opportunity and obligation. *Science*, *346*(6209), 547–

622  549.

623 Leuchs, L., Schneider, M., & Spoormaker, V. I. (2018). Measuring the conditioned response: A

624  comparison of pupillometry, skin conductance, and startle electromyography.

625  *Psychophysiology*. https://doi.org/10.1111/psyp.13283

626 Leventon, J. S., Stevens, J. S., & Bauer, P. J. (2014). Development in the neurophysiology of

627  emotion processing and memory in school-age children. *Developmental Cognitive*

628  *Neuroscience*, *10*, 21–33. https://doi.org/10.1016/j.dcn.2014.07.007

629 Livneh, U., & Paz, R. (2010). An implicit measure of olfactory performance for non-human

630  primates reveals aversive and pleasant odor conditioning. *Journal of Neuroscience*

Methods, *192*(1), 90–95. https://doi.org/10.1016/j.jneumeth.2010.07.027

Livneh, U., & Paz, R. (2012). Amygdala-prefrontal synchronization underlies resistance to

extinction of aversive memories. *Neuron*, *75*(1), 133–142.

https://doi.org/10.1016/j.neuron.2012.05.016

Livneh, U., & Paz, R. (2012). Aversive-Bias and Stage-Selectivity in Neurons of the Primate

Amygdala during Acquisition, Extinction, and Overnight Retention. *Journal of*

Neuroscience, *32*(25), 8598–8610. https://doi.org/10.1523/JNEUROSCI.0323-12.2012

Maren, S. (2001). Neurobiology of Pavlovian Fear Conditioning. *Annual Review of*

Neuroscience, *24*(1), 897–931. https://doi.org/10.1146/annurev.neuro.24.1.897

Mather, M., Clewett, D., Sakaki, M., & Harley, C. W. (2016). GANEing traction: The broad

applicability of NE hotspots to diverse cognitive and arousal phenomena. *The Behavioral*

and Brain Sciences, *39*, e228. https://doi.org/10.1017/S0140525X16000017

Mcdonald, R. J., Devan, B. D., & Hong, N. S. (2004). Multiple memory systems : The power of

interactions. *Neurobiology of Learning and Memory*, *82*, 333–346.

https://doi.org/10.1016/j.nlm.2004.05.009

Michalska, K. J., Shechner, T., Hong, M., Britton, J. C., Leibenluft, E., Pine, D. S., & Fox, N. A.

(2016). A developmental analysis of threat/safety learning and extinction recall during

middle childhood. *Journal of Experimental Child Psychology*, *146*, 95–105.

https://doi.org/10.1016/j.jecp.2016.01.008

Patil, A., Murty, V. P., Dunsmoor, J. E., Phelps, E. A., & Davachi, L. (2017). Reward

retroactively enhances memory consolidation for related items. *Learning and Memory*,

24(1), 65–69. https://doi.org/10.1101/lm.042978.116

Pattwell, S. S., Duhoux, S., Hartley, C. a, Johnson, D. C., Jing, D., & Elliott, M. D. (2012).

Altered fear learning across development in both mouse and human. *Proc Natl Acad Sci*,

109(40), 16318–16323. https://doi.org/10.1073/pnas.1206834109/-

/DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1206834109

Phelps, E. A. (2004). Human emotion and memory: interactions of the amygdala and

hippocampal complex. *Current Opinion in Neurobiology*, *14*(2), 198–202. Retrieved from

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&l

ist_uids=15082325

Pinabiaux, C., Hertz-Pannier, L., Chiron, C., Rodrigo, S., Jambaque, I., & Noulhaine, M. (2013).

Memory for fearful faces across development : specialization of amygdala nuclei and

medial temporal lobe structures. *Frontiers in Human Neuroscience*, *7*, 1–14.

https://doi.org/10.3389/fnhum.2013.00901

R Core team. (2016). R Core Team. *R: A Language and Environment for Statistical Computing.*

R Foundation for Statistical Computing , Vienna, Austria. ISBN 3-900051-07-0, URL

http://www.R-project.org/.R Core Team (2016). R: A Language and Environment for

Statistical Computing. R Foundation . https://doi.org/10.1152/ajpgi.00069.2014

Reis, D. J., & LeDoux, J. E. (1987). Some central neural mechanisms governing resting and

behaviorally coupled control of blood pressure. *Circulation*, *76(Suppl 1*, I2–I9.

Reyes, B. A. S., Carvalho, A. F., Vakharia, K., & Van Bockstaele, E. J. (2011). Amygdalar

peptidergic circuits regulating noradrenergic locus coeruleus neurons: Linking limbic and

arousal centers. *Experimental Neurology*, *230*(1), 96–105.

https://doi.org/10.1016/j.expneurol.2011.04.001

Roozendaal, B., Luyten, L., de Voogd, L. D., & Hermans, E. J. (2016). Importance of amygdala

noradrenergic activity and large-scale neural networks in regulating emotional arousal

effects on perception and memory. *The Behavioral and Brain Sciences*, *39*, e222.

https://doi.org/10.1017/S0140525X15001934

Rudy, J. W. (1993). Contextual Conditioning and Auditory Cue Conditioning Dissociate During

Development. *Behavioral Neuroscience*, *107*(5), 887–891. https://doi.org/10.1037/0735-

7044.107.5.887

Schiele, M. A., Reinhard, J., Reif, A., Domschke, K., Romanos, M., Deckert, J., & Pauli, P.

(2016). Developmental aspects of fear: Comparing the acquisition and generalization of

conditioned fear in children and adults. *Developmental Psychobiology*, *58*(4), 471–481.

https://doi.org/10.1002/dev.21393

Spielberger, C. D., Gorsuch, R. L., & Lushene, R. E. (1988). *STAI-manual for the State Trait*

Anxiety Inventory. Palo Alto (CA): Consulting Psychologists Press.

Stanton, M. E. (2000). Multiple memory systems, development and conditioning. In *Behavioural*

Brain Research (Vol. 110, pp. 25–37). https://doi.org/10.1016/S0166-4328(99)00182-5

Stenson, A. F., Leventon, J. S., & Bauer, P. J. (2019). Emotion effects on memory from

childhood through adulthood: Consistent enhancement and adult gender differences.

Journal of Experimental Child Psychology, *178*, 121–136.

https://doi.org/10.1016/j.jecp.2018.09.016

Vasa, R. A., Pine, D. S., Thorn, J. M., Nelson, T. E., Spinelli, S., Maheu, F. S., … Stewart, H.

(2011). Enhance Right Amygdala Activity in Adolescents during Encoding of Positively-

Valenced Pictures. *Developmental Cognitive Neuroscience*, *1*(1), 88–99.

https://doi.org/10.1016/j.dcn.2010.08.004.Enhanced

Yonelinas, A. P., & Ritchey, M. (2015). The slow forgetting of emotional episodic memories : an

emotional binding account. *Trends in Cognitive Sciences*, *19*(5), 259–267.

https://doi.org/10.1016/j.tics.2015.02.009

Figure Legends

Figure 1. Experimental design. Participants first completed an odor selection procedure (see

Methods) to determine which odorant would be used as the unconditioned stimulus (US).

Immediately afterwards, participants underwent aversive olfactory Pavlovian category

conditioning, using a breath-triggered paradigm. Participants returned 24-hours later and

completed a surprise, self-paced recognition memory test.

Figure 2. Similar effects of aversive learning on recognition memory and skin conductance

response across age. Across age, corrected recognition memory is better for items from the

CS+ versus CS- category (A), driven by better recognition memory for the reinforced items

(CS+US) (B). There was a trend towards higher skin conductance in response to CS+ items

relative to CS- items (C). Participants separated by age group (Teen: 13-17, Adult: 18-25) for

visualization purposes only. The corresponding statistical analyses treat age as a continuous

variable. Different colored dots represent individual participants. Error bars are s.e.m. ** $p < .01$,

~ $p < .1$

Figure 3. Psychophysiological arousal during learning relates to memory 24-hours later.

Participants' average skin conductance response to cue presentation was positively correlated

with their overall recognition memory performance (A). This relationship is observed in the adult

participants but not in the adolescents (B). While trial-evoked responses to the cue did not

predict subsequent memory for that item, higher responses at the time of the outcome were

predictive of better item memory (C).

Figure 4. Better memory for the stimuli from the unreinforced category is associated with higher

trait anxiety. While there was no significant relationship between recognition memory for the

CS+ or CS+US stimuli and trait anxiety, there was a positive relationship between memory for

the CS- stimuli and trait anxiety (A). Follow-up analysis revealed that this positive relationship

was more pronounced in adolescents than adults (B).

728