Research articles

Title of Article: Exercise twice-a-day potentiates markers of mitochondrial biogenesis

in men

Author's Affiliation:
[1]Department of Physical Education and Sports Science, Academic Center of Vitoria,
Federal University of Pernambuco, Vitoria de Santo Antao, Pernambuco, Brazil.
[2]Department of Medicine, University of Missouri School of Medicine, Columbia,
Missouri, USA.
[3]School of Exercise and Nutrition Sciences, Faculty of Health, Deakin University,
Burwood, Victoria, Australia.
[4]Institute for Health and Sport, Victoria University, Melbourne, Victoria, Australia.
[5]Endurance Performance Research Group, School of Physical Education and Sport,
University of São Paulo, Sao Paulo, Brazil.
[6]School of Medical and Health Sciences, Edith Cowan University, Joondalup, Western
Australia, Australia.
[7]Human Performance Research Group, Academic Department of Physical Education,
Technological Federal University of Parana, Curitiba, Parana, Brazil.
20

Authors: Victor A. Andrade-Souza[1], Thaysa Ghiarone[1], Andre Sansonio[1], Kleiton

Augusto Santos Silva[1,2], Fabiano Tomazini[1], Lucyana Arcoverde[1], Jackson Fyfe[3],

Enrico Perri[4], Nicholas Saner[4], Jujiao Kuang[4], Romulo Bertuzzi[5], Carol Gois Leandro[1],

David J. Bishop[4,6]* and Adriano E. Lima-Silva[1,7]

Correspondence and requests for materials should be addressed to:
David John Bishop
Institute for Health and Sport (iHeS), Victoria University. PO Box 14428, Melbourne
VIC   8001,   Australia.   Phone: +61   3   9919   9471.   E-mail   address:
David.Bishop@vu.edu.au

Conflict of Interest: The authors declare no conflict of interest.

Running   Title: Exercise   twice-a-day   potentiates   mitochondrial   biogenesis

Nonstandard Abbreviations

TFEB, transcription factor EB; NAFT, nuclear factor of activated T cells; PGC-1α, peroxisome proliferator-activated receptor-□ coactivator 1 alpha; PPARα, peroxisome proliferator-activated receptor alpha; PPARβ/δ, peroxisome proliferator-activated receptor beta/delta; CHO, carbohydrate; p53, p53 protein; PHF20, PHF20 protein; p38MAPK, p38 mitogen-activated protein kinase; AMPK, 5' adenosine monophosphate-activated protein kinase; COX IV, cytochrome c oxidase subunit IV; CPT1, carnitine palmitoyltransferase 1; NDUF, NADH: ubiquinone oxidoreductase; SDH, succinate dehydrogenase; GLUT4, Glucose transporter type 4; β-HAD, 3-hydroxyacyl-CoA dehydrogenase; CD36, fatty acid translocase cluster of differentiation 36; PFK, phosphofructokinase; CS, citrate synthase; PPARα, peroxisome proliferator-activated receptor alpha; PPAR□, peroxisome proliferator-activated receptor delta; UCP3, uncoupling protein 3; Tfam, mitochondrial transcription factor A; PDK4, pyruvate dehydrogenase kinase 4; CHCHD4, coiled-coil-helix-coiled-coil-helix domain containing 4; p21, p21 protein; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; B2M, β-2-microglobulin; TBP, TATA-box binding protein; 18S, 18S ribosomal RNA; ACTB, actin beta.

Abstract

Endurance exercise begun with reduced muscle glycogen stores seems to potentiate skeletal muscle protein abundance and gene expression. However, it is unknown whether this greater signalling responses is due to performing two exercise sessions in close proximity - as a first exercise session is necessary to reduce the muscle glycogen stores. In the present study, we manipulated the recovery duration between a first muscle glycogen-depleting exercise and a second exercise session, such that the second exercise session started with reduced muscle glycogen in both approaches but was performed either 2 or 15 h after the first exercise session (so-called "twice-a-day" and "once-daily" approaches, respectively). We found that exercise twice-a-day increased the nuclear abundance of transcription factor EB (TFEB) and nuclear factor of activated T cells (NFAT) and potentiated the transcription of peroxisome proliferator-activated receptor-□ coactivator 1 alpha (PGC-1α), peroxisome proliferator-activated receptor alpha (PPARα) and peroxisome proliferator-activated receptor beta/delta (PPARβ/δ) genes, in comparison with the once-daily exercise. These results suggest that part of the elevated molecular signalling reported with previous "train-low" approaches might be attributed to performing two exercise sessions in close proximity. The twice-a-day approach might be an effective strategy to induce adaptations related to mitochondrial biogenesis and fat oxidation.

Keywords: muscle glycogen; high-intensity exercise; mitochondrial biogenesis; molecular signalling; transcription factor

INTRODUCTION

Endurance exercise is a powerful stimulus affecting cytoplasmic and nuclear proteins, and genes encoding mitochondrial proteins, with a subsequent increase in mitochondrial biogenesis (i.e., the generation of new mitochondrial components leading to increased mitochondrial content and respiratory function) (1-8). While these responses are affected by the nature of the exercise (e.g., the exercise intensity (2, 9)), there is evidence substrate availability is also a potent modulator of this response (10-13). It has been hypothesized that initiating endurance exercise with low muscle glycogen stores (the so-called "train-low" approach) results in a greater increase in the content of nuclear proteins (14, 15) and in the transcription of genes (16-19) associated with mitochondrial biogenesis. However, although the "train-low" strategy has been reported to potentiate skeletal muscle signalling responses related to mitochondrial biogenesis (16, 20-22), there are also contrasting findings showing no effects (23, 24) and a consensus is yet to be reached.

Some of the evidence supporting the "train-low" approach is based on performing a first exercise session to reduce muscle glycogen stores, which is followed by a second exercise session 1 to 3 h later – the so-called "twice-a-day" approach (20-23, 25-27). Although the second exercise session will start with reduced muscle glycogen stores, it is difficult to determine if any changes are due to performing the second exercise session with low muscle glycogen or performing the second exercise session soon after the first. The transcriptional responses of many genes associated with mitochondrial biogenesis peak approximately 3 h post exercise and return to basal levels within 8 to 12 h (5, 13, 28-31), including the transcription of peroxisome proliferator-activated receptor-□ coactivator 1 alpha (PGC-1α) - considered a key

regulator of mitochondrial biogenesis (32). Thus, it is possible that reported increases in

gene expression with the twice-a-day approach can be attributed to performing the

second exercise session close to the first, when there is an already increased expression

of genes associated with mitochondrial biogenesis.

The aim of this study was to investigate whether greater exercise-induced

signalling with the "train-low" approach can be attributed to the cumulative effects of

performing two exercise sessions in close proximity. In the present study, we

manipulated the recovery duration between a first muscle glycogen-depleting exercise

and a second exercise session (i.e., a "once-daily" *vs.* a "twice-a-day" approaches). In

the once-daily condition, muscle glycogen content was reduced via evening exercise

(prolonged exercise) followed by a carbohydrate-restricted period, and a second

exercise (i.e., high-intensity interval exercise) was performed the next day (i.e., 15 h

between exercise sessions). In the twice-a-day condition, the same exercises were used

but with a short recovery period between exercise sessions (i.e., 2 h between exercise

sessions). A high-intensity interval exercise session undertaken without a prior muscle

glycogen-depleting exercise served as a control. We hypothesized that the "twice-a-

day" approach would induce a more elevated molecular signalling response during the

second exercise session, compared to the "once-daily" approach.

MATERIALS AND METHODS

Participants

Eight healthy men [age: 30.8 ± 3.5 years, body mass: 78.7 ± 9.9 kg, height: 1.76

± 0.07 m, body fat: $13.6 \pm 5.1\%$, peak oxygen uptake ($\dot{V}O_{2peak}$): 37.1 ± 6.4 mL·kg^{-1}·min^{-}

^{1}, peak power derived from a graded exercise test: 229.6 ± 38.2 W, and first and second

lactate thresholds: 78.1 ± 20.9 and 170.0 ± 37.7 W, respectively] participated in this

study. Participants were recruited by word-of-mouth conversations, online postings, and

flyers posted at the University Campus. Participants were considered eligible if were

physically active (more than 3 times/week) and accustomed to cycling. Participants

were excluded if they had history or signs of cardiac, metabolic, or respiratory diseases

and/or had been using any prescription medications. A medical doctor administered a

medical history screening tool for this purpose. Participants were informed about the

procedures, risks, and benefits associated with the protocol, before they signed a

consent form agreeing to participate in this study, which was approved by the Research

Ethics Committee of Federal University of Pernambuco (approval number:

30378414.8.0000.5208). The study was conducted according to the principles presented

in the Declaration of Helsinki.

Sample size was calculated using a G*Power software

(Heinrich-Heine-University Düsseldorf, version 3.1.9.2, Düsseldorf, Germany),

assuming an effect of exercising with reduced muscle glycogen content on PGC-1α

mRNA expression of 1.34 (16), an alpha of 0.05, and a desired power of at least 0.80.

We used PGC-1α mRNA expression as the main primary outcome as this gene has been

considered a key regulator of mitochondrial biogenesis (32). The effective sample size

necessary to achieve statistical significance was 6 participants; however, sample size

was increased to 8 participants to account for potential dropouts or insufficient muscle

sample size due to technical problems.

Study overview

Each participant completed three experimental trials in a crossover, randomized,

and incomplete balanced Latin Square counterbalanced measure design. The

randomization was performed using software available at http://www.randomized.org

and controlled by an investigator of our laboratory who was not conscious of the aim of

the study. All tests were performed at Vitoria Santo Antao Campus of Federal

University of Pernambuco from March to November 2015.

An overview of the experimental design is shown in Fig. 1. Briefly, in the once-

daily approach participants performed the muscle glycogen-depleting exercise in the

evening (2000 – 2200 h) followed by an overnight fast. The next morning (0800 h),

participants ate a low-carbohydrate breakfast (carbohydrate: 42.7 ± 5.0 kcal, 7%; fat:

365.9 ± 43.1 kcal, 60%; protein: 201.3 ± 23.7 kcal, 33%) and then performed the high-

intensity interval exercise session in the afternoon (1300 h). In the twice-a-day

approach, participants ate a low-carbohydrate breakfast on the morning of the

experimental day (0800 h), and then performed a muscle glycogen-depleting exercise

(0900-1100 h), followed by a 2-h rest period and the high-intensity interval exercise

(1300 h). In the control condition, participants consumed the same low-carbohydrate

breakfast as the two other experimental trials (0800 h), and then performed the same

high-intensity interval exercise session in the afternoon (i.e., 1300 h). Skeletal muscle

biopsies from the *vastus lateralis* and venous blood samples were taken before,

immediately after, and 3 h after completion of the high-intensity interval exercise

sessions. Participants were not allowed to eat any food between the breakfast and the

last muscle biopsy in all experimental trials. Water was provided ad libitum throughout

each trial. Each experimental trial was separated by approximately two weeks to

washout any residual effect of fatigue or damage caused by exercise and the multiple

muscle biopsies. The main primary outcome measure was PGC-1α mRNA, while the

second outcome measures were nuclear/cytosolic proteins and other genes associated

with mitochondrial biogenesis.

176   (PLEASE INSERT FIGURE 1 HERE)

Exercise protocols

Preliminary test

One week prior to the commencement of this study, participants performed a

graded exercise test to volitional fatigue on a cycloergometer (Ergo-Fit 167, Pirmasens,

Germany). The test commenced at 50 W, and thereafter intensity was increased by 25

W every 4 min, with a 1-min break between stages, until volitional exhaustion (33). The

test was interrupted when the participant could no longer maintain the required cadence

(70 rpm). Strong verbal encouragement was provided to each participant.

$\dot{V}O_2$ was measured breath-by-breath throughout the test using an automatic

analyzer (Cortex, Metalyzer 3B$^®$, Saxony, Germany). Before each test, the gas analyzer

was calibrated using ambient air and a cylinder of known gas concentration (12% O_2

and 5% CO_2). The volume was calibrated using a 3-L syringe (Quinton Instruments,

Washington, US). Capillary ear lobe blood samples were taken at rest and immediately

after each 4-min stage of the test for determination of plasma lactate concentration. The

first lactate threshold was visually identified by two experienced investigators as the

first increase in plasma lactate concentration above resting level. The second lactate

threshold was calculated by the modified Dmax method (34). This was determined by

the point on the polynomial regression curve that yields the maximal perpendicular

distance to the straight line connecting the first lactate threshold and the final stage of

the test. $\dot{V}O_{2peak}$ was defined as the highest 30-s average $\dot{V}O_2$ during the test and peak

power was determined as the highest workload reached. If a participant did not

complete the final 4-min stage, then the peak power was determined using the following

equation:

Peak power = P_{last} + (t/240 · 25)   (1)

where P_{last} is the power in Watts of the last completed stage performed by the

participant, t is the time (in seconds) sustained during the last incomplete stage, and 25

corresponds to the increments in power (Watts) at each stage.

Muscle glycogen-depleting exercise

To reduce muscle glycogen stores, participants cycled for 100 min at a power

corresponding to 50% of the difference between their first and second lactate threshold

(124 ± 27 W, 54 ± 5% of peak power). Then, after an 8-min rest, participants performed

six 1-min exercise bouts at 125% of their peak power (287 ± 46 W) interspersed with 1-

min rest periods (35). This protocol has been shown to be effective to reduce muscle

glycogen content (36, 37).

High-intensity interval exercise

The high-intensity interval exercise sessions were preceded by a 5-min warm-up

at 90% of the first lactate threshold. Participants completed ten 2-min intervals at an

intensity of 20% of the difference between the second lactate threshold and their peak

power (182 ± 38 W, 79 ± 5% of peak power). Each 2-min bout was interspersed with a

1-min passive recovery period (33). Participants were required to maintain a pedal

frequency of 70-80 rpm during each 2-min bout. The $\dot{V}O_2$, $\dot{V}CO_2$, RER, and $\dot{V}E$ were

measured breath-by-breath throughout each high-intensity interval exercise session

using the same gas analyzer described for the graded exercise test. Data were then

converted to 30-s intervals and the last 30-s of each 2-min bout were used for further

analysis.

Diet and exercise control before starting experimental manipulation

Participants were asked to register all foods and beverages consumed during the

48-h preceding the start of the preliminary test. The usual macronutrient consumption

was measured from this diet recall using a computer program designed to perform

nutritional calculations (Nutwin, Nutrition Support Program, version 1.5, Department of

Health Informatics, Federal University of São Paulo, Brazil). The daily energy and

macronutrient intakes during the 48-h preceding the first preliminary test were 3330 $\pm$

475 kcal (43.2 $\pm$ 10.0 kcal·kg^{-1}), 49% $\pm$ 6% carbohydrates (409.3 $\pm$ 79.4 g, 5.3 $\pm$ 1.4

g·kg^{-1}), 27% $\pm$ 5% lipids (101.5 $\pm$ 28.2 g, 1.3 $\pm$ 0.5 g·kg^{-1}), and 24% $\pm$ 7% protein

(195.0 $\pm$ 43.8 g, 2.5 $\pm$ 0.5 g·kg^{-1}). Participants were asked to replicate the dinner of two

days before (day 1), and breakfast, lunch and dinner of one day before the experimental

trial (day 2) (Fig. 1). Participants were given verbal and written instructions describing

how to repeat this before each subsequent experimental trial. Checklists were used to

check any deviations from the menu, and analysis of checklists showed no differences

for energy or macronutrient intake between conditions during these meals (Table 1).

Participants were also instructed to avoid any strenuous exercise as well as alcohol and

caffeine consumption for the 24 h prior to each experimental trial.

(PLEASE INSERT TABLE 1 HERE)

Blood collection and analysis

Blood samples were collected from an antecubital vein and separated into three

different tubes. Two millilitres of blood were collected in tubes containing sodium

fluoride and EDTA (Hemogard^{TM} Fluoride/EDTA, BD Vacutainer®, USA). Blood was

centrifuged at 4,000 rev.min^{-1} for 10 min at 4º C with the resulting plasma transferred to

2-mL tubes and immediately analysed for plasma glucose and lactate concentrations.

Plasma glucose and lactate concentrations were analysed with a commercially available

enzymatic kit (Glucose Liquiform and Enzymatic Lactate, respectively, Labtest, Lagoa

Santa, Minas Gerais, Brazil). A further 8 mL of blood was collected in tubes containing

Clot activator and gel for serum separation (SST II Plus, BD Vacutainer®, USA), and

another 3 mL of blood was collected in tubes containing sodium heparin (Sodium

Heparin^{N} Plus, BD Vacutainer®, USA). Both were then centrifuged at 4,000 rev.min^{-1}

for 10 min at 4º C, and the resulting serum/plasma frozen and stored in liquid nitrogen

for the later analysis of free fatty acid, glycerol, and catecholamine concentrations in the

serum/plasma. Serum free fatty acid and glycerol concentrations were determined by an

enzymatic colorimetric method (EFFA-100 and EGLY-200, BioAssay, Hayward,

California, USA). Plasma catecholamine concentrations were determined by using ion-

pairing reverse phase liquid chromatography coupled with electrochemical detection

(35).

Muscle tissue samples and analysis

Nine separate incisions (three per trial) were made into the vastus lateralis under

local anaesthesia (2% Xylestesin®), and a muscle sample taken using a Bergström

needle (38) adapted for manual suction (39). Samples were taken approximately 1 cm

apart from a previous biopsy site. Samples [mean: 118 ± 48 mg; range: 49 to 248 mg]

were immediately snap-frozen in liquid nitrogen, and then stored at -80ºC until

subsequent analyses. Muscle samples were taken at rest (pre), immediately after (post)

and 3 h after the high-intensity interval exercise (3 h post). Biopsies were subsequently

analysed for muscle glycogen content, as well as gene and protein expression (described

subsequently).

Muscle glycogen concentration

Approximately 2 to 3 mg of freeze-dried muscle tissue was powdered and

dissected free of all visible non-muscle tissue. Powdered muscle tissue was then

extracted with 250 µL of 2 M HCl, incubated at 95°C for 2 h (agitated gently every 20

min), and then neutralized with 750 µL of 0.66 M NaOH. Glycogen concentration was

subsequently assayed in triplicate via enzymatic analysis with fluorometric detection

(40) and the mean value reported as millimoles per kilogram dry weight.

Western blotting

Muscle homogenate preparations and protein assays

Approximately 20 mg of frozen muscle tissue was homogenized using a

TissueLyser II (Qiagen, Valencia, CA) in a 1:20 dilution of ice-cold RIPA buffer (pH

7.4) containing: 0.15 M NaCl, 1% Triton-X100, 0.5% sodium deoxycholate, 0.05 M

Tris, 0.1% SDS, 0.1 M EDTA, with the addition of protease/phosphatase inhibitor

cocktail (Cell Signalling Technology [CST], #5872, St. Louis, MI). Homogenates were

rotated end-over-end for 60 min at 4°C. Protein content of muscle homogenate was

measured in triplicate using a Bradford assay (Bio-Rad protein assay dye reagent

concentrate, Bio-Rad Laboratories, Hercules, CA) against bovine serum albumin

standards (BSA, A9647, Sigma-Aldrich). Nuclear and crude cytosolic fractions were

prepared from 40 to 60 mg of wet muscle using a commercially-available nuclear

extraction kit (NE-PER^{TM}, Pierce, USA). Muscle samples were homogenized in

cytoplasmic extraction reagent I buffer containing a protease/phosphatase inhibitor

cocktail (CST, 5872). Following centrifugation (16,000 g for 5 min at 4º C), the

supernatant was taken and pellets containing nuclei were washed five times in PBS to

remove cytosolic contamination, before nuclear proteins were extracted by

centrifugation (16,000 g for 10 min at 4º C) in high-salt nuclear extraction reagent

buffer supplemented with the same inhibitors cocktail following manufacturer's

instruction. Verification of subcellular enrichment is presented in Fig. 3 b and 4 b.

Sufficient muscle was available to prepare subcellular fractions from all eight

participants.

Immunoblotting

RIPA-buffered homogenate was diluted in 4X Laemmli buffer (0. 25 M Tris, 8%

SDS, 40% glycerol, 0.04% bromophenol blue, 20% 2-mercaptoethanol) and equal

amounts of total protein (10 to 20 µg) were loaded on Criterion™ 4-20% TGX Stain-

Free™ Precast Gels (Bio-Rad). All samples for a participant were loaded in adjacent

lanes on the same gel. Four to six different dilutions of a mixed-homogenate internal

standard were also loaded on each gel and a calibration curve plotted of density against

protein content. From the subsequent linear regression equation protein abundance was

calculated from the measured band intensity for each sample on the gel (41). Gel

electrophoresis ran for 90 min at 80-150 V. Proteins were turbo-transferred to a 0.2 µm

PVDF membrane at 25 V for 10 min. Membranes were blocked for 60 min at room

temperature in 5% non-fat dry milk diluted in Tris-buffered saline with 0.1% Tween-20

(TBST). Membranes were then washed in TBST and incubated overnight at 4°C – with

the appropriate primary antibody: Histone H3 (CST, 44995), LDHA (CST, 2012), PGC-

1α (CST, 2178), p-ACC^{Ser79} (CST, 3361), AMPK (CST, 2532), p-AMPK^{Thr172} (CST,

2531), p38 MAPK (CST, 9212), p-p38 MAPK$^{Thr180/Tyr182}$ (CST, 9211), PHF20 (CST,

3934), p53 (CST, 2527), TFEB (CST, 9601), and NFAT2 (CST, 8032) diluted (1:1,000)

in 5% BSA and 0.02% sodium azide in TBST. Following TBST washes the membranes

were incubated in the relevant secondary antibody: Goat anti-rabbit IgG (Perkin

Elmer/NEF812001EA), diluted (1:10,000) in 5% non-fat dry milk in TBST, for 60 min

at room temperature. After further washes, membranes were developed using Clarity

ECL (Bio-Rad) and images were taken with a ChemiDoc Imaging System fitted (Bio-

Rad). Densitometry was performed with Image Lab 5.0 software (Bio-Rad). Images are

typically displayed with at least five bandwidths above and below the band of interest.

Real-Time quantitative PCR

RNA extraction

Total RNA from approximately 10 to 15 mg of frozen muscle was homogenized

in 800 µL of TRIzol reagent (Thermo Fisher Scientific, Waltham, USA) using a

TissueLyser II (Qiagen) (42). The concentration and purity of each sample was assessed

using a NanoDrop One/One^c (Thermo Fisher Scientific). As representative, RNA

integrity of a subset of samples was measured using a Bio-Rad Experion microfluidic

gel electrophoresis system with Experion RNA StdSens Analysis kit (Bio-Rad,

7007104). All tested samples showed good quality (RNA quality indicator > 7). RNA

was stored at −80°C until reverse-transcription was performed.

Reverse transcription

1 µg RNA, in a total reaction volume of 20 µL, was reverse-transcribed to

cDNA using a Thermocycler (Bio-Rad) and iScript RT Supermix (Bio-Rad, 170-8840)

per the manufacturer's instructions. Priming was performed at 25°C for 5 min and

reverse transcription for 30 min at 42°C. All samples, including RT-negative controls,

were performed during the same run. cDNA was stored at −20°C until subsequent

analysis.

qPCR

Relative mRNA expression was measured by qPCR (QuantStudio 7 Flex,

Applied Biosystems, Foster City, CA) using SsoAdvanced Universal SYBR Green

Supermix (Bio-Rad). Primers were designed using Primer-BLAST (43) to include all

splice variants, and were purchased from Sigma-Aldrich (see Supplementary Table 1 for

primer details). All reactions were performed in duplicate on 384-well MicroAmp

optical plates (4309849, Applied Biosystems) using an epMotion M5073 automated

pipetting system (Eppendorf AG). Total reaction volume of 5 µL contained 2 µL of

diluted cDNA template, 2.5 µL of mastermix, and 0.3 µM or 0.9 µM primers. All assays

ran for 10 min at 95°C, followed by 40 cycles of 15 s at 95°C and 60 s at 60°C. The

stability of six potential reference genes were determined by *BestKeeper* (44) and

NormFinder (45) software, and the three most stably expressed genes were TBP, 18S,

and ACTB (Supplementary Table 1). There were no main effects of training approach,

time, nor interaction effects, for the quantification cycle (Ct) values of the three most

stable housekeeping genes ($P > 0.05$). Expression of each target gene was normalized to

the geometric mean of expression of the three reference genes (46), and using the $2^{-\Delta\Delta Ct}$

method (47).

Statistical analysis

Statistical analysis was performed using the GraphPad Prism software version 6.01. All data were checked for normality with the Kolomogorov-Smirnov test. To compare the responses before, immediately after, and 3 h after each high-intensity interval exercise session, data were analysed with two-way repeated measures ANOVA (trial *vs*. time). A Bonferroni post-hoc test was used to locate the differences. All values are expressed as means ± SD. Significance was accepted when p < 0.05.

RESULTS

Muscle glycogen concentration. Prior to the high-intensity interval exercise, muscle glycogen concentration was similarly lower in both the twice-a-day and once-daily conditions, compared to the control condition (~45 and 42%, respectively; Fig. 2). Compared to their respective rest values, muscle glycogen concentration decreased similarly after the high-intensity interval exercise in all three conditions (from 245.0 to 135.7 mmol·kg^{-1} dry mass in the twice-a-day, from 262.2 to 163.7 mmol·kg^{-1} dry mass in the once-daily, and from 449.2 to 312.2 mmol·kg^{-1} dry mass in the control; ~45, 38, and 30% of reduction for twice-a-day, once-daily and control, respectively; *P* < 0.05; Fig. 2), and remained lower 3 h post exercise in both the twice-a-day and once-daily conditions, compared to the control condition (~46 and 43%, respectively; Fig. 2).

(PLEASE INSERT FIGURE 2 HERE)

Cytosolic proteins relative abundance. Representative blots are presented in Fig. 3 a,b. Cytosolic p53 protein relative abundance increased immediately post high-intensity

397   interval exercise in all three conditions (Fig. 3 d), with no differences between

398   conditions. Cytosolic PGC-1α, phosphorylated p53 (p-p53^{Ser15}), PHF20 protein

399   (PHF20), TFEB, p38 mitogen-activated protein kinase (p38 MAPK), phosphorylated 5'

400   adenosine monophosphate-activated protein kinase (p-AMPK^{Thr172}) and NFAT relative

401   abundance were unaffected by the exercise approach or time (Fig. 3 c, 3 e-j).

403   (PLEASE INSERT FIGURE 3 HERE)

Nuclear protein relative abundance. Representative blots are presented in Fig. 4 a,b.

Nuclear PGC-1α, p53, and p-p53^{Ser15} relative abundance increased post high-intensity

interval exercise (Fig. 4 c-e), with no clear differences between the three conditions.

The relative abundance of nuclear PHF20, p38MAPK, and p-AMPK^{Thr172} was

unaffected by either "train-low" approach or time (Fig. 4 f-h). However, nuclear TFEB

relative abundance was greater in the twice-a-day compared to the once-daily condition

both pre and post the high-intensity interval exercise (Fig. 4 i). Moreover, nuclear

NFAT relative abundance was also greater in the twice-a-day compared to both the

once-daily and control condition post the high-intensity interval exercise (Fig. 4 j).

415   (PLEASE INSERT FIGURE 4 HERE)

Mitochondrial-related gene expression. Pre high-intensity interval exercise, total

PGC-1α mRNA content was ~9-fold higher in the twice-a-day compared to both the

once-daily and control conditions (Fig. 5 a). Three hours post the high-intensity interval

exercise, total PGC-1α mRNA content increased in all three conditions compared with

their respective pre-values; however, total PGC-1α mRNA content remained ~10-fold

higher in the twice-a-day compared to both control and once-daily conditions. Similarly,

PGC-1α isoform 4 mRNA content was ~24-fold higher at pre high-intensity interval

exercise in the twice-a-day compared to both the once-daily and control conditions (Fig.

5 c). At 3 h post high-intensity interval exercise, the PGC-1α isoform 4 mRNA content

increased in all three conditions compared with their respective pre-values; however, the

PGC-1α isoform 4 mRNA content remained ~10-fold higher in the twice-a-day

compared to the control and once-daily conditions. Additionally, PGC-1α isoform 1

mRNA content was higher in the twice-a-day compared to the once-daily when all time

points where considered (Fig. 5 b). There was no effect of condition for p53, TFEB,

chromodomain-helicase-DNA-binding protein 4 (CHCHD4), p21, mitochondrial

transcription factor A (Tfam), NADH dehydrogenase subunit β (NDUFβ; mitochondrial

complex I), succinate dehydrogenase subunit β (SDHβ; mitochondrial complex II),

cytochrome c (mitochondrial complex III), and cytochrome c oxidase subunit IV (COX

IV; mitochondrial complex IV) mRNA content post high-intensity interval exercise

(Fig. 5 d-l). However, the mRNA content of representative subunits of mitochondrial

complexes II, III, and IV increased 3 h post high-intensity interval exercise to a similar

extent for all three conditions.

440   (PLEASE INSERT FIGURE 5 HERE)

Fat transport and lipolysis related genes. The content of carnitine

palmitoyltransferase I subunit A (CPT1A) mRNA was higher in both the twice-a-day

and once-daily conditions compared with the control condition at 3 h post the high-

intensity interval exercise; however, there was no difference between the two "train-

low" approaches (Fig. 6 a). There was an increase in mitochondrial uncoupling protein

3 (UCP3) mRNA content 3 h post exercise in all three conditions. The UCP3 mRNA

content was, however, significantly higher only in the twice-a-day compared to the

control condition at 3 h post the high-intensity interval exercise (Fig. 6 b). Pre high-

intensity interval exercise, the PPARα mRNA content was ~11-fold higher in the twice-

a-day compared to both the once-daily and control conditions (Fig. 6 c). Three hours

post the high-intensity interval exercise, the PPARα mRNA content increased ~7- and

9-fold in the once-daily and control conditions, respectively, compared with their

respective pre-values; however, the PPARα mRNA content remained ~16-fold higher in

the twice-a-day compared to the once-daily and control conditions. The PPARβ/δ

mRNA content was higher in the twice-a-day than in the control condition post the

high-intensity interval exercise, and higher than both the once-daily and control

conditions at 3 h post the high-intensity interval exercise (Fig. 6 d). The PPARβ/δ

mRNA content was higher at 3 h post the high-intensity interval exercise compared to

pre- and post- only for the twice-a-day approach. The citrate synthase (CS) mRNA

content was higher 3 h post the high-intensity interval exercise compared to post for all

three conditions. However, the peroxisome proliferator-activated receptor gamma

(PPARγ), β-hydroxyacyl-CoA dehydrogenase (β-HAD), and fatty acid translocase

cluster of differentiation 36 (CD-36) mRNA content were unaffected by the "train-low"

approach or time (Fig. 6 e-h).

467   (PLEASE INSERT FIGURE 6 HERE)

Glycolysis related genes. The mRNA content of phosphofructokinase (PFK) and

glucose transporter 4 (GLUT4) was unaffected by the exercise approach or time. While

pyruvate dehydrogenase kinase isoenzyme 4 (PDK4) mRNA content increased 3 h post

the high-intensity interval exercise, there was no difference for the three different

exercise approaches (Fig. 7 a-c).

(PLEASE INSERT FIGURE 7 HERE)

Physiological responses. The twice-a-day approach was associated with a higher heart

rate, ventilation, and oxygen uptake, and a lower plasma glucose concentration, during

high-intensity interval exercise than both the once-daily and the control condition (Fig.

8 a-c and Table 2). Plasma glucose was also lower during the high-intensity interval

exercise in the once-daily compared to the control condition (Table 2). In addition, the

respiratory exchange ratio was lower and the serum free fatty acid concentration higher

than the control only for the twice-a-day approach (Fig. 8 d and Table 2). Serum

glycerol was higher and plasma lactate lower post the high-intensity interval exercise in

both the twice-a-day and once-daily conditions compared with the control condition

(Table 2). Plasma epinephrine and norepinephrine concentrations were not influenced

by the exercise approach undertaken.

(PLEASE INSERT FIGURE 8 HERE)

(PLEASE INSERT TABLE 2 HERE)

DISCUSSION

There is continued debate about whether beginning exercise with low muscle

glycogen stores potentiates the exercise-induced increase in genes associated with

mitochondrial biogenesis and metabolism (10, 11, 48-52). Some of this controversy

may relate to the observation that some of the evidence supporting the "train-low"

approach is based on performing the experimental exercise session a few hours after a

glycogen-lowering exercise session (20-23, 25-27). Thus, it is difficult to determine if

any observed effects are due to performing the second exercise session with low muscle

glycogen stores and/or performing the second exercise session close to the first. We

investigated this question by performing the same exercise session with similar starting

muscle glycogen stores, but either 2 or 15 h following the previous glycogen-lowering

exercise session. Our results indicate that the greater exercise-induced nuclear protein

abundance (TFEB and NFAT) and transcription of genes involved in mitochondrial

biogenesis (PGC-1α, PPARα, PPARβ/δ) with the so-called "train-low" approach might

be attributed to performing two exercise sessions in close proximity.

Despite the different recovery periods between exercise sessions for the two

"train-low" approaches, muscle glycogen prior to the high-intensity interval exercise

was reduced to a similar extent in both the twice-a-day and once-daily condition

compared with the control condition (Fig. 2). This level of muscle glycogen

concentration is consistent with previous studies that have utilized either the twice-a-

day (21, 22) or once-daily (19) approach. In addition, muscle glycogen concentration

before commencing the high-intensity exercise was below 300 mmol·kg^{-1}·dry mass for

both the twice-a-day and once-daily conditions. Post high-intensity interval exercise

muscle glycogen concentration remained above 100 mmol·kg^{-1}·dry mass. These values

have been suggested as an "upper" and "lower" limit threshold, respectively, for which

muscle glycogen levels may modulate genes related to mitochondrial biogenesis (52).

Thus, the similar low muscle glycogen levels when commencing the high-intensity

interval exercise in both "train-low" conditions allowed us to determine whether any

differences for exercise-induced gene or protein expression could be attributed to

performing high-intensity interval exercise with reduced muscle glycogen stores or to

performing high-intensity interval exercise close to a previous exercise session.

In the present study, both p-AMPK and p-p38 MAPK relative protein abundance

in both the cytosolic and nucleus fractions was not altered with exercise or influenced

by either "train-low" approach (Fig. 4 g,h). It has been suggested that the greater

exercise-induced upregulation of cell-signalling pathways with reduced muscle

glycogen content may be associated with enhanced nuclear and cytosolic abundance and

greater activation of p-AMPK and p-p38 MAPK (14, 15, 53), but contradictory results

have been reported (19, 21, 25, 54). For example, Cochran et al. (25) observed larger

increases in cytosolic p-p38 MAPK but not p-AMPK with the twice-a-day approach,

while Yeo et al. (21) reported larger increases in cytosolic p-AMPK but without

alterations in p-p38 MAPK with the twice-a-day compared to the once-daily approach.

On the other hand, Gejl et al. (54) and Psilander et al. (19) found no effect of either the

twice-a-day or once-daily approach on both p-AMPK and p-p38 MAPK protein content

in cytosolic fractions. In addition, Chan et al. (14) and Steinberg et al. (15) reported an

increase in nuclear abundance of p-p38 MAPK and AMPK, respectively, with the once-

daily approach. The explanation for these contradictory results is unclear, but it is worth

noting both the glycogen-depleting exercise and the "depleted-exercise" varied

considerably in these studies; either high- (25, 54) or low-intensity (14, 15, 19, 21)

exercise was used to reduce muscle glycogen, and high- (21, 25) or low-intensity (14,

15, 19, 54) exercise was implemented during the second exercise session (depleted-

exercise). However, activation of AMPK and p38 MAPK does not seem to be

associated with muscle glycogen levels or a particular train-low regime (55). Thus, it

remains to be solved which protocol or strategy might induce augmented activation of

p-AMPK and p-p38 MAPK.

Although p-AMPK and p-p38 MAPK relative protein abundance in the

cytosolic and nucleus were not altered with exercise or "train-low" approaches, there

was greater PGC-1α, p-p53, and p53 relative protein abundance in the nucleus

immediately post the high-intensity interval exercise (Fig. 4 c-e). There was also a

higher p53 cytosolic protein abundance post the high-intensity interval exercise (Fig 3

d). The increased nuclear and cytosolic p53 abundance are consistent with the well-

accepted notion that cellular stress is associated with accumulation of p53 protein - a

process that has been partly attributed to increased p53 protein stability (56, 57).

Nevertheless, we found no significant differences between either of the "train-low"

approaches and the control condition for p-p53 and p53 relative protein abundance in

either subcellular fraction. The "train-low" approach has been associated with a greater

exercise-induced cytosolic relative protein abundance of p-p53 (16), which is suggested

to influence the gene expression of PGC-1α (58). As differences in neither p-p53 nor

p53 protein content in the nucleus could explain the greater exercise-induced increases

in PGC-1α gene expression with the different train-low approaches in the present study,

we subsequently investigated the nuclear abundance of other proteins that may

contribute to the exercise-induced regulation of this gene.

The nuclear abundance of NFAT increased significantly after performing high-

intensity interval exercise only in the twice-a-day condition, and post-exercise values

were higher than both the control and once-daily conditions (Fig. 4 j). In addition, in

both the pre- and post-exercise muscle samples the nuclear abundance of TFEB was

significantly greater with the twice-a-day compared to the once-daily approach (Fig. 4

i). As activated calcineurin dephosphorylates both NFAT (59, 60) and TFEB (61-63),

leading to their translocation to the nucleus (59-64), this suggests there might be greater

calcineurin activation when high-intensity interval exercise is performed soon after a

prior exercise session. Although we did not have sufficient muscle sample to test this

hypothesis, in the twice-a-day condition we observed significantly greater exercise-

induced increases of genes that have been reported to be regulated by calcineurin (e.g.,

PGC-1α, PPARα and PPARβ/δ; (64, 65)).

In contrast to our findings, some studies have reported greater exercise-induced

increases in PGC-1α mRNA with the once-daily approach (~6-fold from rest) when

compared to a control condition (~3-fold from rest) (16, 18, 19). This previously

reported ~6-fold increase in PGC-1α mRNA with the once-daily approach is of similar

magnitude to the PGC-1α mRNA increase with the once-daily approach in the present

study (~6 fold from rest), although this was not significantly different to the change we

observed in the control condition (~6-fold from rest). However, we did observe a

significantly greater exercise-induced increase in PGC-1α (and PPARα, PPARβ/δ)

mRNA content in the twice-a-day condition, even though muscle glycogen content

before commencing the high-intensity interval exercise was similar between twice-a-

day and once-daily conditions, and both were lower than control. Our findings therefore

suggest that the greater exercise-induced increase in these genes in our study can be

attributed to performing the high-intensity interval exercise soon after the prior exercise

session, rather than beginning the high-intensity interval exercise with lowered muscle

glycogen levels. It should be mentioned, however, that carbohydrate might still play a

role in myocellular signalling as ingesting exogenous carbohydrate between the two

exercise sessions has been reported to block the stimulus for inducing oxidative enzyme

adaptations in skeletal muscle (25, 27).

In the twice-a-day condition we also observed a greater circulating free fatty

acid concentration pre, post, and 3 h post the high-intensity interval exercise (Table 2).

Although respiratory exchange ratio values were always below 1.0 after the third

interval, and systematically lower in the twice-a-day compared to the control condition

(Fig. 8), which could be indicative of greater fat oxidation, this conclusion is limited by

the use of non‑protein respiratory quotient to calculate substrate oxidation during non-

steady state exercise (66). The increase in circulating free fatty acids, however, might

activate calcineurin (67) and regulate skeletal muscle metabolism via coordinated

changes in gene expression (64). As previously mentioned, activated calcineurin will

lead to the translocation of both NFAT and TFEB to the nucleus (59-64) and the

subsequent upregulation of genes related to fat metabolism (64). These exercise-induced

changes are consistent with studies that have reported greater increases in fat oxidation

when training twice-a-day (20, 22). Thus, these results reinforce the hypothesis that

twice-a-day training might be more effective to induce adaptions related to fat

metabolism.

Our study has some limitations that should be mentioned. Firstly, as we were

constrained by the number of muscle biopsies we could take, we elected to take our

final muscle biopsy 3 h post exercise as most of the key genes investigated in the

present study have been reported to peak ~3 h after exercise (e.g., PGC-1α and

PPARβ/δ) (5, 68). However, other genes might peak later (e.g. CD-36) (13) and we may

have underestimated the influence of the two "train-low" approaches on some genes.

Similarly, we did not perform muscle biopsies immediately and 3 h after the muscle

glycogen-depleting exercise; therefore, relevant changes in molecular signalling

following the first exercise session in the once-daily condition may have been missed.

Secondly, consistent with most related research (16, 18, 23, 25-27), we recruited

participants who were active (~3.3 h of aerobic training per week) but who were not

well-trained athletes. While this may limit the applicability of our findings to elite

athletes, it is difficult to perform similar experiments requiring multiple muscle biopsies

in well-trained athletes. Thus, further studies comparing different "train-low"

approached in well-trained athletes are required.

In summary, our findings indicate that the greater exercise-induced signalling

with the so-called "train-low" approach in our study can partially be attributed to the

performance of two exercise sessions in close succession rather than exercising with a

reduced muscle glycogen content. We presented evidence that performing two exercise

sessions separated by a short recovery period increases the nuclear abundance of TFEB

and NFAT and potentiates the transcription of PGC-1α, PPARα and PPARβ/δ.

Although we have proposed some molecular mechanisms by which the twice-a-day

approach might be a more effective strategy to induce adaptations related to

mitochondrial biogenesis and fat oxidation, further research is required to determine if

training using the twice-a-day approach results in greater changes in mitochondrial

content, mitochondrial function, and fat oxidation.

Acknowledgements

This work was supported by the Brazilian National Council for Scientific and

Technological Development (Special visiting researcher of science without borders

program, process Nº 406201/2013-7) and an Australian Research Council Grant

(DP140104165) to D. J. Bishop. V.A. Andrade-Souza is grateful to the Foundation for

the Support of Science and Technology of the State of Pernambuco (FACEPE) for his

scholarship. We thank G. McConnel, X. Yan, C. Granata, A.J. Genders, E.C. Marin, R.

Aragão and C.R. Correia-Oliveira for technical assistance.

Author contributions

V. A. A-S., R. B., D. J. B., and A. E. L-S. designed the experiments. V. A. A-S., T. G.,

A. S., K. A. S. S., F. T., L. A., and A. E. L-S. performed the experiments. V. A. A-S., T.

G., A. S., K. A. S. S., F. T., L. A., and A. E. L-S. conducted the exercise analyses at

Department of Physical Education and Sports Science, Academic Center of Vitoria,

Federal University of Pernambuco and V. A. A-S., J. F., E. P., N. S., and J. K.

conducted qPCR and western blot analyses at Institute for Health and Sport, Victoria

University. V. A. A-S analyzed data and discussed analyses and results with J. F., J. K.,

D. J. B., and A. E. L-S. T.G., K. A. S. S., R. B., and C. G. L. supported data analyses.

V. A. A-S., K. A. S. S., J. K., D. J. B., and A. E. L-S made the figures. All authors

wrote and approved the manuscript.

References

1. Granata, C., Oliveira, R. S., Little, J. P., Renner, K., and Bishop, D. J. (2016)

Training intensity modulates changes in PGC-1alpha and p53 protein content

and mitochondrial respiration, but not markers of mitochondrial content in

human skeletal muscle. *FASEB J.* **30**, 959-970

2. Granata, C., Oliveira, R. S., Little, J. P., Renner, K., and Bishop, D. J. (2017)

Sprint-interval but not continuous exercise increases PGC-1alpha protein

content and p53 phosphorylation in nuclear fractions of human skeletal muscle.

Sci. Rep. **7**, 44227

3. Holloszy, J. O. (1967) Biochemical adaptations in muscle. Effects of exercise on

mitochondrial oxygen uptake and respiratory enzyme activity in skeletal muscle.

J. Biol. Chem. **242**, 2278-2282

4. Hood, D. A., Tryon, L. D., Carter, H. N., Kim, Y., and Chen, C. C. (2016)

Unravelling the mechanisms regulating muscle mitochondrial biogenesis.

Biochem. J. **473**, 2295-2314

5. Little, J. P., Safdar, A., Bishop, D., Tarnopolsky, M. A., and Gibala, M. J.

(2011) An acute bout of high-intensity interval training increases the nuclear

abundance of PGC-1alpha and activates mitochondrial biogenesis in human

skeletal muscle. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **300**, R1303-1310

6. Granata, C., Jamnick, N. A., and Bishop, D. J. (2018) Principles of Exercise

Prescription, and How They Influence Exercise-Induced Changes of

Transcription Factors and Other Regulators of Mitochondrial Biogenesis. *Sports*

Med. **48**, 1541-1559

7.   Granata, C., Jamnick, N. A., and Bishop, D. J. (2018) Training-Induced Changes
in Mitochondrial Content and Respiratory Function in Human Skeletal Muscle.
Sports Med. **48**, 1809-1828

8.   Granata, C., Oliveira, R. S., Little, J. P., Renner, K., and Bishop, D. J. (2016)
Mitochondrial adaptations to high-volume exercise training are rapidly reversed
after a reduction in training volume in human skeletal muscle. *FASEB J.* **30**,
3413-3423

9.   Islam, H., Edgett, B. A., and Gurd, B. J. (2017) Coordination of mitochondrial
biogenesis by PGC-1alpha in human skeletal muscle: a re-evaluation.
Metabolism. **79**, 42-51

10. Baar, K., and Mcgee, S. (2008) Optimizing training adaptations by manipulating
glycogen. *Eur. J. Sport Sci.* **8**, 97-106

11. Bartlett, J. D., Hawley, J. A., and Morton, J. P. (2015) Carbohydrate availability
and exercise training adaptation: Too much of a good thing? *Eur. J. Sport Sci.*
15, 3-12

12. Knuiman, P., Hopman, M. T. E., and Mensink, M. (2015) Glycogen availability
and skeletal muscle adaptations with endurance and resistance exercise. *Nutr*
Metab **12**

13. Pilegaard, H., Osada, T., Andersen, L. T., Helge, J. W., Saltin, B., and Neufer, P.
D. (2005) Substrate availability and transcriptional regulation of metabolic
genes in human skeletal muscle during recovery from exercise. *Metabolism.* **54**,
1048-1055

14. Chan, M. H., McGee, S. L., Watt, M. J., Hargreaves, M., and Febbraio, M. A.
(2004) Altering dietary nutrient intake that reduces glycogen content leads to
phosphorylation of nuclear p38 MAP kinase in human skeletal muscle:

association with IL-6 gene transcription during contraction. *FASEB J.* **18**, 1785-

1787

15.   Steinberg, G. R., Watt, M. J., McGee, S. L., Chan, S., Hargreaves, M., Febbraio,

M. A., Stapleton, D., and Kemp, B. E. (2006) Reduced glycogen availability is

associated with increased AMPKalpha2 activity, nuclear AMPKalpha2 protein

abundance, and GLUT4 mRNA expression in contracting human skeletal

muscle. *Appl. Physiol. Nutr. Metab.* **31**, 302-312

16.   Bartlett, J. D., Louhelainen, J., Iqbal, Z., Cochran, A. J., Gibala, M. J., Gregson,

W., Close, G. L., Drust, B., and Morton, J. P. (2013) Reduced carbohydrate

availability enhances exercise-induced p53 signaling in human skeletal muscle:

implications for mitochondrial biogenesis. *Am. J. Physiol-Reg. I.* **304**, R450-

R458

17.   Lane, S. C., Camera, D. M., Lassiter, D. G., Areta, J. L., Bird, S. R., Yeo, W. K.,

Jeacocke, N. A., Krook, A., Zierath, J. R., Burke, L. M., and Hawley, J. A.

(2015) Effects of sleeping with reduced carbohydrate availability on acute

training responses. *J. Appl. Physiol.* **119**, 643-655

18.   Pilegaard, H., Keller, C., Steensberg, A., Helge, J. W., Pedersen, B. K., Saltin,

B., and Neufer, P. D. (2002) Influence of pre-exercise muscle glycogen content

on exercise-induced transcriptional regulation of metabolic genes. *J Physiol.*

541, 261-271

19.   Psilander, N., Frank, P., Flockhart, M., and Sahlin, K. (2013) Exercise with low

glycogen increases PGC-1 alpha gene expression in human skeletal muscle. *Eur.*

J. Appl. Physiol. **113**, 951-963

20.   Hulston, C. J., Venables, M. C., Mann, C. H., Martin, C., Philp, A., Baar, K.,

and Jeukendrup, A. E. (2010) Training with Low Muscle Glycogen Enhances

Fat Metabolism in Well-Trained Cyclists. *Med. Sci. Sport Exer.* **42**, 2046-2055

21.   Yeo, W. K., McGee, S. L., Carey, A. L., Paton, C. D., Garnham, A. P.,

Hargreaves, M., and Hawley, J. A. (2010) Acute signalling responses to intense

endurance training commenced with low or normal muscle glycogen. *Exp.*

Physiol. **95**, 351-358

22.   Yeo, W. K., Paton, C. D., Garnham, A. P., Burke, L. M., Carey, A. L., and

Hawley, J. A. (2008) Skeletal muscle adaptation and performance responses to

once a day versus twice every second day endurance training regimens. *J. Appl.*

Physiol. **105**, 1462-1470

23.   Cochran, A. J. R., Myslik, F., MacInnis, M. J., Percival, M. E., Bishop, D.,

Tarnopolsky, M. A., and Gibala, M. J. (2015) Manipulating Carbohydrate

Availability Between Twice-Daily Sessions of High-Intensity Interval Training

Over 2 Weeks Improves Time-Trial Performance. *Int. J. Sport Nutr. Exe.* **25**,

463-470

24.   Gejl, K. D., Thams, L., Hansen, M., Rokkedal-Lausch, T., Plomgaard, P., Nybo,

L., Larsen, F. J., Cardinale, D. A., Jensen, K., Holmberg, H. C., Vissing, K., and

Ortenblad, N. (2017) No Superior Adaptations to Carbohydrate Periodization in

Elite Endurance Athletes. *Med. Sci. Sports Exerc.* **49**, 2486-2497

25.   Cochran, A. J. R., Little, J. P., Tarnopolsky, M. A., and Gibala, M. J. (2010)

Carbohydrate feeding during recovery alters the skeletal muscle metabolic

response to repeated sessions of high-intensity interval exercise in humans. *J.*

Appl. Physiol. **108**, 628-636

26.   Hansen, A. K., Fischer, C. P., Plomgaard, P., Andersen, J. L., Saltin, B., and

Pedersen, B. K. (2005) Skeletal muscle adaptation: training twice every second

day vs. training once daily. *J. Appl. Physiol.* **98**, 93-99

27.   Morton, J. P., Croft, L., Bartlett, J. D., Maclaren, D. P., Reilly, T., Evans, L.,

McArdle, A., and Drust, B. (2009) Reduced carbohydrate availability does not

modulate training-induced heat shock protein adaptations but does upregulate

oxidative enzyme activity in human skeletal muscle. *J. Appl. Physiol. (1985).*

106, 1513-1521

28.   Gibala, M. J., McGee, S. L., Garnham, A. P., Howlett, K. F., Snow, R. J., and

Hargreaves, M. (2009) Brief intense interval exercise activates AMPK and p38

MAPK signaling and increases the expression of PGC-1alpha in human skeletal

muscle. *J. Appl. Physiol. (1985).* **106**, 929-934

29.   Pilegaard, H., Ordway, G. A., Saltin, B., and Neufer, P. D. (2000)

Transcriptional regulation of gene expression in human skeletal muscle during

recovery from exercise. *Am. J. Physiol-Endoc. M.* **279**, E806-E814

30.   Russell, A. P., Hesselink, M. K., Lo, S. K., and Schrauwen, P. (2005) Regulation

of metabolic transcriptional co-activators and transcription factors with acute

exercise. *FASEB J.* **19**, 986-988

31.   McGinley, C., and Bishop, D. J. (2016) Distinct protein and mRNA kinetics of

skeletal muscle proton transporters following exercise can influence

interpretation of adaptations to training. *Exp. Physiol.* **101**, 1565-1580

32.   Millay, D. P., and Olson, E. N. (2013) Making muscle or mitochondria by

selective splicing of PGC-1alpha. *Cell Metab.* **17**, 3-4

33.   Edge, J., Bishop, D., and Goodman, C. (2006) Effects of chronic NaHCO3

ingestion during interval training on changes to muscle buffer capacity,

metabolism, and short-term endurance performance. *J. Appl. Physiol. (1985).*

101, 918-925

34.   Bishop, D., Jenkins, D. G., and Mackinnon, L. T. (1998) The relationship

between plasma lactate parameters, W-peak and 1-h cycling performance in

women. *Med. Sci. Sport Exer.* **30**, 1270-1275

35.   Lima-Silva, A. E., Pires, F. O., Bertuzzi, R., Silva-Cavalcante, M. D., Oliveira,

R. S., Kiss, M. A., and Bishop, D. (2013) Effects of a low- or a high-

carbohydrate diet on performance, energy system contribution, and metabolic

responses during supramaximal exercise. *Appl. Physiol. Nutr. Metab.* **38**, 928-

934

36.   Gollnick, P. D., Piehl, K., and Saltin, B. (1974) Selective glycogen depletion

pattern in human muscle fibres after exercise of varying intensity and at varying

pedalling rates. *J. Physiol.* **241**, 45-57

37.   Heigenhauser, G. J., Sutton, J. R., and Jones, N. L. (1983) Effect of glycogen

depletion on the ventilatory response to exercise. *J. Appl. Physiol. Respir.*

Environ. Exerc. Physiol. **54**, 470-474

38.   Bergström, J. (1962) Muscle electrolytes in man determined by neutron

activation analysis on needle biopsy specimens. *Scand. J. Clin. Lab. Invest.* **14**,

1-110

39.   Evans, W. J., Phinney, S. D., and Young, V. R. (1982) Suction applied to a

muscle biopsy maximizes sample size. *Med. Sci. Sports Exerc.* **14**, 101-102

40.   Harris, R. C., Hultman, E., and Nordesjo, L. O. (1974) Glycogen, glycolytic

intermediates and high-energy phosphates determined in biopsy samples of

musculus quadriceps femoris of man at rest. Methods and variance of values.

Scand. J. Clin. Lab. Invest. **33**, 109-120

41.   Murphy, R. M., and Lamb, G. D. (2013) Important considerations for protein

analyses using antibody based techniques: down-sizing Western blotting up-

sizes outcomes. *J. Physiol.* **591**, 5823-5831

42.   Kuang, J., Yan, X., Genders, A. J., Granata, C., and Bishop, D. J. (2018) An

overview of technical considerations when using quantitative real-time PCR

analysis of gene expression in human exercise research. *PLoS One.* **13**,

e0196438

43.   Ye, J., Coulouris, G., Zaretskaya, I., Cutcutache, I., Rozen, S., and Madden, T.

L. (2012) Primer-BLAST: a tool to design target-specific primers for

polymerase chain reaction. *BMC Bioinformatics.* **13**, 134

44.   Pfaffl, M. W., Tichopad, A., Prgomet, C., and Neuvians, T. P. (2004)

Determination of stable housekeeping genes, differentially regulated target

genes and sample integrity: BestKeeper--Excel-based tool using pair-wise

correlations. *Biotechnol. Lett.* **26**, 509-515

45.   Andersen, C. L., Jensen, J. L., and Orntoft, T. F. (2004) Normalization of real-

time quantitative reverse transcription-PCR data: a model-based variance

estimation approach to identify genes suited for normalization, applied to

bladder and colon cancer data sets. *Cancer Res.* **64**, 5245-5250

46.   Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe,

A., and Speleman, F. (2002) Accurate normalization of real-time quantitative

RT-PCR data by geometric averaging of multiple internal control genes.

Genome Biol. **3**, 532-538

47.   Schmittgen, T. D., and Livak, K. J. (2008) Analyzing real-time PCR data by the

comparative C-T method. *Nat. Protoc.* **3**, 1101-1108

48.   Baar, K. (2014) Nutrition and the adaptation to endurance training. *Sports Med.*

835   **44 Suppl 1**, S5-12

836   49.   Burke, L. M. (2010) Fueling strategies to optimize performance: training high or

837   training low? *Scand. J. Med. Sci. Sports.* **20 Suppl 2**, 48-58

838   50.   Hawley, J. A., and Burke, L. M. (2010) Carbohydrate Availability and Training

839   Adaptation: Effects on Cell Metabolism. *Exerc. Sport Sci. Rev.* **38**, 152-160

840   51.   Philp, A., Burke, L. M., and Baar, K. (2011) Altering endogenous carbohydrate

841   availability to support training adaptations. *Nestle Nutr. Inst. Workshop Ser.* **69**,

842   19-31

843   52.   Impey, S. G., Hearris, M. A., Hammond, K. M., Bartlett, J. D., Louis, J., Close,

844   G. L., and Morton, J. P. (2018) Fuel for the Work Required: A Theoretical

845   Framework for Carbohydrate Periodization and the Glycogen Threshold

846   Hypothesis. *Sports Med.* **48**, 1031-1048

847   53.   Wojtaszewski, J. F., MacDonald, C., Nielsen, J. N., Hellsten, Y., Hardie, D. G.,

848   Kemp, B. E., Kiens, B., and Richter, E. A. (2003) Regulation of 5'AMP-

849   activated protein kinase activity and substrate utilization in exercising human

850   skeletal muscle. *Am. J. Physiol. Endocrinol. Metab.* **284**, E813-822

851   54.   Gejl, K. D., Vissing, K., Hansen, M., Thams, L., Rokkedal-Lausch, T.,

852   Plomgaard, P., Meinild Lundby, A. K., Nybo, L., Jensen, K., Holmberg, H. C.,

853   and Ortenblad, N. (2018) Changes in metabolism but not myocellular signaling

854   by training with CHO-restriction in endurance athletes. *Physiol. Rep.* **6**, e13847

855   55.   Hearris, M. A., Hammond, K. M., Seaborne, R. A., Stocks, B., Shepherd, S. O.,

856   Philp, A., Sharples, A. P., Morton, J. P., and Louis, J. B. (2019) Graded

857   reductions in preexercise muscle glycogen impair exercise capacity but do not

augment skeletal muscle cell signaling: implications for CHO periodization. *J.*

Appl. Physiol. (1985). **126**, 1587-1597

56.   Marchenko, N. D., Hanel, W., Li, D., Becker, K., Reich, N., and Moll, U. M.

(2010) Stress-mediated nuclear stabilization of p53 is regulated by

ubiquitination and importin-alpha3 binding. *Cell Death Differ.* **17**, 255-267

57.   Oren, M. (1999) Regulation of the p53 tumor suppressor protein. *J. Biol. Chem.*

274, 36031-36034

58.   Matoba, S., Kang, J. G., Patino, W. D., Wragg, A., Boehm, M., Gavrilova, O.,

Hurley, P. J., Bunz, F., and Hwang, P. M. (2006) p53 regulates mitochondrial

respiration. *Science.* **312**, 1650-1653

59.   Macian, F. (2005) NFAT proteins: key regulators of T-cell development and

function. *Nat. Rev. Immunol.* **5**, 472-484

60.   Masuda, E. S., Imamura, R., Amasaki, Y., Arai, K., and Arai, N. (1998)

Signalling into the T-cell nucleus: NFAT regulation. *Cell Signal.* **10**, 599-611

61.   Mansueto, G., Armani, A., Viscomi, C., D'Orsi, L., De Cegli, R., Polishchuk, E.

V., Lamperti, C., Di Meo, I., Romanello, V., Marchet, S., Saha, P. K., Zong, H.,

Blaauw, B., Solagna, F., Tezze, C., Grumati, P., Bonaldo, P., Pessin, J. E.,

Zeviani, M., Sandri, M., and Ballabio, A. (2017) Transcription Factor EB

Controls Metabolic Flexibility during Exercise. *Cell Metab.* **25**, 182-196

62.   Medina, D. L., Di Paola, S., Peluso, I., Armani, A., De Stefani, D., Venditti, R.,

Montefusco, S., Scotto-Rosato, A., Prezioso, C., Forrester, A., Settembre, C.,

Wang, W., Gao, Q., Xu, H., Sandri, M., Rizzuto, R., De Matteis, M. A., and

Ballabio, A. (2015) Lysosomal calcium signalling regulates autophagy through

calcineurin and TFEB. *Nat. Cell Biol.* **17**, 288-299

63. Settembre, C., De Cegli, R., Mansueto, G., Saha, P. K., Vetrini, F., Visvikis, O.,

Huynh, T., Carissimo, A., Palmer, D., Klisch, T. J., Wollenberg, A. C., Di

Bernardo, D., Chan, L., Irazoqui, J. E., and Ballabio, A. (2013) TFEB controls

cellular lipid metabolism through a starvation-induced autoregulatory loop. *Nat.*

Cell Biol. **15**, 647-658

64. Long, Y. C., Glund, S., Garcia-Roves, P. M., and Zierath, J. R. (2007)

Calcineurin regulates skeletal muscle metabolism via coordinated changes in

gene expression. *J. Biol. Chem.* **282**, 1607-1614

65. Handschin, C., Rhee, J., Lin, J., Tarr, P. T., and Spiegelman, B. M. (2003) An

autoregulatory loop controls peroxisome proliferator-activated receptor gamma

coactivator 1alpha expression in muscle. *Proc. Natl. Acad. Sci. U. S. A.* **100**,

7111-7116

66. Frayn, K. N. (1983) Calculation of substrate oxidation rates in vivo from

gaseous exchange. *J. Appl. Physiol. Respir. Environ. Exerc. Physiol.* **55**, 628-

634

67. Kessen, U., Schaloske, R., Aichem, A., and Mutzel, R. (1999)

Ca(2+)/calmodulin-independent activation of calcineurin from Dictyostelium by

unsaturated long chain fatty acids. *J. Biol. Chem.* **274**, 37821-37826

68. Perry, C. G. R., Lally, J., Holloway, G. P., Heigenhauser, G. J. F., Bonen, A.,

and Spriet, L. L. (2010) Repeated transient mRNA bursts precede increases in

transcriptional and mitochondrial proteins during training in human skeletal

muscle. *J Physiol.* **588**, 4795-4810

Figure legends

Figure 1 | Experimental design. D, dinner; B, breakfast; L, lunch; MG-DE, muscle

glycogen-depleting exercise; HIIE, high-intensity interval exercise. Open dashed circles

indicate that participants replicated their usual diet, while closed circles indicate that

participants ate a low-carbohydrate (CHO) breakfast [CHO: 42.7 ± 5.0 kcal (~7%), fat:

365.9 ± 43.1 kcal (~60%), protein: 201.3 ± 23.7 kcal (~33%)]. Black arrows indicate

time point whenre muscle biopsies and blood samples were taken.

Figure 2 | Muscle glycogen concentration pre, post, and 3 h post the high-intensity

interval exercise. Data are presented as mean ± standard deviation. n = 8. *

significantly lower than control at the same time point ($P < 0.05$); # significantly lower

than pre high-intensity interval exercise for the same condition ($P < 0.05$). Two-way

analysis of variance (ANOVA) with Bonferroni post hoc test.

Figure 3 | Cytosolic protein relative abundance pre and post the high-intensity

interval exercise. (a) Representative immunoblots corresponding to total and

phosphorylated protein relative abundance measured in the cytosolic fraction, pre and

post the high-intensity interval exercise in the control, twice-a-day, and once-daily

approaches; **(b)** LDHA was used as indicators of cytosolic enrichment. N: nuclear

fractions; C: cytosolic fractions; **(c)** cytosolic peroxisome proliferator-activated

receptor-γ coactivator-1 (PGC-1α); **(d)** cytosolic p53 (p53); **(e)** cytosolic

phosphorylated p53 (p-p53^{Ser15}); **(f)** cytosolic PHF20 (PHF20); **(g)** cytosolic

phosphorylated 5' adenosine monophosphate-activated protein kinase (p-AMPK^{Thr172});

(h) cytosolic phosphorylated p38 mitogen-activated protein kinase (p-p38MAPK); **(i)**

cytosolic transcription elongation factor EB (TFEB); **(j)** cytosolic nuclear factor of

activated T cells (NFAT). n = 8 for all proteins. Data are presented as fold changes from

control pre (mean ± standard deviation). # significantly higher than pre high-intensity

interval exercise for the same condition ($P < 0.05$). Two-way analysis of variance

(ANOVA) with Bonferroni post hoc test.

Figure 4 | Nuclear protein relative abundance pre and post the high-intensity

interval exercise. (a) Representative immunoblots corresponding to total and

phosphorylated protein relative abundance measured in the nuclear fraction, pre and

post the high-intensity interval exercise in the control, twice-a-day, and once-daily

approaches; **(b)** H3 was used as indicators of nuclear enrichment. N: nuclear fractions;

C: cytosolic fractions; **(c)** nuclear peroxisome proliferator-activated receptor-γ

coactivator-1 (PGC-1α); **(d)** nuclear p53 (p53); **(e)** nuclear phosphorylated p53 (p-

p53^{Ser15}); **(f)** nuclear PHF20 (PHF20); **(g)** nuclear phosphorylated AMPKThr172 (p-

AMPK^{Thr172}); **(h)** nuclear phosphorylated p38MAPK (p-p38MAPK); **(i)** nuclear

transcription elongation factor EB (TFEB); **(j)** nuclear factor of activated T cells

(NFAT). n = 8 for all proteins. Data are presented as fold changes from control pre

(mean ± standard deviation). * significantly higher than the once-daily and control

condition at the same time point ($P < 0.05$); † significantly higher than the once-daily

condition at the same time point ($P < 0.05$). # significantly higher than pre high-

intensity interval exercise for the same condition ($P < 0.05$). Two-way analysis of

variance (ANOVA) with Bonferroni post hoc test.

Figure 5 | Mitochondrial-related gene expression pre, post, and 3 h post the high-

intensity interval exercise. (a) peroxisome proliferator-activated receptor-γ

coactivator-1 (PGC-1α) total gene expression; **(b)** PGC-1α isoform 1 gene expression;

(c) PGC-1α isoform 4 gene expression; **(d)** p53 (p53) gene expression; **(e)** transcription

elongation factor EB (TFEB) gene expression; **(f)** Chromodomain-helicase-DNA-

binding protein 4 (CHCHD4) gene expression; **(g)** p21 protein (p21) gene expression;

(h) mitochondrial transcription factor A (Tfam) gene expression; **(i)** NADH

dehydrogenase (NDUFβ) (mitochondrial complex I) gene expression; **(j)** succinate

dehydrogenase subunit β (SDHβ) (mitochondrial complex II) gene expression; **(k)**

cytochrome c (mitochondrial complex III) gene expression; **(l)** cytochrome c oxidase

subunit IV (COXIV) (mitochondrial complex III) gene expression. n = 8 for all genes

(except Tfam, n = 7). Data are presented as fold changes from control pre (mean ±

standard deviation). * significantly higher than the once-daily and control condition at

the same time point ($P < 0.05$); # significantly higher than pre high-intensity interval

exercise for the same condition ($P < 0.05$); ☐ significantly higher than post high-

intensity interval exercise for the same condition ($P < 0.05$). Two-way analysis of

variance (ANOVA) with Bonferroni post hoc test.

Figure 6 | Genes related to fat transport and lipolysis pre, post, and 3 h post the

high-intensity interval exercise. (a) carnitine palmitoyltransferase I (CPT1) gene

expression; **(b)** mitochondrial uncoupling protein 3 (UCP3) gene expression; **(c)**

peroxisome proliferator-activated receptor α (PPARα) gene expression; **(d)** peroxisome

proliferator-activated receptor β/δ (PPARβ/δ) gene expression; **(e)** peroxisome

proliferator-activated receptor γ (PPARγ) gene expression; **(f)** citrate synthase (CS)

gene expression; **(g)** β-hydroxyacyl-CoA dehydrogenase (β-HAD) gene expression; **(h)**

fatty acid translocase cluster of differentiation 36 (CD-36) gene expression. n = 8 for all

genes (except CS and UCP3, where n = 7). Data are presented as fold changes from

control pre (mean ± standard deviation). * significantly higher than the once-daily

condition at the same time point ($P < 0.05$); † significantly higher than the control

condition at the same time point ($P < 0.05$); # significantly higher than pre high-

intensity interval exercise for the same condition ($P < 0.05$); ☐ significantly higher than

post high-intensity interval exercise for the same condition ($P < 0.05$). Two-way

analysis of variance (ANOVA) with Bonferroni post hoc test.

Figure 7 | Genes related to carbohydrate metabolism pre, post, and 3 h post the

high-intensity interval exercise. (a) phosphofructokinase (PFK) gene expression; **(b)**

glucose transporter 4 (GLUT4) gene expression; **(c)** pyruvate dehydrogenase kinase

isoenzyme 4 (PDK4) gene expression. n = 8 for all genes. Data presented as fold

changes from control pre (mean ± standard deviation). # significantly higher than pre

high-intensity interval exercise for the same condition ($P < 0.05$). Two-way analysis of

variance (ANOVA) with Bonferroni post hoc test.

Figure 8 | Physiological and systemic responses during the high-intensity interval

exercise. (a) Heart rate (HR); **(b)** Ventilation ($\dot{V}E$); **(c)** Oxygen uptake ($\dot{V}O_2$); **(d)**

Respiratory exchange ratio (RER). n = 8 for all variables. Data are presented as mean ±

standard deviation. Time effect has been omitted for clarity. INT: Interval. Two-way

analysis of variance (ANOVA) with Bonferroni post hoc test.

a

main effect of condition, *P* = 0.01
(twice-a-day higher than once-daily and control)

b

main effect of condition, *P* = 0.03
(twice-a-day higher than once-daily)

c

main effect of condition, *P* = 0.01
(twice-a-day higher than control)

d

main effect of condition, *P* = 0.01
(twice-a-day lower than control)

Table 1 | Analysis of checklists for energy or macronutrient intake between conditions for the dinner eaten two days before, and breakfast, lunch and dinner eaten one day before the experimental trial.

		Dinner two days before the experimental trial	Breakfast one day before the experimental trial	Lunch one day before the experimental trial	Dinner one day before the experimental trial
Energy intake (Kcal)	**Control**	911 ± 348	613 ± 358	955 ± 233	852 ± 215
	Twice-a-day	918 ± 354	602 ± 366	925 ± 236	886 ± 237
	Once-daily	887 ± 334	602 ± 366	948 ± 236	884 ± 236
Carbohydrate (%)	**Control**	45 ± 19	52 ± 12	44 ± 5	50 ± 9
	Twice-a-day	45 ± 19	53 ± 12	46 ± 6	49 ± 9
	Once-daily	43 ± 18	53 ± 12	44 ± 5	49 ± 9
Protein (%)	**Control**	28 ± 17	17 ± 8	32 ± 5	21 ± 12
	Twice-a-day	28 ± 17	17 ± 8	33 ± 6	21 ± 12
	Once-daily	28 ± 17	17 ± 8	34 ± 7	21 ± 12
Lipids (%)	**Control**	27 ± 10	31 ± 7	23 ± 10	29 ± 8
	Twice-a-day	27 ± 10	30 ± 8	21 ± 11	30 ± 9
	Once-daily	30 ± 10	30 ± 8	21 ± 11	30 ± 9

Data are presented as mean $\pm$ standard deviation. n = 8 for all variables.

Table 2 | Plasma lactate, glucose, epinephrine and norepinephrine concentrations, and serum free fatty acid and glycerol concentrations pre, post and 3 h post the high-intensity interval exercise.

	Pre	Post	3 h post
Plasma			
[Lactate] (mmol·L^{-1})			
Control	1.2 ± 0.7	6.2 ± 2.0 □	1.3 ± 0.4
Once-daily	0.9 ± 0.3	3.9 ± 1.0 □†	1.2 ± 0.4
Twice-a-day	1.4 ± 0.5	4.0 ± 1.6 □†	2.0 ± 0.7
[Glucose] (mmol·L^{-1})			
Control	5.2 ± 0.8	5.5 ± 0.9	5.3 ± 0.8
Once-daily	5.2 ± 0.5	5.0 ± 0.8 †	4.8 ± 0.7 †#
Twice-a-day	4.4 ± 0.7 *†	4.8 ± 1.1 □†	4.3 ± 1.0 *†
[Epinephrine] (pg·mL^{-1})			
Control	32.2 ± 18.4	39.3 ± 22.7	36.7 ± 17.8
Once-daily	56.7 ± 25.4	54.2 ± 25.7	35.8 ± 34.8
Twice-a-day	43.1 ± 13.6	43.7 ± 19.0	36.7 ± 10.4
[Norepinephrine] (pg·mL^{-1})			
Control	68.0 ± 34.6	231.1 ± 141.8 □	120.4 ± 90.3
Once-daily	70.7 ± 33.8	350.6 ± 139.8 □	105.5 ± 77.5
Twice-a-day	76.7 ± 49.1	287.5 ± 147.6 □	113.8 ± 44.6
Serum			
[Free Fatty Acid] (μM)			
Control	194.9 ± 118.2	118.6 ± 72.6	130.3 ± 72.1
Once-daily	155.5 ± 79.9	236.5 ± 151.4	158.0 ± 62.8
Twice-a-day	209.9 ± 125.5 †	260.6 ± 151.3 †	230.4 ± 98.2 †
[Glycerol] (mmol·L^{-1})			
Control	0.105 ± 0.021	0.098 ± 0.025	0.098 ± 0.037
Once-daily	0.090 ± 0.037	0.141 ± 0.022 †	0.100 ± 0.039
Twice-a-day	0.106 ± 0.031	0.169 ± 0.046 †	0.102 ± 0.023

Data are presented as mean ± standard deviation. n = 8 for all variables. * significantly different from the once-daily at the same time point ($P < 0.05$); † significantly different from the control condition at the same time point ($P < 0.05$); # significantly different from pre high-intensity interval exercise for the same condition ($P < 0.05$); □ significantly different from pre and post high-intensity interval exercise for the same condition, except for glucose that was significantly different only from post high-intensity interval exercise ($P < 0.05$). Two-way analysis of variance (ANOVA) with Bonferroni post hoc test.