Lotus japonicus symbiosis signaling genes and their role in the establishment of root-
associated bacterial and fungal communities

Rafal Zgadzaj[§ 1,2], Thorsten Thiergart[§ 1], Zoltán Bozsóki [3], Ruben Garrido-Oter [1,2], Simona
Radutoiu[*3], and Paul Schulze-Lefert[*1,2]

[1]Max Planck Institute for Plant Breeding Research, 50829 Cologne, Germany.

[2]Cluster of Excellence on Plant Sciences (CEPLAS), Max Planck Institute for Plant Breeding
Research, 50829 Cologne, Germany.

[3]Department of Molecular Biology and Genetics, Faculty of Science and Technology, Aarhus
University, 8000 C Aarhus, Denmark.

§ joint first authors

* co-corresponding authors, (radutoiu@mbg.au.dk , S.R., schlef@mpipz.mpg.de , P.S.-L.)

Abstract

The wild legume *Lotus japonicus* engages in mutualistic symbiotic relationships with arbuscular mycorrhiza (AM) fungi and nitrogen-fixing rhizobia. Using plants grown in natural soil and community profiling of bacterial 16S *rRNA* genes and fungal internal transcribed spacers (ITS), we examined here the role of the *Lotus* symbiosis genes *RAM1*, *NFR5*, *SYMRK*, and *CCaMK* in structuring bacterial and fungal root-associated communities. We found host genotype-dependent community shifts in the root and rhizosphere compartments that were mainly confined to bacteria in *nfr5* or fungi in *ram1* mutants, whilst *symRK* and *ccamk* plants displayed major changes across both microbial kingdoms. We observed in all AM mutant roots an almost complete depletion of a large number of Glomeromycota taxa that was accompanied by a concomitant enrichment of Helotiales and *Nectriaceae* fungi, suggesting compensatory niche replacement within the fungal community. A subset of Glomeromycota whose colonization is strictly dependent on the common symbiosis pathway was retained in *ram1* mutants, indicating that *RAM1* is dispensable for intraradical colonization by some Glomeromycoyta fungi. However, intraradical colonization by certain Burkholderiaceae taxa is dependent on AM root infection, thereby revealing a microbial interkingdom interaction. Our findings imply a broad role for *Lotus* symbiosis genes in structuring the root microbiota.

Introduction

Mutualistic plant-microbe interactions are essential adaptive responses dating back to plant colonization of terrestrial habitats [1,2]. Endosymbiotic association with obligate arbuscular mycorrhizal (AM) fungi belonging to the phylum Glomeromycota is considered to have enabled early land plants to adapt to and survive harsh edaphic conditions by improving the acquisition of nutrients, especially phosphorus, and water from soil [3]. It is estimated that approximately 80% of extant plant species remain proficient in AM symbiosis (AMS), testifying to its importance for survival in natural ecosystems [4,5,6]. Another more recent endosymbiotic relationship has evolved between plants belonging to distinct lineages of flowering plants (Fabales, Fagales, Cucurbitales, and Rosales) and nitrogen-fixing members of the Burkholderiales, Rhizobiales or Actinomycetales, enabling survival on nitrogen-poor soils. These bacteria fix atmospheric nitrogen under the low oxygen conditions which are provided by plant root nodules.

Studies of mutant legumes deficient in both AM and RNS revealed that a set of genes defined as the common symbiotic signaling pathway (CSSP) are crucial for these symbioses. In the model legume *Lotus japonicus*, Nod factor perception by NFR1 and NFR5 activates downstream signaling through SYMRK, a malectin and leucine-rich repeat (LRR)-containing RLK [7], currently considered to be the first component of the CSSP. SYMRK associates with NFR5 through a mechanism involving intramolecular cleavage of the SYMRK ectodomain, thereby exposing its LRR domains [8]. Signaling from the plasma membrane is transduced to the nuclear envelope where ion channels [9,10], nuclear pore proteins [11,12,13] and cyclic nucleotide-gated channels [14] mediate symbiotic calcium oscillations. These calcium oscillations are interpreted by the calcium- and calmodulin-dependent protein kinase CCaMK, [15,16] that interacts with the DNA binding transcriptional activator CYCLOPS [17,18,19]. Several GRAS transcription factors (NSP1, NSP2, RAM1, RAD1) are activated downstream of CCAMK and CYCLOPS and determine whether plants engage in AMS or RNS symbiosis.

Plants establish the symbioses with AM fungi and nitrogen-fixing bacteria by selecting interacting partners from the taxonomically diverse soil biome. These interactions are driven by low mineral nutrient availability in soil and induce major changes in host and microbial symbiont metabolism [20,21]. Although RNS develops as localized events on legume roots, analysis of *Lotus* mutants impaired in their ability to engage in symbiosis with nitrogen-fixing bacteria revealed that these mutations do not only abrogate RNS, but also impact the composition of taxonomically diverse root- and rhizosphere-associated bacterial communities, indicating an effect on multiple bacterial

taxa that actively associate with the legume host, irrespective of their symbiotic capacity [22]. By contrast, the effect of AMS is known to extend outside the host *via* a hyphal network that can penetrate the surrounding soil and even indirectly affect adjacent plants [23]. In soil, fungal hyphae themselves represent environmental niches and are populated by a specific set of microbes [24]. Although the biology of AMF is well understood, the potential role of AMS on root-associated bacterial and fungal communities is currently unknown.

Here we used mutants impaired in RNS, AM, or both to address the role of symbiosis signaling genes in structuring bacterial and fungal root-associated communities in *Lotus* plants grown in natural soil. We show that genetic disruption of these symbioses results in significant host genotype-dependent microbial community shifts in the root and surrounding rhizosphere compartments. These changes were mainly confined to either bacterial or fungal communities in RNS- or AM-deficient plant lines, respectively, whereas mutants with defects in the CSPP revealed major changes in assemblages of the root microbiota across both microbial kingdoms.

Materials and Methods

Preparation and storage of soil

The soil batches used in this study were collected from Max Planck Institute for Plant Breeding Research agricultural field located in Cologne, Germany (50.958N, 6.865E) in the following seasons: CAS11-spring/autumn 2016, CAS12-spring 2017. The field had not been cultivated in previous years, no fertilizer or pesticide administration took place at the harvesting site. Following harvest, soil was sieved, homogenized and stored at 4 °C for further use.

Soil and plant material

All studied *L. japonicus* symbiosis-deficient mutants, *nfr5-2* [ref. 25], *ram1-2* [ref. 26], *symrk-3* [ref. 7] and *ccamk-13* [ref. 27], originated from the Gifu B-129 genotype.

Plant growth and harvesting procedure

The germination procedure of *L. japonicus* seeds included sandpaper scarification, surface sterilisation in 1% hypochlorite bleach (20 min, 60 rpm), followed by three washes with sterile water and incubation on wet filter paper in Petri dishes for one week (temperature: 20 °C, day/night cycle 16/8h, relative humidity: 60%). For each genotype and soil batch, six to eight biological replicas

were prepared by potting four plants in 7x7x9 cm pot filled with corresponding batch of soil (CAS11 soil six replicates, CAS12 soil eight replicated). Plants were incubated for ten weeks in the greenhouse (day/night cycle 16/8h, light intensity 6000 LUX, temperature: 20 °C, relative humidity: 60%), and were watered with tap water twice per week.

The block of soil containing plant roots was removed from the pot and adhering soil was discarded manually. Three sample pools were collected: complete root systems (harvested 1 cm below the hypocotyl), upper fragments of the root systems (4 cm-long, starting 1 cm below the hypocotyl) and lower root system fragments (harvested from 9 cm below; the latter two were collected from plants grown in the same pot (Fig. 1a). All pools were washed twice with sterile water containing 0.02% Triton X-1000 detergent and twice with pure sterile water by vigorous shaking for 1 min. The rhizosphere compartment was derived by collection of pellet following centrifugation of the first wash solution for 10 min at 1500 g. The nodules and visible primordia were separated from washed root pools of nodulating genotypes (WT and ram1-2) with a scalpel and discarded. In order to obtain the root compartment the root sample pools were sonicated to deplete the microbiota fraction attached to the root surface. It included 10 cycles of 30-second ultrasound treatment (Bioruptor NextGen UCD-300, Diagenode) for complete root systems and upper root fragments, while for the lower root fragments the number of cycles was reduced to three. All samples were stored at -80 °C for further processing. For AM colonisation inspection the whole root system of washed soil-grown plants was stained with 5% ink in 5% acetic acid solution and inspected for intraradical infection.

Generation of 16S rRNA and ITS2 fragment amplicon libraries for Illumina MiSeq sequencing

Root pool samples were homogenized by grinding in a mortar filled with liquid nitrogen and treatment with Precellys24 Tissue lyser (Bertin Technologies) for two cycles at 5600 rpm for 30 sec. DNA was extracted with the FastDNA Spin Kit for Soil, according to the manufacturer's protocol (MP Bioproducts). DNA concentrations were measured fluorometrically (Quant-iT™ PicoGreen dsDNA assay kit, Life Technologies, Darmstadt, Germany), and adjusted to 3.5 ng/µl. Barcoded primers targeting the variable V5-V7 region of the bacterial 16S rRNA gene (799F and 1193R, [28]) or targeting the ITS2 region of the eukaryotic ribosome (fITS7 and ITS4, [29,30]) were

129 used for amplification. The amplification products were purified, pooled and subjected to
130 sequencing with Illumina MiSeq equipment.

Processing of 16S rRNA and ITS2 reads

All libraries from root fractionation experiments as well as from the main experiments were
analyzed together. Due to a very low read count for 16S data in the first experiment in CAS11 soil,
this library was not included in the final analysis. This resulted in an overall lower sample number
for bacteria than for fungi (222 vs. 274 samples). All sets of amplicon reads were processed as
recently described [31], using a combination of QIIME [32] and USEARCH tools [33]. For both
datasets paired end reads were used. For ITS2 data, forward reads were kept, in case that no
paired version was available. Main steps include quality filtering of reads, de-replicating, chimera
detection and OTU clustering at a 97% threshold. 16S reads were filtered against the greengenes
data base [34], whereas for ITS2 the reads where checked with ITSx [35] and compared against a
dedicated ITS database to remove ITS sequences from non-fungal species. Taxonomic
classification was done with uclust (assign_taxonomy from QIIME) for 16S OTUs and rdp classifier
[36] for ITS2 OTUs. For the sake of consistency with NCBI taxonomic classification, the
assignment of the ITS2 sequences was manually corrected so that that all OTUs assigned as
Ilyonectria were assigned as belonging to the Sordariomycetes, Hypocreales and Nectriaceae,
respectively. For 16S data, OTUs assigned as mitochondrial or chloroplast, were removed prior to
analysis.

Statistical analysis

For calculating Shannon diversity indices, OTU tables were rarefied to 1000 reads. Significant
differences were determined using ANOVA (aov function in R) and post-hoc Tukey test (TukeyHSD
in R, $p < 0.05$). For calculating Bray Curtis distances between samples, OTU tables were
normalized using cumulative sum scaling (CSS, [37]). Bray-Curtis distances were used as input for
principal coordinate analysis (PCoA, cmdscale function in R) plots and as input for constrained
analysis of principal coordinate (CPCoA, capscale function, vegan package in R). For the latter, the
analysis was constrained by genotypes (each mutant and WT separately) and corrected for the
effect of soil type and experiment (using the "Condition" function). This analysis has been repeated
with OTU tables from which OTUs that represent known plant symbionts (Phyllobacteriaceae for
16S and Glomeromycota for ITS2) were removed before normalization, distance calculation and
CPCoA. A previously described approach was used to draw ternary plots and for respective

enrichment analysis [22]. Fold change of OTUs between wild type and mutant plants was
calculated as followed. Samples showing a read count <5000 were removed. OTUs with mean RA
>0.1% across all root or rhizosphere samples, respectively, were kept for analysis. Fold change in
RA from WT to mutants was calculated over all WT samples for nfr5-2, ram1-2 and symrk-3,
whereas the change to ccamk-13 was only calculated with WT samples from experiments where
ccamk-13 mutants were present. To avoid zeros in calculation the RA of OTUs missing from
samples was set to 0.001%. The significance of differences in abundance was tested using the
Kruskal-Wallis test (p<0.05).

Results

Root fractionation protocol affects the composition of associated bacterial communities

Earlier physiological studies have shown that only cells of a specific developmental stage, located
in the root elongation zone, respond to Myc and Nod factors, mount symbiotic calcium oscillations
and enable epidermal infection by rhizosphere-derived fungal and bacterial symbionts [38,39].To
explore spatial organization of root-associated bacterial and fungal communities along the
longitudinal axis, we collected samples of the upper and lower root zones as well as the entire root
system of six-week-old Gifu wild-type plants, grown in Cologne soil (2 to 5 cm and >9 cm of the
root system, respectively; Fig. 1A; [40]). Microbial assemblages of these three root endosphere
compartments were compared to the communities in the corresponding rhizosphere fractions, i.e.
soil tightly adhering to the respective root zones, and with the bacterial biome present in unplanted
Cologne soil. 16S rRNA gene amplicon libraries of the V5-V7 hypervariable region and gene
libraries of the Internally Transcribed Spacer 2 (ITS2) region of the eukaryotic ribosome were
generated by amplification [28,29,30]. Information on the number and relative abundance of
operational taxonomic units (OTUs) in each compartment was used to calculate α (Shannon index;
within sample diversity) and β-diversity (Bray-Curtis distances; between samples diversity), OTU
enrichment and taxonomic composition. In bacteria we observed a gradual decrease in α-diversity
from unplanted soil to the rhizosphere and to the root endosphere compartments, a trend which
was similar for each longitudinal root fraction. This suggests that winnowing of root commensals
from the highly complex soil biome occurs in all tested root zones (Fig. S1A). Similar overall results
were obtained for the fungal dataset (Fig. S1B), but the decrease in diversity from unplanted soil
towards the rhizosphere was mild or even lacking. The latter finding is similar to that of a recent
study of root-associated fungi in non-mycorrhizal *A. thaliana* sampled at three natural sites [31].
Analyses of taxonomic composition and β-diversity revealed striking differences in the endosphere

and rhizosphere compartments associated with the upper and lower root longitudinal fractions. The

composition of bacterial and fungal taxa of the whole root closely resembled that of the upper root

fraction (Fig. 1B), with only low numbers of OTUs differentially abundant between these two

compartments (Fig. 1C and 1D). Additionally, we observed a higher sample-to-sample variation in

the taxonomic profiles of the lower root zone compared to the upper whole root fractions (Fig. 1B).

This greater community variation in the developmentally younger region of *L. japonicus* roots might

reflect a nascent root microbiota or greater variation in root tissue and adherent rhizosphere

samples that we recovered from this root zone by our fractionation protocol. Based on the finding

that whole root and upper root compartments host comparable bacterial communities and given

their greater stability we decided to use the former for further analyses.

Host genes needed for symbioses determine bacterial and fungal community composition
of *L. japonicus* root and rhizosphere

For root microbiota analysis, we cultivated wild-type (ecotype Gifu) *L. japonicus* and *nfr5-2*, *symrk-*

3, *ccamk-13* and *ram1-2 (nfr5, symrK, ccamk* and *ram1,* from thereof)* mutant genotypes in parallel

in two batches of Cologne soil, to account for batch-to-batch and seasonal variation at the

sampling site. *nfr5-2* mutant plants are impaired in rhizobial Nod factor perception and signaling,

which prevents initiation of infection thread formation [25]. Mutations in *SymrK* and *CcamK* affect

the common symbiosis pathway downstream of Nod or Myc factor perception, abrogating infection

either by nitrogen fixing rhizobia or AM fungi [7,27]. The RAM1 transcription factor controls

arbuscule formation, and while *ram1-2* mutants of *L. japonicus* are indistinguishable from wild type

and permit incipient AM fungus infection, fungal colonization is terminated with the formation of

stunted symbiotic structures [26]. All plant genotypes appeared healthy (Fig. 2A-E), but the shoot

length and shoot fresh weight of all mutant plants was significantly reduced in comparison to wild

type (Fig. 2F and 2G), suggesting that genetic disruption of either AM or *Rhizobium* symbiosis is

detrimental for the fitness of plants grown in natural soil. Whereas all defects in nitrogen-fixing

symbiosis, validated by the absence of root nodules in *nfr5-2*, *symrk-3* and *ccamk-13* genotypes

(Fig. 2C-E, Table S1), resulted in similarly severe impacts on plant growth (Fig. 2F and 2G), both

shoot length and shoot fresh weight were significantly reduced in *ram1-2* plants, although the

effects were less severe and these plants still formed nodules and, unlike wild-type and *nfr5-2*,

showed impairment in AM symbiosis (Table S1) and a less severe, but significant reduction of both

shoot length and shoot fresh weight in comparison to wild type.

In order to determine the impact of rhizobial and AM symbiosis on root microbiota assembly, we

characterized fungal and bacterial communities of unplanted Cologne soil, rhizosphere, and root

compartments of all aforementioned *L. japonicus* genotypes at bolting stage. Visible nodules and

root primordia were removed from the roots of nodulating wild type and *ram1*-2 genotypes prior to

sample processing for community profiling. We amplified the V5-V7 hypervariable region of the

bacterial 16S rRNA gene and the ITS2 region of the eukaryotic ribosome. High-throughput

sequencing of these amplicons yielded 22,761,657 16S and 21,228,781 ITS reads, distributed in

222 and 274 samples, respectively, which were classified into 5,780 and 3,361 distinct microbial

OTUs. Analysis of α-diversity revealed a general reduction of complexity from unplanted soil to

rhizosphere and lastly in root compartments for bacterial communities, whereas the complexity of

fungal communities was largely similar for the latter two compartments (Fig. S2A and S2B), which

is consistent with a recent study of *A. thaliana* root-associated fungal communities [31]. Bacterial

α-diversity was slightly elevated in the *nfr5*-2 genotype in rhizosphere and root compartments in

comparison to all other genotypes (Fig. S2A). Fungal communities were similarly diverse in the

rhizosphere of all tested plant genotypes, but their diversity in the root compartment was

significantly and specifically reduced in all three AM mutants (*ccamk*-13, *ram1*-2, and *symrk*-3; Fig.

S2B).

Analysis of β-diversity using Principal Coordinate Analysis (PCoA) of Bray-Curtis distances

showed a significant effect of soil batch on soil-resident bacterial and fungal communities (Fig. S2C

and D). In order to account for this technical factor and assess the impact of the different host

compartment and genotypes in community composition, we performed a Canonical Analysis of

Principle Components Coordinates (CAP; [41]). This revealed a clear differentiation of bacterial

and fungal communities in the tested plant genotypes in both root and rhizosphere compartments,

with the host genotype explaining as much as 7.61% of the overall variance of the 16S rRNA, and

13.5% of ITS2 data (Fig. 3; P<0.001). The rhizosphere compartments of wild type and *ram1-2* were

found to harbor similar bacterial communities, but were separate from those of *symrk-3* and

ccamk-13 (Fig. 3A). Further, the rhizosphere communities of each of these four plant genotypes

were found to be significantly different from that of *nfr5* (Fig. 3A). A similar trend was observed for

fungal communities, except that wild-type and *ram1* rhizosphere communities were clearly

separated from each other (Fig. 3C). In the root compartment we found bacterial consortia that

were distinctive for each of the five plant genotypes (Fig. 3B). A pronounced host genotype effect

was also found for the root-associated fungal communities, but in this case the communities of

wild-type and *nfr5* clustered together, indicating a high similarity (Fig. 3D). *L. japonicus* is

nodulated by members of *Mesorhizobium loti* belonging to the Phyllobacteriaceae family, and

engages in AM symbiosis with Glomeromycota fungi. The plant genotypes included in this study

differed in their capacity to accommodate nitrogen-fixing bacteria and/or AM fungi inside roots,

and, consequently, the lack of these symbionts alone might explain the microbial community

separation observed by CAP analysis (Fig. 3). To test whether these symbiotic microbes are the

sole drivers of the observed community separations, we repeated the CAP analysis after *in silico*

removal of *Phyllobacteriaceae* sequencing reads for bacterial communities, and of Glomeromycota

reads for the fungal assemblages. Interestingly, although this reduced the community variance

explained by host genotype across the dataset (Fig. S3 compared to Fig. 3), overall patterns of β-

diversity remained unaltered, suggesting that other community members besides root nodule and

arbuscular mycorrhizal symbionts contribute to the plant genotype-specific community shifts.

Collectively, our analyses of *L. japonicus* symbiotic mutants grown in natural soil show that lack of

AM and/or RNS symbioses has a significant effect on plant growth and on the structures of

bacterial and fungal communities associated with legume roots.

Loss of symbiosis affects specific bacterial and fungal families of the root microbiota

Comparison of bacterial family abundance between wild type and mutants lacking RNS and/or AM

symbiosis identified significant changes in *Comamonadaceae*, *Phyllobacteriaceae*,

Methylophilaceae, *Cytophagaceae* and *Sinobacteracea* in the rhizosphere compartment (Fig. 4A;

top 10 most abundant families). The abundance of *Comamonadaceae* and *Phyllobacteriaceae*

also differed significantly in the root compartment of RNS mutants compared to wild type.

Streptomycetaceae and *Sinobacteraceae* relative abundances were specifically affected by loss of

Nfr5, whereas *Anaeroplamataceae* and *Burkholderiaceae* abundances were affected by the lack of

AM symbiosis in *symrk* and *ccamk* plants (Fig. 4A). The relative abundances of the same two

families were also significantly reduced in *ram1* roots, suggesting that active AM symbiosis

influences root colonization by a subset of bacterial root microbiota taxa.

Six out of the ten most abundant fungal families in the rhizosphere compartment of *Lotus* plants

belonged to Ascomycota (Fig. 4B). By contrast, the root endosphere was dominated by numerous

families of Glomeromycota, which were found to be almost fully depleted from the rhizosphere and

root compartments of *ram1*, *symrk* and *ccamk* mutants, indicating that absence of AM symbiosis

predominantly affects Glomeromycota and does not limit root colonization or rhizosphere

association by other fungal families. However, depletion of Glomeromycota in the AM mutant roots

was accompanied by an increase in the relative abundance of Ascomycota members belonging to

Nectriaceae in both rhizosphere and root compartments and by an increased abundance of

unclassified *Helotiales*, *Leotiomycetes*, and *Sordariomycetes* in the root compartment only (Fig.

4B).

Closer inspection of the microbial community shifts at the OTU level identified 45 bacterial OTUs

and 87 fungal OTUs enriched in the roots of symbiosis mutants compared to those of wild type

(Fig. 5), and 60 bacterial OTUs and 30 differentially abundant fungal OTUs in the rhizosphere

samples (Fig. S4). The absence of RNS in *nfr5-2* roots affected the relative abundance of multiple

OTUs (n=27 in the root, n=23 in the rhizosphere) belonging to diverse taxa. Many of these OTUs

(n=18 in the root, n=16 in the rhizosphere) showed a similar differential relative abundance in

symrk-3 and/or *ccamk-13* mutants when compared to wild type (Fig.5A), indicating that their

contribution to the *Lotus* root communities outside of nodules is affected by active nitrogen fixing

symbiosis. Impairment of both AM and RNS symbioses in *symrk* and/or *ccamk* mutants resulted in

opposite changes in the relative root abundances of OTUs belonging to specific Burkholderiales

families. Depletion of OTUs belonging to *Burkholderiaceae* (n=5) was accompanied by the

enrichment of OTUs from other Burkholderiales families (*Oxalobacteraceae* [n=3],

Comamonadadaceae [n=2], and *Methylophilaceae* [n=2]; Fig. 5A). Only three of the above-

mentioned *Burkholderiaceae* OTUs were depleted in *ram1* roots, suggesting that their enrichment

in *Lotus* roots is dependent on functional AM symbiosis.

Analysis of the ITS2 amplicon sequences from root samples identified a large number of

Glomeromycota OTUs (n=39); thus, *Lotus* Gifu roots grown in natural soil accommodate a

phylogenetically diverse community of AM fungi (Fig. 5B). The majority of these fungal OTUs

(n=31) were depleted in *symrk-3*, *ccamk-13* and *ram1-2* mutant roots, indicating that their

enrichment is dependent on a functional AM symbiosis pathway. Their intraradical colonization

appears to be independent of *RAM1*, as 12 OTUs assigned to Glomeromycota or to unknown taxa,

nine of which define a Glomeromycota sublineage, were depleted in *symrk-3* and *ccamk-13* but not

in *ram1-2* roots. A reduced abundance of Glomeromycota OTUs in the endosphere compartment

was accompanied by an increased abundance of Ascomycota members, especially of members

belonging to the *Nectriaceae* (8 OTUs) and *Helotiales* (7 OTUs) families, suggestive of a mutually

exclusive occupancy of the intraradical niche. In sum, our results reveal that in natural soil CSSP

symbiotic genes are essential for root colonization by a wide range of Glomeromycota fungi;

further, these genes significantly affect the abundances of multiple bacterial taxa, predominantly

belonging to the Burkholderiales and Rhizobiales orders.

Discussion

Here, we investigated the role of host AM and/or RNS genes in establishing structured bacterial and fungal communities in the rhizosphere and endosphere compartments of *L. japonicus* grown in natural soil. Impairment of RNS in *nfr5-2* or AMS in *ram1-2* plants had a significant impact on root microbiota structure, which was mainly confined to the composition of bacterial or fungal communities, respectively.

The shift between the root-associated microbial communities of wild type and *nfr5-2* mutant is in line with both the qualitative and quantitative findings of a previous report on the *Lotus* bacterial root microbiota (Fig. 3A and B in this study; [22]). However, here we observed an enhanced rhizosphere effect in both wild type and *nfr5-2* plants, leading also to a less prominent community shift in this compartment (Fig. S5). These differences in rhizosphere bacterial composition are likely caused by a soil batch effect and, to a lesser extent, possibly also the use of different sequencing platforms (Illumina *versus* 454 pyrosequencing). The nearly unaltered fungal community composition in *nfr5* mutant plants compared to wild type (only 3 out of 39 Glomeromycota OTUs differentially abundant) suggests that NFR5 is dispensable for fungal colonization of *L. japonicus* roots. This finding, together with a massive shoot biomass reduction of *nfr5* plants in natural soil (~4-fold; Fig. 2), further reveals that intraradical colonization by soil-derived fungal endophytes is robust against major differences in plant growth.

A recent microbial multi-kingdom interaction study in *A. thaliana* showed that bacterial commensals of the root microbiota are crucial for the growth of a taxonomically wide range of fungal root endophytes. These antagonistic interactions between bacterial and fungal root endophytes are essential for plant survival in natural soil [31]. We have shown here that an almost complete depletion of diverse Glomeromycota taxa from roots of each of the three AM mutants was accompanied by an enrichment of OTUs belonging to the families *Nectriaceae* and Helotiales (Fig. 4). We speculate that the increased relative abundance of these taxa is caused by intraradical niche replacement as a compensatory effect following the exclusion of Glomeromycota symbionts from the root compartment. Previous mono-association experiments have shown that isolates belonging to *Nectriaceae* and Helotiales can have either mutualistic or pathogenic phenotypes [42,43,44]. Given that all plant genotypes were free of disease symptoms when grown in natural soil (Fig. 2), we speculate that the complex shifts in the composition of the bacterial root microbiota in *nfr5-2, symRK-3,* and *ccamk-13* mutants did not affect the capacity of bacterial endophytes to prevent pathogenic fungal overgrowth. Of note, Helotiales root endophytes were also enriched in

roots of healthy *Arabis alpina*, a non-mycorhyzal plant species and relative of *A. thaliana*, and contribute to phosphorus nutrition of the host when grown in extremely phosphorus-impoverished soil [45]. The enrichment of Helotiales in *Lotus* AM mutants is therefore consistent with potential niche replacement by other fungal lineages to ensure plant nutrition in nutrient-impoverished soils. Although the proposed compensatory effect in AM mutants will need further experimental testing in phosphorus-depleted soils, our hypothesis is consistent with the only mild impairment in plant growth in *ram1-2* mutants (Fig. 2).

We identified three bacterial OTUs, all belonging to the *Burkholderiaceae* family, that are significantly depleted in the roots of each of the three AM mutants compared to wild type. Interestingly, members of the Glomeromycota have been found to contain intracellular endosymbiotic bacteria [46], some belonging to the order Burkholderiales [47]. This finding suggests that these bacteria are either endosymbionts of Glomeromycota fungi that are excluded from the roots of the AM defective genotypes or that their intraradical colonization is indirectly mediated by AM infection. The small changes in the bacterial root microbiota in *ram1-2* plants, which were mainly limited to depletion of the three aforementioned Burkholderia OTUs, revealed that root-associated bacterial community structure is remarkably robust against major changes in the composition of root-associated fungal communities (Fig. 5).

Paleontological and phylogenomic studies established the ancestral origin of genetic signatures enabling AM symbiosis in land plants [1,48]. In monocots and dicots, the extended AM fungal network is primarily recognized as a provider of nutrients, particularly phosphorus [49,50], but the positive impact of AM symbiosis on the host transcends nutrient acquisition [51]. Additionally, phylogenomic studies of the symbiotic phosphate transporter PT4 suggest that this trait evolved late and therefore that phosphorus acquisition might not have been the (only) driving force for the emergence of AM symbiosis [48]. *SymRK* and *Ram1* were identified in the genomes of liverworts, but evolution of *CCaMK* predated the emergence of all land plants, as shown by its presence and conserved biochemical function in advanced charophytes [48]. Together, these findings raise questions regarding the forces driving the evolution of signaling genes enabling intracellular symbioses in land plants. Our study shows that in *L. japonicus*, simultaneous impairment of AM and RN symbioses in *symRK-3* and *ccamk*-13 plants had a dramatic effect on the composition of both bacterial and fungal communities of the legume root microbiota (Fig. 5). Importantly, mutation of *CcamK* and *SymrK* led to an almost complete depletion of a large number of fungal OTUs, mostly belonging to Glomeromycota, indicating that in *Lotus*, these genes predominantly control

the colonization of roots by this particular fungal lineage. The finding that *ram1-2* mutants show
retained accommodation for a subset of fungal root endophytes (n=13; Fig. 5B, and Fig 4B) whose
colonization is dependent on an intact common symbiosis pathway is not surprising based on the
capacity of these mutants to enable fungal colonization but not to sustain a full symbiotic
association [26], and indicates that *RAM1* is dispensable for the intraradical colonization of these
Glomeromycota fungi. Alternatively, these fungal root endophytes may engage in commensal
rather than mutualistic relationships with *L. japonicus* independently of the AM symbiosis pathway
as is the case for multiple species of commensal non-symbiotic rhizobia [22,52]. For instance,
given that *ram1*-2 mutants specifically block AM arbuscule differentiation but not root colonisation
[26], it is conceivable that the Glomeromycota taxa colonizing this plant genotype may not form
arbuscules during root colonization.

Legumes have evolved the capacity to recognize and accommodate both types of intracellular
symbionts, and the large effect of CSSP genes on associated microbiota seen in the present work
could reflect a legume-specific trait. However, in rice, which does not engage in symbiotic
relationships with nodulating rhizobia, mutants lacking *CCaMK* were also found to display
significant changes in root-associated bacterial communities that could be mainly explained by
depletion of Rhizobiales and Sphingomonadales lineages [53]. Thus, our findings based on
comparative microbiota analysis of *Lotus ccamk* and *ram1* mutants suggest a broader role for
common symbiosis signaling genes in microbiota assembly. Future studies on orthologous genes
in basal land plants will contribute to a better understanding of the role of symbiotic signaling in the
evolution of plant-microbiota associations.

Acknowledgements

This work was supported by funds to S. R. from The Danish National Research Foundation (Grant no. DNRF79), by funds to P.S.-L. from the Max Planck Society, a European Research Council advanced grant (ROOTMICROBIOTA), the 'Cluster of Excellence on Plant Sciences' program funded by the Deutsche Forschungsgemeinschaft (DFG), and SPP 2125 DECRyPT from the DFG.

Conflict of interest

The authors declare no conflict of interests.

Data availability

All sequencing reads will be uploaded to the European Nucleotide Archive (ENA). Code and relevant data files (e.g. OTU tables) will be made public via GitHub.

References

1. Remy W, Taylor TN, Hass H, Kerp H. Four hundred-million year-old vesicular arbuscular
mycorrhizae. *Proc Natl Acad Sci USA* 91 1994; 11:841–11.

2. Taylor TN, Remy W, Hass H, Kerp H. Fossil Arbuscular Mycorrhizae from the Early Devonian.
Mycologia 1995; 87(4):560-573.

3. Pirozynski KA, Malloch DW. The origin of land plants: a matter of mycotrophism. *Biosystems*
1975; 6:153-164.

4. Smith SE, Read D J,Vesicular-arbuscular mycorrhizas in agriculture and horticulture. In
Mycorrhizal Symbiosis . Academic press,1997. (pp. 453-469).

5. Wang B, Qiu YL. Phylogenetic distribution and evolution of mycorrhizas in land plants.
Mycorrhiza 2006; 16(5):299-363.

6. Bonfante P, Genre A. Mechanisms underlying beneficial plant-fungus interactions in mycorrhizal
symbiosis. *Nat Commun* 2010; 1:48.

7. Stracke S, Kistner C, Yoshida S, Mulder L, Sato S, Kaneko T, et al. A plant receptor-like kinase
required for both bacterial and fungal symbiosis. *Nature* 2002; 417(6892):959-62

8. Antolin-Llovera M, Ried MK, Parniske M. Cleavage of the SYMBIOSIS RECEPTOR-LIKE
KINASE ectodomain promotes complex formation with Nod factor receptor 5. *Curr Biol* 2014;
24(4):422-427.

9. Charpentier M, Bredemeier R, Wanner G, Takeda N, Schleiff E, Parniske M. Lotus japonicus
CASTOR and POLLUX are ion channels essential for perinuclear calcium spiking in legume root
endosymbiosis. *Plant Cell* 2008; 20(12).

10. Ane JM, Kiss GB, Riely BK, Penmetsa RV, Oldroyd GE, Ayax C. Medicago truncatula DMI1
required for bacterial and fungal symbioses in legumes. *Science* 2004; 303(5662):1364-1367.

11. Kanamori N, Madsen LH, Radutoiu S, Frantescu M, Quistgaard EM, Miwa H. A nucleoporin is
required for induction of Ca2+ spiking in legume nodule development and essential for rhizobial
and fungal symbiosis. *Proc Natl Acad Sci* USA 2006; 103(2):59-364

12. Saito K, Yoshikawa M, Yano K, Miwa H, Uchida H, Asamizu E. NUCLEOPORIN85 is
required for calcium spiking, fungal and bacterial symbioses, and seed production in Lotus
japonicus. *Plant Cell* 2007; 19(2):610-624

13. Groth M, Takeda N, Perry J, Uchida H, Draxl S, Brachmann A. NENA, a Lotus japonicus
homolog of Sec13, is required for rhizodermal infection by arbuscular mycorrhiza fungi and
rhizobia but dispensable for cortical endosymbiotic development. *Plant Cell* 2010; 22(7):2509-
2526.

14. Charpentier M, Sun J, Vaz Martins T, Radhakrishnan GV, Findlay K, Soumpourou E. Nuclear-
localized cyclic nucleotide-gated channels mediate symbiotic calcium oscillations. Science 2016;
352(6289):1102-1105.

15. Levy J, Bres C, Geurts R, Chalhoub B, Kulikova O, Duc G. A putative Ca2+ and calmodulin-
dependent protein kinase required for bacterial and fungal symbioses. Science 2004;
303(5662):1361-1364.

16. Mitra RM, Gleason CA, Edwards A, Hadfield J, Downie JA, Oldroyd GE, et al. A
Ca2+/calmodulin-dependent protein kinase required for symbiotic nodule development: Gene
identification by transcript-based cloning. *Proc Natl Acad Sci U S A* 2004; 101(13):4701-5.

17. Yano K, Yoshida S, Muller J, Singh S, Banba M, Vickers K. et al. CYCLOPS, a mediator of
symbiotic intracellular accommodation. *Proc Natl Acad Sci U S A* 2008; 105(51):20540-20545.

18. Singh S, Katzer K, Lambert J, Cerri M, Parniske M. CYCLOPS, a DNA-binding transcriptional
activator, orchestrates symbiotic root nodule development. *Cell Host Microbe* 2014; 15(2):139-152.

19. Messinese E, Mun JH, Yeun LH, Jayaraman, D, Rouge P, Barre, A, et al. A novel nuclear
protein interacts with the symbiotic DMI3 calcium- and calmodulin-dependent protein kinase of
Medicago truncatula.. *Mol Plant Microbe Interact* 2007; 20(8):912-921.

20. Udvardi M, Poole PS. Transport and metabolism in legume-rhizobia symbioses. *Annu Rev Plant*
Biol 2013; 64:781-805.

21. Gutjahr C, Sawers RJH, Marti G, Andres-Hernandez L, Yang S, Casieri L. Transcriptome
diversity among rice root types during asymbiosis and interaction with arbuscular mycorrhizal
fungi. *Proc Natl Acad Sci* USA 2015; 112(21):6754.

22. Zgadzaj R, Garrido-Oter R, Jensen DB, Koprivova A, Schulze-Lefert P, Radutoiu S. Root
nodule symbiosis in Lotus japonicus drives the establishment of distinctive rhizosphere, root, and
nodule bacterial communities. *Proc Natl Acad Sci U S A* 2016; 113(49):E7996-E8005.

23. Johnson D, Gilbert L. Interplant signalling through hyphal networks. New Phytol 2014;
205(4):1448-53.

24. Hassani MA, Duran, P, Hacquard S. Microbial interactions within the plant holobiont.
Microbiome 2018; 6(1):58.

25. Madsen EB, Madsen, LH, Radutoiu S, Olbryt M, Rakwalska M, Szczyglowski K, et al. A
receptor kinase gene of the LysM type is involved in legumeperception of rhizobial signals. *Nature*
2003; 425(6958):637.

26. Xue L, Cui H, Buer B, Vijayakumar V, Delaux PM, Junkermann S, et al. Network of GRAS
transcription factors involved in the control of arbuscule development in Lotus japonicus. *Plant*
Physiol 2015; 167(3):854-871.

27. Perry J, Brachmann A, Welham T, Binder A, Charpentier M, Groth M, et al. TILLING in Lotus
japonicus identified large allelic series for symbiosis genes and revealed a bias in functionally
defective ethyl methanesulfonate alleles toward glycine replacements. *Plant Physiol* 2009; 151:
1281–1291.

28. Schlaeppi K, Dombrowski N, Oter RG, Ver Loren van Themaat E, Schulze-Lefert P.
Quantitative divergence of the bacterial root microbiota in Arabidopsis thaliana relatives. *Proc Natl*
Acad Sci U S A 2014; 111(2):585-92.

29. Ihrmark K, Bödeker IT, Cruz-Martinez K, Friberg H, Kubartova A, Schenck J,et al. New
primers to amplify the fungal ITS2 region--evaluation by 454-sequencing of artificial and natural
communities. *FEMS Microbiol Ecol* 2012; 82(3):666-77.

30. White TJ, Bruns TD, Lee SB, Taylor JW. Amplification and direct sequencing of fungal
ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, editors.
PCR protocols: a guide to methods and applications. United States: Academic Press, CA, USA,
1990, pp. 315-322.

31. Duran, P., Thiergart T, Garrido-Oter R, Agler M, Kemen E, Schulze-Lefert P, et al. (2018).
Microbial interkingdom interactions in roots promote Arabidopsis survival. *Cell* 2018; 175(4):973-
983.

32. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al. QIIME
allows analysis of high-throughput community sequencing data. *Nat Methods* 2010; 7:335-336.

33. Edgar RC. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat*
Methods 2013; 10:996-998.

34. DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, et al. Greengenes, a
chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl Environ*
Microbiol 2006; 72:5069-5072

35. Bengtsson-Palme J, Ryberg M, Hartmann M, Branco S, Wang Z, Godhe A, De Wit P, et al.
Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences of
fungi and other eukaryotes for analysis of environmental sequencing data. *Methods Ecol Evol* 2013;
4: 914-919.

36. Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid assignment of
rRNA sequences into the new bacterial taxonomy. *Appl Environ Microb* 2007; 73:5261-5267.

37. Paulson, JN, Stine, OC, Bravo HC, Pop M. Differential abundance analysis for microbial
marker-gene surveys. *Nat Methods* 2013; 10:1200-1202.

38. Chabaud M, Genre A, Sieberer BJ, Faccio A, Fournier J, Novero M, et al. Arbuscular
mycorrhizal hyphopodia and germinated spore exudates trigger Ca2+ spiking in the legume and
nonlegume root epidermis. *New Phytol* 2011; 189(1):347-5.

39. Miwa H, Sun J, Oldroyd GE, Downie JA. Analysis of calcium spiking using a cameleon
calcium sensor reveals that nodulation gene expression is regulated by calcium spike number and
the developmental status of the cell. *Plant J* 2006; 48(6):883-94.

40. Bulgarelli D, Rott M, Schlaeppi K, Emiel Ver Loren van Themaat E, Ahmadinejad N, Assenza
F, et al. Revealing structure and assembly cues for Arabidopsis root-inhabiting bacterial microbiota.
Nature 2012; 488:91-95.

41. Anderson MJ, Willis TJ, Canonical analysis of principal coordinates: A useful method of
constrained ordination for ecology. *Ecology* 2003; 84(2):511-525.

42. Lombard L, van der Merwe NA, Groenewald JZ, Crous PW. Generic concepts in Nectriaceae.
Stud Mycol 2015; 80:189-245.

43. Lofgren LA, LeBlanc NR, Certano AK, Nachtigall J, LaBine KM, Riddle J, et al. Fusarium
graminearum: pathogen or endophyte of North American grasses? *New Phytol* 2018; 217(3):1203-
1212.

44. Amselem J, Cuomo CA, van Kan, JA, Viaud M, Benito EP, Couloux A, et al. Genomic analysis
of the necrotrophic fungal pathogens Sclerotinia sclerotiorum and Botrytis cinerea. *PLoS Genet*
2011; 7(8).

45. Almario J, Jeena G, Wunder J, Langen G, Zuccaro A, Coupland G, Bucher M. Root-associated
fungal microbiota of nonmycorrhizal Arabis alpina and its contribution to plant phosphorus
nutrition. *Proc Natl Acad Sci U S A* 2017; 114(44):E9403-E9412.

46. Desiro A, Salvioli A, Ngonkeu EL, Mondo SJ, Epis S, Faccio A, et al. Detection of a novel
intracellular microbiome hosted in arbuscular mycorrhizal fungi. *ISME J* 2014; 8(2): 257-270.

47. Bianciotto V, Lumini E, Lanfranco L, Minerdi D, Bonfante P, Perotto S. Detection and
identification of bacterial endosymbionts in arbuscular mycorrhizal fungi belonging to the family
Gigasporaceae. *Appl Environ Microbiol* 2000;66(10):4503-4509.

48. Delaux PM, Radhakrishnan GV, Jayaraman D, Cheema J, Malbreil M, Volkening JD,et al.
Algal ancestor of land plants was preadapted for symbiosis. *Proc Natl Acad Sci U S A* 2015;
112(43):13390-13395.

49. Javot H, Penmetsa RV, Terzaghi N, Cook DR, Harrison MJ. A Medicago truncatula phosphate
transporter indispensable for the arbuscular mycorrhizal symbiosis. *Proc Natl Acad Sci U S A* 2007;
104(5):1720-5.
553
554 50. Sawers RJ, Svane SF, Quan C, Grønlund M, Wozniak B, Gebreselassie MN, et al. Phosphorus
555 acquisition efficiency in arbuscular mycorrhizal maize is correlated with the abundance of root-
556 external hyphae and the accumulation of transcripts encoding PHT1 phosphate transporters. *New*
557 *Phytol* 2017; 214(2):632-643.
558
559 51. Chen M, Arato M, Borghi L, Nouri E, Reinhardt D. Beneficial Services of Arbuscular
560 Mycorrhizal Fungi - From Ecology to Application. *Front Plant Sci* 2018; 9:1270.

52. Garrido-Oter R, Nakano RT, Dombrowski N, Ma KW, AgBiome T, McHardy AC, et al. Modular Traits of the Rhizobiales Root Microbiota and Their Evolutionary Relationship with Symbiotic Rhizobia. *Cell Host Microbe* 2018; 24(1):155-167 e155.

53. Ikeda S, Okubo T, Takeda N, Banba M, Sasaki K, Imaizumi-Anraku H, et al. The genotype of the calcium/calmodulin-dependent protein kinase gene (CCaMK) determines bacterial community diversity in rice roots under paddy and upland field conditions. *Appl Environ Microbiol* 2011; 77(13):4399-4405.

Main Figures

Fig1, Bacterial & fungal community profile for different root fractions of *L. japonicus*.
A) Cartoon showing the length of the three different root fractions. B) Community profile showing the relative abundance of bacterial (upper panel) and fungal (lower panel) families across compartments and fractions (only samples with >5000 (bacteria) or >1000 (fungi) reads are shown, taxa having average RA < 0.1 (bacteria) or <0.15 (fungi) across all samples are aggregated as low-abundant.). C) Ternary plots showing bacterial OTUs that are enriched in the endosphere of specific root fractions, compared to the soil samples. B) Ternary plots showing fungal OTUs that are enriched in the endosphere of specific root fractions, compared to the soil samples. Circle size corresponds to RA across all fractions. Dark grey circles denote OTUs that are enriched in soil, light grey circles always represent OTUs that are not enriched in any of the fractions.

Fig2, **Phenotypes of WT and mutant plants**. Images depicting *L. japonicus* wild-type (A) and
symbiosis-deficient mutant plants: *ram1-2* (B), *nfr5-2* (C), *symRK-3* (D) and *ccamk-13* (E). Insets
show close-up view of nodules. Scale bars correspond to 1 cm. F) Boxplots display the shoot length
for the identical set of genotypes presented in (A-E). G) Boxplots displaying the shoot fresh mass.
Letters above plots correspond to groups based on Tukey's HSD test (P<0.05). Number of samples
are indicated in brackets.

Fig3, **Constrained PCoA analysis showing genotype effect on microbial communities.**
A) Constrained PCoA plots for bacterial datasets showing rhizosphere samples (n = 100) and B)
root samples (n = 100). C) Constrained PCoA plots for fungal datasets showing only rhizosphere
samples (n = 124) and D) root samples (n = 122).

651
652
Fig4, **Relative abundance for main microbial taxa across plant compartments and genotypes**.
A) RA for bacterial families in rhizosphere (left panel) and root compartment (right panel). B) RA
for fungal families in rhizosphere (left panel) and root compartment (right panel). Taxa are sorted in
decreasing order according to their average RA in wt plants (only first 10 most abundant
taxonomical groups are shown). RA in wt as well as in the respective mutants is displayed.
Significant differences compared to wt are marked with an asterisk in the color of the mutant
(P<0.05, Kruskal-Wallis test). Families that include known symbionts are marked in red
(Phyllobacteriaceae for bacteria and Glomeromyctes for Fungi). For some fungal taxa the next
higher rank is shown, when no family level information was available.

Fig5, **Differential abundance analysis for root associated OTUs**.

A) Bacterial OTUs that are differentially abundant in the roots of mutants compared to wt roots. B) fungal OTUs that are differentially abundant in the roots of mutants compared to wt roots. Only OTUs that have an average RA > 0.1% across all root samples, including mutants, are considered here. For each OTU the fold change in RA from wt to mutant is indicated (P <0.05, Kruskal-Wallis test). Next to each OTU the RA in wt roots is indicated. Phylum and family asscociation (if available) is given for each OTU (Bacterial phyla: Del=Deltaproteobactria, Gem=Gemm-1 , Chl=Chloroflexi, Bet=Betaproteobacteria, Alp=Alphaproteobacteria, Gam=Gammaproteobacteria, Cyt=Cytophagia, Sap=Saprospirae, Ped=Pedosphaerae, Sph= Sphingobacteria, Mol= Mollicutes ; Fungal phyla: Sor=Sordariomyctes, Dot=Dothideomyctes, Mic= Microbotryomyctes, Ust=Ustilaginomyctes, Eur=eurotiomyctes, Leo=Leotiomyctes, Aga=Agaricomyctes, Glo=Glomeromyctes, Pez=Pezizomycota, Muc=Mucoromycotina).

Supplementary Figures

687
688
Supp Fig1, Alpha and beta diversity across root fractions.
A) Shannon diversity index for 16S amplicon data, soil (n=6), lower (n=6), upper (n=6), whole root
fractions (n=6), and respective rhizosphere samples (n=6 each) B) Shannon diversity index for
ITS2 amplicon data, soil (n=6), lower (n=2), upper (n=6), whole root fractions (n=6), and
respective rhizosphere samples (n=6 each, except lower, n=4) (ANOVA with Tukey's post hoc
test, P<0.05). C) Principal coordinate analysis of Bray-Curtis distances for bacterial data. D)
Principal coordinate analysis of Bray-Curtis distances for fungal data.

696
697

698
699
700
701
702 **Supp Fig2, Alpha and beta diversity across plant compartments and genotypes.**
703 A) Shannon diversity indices for the bacterial (16S amplicon) dataset. B) Shannon diversity indices
704 for the fungal (ITS2 amplicon) dataset (ANOVA with Tukey's post hoc test, P<0.05) C) Principal
705 coordinate analysis of Bray-Curtis distances for the bacterial dataset (n=222). D) Principal
706 coordinate analysis of Bray-Curtis distances for the fungal dataset (n=274).

724
725
Supp Fig3, CPCoA, showing the same results as Fig3, but without known symbionts. Results
are separated by compartments. Datasets were constrained by genotype, and filtered for effects of
experiments and soil type. A) Bacterial dataset from which OTUs belonging to the
Phyllobacteriaceae were removed before analysis (root n=100, rhizosphere n=100). B) Fungal
dataset from which OTUs belonging to the Glomeromycota were removed before analysis (root
n=122, rhizosphere n=124).

744
745
Supp Fig4, Differential abundance analysis for rhizosphere-associated OTUs showing
enrichment and depletion in mutants.
A) bacterial OTUs that are differentially abundant in the rhizosphere of mutants compared to wt.
B) fungal OTUs that are differentially abundant in the rhizosphere of mutants compared to wt. Only
OTUs that have an average RA > 0.1% across all rhizosphere samples, including mutants, are
considered here. For each OTU, the fold change in RA from wt to mutant is indicated (P <0.05,
Kruskal-Wallis test). Next to each OTU the RA in wt rhizosphere is indicated. Phylum and family
asscociation is given for each OTU (Bacterial phyla: Del=Deltaproteobacteria, Gem=Gemm-1 ,
Chl=Chloroflexi, Bet=Betaproteobacteria, Alp=Alphaproteobacteria, Gam=Gammaproteobacteria,
Cyt=Cytophagia, Sap=Saprospirae, Ped=Pedosphaerae, Sph= Sphingobacteria, Mol= Mollicutes ;
Fungal phyla: Sor=Sordariomycetes, Dot=Dothideomycetes, Mic= Microbotryomycetes,
Ust=Ustilaginomycetes, Eur=Eurotiomycetes, Leo=Leotiomycetes, Aga=Agaricomycetes,
Glo=Glomeromycetes, Pez=Pezizomycota, Muc=Mucoromycotina).
759
760

761
762
763 **Supp fig5, Ternary plots showing compartment-enriched bacterial OTUs.** Separately for wt
764 and mutant plants. Below each plot the number of enriched OTUs for each compartment is shown.

Supp table1, Symbiotic phenotype of *Lotus japonicus* wild-type and mutants grown in Cologne soil (*n=5*)

Genotype	WT(Gifu)	*nfr5-2*	*ram1-2*	*symrk-2*	*ccamk-13*
Colonization (% of root system with intraradical colonisation)	60-70	60-70	10	0	0
Arbuscules	present	present	present	none	none
Nodulation (% of roots with nodules)	100	0	100	0	0