1   *Vitellogenin* **expression in the ovaries of adult honeybee workers provides insights into**
2   **the evolution of reproductive and social traits**
3
4

5   Carlos Antônio Mendes Cardoso-Júnior[1,2*], Benjamin P. Oldroyd[2], Isobel Ronai[2]
6
7   [1] Departamento de Biologia Celular e Bioagentes Patogênicos, Faculdade de Medicina de
8   Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, Brasil.
    [2] Behaviour and Genetics of Social Insects Laboratory, Ecology and Evolution, School of
    Life and Environmental Sciences A12, University of Sydney, Sydney NSW 2006, Australia.
9
10
11
[*] **Corresponding author**: carloscardoso.bio@usp.br

Contact Information

Carlos A. M. Cardoso-Júnior – carloscardoso.bio@usp.br; Phone: +55 16 3315 3063

Benjamin P. Oldroyd – benjamin.oldroyd@sydney.edu.au

Isobel Ronai – isobel.ronai@sydney.edu.au

Running title: *Vitellogenin* in the ovaries of honeybee workers

Abstract

Social insects are notable for having two female castes that exhibit extreme differences in their reproductive capacity. The molecular basis of these differences is largely unknown. A protein that likely plays a key role in these differences is Vitellogenin (Vg), a powerful antioxidant and insulin-signalling regulator. Here we investigate how Royal Jelly (the major food of honeybee queens) and queen pheromone (a major regulator of worker fertility), affects the longevity and reproductive status of honey bee workers, the expression of *Vg*, its receptor *VgR* and associated regulatory proteins. We find that *Vg* is expressed in the ovaries of workers and that workers fed a queen diet of Royal Jelly have increased *Vg* expression in the ovaries. Surprisingly, we find that *Vg* expression is not associated with worker ovary activation. Our findings provide further support for the 'reproductive ground plan hypothesis' as Vg has acquired non-reproductive functions in honeybee workers.

Keywords: *Apis mellifera*, epigenetic, honey bee, Juvenile hormone, *Vitellogenin receptor* (*VgR*), *Krüppel-homolog 1* (*Kr-h1*), *DNA methyltransferase 3* (*Dnmt3*) and the *Forkhead box O transcription factor* (*FoxO*)

Introduction

Eusocial insects provide an interesting exception to the reproduction-longevity trade-off (Blacher *et al.*, 2017; Heinze and Schrempf, 2008), for queens are highly fecund yet have extraordinarily long life-spans, whereas workers are typically non-reproductive and have a short lifespan (Winston, 1987). Understanding of the molecular basis and evolutionary origins of the phenotypic divergence between queen and worker castes, despite their development from identical genomes, are major outstanding questions in sociobiology (Ronai *et al.*, 2016b).

A likely mechanism by which queens sustain high rates of egg production over their long lifespan is by life-long production of the egg-yolk precursor protein Vitellogenin (Vg) (Corona *et al.*, 2007). The Vg protein is synthesised in the fat body and ovaries of most insects (Hagedorn and Kunkel, 1979), and is taken up by the developing oocyte (Raikhel and Dhadialla, 1992). In eusocial insects Vg seems to have acquired additional functions (Corona *et al.*, 2013; Nelson *et al.*, 2007). Vg acts as an anti-oxidant that helps prevent senescence in queens and workers (Corona *et al.*, 2013). In addition, Vg interacts with components of the insulin signalling pathway (Corona *et al.*, 2007), which is increasingly recognised as an important regulator of lifespan (van Heemst, 2010) and behavioural maturation (Amdam *et al.*, 2004; Nelson *et al.*, 2007).

A significant difference between queens and workers is their diet, which likely contributes to their differential longevity. In honeybees, for example, adult queens are fed glandular secretions that contain Royal Jelly (a nutritious food that is rich in carbohydrates, lipids and protein) (Rembold and Dietz, 1966). In contrast, adult workers feed directly on stored honey and pollen, or on regurgitated crop contents provided by other workers (Winston, 1987). Interestingly, if workers are forced to consume small amounts of Royal Jelly in an artificial diet, they develop a more queen-like phenotype, including ovary activation and increased lifespan (Lin and Winston, 1998; Wang *et al.*, 2014; Yang *et al.*, 2017). The effects of Royal Jelly seem to be associated with the gene networks that regulate reproduction and longevity (Ford, 2013).

Another environmental factor that influences adult honeybee worker reproduction and potentially longevity is their social environment. Worker fertility is influenced by a number of pheromones released by the brood and the queen, one of which is queen mandibular pheromone (QMP). This pheromone signals the queen's presence to workers and supresses activation of worker ovaries (Hoover *et al.*, 2003; Ronai *et al.*, 2016a). Furthermore, QMP regulates the expression of reproductive genes, including *Vg* (Fischer and Grozinger, 2008). When workers are exposed to QMP, *Vg* expression is upregulated (Fischer and Grozinger, 2008). High levels of Vg leads to decreased levels of juvenile hormone (JH) in the haemolymph (Guidugli *et al.*, 2005a), which delays behavioural maturation and the initiation of foraging (Marco Antonio *et al.*, 2008). This suggests that there is an interplay between the levels of JH and Vg within workers, which in turn influence life expectancy (Amdam and Omholt, 2003; Flatt *et al.*, 2013).

The Ovarian Ground Plan Hypothesis argues that the gene networks that were once involved in the regulation of reproduction in the ancestors of social insects have been co-opted to facilitate the regulation of behavioural differences between queen and worker castes (Hunt and Amdam, 2005; West-Eberhard, n.d.). *Vg* appears to have been co-opted in this way (Roy-Zokan *et al.*, 2015). While in honeybee queens Vg acts as a precursor for the synthesis of egg-yolk, in workers it acts as a precursor for the synthesis of Royal Jelly proteins (Amdam *et al.*, 2003) suggesting that the protein's function has changed from direct maternal investment in eggs to alloparental care (the production of larval food for sisters). More recently the Ovarian Ground Plan Hypothesis has been extended to the Reproductive Ground

Plan Hypothesis which argues that genes that once controlled reproduction now influence
behavioural phenotypes between worker sub-castes (Amdam *et al.*, 2006). Again, Vg appears
to have assumed functions that are only indirectly related to reproduction, most notably
behavioural maturation (Nelson *et al.*, 2007).
Here we determine whether ovary state (activated or non-activated) and expression of
Vg in the ovaries of adult honeybee workers are affected by environmental factors. We
investigate the interaction effect of two environmental factors that have opposite effects on
ovary activation: diet (the presence or absence of Royal Jelly (Hoover *et al.*, 2006)) and
social environment (the presence or absence of QMP (Hoover *et al.*, 2003)). We then
investigate how these environmental factors affect the expression of *Vg*'s receptor (*VgR*)
(Dohanik *et al.*, 2018; Guidugli-Lazzarini *et al.*, 2008) and three putative regulators of *Vg*. *A*
priori, *Vg* expression in the ovaries of workers is likely to be influenced by these two
environmental factors because its expression is associated with social functions in addition to
its canonical reproductive function.
141
Material and methods
143
Biological material, cage setup and longevity
We obtained age-matched worker honeybees of standard Australian commercial stock
(primarily *A. m. ligustica*) by incubating frames of emerging brood from two source colonies
overnight at 34.5 °C. Workers from each source colony were randomly allocated to four
cages (Rothenbuhler *et al.*, 1979) (n = 150 per cage, total of 8 cages) in a three-factor (source
colony plus the two treatments) factorial design. Four of the cages contained half a synthetic
queen pheromone strip (Bee Boost Pherotech), which delivered approximately 0.5 queen
equivalents per day (Pankiw and Page, 2003; Ronai *et al.*, 2015), while the other four cages
contained no strip (QMP$^-$).
Pollen and water were provided *ad libitum*. In addition, four of the cages (2 QMP$^+$
and 2 QMP$^-$) were fed a Royal Jelly diet of 50% honey:50% Royal Jelly (Royal Jelly
Australia, stored frozen), whereas the other four cages (2 QMP$^+$ and 2 QMP$^-$) were only fed a
Control diet of pure honey. Food was replenished when necessary, and the number of dead
workers recorded each day (supplementary table 1). Cages were housed in an incubator at
34.5°C in the dark. At seven and fourteen days of age workers were collected and
immediately frozen.
In a parallel experiment to assess longevity, two additional source colonies were
prepared as above (four cages per source colony). Cages were kept under the same
conditions. However, the workers were not collected and instead the number of dead workers
was recorded daily until all workers had died.
164
Assessment of ovary activation
We dissected and imaged the ovaries from workers under a dissecting microscope
(Leica M60). The paired ovaries of the 7-day-old workers were removed and snap frozen.
During dissection the activation state of the ovaries was recorded on a three-point scale: 1 -
deactivated (thin ovarioles, no visible oocytes and Vg not required by the oocytes), 2 - semi-
activated (ridged ovarioles, with oocytes at the vitellarium stage of development and Vg not
yet required for oocyte development) and 3 - activated (mature, white oocytes and Vg
required for normal oocyte development) (Ronai *et al.*, 2016a; Seehuus *et al.*, 2007). Our
scale is based on the reproductive control points that occur during early and mid-oogenesis in
the honeybee worker (Ronai *et al.*, 2016b, 2015).
175
Expression analysis

The ovaries of the 7-day-old workers were macerated in TRIzol (Invitrogen) and total
RNA extracted with the Direct-zol^{TM} RNA Miniprep kit (Zymo Research) according to the
manufacturer's instructions. Each sample consisted of four pairs of non-activated ovaries
(deactivated or semi-activated ovaries), these were pooled to obtain a sufficient amount of
RNA. Activated ovaries were not combined as there were a limited number and individual
ovaries yielded sufficient RNA. The samples were digested with the Turbo DNase Kit
(Thermo Fisher Scientific) to remove DNA. RNA concentration was then quantified using a
Qubit 2.0 Fluorometer system (Invitrogen). Each RNA sample was diluted to a final
concentration of 15 ng/µL with ultrapure water. We used 142.5 ng of each sample to
synthesise cDNA using the SuperScript^{TM} III Reverse Transcriptase Kit (Invitrogen) with
Oligo (dT) primer. The cDNA was diluted to 2 ng/µL in ultrapure water.
To detect the expression of *Vg* and its receptor *VgR* in the ovaries of workers (see
supplementary table 2 for primer sequences and sources) we first performed conventional
PCR. To semi-independently validate *Vg* expression we utilised two sets of *Vg* primers, *Vg*P1
(Lourenco *et al.*, 2012) and *Vg*P2 (Koywiwattrakul *et al.*, 2005), which target different exons
within *Vg* (*Vg*P1: exon 3; *Vg*P2: spans exon 5 and 6). PCR reactions were carried out on three
different cDNA samples of Royal Jelly fed workers. The cDNA (2 ng) was mixed with 0.2
µL TaqTi enzyme (Fisher Biotec Australia), 3.2 nmol dNTPs, 1.25 pmol of each primer,
buffer 1X, 50 nmol MgCl$_2$ in a final volume of 20 µL. The following genes were used as
positive controls: ribosomal protein 49 (*RP49*, also known as *rpl32*) and elongation factor 1
alpha (*Ef1α*). The negative control gene was yellow fluorescent protein (*YFP*), which is not
present in honeybees. Primer sequences and sources are listed in supplementary table 2. The
amplification cycles were: 95 °C for 2 min; 40 cycles of 95°C for 15 sec; 60 °C for 15 sec;
and 72 °C for 30 sec, 72°C for 5 min. RT-PCR products were ran on a 1% agarose gel and
imaged using a Bio-Rad ChemiDoc Touch Imaging System.
We quantified the expression of *Vg* and *VgR* in the ovaries of adult workers with RT-
qPCR. In addition, we quantified the expression of three genes that are potential regulators of
Vg expression in the honeybee: *Krüppel-homolog 1* (*Kr-h1*) (Amdam and Omholt, 2002;
Grozinger *et al.*, 2003; Jindra *et al.*, 2015), *DNA methyltransferase 3* (*Dnmt3*) (Cardoso-
Júnior *et al.*, 2018) and *Forkhead box O transcription factor* (*FoxO*) (Corona *et al.*, 2007;
Hansen *et al.*, 2007; Sheng *et al.*, 2011; Süren-Castillo *et al.*, 2012). RT-qPCR reactions
consisted of 2.5 µL SsoAdvanced™ Universal SYBR® Green Supermix (Bio Rad), 1.25
pmol of each primer, 1 µL diluted cDNA (2 ng) in a total volume of 5 µL, with three
technical replicates per biological sample. The reactions were performed using a CFX384
Real-Time System (Bio-Rad) with the cycle conditions: 95 °C for 10 min; 40 cycles of 95°C
for 10 sec; 60 °C for 10 sec; and 72 °C for 15 sec. For each gene the melt curve was checked
to confirm a single amplification peak. The primer efficiency (supplementary table 2) was
calculated based on an amplification curve of 10 points obtained through serial dilution of
mixed cDNA samples. To normalise the gene expression we used two reference genes (*RP49*
and *Ef1α* (Lourenço *et al.*, 2008)) whose expression was stable according to *BestKeeper*
software (Pfaffl *et al.*, 2004). The relative expression was calculated as in (Brito *et al.*, 2010)
using a relative gene expression formula that takes into account the efficiency of each primer.
219
Statistical analysis
Ovary activation score was calculated, per sample, as the average of the activation
scores of all ovaries included in the sample. Average ovary activation score and gene
expression level were analysed as dependent variables using generalised linear mixed models
(GLMM), with 'colony' as random effect, and diet and QMP treatments as fixed effects. We
used a log function, link = log and family = Gaussian, to model *Vg*, *VgR* and *Kr-h1*

expression irrespective of ovary state. *Dnmt3* and *FoxO* expression and ovary activation
score were modelled without applying a log function. Analyses were performed in *R* (Team,
2018) using the packages lme4, car and lsmeans or GraphPad Prism 7. To compare *Vg* gene
expression in activated and non-activated ovaries, we used a two-tailed Student's *t*-test after
$\log^{10}$ transformation of expression data. Longevity was assessed with survival curves that
were statistically compared using Mantel-Cox test. Two-tailed Spearman correlation test was
used to assess the correlation between *Vg* and *VgR*, *Kr-h1*, *Dnmt3* and *FoxO* expression. A *p*-
value lower than 0.05 was considered significant for all statistical tests. When appropriate,
significance levels for *post-hoc* tests were Bonferroni adjusted for multiple comparisons.

Results

High ovary activation and short lifespan in workers fed Royal Jelly
Ovary activation score was significantly higher in workers fed the Royal Jelly diet
compared to workers fed the Control diet ($p < 0.001$; figure 1a and supplementary table 3). In
addition, the ovary activation score was significantly higher in workers not exposed to QMP
compared to workers exposed (figure 1a, $p < 0.05$, see supplementary table 3 for further
details). There was no significant interaction between the diet and QMP treatments ($p = 0.14$;
figure 1a). For the majority of the treatment combinations there was a significant difference
in ovary activation score when comparing both QMP treatment and diet (comparison of least
square means, $p < 0.001$ after Bonferroni correction). The two exceptions were: (i) QMP$^+$ and
QMP$^-$ of the Control diet ($p = 0.058$) and (ii) QMP$^-$ of the Control diet and QMP$^+$ of Royal
Jelly diet ($p = 0.26$). At 14 days of age, workers with activated ovaries comprised 0% of
QMP$^+$/Control diet, 23% QMP$^-$/Control diet, 40% of QMP$^+$/Royal Jelly diet and 63% QMP$^-$
/Royal Jelly diet (supplementary figure 1).
The Royal Jelly diet reduced the lifespan of workers in both colonies ($p < 0.0001$
Mantel-Cox test followed by Bonferroni correction for multiple comparisons, figure 2).
Differences between workers exposed or not to QMP was only found in bees fed Royal Jelly
from source colony A ($p < 0.0001$ Mantel-Cox test followed by Bonferroni correction for
multiple comparisons, figure 2a).

Vg expression in the ovaries of workers is affected by their diet and social environment
Unlike previous studies (Corona *et al.*, 2007; Guidugli *et al.*, 2005b) we find that *Vg*
is expressed in the ovaries of adult honeybee workers (figure 3a and figure 3b). *Vg*
expression in the ovaries of adult workers is affected by both QMP ($p < 0.05$; figure 3b and
supplementary table 4) and diet ($p < 0.001$; figure 3b and supplementary table 4) with no
significant interaction between the two main effects ($p = 0.8$). In the presence of both QMP
and the Royal Jelly diet, workers had significantly higher expression of *Vg* in their ovaries
compared to the other three treatment groups: QMP$^+$/Control diet (comparison of least square
mean, $p < 0.01$ after Bonferroni correction); QMP$^-$/Control diet ($p < 0.001$) and QMP$^-$/Royal
Jelly diet ($p < 0.01$). In addition, Vg expression was significantly higher in QMP$^-$/Royal Jelly
diet workers compared to workers from QMP$^-$/Control diet treatment ($p < 0.01$). There was
no significant difference in the expression of *Vg* between the two primer sets ($p = 0.76$, *Vg*P1:
figure 3b and *Vg*P2: supplementary figure 2).
VgR expression is affected by both QMP ($p < 0.05$; figure 3c and supplementary table
4) and diet ($p < 0.001$; figure 3c and supplementary table 4) with no significant interaction
between the two main effects ($p = 0.57$). *VgR* expression was significantly higher in the
treatment combination of Royal Jelly diet with no QMP compared to all other treatment
combinations ($p < 0.01$, figure 3c). In addition, *VgR* expression was significantly higher in

the treatment with Royal Jelly diet and QMP$^+$ compared with the Control diet and QMP$^+$ ($p <$
0.001).
277
Vg expression is lower in activated ovaries whereas VgR is higher
We investigated the effect of ovary activation on the expression of *Vg* and *VgR* within the
QMP$^-$/Royal Jelly treatment group to avoid the confounding factors of diet and social
environment. *Vg* expression was significantly lower in activated ovaries compared to non-
activated ovaries ($p = 0.004$; figure 3b inset, supplementary figure 2 inset and supplementary
table 4). In contrast, expression of *VgR* was significantly higher in activated ovaries
compared to non-activated ovaries ($p = 0.0033$; figure 3c inset).
285
Royal jelly alters the expression of putative Vg regulators
Vg expression in the ovaries of workers is not significantly correlated with the
expression level of its receptor, *VgR* ($r^2 = 0.16$, $p = 0.15$, supplementary figure 3). To gain
insights into the regulatory control of *Vg* transcription in the ovaries of workers, we examined
the expression of three additional potential *Vg* regulators. Firstly, the expression of *Kr-h1* was
affected by both QMP exposure ($p < 0.0001$, figure 4a and supplementary table 5) and Royal
Jelly diet ($p < 0.0001$, figure 4a and supplementary table 5). We found no interaction between
the two main effects ($p = 0.16$). Interestingly, the QMP effect was suppressed in the workers
fed Royal Jelly ($p = 0.32$). Secondly, the expression of *Dnmt3* is influenced by QMP
presence ($p < 0.0001$; figure 4b and supplementary table 5). We found an interaction between
QMP and diet ($p < 0.05$). *Dnmt3* is upregulated in the QMP$^+$/Control diet in relation to all
other groups (comparison of least square mean, $p < 0.01$ after Bonferroni correction). Similar
to *Kr-h1*, Royal Jelly suppressed the effects of QMP exposure on *Dnmt3* expression. Thirdly,
the expression of *FoxO* was only affected by diet ($p < 0.0001$, figure 4c) with no significant
interaction between diet and QMP ($p = 0.08$). *FoxO* expression was higher in QMP$^-$/Control
diet workers compared to those fed Royal Jelly (comparison of least square mean $p < 0.05$
after Bonferroni correction). In addition, *FoxO* expression was lower in QMP$^-$/Royal Jelly
diet treatment compared to QMP$^+$/Control diet ($p < 0.0001$ after Bonferroni correction).
We also investigated whether *Vg* expression in the ovaries of workers is correlated
with the expression of its potential regulators (supplementary figure 3). *Vg* expression was
negatively correlated with the expression of *Kr-h1* ($r^2 = -0.28$, $p = 0.01$, supplementary figure
3) but not correlated with expression of *Dnmt3* ($r^2 = 0.12$, $p = 0.28$, supplementary figure 3)
nor expression of *FoxO* ($r^2 = 0.1$, $p = 0.36$, supplementary figure 3).
309
310   **Discussion**
311
Our study demonstrates that *Vg*, a gene associated with longevity and behavioural
maturation, is expressed in the ovaries of adult honeybee workers. Previously, *Vg* expression
was only known in the ovaries of queens, not workers (Corona *et al.*, 2007; Guidugli *et al.*,
2005b). Honeybee workers are therefore similar to many other social and non-social insects
that also express *Vg* in tissues other than the fat body (Jedlicka *et al.*, 2016; Valle, 1993). We
also find that the environment of a worker (diet and exposure to QMP) influences the
expression of *Vg* in its ovaries. While it is known that environmental effects influence *Vg*
expression in the fat body of honeybees and other insects (Amsalem *et al.*, 2014; Roy-Zokan
et al., 2015), this is the first report of changes in *Vg* expression in the ovaries.
The decreased expression of *Vg* in activated ovaries compared to non-activated
ovaries is unexpected for an egg-yolk precursor protein (Raikhel and Dhadialla, 1992). We
propose two plausible explanations. First, in activated ovaries with mature oocytes the Vg
protein is translocated from the haemolymph *via* VgR into the oocytes (Dohanik *et al.*, 2018;

Guidugli-Lazzarini *et al.*, 2008). Translocation of large amounts of Vg into the ovaries to
nourish developing oocytes may supress transcription of *Vg in situ*. Second, and more
speculatively, it is known that Vg influences the population of micro-RNAs in the honeybee
brain and fat body, and by this means Vg may influence gene expression in these tissues
(Nunes *et al.*, 2013). Thus, Vg may be responsible for regulating the microRNA population
in ways that control ovary activation state. We therefore suggest that *Vg* has a signalling role
in the ovaries of workers, providing additional evidence that the ovary regulates the
behaviour of worker honeybees (Corona *et al.*, 2007; Page, 2013; Vergoz *et al.*, 2012; Wang
et al., 2010).

Feeding Royal Jelly to workers has a major effect on *Vg* expression in their ovaries
(approximately 20-fold higher), which provides an opportunity to investigate the pathways
that regulate *Vg* expression in the ovaries. We determined whether genes in physiological,
epigenetic and signalling pathways are influenced by the presence of Royal Jelly in the diet.
First, *Kr-h1*, a downstream target of JH that is frequently used as a reliable proxy of this
hormone (Belles and Santos, 2014; Jindra *et al.*, 2015; Minakuchi *et al.*, 2009), is
downregulated in Royal Jelly-fed workers. This suggests that JH levels are lowered by the
presence of Royal Jelly in the diet. Since JH represses *Vg* expression (Amdam and Omholt,
2003), this effect is likely to be a factor that promotes *Vg* transcription in the ovaries of
workers fed Royal Jelly. In addition, we find that *Kr-h1* is negatively correlated with *Vg*
expression, suggesting that the ovaries play a role in the "double repressor" interaction
between JH and Vg that possibly regulates maturation in honeybee workers (Amdam and
Omholt, 2003; Cardoso-Júnior *et al.*, 2018; Flatt *et al.*, 2013). Second, *Dnmt3*, an enzyme
involved in DNA methylation, is downregulated in Royal Jelly-fed workers and this diet
blocked the effects of QMP. This finding indicates that epigenetic mechanisms may moderate
the effects of Royal Jelly in the diet, for both the adult (this study) and larval (Kucharski *et*
al., 2008) stages of honeybee workers. Third, *FoxO*, a putative regulator of *Vg* expression in
honeybees (Cardoso-Júnior *et al.*, 2018; Corona *et al.*, 2007), is mainly regulated by post-
translational modification and sub-cellular reallocation (van der Heide *et al.*, 2004). We
found that *FoxO* transcription is reduced in adult workers fed Royal Jelly (this study) as has
been previously shown in honeybee larvae (Wheeler *et al.*, 2014). We suggest that FoxO
negatively regulates *Vg* in adult honeybee workers (but see (Corona *et al.*, 2007)). We note
that in general, insects fed highly nutritious diets switch on the insulin/IGF1 pathway, which
leads to the inactivation of FoxO activity by translocating it to the cytoplasm (Essaghir *et al.*,
2009; Sheng *et al.*, 2011). Together, these results suggest that Royal Jelly acts through
epigenetic, physiological and signalling pathways to increase *Vg* expression.

The diet and social environment of adult honeybee workers have an antagonistic
effect on the activation of their ovaries (Lin and Winston, 1998; Ronai *et al.*, 2016a).
Workers fed a diet containing Royal Jelly activate their ovaries despite the presence of QMP.
The Royal Jelly diet appears to counteract the degeneration of the germ cells that normally
occurs in the ovaries of workers exposed to QMP (figure 1b, 1c and (Ronai *et al.*, 2017,
2015)). Our findings indicate that Royal Jelly can block the inhibitory effect of QMP in the
ovaries of workers. As queens have a diet of Royal Jelly, we speculate that queen's ovaries
are protected from her own pheromone by this diet.

The reproduction-longevity trade-off observed in most animals (De Loof, 2011;
Hayflick, 2007; Shorter *et al.*, 2015) clearly does not apply to honeybees (Winston, 1987).
Reproduction may not be costly for queen honeybees since they have unlimited access to a
highly nutritious diet of Royal Jelly. If a diet of Royal Jelly increases *Vg* expression in
queens, as we have shown here for workers, then high levels of the Vg protein are likely to
act as an anti-oxidant (Corona *et al.*, 2007; Seehuus *et al.*, 2006) and for energy storage
(Nilsen *et al.*, 2011), key factors in promoting longevity (Remolina and Hughes, 2008).

While it is difficult to experimentally test the effects of Royal Jelly in the diet of laying queens, since this is their sole food, our study provides a starting point to uncover the nutrigenomic effects of Royal Jelly in honeybees. Curiously, while a low-dose diet of Royal Jelly increases the lifespan of honeybee workers (Wang *et al.*, 2014; Yang *et al.*, 2017) and other invertebrates (Honda *et al.*, 2011; Wang *et al.*, 2015; Xin *et al.*, 2016), we found that workers fed a higher dose of Royal Jelly have shorter lifespans. A likely explanation for these contrasting results is that workers cannot tolerate high concentrations of Royal Jelly (Shorter *et al.*, 2015; Yang *et al.*, 2017). Alternatively, consumption of Royal Jelly may hasten behavioural maturation and workers that are physiologically older are less tolerant of protein in their diets (Paoli *et al.*, 2014).

In conclusion, our finding suggests that the role of *Vg* in the non-activated ovaries of adult honeybee workers is not as an egg-yolk precursor protein but for other functions. Our study also provides further support for the 'reproductive ground plan' hypothesis, which argues that gene networks that once regulated fertility now regulate behavioural maturation in the worker caste among other functions (Amdam *et al.*, 2006; Hunt and Amdam, 2005; West-Eberhard, n.d.).

Acknowledgments

The authors thank Patsavee Utaipanon for helping with the survival experiment and Melanie Hoang for assisting with ovary dissections. This work was supported by grants from Fundac□ão de Amparo à Pesquisa do Estado de São Paulo (2016/15881-0 and 2017/09269-3 to CA) and the Australian Research Council (DP180101696 to BPO and Amro Zayed).

Author's contributions.

C.A. designed the study, carried out the dissections, survival experiment and molecular lab work, analysed the data and wrote the manuscript; B.P.O. participated in the design of the study, analysed the data and wrote the manuscript; I.R. conceived the study, designed the study, analysed the data, performed the cage experiment and wrote the manuscript. All authors gave final approval for publication and declare no conflict of interests.

Figure Legends:

Figure 1. Effect of QMP and Royal Jelly diet on the ovary state of honeybee workers. (*a*) Average ovary activation score from 7-day-old workers treated with QMP and Royal Jelly diet (QMP$^+$/Control diet N = 64; QMP$^-$/Control diet N = 62; QMP$^+$/Royal Jelly diet N = 63; QMP$^-$/Royal Jelly diet N = 75; different letters represent $p < 0.05$). See supplementary table 2 for further statistical details. (*b*) Semi-activated ovary from a QMP$^+$/Royal Jelly diet worker which shows that Royal Jelly allows the young germ cells to develop whereas the older germ cells have degenerated (dashed line delineates transition). (*c*) Activated ovaries from a QMP$^+$/Royal Jelly diet worker.

Figure 2. Effect of Royal Jelly and QMP exposure on the longevity of honeybee workers. Longevity rate of workers from two source colonies, Colony A (*a*) and Colony B (*b*) fed Royal Jelly (purple lines) or control diet (blue lines), exposed to QMP (continuous lines) or not exposed to QMP (dashed lines). Different letters represent statistical significance ($p < 0.05$; see the supplementary tables for further details).

Figure 3. Relative expression of *Vg* and *VgR* in the ovaries of honeybee workers. (*a*) RT-PCR analysis of three samples (S1, S2 and S3) and a negative control (NC) with no cDNA. RP49 and Ef1α were positive control genes and YFP a negative control gene. (*b*) *Vg* (primer set 1) expression in the ovaries of workers exposed to QMP and fed a Royal Jelly or a control diet. Sample size information: QMP$^+$/Control diet N = 16; QMP$^-$/Control diet N = 15; QMP$^+$/Royal Jelly diet N = 18 and QMP$^-$/Royal Jelly diet N = 26. The inset box is the QMP$^-$/Royal Jelly diet samples split into the non-activated ovaries and activated ovaries (non-activated ovaries N = 17; activated ovaries N = 9). (*c*) *VgR* expression under the same treatment groups as *Vg*. Different letters represent statistical significance ($p < 0.05$; see the supplementary tables for further details) and ** represents $p < 0.01$ obtained after two-tailed Student's *t*-tests.

Figure 4. Relative expression of *Kr-h1*, *Dnmt3* and *FoxO* in the ovaries of honeybee workers. The expression of *Kr-h1* (*a*), *Dnmt3* (*b*) and *FoxO* (*c*) in the ovaries of workers exposed to QMP and fed a Royal Jelly or a control diet. Sample size is the same as in figure 3. Different letters represent statistical significance ($p < 0.05$; see the supplementary tables for further details) and **** represents $p < 0.0001$ obtained after two-tailed Student's *t*-tests.

References

Amdam, G., Csondes, A., Fondrk, M.K., Page Jr., R.E., 2006. Complex social behaviour derived from maternal reproductive traits. *Nature*, **439**, 76–78.

Amdam, G., Norberg, K., Fondrk, M.K., Page, R.E., 2004. Reproductive ground plan may mediate colony-level selection effects on individual foraging behavior in honey bees. *Proceedings of the National Academy of Sciences*, **101**, 11350–11355.

Amdam, G., Norberg, K., Hagen, A., Omholt, S.W., 2003. Social exploitation of vitellogenin. *Proceedings of the National Academy of Sciences,* **100**, 1799–1802.

Amdam, G., Omholt, S.W., 2002. The regulatory anatomy of honeybee lifespan. *Journal of Theorical Biology*, **216**, 209–228.

Amdam, G.V., Omholt, S.W., 2003. The hive bee to forager transition in honeybee colonies: The double repressor hypothesis. *Journal of Theorical Biology,* **223**, 451–464.

Amsalem, E., Malka, O., Grozinger, C., Hefetz, A., 2014. Exploring the role of juvenile hormone and vitellogenin in reproduction and social behavior in bumble bees. *BMC Evolutionary Biology*, **14**, 45.

BELLES, X., Santos, C.G., 2014. The MEKRE93 (Methoprene tolerant-Kruppel homolog 1-E93) pathway in the regulation of insect metamorphosis, and the homology of the pupal stage. *Insect Biochemistry and Molecular Biology*, **52**, 60–68.

Blacher, P., Huggins, T.J., Bourke, A.F.G., 2017. Evolution of ageing, costs of reproduction and the fecundity–longevity trade-off in eusocial insects. *Proceedings of the Royal Society B: Biological Sciences*, **284**, 20170380.

Brito, R.M., McHale, M., Oldroyd, B.P., 2010. Expression of genes related to reproduction and pollen foraging in honey bees (*Apis mellifera*) narcotized with carbon dioxide. *Insect Molecular Biology*, **19**, 451–461.

Cardoso-Júnior, C.A.M., Guidugli-Lazzarini, K.R., Hartfelder, K., 2018. DNA methylation affects the lifespan of honey bee (*Apis mellifera* L.) workers – Evidence for a regulatory module that involves vitellogenin expression but is independent of juvenile hormone function. *Insect Biochemistry and Molecular Biology*, **92**, 21–29.

Corona, M., Libbrecht, R., Wurm, Y., Riba-Grognuz, O., Studer, R.A., Keller, L., 2013. Vitellogenin underwent subfunctionalization to acquire caste and behavioral specific expression in the harvester ant *Pogonomyrmex barbatus*. *PLoS Genetics*, **9**, e1003730.

Corona, M., Velarde, R.A., Remolina, S., Moran-Lauter, A., Wang, Y., Hughes, K.A., Robinson, G.E., 2007. Vitellogenin, juvenile hormone, insulin signaling, and queen honey bee longevity. *Proceedings of the National Academy of Sciences,* **104**, 7128–7133.

De Loof, A., 2011. Longevity and aging in insects: Is reproduction costly; cheap; beneficial or irrelevant? A critical evaluation of the "trade-off" concept. *Journal of Insect Physiology*, **57**, 1–11.

Dohanik, V.T., Goncalves, W.G., Oliveira, L.L., Zanuncio, J.C., Serrao, J.E., 2018. Vitellogenin transcytosis in follicular cells of the honeybee, *Apis mellifera* and the wasp *Polistes simillimus*. *Protoplasma*, **255**, 1703–1712.

Essaghir, A., Dif, N., Marbehant, C.Y., Coffer, P.J., Demoulin, J.B., 2009. The transcription of FOXO genes is stimulated by FOXO3 and repressed by growth factors. *Journal of Biological Chemistry*, **284**, 10334–10342.

Fischer, P., Grozinger, C.M., 2008. Pheromonal regulation of starvation resistance in honey bee workers (*Apis mellifera*). *Naturwissenschaften*, **95**, 723–729.

Flatt, T., Amdam, G. V, Kirkwood, T.B., Omholt, S.W., 2013. Life-history evolution and the polyphenic regulation of somatic maintenance and survival. *The Quarterly Review of Biology*, **88**, 185–218.

Ford, D., 2013. Honeybees and cell lines as models of DNA methylation and aging in

response to diet. *Journal of Experimental Gerontology*, **48**, 614–619.

Grozinger, C.M., Sharabash, N.M., Whitfield, C.W., Robinson, G.E., 2003. Pheromone-
mediated gene expression in the honey bee brain. *Proceedings of the National Academy*
of Sciences, **100 Suppl**, 14519–14525.

Guidugli-Lazzarini, K.R., do Nascimento, A.M., Tanaka, E.D., Piulachs, M.D., Hartfelder,
K., Bitondi, M.G., Simoes, Z.L., 2008. Expression analysis of putative vitellogenin and
lipophorin receptors in honey bee (*Apis mellifera* L.) queens and workers. *Journal of*
Insect Physiology, **54**, 1138–1147.

Guidugli, K.R., Nascimento, A.M., Amdam, G., Barchuk, A.R., Omholt, S., Simoes, Z.L.,
Hartfelder, K., 2005a. Vitellogenin regulates hormonal dynamics in the worker caste of
a eusocial insect. *FEBS Letters*, **579**, 4961–4965.

Guidugli, K.R., Piulachs, M.D., Belles, X., Lourenco, A.P., Simoes, Z.L., 2005b.
Vitellogenin expression in queen ovaries and in larvae of both sexes of *Apis mellifera*.
Archives of Insect Biochemistry and Physiology, **59**, 211–218.

Hagedorn, H.H., Kunkel, J.G., 1979. Vitellogenin and vitellin in insects. *Annual Review of*
Entomology, **24**, 475–505.

Hansen, I.A., Sieglaff, D.H., Munro, J.B., Shiao, S.H., Cruz, J., Lee, I.W., Heraty, J.M.,
Raikhel, A.S., 2007. Forkhead transcription factors regulate mosquito reproduction.
Insect Biochemistry and Molecular Biology, **37**, 985–997.

Hayflick, L., 2007. Biological aging is no longer an unsolved problem. *Annals of the New*
York Academy of Sciences, **1100**, 1–13.

Heinze, J., Schrempf, A., 2008. Aging and reproduction in social insects - A mini-review.
Gerontology, **54**, 160–167.

Honda, Y., Fujita, Y., Maruyama, H., Araki, Y., Ichihara, K., Sato, A., Kojima, T., Tanaka,
M., Nozawa, Y., Ito, M., Honda, S., 2011. Lifespan-extending effects of royal jelly and
its related substances on the nematode *Caenorhabditis elegans*. *PLoS One*, **6**, e23527.

Hoover, S.E., Higo, H.A., Winston, M.L., 2006. Worker honey bee ovary development:
seasonal variation and the influence of larval and adult nutrition. *Journal of*
Comparative Physiology A, **176**, 55–63.

Hoover, S.E., Keeling, C.I., Winston, M.L., Slessor, K.N., 2003. The effect of queen
pheromones on worker honey bee ovary development. *Naturwissenschaften*, **90**, 477–
480.

Hunt, J.H., Amdam, G., 2005. Bivoltinism as an antecedent to eusociality in the paper wasp
genus Polistes. *Science*, **308**, 264–267.

Jedlicka, P., Ernst, U.R., Votavova, A., Hanus, R., Valterova, I., 2016. Gene expression
dynamics in major endocrine regulatory pathways along the transition from solitary to
social life in a Bumblebee, *Bombus terrestris*. *Frontiers in Physiology*, **7**, 574.

Jindra, M., Bellés, X., Shinoda, T., 2015. Molecular basis of juvenile hormone signaling.
Current Opinion in Insect Science, **11**, 39–46.

Koywiwattrakul, P., Thompson, G.J., Sitthipraneed, S., Oldroyd, B.P., Maleszka, R., 2005.
Effects of carbon dioxide narcosis on ovary activation and gene expression in worker
honeybees, *Apis mellifera*. *Journal of Insect Science*, **5**, 36.

Kucharski, R., Maleszka, J., Foret, S., Maleszka, R., 2008. Nutritional control of reproductive
status in honeybees via DNA methylation. *Science*, **319**, 1827–1830.

Lin, H., Winston, M.L., 1998. The role of nutrition and temperature in the ovarian
development of the worker honey bee (*Apis mellifera*). *The Canadian Entomologist*,
130, 883–891.

Lourenço, A.P., Mackert, A., Cristino, A.S., Simoes, Z.L., 2008. Validation of reference
genes for gene expression studies in the honey bee, *Apis mellifera*, by quantitative real-
time RT-PCR. *Apidologie*, **39**, 372–385.

Lourenco, A.P., Martins, J.R., Guidugli-Lazzarini, K.R., Macedo, L.M., Bitondi, M.M., Simoes, Z.L., 2012. Potential costs of bacterial infection on storage protein gene expression and reproduction in queenless *Apis mellifera* worker bees on distinct dietary regimes. *Journal of Insect Physiology*, **58**, 1217–1225.

Marco Antonio, D.S., Guidugli-Lazzarini, K.R., do Nascimento, A.M., Simoes, Z.L., Hartfelder, K., 2008. RNAi-mediated silencing of vitellogenin gene function turns honeybee (*Apis mellifera*) workers into extremely precocious foragers. *Naturwissenschaften*, **95**, 953–961.

Minakuchi, C., Namiki, T., Shinoda, T., 2009. Kruppel homolog 1, an early juvenile hormone-response gene downstream of Methoprene-tolerant, mediates its anti-metamorphic action in the red flour beetle *Tribolium castaneum*. *Developmental Biology*, **325**, 341–350.

Nelson, C.M., Ihle, K.E., Fondrk, M.K., Page, R.E., Amdam, G., 2007. The gene vitellogenin has multiple coordinating effects on social organization. *PLoS Biology*, **5**, e62.

Nilsen, K.A., Ihle, K.E., Frederick, K., Fondrk, M.K., Smedal, B., Hartfelder, K., Amdam, G., 2011. Insulin-like peptide genes in honey bee fat body respond differently to manipulation of social behavioral physiology. *Journal of Experimental Biology*, **214**, 1488–1497.

Nunes, F.M., Ihle, K.E., Mutti, N.S., Simoes, Z.L., Amdam, G., 2013. The gene vitellogenin affects microRNA regulation in honey bee (*Apis mellifera*) fat body and brain. *Journal of Experimental Biology,* 216, 3724–3732.

Page, R.E., 2013. The spirit of the hive. The mechanisms of social evolution. Harvard University Press, Cambridge MA.

Pankiw, T., Page, R.E., 2003. Effect of pheromones, hormones, and handling on sucrose response thresholds of honey bees (*Apis mellifera* L.). *The Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and Behavioral Physiology*, **189**, 675–684.

Paoli, P.P., Donley, D., Stabler, D., Saseendranath, A., Nicolson, S.W., Simpson, S.J., Wright, G.A., 2014. Nutritional balance of essential amino acids and carbohydrates of the adult worker honeybee depends on age. *Amino Acids*, **46**, 1449–1458.

Pfaffl, M.W., Tichopad, A., Prgomet, C., Neuvians, T.P., 2004. Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper - Excel-based tool using pair-wise correlations. *Biotechnology Letters*, **26**, 509–515.

Raikhel, A.S., Dhadialla, T.S., 1992. Accumulation of yolk proteins in insect oocytes. *Annual Review of Entomology*, **37**, 217–251.

Rembold, H., Dietz, A., 1966. Biologically active substances in royal jelly. *Vitamines & Hormones*, **23**, 359–382.

Remolina, S.C., Hughes, K.A., 2008. Evolution and mechanisms of long life and high fertility in queen honey bees. *Age (Omaha)*, **30**, 177–185.

Ronai, I., Allsopp, M.H., Tan, K., Dong, S., Liu, X., Vergoz, V., Oldroyd, B.P., 2017. The dynamic association between ovariole loss and sterility in adult honeybee workers. *Proceedings of the Royal Society B: Biological Sciences*, **284**, 20162693.

Ronai, I., Barton, D.A., Oldroyd, B.P., Vergoz, V., 2015. Regulation of oogenesis in honey bee workers via programed cell death. *Journal of Insect Physiology*, **81**, 36–41.

Ronai, I., Oldroyd, B.P., Vergoz, V., 2016a. Queen pheromone regulates programmed cell death in the honey bee worker ovary. *Insect Molecular Biology*, **25**, 646–652.

Ronai, I., Vergoz, V., Oldroyd, B.P., 2016b. The mechanistic, genetic, and evolutionary basis of worker sterility in the social Hymenoptera. *Advances in the Study of Behavior*, **48**, 251–317.

Rothenbuhler, W.C., Kulincevic, J.M., Thompson, V.C., 1979. Successful selection of honey bees for fast and slow hoarding of sugar syrup in the laboratory. *Journal of Apicultural Research*, **18**, 272–278.

Roy-Zokan, E.M., Cunningham, C.B., Hebb, L.E., McKinney, E.C., Moore, A.J., 2015. Vitellogenin and vitellogenin receptor gene expression is associated with male and female parenting in a subsocial insect. *Proceedings of the Royal Society B: Biological Sciences,* **282**, 20150787.

Seehuus, S.-C., Norberg, K., Krekling, T., Fondrk, K., Amdam, G. V., 2007. Immunogold localization of vitellogenin in the ovaries, hypopharyngeal glands and head fat bodies of honeybee workers, *Apis Mellifera. Journal of Insect Science*, **7**, 1–14.

Seehuus, S.C., Norberg, K., Gimsa, U., Krekling, T., Amdam, G., 2006. Reproductive protein protects functionally sterile honey bee workers from oxidative stress. *Proceedings of the National Academy of Sciences,* **103**, 962–967.

Sheng, Z., Xu, J., Bai, H., Zhu, F., Palli, S.R., 2011. Juvenile hormone regulates vitellogenin gene expression through insulin-like peptide signaling pathway in the red flour beetle, *Tribolium castaneum. Journal of Biological Chemistry*, **286**, 41924–41936.

Shorter, J.R., Geisz, M., Özsoy, E., Magwire, M.M., Carbone, M.A., Mackay, T.F.C., 2015. The effects of royal jelly on fitness traits and gene expression in *Drosophila melanogaster. PLoS One*, **10**, e0134612.

Süren-Castillo, S., Abrisqueta, M., Maestro, J.L., Suren-Castillo, S., Abrisqueta, M., Maestro, J.L., 2012. FoxO inhibits juvenile hormone biosynthesis and vitellogenin production in the German cockroach. *Insect Biochemistry and Molecular Biology*, **42**, 491–498.

Team, R.C., 2018. R: A language and environment for statistical computing. R foundation for statistical computing, Vienna, Austria. URL: http://www.R-project.org/.

Valle, D., 1993. Vitellogenesis in insects and other groups: a review. *Memórias do Instituto Oswaldo Cruz*, **88**, 1–26.

van der Heide, L.P., Hoekman, M.F.M., Smidt, M.P., 2004. The ins and outs of FoxO shuttling: mechanisms of FoxO translocation and transcriptional regulation. *Biochemical Journal*, **380**, 297–309.

van Heemst, D., 2010. Insulin, IGF-1 and longevity. *Aging and Disease*, **1**, 147–157.

Vergoz, V., Lim, J., Oldroyd, B.P., 2012. Biogenic amine receptor gene expression in the ovarian tissue of the honey bee *Apis mellifera. Insect Molecular Biology*, **21**, 21–29.

Wang, H., Zhang, S.W., Zeng, Z.J., Yan, W.Y., 2014. Nutrition affects longevity and gene expression in honey bee (*Apis mellifera*) workers. *Apidologie*, 45, 618–625.

Wang, X., Cook, L.F., Grasso, L.M., Cao, M., Dong, Y., 2015. Royal jelly-mediated prolongevity and stress resistance in *Caenorhabditis elegans* is possibly modulated by the interplays of DAF-16, SIR-2.1, HCF-1, and 14-3-3 proteins. *The Journals of Gerontology Series A Biological Sciences and Medical Sciences*, **70**, 827–838.

Wang, Y., Kaftanoglu, O., Siegel, A., Page, R., Amdam, G., 2010. Surgically increased ovarian mass in the honey bee confirms link between reproductive physiology and worker behavior. *Journal of Insect Physiology*, **56**, 1816–1824.

West-Eberhard, M.J., n.d. Wasp societies as microcosms for the study of development and evolution. In: Turillazzi, S., West-Eberhard, M.J. (Eds.), Natural History and Evolution of Paper-Wasps. Oxford University Press, Oxford, pp. 290–317.

Wheeler, D.E., Buck, N.A., Evans, J.D., 2014. Expression of insulin/insulin-like signalling and TOR pathway genes in honey bee caste determination. *Insect Molecular Biology*, **23**, 113–121.

Winston, M.L., 1987. The biology of the honey bee. Harvard University Press, Cambridge.

Xin, X.X., Chen, Y., Chen, D., Xiao, F., Parnell, L.D., Zhao, J., Liu, L., Ordovas, J.M., Lai, C.Q., Shen, L.R., 2016. Supplementation with major royal-jelly proteins increases

645   lifespan, feeding, and fecundity in *Drosophila*. *Journal of Agricultural and Food*
646   *Chemistry*, **64**, 5803–5812.
647   Yang, W., Tian, Y., Han, M., Miao, X., 2017. Longevity extension of worker honey bees
648   (*Apis mellifera*) by royal jelly: optimal dose and active ingredient. *PeerJ*, **5**, e3118.
649

(a) **Colony A**

Percent survival (%)

Stats

— Control Diet / QMP$^+$ ac

···· Control Diet / QMP$^-$ a

— Royal Jelly Diet / QMP$^+$ b

···· Royal Jelly Diet / QMP$^-$ c

Days

(b) **Colony B**

Percent survival (%)

Stats

— Control Diet / QMP$^+$ a

···· Control Diet / QMP$^-$ a

— Royal Jelly Diet / QMP$^+$ b

···· Royal Jelly Diet / QMP$^-$ b

Days