Over-activation of BMP signaling in neural crest cells precipitates heart outflow tract septation

Jean-François Darrigrand[1], Mariana Valente[2], Pauline Martinez[1], Glenda Comai[3], Maxime Petit[4], Ryuichi Nishinakamura[5], Daniel S. Osorio[6], Vanessa Ribes[7], Bruno Cadot[1#].

1. INSERM - Sorbonne Université UMR974 - Center for Research in Myology, 105 boulevard de l'Hôpital, 75634 Paris Cedex 13, France.
2. Cellular, Molecular, and Physiological Mechanisms of Heart Failure team, Paris-Cardiovascular Research Center (PARCC), European Georges Pompidou Hospital (HEGP), INSERM U970, F-75737 Paris Cedex 15, France.
3. Stem Cells and Development, Department of Developmental & Stem Cell Biology, CNRS UMR 3738, Institut Pasteur, 25 rue du Dr. Roux, 75015 Paris, France.
4. Unité Lymphopoïèse – INSERM U1223, Institut Pasteur, 25 rue du Dr. Roux, 75015 Paris, France.
5. Institute of Molecular Embryology and Genetics, Kumamoto University, Japan.
6. Instituto de Investigação e Inovação em Saúde – i3S, Universidade do Porto, and Instituto de Biologia Molecular e Celular – IBMC, Porto, Portugal.
7. Institut Jacques Monod, CNRS UMR7592, Université Paris Diderot, Sorbonne Paris Cité, 75205 Paris Cedex, France.

corresponding author and lead contact: bruno.cadot@inserm.fr

Short title: BMPs and heart outflow tract

Summary

Establishment of separated pulmonary and systemic circulations in vertebrates relies on the key
role of neural crest cells (NCC) for the septation of the embryonic cardiac outflow tract (OFT).
Absence of NCCs induces OFT septation defects, analogous to a loss of Bone Morphogenetic
Proteins (BMPs) activity, though it remains unclear how BMPs control cardiac NCC differentiation
and behaviour. To address this question, we monitored cardiac NCC state upon gain in BMP
signaling, caused by the deletion of *Dullard*, using 3D-imaging and single cell transcriptomics.
Specific loss of *Dullard* in the NCC results in premature OFT septation, pulmonary artery
obstruction and embryonic death. This is caused by uncontrolled NCC convergence towards the
endocardium and asymmetrical myocardial differentiation, promoted by elevated levels of the
guiding cue *Sema3c* and decreased levels in mesenchymal trait markers. Furthermore, we
unraveled the molecular basis of the zipper-like OFT septation where graded Sema3c expression
follow a gradient of BMP activation in NCC along the OFT length.

Keywords

Introduction

The embryonic OFT is a solitary tube called truncus arteriosus, which is remodelled into the aortic (Ao) and pulmonary (Pa) arteries (1) (Fig. 1A). This morphogenesis is orchestrated in time and space by several cross-interacting cell-types including the myocardial progenitors of the second heart field (SHF), the endocardial cells (EC) delineating the OFT lumen, and the cardiac neural crest cells (cNCC) (2,3) (Fig. 1A). Various genetic manipulations or ablations have highlighted the predominant role of cNCC in initiating and controlling OFT septation (4,5). Originally, cNCC delaminate from the dorsal neural tube and migrate through the pharyngeal mesoderm to reach the developing OFT (Fig. 1A). They invade the two cardiac cushions, condense towards the endocardium and trigger its rupture, thereby creating the two great arteries (6,7). The rupture of the endocardium is first detected at the distal levels of the OFT, at E11.5 in mouse, and gets progressively established at more proximal levels.

The intense investigations to identify the molecular cues controlling cNCC stereotyped behaviour and differentiation in the OFT mesenchyme have established the importance of the Bone Morphogenic Proteins (BMP), secreted by the outlying myocardium cells (8–10). Indeed, BMP signaling loss-of-function within the NCC lineage leads to the formation of hypoplastic cushions, a shorter and non-septated OFT, phenocopying the absence of cNCC (11–13). However, little is known on the molecular cascades triggered by BMP signalling responsible for the cNCC mediated OFT septation.

To get insights into these molecular cascades, we decided to dissect the role of Dullard (Ctdnep1)

during OFT morphogenesis, a perinuclear phosphatase uncovered as a BMP intracellular signaling

inhibitor (14–16). Dullard protein is evolutionary conserved from yeast to mammals. In xenopus,

drosophila and mouse embryos, it decreases the levels of phosphorylation of Smad1/5/8, the

downstream transcriptional effectors of BMP signalling (15–17).

We showed here that deletion of *Dullard* in the cNCC increases Smad1/5/8 activity, leading to

premature and asymmetric septation of the OFT, pulmonary artery closure and embryonic death.

This BMP overactivation in the cNCC triggers the downregulation of mesenchymal markers

(*Snai2*, *Twist1, Rac1, Mmp14* and *Cdh2*) and the upregulation of *Sema3c*, which in turn results in

premature cNCC compaction at the endocardium. In addition, asymmetric differentiation of the

distal subpulmonary myocardium contributes to OFT anisotropic rupture. Finally, our data

converge to a model whereby graded BMP activity and *Sema3c* expression in the cNCC along the

OFT axis set the tempo of OFT septation from its distal to its proximal regions. Hence, our findings

reveal that fine tuning of BMP signaling levels in cNCC orchestrate OFT septation in time and

space.

Results

***Dullard* deletion triggers hyper-activation of BMP intracellular signalling in cNCC**

In order to dig into BMP signalling functions during OFT septation, we decided to delete the *Dullard* phosphatase in cNCC. Its role in inhibiting the BMP pathway was first confirmed in vitro using the myogenic cell line C2C12, transfected with *Dullard* WT or *Dullard* with the phosphatase-dead domain (D67E) and exposed for one hour to BMP2 (Fig. S1A). We then used mice carrying floxed versions of *Dullard* allele and expressing the Cre recombinase under the *Pax3* or *Wnt1* loci, which are active within the pre-delaminating cNCC (14,18,19), in combination with a ubiquitous double-fluorescent Cre reporter allele, *Rosa26mTmG,* in which Cre-mediated recombination labels all the NCC-derived cells with membrane-targeted GFP (mGFP+) (20). In agreement with previous lineage analyses the pattern of cell recombination in the heart of embryos carrying either Cre drivers matched that of the cNCC colonizing the cardiac cushions (Fig. S1B) (21,22). We also confirmed by RT-qPCR that Fac-sorted (FACS) single GFP+ cells from *Wnt1^{Cre}; Dullard^{flox/flox};* *Rosa26mTmG* dissected hearts were recombined and did not express *Dullard* (Fig. S1C). We next wanted to assess the activity of intracellular BMP signaling in the cNCC after *Dullard* deletion. To this end, we immunolabelled the phosphorylated forms of Smad1/5/8, on serial cryosections along the OFT axis in E11.5 embryos to follow the distal to proximal progression of cNCC (Fig. 1B, C). Paralleling this time-dependent colonization, quantifications revealed variations of phospho-Smad levels, which were more elevated proximally than distally (Fig. 1C). Furthermore, whatever the levels along the OFT axis, there was a striking increase in Smads activity in both *Pax3^{Cre}* and

Wnt1^{Cre}; Dullard$^{flox/flox}$ mutant cNCC compared to control cNCC (Fig. 1B, C). Altogether, these

results show that, although Dullard is not absolutely required for the establishment of a gradient

of BMP response in cNCC, it is necessary for a tight regulation of its magnitude.

Dullard deletion in cNCC accelerates and jeopardizes OFT septation

The absence of *Dullard* was embryonically lethal in both *Pax3^{Cre}* and *Wnt1^{Cre}* mice. While

Pax3^{Cre}/Wnt1^{Cre}; Dullard$^{flox/flox}$ embryos were recovered at the expected 25% mendelian ratio by

E12.5, this ratio dropped below 10% afterward (Fig. S1D). We first assessed whether this could

be due to defects in cNCC colonization of the OFT. 3D imaging of GFP^{+} cNCC revealed that these

cells reached similar OFT levels in E11.5 control and *Dullard* mutants (Fig. S2A; Movie S1-S4). In

addition, no significant differences in the cNCC number nor in their proliferation and apoptosis

rates were detected between control and mutant OFT, although the mutant OFT tended to be

smaller in diameter (Fig. S2B-E). Thus, *Dullard* is not required for cNCC migration nor for their

amplification and survival in the OFT.

We next examined the heart septation status of *Dullard* mutants, knowing that cNCC are critical

in this morphogenesis process (4–6). For this, we monitored, using 3D lightsheet and confocal

microscopy, E11.5-12 hearts labelled for the arteries marker, Pecam (Fig. 2A; Movie S5, S6). At

distal levels, the OFT of control embryos displayed symmetrical septation and two great arteries

of similar size. In contrast, *Pax3^{Cre}/Wnt1^{Cre}; Dullard$^{flox/flox}$* embryos exhibited asymmetric

breakdown of the endocardium on the pulmonary side and obstruction of the pulmonary artery

(Pa) (Fig. 2A,B). At medial levels in *Pax3*[Cre]/*Wnt1*[Cre]; *Dullard*[flox/flox], the cardiac cushions were

already fused, though on the pulmonary side the Pa was absent compared to controls (Fig. 2B).

At medial levels in controls the endocardium attached to the presumptive pulmonary valve

intercalated-cushion (PV-IC), while in mutants this attachment had broken down and the

endocardium was pushed on the aortic side by converging cNCC. These observations

demonstrate that *Dullard* plays a pivotal role during cNCC-mediated OFT septation through the

modulation of BMP signalling and provided an explanation for the early lethality of *Dullard*

mutants, probably due to haemodynamics complications associated with Pa deficiency.

Dullard controls cNCC mesenchymal state and Sema3c expression
138

We set out to decipher the molecular basis of the defective OFT remodeling observed in

Pax3[Cre]/*Wnt1*[Cre]; *Dullard*[flox/flox] embryos. To address this question, we micro-dissected E11.5

Wnt1[Cre]; *Dullard*[+/flox]; *Rosa26mTmG* and *Wnt1*[Cre]; *Dullard*[flox/flox]; *Rosa26mTmG* OFT and sorted the

cNCC (GFP[+]) and the endocardial cells (CD31[+]) from the other OFT cell-types (RFP[+]) (Fig. 3A). We

then performed single cell RT-qPCR for 44 genes implicated in epithelial-mesenchymal transition

(EMT), migration and/or specification of OFT progenitors (Table S1). Ct values extracted from the

Biomark chips were normalized using the mean value of *Actb* and *Gapdh* housekeeping genes

(Fig. S3A). We next performed an unsupervised hierarchical clustering of all cells according to

their molecular signature. This method allowed the segregation of 6 distinct OFT progenitor

subtypes (Fig. S3B): the Population 1 (Pop1) regrouped endocardial (CD31+) cells, all positive for

the endocardial markers, *Flt1*, *Kdr, Nfatc* and *Tek,* and the Pop5 encompassed epicardial cells

(*Tcf21* and *Wt1*). Importantly, these two populations contained cells derived from both *Wnt1^Cre^;*

*Dullard^+/flox^* and *Wnt1^Cre^; Dullard^flox/flox^* OFT. In contrast, the cNCC coming from *Wnt1^Cre^;*

*Dullard^+/flox^* and *Wnt1^Cre^; Dullard^flox/flox^* OFT were segregated in distinct subpopulations. The

control cNCC were mainly found in Pop2, while most mutant cNCC were in Pop4 and 6 (Fig. S3B).

This indicated that Dullard loss induced a drastic effect on the transcriptional state of cNCC, and

had little impact on the other progenitor cell-types.

We next decided to identify the transcriptomic changes underpinning the segregation between

mutant and control cNCC. By performing unsupervised hierarchical clustering only on cNCC, we

observed that the transcriptional state of cNCC was variable and that five populations of cells

could be defined (Pop1-5)(Fig 3B,C). Both control and mutant cNCC were found in Pop1 and 2,

and thus *Dullard* deletion did not impact the path between these transcriptional signatures (Fig.

S3C). Conversely, barely any mutant cNCC clustered with the Pop3 control cells. This population

of cells presented an upregulation of smooth muscle specification markers *Myh11* and *Acta2*,

indicating their commitment towards a smooth muscle fate, absent in mutant cNCC (Fig. 3D). In

contrast, mutant cNCC clustered in Pop4 and even more strongly in Pop5 (28% of mutant versus

3.95% of control cNCC) (Fig. S3C). Pop5 showed low levels of the mesenchymal markers *Snai2,*

Twist1, Cdh2, Mmp14, Rac1 and high levels of the epithelial marker *Cdh5,* suggesting that mutant

cNCC transitioned towards epithelial-like states (Fig. 3D, Fig. S3B-E). In addition, this population

was characterized by higher levels of *Sema3c* expression, a chemokine implicated in cells

migration, and lower expression of its receptors *Nrp1* and *Plxnd1*. Cells in Pop4 resembled of

those in pop5, although the fold change in the levels of mesenchymal genes, *Cdh5* and *Sema3c*

compared to cNCC was less pronounced. This indicates a close relationship between these two

subsets which might represent two states of the differentiation path of *Dullard* mutated cNCC.

Similar analyses on the RFP$^+$ and CD31$^+$ cells did not show any major effect on RFP$^+$ cells, while

CD31$^+$ sorted from *Wnt1^{Cre}; Dullard$^{Flox/Flox}$* OFT presented a significant decrease in the expression

of *Twist1* and *Snai2* (Fig. S4A-D). In sum, our results suggest that *Dullard* deletion in cNCC mainly

affected cell-autonomously the NCC by changing their mesenchymal properties and their

communication properties via *Sema3c* signaling but had a very limited impact on the

transcriptional signature of the other OFT cell populations.

Sema3c expression is associated with cNCC condensation to the endocardium
and symmetrical differentiation of the SHF.

We then investigated if the specific alterations in the transcriptome of *Wnt1^{Cre}; Dullard$^{flox/flox}$*

cNCC compared to control cells were translated into discrete traits in their behavior. First, by

using a nuclear staining, we analyzed at OFT lower levels, when the endocardium is not yet

ruptured, the condensation pattern of cNCC. In control embryos, sparse and elongated cNCC

nuclei were found at the proximity of the endocardium, whereas away from the endocardium

they harboured a rounded shape and high density (Fig. 4A; S5A). In mutant embryos, the sparse

population of cNCC was barely visible. We thus measured both the shortest distance between

cNCC nuclei and the endocardium as well as between each cNCC. *Wnt1^{Cre}; Dullard$^{flox/flox}$* cNCC

presented a significant decrease both in distance to the endocardium and to their closest

neighbor compared to the *Wnt1^{Cre}; Dullard$^{+/flox}$* cNCC (Fig. 4B). Hence, in absence of *Dullard,* cNCC

are more aggregated and closer to the endocardium than controls (Fig 4A), in coherence with the

the transcriptional transition towards epithelial-like states (Fig. 3). In addition, we noticed that

the orientation of the cNCC nuclei relative to the endocardium, revealing cell orientation, was

different between control and mutant OFT. We consequently determined the angle formed by

the longest axis of the different cNCC with the axis spanning the endocardium and plotted it in a

[0°,90°] referential. While in control embryos, most cNCC nuclei were parallel to the

endocardium, in *Dullard* mutants a significantly higher percentage of cNCC presented nuclei

standing perpendicular to the endocardium (Fig. S5B). In absence of *Dullard*, the aggregation of

cNCC, their convergence towards the endocardium and orientation relative to the endocardium

are enhanced compared to control cells demonstrating that *Dullard* regulates cNCC

morphogenetic cellular properties involved in OFT septation.

Sema3c stands as a cue involved in cNCC convergence towards the endocardium (6), and was

upregulated in our single cell transcriptional studies. We analyzed its distribution at the mRNA

level by *in situ* hybridization (ISH) performed on OFT serial sections and confirmed the increase

in *Sema3c* expression in Pax3/*Wnt1^Cre^;Dullard^flox/flox^* cNCC (Fig. 4C, S5C). The *Sema3c* expression

territories increased distally forming a gradient along the OFT axis of both control and mutant

embryos, parallel to the previously described gradient of BMP signaling activity. Moreover, in the

distal region, *Sema3c* expression territories converged towards the endocardium and thus

associated with areas of cNCC condensation (Fig. 4A,C; S5A,C). Hence, our results highlight a

BMP-regulated gradient of *Sema3c* expression, established along the OFT by the cNCC, that drives

the zipper-like convergence of the cNCC towards the endocardium, and the distal to proximal

OFT septation.

At the distal OFT level, the outer SHF is differentiating in myocardium and starts to express

Myosin Heavy Chain (MyHC, stained by MF20 antibody) (Fig 4D) (23,24). First, we noted a

coregionalization of SHF progenitors expressing MyHC and Sema3c, which is a marker of

subpulmonary myocardium (compare Fig. 4C and Fig. S5C with Fig. S5D) (25,26). Second, the

MyHC myocardium localization differed between *Pax3^{Cre}/Wnt1^{Cre};Dullard$^{flox/flox}$* and

Pax3^{Cre}/Wnt1^{Cre};Dullard$^{flox/+}$ embryos. In the later, the MyHC+ myocardium was located medially

and symmetrically between the Ao and Pa sides, whereas in the former it was asymmetrically

positioned on the pulmonary side (Fig 4D, Fig. S5D). These observations highlight that

symmetrical septation at the distal OFT level is supported by the medial differentiation of

subpulmonary myocardium and that the BMP-dependent regulation of *Sema3c* expression in the

cNCC is crucial for symmetrical positioning of the differentiating subpulmonary myocardium.

Discussion

By looking into the function of the *Dullard* phosphatase during cNCC-mediated OFT septation, we

have uncovered that spatially regulated BMP-dependent aggregation of cNCC set the timing and

symmetrical shaping of the endocardium rupture. We showed that a gradient of BMP activation

exists in the cNCC along the OFT axis, and notably its amplitude, is under the control of the

phosphatase *Dullard*. Other mechanisms are likely to be responsible for its decline along the OFT,

as it is still observed in absence of *Dullard*. In contrast to other developmental systems, this

spatial gradient of BMP activity is unlikely resulting from a gradient of ligand diffusing from a

localized distal source (27). BMP4, particularly, is homogeneously expressed in the myocardium

throughout the entire length of the OFT (8,9,11,28,29). Instead, intracellular signalling inhibitors

could allow a temporal adaptation of cNCC to BMP signals during their proximal ingression.

Prominent expression of *Smad6*, a BMP negative feedback effector, or of the diffusible inhibitor

Noggin is observed in the OFT from E10.5 throughout great arteries formation (30,31).

Importantly, this gradient provides a means by which the zipper-like septation of the OFT can

proceed. The levels of BMP stand as a sufficient parameter to tune the degree of convergence

and aggregation of cNCC to the endocardium. As such, lowering down BMP levels prevents the

expansion of the cardiac cushions and leads to persistent truncus arteriosus (11,12), while

increasing BMP signalling triggers premature cNCC condensation to the endocardium and

accentuated OFT septation (Fig 2). This model is further supported by evidence showing that

altering the ability of cNCC to contract, migrate and adhere, hence to condensate, prevents the

correct formation and positioning of the aorticopulmonary septum (5,6,32).

We have uncovered part of the molecular mechanisms by which BMP controls cNCC behavior. A

BMP over-activation triggers a downregulation of mesenchymal markers, reminiscent of a

transition of the cNCC towards epithelial-like states, with a loss of migratory freedom and

increased cohesiveness between cells (33). Furthermore, out of the transcriptional changes

induced by *Dullard* loss, several lines of evidence support the idea that the up-regulation of

Sema3c is likely to play a predominant role in the increased compaction of cNCC. The secretion

of this guiding cue from the subpulmonary myocardium attracts cNCC towards the OFT, while its

expression in the cNCC is required for their convergence to the endocardium (6,34,35). *Sema3c*

is also promoting the self aggregation of cNCC, as demonstrated in primary cultures (36). Besides,

our results show that increased *Sema3c* levels precipitate OFT septation, not by breaking the

endocardium structure itself as expected by previously shown involvement of *Sema3c* in the NCC

during the endocardium EMT (Plein et al., 2015), but through a rupture in the attachment of the

endocardium to the subpulmonary wall. Finally, our observations indicate that the correct

septation onset, at the distal OFT level, requires the central positioning of differentiating

subpulmonary myocardium. BMP mediated induction of *Sema3c* would control the isotropic

localization of differentiated myocardial cells. On an evolutionary perspective, Keyte et al.

postulated that the emergence of *Sema3c* expression in the pulmonary myocardium has

represented a turning point in the dialogue between SHF and cNCC by attracting cNCC to the

OFT (37). Our results suggest that conversely, cNCC have a favored ability to dialogue with the

subpulmonary myocardium, consolidating the idea of a co-evolution between the cNCC and

subpulmonary myocardium during the evolution of the pulmonary circulation.

Experimental procedures

Mouse strains

All animal experiments were approved by the Animal Ethics Committee of Sorbonne University.

We used the mouse strains described in the following papers with their MGI IDs: *Dullard$^{flox/flox}$*

Acc No. CDB0564K, Web site : http://www.cdb.riken.jp/arg/mutant%20mice%20list.html

(Sakaguchi et al. 2013), *Pax3^{Cre}* ((19), MGI: 3573783), *Wnt1^{Cre}* ((18), MGI:2386570),

Rosa26mTmG ((20), MGI: 3716464), and *C57BL/6JRj* (Janvier Labs).

Immunohistochemistry and imaging

Mouse embryos were collected at E11.5 and dissected in PBS (4°c), incubated 5min in 200mM

KCl to stop heart beating and fixed for 2-3 hours in 4% PFA (Electron Microscopy Science,

#15710S) at 4°c. They were cryoprotected in 20% sucrose overnight at 4°c, embedded in OCT,

and cryosectioned at 12μm thickness.

Sections were permeabilized 10min in PBS/0.5% Triton, incubated for 1h in blocking buffer (5%

goat serum in PBS) and overnight in primary antibody solution (in 1% BSA in PBS). After thorough

washing they were incubated 1h in secondary antibody solution (in 1% BSA in PBS). Lightsheet

acquisition was performed on whole embryos cleared following the 3Disco protocol,

immunostained and imaged using an ultramicroscope (LaVision BioTec) (38). 3D confocal

microscopy was performed on micro-dissected OFT regions immunostained, clarified with BABB

and acquired using a LSM700 confocal microscope (Carl Zeiss) (39). 3D renderings were

generated using the Imaris software.

The primary antibodies used on cryosections include the following: GFP (chicken, Aves Labs, GFP-

1020, 1/500), Pecam (rat monoclonal, Santa-Cruz Biotechnology, sc-18916, 1/200), Phospho-

Smad1/5/8 (rabbit monoclonal, Cell Signaling Technology,13820S,1/500), myosin heavy chain,

MyHC (mouse monoclonal, DSHB, MF20,1/300), Smooth Muscle Actin (mouse monoclonal,

Sigma-Aldrich, A2547, 1/300), Phospho-Histone H3 (rabbit, Cell Signaling Technology, 9701,

1/500), Cleaved Caspase-3 (rabbit monoclonal, Cell Signaling Technology, Asp175, 1/500).

In situ hybridisation

The Sema3c probe was provided by the lab of S. Zaffran (25). Cryosections were processed

following the protocol described in (40).

Image analysis and quantification

Immunostainings were acquired using a Nikon Ti2 microscope and Metamorph (Molecular

Devices), then assembled and analyzed on Fiji (41). Quantification of P-Smad1/5/8 was

performed in cNCC by using the GFP channel as a mask. Measures of cNCC distance and

orientation to the endocardium were performed using Metamorph (Molecular Devices). The

coordinates of the endocardium extremities were registered to create a reference axis as well all

cNCC nuclei longest axis coordinates. From these data we inferred the centroid coordinates of all

cNCC nuclei and calculated their distances and angles to the endocardium axis.

Cell culture and transfection

C2C12 cells were cultivated at 37°C/5% CO2, in growth medium (DMEM, 4.5g/L, D-glucose, 4mM

L-glutamine, 1mM sodium pyruvate, 10% fetal calf serum). Plasmid transfection was performed

using Lipofectamine 2000 (Life Technologies). 24 hours after transfection, BMP2 recombinant

human protein (Thermo Fisher Scientific, #PHC7145) was applied for 1h on cells. Cells were fixed

with 4% PFA and immunostained with the phospho-Smad1/5/8 antibody. Or, cells were washed

in PBS, collected with PBS 1% SDS and passed through Qiashredder columns (Qiagen) to disrupt

nucleic acids. Proteins extracts were then processed for Western blotting using pre-cast gels (Life

Technologies) and transferred on nitrocellulose by semi-dry transfer (Bio-Rad).

 Tissue dissociation and FACS sorting

Mouse embryos were collected at E11.5 and placed in HBSS/1% FBS (HBSS +/+, Invitrogen) during

genotyping. OFT were micro-dissected and dissociated by 15min incubation in collagenase

(0.1mg/ml in H_2O, C2139 Sigma) and thorough pipetting. HBSS (10% FBS) was added to the cells

medium to stop the enzymatic reaction. OFT cell suspensions were centrifuged and resuspended

in HBSS (1% FBS) before immunostaining. The panel of conjugated antibodies used for FACS

included Ter119 Pe-Cy7 (Erythroid Cells, anti-mouse, Sony, Catalog No. 1181110, 1/300), CD45

APC-Cy7 (Rat anti-mouse, BD Pharmingen, Catalog No. 557659, 1/300), CD31 APC (Rat anti-

mouse, BD Pharmingen, Clone MEC 13.3 Catalog No. 561814, 1/300) diluted in HBSS (1% FBS).

Cells were centrifuged and resuspended in the antibody solution for a 25min incubation period

(4°C, dark), washed several times, filtered (Fisher cell strainer, 70μm mesh) and 7AAD PE-Cy7

(1/800) was added in the cells suspension to exclude dead cells. Cells were sorted in a BD

FACSAria™ III into 96-well plates loaded with RT-STA reaction mix (CellsDirect™ One-Step qRTPCR

Kit, Invitrogen) and 0.2x specific TaqMan ® Assay mix (see Table 1 for assays list).

Single-Cell gene expression

Sorted cells in RT-STA reaction mix from the CellsDirect TM One-Step qRT-PCR Kit (Life

Technologies), reverse transcribed and specific target pre-amplified (20 cycles), according to the

manufacturer's procedures. Pre-amplified samples were diluted 5x with low EDTA TE buffer prior

to qPCR analysis using 48.48 Dynamic Array TM IFCs and the BioMark TM HD System (Fluidigm).

The same TaqMan® gene expression assays (20x, Life Technologies) were individually diluted 1:1

with 2x assay loading reagent (Fluidigm). Pre-amplified samples were combined with TaqMan

Universal Master Mix (Life Technologies) and 20x GE sample loading reagent (Fluidigm). Loading

of the 48.48 Dynamic Array TM IFCs and qPCR cycling conditions followed the Fluidigm procedure

for TaqMan ® gene expression assays.

Bioinformatic analysis

The analytic framework used followed the one described in (42). For visualisation of the single

cell multiplex qPCR, done on 44 genes, we generated a heatmap using the pHeatmap (v1.0.10) R

package (v3.2.2).

Statistical analysis

Statistical analysis was performed with the Student's t-test or Mann-Whitney test depending on

normality. The analysis was performed using Prism Software (GraphPad). Statistical significance

is represented as follows: ***$p < 0.001$. All results are shown as mean ± standard deviation.

Acknowledgments

We thank the Cadot, Bitoun and Ribes Laboratories for discussions, Sigolène Meilhac for her help on understanding heart development concepts, Edgar Gomes laboratories for useful comments, Stéphane Zaffran for the *Sema3c* ISH probe and Morgane Belle for Lightsheet microscopy. The *Pax3^Cre* and *Rosa26mTmG* transgenic lines were kindly provided by F. Relaix, and the *Wnt1^Cre* line by A. Pierani. This work was supported by Agence Nationale pour la Recherche (ANR-14-CE09-0006-04) to BC; Association Institut de Myologie to BC. V.R. is an INSERM researcher, and work in her lab is supported by CNRS/INSERM ATIP-AVENIR program, as well as by a Ligue Nationale Contre le Cancer grant (PREAC2016.LCC). MV has a postdoctoral fellowship from the Laboratoire d'Excellence Revive (Investissement d'Avenir; ANR-10-LABX-73).

Authors contributions

JFD, PM, GC, MV and BC performed the experiments; MP analyzed the transcriptomics data; JFD, VR and BC conceived and designed the experiments; JFD, VR and BC wrote the manuscript with assistance from MV and GC; DSO and RN provided reagents.

Figures legends

Figure 1: Dullard is required to downregulate cNCC response to myocardial BMP signals. A: On the left, schematic representation of cNCC (green) migrating from the neural tube, through the somites (S) 1 to 3, towards the heart region (red). On the right, schematic transversal sections of the OFT showing cardiac cushions colonization by cNCC supporting OFT septation. **B:** Transversal sections of E11.5 control *Wnt1^{Cre}/Pax3^{Cre}; Dullard$^{flox/+}$; Rosa26mTmG* and mutant *Wnt1^{Cre}/Pax3^{Cre}; Dullard$^{flox/flox}$; Rosa26mTmG* hearts labelled for phospho-Smad1/5/8 (PSmad1/5/8) and cNCC-derived GFP+ (green). Green dotted lines delineate the cNCC. **C:** Heart schema displaying the position along the OFT distal-proximal axis of transverse sections used to measure PSmad1/5/8 levels and graph showing PSmad1/5/8 intensity levels in the cNCC (mean+/-s.e.m.)

Figure 2: Dullard deletion in cNCC causes asymmetric and premature OFT septation. A. Three-dimensional rendering of the OFT from E11.5 *Pax3^{Cre}; Dullard$^{flox/+}$* and *Pax3^{Cre}; Dullard$^{flox/flox}$* embryos (Pa width represented with an oblique white line). **B.** Transversal sections along the OFT of E11.5 *Wnt1^{Cre}; Dullard$^{flox/+}$; Rosa26mTmG* and mutant *Wnt1^{Cre}; Dullard$^{flox/flox}$; Rosa26mTmG* embryos stained for Pecam (red) and GFP (green). Brackets highlight the asymmetry between the Ao and Pa poles in mutants compared to controls. Arrowheads point at the ruptured endocardium in mutant embryos contrasting with its unruptured state in controls. Ao: Aorta; Pa: pulmonary artery; PV-IC: Pulmonary valve intercalated-cushion

Figure 3: Single-cell analyses unravel a role for Dullard in regulating transcriptomic states of

cNCC. A: Main steps performed to profile gene expression in OFT progenitor cells sorted from

Wnt1^{Cre}; Dullard$^{flox/+}$; Rosa26mTmG and *Wnt1^{Cre}; Dullard$^{flox/flox}$; Rosa26mTmG* E11.5 embryos. **B:**

Unsupervised hierarchical clustering of GFP$^+$ cNCC based on the levels of expression of the 44

genes included in the panel (Table S1). **C:** Diffusion maps of the cNCC according to their cluster

of appartenance and their genotype. **D:** Boxplot representation of the expression level of genes

differentially expressed between cNCC Pop3 and 5.

Figure 4: BMP signaling regulates *Sema3c* expression and cNCC condensation, as well as distal

myocardialization. **A:** Transversal sections of the OFT stained with DAPI. The endocardium is

delineated with a red line and the inner side of the myocardium with a greyline. Areas where

cNCC are of high density and rounded nuclear shape are surrounded with a green dotted line,

while areas where cNCC are of low density and elongated nuclear shape are surrounded with a

yellow dotted line. **B:** Measure of the shortest distance between cNCC nuclei and to the

endocardium in both genotypes (5 embryos analyzed per genotype, Mann-Whitney Test, P-value

< 0.0001). **C:** Sema3c ISH and Pecam/GFP immunostaining on OFT transversal sections at distal,

medial and proximal levels, from the indicated genotypes. **D:** Distal sections of the OFT stained

for Myosin Heavy Chain (MF20) to mark differentiated myocardium in control and mutant

embryos.

Figure S1: Dullard is a regulator of BMP signaling and its deletion in Neural Crest Cells induces

embryonic death around E12.5. A: Top: Western blot of the phosphorylation of Smad1/5/8 in

the muscle cell line C2C12 after exposure to BMP2 for 1H. Cells transfected with Dullard (Dull)

do not show any phosphorylation of Smad1/5/8 whereas a mutation in the phosphatase domain

(D67E) retains this phosphorylation. Bottom-left: A similar result was obtained by

immunofluorescence showing that only cells transfected with Dullard do not have nuclear

phosphorylated Smad1/5/8. Right: A quantification of three different experiments was

performed on transfected cells. **B:** *Wnt1^{Cre}* and *Pax3^{Cre}* mouse driver lines present the same

pattern of cNCC colonization in the OFT as shown by the reporter line *Rosa26mTmG* . **C:** Results

of the RT-qPCR on single cells from the OFT showing that only the GFP+ cNCC were effectively

depleted of Dullard. Few RFP+ cells present a down regulation of Dullard, probably due to an

ineffective recombination of the *Rosa26mTmG* reporter (but not of the Dullard floxed allele) by

the Cre driver. **D:** Percentage of living *Pax3^{Cre}/Wnt1^{Cre}; Dullard$^{flox/flox}$* embryos before and after

E12.5 (percentage calculated on 25 *Pax3^{Cre}* and 40 *Wnt1^{Cre}* litters recovered at embryonic stages

spanning embryonic gestation).

Figure S2: Absence of Dullard in cNCC does not affect their colonization and survival in the OFT.

A: Left: Three dimensional rendering of cNCC (green) over Pecam (white) after BABB clearing and

confocal acquisition of *Wnt1^{Cre}; Dullard$^{flox/+}$; Rosa26mTmG* and *Wnt1^{Cre}; Dullard$^{flox/flox}$;*

Rosa26mTmG E11.5 embryos. Right: Three dimensional rendering of cNCC (green isosurface)

over Pecam (white) after 3Disco clearing and Lightsheet acquisition of *Pax3^{Cre}; Dullard$^{flox/+}$;*

Rosa26mTmG and *Pax3^{Cre}; Dullard$^{flox/flox}$; Rosa26mTmG* E12 embryos. No defect in the

colonization of the cNCC in the OFT was observed. **B, C, D:** Quantification of cNCC total number

per section of OFT, proliferative cNCC through PH3 staining and apoptotic cNCC through cleaved-

caspase-3 staining. We do not observe any significant difference between genotypes. **E:**

Representation of the OFT and the different levels of sectioning. Quantification of the OFT

transversal area along its proximal-distal axis, which shows a slight reduction in E11.5 *Wnt1^{Cre}*;

*Dullard^{flox/flox}*OFT compared to controls.

Figure S3: Single-cell transcriptional analyses of OFT progenitors at E11.5. A: Graph showing the

linear regression between *Actb* and *Gapdh* Ct values. **B:** Unsupervised clustering heatmap of the

OFT progenitors from *Wnt1^{Cre}; Dullard^{+/flox}; Rosa26mTmG* and *Wnt1^{Cre}; Dullard^{flox/flox}*;

Rosa26mTmG based on the gene expression level of the 44 genes included in the panel (Table

S1). Six different populations can be discriminated, among which the endocardial (Pop1)

progenitors expressing high levels of *Flt1*, *Kdr*, *Nfatc1* and *Tek*, and the pericardial (Pop5)

progenitors expressing high levels of *Wt1*, *Tcf21.* **C:** Statistical distribution of *Wnt1^{Cre}; Dullard^{+/flox}*;

Rosa26mTmG and *Wnt1^{Cre}; Dullard^{flox/flox}; Rosa26mTmG* cNCC in the 5 cNCC clusters defined by

their distinct transcriptional signatures. **D:** Genetic composition of the diffusion components (DC)

1 and 2, on which are based the diffusion maps presented in Fig. 3C. **E:** Expression levels of genes

highlighted in DC1 and 2 and not presented in Fig. 3D.

Figure S4: Analysis of the transcriptional signatures of control and mutant RFP+ CD31- and RFP+

CD31+ (endocardial) progenitors. A: Left: Unsupervised clustering heatmap of the RFP+ CD31-

progenitors from *Wnt1^{Cre}; Dullard^{+/flox}; Rosa26mTmG* and *Wnt1^{Cre}; Dullard^{flox/flox}*;

Rosa26mTmG based on the gene expression level of the 44 genes included in the panel (Table

S1). Five different populations can be discriminated, all composed of both control and mutant

RFP+ CD31+ cells. Right: Unsupervised clustering heatmap of the RFP+ CD31+ (endocardial)

progenitors. Six populations with distinct transcriptional signatures (Pop1-6) emerged from the

clustering, and Pop1 and 6 presented clear segregation of control and mutant endocardial cells.

B: Diffusion maps of the endocardial progenitors according to their cluster of appartenance and

their genotype, and based on the diffusion components (DC) 1 and 2 presented in C. **C:** Genetic

composition of DC 1 and 2, on which are based the diffusion maps presented in B. **D:** Expression

levels of genes highlighted in DC1 and 2.

Figure S5: BMP signaling regulates *Sema3c* expression, cNCC compaction and orientation, as

well as distal myocardialization. A: Transversal sections of *Pax3^{Cre}; Dullard$^{flox/flox}$* and *Pax3^{Cre};*

Dullard$^{+/flox}$ OFT stained with DAPI. The endocardium is delineated with a red line and the inner

side of the myocardium with a greyline. Areas where cNCC are of high density and rounded

nuclear shape are surrounded with a green dotted line, while areas where cNCC are of low

density and elongated nuclear shape are surrounded with a yellow dotted line. In *Wnt1^{Cre};*

Dullard$^{flox/flox}$ mutants, the region of high cNCC density is closer to the endocardium compared to

controls. **B:** Left: Measure, in *Wnt1^{Cre}; Dullard$^{flox/flox}$* and *Wnt1^{Cre}; Dullard$^{+/flox}$* cardiac cushions, of

the angles formed by cNCC nuclei longest axis with the endocardial axis, plotted in a [0;90°]

referential. Right: Mapping of the cNCC in the cushions of representative control and mutant

embryos, color-coded according to their angle to the endocardium. Dark blue represents angles

parallel to the endocardium, while dark red are perpendicular. 5 embryos were analyzed for both

genotypes. **C:** Sema3c ISH and Pecam/GFP immunostaining on OFT transversal sections at distal,

medial and proximal levels, from the indicated genotypes. The areas of *Sema3c* expression

(delineated in green) are highly increased in the *Pax3^{Cre}; Dullard$^{flox/flox}$* embryos compared to

controls. These areas are bigger and closer to the endocardium (delineated in red) at distal levels

in both genotypes, and highest *Sema3c* expression correlates with endocardial breakdown. **D:**

Distal sections of the OFT stained for Myosin Heavy Chain (MF20) show an asymmetric

localization of differentiated myocardium in *Wnt1^{Cre}; Dullard$^{flox/flox}$* embryos. The differentiating

MF20+ myocardium is located in Sema3c+ areas of the myocardium (compare Fig. 4C and Fig.

S5D), which indicates that it belongs to the subpulmonary lineage.

Movie S1/S2: Three dimensional rendering of cNCC (green) over Pecam (white) after BABB

clearing and confocal acquisition of *Wnt1^{Cre}; Dullard$^{flox/+}$; Rosa26mTmG* and *Wnt1^{Cre};*

Dullard$^{flox/flox}$; Rosa26mTmG E11.5 embryos. No defect in the OFT colonization of mutant cNCC is

observed.

Movie S3/S4: Three dimensional rendering of cNCC (green isosurface) over Pecam (white) after

3Disco clearing and Lightsheet acquisition of *Pax3^{Cre}; Dullard$^{flox/+}$; Rosa26mTmG* and *Pax3^{Cre};*

Dullard$^{flox/flox}$; Rosa26mTmG E12 embryos. No defect in the OFT colonization of mutant cNCC is

observed.

Movie S5/S6: Three dimensional rendering of cNCC (green isosurface) over Pecam (red

isosurface) after 3Disco clearing and Lightsheet acquisition of *Pax3^{Cre}; Dullard$^{flox/+}$; Rosa26mTmG*

and *Pax3^{Cre}*; *Dullard$^{flox/flox}$*; *Rosa26mTmG* E12 embryos. An atrophy of the pulmonary artery is

observed.

Table S1: List of TaqMan® gene expression assays (20x, Life Technologies) used for single-cell

qPCRs experiments.

524
525

References

527
1. Brickner ME, Hillis LD, Lange RA. Congenital heart disease in adults. First of two parts. N Engl J Med.
2000 Jan 27;342(4):256–63.

2. Kelly RG. Chapter two - The Second Heart Field. In: Bruneau BG, editor. Current Topics in
Developmental Biology [Internet]. Academic Press; 2012 [cited 2018 Jul 6]. p. 33–65. (Heart
Development; vol. 100). Available from:
http://www.sciencedirect.com/science/article/pii/B9780123877864000026

3. Keyte A, Hutson MR. The Neural Crest in Cardiac Congenital Anomalies. Differ Res Biol Divers. 2012
Jul;84(1):25–40.

4. Bockman DE, Redmond ME, Waldo K, Davis H, Kirby ML. Effect of neural crest ablation on
development of the heart and arch arteries in the chick. Am J Anat. 1987 Dec;180(4):332–41.

5. Phillips HM, Mahendran P, Singh E, Anderson RH, Chaudhry B, Henderson DJ. Neural crest cells are
required for correct positioning of the developing outflow cushions and pattern the arterial valve
leaflets. Cardiovasc Res. 2013 Aug 1;99(3):452–60.

6. Plein A, Calmont A, Fantin A, Denti L, Anderson NA, Scambler PJ, et al. Neural crest–derived SEMA3C
activates endothelial NRP1 for cardiac outflow tract septation. J Clin Invest. 2015 Jul 1;125(7):2661–
76.

7. Waldo K, Miyagawa-Tomita S, Kumiski D, Kirby ML. Cardiac Neural Crest Cells Provide New Insight
into Septation of the Cardiac Outflow Tract: Aortic Sac to Ventricular Septal Closure. Dev Biol. 1998
Apr 15;196(2):129–44.

8. Jiao K, Kulessa H, Tompkins K, Zhou Y, Batts L, Baldwin HS, et al. An essential role of Bmp4 in the
atrioventricular septation of the mouse heart. Genes Dev. 2003 Oct 1;17(19):2362–7.

9.   Liu W, Selever J, Wang D, Lu M-F, Moses KA, Schwartz RJ, et al. Bmp4 signaling is required for
outflow-tract septation and branchial-arch artery remodeling. Proc Natl Acad Sci U S A. 2004 Mar
30;101(13):4489–94.

10.   McCulley DJ, Kang J-O, Martin JF, Black BL. BMP4 is required in the anterior heart field and its
derivatives for endocardial cushion remodeling, outflow tract septation, and semilunar valve
development. Dev Dyn. 2008 Nov 1;237(11):3200–9.

11.   Jia Q, McDill BW, Li S-Z, Deng C, Chang C-P, Chen F. Smad signaling in the neural crest regulates
cardiac outflow tract remodeling through cell autonomous and non-cell autonomous effects. Dev
Biol. 2007 Nov 1;311(1):172–84.

12.   Stottmann RW, Choi M, Mishina Y, Meyers EN, Klingensmith J. BMP receptor IA is required in
mammalian neural crest cells for development of the cardiac outflow tract and ventricular
myocardium. Dev Camb Engl. 2004 May;131(9):2205–18.

13.   Tang S, Snider P, Firulli AB, Conway SJ. Trigenic neural crest-restricted Smad7 over-expression results
in congenital craniofacial and cardiovascular defects. Dev Biol. 2010 Aug 1;344(1):233–47.

14.   Sakaguchi M, Sharmin S, Taguchi A, Ohmori T, Fujimura S, Abe T, et al. The phosphatase Dullard
negatively regulates BMP signalling and is essential for nephron maintenance after birth. Nat
Commun. 2013 Jan 29;4:1398.

15.   Satow R, Kurisaki A, Chan T, Hamazaki TS, Asashima M. Dullard Promotes Degradation and
Dephosphorylation of BMP Receptors and Is Required for Neural Induction. Dev Cell. 2006
Dec;11(6):763–74.

16.   Urrutia H, Aleman A, Eivers E. Drosophila Dullard functions as a Mad phosphatase to terminate BMP
signaling. Sci Rep. 2016 Aug 31;6:32269.

17.   Bruce DL, Sapkota GP. Phosphatases in SMAD regulation. FEBS Lett. 2012 Jul 4;586(14):1897–905.

18.   Danielian PS, Muccino D, Rowitch DH, Michael SK, McMahon AP. Modification of gene activity in
mouse embryos in utero by a tamoxifen-inducible form of Cre recombinase. Curr Biol. 1998 Dec
3;8(24):S1–2.

19.   Engleka KA, Gitler AD, Zhang M, Zhou DD, High FA, Epstein JA. Insertion of Cre into the Pax3 locus
creates a new allele of Splotch and identifies unexpected Pax3 derivatives. Dev Biol. 2005 Apr
15;280(2):396–406.

20.   Muzumdar MD, Tasic B, Miyamichi K, Li L, Luo L. A global double-fluorescent Cre reporter mouse.
genesis. 2007 Sep 1;45(9):593–605.

21.   Brown CB, Feiner L, Lu M-M, Li J, Ma X, Webber AL, et al. PlexinA2 and semaphorin signaling during
cardiac neural crest development. Development. 2001 Aug 15;128(16):3071–80.

22.   Jiang X, Rowitch DH, Soriano P, McMahon AP, Sucov HM. Fate of the mammalian cardiac neural
crest. Development. 2000 Apr 15;127(8):1607–16.

23. van den Hoff MJB, Moorman AFM, Ruijter JM, Lamers WH, Bennington RW, Markwald RR, et al.
Myocardialization of the Cardiac Outflow Tract. Dev Biol. 1999 Aug 15;212(2):477–90.

24. Phillips HM, Murdoch JN, Chaudhry B, Copp AJ, Henderson DJ. Vangl2 Acts via RhoA Signaling to
Regulate Polarized Cell Movements During Development of the Proximal Outflow Tract. Circ Res.
2005 Feb 18;96(3):292–9.

25. Bajolle F, Zaffran S, Kelly RG, Hadchouel J, Bonnet D, Brown NA, et al. Rotation of the Myocardial
Wall of the Outflow Tract Is Implicated in the Normal Positioning of the Great Arteries. Circ Res
[Internet].   2006   Feb   17   [cited   2018   Sep   21];   Available   from:
https://www.ahajournals.org/doi/abs/10.1161/01.RES.0000202800.85341.6e?url_ver=Z39.88-
2003&rfr_id=ori:rid:crossref.org&rfr_dat=cr_pub%3dpubmed

26. Rana MS, Théveniau-Ruissy M, Bono CD, Mesbah K, Francou A, Rammah M, et al. Tbx1 Coordinates
Addition of Posterior Second Heart Field Progenitor Cells to the Arterial and Venous Poles of the
HeartNovelty and Significance. Circ Res. 2014 Oct 10;115(9):790–9.

27. Bier E, De Robertis EM. BMP gradients: A paradigm for morphogen-mediated developmental
patterning. Science. 2015 Jun 26;348(6242):aaa5838–aaa5838.

28. Jones CM, Lyons KM, Hogan BL. Involvement of Bone Morphogenetic Protein-4 (BMP-4) and Vgr-1
in morphogenesis and neurogenesis in the mouse. Development. 1991 Feb 1;111(2):531–42.

29. Zhang J, Chang JYF, Huang Y, Lin X, Luo Y, Schwartz RJ, et al. The FGF-BMP Signaling Axis Regulates
Outflow Tract Valve Primordium Formation by Promoting Cushion Neural Crest Cell
DifferentiationNovelty and Significance. Circ Res. 2010 Nov 12;107(10):1209–19.

30. Choi M, Stottmann RW, Yang Y-P, Meyers EN, Klingensmith J. The Bone Morphogenetic Protein
Antagonist Noggin Regulates Mammalian Cardiac Morphogenesis. Circ Res. 2007 Feb 2;100(2):220–
8.

31. Galvin KM, Donovan MJ, Lynch CA, Meyer RI, Paul RJ, Lorenz JN, et al. A role for Smad6 in
development and homeostasis of the cardiovascular system. Nat Genet. 2000 Feb;24(2):171–4.

32. Luo Y, High FA, Epstein JA, Radice GL. N-cadherin is required for neural crest remodeling of the
cardiac outflow tract. Dev Biol. 2006 Nov 15;299(2):517–28.

33. Kim HY, Jackson TR, Davidson LA. On the role of mechanics in driving mesenchymal-to-epithelial
transitions. Semin Cell Dev Biol. 2017 Jul 1;67:113–22.

34. Feiner L, Webber AL, Brown CB, Lu MM, Jia L, Feinstein P, et al. Targeted disruption of semaphorin
3C leads to persistent truncus arteriosus and aortic arch interruption. Development. 2001 Aug
15;128(16):3061–70.

35. Toyofuku T, Yoshida J, Sugimoto T, Yamamoto M, Makino N, Takamatsu H, et al. Repulsive and
attractive semaphorins cooperate to direct the navigation of cardiac neural crest cells. Dev Biol.
2008 Sep 1;321(1):251–62.

36. Kodo K, Shibata S, Miyagawa-Tomita S, Ong S-G, Takahashi H, Kume T, et al. Regulation of Sema3c
and the Interaction between Cardiac Neural Crest and Second Heart Field during Outflow Tract
Development. Sci Rep [Internet]. 2017 Jul 28 [cited 2018 Jan 8];7. Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5533775/

37. Keyte AL, Alonzo-Johnsen M, Hutson MR. Evolutionary and Developmental Origins of the Cardiac
Neural Crest: Building a Divided Outflow Tract. Birth Defects Res Part C Embryo Today Rev. 2014
Sep;102(3):309–23.

38. Belle M, Godefroy D, Couly G, Malone SA, Collier F, Giacobini P, et al. Tridimensional Visualization
and Analysis of Early Human Development. Cell. 2017 Mar;169(1):161-173.e12.

39. Gopalakrishnan S, Comai G, Sambasivan R, Francou A, Kelly RG, Tajbakhsh S. A Cranial Mesoderm
Origin for Esophagus Striated Muscles. Dev Cell. 2015 Sep;34(6):694–704.

40. Chotteau-Lelièvre A, Dollé P, Gofflot F. Expression Analysis of Murine Genes Using In Situ
Hybridization With Radioactive and Nonradioactively Labeled RNA Probes. In: Darby IA, Hewitson
TD, editors. In Situ Hybridization Protocols [Internet]. Totowa, NJ: Humana Press; 2006 [cited 2019
Jan 29]. p. 61–87. (Methods in Molecular Biology™). Available from: https://doi.org/10.1385/1-
59745-007-3:61

41. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source
platform for biological-image analysis. Nat Methods. 2012 Jul;9(7):676–82.

42. Valente M, Resende TP, Nascimento DS, Burlen-Defranoux O, Dupont B, Cumano A, et al. HSA+
immature cardiomyocytes persist in the adult heart and expand after ischemic injury. bioRxiv. 2019
Jan 13;518530.

A

B

C

A

$Pax3^{Cre}:Dullard^{+/flox}$ $Pax3^{Cre}:Dullard^{flox/flox}$

Pecam   Pecam

Ao   Pa   Ao   Pa

Distal   Distal
Medial   Medial
Proximal   Proximal

500µm   E12   500µm   E12

B

$Wnt1^{Cre}:R26^{mTmG}$ $Pax3^{Cre}:R26^{mTmG}$

$Dullard^{flox/+}$ $Dullard^{flox/flox}$ $Dullard^{flox/+}$ $Dullard^{flox/flox}$

DAPI   cNCC (GFP)   Pecam

Distal

Medial

Proximal

50µm   E11.5

A

*Wnt1^Cre^:_R26^mTmG_*

Medial

*Dullard^flox/+^*

DAPI

Ao

Pa

50μm

E11.5

*Dullard^flox/flox^*

DAPI

Ao

Pa

Ao — Pa

Ao — Pa

B

cNCCs distance to endocardium (μm)

***

Wnt1^Cre^:Dullard^+/flox^   Wnt1^Cre^:Dullard^flox/flox^

cNCCs distance to closest nuclei (μm)

***

Wnt1^Cre^:Dullard^+/flox^   Wnt1^Cre^:Dullard^flox/flox^

C

*Wnt1^Cre^:_R26^mTmG_*

*Dullard^flox/+^*

Sema3c   DAPI cNCC (GFP) Pecam

*Dullard^flox/flox^*

Sema3c   DAPI cNCC (GFP) Pecam

Distal

Pa

Ao

Pa

Ao

50μm

E11.5

Pa

Ao

Pa

Ao

Medial

Ao

Pa

Ao

Pa

Ao

Pa

Ao

Pa

Proximal

Ao

Pa

Ao

Pa

Ao

Pa

Ao

Pa

D

*Wnt1^Cre^:_R26^mTmG_*

*Dullard^flox/+^*

DAPI cNCC (GFP)   MF20

Ao

Pa

50μm

E11.5

*Dullard^flox/flox^*

DAPI cNCC (GFP)   MF20

Ao

Pa