Reassessment of the basis of cell size control based on analysis of cell-to-cell variability

Giuseppe Facchetti[1,*], Benjamin Knapp[2], Fred Chang[2,*], Martin Howard[1,*]

[1]John Innes Centre, Computational and Systems Biology, Norwich, United Kingdom

[2]University of California San Francisco, Cell and Tissue Biology, San Francisco, CA, USA

* Corresponding authors

Giuseppe Facchetti   Giuseppe.Facchetti@jic.ac.uk

Fred Chang   Fred.Chang@ucsf.edu

Martin Howard   Martin.Howard@jic.ac.uk

ABSTRACT

Fundamental mechanisms governing cell size control and homeostasis are still poorly understood. The relationship between sizes at division and birth in single cells is used as a metric to categorize the basis of size homeostasis [1-3]. Cells dividing at a fixed size regardless of birth size (sizer) are expected to show a division-birth slope of 0, whereas cells dividing after growing for a fixed size increment (adder) have an expected slope of +1 [4]. These two theoretical values are, however, rarely experimentally observed. For example, rod-shaped fission yeast *Schizosaccharomyces pombe* cells, which divide at a fixed surface area [5, 6], exhibit a division-birth slope for cell lengths of 0.25±0.02, significantly different from the expected sizer value of zero. Here we investigate possible reasons for this discrepancy by developing a mathematical model of sizer control including the relevant sources of variation. Our results support *pure* sizer control and show that deviation from zero slope is exaggerated by measurement of an inappropriate geometrical quantity (e.g., length instead of area), combined with cell-to-cell radius variability. The model predicts that mutants with greater errors in size sensing or septum positioning paradoxically appear to behave as better sizers. Furthermore, accounting for cell width variability, we show that pure sizer control can in some circumstances reproduce the apparent adder behaviour observed in *E. coli*. These findings demonstrate that analysis of geometric variation can lead to new insights into cell size control.

SIGNIFICANCE

How cells control their size is a fundamental but still open question. It has been assumed that the two principle methods of control, namely sizer (where cells always divide at a fixed size) and adder (where cells always add a fixed size increment), can be identified because the correlation between size at birth and size at division is equal to 0 and to +1, respectively. However, experiments mainly provide numbers in between these two values. We show that use of cell length as the size measure, together with cell-to-cell radius variability, can explain this discrepancy and support pure sizer control in fission yeast. Surprisingly, the same phenomenon might exclude adder control in bacteria like *E. coli*.

INTRODUCTION

Cell growth and cell division are coordinated during the cell cycle such that the size of a cell is maintained around a target value specific for each organism and cell type. However, how this outcome is achieved is still largely an open question. Investigations of cell size control intrinsically require a quantitative experimental approach, with many investigations focusing on the simple geometries of yeasts and bacteria. In fission yeast, analyses of individual cells show that they grow to a surface area of about 160 μm^2 before entering mitosis and dividing [6]. Phenomenological characterization of size homeostasis, measuring birth and division sizes of individual single cells, has shown that cells grow to a specified size regardless of birth size. These findings suggest that these cells monitor their own size, dividing at the same size regardless of the size at birth [1, 4, 7]. This principle is called sizer control. On the other hand, the same analysis on *E. coli* data reveals that these cells appear to add a constant size increment during each cell cycle [3, 8], so-called adder control. These cells show a positive correlation between size at birth and size at division [4, 9], so that shorter (longer) cells tend to divide shorter (longer). Theoretical studies have further investigated adder control in terms of robustness to stochastic perturbations and their consequences for the duration of different cell cycle phases [10, 11].

The interpretation of these measurements assumes an unambiguous correspondence between the observed behaviour (division-birth size correlation) and the underlying basis of size control. No correlation implies cells with pure sizer control; a perfect correlation implies cells with pure adder control. However, experimental data has revealed correlations that lie in between these two cases, results which have challenged the notion of a simple basis for size control. As a result, controversies over the basis of size control persist even in *E. coli* (sizer versus adder [12]), as well as in budding yeast (sizer versus adder [2, 13]), whereas a recent study has proposed a combination of a timer (fixed time duration cell cycle) and an adder for *C. crescentus* [14]. Because of its stereotypical shape, and greater available understanding, this work considers fission yeast as a reference model. Even in this case, the measured division-birth correlation is significantly different from zero, casting some doubt on the sizer hypothesis [15].

Previous work showed that size homeostasis in fission yeast is based on total surface area sensing (rather than on cell length or volume sensing) [5, 6]. Quantitative measurements support the idea that this surface area control is achieved by accumulation of Cdr2 in protein clusters ("nodes") in a cortical band around the nucleus. As a result of this localization, the Cdr2 local nodal density scales with cell surface area, which can then trigger mitosis via thresholding [5, 6]. Moreover, through use of a *cdr2-T166A rga2*Δ mutant, cell size homeostasis was successfully switched to length-based size control, confirming the key role of Cdr2 protein in the mechanism [6]. Critical to these conclusions were analyses of mutant cells with altered widths, using *rga2*Δ (thinner) and *rga4*Δ (fatter) mutants [6, 16, 17], which allowed for a robust distinction to be made between size controls based on length, area or volume. However, most data from the literature uses length as

the measure of cell size [1, 7, 15] and for wild-type cells shows a significantly positive division-birth slope

(approximately from 0.2 to 0.3), suggesting that cells might inherit and preserve some elements of size

information from the previous cell cycle, similar to adder behaviour. Our data (Fig. 1A) shows a similarly

positive division-birth slope of around 0.25. Clearly, these findings are not straightforwardly consistent with

pure sizer control [6].

This non-zero slope raises the possibility that cells may display pure size control that is obscured by

measurement errors or other sources of variability (Fig. 1B). Alternatively, cells might adopt an imprecise

implementation or some "impure" mixture of adder and sizer controls (Fig. 1C). To test these possibilities, we

sought to determine what division-birth correlations would be generated by pure sizer (or adder) control, if

the relevant sources of variation were properly accounted for. Our first result is that we can explain all the

correlations experimentally observed for fission yeast in terms of pure sizer control only. Experimentally

measuring a geometrical measure of cell size (length) that is different to that intrinsically used by the cells in

their size control (area), together with radius variability within the cell population, can generate a large shift

of the division-birth slope to higher values. We surprisingly found that a reduction in size-sensing precision by

the cell, or in the precision of placement of the division plane, can paradoxically appear to generate better

sizer behaviour in a division-birth plot. These insights also apply to size control in *E. coli*, indicating that our

approach may improve the assessment of cell size homeostasis control in many cell types.

RESULTS

Sources of variation and mathematical model

In order to improve our understanding of the variety of size homeostasis behaviours, we developed a simple

mathematical model of pure area-based sizer control in fission yeast, including the relevant sources of

variation in cell size regulation and measurement (Materials and Methods). Since a sizer mechanism is based

on size sensing only, the model does not contain time as a variable and, therefore, we do not need to include

any error in the duration of the cell cycle. We aimed to test whether such a model is sufficient to reproduce

the apparently imperfect sizer behaviour observed above. We assumed that each source of variation is

Gaussian distributed, with the free parameters being the following standard deviations: σ_μ (error μ that cells

make in sensing size and therefore in mitotic commitment), σ_α (error giving asymmetric placement α of the

division plane with respect to the exact mid-plane of the cell) and σ_ρ (natural cell-to-cell variability ρ of the

radius). We also included σ_ε as the standard deviation of the measurement error ε (µm) due to image

acquisition and segmentation (see Table 1 for the entire list of variables and parameters of the model). As an

example, the measured cell radius (R^*, where in the following the mark * will denote a measured quantity) is

given by $R^* = R(1 + \rho) + \varepsilon$, where R is the mean radius of the cell population (see Materials and Methods).

Similar expressions can also be derived for the length at birth and length at division. From these equations, we

can calculate the coefficient of variation (CV) for all geometrical quantities at division (Materials and

Methods). For instance, in case of the division length, the CV is given by the sum of three contributions:

117
$$\mathrm{CV}_{L_d^*}^2 \approx \sigma_\mu^2 + \sigma_\rho^2 + \left(\frac{2\pi R}{A}\right)^2 \sigma_\varepsilon^2. \tag{1}$$

Compared to simple probabilistic sizer models for which cells divide according to a probability of division as a

function of cell size [2, 18], our equation describes the distribution of sizes at division as a combination of

well-defined sources of error.

We first estimated the model parameters. All parameters, except for the noise σ_α due to asymmetry in

division, were obtained by fitting to the measured CV of radius, length, area and volume at division (which are

all clearly unaffected by σ_α). We utilised images of wild-type cells acquired at different z-planes (to guarantee

good focus). We then generated (and re-segmented) images at four different resolutions, i.e. with different

values of the experimental error σ_ε. This was achieved by binning a square of *n* by *n* pixels into a single pixel,

with *n* ranging from 1 to 4, i.e. four different values of σ_ε (Fig. 2A). Cells were then segmented using an

algorithm specific for rod or quasi-rod shaped cells (Materials and Methods). For each of the four geometrical

quantities (radius, length, area and volume), and for each resolution, we measured the coefficient of variation

of division size. We then fitted these data points with the model equations by tuning the 6 model parameters

(σ_μ, σ_ρ and the four values of σ_ε). The minimal Root Mean-Square Deviation (RMSD) was obtained with $\sigma_\mu =$

6.5% and $\sigma_\rho = 2.7\%$ (Fig. 2B and S1). Finally, through measurement of the localization of the septum, we

directly estimated the asymmetry in division in wild-type cells (Materials and Methods). By knowing σ_ε , we

obtained the value of $\sigma_\alpha = 3.2\%$, consistent with literature reports [19] (Fig. 2C).

From the above estimation, we note the value of 6.5% for the area sensing error, which is smaller than the CV

of the division length (about 7-8%) normally used to indicate the accuracy of size sensing [7]. This initial result

indicates that the sizer mechanism in fission yeast is slightly more precise than previously thought.

Alteration of cell radius variability supports pure sizer control

Most of the analysis reported in the literature on rod-shaped cells is based on length, because length is the

easiest quantity to measure. Therefore, it is important to find an appropriate way to interpret the outcomes

from these measurements. In this case, our data for the wild-type alone shows a positive birth-division slope:

0.25±0.02 (Fig. 1A, S2, consistent with available values [1, 7, 15]). We next investigated whether these

discrepancies were due to errors, and whether this data could support the existence of pure sizer control in

fission yeast.

From the expressions of the measured length at birth (L_b^*) and measured length at division (L_d^*), while still

having underlying area-based size control, we can derive an equation for the theoretical value of the division-

birth slope (Materials and Methods):

$$\text{slope}(L_d^*, L_b^*) = \frac{2\sigma_\rho^2}{H^2 \sigma_\rho^2 + K^2},\tag{2}$$

where $H^2 = 1 - \frac{4\pi R^2}{3A}$ and $K^2 = \left(\sigma_\mu^2 + \sigma_\alpha^2\right)\Big/\left(1 - \frac{4\pi R^2}{3A}\right) + \left(\frac{4\pi R}{A}\right)^2 \sigma_\varepsilon^2 \Big/ \left(1 - \frac{4\pi R^2}{3A}\right)$. The expression defines a

sigmoidal curve , with the radius variability having an important effect (Fig. S3A). The reasons for these

dependencies are as follows. For surface-area based size control, both birth and division lengths strongly

depend on the true value of the radius and on its variability from cell to cell, as cells with a bigger (smaller)

radius divide shorter (longer) and give birth to shorter (longer) daughter cells, like in adder control. A

consequence of this correlation is a positive division-birth slope (Fig. 3A), with a theoretical slope given by Eq.

(2). This effect was already visible in Fig. 1A: due to area sensing, wider mutants have shorter birth and

division lengths (large circles at the bottom-left of the plot), whereas thinner mutant have longer birth and

division lengths (small circles at the top-right of the plot). On the contrary, when all the cells have the same

radius (i.e. $\sigma_\rho = 0$), the true division length will be perturbed by the sensing error only, and will thus be

independent from the length at birth, giving a zero division-birth slope.

To verify whether radius variation can explain the observed division-birth slope based on length, we exploited

the natural variability of the radius that is present in wild-type cells. We modulated the variability of the radii

of a population of wild-type cells by selecting subsets with a measured radius R^* in the range $R \pm w$. For a

given w, we extracted the division-birth slope and radius natural variability of the selected cells (Fig. S3B and

Materials and Methods). By gradually reducing w (i.e. by reducing the radius variability), the experimental

division-birth slope decreased towards zero as predicted by the model, without any further parameter

adjustment (Fig. 3B). We also analysed different combinations of wild-type, thin (*rga2Δ*) and fat (*rga4Δ*) cells.

Each combination had a different variability of the radius and, according to our analysis, therefore also a

different division-birth slope for length. Model predictions, again without any further parameter adjustment,

were in good agreement with the experimental data (Fig. 3C). Overall, these results again support a *pure* area-

based sizer control in fission yeast, where variations in the radius cause division-birth slopes based on length

to deviate far from zero.

Wild-type data alone supports area-based size control

So far, our analysis has assumed surface area sensing. The previous experiments underpinning this result were

based on variability of the cell radius introduced by thin (*rga2Δ*) and fat (*rga4Δ*) mutants [5, 6]. We attempted

to overcome this limitation by again exploiting the natural variability of the radius in wild-type cells to verify

that area sensing was preserved and was not an artefact due to use of the two mutants.

Following the same procedure used for area sensing, we developed a model based on a length sensing

assumption and a model based on a volume sensing assumption (Materials and Methods). For each of the two

models, we derived expressions for the CVs at division and the division-birth slopes for the various geometric

quantities (length, area, volume). As for area sensing (Fig. 2B), the CV equations were used to estimate the

parameters σ_ε, σ_μ and σ_ρ: by tuning their values (separately for length and volume), we found the best fit for

the experimental division size CVs for wild-type cells (Fig. 3D). With the obtained parameter values (Table S1),

the equations for the division-birth slopes were then used to calculate the theoretical value of the slopes. We

found that the experimental CVs but not the experimental wild-type slopes can be reproduced by the model

based on a volume sensing assumption, whereas the model based on a length sensing assumption was unable

to reproduce either (Figs. 3D-E). On the other hand, the model based on area sensing provided a good fit to

both.

Furthermore, for area-based size sensing, one would naively expect that the CV of area at division (A_d^*) should

be smaller than that for length (L_d^*), i.e. $CV_{A_d^*}^2 < CV_{L_d^*}^2$. Our data in Fig. 2B shows that this is only true for

sufficiently small experimental measurement errors, σ_ε. This is again expected, since errors in measurement

affect the area calculation more than for length, as radius only enters the area estimation. Only when σ_ε is

sufficiently small ($\sigma_\varepsilon < R\sigma_\rho$, as predicted by the model, see Materials and Methods) does area sensing

emerge as being less variable.

These results indicate both the importance of precise measurements in order to correctly infer the basis of

cell size control and, without reliance on mutants, that fission yeast sizer control is indeed based on surface

area.

Asymmetric division and defective size sensing can paradoxically appear to enhance sizer behaviour

Besides experimental error and radius variability, the model also includes two further biological sources of

variability: asymmetry and sensing precision. We next studied their role in size homeostasis behaviour, based

on length measurements.

We showed above how variability in the radius can mask sizer behaviour, bringing the division-birth slope

based on length far from zero (Fig. 3B-C). However, for a given value of the radius variability σ_ρ, Eq. 2

indicates that the slope decreases when the asymmetry (σ_α) or the sensing (σ_μ) error increases. This result

means that increased errors in the division process (in term of septum positioning and/or size sensing)

paradoxically generate better sizing behaviour in a division-birth plot, with a slope closer to zero (Fig. S3A).

These counter-intuitive behaviours can be rationalised as follows: in the case of a perfect sizer, an increase in

the asymmetry error does not affect the distribution of the division length. As a consequence, the cloud of

points of the division-birth size homeostasis plot is stretched along the x-direction only, causing a reduction of

the slope (Fig. 4A). A bigger sensing error produces a wider distribution of both division and birth lengths, i.e.

a stretch along both x- and y-directions. However, since the slope value is affected by (reduced) only by the

horizontal stretch due to a wider distribution of birth lengths (Materials and Methods), this case turns out to

be similar to the asymmetry case. In both cases, the wider spread of the data points reduces the statistical

confidence of the slope value. Therefore, to attain similar levels of precision for the slope values as before, a

higher number of segmented cells are required (Fig. S4A).

To verify these counter-intuitive predictions about the asymmetry (σ_α) or the sensing (σ_μ) error, we re-

examined the case of the *pom1Δ* mutant cells, which divide shorter and often display asymmetric divisions as

compared to wild-type cells ($\mathrm{CV}_{L_b^*}$=22% vs 8.4% for the wild-type) [20, 21]. Although formally we cannot

exclude that the *pom1* deletion might induce different geometric size sensing, we assume here that *pom1Δ*

cells still use surface area for triggering mitosis. Moreover, because of the incompressibility of the G2 phase,

cells that have a length at birth above 7-8 μm follow an adder/timer mechanism rather than a sizer

mechanism [7] (grey colour in Fig. 4B). Therefore, to avoid interference from this altered size control and to

exclude the effects of highly asymmetric divisions, we limited our analysis to cells that start the cell cycle at

lengths below ~7 μm (green colour in Fig. 4B). For this subset, the CV of the division length is 9.0%,

significantly higher than the wild-type (7.2%), indicating a higher sensing error by this mutant (estimated at

$\sigma_\mu \approx 8.6\%$). In agreement with the value of 0.14 predicted by the model, the experimental division-birth

slope decreases from +0.25 to +0.15±0.03 (Fig. 4B), making the mutant erroneously appear as a better sizer.

For a repeat experiment on agar (Fig. S4B), with a division length CV of 12.2%, the model predicts a slope of

0.08 in good agreement with the experimental measurement of 0.09±0.07.

Model predictions match length- and volume-based sizer control experiments

In a previous work we manipulated fission yeast size sensing in order to generate Cdr2-dependent cell size

sensing according to cell length, using the *cdr2-T166A rga2Δ* background. Following this reprogramming, cell

division indeed occurred based on length rather than surface area [6]. According to our model for length-

based size control, the division-birth slope for length should be zero regardless of the natural variability of the

radius and of the segmentation error (Materials and Methods). The experimental value for *cdr2-T166A rga2Δ*

of +0.03±0.05 confirms our model prediction and provides additional support for length sensing in this mutant

(Fig. 4C).

In the same work, we showed that division in the *cdr2Δ* mutant is no longer controlled by area and is instead

based on a secondary Cdr2-independent mechanism that moves towards volume control [6]. Similar to

pom1Δ, data from this mutant can be split into two regimes: a sizer-like regime at shorter cell birth lengths

(less than 10 μm) and an adder/timer-like regime for longer cell birth lengths. The division-birth slope

obtained for the former regime is 0.47 ±0.04 (Fig. 4D), which, although still (just) closer to sizer than adder

behaviour, is even further away from the ideal sizer behaviour than the wild-type. Although the CV of the

division length (7.0%) is similar to the wild-type value, an important question to answer is whether *cdr2Δ* is

still implementing a sizer.

For a model based on pure volume sensing, we can again derive the equation for the division-birth slope

(Materials and Methods). The equation and the result indicate that the effect of natural variability of the

radius is bigger for volume sensing than for area sensing (Fig. 3F). Intuitively, this result follows because cell

volume has a quadratic dependence on the radius, whereas area has a linear dependence only. For this

reason, the shift of the division-birth slope away from zero will be more pronounced. Without any parameter

changing, the model prediction is a slope of +0.62, close to the observed value of around +0.5. Therefore, for

cdr2Δ, as for all the cases analysed in this study (Fig. 4E), the appearance of more adder-like behaviour is likely

due to the use of an incorrect geometrical feature (length) and to the effect of variability in the cell radius.

Incorporating a varying radius into the mathematical model

So far, the model was based on the assumption that cell radius does not change from birth to division.

Experimental data, however, show a strong, but not perfect, positive correlation between the two

measurements ($q = 0.72$, Fig. S5A), indicating that some adjustments in the cell width might occur during the

cell cycle. As an additional test, we therefore relaxed the constant radius assumption and developed a

generalised mathematical model varying radius from $R_b = R(1 + \rho)$ at birth to $R_d = q \cdot R_b + R(1 - q)$ at

division. These dynamics guarantee the preservation through generations of cell radius around the average

value *R*. Numerical simulations show that the key features of the model outcomes are preserved. Moreover,

since the variation of radius within one cell cycle is relatively small (Fig. S5A), we did not find significant

changes in our results. As a result, the agreement with the experimental evidence is still good (Fig. S5B).

Adder behaviour can emerge from pure sizer control

With the exception of length sensing in the *cdr2-T166A rga2Δ* mutant, utilising length as the experimental size

measure partially masks pure sizer behaviour by moving the division-birth slope towards adder behaviour, i.e.

towards a value of +1. We therefore asked whether the same effect is present in a pure adder mechanism. In

particular, we focus on an adder mechanism based on cell volume. This assumption allowed us to develop a

model (Materials and Methods) for the widely studied bacterium *E. coli* which shares the same rod-shape as

fission yeast and which is believed to utilize cell volume as the key geometrical quantity for size control [22].

In contrast with other models [23], also in this adder model we did not include any direct errors on the time

durations: rather, variation in the duration of the cell cycle is a consequence of the error in sensing the added

size, which is the fundamental quantity monitored by a cell which implements adder control. Similar to our

observations for pure sizer control based on volume sensing, the effect of natural cell radius variability causes

a monotonic increase of the division-birth length slope from the theoretical adder value of +1 up to +2 (Fig.

S6A). By using 3.5% as an estimation of the radius variability [9], our adder model generates a division-birth

length slope of +1.51±0.03 (Fig. 5A). Robustness of the outcome was tested by additively

increasing/decreasing each standard deviation by up to 1% (or 2%): slope values remained entirely above +1

(Fig. S6B). However, all the experimentally reported division-birth length slopes for *E. coli* range from +0.7 to

+1 [8, 9], i.e. they never overshoot the theoretical pure adder value. This result may suggest that an adder

mechanism is incompatible with the observed behaviour.

Our results on the *cdr2Δ* mutant revealed that, in the case of a volume-based pure sizer, the effect of radius

variability is more pronounced than in the case of area sensing and moves the division-birth length slope

much closer to adder-like behaviour. Running a numerical simulation of such a volume-based sizer model with

$\sigma_\rho = 3.5\%$ (slightly higher than the radius variability in fission yeast) on *E. coli* sized cells [9], we obtained a

division-birth length slope of +0.81±0.02 (Fig. 5B), in agreement with the experimental results from this

bacterium. This finding strongly suggests that sizer behaviour in *E. coli* cannot currently be ruled out.

DISCUSSION

Phenomenological characterization of size homeostasis is a common approach for extracting information

about cell size regulation. Although initial results previously suggested an imperfect or possibly mixed sizer-

like control in fission yeast, we have shown here that *pure* sizer control based on *accurate* sensing of cell size

is implemented in this organism, using a Cdr2-dependent mechanism with area-based control in the wild-type

and length-based control in *cdr2-T166A rga2Δ* [6]. Moreover, we have identified three generalizable aspects

in the analyses of size homeostasis: (i) high accuracy in the measurement of the cell size is necessary to avoid

misleading data (Fig. 2B); (ii) use of the wrong geometrical quantity (e.g., length instead of area), combined

with natural variability of the cell radius, distorts the observed size control behaviours (Fig. 3B-C). While this

effect disappears in the presence of length-based sizer sensing (see *cdr2-T166A rga2Δ*, Fig. 4C), the behaviour

is even more markedly perturbed when sizer control is based on cell volume (see *cdr2Δ* in Fig. 4D and *E. coli* in

Fig. 5B); (iii) asymmetric cell division and, more surprisingly, a reduction in the precision of size sensing, can

paradoxically appear to enhance sizer behaviour. As shown by the case of *pom1Δ* in fission yeast, an

apparently stronger sizer behaviour (lower division-birth slope) may be due to a loss of precision in size

control (Fig. 4B). These demonstrations show that correct interpretation of size homeostasis behaviour

requires accurate measurements of multiple geometrical quantities. Furthermore, "intermediate" behaviour

(as in wild-type cells) may not necessarily be due to imperfect size control, nor to a combination of two or

more distinct size controls (e.g. sizer and adder). Even limited variability can move the division-birth length

slope from 0 towards +1 in the case of pure sizer control and from +1 towards +2 in case of pure adder

control, thus completely obscuring the underlying fundamental basis of size control (Fig. 6).

An important aspect that has emerged in our analysis is the effect of radius variability. One might expect that

the small variations found from cell to cell might have only a marginal effect on size control behaviour.

Surprisingly, the curve and data in Fig. 3C show that, in the case of area-based sizer sensing in fission yeast, a

radius variability of only 7% is sufficient to move the division-birth slope close to +1, making pure sizer control

wrongly appear as a pure adder. The equations of the model allow us to understand how any particular size

homeostasis behaviour can emerge from antagonism between inaccuracy in the division process (asymmetry

and size sensing) and radius variability. In the approximation of the slope expression given in Eq. (2), the first

non-zero term is $2\sigma_\rho^2/K^2$. If we ignore the contribution of the experimental error σ_ε^2, this term can be

approximated by $2\sigma_\rho^2/(\sigma_\alpha^2 + \sigma_\mu^2)$, which clearly displays the antagonism between the two effects. In our case,

the combination of the estimated values for asymmetry ($\sigma_\alpha = 3.2\%$) and size sensing ($\sigma_\mu = 6.5\%$) errors

gives a total standard deviation of 7.3%. Therefore, we can expect that a radius variability of the same order

of magnitude can neutralize the effect of these two sources of variation and lead to adder-like behaviour.

For the same reason, despite the adder phenotype, sizer control may still not be ruled out in *E. coli* (Fig. 5),

consistent with some alternative theories [12, 22, 24]. Furthermore, according to our analysis, adder control

would tend to show behaviour closer to a timer (slope significantly above +1). One caveat is that our adder

model considers the simple case of a constant cell size increment from birth to division. It has been

alternatively suggested that a constant volume (per origin) is added between two initiations and that cells

then divide a constant time after initiation [22]. Although further investigation is needed to better understand

this misalignment between the basis of size control and the resulting division-birth slope behaviour, our

results suggests that *E. coli* might not implement adder control.

Overall, the analysis presented here shows that pure sizer control is able to reproduce a large variety of cell

size homoeostasis behaviours. Our results, with careful attention to both sources of variation in size control

and to geometrical aspects preserved by the cell, could be widely relevant in deciphering size homeostasis

behaviour in many cell types. Such accurate analyses of cell size behaviours will be critical for dissecting the

underlying molecular mechanisms responsible.

360
361
Figure 1: Mapping between cell behaviour and underlying basis of size control. (**A**) Division-birth correlation for wild-type (FC15, n=326) based on measurement of cell length. Size of circles proportional to cell radius. Binned data, with mean value ± standard error, also shown (dark circles), together with best fit line, with indicated slope. (**B**) Theoretical assumption of a one-to-one mapping between division-birth slope and size control, where slope=0 implies sizer control and slope=+1 implies adder control. No intermediate values for the slope are expected. (**C**) Slope values between 0 and +1 could be interpreted as a mix between sizer and adder control (or an imprecise sizer or imprecise adder).
369

Figure 2: Motivation and model parameter estimation. (**A**) Reduction of image resolution by pixel binning:
reported value of resolution is res=1/n, (where n x n pixels have been averaged and condensed into a single
pixel; res=1 is the experimentally acquired image). (**B**) Fitting of the observed Coefficient of Variations at
division (symbols) by tuning the parameters of the model (curves), at different image resolutions and
therefore different experimental errors, σ_ε (wild-type strain FC15, n=326). See also Fig. S1. Dashed line is
located where $\sigma_\varepsilon = R\sigma_\rho$: for smaller errors, area sensing becomes apparent, i.e. $CV^2_{A_d^*} < CV^2_{L_d^*}$. (**C**)
Measurement of division asymmetry and its fitting with a Gaussian distribution to determine σ_α (n=106),
where $L^*_{s\text{-}t}$ is the measured distance between the division septum and cell tip. Experimental data are fitted
with $\sigma_{\alpha*} = 0.035$. Since $\sigma^2_{\alpha*} \approx \sigma^2_\alpha + 5\sigma^2_\varepsilon/L^2$, we can extract $\sigma_\alpha = 0.032$ (Materials and Methods).

Figure 3: Fission yeast implements a pure area-based sizer (**A**) Intuitive explanation for how, for area sensing, variability in cell radius causes deviation from perfect sizer behaviour. Ellipses represent cloud of data points and dashed lines their linear fitting. (**B**) Inset: radius distribution in red and example of selection of cells with radius in range of ± w = ± 0.1 μm around the mean value (grey shadow). Main plot: for given w (value in squared brackets on x-axis), division-birth slope for length is measured on subset of selected cells (black circle, slope ± SD) and radius variability is calculated (Materials and Methods and Fig. S3B). Predicted slope based on area sensing is calculated with Eq. 2 (green curve). (**C**) Predicted area-based (green curve) and measured division-birth slope for length (mean value ± SD) obtained by combining mutants with different radii. (**D**) Fitting of the observed CV at division (symbols) by tuning the parameters of the model (curves), at different image resolutions and therefore different experimental errors, σ_ε. Left panels: length sensing model. Right panel: volume sensing model (see also Table S1). (**E**) Bars report the theoretical values of division-birth slope for length for each possible geometric size sensing model (based on length, area or volume). Area sensing is the one that provides the best agreement with the experimental data (black line ± SD, continuous and dashed lines). Panel D uses the same data as in Fig. 1A-C. For all other panels data for wildtype strain on agar (FC15, n=326) with automated segmentation.

Figure 4: Counter-intuitive predictions arising from asymmetric division errors and consequences of length or volume sensing. (**A**) Possible explanation of how an increase in the asymmetry error at division brings the birth-division slope closer to zero. Black bars indicate the width of the distribution of the division length. (**B**) Size homeostasis plot for *pom1Δ* (FC2063, n=1757, data from [6]). Slopes of reported fit lines are 0.15±0.03 for the sizer regime (green colour, length at birth less than 7 μm) and +0.86±0.03 for the adder/timer-like regime (grey colour, length at birth greater than 7 μm). In the sizer regime, division length has a CV of 9.0%, higher than the wild-type, but has a lower birth-division slope (+0.15 versus +0.25 for the wild-type as reported in Fig. 1A), as predicted by the model. (**C**) Size homeostasis data for the *cdr2-T166A rga2Δ* mutant which implements length sensing shows near perfect sizer behaviour (FC3218, n=1785, data from [6]). (**D**) Size homeostasis data for the *cdr2Δ* mutant, which implements volume sensing. Regression line and reported slope of 0.47±0.04 refers to the binned data with length at birth smaller than 10 μm (green colour). Cells with length at birth greater than 10 μm (grey colour) show a slope of +0.88 ± 0.16 (FC3161, n=1046). Binned data (with mean value ± standard error) shown as bold circles in panels B-D. (**E**) Summary of the agreement between experimental slopes and values obtained by the model. Data points show the mean value ± standard error (see also Fig. S5B).

Figure 5: Radius variability and sizer control reproduce the apparent adder behaviour in *E. coli*. (**A**) Simulation of division-birth correlation based on length measurements for *E. coli* utilising pure adder control based on cell volume. (**B**) Same plot in case of pure sizer control based on cell volume. Both plots of panels A and B have been obtained with R=0.55 μm, V=3.77 μm³, 3.5% as the natural radius variability and the same noise used for fission yeast (Fig. S6B shows the effect of perturbations of these values). Binned data, with mean value ± standard error, also shown (dark circles), as well as best fit line, with indicated slope.

Our results

Figure 6: Summary. How a pure sizer mechanism can show a wide range of behaviours. Noise and error lead to an increase of the slope for sizer control (and also for adder control).

441 **TABLE 1: Variables and parameter values of the model**.

Symbols	Description	units	Value for each strain				
			WT	*pom1Δ* (subset of short cells only)	*cdr2-T166A rga2Δ*	*cdr2Δ*	*E. coli*
ρ, σ_ρ	Gaussian random variable with standard deviation σ_ρ (value reported here) describing cell-to-cell radius variability	%	2.7	2.7	2.7	2.7	3.5
μ_b, μ_d, σ_μ	Gaussian random variables with standard deviation σ_μ (value reported here) describing cell error in sensing its size	%	6.5	8.6	6.5	6.5	6.5 [a]
α, σ_α	Gaussian random variable with standard deviation σ_α (value reported here) describing the asymmetry in division, defined as $$(2L^*_{s-t} - L^*_d)/L^*_d$$ where L^*_{s-t} is measured distance between the division septum and cell tip.	%	3.2	3.2	3.2	3.2	3.2 [a]
$\varepsilon_b, \varepsilon_d, \sigma_\varepsilon$	Gaussian random variables with standard deviation σ_ε (value reported here) describing experimental error in the measurement at different resolutions res.=1 (0.0635 µm/pixel) res.=0.5 (0.127 µm/pixel) res.=0.33 (0.190 µm/pixel) res.=0.25 (0.254 µm/pixel)	µm	0.045 0.052 0.067 0.078	0.06 [b]	0.06 [b]	0.06 [b]	0.04 [b]
R	Mean cell radius	µm	1.96	1.89	1.58	1.95	0.55
L	Target cell length at division in case of length sensing (mean of the experimental measurement)	µm	n.a.	n.a.	18.3	n.a.	n.a.
A	Target cell area at division in case of area sensing (mean of the experimental measurement)	µm²	164	158	n.a.	n.a.	n.a.
V	Target cell volume at division in case of volume sensing (mean of the experimental measurement)	µm³	n.a.	n.a.	n.a.	183	3.77
q	Persistence of the cell radius from birth to division	-	0.72	0.72	0.72	0.72	-

443 [a]: Simulations with additively increasing/decreasing each standard deviation by up to 1% (or 2%) are reported
444 in Fig. S6.
445 [b]: Estimation based on the resolution of the images.

REFERENCES

1. Jun, S., and Taheri-Araghi, S. (2015). Cell-size maintenance: universal strategy revealed. Trends in microbiology *23*, 4-6.

2. Chandler-Brown, D., Schmoller, K.M., Winetraub, Y., and Skotheim, J.M. (2017). The adder phenomenon emerges from independent control of pre-and post-start phases of the budding yeast cell cycle. Current Biology *27*, 2774-2783. e2773.

3. Sauls, J.T., Li, D., and Jun, S. (2016). Adder and a coarse-grained approach to cell size homeostasis in bacteria. Current opinion in cell biology *38*, 38-44.

4. Facchetti, G., Chang, F., and Howard, M. (2017). Controlling cell size through sizer mechanisms. Current Opinion in Systems Biology *5*, 86-92.

5. Pan, K.Z., Saunders, T.E., Flor-Parra, I., Howard, M., and Chang, F. (2014). Cortical regulation of cell size by a sizer cdr2p. Elife *3*, e02040.

6. Facchetti, G., Knapp, B., Flor-Parra, I., Chang, F., and Howard, M. (2019). Reprogramming Cdr2-Dependent Geometry-Based Cell Size Control in Fission Yeast. Current Biology.

7. Wood, E., and Nurse, P. (2013). Pom1 and cell size homeostasis in fission yeast. Cell cycle *12*, 3417-3425.

8. Campos, M., Surovtsev, I.V., Kato, S., Paintdakhi, A., Beltran, B., Ebmeier, S.E., and Jacobs-Wagner, C. (2014). A constant size extension drives bacterial cell size homeostasis. Cell *159*, 1433-1446.

9. Taheri-Araghi, S., Bradde, S., Sauls, J.T., Hill, N.S., Levin, P.A., Paulsson, J., Vergassola, M., and Jun, S. (2015). Cell-size control and homeostasis in bacteria. Current Biology *25*, 385-391.

10. Lin, J., and Amir, A. (2017). The effects of stochasticity at the single-cell level and cell size control on the population growth. Cell systems *5*, 358-367. e354.

11. Barber, F., Ho, P.-Y., Murray, A.W., and Amir, A. (2017). Details matter: noise and model structure set the relationship between cell size and cell cycle timing. Frontiers in cell and developmental biology *5*, 92.

12. Wallden, M., Fange, D., Lundius, E.G., Baltekin, Ö., and Elf, J. (2016). The synchronization of replication and division cycles in individual E. coli cells. Cell *166*, 729-739.

13. Soifer, I., Robert, L., and Amir, A. (2016). Single-cell analysis of growth in budding yeast and bacteria reveals a common size regulation strategy. Current Biology *26*, 356-361.

14. Banerjee, S., Lo, K., Daddysman, M.K., Selewa, A., Kuntz, T., Dinner, A.R., and Scherer, N.F. (2017). Biphasic growth dynamics control cell division in Caulobacter crescentus. Nature microbiology *2*, 17116.

15. Fantes, P. (1977). Control of cell size and cycle time in Schizosaccharomyces pombe. Journal of cell science *24*, 51-67.

16. Villar-Tajadura, M.A., Coll, P.M., Madrid, M., Cansado, J., Santos, B., and Pérez, P. (2008). Rga2 is a Rho2 GAP that regulates morphogenesis and cell integrity in S. pombe. Molecular microbiology *70*, 867-881.

17. Das, M., Wiley, D.J., Medina, S., Vincent, H.A., Larrea, M., Oriolo, A., and Verde, F. (2007). Regulation of cell diameter, For3p localization, and cell symmetry by fission yeast Rho-GAP Rga4p. Molecular biology of the cell *18*, 2090-2101.

18. Patterson, J.O., Rees, P., and Nurse, P. (2019). Noisy Cell-Size-Correlated Expression of Cyclin B Drives Probabilistic Cell-Size Homeostasis in Fission Yeast. Current Biology.

19. Saha, S., and Pollard, T.D. (2012). Anillin-related protein Mid1p coordinates the assembly of the cytokinetic contractile ring in fission yeast. Molecular biology of the cell *23*, 3982-3992.

20. Padte, N.N., Martin, S.G., Howard, M., and Chang, F. (2006). The cell-end factor pom1p inhibits mid1p in specification of the cell division plane in fission yeast. Current Biology *16*, 2480-2487.

21. Huang, Y., Chew, T.G., Ge, W., and Balasubramanian, M.K. (2007). Polarity determinants Tea1p, Tea4p, and Pom1p inhibit division-septum assembly at cell ends in fission yeast. Developmental cell *12*, 987-996.

22. Zheng, H., Ho, P.-Y., Jiang, M., Tang, B., Liu, W., Li, D., Yu, X., Kleckner, N.E., Amir, A., and Liu, C. (2016). Interrogating the Escherichia coli cell cycle by cell dimension perturbations. Proceedings of the National Academy of Sciences *113*, 15000-15005.

23. Eun, Y.-J., Ho, P.-Y., Kim, M., LaRussa, S., Robert, L., Renner, L.D., Schmid, A., Garner, E., and Amir, A.
(2018). Archaeal cells share common size control with bacteria despite noisier growth and division.
Nature microbiology *3*, 148.
24. Si, F., Li, D., Cox, S.E., Sauls, J.T., Azizi, O., Sou, C., Schwartz, A.B., Erickstad, M.J., Jun, Y., and Li, X.
(2017). Invariance of initiation mass and predictability of cell size in Escherichia coli. Current Biology
27, 1278-1287.